CRF neurons establish resilience via stress-history dependent BNST modulation

Authors:

Sherod E Haynes[1,2,3], Anthony Lacagnina[4*], Hyun Seo Seong[1*], Muhammad Afzal[5], Carole Morel[3], Aurelie Menigoz[1], Kanaka Rajan[5], Roger L Clem[4], Helen S Mayberg[5], Donald G. Rannie[1], Larry J. Young[1], Ming-Hu Han[3,4,6†]

Affiliations:

[1]Center for Translational Social Neuroscience, Yerkes National Primate Research Center, Department of Psychiatry and Behavioral Sciences, Emory University, Atlanta, GA 30322, USA

[2]Neuroscience Graduate Program, Emory University, Atlanta, GA 30322, USA

[3]Department of Pharmacological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[4]Nash Family Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[5]Nash Family Center for Advanced Circuit Therapeutics, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[6]Department of Mental Health and Public Health, Faculty of Life and Health Sciences, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, Guangdong, China

*These authors contributed equally

†Corresponding electronic address: ming-hu.han@mssm.edu.

Abstract

34
35
36
Cumulative stress is a major risk factor for developing major depressive disorder (MDD), yet not

everyone experiencing chronic stress develops MDD. In those who do not, it is unclear at what

point, or by what mechanism, a trajectory of stable resiliency emerges. Utilizing a 10-day

repeated social defeat stress model (RSDS) for MDD, we observed that a critical period between

7 and 10 daily defeats marks the phenotypical divergence of resilient from susceptible mice. In

response to ongoing stress, corticotropin-releasing factor (CRF) neurons of the oval nucleus of

the bed nucleus of the stria terminalis (BNSTov) display a sustained increased firing rate in

resilient, but not susceptible mice. This neurophysiological adaptation was self-sustaining, but

only after 7 critical stress exposures, indicating that the process of developing resilience is

dependent on stress history. Our study reveals a novel process by which individuals might persist

in the face of adversity by way of stress-provoked activation, not inhibition of a key CRF limbic

region that establishes a pathway to resilience.

Major depressive disorder (MDD) is a crippling heterogeneous neuropsychiatric condition with

high morbidity and lifetime prevalence, exacerbated by the ongoing global pandemic[1]. MDD is

episodic in nature, with treatment success diminishing with each subsequent depressive

episode[2]. Major life stressors are key risk factors for precipitating the onset of an episode[3,4].

While most people will report at least one major life stressor at some point in their lives, not

everyone will go on to develop MDD. It is also known that depression can emerge after

repeated significant stressor exposure, suggesting that cumulative exposure to stress is

associated with increased depression risk in vulnerable individuals[5–8]. Cumulative stress

causes numerous psychological insults that predispose neuropsychiatric conditions[5,9,10]. Many

studies have explored the individual differences in stress susceptibility or resiliency, but the

process critical to driving the divergence in phenotypes remain elusive. Recent studies suggest

mechanisms involved in the emergence of depression may be distinct from those that maintain

the depressive state[11,12]. With the odds of treatment failure increasing with subsequent

depressive episodes,[2,13] and increased depression risk of cumulative stress, a potentially more

effective strategy would be to target the mechanisms mediating resilience, enhancing the ability

to cope with cumulative stress.

The ability to engage in social and hedonic-like behaviors in the face of psychosocial stress is

an adaptive behavioral strategy, and failing to do so is associated with high morbidity[14–16].

Resiliency is defined as the "process of adapting well in the face of adversity… [to] threats, or

significant sources of stress[15]." Resiliency is unlikely to emerge from the absence of a

pathological stress response but instead is an active process involving a myriad of molecular,

cellular and circuit-level changes in the brain. Repeated social defeat stress (RSDS) induces

robust depression-like behavioral phenotypes in roughly 2/3 of mice[6,10]. The standard 10-day

RSDS protocol has been employed widely to identify and study neurobiological features of

susceptible and resilient subpopulations[6,7,17–23]. Studies on stress vulnerability often test

hypotheses after establishing vulnerable subgroups, making the processes that led to such

phenotypic diversification unclear.

The Bed Nucleus of the Stria Terminalis (BNST) is an important node for integrating sensory

cues, interoception, cognition, and motivational states to enact adaptive responses[24,25]. The

BNST is well-positioned in the social salience network to integrate external cues with internal

states to influence the outcome and context of social interactions[24–27]. The oval nucleus of the

BNST (BNSTov) is a stress-sensitive subregion that is a key candidate region for encoding

stress modulation of social behavior through its projections to areas such as the ventral

tegmental area (VTA)[28–31] and dorsal raphe[32,33]. Corticotrophin-releasing factor (CRF) neurons

of the BNSTov (BNSTov[CRF]) are a significant output of this region and can influence affective

states[34–37]. Since BNSTov[CRF] neurons are thought to promote arousal and adaptive responding

according to stress-related changes in internal state[24,25,27,38–44], we hypothesized they might play

a role in establishing resiliency to repeated social stress.

Using the 10-day RSDS paradigm, we used cell-type selective *ex vivo* electrophysiology,

chemogenetics, optogenetics, and *in vivo* fiber photometry to interrogate the BNSTov[CRF] system

to explore its role in the divergence of susceptible and resilient phenotypes. We observed that

BNSTov[CRF] neurons encode repetitive social stress and undergo adaptation that coincides with

the divergence of resilient and susceptible phenotypes. Unexpectedly, we observed that

resiliency entails cumulative stress-history dependent neuroadaptive changes, whereas

susceptibility emerges in the absence of similar adaptations. Activation of BNSTov[CRF] neurons

using Designer-Receptors-Exclusively-Activated-by-Designer-Drugs (DREADDs) during social

defeat led to a resilient phenotype while inhibition caused susceptibility. Finally, using

RNAscope and optogenetics, we provide intriguing evidence for a potential role of *Crfr1* in

BNSTov[CRF] neurons in mediating resilience.

53
Results:

Susceptible vs. resilience phenotypes emerge between 7 and 10 daily episodes of social

defeat stress.

RSDS produces divergent and enduring resilient/susceptible phenotypes following 10 episodes

of social defeat stress[6,21] has high ethological validity and enables the investigation of the

consequences of stress accumulation[5,7]. To examine the temporal divergences of resilient and

susceptible phenotypes in mice, a modified RSDS stress protocol was employed in which mice

were subjected to discrete numbers of social defeat episodes (SDEs) interspersed with social

interaction (SI), and sucrose preference (SP) tests administered after 1, 4, 7, and 10 SDEs (Fig.

1a-d). Social Interaction (SI) Ratio is a behavioral score of the SI test and measures the time

spent in the area proximal to the enclosure of a novel social target (social interaction zone). To

generate the SI ratio, interaction zone time is calculated both in the absence and presence of

the social target, such that SI ratio $\geqq$ 1 is resilient and <1 is susceptible[21]. Surprisingly, we

found that the susceptible phenotype emerged between 7 and 10 SDEs, reflected by a robust

decrease in social interaction on day 10 compared to day 7 (Fig. 1e, Extended Data Fig 1a) (SI

ratio: 2.062 +/- 0.226 after 7 SDEs to 0.559 +/- 0.093, p<0.0001, n = 11 for susceptible and

1.967 +/- 0.188 (7 SDEs) to 1.554 +/- 0.163 (10 SDEs), p=0.2929, n=11 for resilient). There

were no significant differences in SI ratios after 7 SDEs between mice that went on to become

susceptible vs resilient after 10 SDEs (mean: 2.062 +/- 0.226 vs 1.967 +/- 0.188, two-tailed t-

test, p=0.7483, n= 11/group) (Extended Data Fig. 1b-c). To test whether this was a result of

repeated SI tests, in a separate experiment, we subjected independent groups of mice to either

1, 4, 7, or 10 SDEs, conducting the SI and SP tests only once, and observed a similar effect of

susceptible/resilient subgroups emerging after the seventh episode of social defeat (Extended

Data Fig. 2d-f). Resilient (10 SDEs) mice had an indistinguishable SI ratio from all mice after 7

SDEs, while susceptible (10 SDEs) mice had an SI ratio significantly lower than both groups

(Fig. 1f). In the SP test, resilient mice showed a similar preference for sucrose to all 7 SDE and

stress-naïve control mice. Susceptible mice, by contrast, exhibited a decreased SP relative to 7

SDE and resilient groups (Fig. 1g). Previous studies reported that RSDS produces susceptible

and resilient phenotypes after 10 SDEs in a bimodal distribution[21]. Here, we observed a

unimodal distribution of social interaction toward a novel conspecific in 7 SDE mice but a

bimodal distribution in 10 SDE mice, consistent with the emergence of distinct resilience and

susceptible phenotypes (Fig. 1h,i).   The susceptible population of mice represents a statistically

distinct group compared to 7 SDE and 10 SDE resilient mice (Kolmogorov-Smirnov, p<0.0001;

Fig. 1j). The time spent interacting with a novel conspecific is significantly less in susceptible 10

SDE mice compared to the 7th SDE and 10th SDE resilient mice, which were indistinguishable

from each other (Fig.1k). The divergence into susceptible and resilient phenotypes was

dependent upon the number of SDEs, rather than the passage of time. Mice subjected to 7

consecutive days of social defeat stress displayed social approach behavior when tested on day

11 and day 30, yet when subjected to three additional defeat episodes, social avoidance

emerged (Extended Data Fig. 1g). Thus, it is the number of episodes (SDEs #8-10), not the

duration of time after the 7th SDE, that is responsible for the emergence of susceptible

phenotypes. To gain insight into potential neural mechanisms involved in the phenotypic

divergence, we used c-FOS immunohistochemistry as a proxy of neural activation in mice

subjected to 10 SDEs following a 30-minute social interaction with a non-aggressive novel

conspecific (Extended Data Fig. 2a). While several brain regions implicated in social behavior

and motivation were assessed, only the anterior dorsal BNST showed significant differences in

c-FOS immunoreactivity (c-FOS-ir) between the stress-naïve, susceptible, and resilient mice

(Extended Data Fig. 2b). Given its role in responding to chronic stress, valence, and

motivational processes, we more specifically examined the BNSTov[24]. While no difference in c-

FOS-ir was detected between stress-naïve and 7 SDE mice, there was a significant decrease in

c-FOS-ir in susceptible (10 SDEs) and a significant increase in resilient (10 SDEs) mice,

respectively (Extended Data Fig. 2c-e). Thus, the c-FOS-ir data suggests a neural divergence

in BNSTov in response to social threat that paralleled the divergence in SI defining resilient and

susceptible phenotypes.

Divergence in BNSTov^{CRF} neuronal firing rates tracks the emergence of resilient and

susceptible phenotypes.

CRF neurons of the BNST are a major output source of the oval BNST that are sensitive to

chronic stressors[27,45–47]. Thus, we hypothesized that chronic stress would alter BNSTov^{CRF}

neuronal activity coinciding with the divergence in resilient and susceptible behavioral

phenotypes. To test this hypothesis, *Crf*-ires-*Cre;*ai14 (tdTomato) mice[30,48–51] were subjected to

either 7 or 10 daily SDEs, and cell-attached *ex vivo* electrophysiological recordings were

conducted in the BNSTov (Fig. 2a,b). CRF$^+$, but not CRF$^-$, BNSTov neurons significantly

increased firing rate in 7 SDE mice compared to stress-naïve control mice. By contrast, CRF$^+$

firing in susceptible 10 SDE mice was indistinguishable from that of controls and was lower than

that of 7 SDE and resilient 10 SDE mice. CRF$^-$ neuron firing did not differ between groups (Fig.

2c,d). Moreover, there was a strong correlation between firing rate and social interaction ratio in

CRF$^+$ but not CRF$^-$ neurons in 10 SDE mice (CRF$^+$, R^2=0.5725, *p=0.0113; CRF- R^2=0.06096,

p=0.5219, Fig. 2e). CRF neurons display both burst and non-burst firing patterns[40,52,53] (Fig. 2f).

Burst firing patterns were prominent in resilient and 7 SDE mice, but few were seen in

susceptible and control mice (Fig. 2h). Additionally, percentage spikes within burst were higher

in 7 SDE and resilient compared to susceptible 10 SDE mice (***p=0.0004, one-way ANOVA,

Fig. 2i), but not susceptible or control mice. The number of spikes per burst and number of

bursts per cell was not significantly different amongst the groups (Fig. 2j,k). These data and

correlation analysis suggest the possibility that there is a causal link between the neuronal

activity of BNSTov[CRF] neurons and the divergence of behavioral phenotypes.

BNSTov[CRF] neurons bidirectionally modulate the emergence of resiliency.

To test the hypothesis that BNSTov[CRF] neurons are regulating the maintained resiliency over the

last 3 episodes of RSDS, during which susceptible and resilient phenotypes emerge, we

injected *Crf*-ires-*Cre* mice with AAVs encoding Cre-dependent excitatory (DIO-hM3Dq),

inhibitory (DIO-hM4Di) DREADDs, or a control mCherry construct into the BNSTov and

administered CNO via drinking water[54–56] (Fig. 3a,b). We presaged that this experimental

design afforded the opportunity to modulate neurons over a longer time span than

intraperitoneal (IP) injections would allow with minimal invasiveness, particularly because it was

not clear over when neuroadaptation occurs within the 3-day window. mCherry control mice

exhibited both susceptible and resilient phenotypes (SI ratio <1.0 and $\geqq$1.0, respectively) in

roughly the 60/40 ratio as expected[6,21] (Fig. 3c,f). Interestingly, mice injected with inhibitory DIO-

hM4Di displayed a robust susceptible phenotype, while DIO-hM3Dq mice displayed resilient

phenotypes following CNO drinking water administration (Fig. 3c). Moreover, none of the DIO-

hM4Di + CNO mice went on to develop resilience (0/10, SI >1.0) while 89% (8/9, SI $\geqq$1.0) of

the DIO-hM3Dq were resilient (Fig 3c,f). Notably, the social defeat experience was not affected

by the DREADDs manipulations (Extended Data Fig. 3a, b). The SP test — a test of hedonic

behavior — revealed differences between mCherry susceptible (mCherry (S)) and resilient

(mCherry (R)) mice that were mirrored in DIO-hM4Di and hM3Dq mice, respectively. mCherry

(S) and hM4Di mice displayed a significant decrease in SP relative to mCherry (R) and hM3Dq-

injected mice (Fig. 3d). There were no significant differences in locomotion (Fig. 3e). The

chemogenetic manipulation also produced bidirectional effects on anxiety-like behavior in

elevated plus maze (EPM) and open-field tests (Extended Data Fig. 4). Surprisingly, mice

injected with hM3Dq-DREADDs went on to become resilient (Fig 3c) although activation of *Crf*

neurons of the BNST have previously been shown to produce depressive- and anxiogenic-like

responses[27,34,39,46,49,57]. Indeed in our hands, BNSTov[CRF] activation produced social avoidance

in mice with low (0-1 social defeat) stress exposure replicating previous studies (Extended Data

Fig. 5). Interestingly, resilience only occurred as a result of BNSTov[CRF] chemogenetic activation

between 7 and 10 episodes of social defeat; excitatory hM3Dq-DREADDs activation during

episodes 4-7 or 10-13 failed to display resiliency (Fig. 3g-i), suggesting stress history plays a

critical role in the behavioral outcomes of CRF modulation. Modulating *Crf* neurons with

inhibitory hM4Di- or excitatory hM3Dq- DREADDs between 8 and 10 SDEs led to enduring

susceptible or resilient phenotypes, respectively, up to 6 weeks after CNO manipulation

(Extended Data Fig. 6). These observations strongly support that the behavioral outcomes

induced by the activation of BNSTov[CRF] neurons are stress-history dependent.

Calcium dynamics underlying stress adaptation

We combined fiber photometry with excitatory or inhibitory DREADDs to determine whether

SDE-induced neuroadaptation in BNSTov[CRF] neurons, *in vivo*, reflects behavioral responses

consistent with resilient and susceptible phenotypes. Mice were co-injected with Cre-dependent

gCAMP7f and either DIO-hM4Di, -hM3Dq DREADDs or -mCherry viral vectors (Fig. 4a,b).

Clozapine-N-oxide (CNO) was administered via drinking water over the last 3 episodes of

RSDS, concomittant with the observed divergence of susceptible and resilient phenotypes.

In the control mice, we measured the calcium dynamics encoding social interaction with a novel

conspecific. In mice subjected to 7 SDEs, there was a decrease in neural activation upon

interaction in the mCherry (S/R) mice (Extended Data Fig. 7a). This pattern of activity occurred

in the presence and absence of a social target (no target/target trials) (Extended Data Fig. 7b).

Given that mice with shared stress-history of 7 SDEs appeared to have unimodal distribution on

a variety of measures, we hypothesized that calcium-encoded neural activity may diverge in

susceptible/resilient mice concomitant with the display of their respective behaviors on SI test.

Therefore, fiber photometric recordings were taken during SI Test 2 after mice after 10 SDEs.

Indeed, we observed mCherry (S) mice similarly had no change in the pattern of neural activity

between no target/ target trials, whereas mCherry (R) mice exhibited an increase in activity

upon social interaction (Extended Data Fig 7c). When comparing changes in calcium-encoded

neural dynamics between SI Test 1 and 2, mCherry (S) mice showed no difference, whereas

mCherry (R) mice developed an increase concomitant with the display of resilience (Fig. 4c-e).

These findings suggest that the persistence of neural activity across SDEs #8-10 is associated

with resiliency.

While we showed that we could bidirectionally drive the development of resilience/susceptibility,

this manipulation offers limited insight into the *in vivo* neural dynamics that reflect this

divergence point. To draw a link between neural changes and behavior *in vivo*, we multiplexed

chemogenetics and fiber photometry to bidirectionally drive the development of

resilient/susceptibility and mimic the neuroadaptive changes in BNSTov^{CRF} neurons that

produce behavioral divergence. During SI Test 1, similar to control mice, DREADDs injected

mice experienced a decreased in calcium-related neural activity upon initiating social interaction

with a novel conspecific (Extended Data Fig 7a), whose neural pattern was observed in both no

target and target trials (Extended Data Fig 7b). In hM4Di-injected mice, there were no

significant differences in neural activity during SI Test 1 and 2 trials, similar to control mCherry

(S) mice (Fig. 4f-h; Extended Data Fig. 7a-c). As was observed in the resilient (mCherry (R))

mice, the hM3Dq-injected group showed an increase in calcium-encoded neural activity (Fig 4f-

h). Mice injected with cre-dependent hM4Di showed a significant decrease in calcium-related

neural activity. In contrast, social interaction initiation led to increased activity in DIO-hM3Dq

injected mice (Fig. 4f-h). There were no significant differences in SI of mice subjected to 7

SDEs, yet differences emerge following 3 additional SDEs with a subset of mice becoming

susceptible and resilient (mCherry(R)/(S)) or activating/inhibiting DREADD promoting resiliency

and susceptibility in the direction predicted (Fig. 4i). We observed a strong correlation between

SI ratio and calcium activity only after 10 SDEs (Fig. 4j). There were no significant differences

observed in calcium-based neuronal activity upon social interaction with a novel conspecific in

mice subjected to 7 SDEs (Extended Data Fig. 7a,b), or in the absence of social interaction

(Extended Data Fig. 7d). In contrast, after 10 SDEs, mCherry (R) and DIO-hM3Dq injected

mice displayed significantly greater neuronal activation upon social contact relative to mCherry

(S) and DIO-hM4Di injected mice (Extended Data Fig. 7d). Despite the difference in time spent

interacting with a novel conspecific, the number of interaction zone entries were not significantly

different, nor differences in distance traveled (Fig. 4k, Extended Data Fig. 7e). These data

suggest that BNSTov^{CRF} neuronal dynamics are differentially altered by stress modulation in

accordance with phenotypical display resiliency/susceptibility.

***Crfr1* expression in BNSTov^{CRF} neurons mirrors the behavioral emergence of resilience.**

We observed a stress-induced enhancement in firing rates of BNSTov^{CRF} neurons in resilient

mice. To explore the effect of this BNSTov^{CRF} stress modulation on CRF receptor transmission,

we used RNAScope *in situ* hybridization to quantify *Crfr1* and *Crfr2* in accordance with stress

history. Mice were subjected to either 7- or 10-daily episodes of social defeat, and the BNST

and *Crfr1* and *Crfr2* genes were assessed (Fig. 5a-c) due to their reported role in mediating

stress responses[30,58–60]. The percentage of CRFR1-expressing neurons among CRF-

expressing neurons was higher in mice subjected to 7 SDEs than in susceptible mice but not

significantly different than resilient mice (Fig. 5d). In contrast, there were no significant

differences in BNSTov neurons co-expressing *Crfr2 and Crf mRNA* across groups of mice (Fig.

5e). The overlap between *Crfr1* and *Crf* in the BNST was significantly greater in the oval

nucleus than in anterolateral, anteromedial, and ventral subregions of the anterior dorsal BNST

(Fig. 5c,f).

To explore the role of firing rate changes on gene expression and the development of resiliency,

we optogenetically stimulated BNSTov^{CRF} neurons using transgenic *Crf-Cre*::ai32 mice, which

express channelrhodopsin-2 (ChR2) in *Crf*-containing neurons [61] (Fig. 5g). For control

experiments, transgenic mice of a similar background were used, except instead of ai32, mice

expressed the tdTomato fluorophore in CRF neurons (*Crf-tdTomato*). 5 Hz stimulation

frequency was used in order to mirror the average firing rate observed in the spontaneous firing

rate of resilient mice (Fig. 2d). Mice received 15-minute 5 Hz photo-stimulation of BNSTov^{CRF}

neurons following physical stress on SDEs #8-10. Mice were placed in the adjacent

compartment separating the aggressor, preventing physical contact. The compartment divides

the aggressor cage in half by a clear plexiglass that allows for continuous sensory cues (Fig.

5h,i). Photo-stimulation of *Crf*::ChR2 mice led to a significantly higher SI ratio and greater

percentage resilient than *Crf::tdTomato* mice (89% vs. 33%) when stimulated during SDEs #8-

10 (Fig. 5j). Surprisingly, *Crf::ChR2* mice that received photo-stimulation but were not subjected

to SDEs #8-10 showed a significant decrease in SI ratio and 100% of the mice, becoming

susceptible (0/6, SI $\geqq$ 1.0) (Fig. 5j). Photo-stimulation paired with SDEs #8-10 increased the

percentage of cells co-expressing *Crfr1* mRNA in CRF+ neurons relative to mice that

experienced photo-stimulation in the absence of additional stress (Fig. 5k, l). Additionally, we

obtained slice preparation from the optogenetically-induced resilient mice, and performed cell-

attached single unit recordings from BNSTov ChR2-expressing CRF neurons (Extended Data

Fig. 8a). Our recording data showed that optogenetic stimulation reliably induced 2.5 Hz, 5 Hz

and 10 Hz spike responses as expected (Extended Data Fig. 8b), and interestingly triggered

burst firing after the 5 Hz optical stimulation (Extended Data Fig. 8c,d). Importantly, Bath applied

CRFR1-selective antagonist NBI 27914 significantly decreased the firing rate of tested BNSTov

ChR2-expressing CRF neurons (Extended Data Fig. 8e,f). In summary, these data show that

maintenance of resiliency requires social defeat stress BNSTov^{CRF} activation and is correlated

with the upregulation of *Crfr1* expression in a stress-history dependent manner.

Discussion

The RSDS paradigm was modified to observe the effects of cumulative stress on neuroplasticity

in regions critical for mood regulation. Indeed, our work has uncovered a discrete window of

neuronal and behavioral plasticity between 7 and 10 SDEs during which susceptible and

resiliency phenotypes are established. Many of the studies investigating mechanisms of

susceptibility and resiliency have explored changes occurring after the 10 SDEs, when stable

susceptible/resilient phenotypes have already emerged. By capturing behavioral,

electrophysiological, and *in vivo* fiber photometric measures during the intra-social defeat stress

period, we uncover the mechanisms underlying the establishment of susceptible and resiliency

phenotypes, which may differ from those involved in its maintenance.

The BNST has been implicated in prosocial behaviors, and is a complex structure consisting of

16 different subnuclei serving separate functions[24,27,52,53]. While the oval nucleus has been

investigated in stress responses, we hypothesized that the region would be instrumental in

processing contexts associated with social stress. Indeed, we observed that individual

differences of stress effects on social behavior are encoded by BNSTov^{CRF} neurons. Prior

studies have implicated the overactivation of CRF neurons in the BNST as being pro-depressive

and anxiogenic[31,39,46,47,49], therefore, we were surprised to observe that BNSTov[CRF] neurons

exhibited increased activity in resilient mice. Moreover, cell-type-specific activation of this CRF

neuronal population yielded resiliency in mice. Importantly, we were able to replicate the effects

of BNSTov[CRF] activation, achieving a social avoidant response in line with previous studies, in

stress-naïve mice, suggesting that stress accumulation or stress history may play a critical role.

To our knowledge, this study is the first to target this neuronal population in a stress history-

dependent manner in the context of social behavior.   While CRF or CRF neuronal involvement

in stress responding have been reported [20,62–64], it is unknown about the contribution of this

neuronal population to resilience establishment in the context of social behavior when stress is

increasingly accumulated. Indeed, we observed that the activating Cre-dependent hM3Dq

manipulation only produced resilient mice after a certain "dose" of daily stressors (7 SDEs),

suggesting that the BNSTov may be tightly modulated based on stress-history resulting in

adaptive responses in contexts where they may be most advantageous. Our findings support a

view that resilience is a stress history-dependent state, which does not exist prior to stress.

This is highly consistent with previous demonstrations that resilience is a status achieved by

active regulation of genes and ion channel functions in this group more than in susceptible

animals[21,65–67].

CRF neurons have been observed to influence the salience of stressful contexts according to

stress exposure[24,27,60,64,68]. One feasible mechanism by which this occurs is through CRF

receptor dynamics. We hypothesized that the development of resiliency or susceptibility occurs

via CRF-CRFR interactions in the BNST. We were surprised to observe *Crfr1* mRNA

expression in CRF neurons corresponded to the establishment of resilient/susceptible

phenotypes. CRFR1 has been found largely on non-CRF neurons in the BNST[68,69], but our

finding suggests that stress promotes *Crfr1* mRNA expression on CRF neurons themselves in a

stress-history-dependent manner. This is evidenced by there being a higher overlap of *Crfr1/Crf*

mRNA expression in mice subjected to 7 SDEs than in severely stressed mice that underwent

10 SDEs and became susceptible.

CRFR1 has been shown to be selectively activated in settings of stress following a chronic

stressor, serving as a signaler of ongoing stress[70]. Several studies using viral constructs to

overexpress CRF in the dorsolateral BNST led to a decrease in CRFR1, presumably as a

compensatory response, yet the behavioral consequences of this were not explored in the

context of susceptibility and resiliency to stress[34,35]. CRFR1 is a G_s-coupled receptor that leads

to strong depolarization promoting cell activation[36,46,70–72]. Therefore, these neurons establish

resiliency by way of prolonged activation of CRF neurons, in part, by the maintenance of

CRFR1 expression.   This auto receptor-like ability has also been observed in terms of neuronal

activation. Chronic stress was shown to shift connectivity of local CRF$^+$ neurons from CRF$^+$-

CRF$^-$ to a larger percentage of CRF$^+$-CRF$^+$ cells[73]. Studies using prolonged overactivation (over

weeks to months) of CRF activity have yielded antidepressant and anxiolytic results[29,34,35]. Here

we identify a more discrete timeline in the span of days to capture the transition of when

BNSTov^{CRF} activation becomes pro-resilient. By optogenetically activating BNSTov^{CRF} neurons,

we observed an increase in CRFR1 expression. Though correlative, this exquisite regulation of

CRFR1 according to stress-history and at times demonstrating an opposing effect on

depressive-like behavior, may underlie why clinical trials of CRFR1 antagonists for MDD have

been met with variable success[59,74–76]. Although beyond the scope of the current study, future

experiments using siRNA or CRISPR knockdown approaches should be performed to determine

whether upregulation of CRFR1 is a critical facet of neuroadaption leading to resiliency.

Stress-sensitive regions such as the BNST have been found to be of critical importance in

stress coping and reactivity[77–79]. Stress resilience has long been considered a response

separate from or in the absence of stimuli that gives rise to stress susceptibility, mediated by

parallel circuits or cell-types in a particular brain region[14,15,80]. Here, we observe that activity

dynamics of CRF neurons can shape and influence resiliency to stress, potentially through

(auto)regulation of *Crfr1* mRNA. Other substrates may be used to classify these neurons either

on the basis of dual-neuropeptide or circuit-specific identities and should be the object of future

exploration.

Previous work has shown that resiliency is influenced by dopaminergic VTA neurons in the

nucleus accumbens (NAc)[29], in part, through the actions of brain-derived-neurotrophic-factor

(BDNF)[21,81,82]. CRF peptide has been important for BDNF release in the NAc as a stress-

coincidence sensor[20], yet the sources of CRF important for altering stress effect on social and

hedonic behavior has not been extensively characterized. While long-range GABAergic BNST

neurons projecting to the VTA have been shown to influence reward and anxiety-like

behavior[48,83–86], it is unclear to what degree these cells comprise the oval nuclear BNST

population. The BNST also sends projections to the dorsal raphe, lateral and paraventricular

hypothalamus, and ventrolateral periaqueductal gray[27,31–33,42,77,87–89], among others, whose

modulation have been linked to stress on affect and social motivation. In this way, the BNST

acts as a node for integrating information regarding stress history and determining socio-

affective outcomes according, possibly due to CRFR1 receptor dynamics occurring on CRF

neurons of the oval nucleus, thereby shaping the long-lasting outcome of resiliency.

Our study highlights a previously unknown mechanism by which the BNST encodes cumulative

social stress and effectuates susceptible or resilient outcomes. Importantly, there are currently

no Food and Drug Administration-approved drugs aimed at preventing a depressive episode

from occurring. By targeting mechanisms involved in establishing resiliency, the possibility may

exist to therapeutically leverage windows of plasticity to effectuate resilience and evade the

development of MDD.

360
Methods

362
Mice
The study used wild-type, Crf-ires-Cre (Jackson labs: 011087), ai14 (Cre-responsive tdTomato

reporter mouse; Jackson labs: 007915), ai32 (Cre-responsive channelrhodopsin-2/fused with

eYFP; Jackson labs: 012569) mice on C57BL/6J background that were bred at Icahn School of

Medicine at Mount Sinai and used were between 6-7 weeks at the start of experimental

manipulations. Upon receipt from Jackson Laboratories, mice were acclimated to the housing

facility for 1-2 weeks prior to the start of experiments. All mice were group-housed and

maintained on a 12/12 light-dark cycle with ad libitum access to food and water. Following

social defeat stress, mice were singly housed and maintained on a 12/12 light-dark cycle with

ad libitum access to food and water. All experiments were approved by the Institutional Animal

Care and Use Committee and comply with institutional guidelines for the Animal Care and Use

Committee set forth by Icahn School of Medicine at Mount Sinai.

Repeated social defeat stress paradigm

The repeated social defeat stress paradigm was performed according to published

protocols[6,21,23,82,90–92]. Briefly, CD1 aggressors were singly housed in cages (26.7 cm width x

48.3 cm depth x 15.2 cm height; Allentown Inc) at least 24 hours before the start of the

experiment on one side with a clear perforated Plexiglas divider. Social stress consists of

physical and sensory stress components. During the period which marks the physical stress,

C57BL/6J mice are placed on the ipsilateral side of the cage as the CD1 aggressor for 10

minutes. Following this, the intruder mouse is placed in the contralateral side of the perforated

Plexiglas divider for the remainder of the 24-hour period, marking the sensory-stress period.

Every 24 hours, the intruder mouse is paired with a new aggressor for 10 episodes. Control

mice were housed two mice per cage divided by a perforated Plexiglas divider and rotated and

handled daily like the socially defeated mice.

Social interaction test

Social interaction testing was performed as described[6,20,21,65,66,81,82,93]. Briefly, a novel

conspecific of CD1 strain is placed in an interaction zone of a standard open-field arena, and

the time the intruder spends in the interaction zone is measured. The mice spend a total of 5

minutes in the open arena (2.5 min with and without the novel CD1). Ethovision (Noldus

Information Technology) video-tracking software is used to track interaction time. All social

interaction testing takes place 24 hours after the last defeat. Social interaction (SI) is measured

by time spent in the interaction zone during first (CD1 absent) over second (CD1 present) trials.

Mice are categorized according to SI ratios; an SI ratio $\geqq 1$ defines resilient, whereas an SI ratio

<1 is susceptible as described previously[5,17,19,21].

Sucrose preference test

For sucrose preference testing, a solution of 1% sucrose or diluent alone (drinking water) is filled

in 50 ml tubes with ball-pointed sipper nozzles (Ancare). Animals are acclimatized to two-bottle

choice conditions prior to testing. The bottles are weighed, and positions interchanged daily.

Sucrose preference is calculated as a percentage [100 x volume of sucrose consumed (in bottle

A)/total volume consumed (bottles A and B)] and averaged over 2 days of testing.

Cell-attached electrophysiology

Acute coronal brain slices of the BNSTov were prepared according to previously published

protocols[65,66,90,94]. All recordings were carried out blind to stress history, behavioral phenotype,

or drug treatment. Male 8-12 weeks old mice were perfused with cold artificial cerebrospinal

fluid (aCSF) containing (in mM): 128 NaCl, 3 KCl, 1.25 NaH_2PO_4, 10 D-glucose, 24 $NaHCO_3$, 2

$CaCl_2$, and 2 $MgCl_2$ (oxygenated with 95% O_2 and 5% CO_2, pH 7.35, 295-305 mOsm). Acute

brain slices containing BNSTov were cut using a microslicer (DTK-1000, Ted Bella) in sucrose-

ACSF, derived by replacing NaCl with 254 mM sucrose, and saturated by 95% O_2 and 5% CO_2

(2.5 ml/min) and 35ºC. Glass recording pipettes (2-4 MΩ) were filled with an internal solution

containing (mM): 115 potassium gluconate, 20 KCl, 1.5 $MgCl_2$, 10 phosphocreatine, 10 HEPES,

2 magnesium ATP and 0.5 GTP (pH 7.2, 285 mOSm). BNSTov CRF neurons were identified by

location, and infrared differential interference contrast microscopy and recordings were made

from CRF-positive neurons as indicated by the presence of tdTomato (in *Crf*::tdTomato mice) or

eYFP in the case of (*Crf*::ChR2) mice. Spontaneous firing rates were recorded in cell-attach

mode, and data acquisition and online analysis of firing rates were collected using a Digidata

1440 digitizer and pClamp 10.2 (Axon Instruments). The timing of recordings was made

consistent throughout treatment groups. A burst was defined as containing at least three

spikes, with interspike intervals <15 ms[95]. Neuronal firing rates were considered bursting or

non-bursting if they had undergone a statistically significant change with a P<0.05 on the rank-

sum test.

427
428

Qualitative defeat assessment

Social defeat encounters were recorded, and BORIS (Behavioral Observation Research

Interactive Software) was used to code behaviors. Five behaviors (cage exploration, aggressor

grooming, flight, motionless, rearing/defensive posturing) were scored per published reports on

typical social defeat behaviors[9,18,37,96]. The total time engaging in each behavior was tallied to a

blinded observer, and scores were aggregated.

Viral constructs

For DREADD experiments in CRF neuronal populations, CRF-ires-Cre animals were injected

with AAV5-hSyn-DIO-hM4D(Gi)-mCherry (Addgene: 44362-AAV5) (≥7x10^12), AAV5-hSyn-

DIO-hM3D(Gq)-mCherry (Addgene: 44361-AAV5) (≥7x10^12), and AAV5-hSyn-DIO-mCherry

(Addgene: 50459). For fiber photometric recordings, CRF-ires-Cre animals were injected with

AAV9-syn-FLEX-jGCaMP7f-WPRE (addgene: 104492-AAV9) (>1x10^12). All viruses were

purchased from Addgene.

Stereotaxic virus and optic fiber implantation

Under ketamine (80 mg/kg)/xylazine (10 mg/kg) anesthesia, mice were placed in a stereotaxic

frame (Kopf Instruments) and the BNSTov was targeted (coordinates: anterior/posterior +0.20,

media/lateral: +/- 2.15, dorsal/ventral, –4.0 mm; 15-degree angle). Hair was shaved around the

crown of the head, alcohol, and betadine were applied to the scalp. Ophthalmic ointment was

applied to the eyes to prevent dryness, a midline incision was made down the scalp, and a

craniotomy was made using a dental drill. A 10 ul Nanofil Hamilton syringe (WPI, Sarasota, Fl)

with a 34-gauge beveled metal needle was used to infuse 0.5 ul virus at a rate of 85 nl/minute.

Following infusion, the needle was kept at the injection site for 10 minutes and slowly

withdrawn. For the optogenetic experiments, chronically implantable optic fibers constructed

with 1.25 mm diameter, 200 um core, 0.39 numeral aperture (NA) were used (RWD).   For the

fiber photometry experiments, chronically implantable optic fibers constructed with 1.25 mm

diameter, 400 um core 0.48 numerical aperture (NA) optic fiber and unilaterally implanted into

the BNSTov (coordinates: anterior/posterior +0.20, media/lateral: +/- 2.15, dorsal/ventral, –3.9

mm; 15º angle) (thor labs). Fiber optical ferrules were cemented to the skull using dental acrylic

(Parkell C&B Metabond). All optical stimulation and fiber photometric recording experiments

were conducted a minimum of 3-4 weeks post-implantation.

CNO-drinking water construct

Clozapine-N-Oxide (CNO) was obtained from (Hello Bio, catalog no HB6149). The dry chemical

was dissolved in drinking water obtained from the vivarium and diluted such that each mouse

received 5 mg/kg/day based on previous studies. CNO was made fresh each day for the three

days it was administered. CNO solutions were protected from light throughout the experimental

procedure. On average, mice consumed ~4-5 mL of water per day. Water bottles and mice

were weighed daily. CNO water bottles were exchanged for normal drinking water after the

social interaction test 1 and 8-12 hours before social defeat stress to ensure neuronal

modulation during stress exposure.

Elevated plus-maze

Mice began testing by being placed in the center of the maze (Model ENV-560A, Med

Associated, Fairfax, VT, USA), where movement was tracked using Ethovision behavioral

tracking software (Noldus). The test lasted 5 minutes.

Open field test

Mice were placed in an arena (42 cm (w) x 42 cm (d) x 42 cm (h); Nationwide Plastics, custom

order). Mice were tracked using behavioral tracking software (Ethovision, Noldus), and time

spent in the designated center and surround zone, as well as locomotor activity, were

measured.

Optogenetic manipulation of BNSTov^{CRF} neurons

Optical fibers were implanted on Ai32 (transgenic mouse line expressing light-gated cation

channel channelrhodopsin-2 (ChR2). Optical fiber   Optogenetic stimulation was conducted via

the use of a diode-pumped solid-state (DPSS) 473-nm blue laser (Crystal Laser, BCL-473-050-

M), using a patch cord with an FC/PC adaptor (Doric Lenses, MFP_200/240/900-0.22_4m_FC-

MF2.5). A functional generator (Agilent Technologies; 33220A) was used to generate a 5 Hz

frequency, pulse width of 10 ms for 15 minutes[97,98], and power density between 7-9 mW mm$^{-2.}$

Experimenters were blinded to the stimulation group.

Fiber photometry calcium imaging

Optical recordings of GCaMP7f fluorescence were acquired using two LEDs at 490 and 405

(Thor Labs), reflected off dichroic mirrors (Semrock, FF495) and coupled into a 400 micro 0.48

NA optical fiber (Thorlabs BFH48-600) using a 40 x 0.48 NA microscope objective and fiber

launch with the pat chord linked to an implanted 400 um optical fiber with zirconia sheath.

Signals in both 470 and 405 nm channels are monitored throughout the recordings, whereas the

405 nm is used as an isosbestic control for ambient fluorescents and motion artifacts caused by

movement or torque about the fiber optic implant. Wavelengths were modulated at frequencies

of 210-220 and 330 Hz, respectively, with power output maintained at 20 mA and a DC offset of

3 mA for both light sources. All signalers were acquired at 1 kHz and lowpass filtered at 3 Hz.

Prior to social interaction testing, mice were handled for and connected via a patch cable and

placed in the home cage for 5-7 minutes for basal BNSTov recording and habituation prior to

the start of the no-target trial.

Fiber photometry analysis

A custom-written MATLAB code (modified from previously published work)[99] was used to

analyze the GCaMP7f signal. Firstly, the bulk fluorescent signal from both the 470 nm and 405

nm channels were normalized. A linear regression (slope of the 405 nm fitted against the 470

nm signal) was applied over the data to correct for bleaching of signal of the duration at each

recording. The Initial 3 seconds for the signals were discarded because of the

photoreceiver/LED rise time artifact. Detection of GCaMP7f signal is calculated as a change in

the 470 nm/fitted 405 nm signal over the fitted 405 signal (ΔF/F). For the calculation of intensity

values around events of interest (PC2), a 6-second window was used. We used 3 seconds

before and after the PC2 event onset and computed z-scores for all such event windows.

These z-scored intensity values were then averaged within and between animals in particular

groups, and 95% confidence intervals for the averaged intensities were computed.

RNAScope *in-situ* hybridization

To prepare frozen sections, mice were placed in an air-tight chamber with 2.5 ml of isoflurane-

infused cotton balls. After 45-60 seconds (after respiratory depression and loss of

consciousness as evidenced by the negative tail and paw pinch), brains were acutely harvested

and placed in a -80°C storage chamber until RNAScope *in-situ* hybridization protocol

commenced. 16-µm coronal sections were collected on a cryostat (Leica Biosystems) at -20°C

and mounted directly onto ColorFrost Plus microscope slides (Fischer Scientific). Slides were

stored at -80°C until *in situ* hybridization (ISH) processing. ISH was performed using the

RNAscope multiplex fluorescent kit (Advanced Cell Diagnostics, ACD) according to the

manufacturer's instructions. The tissue was fixed in 4% paraformaldehyde (PFA) in PBS chilled

to 4°C for 15 minutes, washed twice briefly in 1x PBS, then dehydrated in 50% ethanol, 70%

ethanol, and twice in 100% ethanol for 5 minutes each at room temperature (RT). A

hydrophobic barrier was traced around sections of interest using an ImmEdge pen (Fischer

Scientific), and once the barrier dried, sections were incubated in Protease IV reagent for 30

minutes at RT, then rinsed twice in 1x PBS for 5 minutes. Probes for CRH and either CRHR1 or

CRHR2 (ACD) were warmed, mixed, and placed onto sections for hybridization at 40°C for 2

hours in a HybEZ II oven (ACD), which was used for all subsequent incubations. Sections were

washed twice in 1x wash buffer (ACD) for 2 minutes each, then incubated with a series of

amplification reagents at 40°C, Amp 1-FL for 30 minutes, Amp 2-FL for 15 minutes, Amp 3-FL

for 30 minutes, and Amp 4-FL Alt A or C for 15 minutes, washing twice in 1x wash buffer for 2

minutes each between steps. DAPI was applied to the sections for 30 s, then immediately

coverslipped with ProLong Gold mounting medium (Invitrogen).

Images were collected on a Zeiss LSM 780 confocal microscope at 20x, 40x, or 63x

magnification. mRNA puncta were quantified manually using FIJI 1.0 software (Image J), all

experimental conditions were blinded to the imager, and all cell analysis and quantification.

Immunohistochemistry

Mice were quickly anesthetized with urethane and perfused with cold 1x-PBS (OmniPure, Bio-

Rad) followed by cold 4% PFA (Fisher Hamilton Scientific). Brains were placed in PFA at 4ºC

for at least 24 hours, followed by being replaced with a 15% sucrose/PBS solution at 4ºC for 24

hours and brain sinking. The next day, brains were placed in a 30% sucrose/PBS solution at

4ºC for 24 hours. Once the brains sunk, it was washed in 1xPBS three times and mounted on a

freezing microtome for sectioning at 35 micrometers.

Sections were washed three times for 10 minutes in 1x PBS, then blocked in 5% bovine serum

albumin (Sigma-Aldrich) in 1x PBS with 0.3% Triton-X (Sigma-Aldrich) for 1 hour. Tissues were

incubated with primary antibodies for rabbit anti-mCherry ((1:1000), Invitrogen), rabbit anti-c-Fos

antibody (1:250, Milipore C-10, sc-271243) and goat anti-GFP (1:500, abcams) overnight at

4ºC. Sections were washed three times with 1xPBS and blocked with 0.3% Triton-X for one

hour. Tissue was blocked with secondary antibodies goat anti-rabbit/568 (1:1000, Abcams) and

donkey anti-goat Alexa-488 (1:500, Abcams) at room temperature for 1 hour. Sections were

washed three times with 1x PBS for 10 minutes and mounted on slides with ProLong Gold

antifade reagent with DAPI (invitrogen, P36931). Z-stacked images were acquired with a Zeiss

LSM780 multi photon confocal system and z-stacks were collated using ImageJ (Fiji). Cell

counters were obtained using the Cell Counter feature on ImageJ.

Statistical analysis

Animals were randomly assigned to control and experimental groups and all experimenters were blinded. Mice were excluded if viral infection was off-target. No data was excluded for other reasons. Student's two-tailed t-tests were used for comparisons of two experimental groups. For parametric data sets, comparisons among three or more groups were performed using one or two-way ANOVA tests followed by Tukey's or Bonferroni posthoc tests. For all tests, $p<0.05$ was determined to be significant. Statistical analyses were performed using Graph Pad Prism 9 .3.1 software (La Jolla, CA, USA). For data not normally distributed, non-parametric analyses were performed.

Data & Code availability

All source data for this manuscript and MATLAB code used to analyze photometry data is available upon request.

Acknowledgements

We would like to thank S.M. for generous sharing for input on the manuscript. We would like to thank S.R. and L.L. for lending viral constructs for behavioral validation. We would like to thank the animal care staff at ISMMS for the animal care staff and technical support. This research was funded by the National Institutes of Health grants R01MH072908 to DGR and LJY, R01MH1206387 to MHH and SJR, R21MH112081 to MHH and SJR, F31MH114624 to SEH, and P51 OD011132 to YNPRC, by the National Key R&D Program of 1034 China No. 2021ZD0202900 to MHH.

References

1.   Veer, I. M. *et al.* Psycho-social factors associated with mental resilience in the Corona

lockdown. *Transl. Psychiatry* **11**, 67 (2021).

2.   Post, R. M. Transduction of psychosocial stress into the neurobiology of recurrent

affective disorder. *Am. J. Psychiatry* **149**, 999–1010 (1992).

3.   Gold, P. W. Evidence that stress per se has a role in the precipitation and natural history

of depressive illness. *Mol. Psychiatry* **18**, 954–956 (2013).

4.   Kendler, K. S., Karkowski, L. M. & Prescott, C. A. Causal relationship between stressful life

events and the onset of major depression. *Am. J. Psychiatry* **156**, 837–841 (1999).

5.   Barthas, F. *et al.* Cumulative Effects of Social Stress on Reward-Guided Actions and

Prefrontal Cortical Activity. *Biol. Psychiatry* **88**, 541–553 (2020).

6.   Golden, S. A., Covington, H. E., Berton, O. & Russo, S. J. A standardized protocol for

repeated social defeat stress in mice. *Nat. Protoc.* **6**, 1183–1191 (2011).

7.   Martinez, M., Phillips, P. J. & Herbert, J. Adaptation in patterns of *c-fos* expression in the

brain associated with exposure to either single or repeated social stress in male rats.

European Journal of Neuroscience **10**, 20–33 (1998).

8.   Zheng, Z. *et al.* Hypothalamus-habenula potentiation encodes chronic stress experience

and drives depression onset. *Neuron* (2022) doi:10.1016/j.neuron.2022.01.011.

9.   Kudryavtseva, N. N., Bakshtanovskaya, I. V. & Koryakina, L. A. Social model of depression

608   in mice of C57BL/6J strain. *Pharmacol. Biochem. Behav.* **38**, 315–320 (1991).

609   10.   Laine, M. A. *et al.* Brain activation induced by chronic psychosocial stress in mice. *Sci.*

610   *Rep.* **7**, 15061 (2017).

611   11.   Hultman, R. *et al.* Brain-wide Electrical Spatiotemporal Dynamics Encode Depression

612   Vulnerability. *Cell* **173**, 166-180.e14 (2018).

613   12.   Drysdale, A. T. *et al.* Resting-state connectivity biomarkers define neurophysiological

614   subtypes of depression. *Nat. Med.* **23**, 28–38 (2017).

615   13.   Samuels, B. A. *et al.* Modeling treatment-resistant depression. *Neuropharmacology* **61**,

616   408–413 (2011).

617   14.   Franklin, T. B., Saab, B. J. & Mansuy, I. M. Neural mechanisms of stress resilience and

618   vulnerability. *Neuron* **75**, 747–761 (2012).

619   15.   Russo, S. J., Murrough, J. W., Han, M.-H., Charney, D. S. & Nestler, E. J. Neurobiology of

620   resilience. *Nat. Neurosci.* **15**, 1475–1484 (2012).

621   16.   Karatsoreos, I. N. & McEwen, B. S. Psychobiological allostasis: resistance, resilience and

622   vulnerability. *Trends Cogn Sci (Regul Ed)* **15**, 576–584 (2011).

623   17.   Laine, M. A. *et al.* Genetic Control of Myelin Plasticity after Chronic Psychosocial Stress.

624   *eNeuro* **5**, (2018).

18. Markham, C. M., Norvelle, A. & Huhman, K. L. Role of the bed nucleus of the stria

terminalis in the acquisition and expression of conditioned defeat in Syrian hamsters.

Behav. Brain Res. **198**, 69–73 (2009).

19. Gururajan, A., Bastiaanssen, T. F. S., Ventura Silva, A. P., Moloney, G. M. & Cryan, J. F.

The impact of psychosocial defeat stress on the BNST transcriptome in adult male mice.

Eur. J. Neurosci. (2021) doi:10.1111/ejn.15567.

20. Walsh, J. J. *et al.* Stress and CRF gate neural activation of BDNF in the mesolimbic reward

pathway. *Nat. Neurosci.* **17**, 27–29 (2014).

21. Krishnan, V. *et al.* Molecular adaptations underlying susceptibility and resistance to

social defeat in brain reward regions. *Cell* **131**, 391–404 (2007).

22. Matsuda, S. *et al.* Persistent c-fos expression in the brains of mice with chronic social

stress. *Neurosci. Res.* **26**, 157–170 (1996).

23. Berton, O. *et al.* Essential role of BDNF in the mesolimbic dopamine pathway in social

defeat stress. *Science* **311**, 864–868 (2006).

24. Waraczynski, M. Toward a systems-oriented approach to the role of the extended

amygdala in adaptive responding. *Neurosci. Biobehav. Rev.* **68**, 177–194 (2016).

25. Flanigan, M. E. & Kash, T. L. Coordination of social behaviors by the bed nucleus of the

stria terminalis. *Eur. J. Neurosci.* (2020) doi:10.1111/ejn.14991.

26.   Lebow, M. A. & Chen, A. Overshadowed by the amygdala: the bed nucleus of the stria

terminalis emerges as key to psychiatric disorders. *Mol. Psychiatry* **21**, 450–463 (2016).

27.   Daniel, S. E. & Rainnie, D. G. Stress modulation of opposing circuits in the bed nucleus of

the stria terminalis. *Neuropsychopharmacology* **41**, 103–125 (2016).

28.   Jennings, J. H. *et al.* Distinct extended amygdala circuits for divergent motivational

states. *Nature* **496**, 224–228 (2013).

29.   Dedic, N. *et al.* Chronic CRH depletion from GABAergic, long-range projection neurons in

the extended amygdala reduces dopamine release and increases anxiety. *Nat. Neurosci.*

21, 803–807 (2018).

30.   Chen, Y., Molet, J., Gunn, B. G., Ressler, K. & Baram, T. Z. Diversity of Reporter

Expression Patterns in Transgenic Mouse Lines Targeting Corticotropin-Releasing

Hormone-Expressing Neurons. *Endocrinology* **156**, 4769–4780 (2015).

31.   Dabrowska, J., Martinon, D., Moaddab, M. & Rainnie, D. G. Targeting Corticotropin-

Releasing Factor Projections from the Oval Nucleus of the Bed Nucleus of the Stria

Terminalis Using Cell-Type Specific Neuronal Tracing Studies in Mouse and Rat Brain. *J.*

Neuroendocrinol. **28**, (2016).

32.   Matthews, G. A. *et al.* Dorsal raphe dopamine neurons represent the experience of

social isolation. *Cell* **164**, 617–631 (2016).

33.   Marcinkiewcz, C. A. *et al.* Serotonin engages an anxiety and fear-promoting circuit in the

extended amygdala. *Nature* **537**, 97–101 (2016).

34.   Regev, L. *et al.* Prolonged and site-specific over-expression of corticotropin-releasing

factor reveals differential roles for extended amygdala nuclei in emotional regulation.

Mol. Psychiatry **16**, 714–728 (2011).

35.   Sink, K. S. *et al.* Effects of continuously enhanced corticotropin releasing factor

expression within the bed nucleus of the stria terminalis on conditioned and

unconditioned anxiety. *Mol. Psychiatry* **18**, 308–319 (2013).

36.   Konishi, S., Kasagi, Y., Katsumata, H., Minami, S. & Imaki, T. Regulation of corticotropin-

releasing factor (CRF) type-1 receptor gene expression by CRF in the hypothalamus.

Endocr. J. **50**, 21–36 (2003).

37.   Jasnow, A. M., Banks, M. C., Owens, E. C. & Huhman, K. L. Differential effects of two

corticotropin-releasing factor antagonists on conditioned defeat in male Syrian hamsters

(Mesocricetus auratus). *Brain Res.* **846**, 122–128 (1999).

38.   Avery, S. N., Clauss, J. A. & Blackford, J. U. The human BNST: functional role in anxiety

and addiction. *Neuropsychopharmacology* **41**, 126–141 (2016).

39.   Kim, S.-Y. *et al.* Diverging neural pathways assemble a behavioural state from separable

features in anxiety. *Nature* **496**, 219–223 (2013).

40.   Rodríguez-Sierra, O. E., Goswami, S., Turesson, H. K. & Pare, D. Altered responsiveness of

680   BNST and amygdala neurons in trauma-induced anxiety. *Transl. Psychiatry* **6**, e857

681   (2016).

682   41.   Ventura-Silva, A. P. *et al.* Stress shifts the response of the bed nucleus of the stria

683   terminalis to an anxiogenic mode. *Eur. J. Neurosci.* **36**, 3396–3406 (2012).

684   42.   Giardino, W. J. *et al.* Parallel circuits from the bed nuclei of stria terminalis to the lateral

685   hypothalamus drive opposing emotional states. *Nat. Neurosci.* **21**, 1084–1095 (2018).

686   43.   Duvarci, S., Bauer, E. P. & Paré, D. The bed nucleus of the stria terminalis mediates inter-

687   individual variations in anxiety and fear. *J. Neurosci.* **29**, 10357–10361 (2009).

688   44.   Duque-Wilckens, N. *et al.* Extrahypothalamic oxytocin neurons drive stress-induced

689   social vigilance and avoidance. *Proc Natl Acad Sci USA* **117**, 26406–26413 (2020).

690   45.   Dabrowska, J. *et al.* Striatal-enriched protein tyrosine phosphatase-STEPs toward

691   understanding chronic stress-induced activation of corticotrophin releasing factor

692   neurons in the rat bed nucleus of the stria terminalis. *Biol. Psychiatry* **74**, 817–826

693   (2013).

694   46.   Hu, P. *et al.* Chronic Stress Induces Maladaptive Behaviors by Activating Corticotropin-

695   Releasing Hormone Signaling in the Mouse Oval Bed Nucleus of the Stria Terminalis. *J.*

696   *Neurosci.* **40**, 2519–2537 (2020).

47. Daniel, S. E. *et al.* Chronic stress induces cell type-selective transcriptomic and

electrophysiological changes in the bed nucleus of the stria terminalis.

Neuropharmacology **150**, 80–90 (2019).

48. Wanat, M. J., Bonci, A. & Phillips, P. E. M. CRF acts in the midbrain to attenuate

accumbens dopamine release to rewards but not their predictors. *Nat. Neurosci.* **16**,

383–385 (2013).

49. Salimando, G. J., Hyun, M., Boyt, K. M. & Winder, D. G. BNST GluN2D-Containing NMDA

Receptors Influence Anxiety- and Depressive-like Behaviors and ModulateCell-Specific

Excitatory/Inhibitory Synaptic Balance. *J. Neurosci.* **40**, 3949–3968 (2020).

50. Hartley, N. D. *et al.* Dynamic remodeling of a basolateral-to-central amygdala

glutamatergic circuit across fear states. *Nat. Neurosci.* **22**, 2000–2012 (2019).

51. Sanford, C. A. *et al.* A Central Amygdala CRF Circuit Facilitates Learning about Weak

Threats. *Neuron* **93**, 164–178 (2017).

52. Rodríguez-Sierra, O. E., Turesson, H. K. & Pare, D. Contrasting distribution of

physiological cell types in different regions of the bed nucleus of the stria terminalis. *J.*

Neurophysiol. **110**, 2037–2049 (2013).

53. Gungor, N. Z. & Paré, D. Functional heterogeneity in the bed nucleus of the stria

terminalis. *J. Neurosci.* **36**, 8038–8049 (2016).

54.   Zhan, J., Komal, R., Keenan, W. T., Hattar, S. & Fernandez, D. C. Non-invasive Strategies

for Chronic Manipulation of DREADD-controlled Neuronal Activity. *J. Vis. Exp.* (2019)

doi:10.3791/59439.

55.   Schalbetter, S. M. *et al.* Oral application of clozapine-N-oxide using the micropipette-

guided drug administration (MDA) method in mouse DREADD systems. *Lab Anim (NY)*

50, 69–75 (2021).

56.   Wess, J., Nakajima, K. & Jain, S. Novel designer receptors to probe GPCR signaling and

physiology. *Trends Pharmacol. Sci.* **34**, 385–392 (2013).

57.   Walker, D. L., Miles, L. A. & Davis, M. Selective participation of the bed nucleus of the

stria terminalis and CRF in sustained anxiety-like versus phasic fear-like responses. *Prog*

Neuropsychopharmacol Biol Psychiatry **33**, 1291–1308 (2009).

58.   Tran, L., Schulkin, J. & Greenwood-Van Meerveld, B. Importance of CRF receptor-

mediated mechanisms of the bed nucleus of the stria terminalis in the processing of

anxiety and pain. *Neuropsychopharmacology* **39**, 2633–2645 (2014).

59.   Lee, Y., Fitz, S., Johnson, P. L. & Shekhar, A. Repeated stimulation of CRF receptors in the

BNST of rats selectively induces social but not panic-like anxiety.

Neuropsychopharmacology **33**, 2586–2594 (2008).

60.   Dedic, N. *et al.* Deletion of CRH From GABAergic Forebrain Neurons Promotes Stress

Resilience and Dampens Stress-Induced Changes in Neuronal Activity. *Front. Neurosci.*

13, 986 (2019).

61.   Wang, Y. *et al.* Single-cell morphological characterization of CRH neurons throughout the

whole mouse brain. *BMC Biol.* **19**, 47 (2021).

62.   Chen, P. *et al.* Prefrontal Cortex Corticotropin-Releasing Factor Neurons Control

Behavioral Style Selection under Challenging Situations. *Neuron* **106**, 301-315.e7 (2020).

63.   Elliott, E., Ezra-Nevo, G., Regev, L., Neufeld-Cohen, A. & Chen, A. Resilience to social

stress coincides with functional DNA methylation of the Crf gene in adult mice. *Nat.*

Neurosci. **13**, 1351–1353 (2010).

64.   Lemos, J. C. *et al.* Severe stress switches CRF action in the nucleus accumbens from

appetitive to aversive. *Nature* **490**, 402–406 (2012).

65.   Zhang, H. *et al.* α1- and β3-Adrenergic Receptor-Mediated Mesolimbic Homeostatic

Plasticity Confers Resilience to Social Stress in Susceptible Mice. *Biol. Psychiatry* **85**, 226–

236 (2019).

66.   Friedman, A. K. *et al.* Enhancing depression mechanisms in midbrain dopamine neurons

achieves homeostatic resilience. *Science* **344**, 313–319 (2014).

67.   Bagot, R. C. *et al.* Circuit-wide Transcriptional Profiling Reveals Brain Region-Specific

Gene Networks Regulating Depression Susceptibility. *Neuron* **90**, 969–983 (2016).

68.   Dabrowska, J., Hazra, R., Guo, J.-D., Dewitt, S. & Rainnie, D. G. Central CRF neurons are

752   not created equal: phenotypic differences in CRF-containing neurons of the rat

753   paraventricular hypothalamus and the bed nucleus of the stria terminalis. *Front.*

754   *Neurosci.* **7**, 156 (2013).

755   69.   Justice, N. J., Yuan, Z. F., Sawchenko, P. E. & Vale, W. Type 1 corticotropin-releasing

756   factor receptor expression reported in BAC transgenic mice: implications for reconciling

757   ligand-receptor mismatch in the central corticotropin-releasing factor system. *J. Comp.*

758   *Neurol.* **511**, 479–496 (2008).

759   70.   Ramot, A. *et al.* Hypothalamic CRFR1 is essential for HPA axis regulation following

760   chronic stress. *Nat. Neurosci.* **20**, 385–388 (2017).

761   71.   McClard, C. K. *et al.* POU6f1 Mediates Neuropeptide-Dependent Plasticity in the Adult

762   Brain. *J. Neurosci.* **38**, 1443–1461 (2018).

763   72.   Hillhouse, E. W. & Grammatopoulos, D. K. The molecular mechanisms underlying the

764   regulation of the biological activity of corticotropin-releasing hormone receptors:

765   implications for physiology and pathophysiology. *Endocr. Rev.* **27**, 260–286 (2006).

766   73.   Partridge, J. G. *et al.* Stress increases GABAergic neurotransmission in CRF neurons of

767   the central amygdala and bed nucleus stria terminalis. *Neuropharmacology* **107**, 239–

768   250 (2016).

74.   Spierling, S. R. & Zorrilla, E. P. Don't stress about CRF: assessing the translational failures

of CRF1antagonists. *Psychopharmacology (Berl)* **234**, 1467–1481 (2017).

75.   Waters, R. P. *et al.* Evidence for the role of corticotropin-releasing factor in major

depressive disorder. *Neurosci. Biobehav. Rev.* **58**, 63–78 (2015).

76.   Reul, J. Corticotropin-releasing factor receptors 1 and 2 in anxiety and depression. *Curr.*

Opin. Pharmacol. **2**, 23–33 (2002).

77.   Johnson, S. B. *et al.* A basal forebrain site coordinates the modulation of endocrine and

behavioral stress responses via divergent neural pathways. *J. Neurosci.* **36**, 8687–8699

(2016).

78.   Crestani, C. C., Alves, F. H. F., Correa, F. M. A., Guimarães, F. S. & Joca, S. R. L. Acute

reversible inactivation of the bed nucleus of stria terminalis induces antidepressant-like

effect in the rat forced swimming test. *Behav. Brain Funct.* **6**, 30 (2010).

79.   Fiedler, D. *et al.* Brain-Derived Neurotrophic Factor/Tropomyosin Receptor Kinase B

Signaling Controls Excitability and Long-Term Depression in Oval Nucleus of the BNST. *J.*

Neurosci. **41**, 435–445 (2021).

80.   Cathomas, F., Murrough, J. W., Nestler, E. J., Han, M.-H. & Russo, S. J. Neurobiology of

resilience: interface between mind and body. *Biol. Psychiatry* **86**, 410–420 (2019).

81.   Koo, J. W., Chaudhury, D., Han, M.-H. & Nestler, E. J. Role of Mesolimbic Brain-Derived

Neurotrophic Factor in Depression. *Biol. Psychiatry* **86**, 738–748 (2019).

82.   Wook Koo, J. *et al.* Essential Role of Mesolimbic Brain-Derived Neurotrophic Factor in

Chronic Social Stress-Induced Depressive Behaviors. *Biol. Psychiatry* **80**, 469–478 (2016).

83.   Takahashi, D. *et al.* Tonic Suppression of the Mesolimbic Dopaminergic System by

Enhanced Corticotropin-Releasing Factor Signaling Within the Bed Nucleus of the Stria

Terminalis in Chronic Pain Model Rats. *J. Neurosci.* **39**, 8376–8385 (2019).

84.   Caillé, S., Guillem, K., Cador, M., Manzoni, O. & Georges, F. Voluntary nicotine

consumption triggers in vivo potentiation of cortical excitatory drives to midbrain

dopaminergic neurons. *J. Neurosci.* **29**, 10410–10415 (2009).

85.   Silberman, Y., Matthews, R. T. & Winder, D. G. A corticotropin releasing factor pathway

for ethanol regulation of the ventral tegmental area in the bed nucleus of the stria

terminalis. *J. Neurosci.* **33**, 950–960 (2013).

86.   Rodaros, D., Caruana, D. A., Amir, S. & Stewart, J. Corticotropin-releasing factor

projections from limbic forebrain and paraventricular nucleus of the hypothalamus to

the region of the ventral tegmental area. *Neuroscience* **150**, 8–13 (2007).

87.   Dong, H. W., Petrovich, G. D., Watts, A. G. & Swanson, L. W. Basic organization of

projections from the oval and fusiform nuclei of the bed nuclei of the stria terminalis in

adult rat brain. *J. Comp. Neurol.* **436**, 430–455 (2001).

88.   Maita, I., Bazer, A., Blackford, J. U. & Samuels, B. A. Functional anatomy of the bed

nucleus of the stria terminalis-hypothalamus neural circuitry: Implications for valence

surveillance, addiction, feeding, and social behaviors. *Handb. Clin. Neurol.* **179**, 403–418

(2021).

89.   Kaouane, N., Ada, S., Hausleitner, M. & Haubensak, W. Dorsal Bed Nucleus of the Stria

Terminalis-Subcortical Output Circuits Encode Positive Bias in Pavlovian Fear and

Reward. *Front. Neural Circuits* **15**, 772512 (2021).

90.   Cao, J.-L. *et al.* Mesolimbic dopamine neurons in the brain reward circuit mediate

susceptibility to social defeat and antidepressant action. *J. Neurosci.* **30**, 16453–16458

(2010).

91.   Morel, C. *et al.* Midbrain projection to the basolateral amygdala encodes anxiety-like but

not depression-like behaviors. *Nat. Commun.* **13**, 1532 (2022).

92.   Kumar, S. *et al.* Prefrontal cortex reactivity underlies trait vulnerability to chronic social

defeat stress. *Nat. Commun.* **5**, 4537 (2014).

93.   Friedman, A. K. *et al.* KCNQ channel openers reverse depressive symptoms via an active

resilience mechanism. *Nat. Commun.* **7**, 11671 (2016).

94.   Chaudhury, D. *et al.* Rapid regulation of depression-related behaviours by control of

midbrain dopamine neurons. *Nature* **493**, 532–536 (2013).

95.   Xiao, Q. *et al.* A new GABAergic somatostatin projection from the BNST onto accumbal

parvalbumin neurons controls anxiety. *Mol. Psychiatry* **26**, 4719–4741 (2021).

96.   Cooper, M. A. & Huhman, K. L. Blocking corticotropin-releasing factor-2 receptors, but

not corticotropin-releasing factor-1 receptors or glucocorticoid feedback, disrupts the

development of conditioned defeat. *Physiol. Behav.* **101**, 527–532 (2010).

97.   Challis, C. *et al.* Raphe GABAergic neurons mediate the acquisition of avoidance after

social defeat. *J. Neurosci.* **33**, 13978–88, 13988a (2013).

98.   Challis, C., Beck, S. G. & Berton, O. Optogenetic modulation of descending

prefrontocortical inputs to the dorsal raphe bidirectionally bias socioaffective choices

after social defeat. *Front. Behav. Neurosci.* **8**, 43 (2014).

99.   Calipari, E. S. *et al.* Dopaminergic dynamics underlying sex-specific cocaine reward. *Nat.*

Commun. **8**, 13877 (2017).

a Repeated x10

Social defeat 10 min → Social contact ~24 h

b Repeated social defeat stress

RSDS episodes
1   4   7   10
SI / SP Tests   SI / SP Tests   SI / SP Tests   SI / SP Tests

c Social interaction test

no target

target

d Sucrose preference test

e

Control ----
Susceptible ——
Resilient ——

Social interaction ratio vs RSDS episodes (#)

f SI ratio — Control, Stressed x7, Susceptible, Resilient

g % Sucrose preference — Control, Stressed x7, Susceptible, Resilient

h RSDS x 7 — Stressed x7

# of mice vs Social interaction time (s)

i RSDS x 10 — Susceptible, Resilient

# of mice vs Social interaction time (s)

j Stressed x7, Susceptible, Resilient — Cumulative distribution % vs Social interaction time (s)

k Stressed x7, Susceptible, Resilient — Social interaction time (s)

Susceptible and resilient subgroups emerge between 7 and 10 daily episodes of social defeat stress

Figure 1

Firing rate alterations in BNSTov^CRF neurons occur as adaptation to social stress, persisting in resilient, not susceptible mice.

Figure 2

Chemogenetic Modulation of BNSTov^{CRF} Neurons Recapitulates Behavioural Tipping Point

Figure 3

BNSTov^CRF calcium-dynamics encode stress effect on social interaction

Figure 4

BNSTov *Crfr1* is associated with the emergence of resiliency

Figure 5

Figure 1. Susceptible and resilient subgroups emerge between 7 and 10 daily episodes of social defeat stress.

a, Experimental design for repeated social defeat stress (RSDS). **b,** Experimental timeline of social defeat stress and behavioral tests: social interaction (SI) and sucrose preference (SP). **c,** Social interaction test schema involving target and no target trials. **d,** Sucrose preference test schematic. **e,** Effect of cumulative social defeat stress on social interaction measured as SI ratios, after 1, 4, 7, and 10 social defeat episodes (SDEs) (n=11-18 mice/group), two-way ANOVA interaction $F_{(6,134)}$=5.783, ****P<0.0001, row factor $F_{(3,134)}$=12.11, ****P<0.0001, $F_{(2,134)}$=4.086, column factor *P<0.05, Tukey's post-hoc test susceptible vs resilient ****P<0.0001, susceptible vs control **P<0.01, resilient vs control P=0.2751. SI test susceptible (SI test stressed x7 vs stressed x10) ****P<0.0001. **f,** Aggregated data on social interaction test across experiments. One-way ANOVA treatment $F_{(3,109)}$=14.61, ****P<0.0001. Tukey's post-hoc test control vs stressed x7 *P<0.05, control vs susceptible **P<0.01, stressed x7 vs susceptible ****P<0.0001, susceptible vs resilient ****P<0.0001 (n=25-37 mice). **g,** Sucrose preference test. One-way ANOVA treatment $F_{(3,93)}$=24.06, ****P<0.0001). Tukey's post-hoc test control vs stressed x7 P=0.77, control vs susceptible ****P<0.0001, control vs resilient P=0.9387, stressed x7 vs susceptible ****P<0.0001, stressed x7 vs resilient P=0.984, susceptible vs resilient ****P<0.0001 (n=23-25 mice). **h,** Distribution of stressed mice who underwent 7 SDEs. **i,** Distribution of mice who underwent 10 SDEs and sorted into susceptible and resilient mice. **j,** Cumulative distribution of all stressed mice. Kolmogorov-Smirnov (distance) 0.2054 Susceptible vs Resilient, ****P<0.0001. **k,** Time spent interaction socially with a novel conspecific. One-Way ANOVA $F_{(2,201)}$=76.63, ****P<0.0001. Tukey's

post-hoc susceptible vs resilient ***P<0.001, susceptible vs stressed x7 ****P<0.0001, stressed vs resilient P=0.4984 (n=72 susceptible, 64 resilient mice). All data represent means $\pm$ SEM. *P < 0.05, **P < 0.01, ***P < 0.001, ****P<0.0001, ns = not significant.

Figure 2. Firing rate alterations in BNSTov^{CRF} neurons occur as adaptation to social stress, persisting in resilient, not susceptible mice.

a, Mouse genotype and timeline of cell-attached electrophysiology experiments. **b,** Fluorescence-guided cell-attached electrophysiology setup, brain slice of the BNST (CRF cells, tdTomato) and DIC image of CRF neurons, scale bar 0.63 mm. **c,** Representative trace of BNSTov^{CRF} positive and negative neurons of control, stressed (SDEs x 7), susceptible, and resilience mice. **d,** Firing rate of CRF$^+$ neurons (n=9-31 cells/group), ****P<0.0001, one-way ANOVA, Tukey's multiple comparison's test, control vs stressed (SDEs x 7) **P<0.01, susceptible vs resilient **P<0.01, susceptible vs stressed (SDEs x7), ****P<0.0001. Firing rate of CRF$^-$ neurons (n=7-15 cells per 4-6 mice/group), one-way NOVA, $F_{(3, 45)}$=3.113, *P<0.05, Tukey's multiple comparison's test, control vs stressed x7 P=0.9192, control vs susceptible P=0.0769, control vs resilient P=0.1742, stressed (SDEs x 7) vs susceptible P=0.1370, stressed x7 vs resilient P=0.3219, susceptible vs resilient P=0.9673. **e,** Correlation of firing rate with social interaction ratio: CRF$^+$, R^2=0.5726, *P<0.05; CRF$^-$, R^2=0.0609, P=0.5219. **f,** Representative sample of burst firing. **g,** Representative sample of tonic firing. **h,** Percentage of bursting cells per animal group. **i**, Percentage of spikes within burst. **j,** Number of spikes per burst. **k,**

Number of bursts per cell. (n=10-52 cells per 4-6 mice/group). All data represent means $\pm$ SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, ns = not significant.

Figure 3. Chemogenetic modulation of BNSTov^{CRF} neurons bidirectional recapitulates behavioural tipping point.

a, Viral targeting of BNSTov. **b,** Experimental design of chemogenetic manipulation of BNSTov neurons with CNO-drinking water construct. **c,** Social interaction test, two-way ANOVA $F_{(3,68)}$=18.01, row ****P<0.0001, row factor $F_{(3, 68)}$=9.965, P=0.2616 (time), row factor x time $F_{(1,68)}$=1.281, ****P<0.0001, Sidak's multiple comparisons test mCherry (s), **P=0.0017 (n=10 mice); mCherry (r), P=0.1322 (n=9 mice); hM4Di, ****P<0.0001 (n=10 mice); hM3Dq, **P=0.0018 (n=9 mice). **d,** Sucrose preference test. one-way ANOVA treatment $F_{(3,27)}$=8.310, ***P<0.001, Tukey's post-hoc testing susceptible vs hM4Di, P=9619; susceptible vs resilient, **P<0.01; susceptible vs hM3Dq, **P<0.01; hM4Di vs resilient, **P<0.01; hM4Di vs hM3Dq, *P<0.05; resilient vs hM3Dq P=0.9368 (7-9 mice per group). **e,** Distance traveled, one-way ANOVA $F_{(3,27)}$=1.031, P=0.3944. **f,** Percentage of susceptible or resilient mice: DIO-hM4Di, 100% susceptible; DIO-hM3Dq, 89% resilient and 11% susceptible; mCherry, 40% resilient and 60% susceptible. Chi-square test=15.66, df=2, ***P<0.001 (n=21 mCherry, 10 hMDi, 9 hM3Dq); Fisher's post-hoc test mCherry vs hM4Di, *P<0.05; mCherry vs hM3Dq, *P<0.05; hM4Di vs hM3Dq, ****P<0.0001. **g,** Repeated social defeat DREADDs manipulation (4-7 episodes of stress), 2-way ANOVA treatment $F_{(1,16)}$=0.01749, P=0.8964; SI Test 1 vs SI Test 2,

$F_{(1,16)}=0.2377$, P=0.6325; interaction $F_{(1,16)}=0.07055$, P=0.6218 (n=5-6 mice/group). **h,** RSDS DREADDs manipulation (7-10 episodes of stress), 2-way ANOVA treatment, $F_{(1,22)}=8.675$, **P< 0.01; SI Test 1 vs SI Test 2, $F_{(1,22)}=2.549$, P=0.1247; interaction $F_{(1,22)}=2.549$, P=0.1247. Sidak's post-hoc test SI Test 1 vs SI Test 2, control, P=0.5533; DIO-hM3Dq, *P<0.05 (n=6-7 mice/group). **i,** RSDS DREADDs manipulation (10-13 episodes of stress), 2-way ANOVA $F_{(1,30)}=1.192$, treatment, P=0.2623. SI Test 1 vs Si Test 2, $F_{(1,30)}=0.3132$, P=0.2836. Interaction $F_{(1,30)}=0.2623$, P=0.5799. All data represent means ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, ns = not significant.

Figure 4. BNSTov[CRF] calcium-dynamics encode stress effect on social interaction

a, Viral targeting of the BNSTov. **b**, Experimental design of multiplexed chemogenetics with drinking water-CNO delivery and fiber photometry. **c,f,** Representative calcium recordings of mCherry susceptible (S), mCherry resilient (R), hM3Dq, and hM4Di groups, respectively. Vertical scale bar is equal to z-score of 1, and horizontal scale bar is equal to 10 s. **d**, **g**, Representative averaged trace centered around interaction bout for the four groups stated above. **e,** Two-way RM ANOVA row factor $F_{(3,14)}=1.401$; pre vs post-CNO $F_{(1,14)}=0.8179$; subject $F_{(14,14)}=1.195$; Row x SI Test 1/SI Test 2, $F_{(3, 14)}=9.343$, **P<0.01. Sidak's post-hoc test, susceptible vs resilient, P=0.137 vs *P=0.05, respectively (n=4 mice/group). **h,** hM4Di vs hM3Dq, P=0.3069 vs *P<0.05, respectively (n=4-5 mice/grp). **i,** SI Test 1 social interaction test. One-way ANOVA $F_{(3,16)}=0.291$, P=0.8308 (n=5 mice/group). SI Test 2 social interaction test, One-Way ANOVA $F_{(3,16)}=17.83$, ****P<0.0001; Tukey's post-hoc test, susceptible vs resilient, ****P<0.0001; susceptible

vs hM3Dq, ***P<0.001; susceptible vs hM4Di, P=0.0762; resilient vs hM3Dq, P=0.8506; resilient vs HM4Di, **P<0.01; hM3Dq vs hM4Di, *P<0.05 (n=5 mice/group). **j**, Correlation of SI ratio and z-score delta F/F upon social entry, simple linear regression pre-CNO $F_{(1,15)}$=0.7674, P=0.3948; SI test 2 $F_{(1,17)}$=7.268, *P<0.05. Intersection of lines, $F_{(1,32)}$=6.896, *P<0.05 (n=18 mice). **k**, Number of social interaction bouts, one-way ANOVA $F_{(3,16)}$=1.346, P=0.2949 (n=5 mice/group). All data represent means ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, ns = not significant.

Figure 5. BNSTov *Crfr1* is associated with the emergence of resiliency

a, Experimental timeline of RNAScope *in situ* hybridization of mice that underwent social defeat stress. **b**, Representative images of control, stressed x7, susceptible, and resilient. 20x magnification, scale bar 0.64 mm. **c**, Schematic of anterior dorsal BNST. **d**, *Crfr1* mRNA colocalization, one-way ANOVA $F_{(3,18)}$=20.91, ****P<0.0001; Tukey's posthoc test, control vs stressed x7, P=0.0701; control vs susceptible, P=0.3336; control vs resilient, ****P<0.0001; stressed x7 vs susceptible, **P<0.01; stressed x7 vs resilient, *P<0.05; susceptible vs resilient, ****P<0.0001 (n=4-6 BNST brain samples). **e**, *Crfr2* mRNA colocalization, one-way ANOVA $F_{(3,10)}$=1.790, P=0.2166 (n=3 BNST brain samples). **f,** RNAscope BNST subregion comparison, two-way ANOVA interaction $F_{(6,24)}$=1.057, P=0.4147; row factor $F_{(2,24)}$=0.6092, P=0.5520; column factor $F_{(3,24)}$=1.549, P=0.2275 (n=3 BNST brain samples). **g**, Viral injection site. **h**, Experimental timeline. **i**, Optogenetic and social behavioral setup. **j**, Optogenetics social interaction, two-way ANOVA interaction $F_{(1,28)}$=5.059, *P<0.05; row factor $F_{(1,28)}$=1.061e-005, P=0.9974; column factor

$F_{(1,28)}$=1.419, P=0.2436. Sidak's post-hoc test, *Crf*::tdT, *P<0.05; *Crf*::ChR2, P=0.7233

(n=7-9 mice/group). Optogenetic manipulation unpaired t-test, *Crf*::tdT, t=2.556, df=40,

*P<0.05 (n=8-9 mice); *Crf*::ChR2, t=0.7758, df=40, P=0.442457 (n=8-9 mice); *Crf*::ChR2,

t=2.369, df=40, *P<0.05 (n=6 mice). Holm-Sidak method for multiple comparisons. **k**,

Representative image, scale bar 0.64mm. **l**, Optogenetics-RNAScope experiment. One-

Way ANOVA $F_{(3,9)}$=13.53, **P<0.01. Tukey's post-hoc test, *Crf*::ChR2 (stress + stim) vs

Crf::tdT (resilient), P=0.7872; *Crf*::ChR2 (stress + stim) vs. *Crf*::tdTomato (s), **P<0.01;

Crf::ChR2 (stress + stim) vs. *Crf*::chR2 (stim), **P<0.01; *Crf*::tdTomato (r) vs.

Crf::tdTomato (s), *P<0.05; *Crf*::tdTomato (r) vs. *Crf*::ChR2 (stim), *P<0.05; *Crf*::tdTomato

(s) vs. *Crf*::ChR2 (stim), P=0.9988. All data represent means ± SEM. *P<0.05, **P<0.01,

***P<0.001, ****P<0.0001, ns = not significant.