Excitation creates a distributed pattern of cortical suppression due to varied recurrent input

Jonathan F O'Rawe[1], Zhishang Zhou[1], Anna J Li[1,4], Paul K LaFosse[1,2,3], Hannah C Goldbach[1,5], Mark H Histed[1,*]

1. National Institute of Mental Health Intramural Program, National Institutes of Health, Bethesda MD USA.
2. National Institutes of Health-University of Maryland Graduate Partnerships Program, Bethesda, MD USA
3. Neuroscience and Cognitive Science Program, University of Maryland College Park, College Park, MD USA

Current affiliations:
4. Department of Biological Structure, University of Washington, Seattle WA USA
5. National Eye Institute Intramural Program, National Institutes of Health, Bethesda MD USA

Summary

Dense local, recurrent connections are a major feature of cortical circuits, yet how they affect neurons' responses is unclear, with some studies reporting weak recurrent effects, some amplification, and others showing instead local suppression. Here, we show that optogenetic input to mouse V1 excitatory neurons generates salt-and-pepper patterns of both excitation and suppression. Responses in individual neurons are not strongly predicted by that neuron's direct input. A balanced-state network model reconciles a set of diverse observations: the observed dynamics, suppressed responses, decoupling of input and output, and long tail of excited responses. The model shows recurrent excitatory-excitatory connections are strong and also variable across neurons. Together, these results demonstrate that excitatory recurrent connections can have major effects on cortical computations, by shaping and changing neurons' responses to input.

---

[*] Correspondence and Lead Contact: mark.histed@nih.gov

Introduction

The cerebral cortex of mammals is specialized into areas that perform different functions[1].
Animals from rodents to primates have several different visual cortical areas, each containing
neurons with different types of selectivity[2–4]. In principle, these different representations in
different visual areas could be created purely by feedforward mechanisms, where
transformations happen via projections from one area or layer to the next, without outputs of a
neuron feeding back (directly or indirectly) to influence that neuron's activity. In fact, in a variety
of artificial neural networks, much or all computation is provided by feedforward mechanisms[5].

Yet in the brains of animals and humans, cortical recurrent connectivity is extensive. Most
excitatory connections that a cortical neuron receives originate within a few hundred microns of
their cell bodies[6–8]. Such recurrent connections can in principle have large effects on neural
computation[9], dramatically changing how cortical neurons respond to input.

How recurrent connections affect cortical computation is not fully understood, but important
aspects of the structure of cortical recurrent connectivity have been determined. Some features
of cortical network activity, such as irregular firing, are well-described by balanced-state models
which assume strong recurrent coupling between excitatory and inhibitory neurons (either
moderately strong, yielding 'loose balance', or very strong, yielding 'tight balance'[10]). Work using
inhibitory perturbations has shown that not just excitatory-inhibitory connectivity is strong, but
the average excitatory-excitatory connectivity is strong as well. More precisely, cortical recurrent
excitatory coupling is strong enough that the excitatory network is unstable and self-amplifying,
a phenomenon described by inhibition-stabilized network models (ISNs)[11–14].

While some consensus has developed on these average cortical connectivity properties (but
see[15]), the effect recurrent connections have on transforming sensory or input signals has been
less clear. For example, some recent studies have shown that certain patterns of excitatory
input can be amplified by the cortical network[16,17], consistent with some theoretical
predictions[18,19]. On the other hand, however, some studies have shown that nearby neurons can
be substantially suppressed by stimulation that excites a single or a small ensemble of
excitatory cortical neurons[20,21]. How excitatory and inhibitory neurons might interact through
recurrent connections to create such suppression has not been determined.

Here, to understand how cortical neurons' responses are shaped by the cortical recurrent
network, we stimulate excitatory cells in the visual cortex optogenetically and record responses
of local neurons with electrophysiology and two-photon imaging. First, we find that stimulation of
excitatory cells leads to a salt-and-pepper pattern of local suppression, consistent with the
pattern of excited and suppressed cells produced when animals see a strong visual stimulus. To
understand how this suppression effect might arise from cortical recurrent circuitry, we examine
both the patterns of firing rate changes and the dynamics of responses. Recent theoretical work
has shown that cortical visual responses can be "reshuffled" by additional excitatory input[22] —
that is, strong average recurrent coupling allows individual neurons' firing to change significantly
in response to input while the distribution of population activity is little-changed[23,24]. We
implement this scenario in a conductance-based simulation and find that it can explain the

suppression we observe. In addition, our data is consistent with substantial variability in local
recurrent connectivity, with some neurons receiving large net recurrent excitation and others
smaller or net suppressive recurrent input. Our results go beyond prior work that found strong
average recurrent connectivity, showing that *variance* in excitatory-excitatory connectivity must
also be substantial, and further show that this variance in recurrent connectivity can decouple
neurons' firing rate responses from the direct input they receive.

The suppression we observe during excitatory cell stimulation occurs in individual cells, but the
mean response is elevated. This increase in mean, however, seems at odds with the prior
finding that single-cell stimulation leads to inhibition on average[20]. To resolve this, we simulate
the effect of single cell stimulation and find that the difference in the two results can be
explained by the activation state of the cortical network. Increasing activity in the network with
visual stimulation results in a slight decrease in mean responses to stimulation, showing the
prior results and our current results can be described in the same model framework.

Thus, a balanced-state cortical model, with strong average coupling and variability in recurrent
connectivity, explains many features of our data, including dynamics and neural response
distributions. These results show how cortical neural suppression can be generated from
excitatory input: variability in recurrent input means that firing rate responses are decoupled
from (are only weakly affected by) the level of excitatory input we provide to that cell. This arises
because much of the input a cell receives comes from recurrent sources. Because recurrent
input varies from cell to cell, the result is many excited cells, but also a substantial number of
suppressed neurons.

Results

Strong visual input leads to salt-and-pepper distributed suppression in primary visual cortex

We first measured local patterns of suppression in visual cortex in response to visual stimuli. We presented small high-contrast visual stimuli to headfixed mice while measuring activity in V1 layer 2/3 neurons via two-photon imaging (Fig. 1A). We expressed GCaMP7s in all neurons via viral injection (AAV-hSyn-GCaMP). Animals were kept awake and in an alert state[25] with occasional drops of water reward.

Figure 1: V1 neurons show salt-and-pepper suppression to strong visual stimuli. (A) Experimental setup. Awake mice viewed a small (15 degree diameter) visual stimulus with rapidly changing frames of oriented noise (Methods). **(B)** Example 2-photon imaging data from layer 2/3 of V1 in response to the stimulus, during the transient and **(C)** steady-state periods. Time intervals used for averaging in (B-D) displayed in green and purple in (G). Intermixed (salt-and-pepper) elevated and suppressed responses emerge during the steady-state period. **(D)** Deconvolved responses from (C), projected onto segmented cell masks (Methods). **(E)** Example dF/F trace for one elevated and one suppressed cell. Shaded regions: SEM across trials. Shaded red: optogenetic stimulation duration. **(F)** Deconvolution of the traces in (E) reveals an initial transient period and then a steady-state response. **(G)** Average response for all elevated and suppressed cells in (B-D, N = 1, pos. neurons = 42, neg. neurons = 28). **(H)** Spatial distribution of elevated (red) and suppressed (blue) cells collapsed across animals (N = 3; 339 neurons), showing random distribution of neurons across the cortex (statistical analysis; Fig. S1F-H). **(I)** Visual response amplitudes are similar across animals. Thin lines: CDFs for individual animals, thick line: population CDF. Inset: medians are near zero, m_1-m_3: individual animals, error bars: ± SEM. **(J)** Proportion of cells suppressed in each mouse. Error bars: Wilson score 95% confidence intervals. Black line: group mean (44% ± 7%). See also   Fig. S1.

We imaged responses to two types of high-contrast visual stimuli, a fast-changing stimulus
designed to minimize adaptation ("oriented noise", Fig. 1A)[26–28] and a drifting grating (Fig. S1D).
We found a salt-and-pepper mix of suppressed and excited cells (Fig. 1B,C), with suppression
stronger after the initial stimulus response (Fig. 1C). In other words, in response to both types of
visual stimuli, we found some cells that responded with strongly elevated steady-state
responses, and other cells that showed suppressed responses (Fig. 1C–F, Fig. S1A-E).

Deconvolving fluorescence responses to yield a proxy measure of spike rate confirmed this salt-
and-pepper pattern, with substantial numbers of suppressed and excited neurons intermingled
(Fig. 1D). The deconvolution revealed an initial transient response in excited cells (Fig 1F,G),
followed by either an elevated or suppressed steady state.

We confirmed that the spatial distribution of elevated and suppressed neurons was randomly
scattered across the cortex (Fig. 1H). We found our data was consistent with random scatter
(data vs 2d Poisson process model for spatial randomness, $p > 0.05$, Bonferroni correction, Fig.
S1F-H).

The viral expression strategy we used for these experiments results in both excitatory and
inhibitory neurons that express GCaMP. However, the large fraction of suppressed neurons
(Fig. 1D,H-J; proportion suppressed 44% ± 7%, N=3 animals, mean ± standard error) implies
that it is not that a group of inhibitory neurons was suppressed by stimulation, but that many
excitatory neurons were suppressed. Below, we confirm with electrophysiology and imaging that
optogenetic excitatory input produces suppression in many excitatory cells.

Optogenetic excitatory drive also results in sparse and distributed suppression

To examine the influence of recurrent excitatory-inhibitory circuits on local response properties,
we next measured V1 responses while optogenetically stimulating excitatory cortical cells (Fig.
2A). Direct stimulation allows us to exclude some feedforward mechanisms for
suppression — for example, to argue against the possibility that cortical suppression is
generated principally by suppression of thalamic inputs[29].

We injected a Cre-dependent excitatory opsin (soma-targeted ChrimsonR, or stChrimsonR) in
layer 2/3 of a mouse expressing Cre in excitatory neurons only (*Emx1-Cre*[30]), and expressed
GCaMP7s in all neurons with a second virus (AAV-hSyn-GCaMP7s) (Fig. 2B,E).

With optogenetic stimulation we also found a clear salt-and-pepper distribution of elevated and
suppressed responses (Fig. 2C,F,H; short stimulation pulses Fig. 2B-D, long pulses with
imaging of steady-state during stimulation, Fig. 2E-G). Neural responses to stimulation increase
as power increases (Fig. 2D; asymptote may be due to opsin saturation.) As in the case of
visual responses, we confirmed that the spatial patterns of responses were compatible with
random scattering (all p's > 0.05, Fig. S1F-H). The proportion of suppressed neurons with
optogenetic stimulation (Fig. 2I,J; 38% ± 8%, mean ± SEM) was comparable to that seen with
visual stimulation (Fig. 1IJ). These optogenetic data suggest that the network is being driven to
a new steady state or fixed point by input. While there was a slight decay in the excited
population's response at high power (perhaps due to network effects, spike rate adaptation, or

opsin desensitization), at moderate stimulation power (1 mW, Fig. 2G), deconvolved firing rates
are largely constant while stimulation is on.

Figure 2: Salt-and-pepper elevation and suppression to optogenetic excitation. (A) Experimental setup, using
two-photon imaging (GCaMP7s, all cells, 920nm) and optogenetic excitation of excitatory neurons (stChrimsonR,
595nm). **(B)** Example field of view. **(C)** Deconvolved steady-state response (scaled to match dF/F %) to optogenetic
stimulation (200 ms duration) from (B). Red: elevation of firing rate relative to baseline, blue: suppression. **(D)**
Increasing power leads to stronger elevation and suppression (steady-state response) in their respective populations.
Shaded region: SEM across cells. **(E)** Field of view from an example animal stimulated with long (4 sec) optogenetic
pulses; stimulation during imaging flyback (Methods). Gray: areas omitted from analysis to exclude stimulation
artifact. **(F)** Deconvolved response to stimulation, conventions same as (C). **(G)** Population timecourses for cells in
(F)**.** Red region: optogenetic stimulation period. Steady-state response averaging period: 200- 3750 ms. Light lines:
individual cell traces, heavy lines: population averages. Shaded region (largely obscured by thick lines): SEM across
cells. **(H)** Spatial distribution of elevated (red) and suppressed (blue) cells collapsed across all animals (N = 3), same
conventions as Fig 1H. Statistical analysis: Fig. S1F-H. **(I)** Optogenetic response amplitudes are similar across
animals. Conventions as in Fig. 1I**. (J)** Proportion of cells suppressed by optogenetic stimulation in each mouse. Error
bars: Wilson score 95% confidence intervals. Black line: group mean (38% ± 8%).

We confirmed the opsin we used was expressed only in excitatory cells using fluorescence in-
situ hybridization. We labeled excitatory, inhibitory, and stChrimsonR-expressing neurons
(RNAScope, ACD Inc; Fig. S2A,B). Excitatory neurons expressed the opsin (Fig. 2I), but as
expected for AAV expression[31], not all excitatory neurons were opsin-positive (59%, N =
115/195, Wilson score 95% CI: [52.0%,65.7%], Fig. S2A). None of the inhibitory neurons (24%
of neurons in the sample, N = 62/257) showed expression of the opsin (Fig. S2B).

The two-photon imaging experiments showed a salt-and-pepper pattern of excitation and
suppression within the imaging fields of view. To examine whether this salt-and-pepper pattern
exists at larger distances from the stimulation site, we used electrophysiology. We recorded
neural responses to stChrimsonR stimulation using a silicon electrode array (Fig. 3A,E) and the
same viral strategy for opsin expression as we used with imaging.

We found both elevation and suppression across all distances (Fig. 3B–D) and depths (Fig. 3F–
H) from the stimulation site, suggesting a similar salt-and-pepper organization of elevated and
suppressed cells extends over distance. Across the population of recorded neurons, 56.6% (77
of 136) showed an elevated steady-state response to the optogenetic stimulation, and 36.0%
(49 of 136) showed a suppressed steady-state response, comparable to our two-photon
measurements (Fig. 2). Both elevated and suppressed cells on average showed an initial
(positive) transient followed by a (positive or negative) steady-state response (Fig. 3I,J).

The electrophysiological recordings show similar dynamics as the deconvolved imaging
timecourses (Fig. 2H), except for one feature: the recordings show an initial brief positive
transient in the suppressed cells (Fig. 3B–D, F–H, blue lines; Fig. 3J) not just in the elevated
cells as in the imaging data. This transient is likely concealed in the imaging data due to the
slower timescale of imaging. The imaging frame rate (30 Hz; 33 ms frames) is slower than the
transient, so within one frame the positive transient would be averaged with suppression,
yielding a result near zero. In the case of elevated cells, the positive transient is averaged with
an elevated steady state, and so the response in that frame remains positive.

Global spatial patterns arise from trends in local salt-and-pepper suppression

The neurophysiology data showed some evidence of a larger-scale organization on top of the
local salt-and-pepper distribution of elevation and suppression. Over distances of more than a
millimeter from the stimulation site, we found that the number of elevated units gradually
decreased (Fig. 3K,L; Pearson's chi-squared test, χ^2 = 51.31, df = 3, p < 0.001) and the number
of suppressed units gradually increased (Fig. 3M,N; χ^2 = 44.83, df = 3, p < 0.001; see Fig. S2E-
H for unit counts as a proportion of total units). There was also a similar trend in neurons' firing
rates (Fig. S2I,J). Elevated single units showed less elevated firing rate with distance from the
stimulation site, and suppressed single units showed more suppression with distance from the
stimulation site, though the linear trend between distance and population response was stronger
in unit counts than in average population firing rates (Pearson's r = -0.11, df = 29, p = 0.56,
Pearson's r = 0.32, df = 46, p < 0.05; Fig. S2I,J). Notably, however, the number of elevated
neurons did not go to zero even at 1.2 mm from the stimulation site: only the relative numbers of

elevated and suppressed neurons changed. This suggests that the salt-and-pepper organization
we saw with imaging persists across the cortex.

Figure 3: Stimulation of V1 excitatory neurons yields salt-and-pepper organization across the cortex. (A)
Neural responses recorded across the cortex. Recordings in vivo from awake mice. **(B, C, D)** Example neurons at
three distances from stimulation light (0 µm, 400 µm, 1200 µm), showing elevated and suppressed cells at all
distances. **(E)** Neural responses recorded through cortical depth. **(F, G, H)** Example neurons recorded at three
depths (250 µm, 550 µm, and 800 µm), showing elevated and suppressed cells at different depths. **(I)** Population
average timecourses of elevated cells. Blue bar: interval for steady-state rate calculation. Shaded regions: SEM
across cells. **(J)** Population time courses of suppressed cells, same conventions as (I). **(K)** Counts of elevated units
(single and multi-units) by distance and depth, smoothed with a Gaussian kernel for display. **(L)** Distribution of
elevated steady-state responses across horizontal distance, summed across depth. Shaded region: Wilson score
95% CIs. Note lower limit of y-axis not zero. **(M-N)** Same as (K-L), but for units with suppressed steady-state
responses. See also Fig. S2.

The trends over distance we saw with physiology, however, give only a partial view into how
population responses varied with distance from the stimulation site. To measure the extent of
suppression across the cortex, we turned to widefield, mesoscale calcium imaging. For these
experiments, we expressed GCaMP in all excitatory cells using a genetic mouse line (to
maximize consistency of GCaMP expression across cortical distance; Fig. 4A; Ai148::Cux2-

CreERT2, or Ai162::Cux2-CreERT2, see Methods). We restricted expression of stChrimsonR to
excitatory cells using the CamKIIa promoter (AAV-CamKIIa-stChrimsonR) and stimulated while
simultaneously imaging responses.

We saw clear spatial patterns in widefield imaging, broadly consistent with the spatial trends we
saw in the electrophysiology data. During the initial frame of stimulation (~7 Hz imaging, 140 ms
frame period), we saw an increase in activity both at the center of the stimulation light and
extending some distance outside the center of expression (Fig. 4B,E,H).

Figure 4: Widefield imaging of excitatory neurons shows average center-surround organization during
steady-state periods. (A) Experimental setup: stChrimsonR in excitatory neurons via viral transfection (AAV-
CamKIIa-stChrimsonR), expression of GCaMP via mouse line (either Ai148::Cux2-creERT2, GCaMP6f, or
Ai162::Cux2-creERT2, GCaMP6s; induced with tamoxifen as adult; Methods). Right: imaging field of view for one
animal. **(B-C)** Mean deconvolved response (see Fig. S3) during first frame (B) and during the late stimulation period
(C) in an example animal (Fiber for light delivery slightly obstructs the imaging field, see Fig. S4D-F). **(D)** Average
response to stimulation over time (N = 3). Red shaded region: stimulation period, orange bar: first frame, maroon bar:
late stimulation (steady-state) time period. **(E,F)** Average responses, N = 3 animals. Responses for each animal were
aligned spatially to the peak during the late stimulation period (Methods), smoothed for visualization. **(G)** Response
as a function of distance, averaged from data in E, F. Smoothing: LOWESS. Shaded regions: bootstrapped 95% CIs.
Vertical lines: zero crossings and inflection points. Zero crossings defined by shortest distance at which 95% CI
included zero. Black lines: late stimulation period. Solid black: first zero crossing, dotted black: local minimum. Red
dashed line: early response, first zero crossing. **(H, I)** Same as (E,F) but with superimposed circles whose radii
correspond to lines in (G). See also Figs. S3-4.

A center-surround pattern emerged later in the stimulation pulse (Fig. 4C,F,I) consistent with the
large-scale patterns in the electrophysiological recordings. The area with maximum
stChrimsonR expression continued to show an elevated response, while a donut-shaped region
around it was suppressed (see Fig. S4G-I for spatiotemporal response). The activated area in
the center reflected the area of expression, measured with fluorescence imaging of the cortical
surface (Fig. S4A-F). To examine these timecourses (Fig. 4D), we deconvolved imaging
responses to yield approximations to spike rate changes. We compared several different
deconvolution methods and found suppression in all cases (Fig. S3). The suppression was
strongest about 500 µm from the center of our laser stimulus, and extended over 1 mm from the
stimulation center (Fig. 4 G–I). In electrophysiology, the number of suppressed cells increases
by a factor of two over approximately this distance (Fig. 3K-N), and therefore the increased
number of suppressed individual neurons may be the substrate for the suppression in this
imaging data.

In summary, the physiology and imaging data together support the idea that suppressed and
elevated neurons are locally organized in a salt-and-pepper pattern, and that the proportion of
suppressed to elevated neurons increases with distance from the stimulation site. This change
in the proportion of suppressed cells results in a center-surround pattern that can be seen with
population-level imaging, with net suppression in excitatory cells emerging, after an initial
positive transient, about 500 µm away from the stimulation site.

Response dynamics support a balanced-state excitatory-inhibitory network that is
driven to a new steady state by input

If suppression is due to local recurrent network effects, we would expect excitatory cells to be
recruited first by stimulation, and then inhibitory cells should receive inputs from excitatory cells
and respond slightly later. After this first few milliseconds, balanced-state models predict that
excitatory and inhibitory cells should later show similar response distributions[10,11,32,33]. This is in
contrast to weakly-coupled models, or a feedforward inhibition framework, where excitatory and
inhibitory populations change firing rates in opposite directions: that is, input drives inhibitory
cells to increase their rates, inhibiting excitatory cells, which then decrease their rates.

Figure 5: Response dynamics are consistent with steady-state suppression shaped by a recurrent network mechanism. Wide-waveform (excitatory) units have slightly earlier onset latencies than narrow (inhibitory) units, but other quantities do not differ across neural populations. **(A)** Wide/narrow sorting approach. Bimodal widths (classification threshold: 0.445 ms). **(B)** Average traces for narrow- and wide-waveform units (Methods). Inset: enlarged view of laser onset, highlighting latency difference between narrow and wide units. Green markers: time to half peak. **(C)** Onset latencies slightly shorter for wide-waveform units. **(D)** Peak firing rate does not differ between wide and narrow units (medians, wide: 24.5 spk/s, narrow: 31.2 spk/s). **(E)** Steady-state firing rate does not differ (medians, wide: 4.0 spk/s, narrow: 5.1 spk/s). **(F)** Onset latencies do not differ for elevated and suppressed populations. Conventions as in C. **(G)** Time to steady state does not differ for elevated and suppressed populations. See also Fig. S5.

Our data supports the balanced-state recurrent model (Fig. 5A) — we saw differences in excitatory and inhibitory responses in the first few milliseconds, but at later times distributions of excitatory and inhibitory rates were similar.

We classified cells into putative excitatory and inhibitory classes by waveform (Fig. 5A). We have previously confirmed[13] with *in vivo* pharmacology that narrow-waveform cells are inhibitory interneurons, likely PV-positive fast-spiking cells, while wide-waveform cells are primarily excitatory neurons. We saw here that wide-waveform (largely excitatory) neurons have a slightly faster onset latency than narrow-waveform inhibitory cells, faster by approximately 2.5 ms (Fig. 5B, inset, 5C; narrow latency 7.9 ms, wide latency 5.4 ms, difference 2.5 ms, Mann-Whitney U = 1256.0, p < 0.01; onset latencies computed via curve-fitting to rising phases, see Fig. S5A for details).

If subgroups of excitatory and inhibitory cells composed the suppressed and elevated populations, we might expect to see differences in the dynamics of elevated and suppressed

cells. But we found no significant differences in onset time or time to steady state for elevated
and suppressed neurons (Fig. 5F,G, onset time Mann-Whitney U = 1741.0, p = 0.17, time to
steady state, Mann-Whitney U = 801.0, p = 0.32). This was also true when restricting the
analysis to only wide-waveform cells (Fig. S5C,D). Another possibility could have been that the
neurons with suppressed steady-state responses were cells that did not express opsin. But the
similar onset latencies of the elevated and suppressed cells (Fig. 5F) excludes that possibility,
and provides further support to the idea that instead a balanced-state recurrent network
explains the suppression.

Beyond the differences in onset latency, we found other response dynamics were not different
between excitatory and inhibitory cells. Consistent with a recurrent network with strongly
coupled excitation and inhibition, we found that both excitatory and inhibitory cell populations
increase their average firing rate when excitatory cells are stimulated (wide mean Δ: 14.54
spk/s, t = 5.52, df = 93, p < 0.001, narrow mean Δ: 13.53 spk/s, t = 3.07, df = 41, p < 0.01). That
is, both excitatory and inhibitory populations contain elevated and suppressed neurons, though
elevated cells dominate both averages (Fig. 5E). Further, the initial transient and steady-state
firing rate medians were not detectably different between inhibitory and excitatory cells
(transient: Mann-Whitney U = 1816.0, p = 0.23, steady-state Mann-Whitney U = 1866.0, p =
0.31, Fig. 5D,E). Also, time to steady state for wide-waveform and narrow-waveform cells did
not differ (Fig. S5B), consistent with the idea that the steady-state dynamics emerge from
integration of both inhibitory and excitatory inputs. Overall, the response distribution and
dynamics we observed in inhibitory and excitatory cells are consistent with a strongly-coupled
recurrent network.

A neuron's response is only weakly predicted by optogenetic input to that neuron

We used the imaging data to determine if the suppression we observed was explained by
variation across cells in optogenetic drive. We found that while indeed there was variability in
different cells' responses, there was very little relationship between opsin expression and cells'
firing rate changes. To estimate the optogenetic drive to individual neurons, we measured
fluorescence of mRuby2 (fused to stChrimsonR) in donut-shaped regions around each cell's
membrane using two-photon imaging (Fig. 6, Fig. S6). The measured *in* vivo distribution of
opsin expression was well-fit by a lognormal distribution after excluding the 16.8% (Wilson score
95% confidence interval: [12.7%, 22.1%]) of cells with low fluorescence (Fig. 6C, see Methods).
The *in vivo* estimate of the percentage of non-expressing cells was lower than what we
observed with histology, perhaps because we selected FOVs with dense opsin for *in vivo*
imaging. At the same time, however, our *in vivo* observations are consistent with past work that
finds that AAV transfects adult neurons in a non-uniform way*, finding substantial variability in
opsin expression from cell to cell[13,31].

We found that the amount of opsin-related fluorescence explained little of the variance in
steady-state responses (Fig. 6A-D, Pearson's r = 0.21, df = 106, p < 0.05, high fluorescence
neurons excluded [> 0.5]; Fig. 6D-E, population: at 1 mW: Pearson's r = 0.18, df = 219, p <
0.01, at 2 mW: Pearson's r = 0.17, df = 219, p < 0.01).

This striking decoupling effect — that the amount of opsin input barely predicts how cells' firing
rates are modulated by stimulation — suggests that a given cell's response may not be dictated
by input to that cell, but instead by recurrent inputs.

Notably, both high-expressing cells and low-expressing cells showed little relationship between
opsin expression and response (Fig. 6E, red and gray lines). This supports the idea that the
decoupling is not due to cell-autonomous intrinsic properties but indeed due to recurrent
network inputs. To further test this, we measured the variability of neural responses as a
function of stimulation intensity. If a neuron's response were in fact controlled primarily by its
opsin expression (the optogenetic input to that neuron) and not network input, increasing the
input intensity should keep the variance in response the same, or reduce it, because the fixed
opsin level is the principal source of response drive (Fig. S2K). Or, if response was dictated by
opsin level, increasing intensity might produce a bimodal response distribution as the
optogenetically-driven neurons separate from non-expressing neurons (Fig. S2L). We found
support for none of these possibilities. Instead, the response pattern increased in variance as
stimulation grew stronger (Fig. S2M-O), supporting the idea that it was network input, not opsin
level, that controlled cells' responses.

We next turned to simulations, fit to our data and building on the recent theoretical advances of
Sanzeni et al. (in press), to more completely characterize recurrent network influences on
neurons' responses.

Figure 6: stChrimsonR expression only weakly predicts 2-photon steady-state response. (A) Left: Red
(stChrimsonR-mRuby2) and green (GCaMP7s) fluorescence of an example cell with a suppressed steady-state
response during optogenetic stimulation. Right: donut-shaped region of interest (ROI), inner and outer boundary
calculated by shrinking or expanding the cell border (CaImAn, Methods) **(B)** Same as (A), except for an example cell
that shows an elevated steady-state response. The suppressed cell shows brighter red fluorescence than the
elevated cell, quantified in D. **(C)** Distribution of opsin fluorescence intensity (N=3 animals). Orange: Lognormal fit,
excluding non-expressing cells (gray; see Fig. S6). **(D)** Example relationship between opsin expression and
response; only a weak relationship is seen. x-axis: red fluorescence in donut-shaped ROIs (n = 113 cells, N=1

animal; 2 mW stimulation power). Example cells are highlighted (colored markers, letters). **(E)** Population data: same
as D for N=3 animals (N=244 neurons). Two laser intensities, 1 mW (brown), 2 mW (red). Heavy lines: LOWESS fits;
shaded regions: bootstrapped standard error. Slight decline at high values may be due to response saturation or
overexpression of opsin in a few cells. Inset: Zoomed view of area indicated by dashed box, cells with the least opsin
expression show a slightly smaller response on average than other cells. See also Fig. S6.

Input from the recurrent network dominates responses, as explained by a balanced-state model

Thus far, a moderately- or strongly-coupled balanced-state network seems consistent with both
the response distributions and dynamics we observe. Indeed, recent theoretical work in rate-
based models[22] has shown that this kind of heterogeneous network response ("reshuffling")
occurs in strongly-coupled cortical networks. To understand if our experimental data could be
explained by this reshuffling mechanism, we examined recurrent network models with features
reflecting our data, and determined which features of the recurrent network models were
important to explain the suppression.

Figure 7: Strongly-coupled recurrent neural network model with heterogeneous connectivity describes the data. (A,B) Simulation design: (A) conductance-based spiking network model with 8000 excitatory cells and 2000 inhibitory cells. (B) Network mean recurrent strength is varied to measure effects on neural responses. **(C)** Optogenetic input strengths sampled from a lognormal distribution fit from *in vivo* 2p measurements (Fig. 6C). **(D)** Schematic of data features simulations describe. **(E)** The tightly-balanced model fits the long tail of excitation and the proportion of suppressed neurons. **(F-H)** Responses to stimulation during (F) electrophysiology, (G) 2-photon experiments**,** and (H) simulations. **(I)** Left, Mean timecourse, elevated cells, strongest recurrent network. Right, same: but for suppressed cells. **(J)** Balance index (Ahmadian and Miller, 2021) of the 3 networks (B). The strongest-coupled network (purple) has a median index in the tight balance regime. **(K)** Schematic of types of input variability. Variation in input can arise from variation across cells in number of recurrent inputs, strength of recurrent inputs, or optogenetic input strength. **(L)** Simulated neural responses to optogenetic input, with (left, same as Fig. 7F) and without (center) variability in number of recurrent inputs. Relationship between optogenetic input and response strengthens when variance sources are removed (R^2 original = 0.50, R^2 reduced conn. Num. var. = 0.66; R^2 reduced conn num and reduced strength var. = 0.77) **(M)** Steady-state firing rate distributions when input variability components are removed. Purple: network with parameters as in panels E (tight),H,I,L. Right: Same data, zoom to the suppressed portion of the distribution. Some suppression exists if either source of variance is removed, but

suppression nearly abolished when both sources of variance are removed. **(N)** Finally, reducing variance in synaptic
weights (by a factor of 10) nearly removes response variability and suppression. See also Figs. S7-8.

We simulated conductance-based spiking neural networks, varying network connectivity and
opsin drive across neurons in these models, and measured network responses to excitatory cell
stimulation (Fig. 7A,B).

Each simulation consisted of two sparsely connected populations of conductance-based spiking
neurons, one excitatory (80%) and one inhibitory population (20%). For each set of network
parameters, we adjusted a background input current to either excitatory or inhibitory neurons to
hold the spontaneous firing rate of the neurons at a value (~5.4 spk/s) consistent with the data
(Fig. S8F). We drew the opsin input strength for each neuron from a distribution fit to the
imaging data, and scaled that distribution until the 75[th] percentile of the network response
matched the electrophysiology data (lognormal distribution, with 16.8% nonexpressing, Fig. 7C).
We also ran these simulations using the percentage of nonexpressing neurons as estimated
from the histology data (41%) and found no qualitative differences (Fig. S8I-M).

We first manipulated the mean connectivity strength of recurrent connections. We constructed
three different models, varying the average strength of recurrent coupling in each (schematic,
Fig. 7B). The "tightly balanced" network had the strongest recurrent coupling, and we scaled
down the synaptic weights by a factor of 2 or 4 to create more weakly-coupled "50%" and "25%
strength" network simulations. We confirmed that each of these simulations showed paradoxical
suppression of inhibitory cells, a sign of strong recurrent coupling within the excitatory network
and the ISN regime, as observed in visual cortex[11,13,34]. We stimulated the inhibitory cells in
each network and found, as expected, paradoxical suppression (Fig. S7A,B).

Recurrent excitatory-inhibitory networks can be tightly or loosely balanced[10], depending on the
total amount of recurrent excitatory and inhibitory input to network neurons. To classify the
networks, we calculated the balance index, a ratio that measures how completely inhibitory
input cancels out the excitatory input for each neuron in the networks[10] (see Methods). We
found that all three networks we constructed are balanced, as expected due to their irregular
spontaneous activity (balance index << 1), and the networks span a range from loose to tight
balance (Fig. 7J).

The model replicates the long tail of positive responses, suppressed responses, and dynamics

Two characteristic features of the response data we observed are the long-tailed positive
response and the substantial proportion of suppressed cells (Fig. 7D). All three simulated
networks showed a long tail of elevated responses as in the data, with many neurons showing
increases in firing rate to stimulation, and a few showing large increases (Fig. 7H). However, the
amount of suppression depended on recurrent coupling strength. Increasing the total excitatory
and inhibitory recurrent input by varying the mean coupling strength leads to more suppressed
neurons when other network parameters are held constant (Fig. 7E). The network that best fit
the fraction of suppressed cells we observed was the most strongly-coupled network, which was
just inside the tight balance regime (Fig. 7E,J; suppression sensitivity to baseline rate and

coupling strength characterized in Fig. S8F-H). Additional model components could lead to
similar results with networks of higher or lower balance index estimates. For instance, adding
structured connectivity may reduce the required coupling strength[22]. However, our data
underline that the recurrent coupling should be strong enough so that when input arrives to a
population of neurons, many neurons' responses are substantially controlled by their recurrent
input.

Excitatory and inhibitory cells' response distributions were similar in the model, as also seen as
in the data (Fig. S7E,F).

Finally, to further demonstrate how well the model could reproduce the features of the observed
response distributions, we simulated responses while parametrically increasing the strength of
the input, and compared the results to the electrophysiological and 2-photon responses to
increasingly strong experimental optogenetic stimulation. The shapes of the response
distributions in the most tightly coupled model and the data were similar (Fig. 7F–H).

Given the ability of a balanced-state model to describe the suppression, we checked if the
model dynamics were consistent with the data. We found that model responses were
qualitatively similar (Fig. 7I) to the timecourses of responses seen in the data (Figs. 2–3).
Excitatory cells first showed a brief, positive transient response before the network settled into a
new steady state, with some cells excited and some suppressed. The initial positive transient in
suppressed cells is a key observation, as it suggests a network mechanism where input initially
excites many excitatory neurons, but later recurrent inputs lead to suppression in many of the
same neurons. A second similar feature of the dynamics in model and data is the offset
dynamics in both elevated and suppressed cells: after stimulation ends, both show a slight
suppression before returning to baseline. Finally, excitatory cells have earlier onset times,
indicating that E cells were directly stimulated and I cells were recruited just a few milliseconds
later (Fig. S7D), before both populations then evolved to a new steady state.

One feature of the dynamics seen in the data but not the model is that for high stimulation
powers there is a slight decay during the tonic or steady-state period (Fig. 3I). However, this
decay effect is not seen at lower stimulation intensities, suggesting it arises from known opsin
dynamics (inactivation at high light power, e.g.[35]) or other known biophysical, non-network,
effects like spike-rate adaptation.

In sum, this model recapitulates many of the features of our observations, suggesting that a
excitatory-inhibitory mechanism with strong and variable recurrent coupling explains how V1
neurons respond to input.

Variability in recurrent input creates different responses in different cells, and

explains the decoupling of a neuron's response from its optogenetic input

In our data, we saw strikingly little correlation between opsin expression and neural response to
stimulation (Fig. 6), suggesting recurrent input strongly governs the response. In fact, the tightly-
coupled model showed the same pattern of responses (Fig. 7L, left).

We therefore asked which sources of variability were important to explain why neural responses
were weakly related to optogenetic input. To do this we varied sources of input variability in the
model (Fig. 7K). First, we reduced variability in either the number or strength of recurrent inputs
and found that this created a stronger correlation between optogenetic input strength and
response (Fig. 7L, middle and right) which made the model a worse fit for the data (Fig. 6). The
original relationship was not recovered by increasing the recurrent strength of the network (Fig.
7 Supp 3), implying recurrent connection variability was required to produce this effect and
higher recurrent strength could not substitute for it. Next, we asked whether variability in opsin
input across cells was also essential to explain the suppression we observed. Removing the
variance in optogenetic input across cells (so that each excitatory cell with opsin received the
same input) significantly reduced the number of suppressed neurons and also produced a
worse fit to the data (Fig. 7M; right inset highlights suppressed neurons).

If input variability was the primary source of variability in neural responses, then removing
variability from both kinds of input — both optogenetic input variability and recurrent input
variability — should substantially reduce the amount of suppression observed. This is what we
found. Removing or reducing both types of variability produced a set of neural responses
clustered tightly around the mean response (Fig. 7N) with no suppressed neurons. Thus, both
variability in recurrent input and in optogenetic input are required to explain the data.

Figure 8: Single cell stimulation produces elevated firing rates in a small subset of cells, but widespread weak suppression across the population. (A) Simulation schematic: one cell stimulated ('tight' network, Fig. 7B,E,H). **(B)** Single cell stimulation weakly modulates other cells. **(C)** Single cell stimulation reshuffles the distribution of responses; individual neurons change response (black, note variance of distribution), mean/median remain near zero. **(D)** Densities, same data as (C). **(E)** Defining cells by their connectivity to/from the stimulated cell (direct input from stimulated cell, FF, brown; input from an inhibitory cell receiving FF input, blue) reveals a small number of excited cells. N=1 instantiation of network (weight choice). **(F)** Means of E across many instantiations. Black: full population of E cells, Brown, blue: same conventions as E. Error bars: SEM. **(G)** Simulated visual input during stimulation leads to mean suppression. Red: lognormally distributed input + single cell stimulation, black: single cell stimulation alone. Red mean is negative. **(H)** Mean suppression increases with stronger input.

Together these results show that both optogenetic input variability and recurrent connection
variability help create the variability in different neurons' responses. Each neuron's firing rate is
affected not just by the optogenetic input that particular cell receives, but also by recurrent input
received from other neurons, and the other neurons themselves receive different amounts of
optogenetic and recurrent input. When optogenetic input is delivered, the whole network
changes state to a new set of firing rates, and each neuron's new firing rates are only weakly

related to the optogenetic input to that neuron. Thus, the recurrent network explains the
unexpected decoupling of optogenetic input strength from neural response strength that we
observed experimentally (Fig. 6).

A balanced-state network model with connection variability also explains expected
responses to single cell stimulation

Past work has found that stimulating a single cell in visual cortex leads to mean suppression in
the surrounding population[20] Our results seem initially to contradict this finding, because our
data and simulations both find a net positive response across the population when we stimulate
many excitatory cells.

To determine if the effects of single-cell stimulation could also be explained by the balanced-
state simulation that describes our data, we performed simulations of single cell stimulation in
the same tightly-coupled network (Fig. 7A,B), measuring the response of the non-stimulated
population (Fig 8A,B). We found that, while single cell stimulation produced a range of individual
cell responses (i.e. reshuffling, Fig. 8C), the mean response was not negative, but instead close
to zero (Fig. 8C,D; mean firing rate 95% CI [-0.006, 0.021], t (9998) for nonzero mean = 1.13, p
= 0.26). The excitatory cells that received a direct connection from the stimulated cell
(feedforward, FF, cells, n=107 neurons) had an elevated response. Those that received a
connection from the inhibitory cells which received a monosynaptic input from the directly
connected E cells had a very slightly suppressed response (i.e. E-I-E connections, or feedback
(FB) cells, n=4380 neurons; Fig. 8E). The small set of strongly excited cells average with the
large number of weakly suppressed cells to lead to a mean response near zero (Fig. 8F,G).

Single-cell stimulation in the model we fit, therefore, could not account for the mean suppression
observed in previous studies. We hypothesized that this difference could be due to difference in
the activation state of the network. Chettih and Harvey (2019) stimulated during visual input,
while here we delivered optogenetic input during spontaneous activity. Such effects can be seen
in balanced-state models: Sanzeni et al. (in press) found that increasing the firing rate of a
similar network to our model reduced the mean response of the network. Further, in models of
visual cortex with subnetwork connectivity (e.g. higher connectivity between neurons with similar
orientation tuning[36], it has also been shown that visual input can shift the network response to
be more negative[37]. Therefore, to test whether additional network drive could reproduce mean
suppressive responses, we simulated single-cell stimulation paired with an input that mimics
visual drive (Fig. 8G; Methods). Indeed, this shifted the mean population response negative
(Fig. 8G; firing rate change: mean: -0.05; 95% CI: [-0.07, -0.04]). This effect is quantitatively
dependent on the strength of the simulated visual input: as simulated visual input grows
stronger, the more negative the mean response to optogenetic input becomes (Fig. 8H).

Thus, a strongly-coupled balanced state model is consistent with not just our data, but with past
results on single-cell stimulation. Strong mean connectivity, as well as variability in recurrent
connectivity, shape the responses of the network.

Discussion

We found robust suppression in visual cortex in response to direct optogenetic drive to excitatory neurons, with intermixed elevated and suppressed neurons. This salt-and-pepper distribution of responses resembles what is observed during visual input, and arises without input to inhibitory neurons. The firing rate distributions and response dynamics suggest a network mechanism for the observed suppression: that recurrent input variability, combined with external input variability, decouples the optogenetic input strength from the firing rate response in individual cells. This yields a weak correlation between input and response (Fig. 6E), so that a high level of opsin in a cell does not necessarily mean that cell fires strongly in response to stimulation. This recurrent network mechanism seems likely to create variability in visual responses as well (Fig. 1), because these recurrent connections are present in the cortical network for all kinds of input, and so shape responses to visual input also.

Intuitively, the network mechanism that creates the salt-and-pepper excitation and suppression is that external inputs first elevate the firing rates of excitatory cells (Fig. 5A-C), some more than others. That activation excites inhibitory cells, also some more than others. The result is the network settles into a new steady state (Fig. 5E–G) with a very broad distribution of excitatory cell firing rate changes (Figs. 6,7). Our measurements, showing a long tail of excited responses (Fig. 2I), a substantial number of suppressed cells, and response dynamics with initial transients followed by steady-state excitation and suppression (Fig. 3I,J), all confirm that recurrent inputs can explain the response patterns we see.

The salt-and-pepper pattern of responses varies gradually over space, with suppressed cells becoming a larger proportion of neurons with distance from the stimulation site (Fig. 3,4). The salt-and-pepper distribution of responses we observe is therefore overlaid on top of the global trends we observed in widefield imaging. This global suppression, in a concentric surround region similar to surround suppression during vision (e.g.[38]), is driven by direct excitatory inputs, suggesting that visual surround suppression is not inherited from other regions but also arises from recurrent interactions.

The role of inhibition and suppression in the cortex: sharpening or high-dimensional pattern modification?

In principle, one role of suppression in the cortex could be to sharpen responses to input via attenuating responses in non-driven cells. The finding of distance-dependent suppression in our widefield data (Fig. 4) implies exactly this conclusion. Pioneering work using single-cell stimulation[20] also found the same sort of suppression in non-stimulated neurons. They showed that suppression falls off with distance by averaging across recorded neurons. (Note that this is true across tuning properties: while like-tuned cells in Chettih and Harvey (2019) show less suppression than other neurons, the average effect in like-tuned cells is still suppression.)

Our data and model extend this to show that sharpening is not the only, or likely even the primary, effect of cortical suppression. Using individual cells' responses with 2p imaging and electrophysiology combined with simulations, we demonstrate that cortical stimulation generates large response variability even in cells directly receiving input. That statement has significant

593   consequences for how the cortex transforms its input — it is not just that recurrent connectivity
594   sharpens responses, but it can create much more complex and high-dimensional
595   transformations[21]. Such transformations are central to neural coding and how neural codes are
596   created from input.

Variability in recurrent connectivity in the cortex: experimental evidence

We find that variability in connection strength between L2/3 excitatory neurons is necessary to
create the heterogeneous responses to input we, and others[39], observe. Several observations
suggest that the brain has recurrent variability at least as large, and possibly larger, than we use
in the simulations. First, electrophysiological studies often find a long tail of synaptic strengths
between pairs of neurons, with a few very large connections[40–42]. The variance of individual
synaptic weights may be lower[43], with the larger connection strengths due to multiple synaptic
contacts between neurons (though see[44] for evidence of long-tailed synaptic bouton sizes.) If
there is a long tail in synaptic connection strengths, this would still support our finding of high
recurrent variance, as it would increase the recurrent variability even beyond the weight
distributions we used, which are truncated Gaussians with mean and variance equal. Second,
we used a connection sparsity of 2%. We set the number of inputs a cell received from the
recurrent network according to a binomial sum, with fixed connection probability between
neurons. Connection probability in the brain may be higher, as for example paired recording
studies have found connection probabilities of 10% or higher[41,42]. And higher connection
probability will produce greater variance in net input into different cells, as binomially-distributed
sums have a larger variance as connection probability increases (in the 0–50% range). Finally,
patterned or subnetwork-specific connections, which we did not include, would also only
increase variance, though specific connections seem to have just a moderate effect on
connection probability — shared tuning changes the connection probability from 10–20% on
average to 30–50% for like-tuned neurons, in some cases[36]. Taken together, the substantial
recurrent variability that explains our data is consistent with experimental measurements of
recurrent connection variability.

Strong balance, loose balance: implications for models that describe cortical networks

We find that a two-population excitatory-inhibitory model is sufficient to explain the data we
observe. *A priori*, it could have been that a model with multiple inhibitory subtypes[15,45] would be
needed to reproduce the dynamics and population statistics we saw. Recent work has argued
for particular roles for cortical inhibitory subtypes: that parvalbumin-positive (PV) neurons are
the primary class providing inhibition stabilization[13,46], while somatostatin-positive (SOM) cells
are involved in gain control[46]. These separate roles are still consistent with our findings. PV cells
are likely to be the primary inhibitory cell class in our data and model, as PV cells are the
narrow-waveform cells that we identify in electrophysiology[13] (see Fig. 5). Those cells show
dynamic and response firing rate changes expected for the inhibitory population in an E-I model
(slightly delayed onset latency, similar distribution of firing rate change as E cells). It is also
plausible that stimulating cortical excitatory cells as we did does not cause gain to vary, so that
a separate gain role of SOM neurons was not evident in our experiments. Thus, a two-
population inhibitory model (with PV cells likely making up a large part of the I population in the
model) is sufficient to explain our data.

In addition to supporting the idea that recurrent connections between neurons have substantial
variability, our results also confirm that the mean V1 recurrent connectivity is strong — i.e. V1
operates as an inhibitory-stabilized network, meaning that the excitatory network is unstable if
inhibition could be frozen[11,13,14,22]. Within the class of balanced networks, two sorts of balance
have been distinguished: "loose" and "tight" balance[10]. The best network in our results (Fig. 7) is
on the border of the tight- and loose-balance regimes, with individual cells falling in either the
tight or loose-balance regimes. A network near the transition from loose to tight balance is
broadly consistent with past experimental data ([22], reviewed in ref. [10]) which do not suggest a
very tightly-balanced regime for the cortex (Fig. 7). Recent work has shown that adding
structured (tuned subclass) connectivity allows substantial recurrent effects with looser
balance[22], further supporting the idea that our data support loose or moderate balance.

The mechanism we find for suppression is strikingly different than paradoxical suppression in an
ISN when inhibitory cells are stimulated[12–14]. In both cases, suppression is paradoxical: here we
excite excitatory cells and see suppression of excitatory cells, and in an ISN, exciting inhibitory
cells causes suppression in inhibitory cells. But in paradoxical inhibitory suppression, the *mean*
firing rate of the inhibitory population decreases[14]. Here with excitatory cell stimulation, the
mean firing rate change is non-paradoxical, as excitatory cell average rates *increase*. It is the
substantial variability or heterogeneity of recurrent connections in combination with variability of
input that causes many cells to be suppressed as others increase their firing. However, both
types of paradoxical suppression, when excitatory or inhibitory cells are stimulated, are only
present when the network operates as an ISN – that is, both effects happen in a network with
strong average recurrent coupling[13,22]. The observed paradoxical suppression of excitatory cells,
however, requires variability around that strong average recurrent coupling.

Future: subnetworks, computation, and interactions between areas

These results could be extended in a few ways. First, here we did not consider how subnetwork
connectivity between excitatory neurons in the cortex might influence the effects. Ko and
colleagues (2011) showed approximately a 2–3 fold increase in probability of connection
between V1 excitatory neurons that had similar tuning (orientation or direction) compared to
those with dissimilar tuning. Adding subnetwork connectivity would not qualitatively change our
conclusions: that suppression results from recurrent influences, and that it depends on
variability of connectivity within the network. However, future work stimulating within or across
subnetworks might change the fraction of cells suppressed, given that input patterns would drive
neuron populations with somewhat more or less connectivity with each other and the rest of the
network. Cell-specific two-photon holographic stimulation[16,47–49] seems well-placed to study how
patterned activity in one subnetwork affects activity in another subnetwork.

While local collaterals probably contribute the majority of recurrent cortical input, meaning
nearby neurons influence each other via direct synapses, it is possible that long-range, inter-
areal, connections could contribute to the experimental results we observe. Estimates of
connectivity falloff show most connections to a given neuron come from local neurons[6,7]. But in
principle, cells in other areas could form part of the recurrent population. This could happen for
example if projections from V1 to the thalamus recruited neurons there which connect back to
the cortex. However, our widefield imaging data (Fig. 4) shows that the suppression peaks a few

hundred microns from the stimulation site, suggesting relatively local influence. Therefore, it
seems likely that the recurrent connections in the simulations primarily reflect local connections
within V1 to nearby neurons.

Conclusion

Here we find paradoxical suppression of excitatory cells in the cortex when excitatory cells are
stimulated. These results suggest that a primary purpose of recurrent connectivity in visual
cortex is to change the steady-state firing rate of network neurons, beyond just how inputs are
transformed by feedforward connections. Our results are a step forward in explaining how
cortical networks change their firing in response to different patterns of input — a fundamental
building block of neuronal computation.

Acknowledgements

688 We thank Kaya Matson for help with RNAscope, Aanika Kashyap for histological analysis.and
689 Nicolas Brunel, Alessandro Sanzeni, and Ken Miller for helpful comments on the manuscript
690 and/or discussion. This work was funded by the National Institutes of Health (BRAIN
691 U01NS108683 and intramural support ZIAMH002956). This work utilized the computational
692 resources of the NIH HPC Biowulf cluster (http://hpc.nih.gov).

Author contributions

694 Electrophysiology data was collected by ZZ. 2-photon data was collected by PKL, AJL, and
695 JFO. Widefield data was collected by JFO. Histological data collection and analysis was
696 performed by ZZ and HCG. JFO and MHH designed and implemented simulations. JFO, MHH,
697 and ZZ curated and analyzed data. JFO and MHH wrote and edited the manuscript.

Declaration of interests

699 The authors declare no competing interests.

700 # STAR Methods

701 ## *Key resources table*

Reagent or resource	Source	Identifier	Additional information
Chemicals, peptides, and recombinant proteins			
Tamoxifen	Sigma-Aldrich	T5648-5G	
Experimental models: organisms/strains			
Ai148	The Jackson Laboratory	RRID:IMSR_JAX:030328	
Ai162	The Jackson Laboratory	RRID:IMSR_JAX:022731	
Cux2-CreERT2	MMRRC	RRID:MMRRC_032779-MU	
Emx1-cre	The Jackson Laboratory	RRID:IMSR_JAX:005628	
Bacterial and virus strains			
AAV9-hSyn-jGCaMP7s-WPRE	Addgene	RRID:Addgene_104487	
AAV9-Syn-DIO-stChrimsonR-mRuby	Addgene	RRID:Addgene_105448	
AAV9-CamKIIa-ChrimsonR-mScarlet-KV2.1	Addgene	RRID:Addgene_124651	
Software and algorithms			
Mworks	The Mworks Project		mworks.github.io
Other			
C and B Metabond	Parkell	S380	
Kwik-sil	World Precision Instruments	KWIK-SIL	

Resource Availability

- *Lead Contact*
Additional information and requests for resources should be directed to the lead contact,
Mark Histed (mark.histed@nih.gov)
- *Materials availability*
This work did not produce novel reagents.
- *Data and code availability*
Data and code will be published in a GitHub repository or on DANDI on acceptance for
publication.

Experimental Model and Subject Details

All procedures were approved by the NIMH Institutional Animal Care and Use Committee (IACUC) and conform to relevant regulatory standards. Emx1-cre animals[30] of both sexes (N = 14; https://www.jax.org/strain/005628) were used for 2-photon and electrophysiology experiments (5 for electrophysiology, 3 for visual stimulation imaging, 6 for optogenetic stimulation imaging). For widefield imaging experiments, Ai162 (N = 2; https://www.jax.org/strain/031562) and Ai148 (N = 1; https://www.jax.org/strain/030328) animals[50] were crossed with the Cux2-CreERT2 line[51] (https://www.mmrrc.org/catalog/sds.php?mmrrc_id=32779), and GCaMP6f or GCaMP6s was induced via tamoxifen injection during adulthood (P22 or later, tamoxifen 2 mg intraperitoneally daily for 3 days). All animals were singly housed on a reversed light cycle. During experiments animals were water scheduled and given occasional water rewards to keep them awake and alert. To ensure animals did not drift into a quiet wakefulness or quiescent state, we monitored animals during data collection to verify they continued to drink the delivered reward.

Methods Details

Implants and injections

Details of the headplate and window procedures are described in previous studies[13,52]. Optical glass windows (3 mm diameter) were placed over V1 (center: -3 mm ML, +1.5 mm AP, relative to lambda) for 2p and widefield imaging. Windows were also used before electrophysiology for imaging to localize V1.

For Emx1-Cre animals, 300 nL of AAV9-syn-jGCaMP7s-WPRE (RRID:Addgene_104487) and/or AAV9-Syn-DIO-stChrimsonR-mRuby (RRID:Addgene_105448) were injected 250 μm below the dura (200 nL/min) prior to cementing the cranial window. For Ai148 and 162 animals, AAV9-CamKIIa-stChrimsonR-mRuby2 was generated by cloning the CaMKIIa promoter (RRID:Addgene_120219) into a pAAV backbone containing stChrimsonR-mRuby2 (RRID:Addgene_105447) and packaged into an AAV (Vigene, Inc.). This was injected at the same depth as the hSyn-DIO-stChrimsonR virus, but with 100 nL volume at 100 nL/min.

Electrophysiology

Electrophysiological methods are described in detail in previous studies[13], and are summarized here. Animals were head-fixed during recording. Before the first session of electrophysiology, the animal's cranial window was removed and the craniotomy was flushed with saline. Between recording sessions, the craniotomy was covered using Kwik-Sil polymer (WPI, Inc.). A fiber optic cannula (400 μm diameter, 0.39 NA, Thorlabs) was placed to center light output at the center of stChrimsonR expression. For light intensity calculations, spot area was defined as the area inside the 50% contour of light spot intensity on the cortex, measured with a camera by imaging the spot on the brain surface. 1–2% agarose (Type IIIA, Sigma) was placed over the dura at the start of each session, and an array of four electrodes (4 probes, 32 sites in total, part #A4x8-5mm-100-400-177-A32, NeuroNexus, Inc.) were lowered into the cortex using a micromanipulator (Sutter MPC-200). One probe was placed at the center of the light spot.

Probes were advanced 600–1000 μm below the point in which the first probe touched the dura.
Probes were not moved for 1 hour prior to recording, as we found this improved recording
stability. Recording data was sampled at 30 kHz (Cerebus, Blackrock Microsystems.)

Optogenetic stimulation was performed with randomly interleaved stimulation light pulses with
several intensities over the range 0.2 mW/mm^2 to 15 mW/mm^2. Stimulation pulses were 600 ms
long and delivered with a 4 s period.
757
For spike recordings, waveforms (bandpass filtered, 750 Hz – 7.5 kHz) were digitized and saved
by storing a short data section around points where amplitude exceeded 3 times the RMS noise
on that channel. Single units were identified (OfflineSorter, Plexon, Inc) based on clusters in the
waveform PCA that were separate from noise and other clusters, had unimodal spike width
distributions, and inter-spike intervals consistent with cortical neuron absolute and relative
refractory periods. A single-unit score was assigned to each unit manually based on these
factors [13,23]. To compare these populations quantitatively, we calculated SNR for both single and
multiunits[53,54]. Median SNR for single units was larger than median SNR for multiunits (SU:
3.32, MU: 2.26; Fig. S2C), consistent with prior reports[13,53,55].

Histology
Following completion of electrophysiology experiments, mice were anesthetized with isoflurane
and injected intraperitoneally with pentobarbital sodium (150 mg/kg), and perfused transcardially
with cold (4°C) PBS followed by cold 4% paraformaldehyde. Brains were extracted and fixed in
4% paraformaldehyde for 6–12 hr and then cryoprotected in a 30% (% w/v) sucrose solution in
PBS until they sank. Tissue was cryosectioned at 10μm and mounted on charged slides.
Fluorescence in situ hybridization was done using RNAscope's Multiplex Fluorescent Assay
v2[56]. Inhibitory neurons were labeled with a VGAT probe (Slc32a1, #319191-C2; Alexa Fluor
488), excitatory neurons were labeled with a VGLUT1 probe (Slc17a7, #416631-C3; Cy5), and
an mRuby2 probe (#487361; Cy3) labeled stChrimsonR-mRuby2 expressing neurons. Slides
were coverslipped with DAPI. We imaged slides on a Zeiss LSM780 confocal microscope with a
40x oil immersion objective. We imaged each fluorophore separately with a single excitation
laser, and collected all three emission channels. To compensate for bleedthrough where the
other two fluorophores might be weakly excited by a laser selected for another fluorophore, we
subtracted a scaled version of the primary emission channel image for each non-selected
fluorophore from the primary channel for the selected fluorophore. Five representative areas
were quantified independently by two observers.

Visual stimulation
Visual stimuli were presented using MWorks (https://mworks.github.io/). Grating stimuli
(sinusoidal contrast variation, 0.1 cyc/deg, orientation = 0 deg) were masked with a circular
raised-cosine envelope (15 deg FWHM). Visual stimuli were displayed on an LCD display, with
center positioned 0-10 degrees of visual angle temporal to the central meridian. Oriented noise
stimuli were generated by filtering white noise pixel arrays (each pixel drawn independently from
a uniform distribution) with a spatial band-pass filter (peak orientation = 0 deg, orientation
bandwidth = 10 deg, peak spatial frequency = 0.05, frequency bandwidth = 0.05). Frames were

generated at 60 Hz and the noise pattern was independent from frame to frame[26–28]. Visual
stimuli were presented for 3 or 5 seconds, depending on the experiment.

2-photon imaging
During 2-photon experiments, animals were awake, water-scheduled, and given periodic water
rewards (20% probability per trial, reward once every 30 s on average). If animals stopped
licking in response to the rewards, data collection was ended. We imaged GCaMP7s responses
(920 nm excitation) with either a galvo-galvo (5 Hz) or resonant scanning (30 Hz) two-photon
microscope. stChrimsonR-mRuby2 expression was imaged at 1000 nm. The microscopes used
for imaging were built using MIMMS components (https://www.janelia.org/open-science/mimms-
21-2020) and other custom components, built in-house or provided by Sutter Inc. A second light
path, combined with the 2p stimulation light path before the tube lens using a dichroic, was used
to stimulate stChrimsonR using 530nm light (CoolLED, pE-4000). For 200 ms long optogenetic
pulses, we measured responses in the first frame after stimulation. For 4 s long optogenetic light
pulses (6 s period), we imaged while stimulation was ongoing. To do this, we avoided
stimulation artifacts by stimulating only during horizontal flyback (approximate pulse duration 19
µs, off time 44 µs, duty cycle 30%, line rate 8kHz).

Widefield imaging
For widefield imaging experiments, we used Ai162;Cux2-creERT2 or Ai148;Cux2-creERT2
animals, expressing GCaMP6f or 6s in L2/3 excitatory cells. Animals were head-fixed and
awake during widefield imaging experiments. Prior to imaging, a fiber optic cannula was aimed
at the center of the focal stChrimsonR expression. Images were collected using a Zeiss
microscope (Discovery V12) with a 1.0x objective using excitation light with wavelength
centered at 475 nm (Xylis X-Cite XT720L). A Zyla 4.2 sCMOS camera (Oxford Instruments)
collected images (100 ms exposure time, approximately 140 ms frame period) with 4-pixel
binning. Laser powers were randomly interleaved, with 50 repetitions per laser power. Laser
pulses were 600 ms long, and presented with 6 s period.

Analysis of electrophysiology data
For spike rate plots, spike counts were binned (1 ms bins), and smoothed via LOWESS[57]. To
classify units as having elevated or suppressed responses, we measured spike rate over 145–
400 ms after stim onset, relative to baseline (-1020 ms–0 ms relative to stim onset) for 6
mW/mm^2 stimulation intensity. To classify cells as wide- or narrow-waveform, we used a spike
width threshold of 0.445 ms based on the bimodal distribution of waveform widths (Fig. 5B).
This threshold is consistent with pharmacological segregation of inhibitory and excitatory cells[13].
825
826   For analysis of onset times, we fit a sigmoid (logistic function) to each cell's response from 100
827   ms before to 100ms after laser pulse onset:

828
$$f(x) = \frac{L}{1 + e^{-k(x-x_0)}} + b$$

L: upper asymptote, b: lower asymptote, k: slope, x_0: onset latency. X_0 was constrained to the
range [onset+0.5 ms, onset + 30 ms]. We defined onset latency as x_0, the time to half-max. To
estimate the time to steady-state, the same function was fit to data from 500 ms before and after
the laser onset, with the spike rates within a 50 ms window around the initial transient blanked
by setting to the baseline firing rate. Each cell's time to steady-state was computed as the
difference between the steady-state onset and the initial onset (difference between the x_0
parameters of the two fits).

Analysis of 2-photon data
For short optogenetic stimulation (200 ms pulses) during two-photon imaging, we avoided
stimulation light influencing imaging responses by measuring responses in the frame after the
stimulus offset. For long pulses (4 s), we stimulated during imaging frames by restricting
stimulation to imaging line flyback and intensities we give are the average intensity, corrected
for the 30% stimulation duty cycle. Because we found that the LED device we used for
stimulation (pE-4000, CoolLED Ltd; specified bandwidth 100 kHz) had some variability in
onset/offset for each line, we removed pixels (~40% of frame) at left and right edges of field of
view to ensure no stimulation light could affect images. Image frames were motion corrected
using NoRMCorre through CaImAn[58]. Deconvolution was done with OASIS[59] via CaImAn. To
ease interpretability of the deconvolution signals, each neuron's deconvolved signal was
normalized to have the same maximum value as the dF/F of the corresponding fluorescence
trace. To separate populations into elevated and suppressed cells, we performed a one-sample
t-test ($\alpha = 0.05$, different from zero, two-tailed) on the deconvolved dF/F during the stimulation
period (long pulses) or the frame just after the stimulation period (short pulses). For the short
pulses, we used the frame just after stimulation to estimate responses for each neuron per trial.
For the long pulses, we averaged data within the period 750 ms after stimulus onset to the
stimulus offset in order to capture the steady-state response. For visual response data, data
were preprocessed in the same manner as the short optogenetic stimulation experiments. We
averaged steady-state responses from 750 ms after stimulus onset to stimulus offset.

For spatial analyses, we used the spatstat package[60] in R (ver. 4.2.3). For each individual
animal, we tested the spatial distributions of elevated and suppressed responses against an
inhomogeneous Poisson process model using the *Linhom* and *Lcross.inhom* functions. We
used an inhomogeneous process as signal properties (e.g. slight tilt of imaging field) and
biological properties (e.g. vasculature) may produce inhomogeneities in rate/intensity that could
be mistaken for clustering. The L function estimates the expected number of discovered
neurons for different diameter circular search areas centered on each neuron, given the
modeled Poisson process. We corrected for windowing in the selected field-of-view using
Ripley's isotropic correction. Global envelopes were generated (using the *envelope* function)
with p < 0.05, Bonferroni-corrected.

For 2-photon opsin measurements, for each field of view we corrected for neuropil signal by
manually selecting a region of neuropil with no visible cell bodies/processes and subtracting that
intensity. We measured red fluorescence in donut-shaped regions of interest around the border
of each cell mask. Each animal's distribution of opsin was normalized by dividing by their

maximum opsin fluorescence, and then combined. We fit a lognormal distribution via least-
squares (details in Fig. S6).

Analysis of widefield imaging data

Widefield fluorescence images were motion corrected for rigid translation, and any linear trend
across the full imaging session was estimated via regression and subtracted. Deconvolution
was done via Widefield Deconvolution[61], which differs from single-neuron deconvolution
algorithms like OASIS by dropping the sparsity assumption useful for spike trains of single
neurons. This algorithm produces better results for aggregated signals, such as that from a
single pixel during widefield imaging[61]. We rescaled the deconvolved signals to the maximum
dF/F of the imaging data, as with the two-photon deconvolution. Comparison of Widefield
Deconvolution, OASIS, and first-differencing is given in Fig. S3. For timecourse analyses, center
and surround ROIs were defined as as the top 30% of elevated or suppressed pixels within a 1
mm radius of the center of response. To average images across animals, images were aligned
on the basis of their maximum response during the late stimulation period. For quantification of
spatial falloff (Fig 4G–I), we found the peak, averaged the responses radially, and then fit a
curve to the responses vs. distance (LOWESS; 95% CI via bootstrap). Crossing points are the
minimum distance at which the 95% confidence interval contains zero.

Spiking network model

We simulated a conductance-based neural network model with 10000 neurons (8000 excitatory,
2000 inhibitory) to understand the recurrent features that contribute to the response properties
we observe during excitatory cell stimulation. Simulations were performed using Brian2[62].

Membrane and synaptic dynamics evolve according to the following equations:

$$(1)\ C\frac{dV}{dt} = g_L(E_L - V) + g_E(E_E - V) + g_I(E_I - V) + I_{background} + c \cdot I_{opto}(t)$$

$$(2)\ \frac{dg_E}{dt} = -g_E/\tau_E$$

$$(3)\ \frac{dg_I}{dt} = -g_I/\tau_I$$

Each synapse was stepped by its corresponding connection weight for every presynaptic spike.

Connections between neurons were made with 2% probability, independently for each potential
connection[36,41,63].

Parameter	Value		Parameter	Value
τ_E	5 ms		Mean W_{II}	4.0 nS
τ_I	10 ms		Variance W_{II}	4.0 nS
E_L	-60 mV		Mean W_{IE}	5.0 nS
E_I	-80 mV		Variance W_{IE}	5.0 nS

E_E	0 mV		E cell $I_{background}$ Full Network	260 pA
g_L	10.0 nS		I cell $I_{background}$ Full Network	140 pA
Mean W_{EE}	0.4 nS		E cell $I_{background}$ 50% Network	227.5 pA
Variance W_{EE}	0.4 nS		I cell $I_{background}$ 50% Network	172.5 pA
Mean W_{EI}	0.8 nS		E cell $I_{background}$ 25% Network	208.5 pA
Variance W_{EI}	0.8 nS		I cell $I_{background}$ 25% Network	191.5 pA

Table 1: Spiking neural network model parameters

Synaptic weights were drawn from truncated (rectified) Gaussian distributions. Mean
connectivity parameters were based on published measurements, with excitatory connection
strength an order of magnitude weaker than inhibitory connection strength[36,64,65] and I-to-E
connectivity stronger than I-to-I connectivity[65]. Background currents were chosen for inhibitory
and excitatory cell populations to fix baseline firing-rates for each constructed network to data
(Fig S8F). Network parameters shown in Table 1.

Optogenetic stimulation simulations

Optogenetic stimulation was an additional constant current for the length of the stimulation
period, with onset and offset ramped linearly over 3 ms. The strength of the optogenetic
stimulation (c in Eq. 1 was chosen from a lognormal distribution derived from data (Fig. 6), or
held constant (Fig. 7M,N). For each simulation, this stimulation distribution was scaled by a
constant to reproduce the response rate from data, at the 75^{th} percentile across excitatory cells.
Steady-state response was measured for each cell as their firing rate during the 1 s baseline
period subtracted from the firing rate during the last 500 ms of the stimulation period. To reduce
connection strength variability (Fig. 7N), we reduced the variability of the truncated Gaussians
that define connection strength by a factor of 100 (setting both the synaptic strength variability
and connection number variability to zero produced a network that was less stable).

Single cell stimulation simulations

A single cell was stimulated with intensity from maximum of input distribution (Fig. 7H). Controls:
same parameters but no stimulation. To simulate single cell stimulation with visual input, we
provided single cell stimulation during either the lognormal optogenetic stimulation, as
previously described, or during uniform input of both excitatory and inhibitory cells.

Balance index

We computed the balance index as described by Ahmadian and Miller (2021). For each neuron,
we computed this index as the net current (excitatory + inhibitory) divided by the excitatory
current. The index becomes smaller as balance becomes tighter, with component currents
becoming larger, and the index becomes larger as inhibitory input from the network shrinks.

Quantification and Statistical Analysis

All analyses, unless specifically noted in Methods Details, were performed in python using

NumPy and SciPy packages[57,66]. Degrees of freedom and statistical tests are described in the

results text. Error metrics plotted in figures are listed in the figure legends. Significance was

adjusted for multiple testing using a Bonferroni correction when appropriate.

References

1. Van Essen, D.C., Anderson, C.H., and Felleman, D.J. (1992). Information Processing in the Primate Visual System: An Integrated Systems Perspective. Science *255*, 419–423. 10.1126/science.1734518.

2. Glickfeld, L.L., and Olsen, S.R. (2017). Higher-Order Areas of the Mouse Visual Cortex. Annu. Rev. Vis. Sci. *3*, 251–273. 10.1146/annurev-vision-102016-061331.

3. Kravitz, D.J., Saleem, K.S., Baker, C.I., Ungerleider, L.G., and Mishkin, M. (2013). The ventral visual pathway: an expanded neural framework for the processing of object quality. Trends Cogn. Sci. *17*, 26–49. 10.1016/j.tics.2012.10.011.

4. Van Essen, D.C., and Gallant, J.L. (1994). Neural mechanisms of form and motion processing in the primate visual system. Neuron *13*, 1–10. 10.1016/0896-6273(94)90455-3.

5. Krizhevsky, A., Sutskever, I., and Hinton, G.E. (2012). ImageNet Classification with Deep Convolutional Neural Networks. In Advances in Neural Information Processing Systems (Curran Associates, Inc.).

6. Binzegger, T., Douglas, R.J., and Martin, K.A.C. (2004). A Quantitative Map of the Circuit of Cat Primary Visual Cortex. J. Neurosci. *24*, 8441–8453. 10.1523/JNEUROSCI.1400-04.2004.

7. Hellwig, B. (2000). A quantitative analysis of the local connectivity between pyramidal neurons in layers 2/3 of the rat visual cortex. Biol. Cybern. *82*, 111–121. 10.1007/PL00007964.

8. Rossi, L.F., Harris, K.D., and Carandini, M. (2019). Excitatory and inhibitory intracortical circuits for orientation and direction selectivity. 556795. 10.1101/556795.

9. Douglas, R.J., Koch, C., Mahowald, M., Martin, K.A., and Suarez, H.H. (1995). Recurrent excitation in neocortical circuits. Science *269*, 981–985. 10.1126/science.7638624.

10. Ahmadian, Y., and Miller, K.D. (2021). What is the dynamical regime of cerebral cortex? Neuron. 10.1016/j.neuron.2021.07.031.

11. Ozeki, H., Finn, I.M., Schaffer, E.S., Miller, K.D., and Ferster, D. (2009). Inhibitory Stabilization of the Cortical Network Underlies Visual Surround Suppression. Neuron *62*, 578–592. 10.1016/j.neuron.2009.03.028.

12. Sadeh, S., and Clopath, C. (2021). Inhibitory stabilization and cortical computation. Nat. Rev. Neurosci. *22*, 21–37. 10.1038/s41583-020-00390-z.

13. Sanzeni, A., Akitake, B., Goldbach, H.C., Leedy, C.E., Brunel, N., and Histed, M.H. (2020). Inhibition stabilization is a widespread property of cortical networks. eLife *9*, e54875. 10.7554/eLife.54875.

14. Tsodyks, M.V., Skaggs, W.E., Sejnowski, T.J., and McNaughton, B.L. (1997). Paradoxical Effects of External Modulation of Inhibitory Interneurons. J. Neurosci. *17*, 4382–4388. 10.1523/JNEUROSCI.17-11-04382.1997.

15. Mahrach, A., Chen, G., Li, N., van Vreeswijk, C., and Hansel, D. (2020). Mechanisms
underlying the response of mouse cortical networks to optogenetic manipulation. eLife *9*,
e49967. 10.7554/eLife.49967.

16. Marshel, J.H., Kim, Y.S., Machado, T.A., Quirin, S., Benson, B., Kadmon, J., Raja, C.,
Chibukhchyan, A., Ramakrishnan, C., Inoue, M., et al. (2019). Cortical layer–specific critical
dynamics triggering perception. Science *365*, eaaw5202. 10.1126/science.aaw5202.

17. Peron, S., Pancholi, R., Voelcker, B., Wittenbach, J.D., Ólafsdóttir, H.F., Freeman, J., and
Svoboda, K. (2020). Recurrent interactions in local cortical circuits. Nature, 1–4.
10.1038/s41586-020-2062-x.

18. Goldman, M.S. (2009). Memory without Feedback in a Neural Network. Neuron *61*, 621–
634. 10.1016/j.neuron.2008.12.012.

19. Murphy, B.K., and Miller, K.D. (2009). Balanced Amplification: A New Mechanism of
Selective Amplification of Neural Activity Patterns. Neuron *61*, 635–648.
10.1016/j.neuron.2009.02.005.

20. Chettih, S.N., and Harvey, C.D. (2019). Single-neuron perturbations reveal feature-specific
competition in V1. Nature *567*, 334–340. 10.1038/s41586-019-0997-6.

21. Oldenburg, I.A., Hendricks, W.D., Handy, G., Shamardani, K., Bounds, H.A., Doiron, B., and
Adesnik, H. (2022). The logic of recurrent circuits in the primary visual cortex.
2022.09.20.508739. 10.1101/2022.09.20.508739.

22. Sanzeni, A., Palmigiano, A., Nguyen, T.H., Luo, J., Nassi, J.J., Reynolds, J.H., Histed, M.H.,
Miller, K.D., and Brunel, N. (2022). Mechanisms underlying reshuffling of visual responses
by optogenetic stimulation in mice and monkeys. 2022.07.13.499597.
10.1101/2022.07.13.499597.

23. Histed, M.H. (2018). Feedforward Inhibition Allows Input Summation to Vary in Recurrent
Cortical Networks. eNeuro *5*, ENEURO.0356-17.2018. 10.1523/ENEURO.0356-17.2018.

24. Nassi, J.J., Avery, M.C., Cetin, A.H., Roe, A.W., and Reynolds, J.H. (2015). Optogenetic
Activation of Normalization in Alert Macaque Visual Cortex. Neuron *86*, 1504–1517.
10.1016/j.neuron.2015.05.040.

25. McGinley, M.J., Vinck, M., Reimer, J., Batista-Brito, R., Zagha, E., Cadwell, C.R., Tolias,
A.S., Cardin, J.A., and McCormick, D.A. (2015). Waking State: Rapid Variations Modulate
Neural and Behavioral Responses. Neuron *87*, 1143–1161. 10.1016/j.neuron.2015.09.012.

26. Beaudot, W.H.A., and Mullen, K.T. (2006). Orientation discrimination in human vision:
Psychophysics and modeling. Vision Res. *46*, 26–46. 10.1016/j.visres.2005.10.016.

27. Bondy, A.G., Haefner, R.M., and Cumming, B.G. (2018). Feedback determines the structure
of correlated variability in primary visual cortex. Nat. Neurosci. *21*, 598–606.
10.1038/s41593-018-0089-1.

28. Rolfs, M., and Carrasco, M. (2012). Rapid Simultaneous Enhancement of Visual Sensitivity
and Perceived Contrast during Saccade Preparation. J. Neurosci. *32*, 13744–13752a.
10.1523/JNEUROSCI.2676-12.2012.

29. Nakajima, M., Schmitt, L.I., and Halassa, M.M. (2019). Prefrontal Cortex Regulates Sensory
Filtering through a Basal Ganglia-to-Thalamus Pathway. Neuron *103*, 445-458.e10.
10.1016/j.neuron.2019.05.026.

30. Gorski, J.A., Talley, T., Qiu, M., Puelles, L., Rubenstein, J.L.R., and Jones, K.R. (2002).
Cortical Excitatory Neurons and Glia, But Not GABAergic Neurons, Are Produced in the
Emx1-Expressing Lineage. J. Neurosci. *22*, 6309–6314. 10.1523/JNEUROSCI.22-15-
06309.2002.

31. Watakabe, A., Ohtsuka, M., Kinoshita, M., Takaji, M., Isa, K., Mizukami, H., Ozawa, K., Isa,
T., and Yamamori, T. (2015). Comparative analyses of adeno-associated viral vector
serotypes 1, 2, 5, 8 and 9 in marmoset, mouse and macaque cerebral cortex. Neurosci.
Res. *93*, 144–157. 10.1016/j.neures.2014.09.002.

32. Ahmadian, Y., Rubin, D.B., and Miller, K.D. (2013). Analysis of the Stabilized Supralinear
Network. Neural Comput. *25*, 1994–2037. 10.1162/NECO_a_00472.

33. Renart, A., de la Rocha, J., Bartho, P., Hollender, L., Parga, N., Reyes, A., and Harris, K.D.
(2010). The Asynchronous State in Cortical Circuits. Science *327*, 587–590.
10.1126/science.1179850.

34. Li, N., Chen, S., Guo, Z.V., Chen, H., Huo, Y., Inagaki, H.K., Chen, G., Davis, C., Hansel,
D., Guo, C., et al. (2019). Spatiotemporal constraints on optogenetic inactivation in cortical
circuits. eLife *8*, e48622. 10.7554/eLife.48622.

35. Schneider, F., Gradmann, D., and Hegemann, P. (2013). Ion Selectivity and Competition in
Channelrhodopsins. Biophys. J. *105*, 91–100. 10.1016/j.bpj.2013.05.042.

36. Ko, H., Hofer, S.B., Pichler, B., Buchanan, K.A., Sjöström, P.J., and Mrsic-Flogel, T.D.
(2011). Functional specificity of local synaptic connections in neocortical networks. Nature
473, 87–91. 10.1038/nature09880.

37. Podlaski, W., Russell, L.E., Roth, A., Bicknell, B., Häusser, M., and Machens, C. (2022).
The dynamical regime of mouse visual cortex shifts from cooperation to competition with
increasing visual input | Cosyne 2022. https://www.world-wide.org/cosyne-22/dynamical-
regime-mouse-visual-cortex-5af75de1/.

38. Angelucci, A., and Bressloff, P.C. (2006). Contribution of feedforward, lateral and feedback
connections to the classical receptive field center and extra-classical receptive field
surround of primate V1 neurons. In Progress in Brain Research Visual Perception., S.
Martinez-Conde, S. L. Macknik, L. M. Martinez, J.-M. Alonso, and P. U. Tse, eds. (Elsevier),
pp. 93–120. 10.1016/S0079-6123(06)54005-1.

39. Pancholi, R., Sun-Yan, A., and Peron, S. (2023). Microstimulation of sensory cortex
engages natural sensory representations. Curr. Biol. *33*, 1765-1777.e5.
10.1016/j.cub.2023.03.085.

40. Arellano, J., Benavides-Piccione, R., DeFelipe, J., and Yuste, R. (2007). Ultrastructure of
1048   dendritic spines: correlation between synaptic and spine morphologies. Front. Neurosci. *1*,
1049   10. 10.3389/neuro.01.1.1.010.2007.

1050   41. Holmgren, C., Harkany, T., Svennenfors, B., and Zilberter, Y. (2003). Pyramidal cell
1051   communication within local networks in layer 2/3 of rat neocortex. J. Physiol. *551*, 139–153.
1052   10.1111/j.1469-7793.2003.00139.x.

1053   42. Song, S., Sjöström, P.J., Reigl, M., Nelson, S., and Chklovskii, D.B. (2005). Highly
1054   Nonrandom Features of Synaptic Connectivity in Local Cortical Circuits. PLOS Biol. *3*, e68.
1055   10.1371/journal.pbio.0030068.

1056   43. Chapeton, J., Fares, T., LaSota, D., and Stepanyants, A. (2012). Efficient associative
1057   memory storage in cortical circuits of inhibitory and excitatory neurons. Proc. Natl. Acad.
1058   Sci. *109*, E3614–E3622. 10.1073/pnas.1211467109.

1059   44. Loewenstein, Y., Kuras, A., and Rumpel, S. (2011). Multiplicative Dynamics Underlie the
1060   Emergence of the Log-Normal Distribution of Spine Sizes in the Neocortex In Vivo. J.
1061   Neurosci. *31*, 9481–9488. 10.1523/JNEUROSCI.6130-10.2011.

1062   45. Litwin-Kumar, A., Rosenbaum, R., and Doiron, B. (2016). Inhibitory stabilization and visual
1063   coding in cortical circuits with multiple interneuron subtypes. J. Neurophysiol. *115*, 1399–
1064   1409. 10.1152/jn.00732.2015.

1065   46. Bos, H., Oswald, A.-M., and Doiron, B. (2020). Untangling stability and gain modulation in
1066   cortical circuits with multiple interneuron classes. 2020.06.15.148114.
1067   10.1101/2020.06.15.148114.

1068   47. Dalgleish, H.W., Russell, L.E., Packer, A.M., Roth, A., Gauld, O.M., Greenstreet, F.,
1069   Thompson, E.J., and Häusser, M. (2020). How many neurons are sufficient for perception of
1070   cortical activity? eLife *9*, e58889. 10.7554/eLife.58889.

1071   48. Packer, A.M., Peterka, D.S., Hirtz, J.J., Prakash, R., Deisseroth, K., and Yuste, R. (2012).
1072   Two-photon optogenetics of dendritic spines and neural circuits. Nat. Methods *9*, 1202–
1073   1205. 10.1038/nmeth.2249.

1074   49. Sadeh, S., and Clopath, C. (2020). Theory of neuronal perturbome in cortical networks.
1075   Proc. Natl. Acad. Sci. *117*, 26966–26976. 10.1073/pnas.2004568117.

1076   50. Daigle, T.L., Madisen, L., Hage, T.A., Valley, M.T., Knoblich, U., Larsen, R.S., Takeno,
1077   M.M., Huang, L., Gu, H., Larsen, R., et al. (2018). A Suite of Transgenic Driver and Reporter
1078   Mouse Lines with Enhanced Brain-Cell-Type Targeting and Functionality. Cell *174*, 465-
1079   480.e22. 10.1016/j.cell.2018.06.035.

1080   51. Franco, S.J., Gil-Sanz, C., Martinez-Garay, I., Espinosa, A., Harkins-Perry, S.R., Ramos, C.,
1081   and Müller, U. (2012). Fate-Restricted Neural Progenitors in the Mammalian Cerebral
1082   Cortex. Science *337*, 746–749. 10.1126/science.1223616.

1083   52. Goldbach, H.C., Akitake, B., Leedy, C.E., and Histed, M.H. (2021). Performance in even a
1084   simple perceptual task depends on mouse secondary visual areas. eLife *10*, e62156.
1085   10.7554/eLife.62156.

53. Kelly, R.C., Smith, M.A., Samonds, J.M., Kohn, A., Bonds, A.B., Movshon, J.A., and Lee,
T.S. (2007). Comparison of Recordings from Microelectrode Arrays and Single Electrodes in
the Visual Cortex. J. Neurosci. *27*, 261–264. 10.1523/JNEUROSCI.4906-06.2007.

54. Nordhausen, C.T., Maynard, E.M., and Normann, R.A. (1996). Single unit recording
capabilities of a 100 microelectrode array. Brain Res. *726*, 129–140. 10.1016/0006-
8993(96)00321-6.

55. Wissig, S.C., and Kohn, A. (2012). The influence of surround suppression on adaptation
effects in primary visual cortex. J. Neurophysiol. *107*, 3370–3384. 10.1152/jn.00739.2011.

56. Wang, F., Flanagan, J., Su, N., Wang, L.-C., Bui, S., Nielson, A., Wu, X., Vo, H.-T., Ma, X.-
J., and Luo, Y. (2012). RNAscope: a novel in situ RNA analysis platform for formalin-fixed,
paraffin-embedded tissues. J. Mol. Diagn. JMD *14*, 22–29. 10.1016/j.jmoldx.2011.08.002.

57. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D.,
Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental
algorithms for scientific computing in Python. Nat. Methods *17*, 261–272. 10.1038/s41592-
019-0686-2.

58. Giovannucci, A., Friedrich, J., Gunn, P., Kalfon, J., Brown, B.L., Koay, S.A., Taxidis, J.,
Najafi, F., Gauthier, J.L., Zhou, P., et al. (2019). CaImAn an open source tool for scalable
calcium imaging data analysis. eLife *8*, e38173. 10.7554/eLife.38173.

59. Friedrich, J., Zhou, P., and Paninski, L. (2017). Fast online deconvolution of calcium
imaging data. PLOS Comput. Biol. *13*, e1005423. 10.1371/journal.pcbi.1005423.

60. Baddeley, A., Rubak, E., and Turner, R. (2015). Spatial Point Patterns: Methodology and
Applications with R (CRC Press).

61. Stern, M., Shea-Brown, E., and Witten, D. (2020). Inferring the Spiking Rate of a Population
of Neurons from Wide-Field Calcium Imaging 10.1101/2020.02.01.930040.

62. Stimberg, M., Brette, R., and Goodman, D.F. (2019). Brian 2, an intuitive and efficient neural
simulator. eLife *8*, e47314. 10.7554/eLife.47314.

63. Song, S., Sjöström, P.J., Reigl, M., Nelson, S., and Chklovskii, D.B. (2005). Highly
Nonrandom Features of Synaptic Connectivity in Local Cortical Circuits. PLOS Biol. *3*, e68.
10.1371/journal.pbio.0030068.

64. Xue, M., Atallah, B.V., and Scanziani, M. (2014). Equalizing excitation–inhibition ratios
across visual cortical neurons. Nature *511*, 596–600. 10.1038/nature13321.

65. Jiang, X., Shen, S., Cadwell, C.R., Berens, P., Sinz, F., Ecker, A.S., Patel, S., and Tolias,
A.S. (2015). Principles of connectivity among morphologically defined cell types in adult
neocortex. Science *350*, aac9462. 10.1126/science.aac9462.

66. Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau, D.,
Wieser, E., Taylor, J., Berg, S., Smith, N.J., et al. (2020). Array programming with NumPy.
Nature *585*, 357–362. 10.1038/s41586-020-2649-2.

1123

1124

Supplemental Figures

1126 Figure S1

1127

Figure S1: **(A-E): Both grating patches and oriented noise stimuli produce steady-state**
elevation and suppression in layer 2/3 of V1. (A) dF/F response at each imaging pixel to
oriented noise stimuli (small stimulus, FWHM = 15 deg, stimulus approximately aligned to cells'
receptive fields measured outside this experiment, same animal as in Fig. 1), corresponding to
the deconvolved cell responses shown in Fig. 1B. Here and in Fig. 1B, responses are measured
beginning 750 ms after stimulus onset to focus on steady-state response (Methods.) Evidence
of suppression is seen here but is more evident when data is deconvolved (compare this panel
to Fig. 1B), as expected for sustained suppression preceded by a transient, as the initial
transient seen in Figs. 2, 3, 7. **(B)** dF/F response to a drifting grating (Gabor patch, spatial
frequency 0.1 cpd, FWHM 15 deg), from the same animal, showing cells that are elevated and
suppressed in response to drifting gratings. Overall pattern of responses to noise stimulus and
grating is similar. **(C)** Deconvolved population response to oriented noise stimulus (replicated
from Fig. 1F for comparison.) Stimulus on during time indicated by light red shaded box. **(D)**
Deconvolved cell responses to Gabor patch, same data as in (B). Gabor patches drive both
steady-state elevation and suppression, though show signs of stronger off responses and
potentially a larger onset transient. **(E)** Population deconvolved response to oriented noise
stimulus in two additional animals, consistent with effects from example animal. **(F-H): Spatial**
distributions of elevated and suppressed cells are consistent with an inhomogeneous

spatial Poisson process, independent within and across classes. (F-H) Example L-

functions (Baddeley et al., 2015) from a typical animal (blue: data, black: expectation from

Poisson process model, error bars: global envelopes of Poisson process model), showing

agreement with the Poisson process model within elevated, suppressed, and across

populations, respectively (all $p > 0.05$, Bonferroni correction). See Methods for analysis details.

1152

Figure S2: **(A-B): Viral approach expresses opsin in only excitatory cells. (A)** Selective
expression of opsin in excitatory cells only, as expected for the double-inverted lox-site AAV
vector and excitatory Cre mouse line (Emx1-Cre). The stChrimsonR opsin was fused to
mRuby2, so we measured mRuby2 mRNA (red) and VGLUT1 mRNA (white), a marker of
excitatory cells, via fluorescent in situ hybridization (RNAscope; Methods). Cell counting showed
76% of neurons are VGLUT1 positive (N = 195/257). Arrows highlight a few example neurons.
As expected, all cells that express the opsin are excitatory, but not all excitatory neurons
express the opsin (59% of VGLUT1 cells are mRuby2 positive: N = 115/195). **(B)** mRuby2
mRNA (red) and VGAT mRNA (green), a marker of inhibitory neurons. 24% of neurons are
VGAT positive (N = 62/257), and zero express the opsin. **(C-D): Sorting and quality of**

electrophysiology data (C) Single units demonstrate higher SNR (N=136, median = 3.32) than
multi-units (N=184, median = 2.26). **(D)** Mean spike-waveform of putative excitatory units (wide)
in red (N = 94), mean spike-waveform of putative inhibitory units (narrow) in blue (N = 42.)
Bimodal histogram of spike widths is shown in Fig. 5B. **(E-H): Number of detected elevated**
and suppressed units by depth and horizontal distance, presented in terms of proportion
of the units in the population. (E-F) Fraction of neurons found at each depth and horizontal
distance for elevated (red) and suppressed (blue) neuron populations. **(G-H)** Same as A-B, but
summed across depth. Error bars: Wilson score 95% CIs. **Steady-state firing rates of**
neurons in layer 2/3 follow a weak spatial gradient with similar trends as the spatial
distribution observed in cell counts. (I) Elevated cell steady-state rates, with the highest and
lowest powers for comparison. Rate is the difference in firing rate during stimulation relative to
baseline. **(J)** Suppressed cell steady-state rates, with the highest and lowest powers for
comparison, measured in relation to decreases from baseline. **(K-O): Shape and variance of**
response distributions are inconsistent with cell-autonomous effects. (K) Competing
hypotheses for response distribution shape. If the variance in responses is driven by network
input, we would expect that responses would not be strongly correlated to opsin expression
levels, and also as stimulation increases, response variance would also increase. **(L)** If cell-
autonomous features like opsin expression levels drive the responses at high powers, the opsin
input should dominate network input, leading to variance decreases and/or a bimodal response
distribution. **(M)** Histograms of the electrophysiological response for increasing laser intensities.
(N) Variance of the distributions in (A), plotted across laser intensity. Shaded blue: standard
error. **(O)** Two-photon response variance to optogenetic stimulation, sorted by estimates of
opsin expression. We see an increase in variance in both the opsin positive and negative cells,
which does not support the cell-autonomous account.

1187

1189

Figure S3: **Center-surround organization is present regardless of deconvolution method.**
(A) Mean whole-frame dF/F GCaMP response in an example animal. **(B)** We tested 3 different
methods of deconvolution, OASIS [59], first-differences (i.e. subtracting one frame from the
previous), and Widefield Deconvolution [61]. Widefield Deconvolution is expected to be the best
method, as it is designed for data like this and does not incorporate the sparse-event constraints
of OASIS, which is designed for single neurons. We found similar time-series results for each of
the methods. The first-differences method (i.e. deconvolution with an kernel that decays
immediately) seems to overestimate decreases in firing rate, as might be expected. All panels
use the same dataset. **(E, F, G)** Spatial distribution of responses during the early laser period.
All deconvolution methods produce a qualitatively similar excitatory response during this early
period. **(H, I, J)** Spatial distributions of responses during late laser period demonstrates slight
differences in size of surround, but overall a qualitatively similar center-surround organization
with all methods. **(K, L, M)** Spatial distributions of response during the late laser period, but with
dashed contours depicting the manually-drawn regions of interest (ROIs) that we used to
produce the time-series data in **(B, C, D)**, with red dashed contours representing the center
ROIs, and blue dashed contours representing the surround ROIs.

Figure S4: **(A-C): Stimulation response correlates to the pattern of stChrimsonR**
expression. (A) Example animal's response during the late stim period. **(B)** Example animal's
stChrimsonR expression pattern (gray: fluorescence) with overlaid contours of fluorescence
intensity. **(C)** Example animal's response during the late stim period overlaid with their
stChrimsonR contours. **(D-F): Expression and surround response of each mouse in the**
widefield dataset. (D-F) Field-of-view showing GCaMP expression (green image),

stChrimsonR expression (red image). Contours: black = 80% of maximum illumination, red =
80% of maximum expression, blue = local minimum of the surround suppression. **(G-I):**
Spatiotemporal response pattern of widefield response to excitatory cell stimulation. (G)
Response over time, each frame corresponding to a timepoint in the timecourse in (B). Group
ROIs were selected as the top 30% of positively or negatively responding pixels within 1 mm of
the center of response and were used to compute the timecourses in (B); Methods. **(H)**
Timecourse of the response in the center and surround in the group-averaged signal. Error:
standard deviation across pixels. **(I)** Reproduction of Fig. 4D. Same as (B), but each animal's
timecourse was generated from their individual data and then averaged, resulting in less
smoothing between center and surround due to small variations in optogenetic expression
region size across animals. Error: standard error across animals.

1226

 Figure S5

1228

Figure S5: **Differences in dynamics are restricted to those seen between the onset of wide- and narrow-waveform cells.** The excitatory and inhibitory (wide- and narrow-) onset latency difference is shown in Fig. 7C. Other quantities shown here do not differ: wide- vs narrow (excitatory vs inhibitory) time to steady state (B), and onset time and time to steady state (C,D) for elevated and suppressed groups of wide-waveform excitatory cells. **(A)** Example single neuron firing rate with fits. To obtain the onsets for individual cells, each cell's mean timecourse was smoothed with width dependent on the detectability of the transient signal (SNR; Methods), then a logistic function was fit to data from time range [-100ms, 100ms]. The onset time (latency) was defined as the time to half-max of the logistic function. **(B)** No difference in median time to steady state was found across narrow-spiking and wide-spiking cells. **(C)** No difference in median onset time for elevated and suppressed groups of wide-waveform (excitatory) cells. **(D)** Same as B, but difference in median time to steady-state.

1241

1242

Figure S6: **Distributions of opsin fluorescence measured in vivo. (A)** Red channel after
neuropil correction (Methods). Y-axis, first derivative of CDF, smoothed with LOWESS; point
separating non-expressing neurons (left, below dashed line) and expressing (above dashed
line) is set at the local minimum. Error bars (light blue): bootstrapped standard error (N=244
neurons, N=3 animals). **(B)** Histogram, same data. The log-likelihoods (LLs) indicate that a
lognormal distribution (orange) fits the observed distribution better than a Gaussian (green).
Shown: fits used for simulations, excluding non-expressing neurons (gray). LLs, lognormal =
109.0, Gaussian 49.0. (LLs when including all neurons: lognormal = 95.3, Gaussian = 47.5).

1252
Figure S7: **(A-B): Networks at all tested recurrent strengths operate within the ISN regime.**
(A) To examine paradoxical suppression, we record the steady-state responses of the I cells in
response to different levels of stimulation. We performed this experiment on all networks
presented in Figure 8. **(B left)** Steady-state responses of the I cell population to 5 levels of
stimulation. Error bars are standard error to the mean. Graph has been zoomed into the region
which clearly shows paradoxical suppression in all 3 networks. This paradoxical suppression is
predicted for both loosely and tightly balanced networks. Our simulations used three recurrent
strength values, one in the tight-balance regime and two in the loose-balance regime, and we
confirmed that all three showed paradoxical effects of suppression when I cells are stimulated
(B right) Same as (B left) but input normalized by the input value calculated in Fig. 8 to drive
each network to the same firing rate (input level that achieves same value of the 75th percentile
of evoked rates; see Fig. 8). **(C-F): Inhibitory neurons in balanced state model show similar**
responses to excitatory neurons but are recruited after initial stimulation. (C) Mean
timecourses for elevated and suppressed inhibitory cells (left and right, respectively) show the
same characteristic transient response followed by steady-state responses. **(D)** Cross-
correlation analysis of E- and I-cell response. Network has no synaptic delays built into the
model. E-cells respond to direct stimulation, and then I-cells are recruited after. **(E)** Population
distribution of steady-state responses is similar across E- and I-cells, though excitatory cells
show a slightly longer-tailed positive response (true in the data as well; Fig. 5E), as seen
through the distribution of responses or their corresponding CDFs **(F)**

1274

Figure S8: **(A-E): Increasing strength of recurrent connections does not substitute for**
recurrent connection variability. (A) Cumulative distribution of responses to optogenetic
stimulation in model with 1x recurrent strength, matched to the 75[th] percentile of the response
measured using electrophysiology. Negligible recurrent variability in this simulation (same
number of recurrent connections to each neuron, variability in recurrent strength ~1% of mean,
see Methods), and so spread in responses as a function of input is due to optogenetic input
variability. Distribution of input is inferred from data in Fig. 6 (lognormal fit; Methods.) **(B)** Same
as (A) but in model with 2.5x recurrent strength. **(C)** Relationship between input and steady-
state response in the model with 1x recurrent strength. Marginal distribution of response show
on the right. **(D)** Same as (C), but in model with 2.5x recurrent strength.
Note that both stimulations produce similar variability between input strength and firing rates.
This variability is seen as spread in the red cloud of points around an imagined curve that could
be fit through the points. **(E)** Estimated mean absolute error of the relationship between the

input and output as measured by a LOWESS fit across all recurrent strength manipulations. **(F-**
H): Sensitivity analysis demonstrates that matching suppression is achievable within
confidence bounds observed baseline firing rates, but only in the network with the
strongest recurrent connectivity strength. (F) Bootstrapped distribution of baseline firing rate
estimated from electrophysiology data. 95% confidence intervals are drawn from the
bootstrapped distribution. **(G)** Bootstrapped distribution of the proportion of the population that is
suppressed following optogenetic stimulation, estimated from the electrophysiology data. **(H)**
The network baseline firing rate and recurrent strength were systematically manipulated, finding
that the only networks that can replicate the proportion of suppression we observe within the
baseline firing rate we observe are networks with strong recurrent connectivity. **(I-M): Models**
with 41% of cells without opsin replicate steady-state dynamics, noisy relationship
between opsin input and steady-state response, and response distribution. (I) Distribution
of opsin input was generated by sampling from a lognormal distribution fit to our observed opsin
fluorescence, and in order to replicate the sparse expression we observed in histology we set
41% of cells to 0 at random. **(J)** Mean timecourse of response to stimulation in elevated cells
maintains the same transient and steady state dynamics observed in the main simulations. **(K)**
Same as (B), but in suppressed cells. **(L)** Relationship between opsin input and steady state
response remains weak but positive, marginal distribution shown on right. **(M)** Cumulative
response distribution to stimulation shows typical long tail and large proportion of suppressed
responses.