Transcriptional response of circadian clock genes to an 'artificial light at night' pulse in the cricket *Gryllus bimaculatus*

Levy, Keren[1], Fishman, Bettina[2], Barnea, Anat[3], Ayali, Amir[1,4*], Tauber, Eran[2*]

[1]*School of Zoology, Tel Aviv University, Tel-Aviv 6997801, Israel;kerenlevy@mail.tau.ac.il; ayali@post.tau.ac.il*

[2]*Department of Evolutionary and Environmental Biology and Institute of Evolution, University of Haifa, Haifa 3498838, Israel; bfishman@univ.haifa.ac.il; etauber@univ.haifa.ac.il*

[3]*Department of Natural and Life Sciences, The Open University of Israel, Raanana 43107, Israel; anatba@openu.ac.il*

[4]*Sagol School of Neuroscience, Tel Aviv University, Tel-Aviv 6997801, Israel; ayali@post.tau.ac.il*

* Correspondence: AA: ayali@post.tau.ac.il; ET: etauber@univ.haifa.ac.il

Keywords: Light pollution, Artificial light at night, ALAN, Masking, Insects, Circadian rhythm, Extracellular RNA

ABSTRACT

Light is the major signal entraining the circadian clock that regulates physiological and behavioral rhythms in most organisms, including insects. Artificial Light at Night (ALAN) disrupts the natural light-dark cycle and negatively impacts animals at various levels. We simulated ALAN using dim light stimuli and tested their impact on gene expression in the cricket *Gryllus bimaculatus*, a model of insect physiology and chronobiology. At night, adult LD crickets were exposed for 30 minutes to a light pulse of 2-40 lx. The relative expression of five circadian clock associated genes was compared using qPCR. A dim ALAN pulse elicited tissue-dependent differential expression in some of these genes. The strongest effect was observed in the brain and in the optic lobe, the cricket's circadian pacemaker. Expression of *opsin-Long Wave* (*opLW)* was upregulated, as well as *cryptochrome1-2 (cry),* and *period* (*per*).

Our findings demonstrate that even a dim ALAN exposure may affect insects at the molecular level, underscoring the detrimental impact of ALAN on the circadian clock system.

INTRODUCTION

For most animal species, light detection is vital for their temporal adaptation to the earth's diel cycles. Monitoring the annual change in the daily light duration (photoperiod) is also used by many organisms in adapting to the seasonal cycle and conditions [1–3]. Many behaviors depend on such cycles: daily periods of activity and rest, sleep, foraging, courtship and mating, ecdysis in insects, migration, as well as diapause, to name just a few [1,4–7]. Consequently, many living organisms are highly sensitive to the detection of light and utilize light for other functions in addition to vision, such as entrainment of the circadian clock, timing of gene expression, sexual maturation, and hormonal regulation [4,5,8–10].

Artificial light at night (ALAN) is a fast-growing worldwide phenomenon [11,12]. ALAN refers to both an increase in light intensity and changes in the naturally occurring light spectrum. ALAN causes changes in the behavior and temporal activity of many animals, such as birds [13,14], rodents [15], anurans [16], and insects [17,18]. ALAN impairs sleeping behavior [19], camouflage, and population synchronization [17,20,21]. Moreover, it affects predation [22,23], orientation [24], and pollination [25,26], causes high insect mortality [27–29], changes in community structures, and changes in biodiversity [30–32]. Although the specific impacts of ALAN on insects have recently received considerable attention [33], the underlying molecular mechanisms are still poorly understood. Here, we specifically focus on the effect of dim ALAN on the circadian system, which has been well studied in vertebrates [34–36], but to a lesser extent in insects.

The circadian clock and its light input pathways in insects have been studied extensively in the fruit-fly, *Drosophila melanogaster* [37,38], as well as in cockroaches, crickets, and bees [reviewed by 39]. The cricket *Gryllus bimaculatus* has been widely used as a model for behavior and circadian activity [9,40]. In contrast to the fruit-fly, whose clock neurons are located in the brain, the circadian pacemaker of the cricket is located in the optic lobes [39,41–48]. Light entrainment of the circadian system in the cricket is mediated by green-sensitive opsins in the insect's compound

eyes [43]. The pacemaker consists of two major feedback loops, one of which is based on the *period* (*per*) and *timeless* (*tim*) genes, repressing their own transcription by inhibiting the transcription of *Clock* (*Clk*) and *cycle* (*cyc*). The other feedback loop is based on the two *cry* genes, *cry1* (a *Drosophila*-type *cry*), which is a light sensitive photopigment, and *cry2*, a light insensitive mammalian-type *cry* [39,49]. This second feedback loop involves the upregulation of *PAR domain protein 1* (*Pdp1*) and *c-fosB* [49–51].

Recently, we have demonstrated that lifelong exposure of male crickets to dim ALAN as low as 2 lx leads to a loss of rhythmicity and desynchronization of stridulation and locomotion behavior [17]. Stridulation serves to attract females and is therefore crucial for the species' fitness. We have reasoned that these behavioral changes are mediated by changes in gene expression. Indeed, recent studies on birds [52,53], amphibians [54], and glow-worms [55] have revealed altered gene expression following exposure to dim ALAN, an effect that was present in both the visual system and various other tissues. Here, we sought to investigate the effect of exposure to dim ALAN on gene expression in four different tissues of the cricket G. bimaculatus. Due the effect of ALAN on the male behavior (locomotion, stridulation), we have focused on transcriptional response in males. Understanding the effect of dim ALAN on internal processes and pathways is crucial for assessing the threats of ALAN to pollinators and ground-dwelling insects, as well as for predicting the long-term effects on ecosystems.

RESULTS

The relative levels of gene expression in the different cricket tissues and hemolymph samples following a 30 min light pulse of 2, 5, or 40 lx, and a no-pulse control, are shown in Figure 1. As can be seen, in the brain the expression of most genes was affected by the different light intensities (Figure 1A). The expression levels of both *per* and *cry2* significantly increased with increasing light pulse intensity ($F_{3,21}= 3.70$, $p = 0.028$, and $F_{3,24}= 3.04$, $p = 0.048$, respectively, Figure 1A). A dependence on light pulse intensity can also be seen for *cry1*, although the effect was only marginally significant ($F_{3,21}= 2.96$, $p = 0.056$). The expression of *opLW* differed significantly only when light treatment experiments were pooled and compared with the control

($F_{1,26}$= 4.90, p = 0.036). The *c-fosB* transcription in the brain did not differ among the different treatments ($F_{3,24}$= 1.15, p = 0.35).

In the optic lobe, the expression of opLW was 100-fold higher compared to that seen in the brain, but did not change significantly following exposure to light (Figure 1B). In contrast to the brain (Figure 1A), in which most genes exhibited light-induced upregulation, in the optic lobe the light stimuli elicited a decrease in the expression of most genes. Downregulation was observed in cry2 although only marginally significant (F3,16= 2.86, p = 0.07) and a similar trend was observed in cry1 (Figure 1B). No change in per and c-fosB expression was observed (F3,17= 1.45, p = 0.26, F3,18= 0.98, p = 0.42, respectively).

In the Malpighian tubules most gene expression did not change with increased light pulse intensity (Figure 1C). A marginally significant effect was observed for *opLW* ($F_{3,42}$= 2.58, p = 0.066) and no differences were found for *cry2* ($F_{3,43}$= 0.80, p = 0.501), *per* ($F_{3,40}$= 1.14, p = 0.34), or *c-fosB* ($F_{3,42}$= 0.18, p = 0.91). However, *opLW* expression was higher in the control, compared to the light treatments pooled together ($F_{1,44}$= 6.84, p = 0.012), and the expression of *cry1* significantly diminished following exposure to any of the light pulses ($F_{3,41}$= 7.21, p = 0.0005, Figure 1C).

Interestingly, we found traces of expression of all genes in the hemolymph (Figure 1D). Gene expression did not differ significantly among samples. However, when light treatments were pooled together, upregulation in *per*, was observed ($F_{1,18}$= 5.83, p = 0.027, Figure 1D).

In order to enable a better comparison of gene expression among tissues, we repeated the analysis using within-tissue normalization of the expression values (relative to the control mean expression values in each tissue, Figure 2). Gene expression patterns varied substantially among tissues. The expression of *opLW* was strongly upregulated in the brain and in the optic lobe throughout all three light treatments, while being downregulated in the Malpighian tubules (Figure 2A). The light stimuli elicited upregulation of *cry2* in the brain and downregulation in the optic lobe (Figure 2B). The transcriptional response of *per* showed a similar pattern, being upregulated in the brain and downregulated in the Malpighian tubules (Figure 2C). The expression of *c-fosB* showed no significant changes in the brain, while being downregulated in the optic lobe. *c-fosB* was the only gene that was upregulated in the Malpighian tubules

(Figure 2D). Expression of *cry1* was downregulated in all tissues except the brain, where it was upregulated under 40 lx only ($F_{3,23}= 7.77$, $p < 0.001$. Figure 2E). The PCA of the relative gene expression indicated the necessity for separate calculations for each tissue, as it clearly separated the optic lobe and the haemolymph from the brain and the Malphigian tubules, while the latter two mostly overlapped (Figure 2F).

To simultaneously analyze the transcriptional differences between the treatments, we used Linear Discriminant Analysis (LDA). The first two LD functions explained 67.12% of the variation in the brain, 73.29% in the optic lobes, 66.87% in the Malpighian tubules and, 73.76% in the haemolymph (Figure 3). In the brain, the control group was clearly separated from the 5 and 40 lx treatments, but not from the 2 lx one (Figure 3A). In contrast, the clustering of the treatments in the optic lobe revealed a substantial difference between the control and all ALAN treatments (Figure 3B). In the Malpighian tubules and in the haemolymph, the discriminant map did not reveal any significant difference between the control and the ALAN treatments (Figure 3C,D).

DISCUSSION

Insects have served for almost a century as successful models in laboratory studies on the role of light in regulating circadian behavior [7], and in raising questions about the underlying molecular mechanisms. Here, we investigated the effect of a short, dim ALAN pulse on the gene expression of five genes associated with the circadian system of the cricket *Gryllus bimaculatus*.

This work substantiates the findings of previous studies [56] that uncovered the heterogeneity of circadian pacemakers across various tissues in *G. bimaculatus* (see PCA analysis, Figure 2F), and demonstrates the tissue-specific effects of ALAN on transcription. In the optic lobe, where the cricket's central pacemaker is located [46,48,57,58], the response to ALAN exposures was clearly distinct from that of the control, presumably due to the strong effect of dim ALAN on *opLW* expression. In the brain, transcriptional responses of the control and the 2 lx treatment were largely overlapping, and to a lesser extent also with those of the 5 lx and 40 lx treatments. The Malpighian tubules and haemolymph seemed less affected by the light exposures.

Indeed, a previous study [56] suggested that *G. bimaculatus* tubules do not harbor a circadian pacemaker, as *per* transcript levels lacked any diurnal rhythm (in contrast to in *Drosophila* [59]).

The expression of circadian clock genes in *G. bimaculatus* in tissues other than the optic lobes has been previously reported [56]. Particularly interesting is the role of these genes in the brain. Expression of *per* oscillates under constant darkness (DD), and this oscillation persists after removal of the optic lobes. However, in the absence of the optic lobes, the phase of the brain *per* mRNA rhythm is aberrant. These results allude to a brain circadian pacemaker that is subordinate to the central clock in the optic lobes [56].

In the hemolymph, we found minute levels of circadian clock transcripts, which to the best of our knowledge is the first report of extracellular RNA (exRNA) in circadian clock transcripts. exRNA is emerging as a newly discovered form of intracellular signaling [60]. It is secreted to the biofluids, encapsulated by protecting extracellular vesicles. The exRNA is largely composed of small non-coding RNA, and to a lesser extent messenger RNA [60]. exRNA have been shown to be an effective biomarker for diagnostic purposes, but their functional role awaits further study.

The rather rapid transcriptional response to light of clock genes such as *per* or *cry*, is intriguing, and the underlying molecular mechanism warrants further investigation. In *Drosophila*, a similar rapid change of *per* mRNA following a light pulse at *zeitgeber* time (ZT) 15 has been previously reported [61,62], although, in contrast to the cricket, light evoked downregulation of *per.* It was suggested that the response to light is mediated by an histone acetylation mechanism. In accord with reports pertaining to various other insects (e.g. ants [63] and lepidoptera [64]), we observed very high levels of *opLW* in the crickets' optic lobe. All ALAN treatments induced some upregulation in *opLW* expression in the optic lobe, although not significant (we sampled 4.5 h post lights-off, while 6 h after onset of darkness a substantial downregulation has been reported [43], Figure 1 therein).Given that the circadian photosensitivity of *G. bimaculatus* varies during the day, as was shown by the phase response curve of this species [65,66], one can predict that the effect of transient ALAN would be time dependent.

The disruption caused by ALAN to the circadian transcriptional cycle (Figure 1) is likely to disrupt the diurnal pattern of behavior (locomotor, stridulation), but the specific behavioral output awaits further study.

The circadian disruption caused by dim (2-5 lx) ALAN may have ecological implications, which should be taken into consideration when designing regulations concerning suitable light intensity for park and city lights [67]. Not only flying insects, which are attracted to light [28], are threatened by ALAN, but also ground-dwelling insects [21,24,68]. The deep-level effect of ALAN may therefore threaten many more species of insects than currently assumed, and add to the ongoing insect decline described in the last century [69,70]. Our own work in crickets [17] demonstrated that ALAN can lead to arrhythmicity in locomotory behavior and male stridulation and therefore may affect the survival and reproduction of exposed individuals.

It should also be noted, that outdoors, especially in proximity to cities and streetlights, insects are not exposed to a single light pulse, but rather to night-long, and even lifelong ALAN [17]. Populations of the small ermine moth for example, were found to reduce their flight-to-light behavior following long-term exposure to ALAN, which may result in lower predation risk and mortality in this population [71]. It remains unknown whether crickets, and indeed other insects, have developed similar adaptations to chronic ALAN exposure.

MATERIALS AND METHODS

Insect rearing conditions

Gryllus bimaculatus crickets were reared under a constant temperature of 26±1°C and white fluorescent light (5W white CFL bulb, NeptOn, 6500 K, 380-780 nm, peak: 547 & 612 nm, Figure S1), under a 12h light:12h dark cycle. Actual light intensity measured at the top of the containers ranged from 250 to 350 lx, while the intensity under the provided shelter (egg cartons) ranged from 20 to 60 lx. Crickets were fed three times a week with dog-food pellets and vegetables. The rearing boxes contained water flasks with absorbent cotton wool.

Light-pulse experiments and sample preparation

Individual males, 3-5 days post adult emergence, were removed from the breeding colony and maintained individually, under similar 12h light:12h dark conditions as above, for three days. On the fourth night, three hours post lights-off (i.e. at *zeitgeber* time ZT 15), the experimental animals were exposed to a 30 min long light pulse of 2, 5, or 40 lx (by partially covering the same 5W white CFL bulb as above). An additional group (no pulse) served as control.

One hour after pulse termination (ZT 16.5), the insects were decapitated under red light (led, 600-670 nm, peak: 642 nm), and their brain, optic lobes, and Malpighian tubules were removed over ice. Hemolymph (25 µl) was also extracted from each animal. Samples were placed separately into individually marked PCR tubes containing 100 µl RNAlater (Thermo Fisher Scientific), and immediately frozen at -20°C.

Total RNA was extracted from each sample using the Purelink RNA mini kit (Invitrogen), according to the manufacturer's instructions. Samples with genomic DNA residuals were treated with the PureLink™ DNase Set (Invitrogen). RNA concentration was determined by NanoDrop 2000 (Thermo Fisher Scientific). First strand cDNA synthesis was carried out using the RT Random Primers of the High-Capacity cDNA Reverse Transcription kit (Applied Biosystems™) and treated with RNase Inhibitor (Applied Biosystems™) according to the manufacturer's instructions. cDNA was synthesized out of 20 ng/µl RNA and was then 10x diluted.

The number of successfully collected samples per experimental group varied as follows: 5–11 brain samples, 4–6 optic lobe samples, 9–15 Malpighian tubules, and 5-7 haemolymph samples. Technical difficulties resulted in not all samples yielding useful results (a detailed list of the final sample sizes is given in Table S1).

Primers and qPCR

The expression of two housekeeping genes (*Actin* and *Rpl18a*) and five target genes (*opLW*, *per*, *cry1*, *cry2*, *c-fosB*) was determined in all samples. Primer sequences are listed in Table S2. A standard curve was generated for each of the genes by using a serial dilution (1:2) of the pooled cDNA [72]. qPCR (Quantstudio 3, Thermo Fisher Scientific) was run using 59°C as annealing temperature. Fast SYBR® Green Master

Mix (Applied Biosystems™) was used. The cycle point was calculated using QuantStudio™ Design and Analysis Desktop Software (Applied Biosystems).

Data-processing and statistical analysis

Gene expression was quantified using the relative standard curve method and following primer efficiencies: *opLW*: 94.38%, *cry2*: 105.95%, *per*: 100.16%, *c-fosB*: 105.02%, and *cry1*: 93.57%. The target gene expression was normalized by dividing its values by the corresponding geometric mean of both housekeeping genes. The expression values were log transformed and a nested ANOVA was applied using R version 4.1.3 [73] and the *nlme* package [74]. A mixed model was used, with light treatment and cDNA replicate as fixed and random effects, respectively. A principal component analysis (PCA) was applied to all expression values and the first two principal components (PC1 and PC2) of the PCA were used as explanatory variables for a linear discriminant analysis (LDA), with the treatments as response variables.

AKNOWLEDGEMENTS

We thank Odelia Pizanti, Yael Diesendruck, Daniel Knebel, David Burg, Alex Slavenko, Lilach Raijman, Omer Lavy, Michaela Kolker-Ghatan, and Talma and Rami Zelingher for their assistance. This work is dedicated to our mentor Prof. Meir Paul Pener (1930-2021).

FUNDING

This research was funded by The Open University of Israel Research Fund and by the Constantiner Institute for Molecular Genetics of Tel Aviv University.

AUTHOR CONTRIBUTION

Conceptualization, A.B., A.A., E.T. and K.L.; methodology, A.A., E.T., and B.F.; validation, K.L., B.F., and E.T.; formal analysis, K.L., and E.T.; investigation, K.L.; resources, A.B., E.T.; writing—original draft preparation, K.L.; writing—review and editing, E.T., A.A., and A.B.; funding acquisition, A.B.

The authors have no competing interests related to this study.

Supplementary material is available online on FigShare: DOI: 10.6084/m9.figshare.20721265

REFERENCES

1.   Borges, R.M. Impacts of Artificial Light at Night on Biological Timings. *Indian J. Entomol.* **2022**, *84*, 483–492, doi:10.55446/IJE.2022.182.

2.   Saunders, D.S. Insect Photoperiodism: Seeing the Light. *Physiol. Entomol.* **2012**, *37*, 207–218, doi:10.1111/j.1365-3032.2012.00837.x.

3.   Pittendrigh, C.S. Circadian Rhythms and the Circadian Organization of Living Systems. *Cold Spring Harb. Symp. Quant. Biol.* **1960**, *25*, 159–184, doi:10.1101/SQB.1960.025.01.015.

4.   Kronfeld-Schor, N.; Visser, M.E.; Salis, L.; van Gils, J.A. Chronobiology of Interspecific Interactions in a Changing World. *Philos. Trans. R. Soc. B Biol. Sci.* **2017**, *372*, 20160248, doi:10.1098/rstb.2016.0248.

5.   Ampleford, E.J.; Steel, C.G.H. Circadian Control of Ecdysis in *Rhodnius prloixus* (Hempitera). *J. Comp. Physiol. A* **1982**, *147*, 281–286, doi:10.1007/BF00609852.

6.   Helm, B.; Visser, M.E.; Schwartz, W.; Kronfeld-Schor, N.; Gerkema, M.; Piersma, T.; Bloch, G. Two Sides of a Coin: Ecological and Chronobiological Perspectives of Timing in the Wild. *Philos. Trans. R. Soc. B Biol. Sci.* **2017**, *372*, 20160246, doi:10.1098/rstb.2016.0246.

7.   Saunders, D.S.; Steel, C.G.H.; Vafopoulou, X.; Lewis, R.D. Circadian Rhythms of Activity in Individual Insects. In *Insect Clocks*; Elsevier B.V: Amsterdam, 2002; pp. 7–42.

8.   Merrow, M.; Spoelstra, K.; Roenneberg, T. The Circadian Cycle: Daily Rhythms from Behaviour to Genes. *EMBO Rep.* **2005**, *6*, 930–935, doi:10.1038/sj.embor.7400541.

9.   Horch, H.W.; Mito, T.; Popadi, A.; Ohuchi, H.; Noji, S. *The Cricket as a*

Model Organism; Springer: Tokyo, 2017; ISBN 978-4-431-56476-8.

10. Moaraf, S.; Vistoropsky, Y.; Pozner, T.; Heiblum, R.; Okuliarová, M.; Zeman, M.; Barnea, A. Artificial Light at Night Affects Brain Plasticity and Melatonin in Birds. *Neurosci. Lett.* **2020**, *716*, 134639, doi:10.1016/j.neulet.2019.134639.

11. Falchi, F.; Furgoni, R.; Gallaway, T.A.; Rybnikova, N.A.; Portnov, B.A.; Baugh, K.; Cinzano, P.; Elvidge, C.D. Light Pollution in USA and Europe: The Good, the Bad and the Ugly. *J. Environ. Manage.* **2019**, *248*, 109227, doi:10.1016/j.jenvman.2019.06.128.

12. Hölker, F.; Wolter, C.; Perkin, E.K.; Tockner, K. Light Pollution as a Biodiversity Threat. *Trends Ecol. Evol.* **2010**, *25*, 681–682, doi:10.1016/j.tree.2010.09.007.

13. Dominoni, D.; Quetting, M.; Partecke, J. Artificial Light at Night Advances Avian Reproductive Physiology. *Proc. R. Soc. B Biol. Sci.* **2013**, *280*, 20123017–20123017, doi:10.1098/rspb.2012.3017.

14. Amichai, E.; Kronfeld-Schor, N. Artificial Light at Night Promotes Activity Throughout the Night in Nesting Common Swifts (*Apus apus*). *Sci. Rep.* **2019**, *9*, 1–8, doi:10.1038/s41598-019-47544-3.

15. Kumar, D.; Soni, S.K.; Kronfeld-Schor, N.; Singaravel, M. Wheel-Running Activity Rhythms and Masking Responses in the Diurnal Palm Squirrel, *Funambulus pennantii*. *Chronobiol. Int.* **2020**, *37*, 1693–1708, doi:10.1080/07420528.2020.1826959.

16. Buchanan, B.W. Observed and Potential Effects of Artificial Night Lighting on Anuran Amphibians. *Ecol. Consequences Artif. Night Light.* **2006**, *09*, 192–220.

17. Levy, K.; Wegrzyn, Y.; Efronny, R.; Barnea, A.; Ayali, A. Lifelong Exposure to Artificial Light at Night Impacts Stridulation and Locomotion Activity Patterns in the Cricket *Gryllus bimaculatus*. *Proc. R. Soc. B Biol. Sci.* **2021**, *288*, 20211626, doi:10.1098/rspb.2021.1626.

18. Durrant, J.; Michaelides, E.B.; Rupasinghe, T.; Tull, D.; Green, M.P.; Jones,

T.M. Constant Illumination Reduces Circulating Melatonin and Impairs Immune Function in the Cricket *Teleogryllus Commodus*. *PeerJ* **2015**, *3*, e1075, doi:10.7717/peerj.1075.

19. Raap, T.; Pinxten, R.; Eens, M. Light Pollution Disrupts Sleep in Free-Living Animals. *Sci. Rep.* **2015**, *5*, 1–8, doi:10.1038/srep13557.

20. Gaston, K.J.; Davies, T.W.; Nedelec, S.L.; Holt, L.A. Impacts of Artificial Light at Night on Biological Timings. *Annu. Rev. Ecol. Evol. Syst.* **2017**, *48*, 49–68, doi:10.1146/annurev-ecolsys-110316-022745.

21. Owens, A.C.S.; Lewis, S.M. The Impact of Artificial Light at Night on Nocturnal Insects: A Review and Synthesis. *Ecol. Evol.* **2018**, *8*, 11337–11358, doi:10.1002/ece3.4557.

22. Sanders, D.; Kehoe, R.; Cruse, D.; van Veen, F.J.F.; Gaston, K.J. Low Levels of Artificial Light at Night Strengthen Top-Down Control in Insect Food Web. *Curr. Biol.* **2018**, *28*, 2474-2478.e3, doi:10.1016/j.cub.2018.05.078.

23. Bolliger, J.; Hennet, T.; Wermelinger, B.; Bösch, R.; Pazur, R.; Blum, S.; Haller, J.; Obrist, M.K. Effects of Traffic-Regulated Street Lighting on Nocturnal Insect Abundance and Bat Activity. *Basic Appl. Ecol.* **2020**, *47*, 44–56, doi:10.1016/j.baae.2020.06.003.

24. Foster, J.J.; Tocco, C.; Smolka, J.; Khaldy, L.; Baird, E.; Byrne, M.J.; Nilsson, D.E.; Dacke, M. Light Pollution Forces a Change in Dung Beetle Orientation Behavior. *Curr. Biol.* **2021**, *31*, 3935-3942.e3, doi:10.1016/j.cub.2021.06.038.

25. Giavi, S.; Blösch, S.; Schuster, G.; Knop, E. Artificial Light at Night Can Modify Ecosystem Functioning beyond the Lit Area. *Sci. Rep.* **2020**, *10*, 1–12, doi:10.1038/s41598-020-68667-y.

26. Knop, E.; Zoller, L.; Ryser, R.; Gerpe, C.; Hörler, M.; Fontaine, C. Artificial Light at Night as a New Threat to Pollination. *Nature* **2017**, *548*, 206–209, doi:10.1038/nature23288.

27. Rich, C.; Longcore, T. *Ecological Consequences of Artificial Night Lighting*; Island Press: Washington DC, 2006; ISBN 9781559631297.

28. Eisenbeis, G.; Hänel, A. Light Pollution and the Impact of Artificial Night Lighting on Insects. In *Ecology of Cities and Towns: A Comparative Approach*; McDonnell, M.J., Hahs, A.K., Breuste, J.H., Eds.; Cambridge University Press: Cambridge, 2009; pp. 243–263 ISBN 9780511609763.

29. Perkin, E.K.; Hölker, F.; Tockner, K. The Effects of Artificial Lighting on Adult Aquatic and Terrestrial Insects. *Freshw. Biol.* **2014**, *59*, 368–377, doi:10.1111/fwb.12270.

30. Davies, T.W.; Bennie, J.; Gaston, K.J. Street Lighting Changes the Composition of Invertebrate Communities. *Biol. Lett.* **2012**, *8*, 764–767, doi:10.1098/rsbl.2012.0216.

31. Sanders, D.; Frago, E.; Kehoe, R.; Patterson, C.; Gaston, K.J. A Meta-Analysis of Biological Impacts of Artificial Light at Night. *Nat. Ecol. Evol.* **2021**, *5*, 74–81, doi:10.1038/s41559-020-01322-x.

32. Sanders, D.; Gaston, K.J. How Ecological Communities Respond to Artificial Light at Night. *J. Exp. Zool. Part A Ecol. Integr. Physiol.* **2018**, *329*, 394–400, doi:10.1002/jez.2157.

33. Owens, A.C.S.; Cochard, P.; Durrant, J.; Farnworth, B.; Perkin, E.K.; Seymoure, B. Light Pollution Is a Driver of Insect Declines. *Biol. Conserv.* **2020**, *241*, 108259, doi:10.1016/j.biocon.2019.108259.

34. Bachleitner, W.; Kempinger, L.; Wülbeck, C.; Rieger, D.; Helfrich-Förster, C. Moonlight Shifts the Endogenous Clock of *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 3538–3543, doi:10.1073/pnas.0606870104.

35. Kempinger, L.; Dittmann, R.; Rieger, D.; Helfrich-Förster, C. The Nocturnal Activity of Fruit Flies Exposed to Artificial Moonlight Is Partly Caused by Direct Light Effects on the Activity Level That Bypass the Endogenous Clock. *Chronobiol. Int.* **2009**, *26*, 151–166, doi:10.1080/07420520902747124.

36. Kronfeld-Schor, N.; Dominoni, D.; Iglesia, H.D. la; Levy, O.; Herzog, E.D.; Dayan, T.; Helfrich-Forster, C. Chronobiology by Moonlight. *Proc. R. Soc. B Biol. Sci.* **2013**, *1765*, 20123088, doi:10.1098/rspb.2012.3088.

37. Helfrich-Förster, C. The Circadian Clock in the Brain: A Structural and Functional Comparison between Mammals and Insects. *J. Comp. Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol.* **2004**, *190*, 601–613, doi:10.1007/s00359-004-0527-2.

38. Tomioka, K.; Matsumoto, A. The Circadian System in Insects: Cellular, Molecular, and Functional Organization. In *Advances in Insect Physiology*; Academic Press Inc., 2019; Vol. 56, pp. 73–115 ISBN 9780081028421.

39. Helfrich-Förster, C. Light Input Pathways to the Circadian Clock of Insects with an Emphasis on the Fruit Fly *Drosophila melanogaster*. *J. Comp. Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol.* **2020**, *206*, 259–272, doi:10.1007/s00359-019-01379-5.

40. Tomioka, K.; Chiba, Y. Post-Embryonic Development of Circadian Rhythm in the Cricket, *Gryllus bimaculatus*: A Rhythm Reversal. *J. Comp. Physiol. □ A* **1982**, *147*, 299–304, doi:10.1007/BF00609663.

41. Tomioka, K.; Chiba, Y. Circadian Rhythm in the Neurally Isolated Lamina-Medulla-Complex of the Cricket, *Gryllus bimaculatus*. *J. Insect Physiol.* **1986**, *32*, 747–755, doi:10.1016/0022-1910(86)90077-6.

42. Abe, Y.; Ushirogawa, H.; Tomioka, K. Circadian Locomotor Rhythms in the Cricket, *Gryllodes sigillatus I*. Localization of the Pacemaker and the Photoreceptor. *Zoolog. Sci.* **1997**, *14*, 719–727, doi:10.2108/zsj.14.719.

43. Komada, S.; Kamae, Y.; Koyanagi, M.; Tatewaki, K.; Hassaneen, E.; Saifullah, A.; Yoshii, T.; Terakita, A.; Tomioka, K. Green-Sensitive Opsin Is the Photoreceptor for Photic Entrainment of an Insect Circadian Clock. *Zool. Lett.* **2015**, *1*, 1–10, doi:10.1186/s40851-015-0011-6.

44. Tomioka, K. Chronobiology of Crickets: A Review. *Zoolog. Sci.* **2014**, *31*, 624–632, doi:10.2108/zs140024.

45. Loher, W. Temporal Organization of Reproductive Behavior. In *Cricket Behavior and Neurobiology*; Huber, F., Moore, T.E., Loher, W., Eds.; Cornell University Press.: Ithaca, London, 1989; pp. 83–113.

46. Moriyama, Y.; Kamae, Y.; Uryu, O.; Tomioka, K. Gb'clock Is Expressed in the Optic Lobe and Is Required for the Circadian Clock in the Cricket *Gryllus bimaculatus*. *J. Biol. Rhythms* **2012**, *27*, 467–477, doi:10.1177/0748730412462207.

47. Masayuki, I.; Kenji, T. Temperature Dependency of the Circadian Locomotor Rhythm in the Cricket *Gryllus bimaculatus*. *Zoolog. Sci.* **1993**, *10*, 597–604, doi:10.34425/zs001120.

48. Tomioka, K. Analysis of Coupling between Optic Lobe Circadian Pacemakers in the Cricket *Gryllus bimaculatus*. *J. Comp. Physiol. A* **1993**, *172*, 401–408, doi:10.1007/BF00213522.

49. Tokuoka, A.; Itoh, T.Q.; Hori, S.; Uryu, O.; Danbara, Y.; Nose, M.; Bando, T.; Tanimura, T.; Tomioka, K. Cryptochrome Genes Form an Oscillatory Loop Independent of the per / Tim Loop in the Circadian Clockwork of the Cricket *Gryllus bimaculatus*. *Zool. Lett.* **2017**, *3*, 1–14, doi:10.1186/s40851-017-0066-7.

50. Kutaragi, Y.; Tokuoka, A.; Tomiyama, Y.; Nose, M.; Watanabe, T.; Bando, T.; Moriyama, Y.; Tomioka, K. A Novel Photic Entrainment Mechanism for the Circadian Clock in an Insect: Involvement of c-Fos and Cryptochromes. *Zool. Lett.* **2018**, *4*, 1–12, doi:10.1186/s40851-018-0109-8.

51. Moriyama, Y.; Takeuchi, K.; Shinohara, T.; Miyagawa, K.; Matsuka, M.; Yoshii, T.; Tomioka, K. Timeless Plays an Important Role in Compound Eye-Dependent Photic Entrainment of the Circadian Rhythm in the Cricket *Gryllus bimaculatus*. *Zoolog. Sci.* **2022**, *39*, 1–9, doi:10.2108/zs220011.

52. Alaasam, V.J.; Liu, X.; Niu, Y.; Habibian, J.S.; Pieraut, S.; Ferguson, B.S.; Zhang, Y.; Ouyang, J.Q. Effects of Dim Artificial Light at Night on Locomotor Activity, Cardiovascular Physiology, and Circadian Clock Genes in a Diurnal Songbird. *Environ. Pollut.* **2021**, *282*, 117036, doi:10.1016/j.envpol.2021.117036.

53. Dominoni, D.M.; de Jong, M.; van Oers, K.; O'Shaughnessy, P.; Blackburn, G.J.; Atema, E.; Mateman, A.C.; D'Amelio, P.B.; Trost, L.; Bellingham, M.; et

al. Integrated Molecular and Behavioural Data Reveal Deep Circadian Disruption in Response to Artificial Light at Night in Male Great Tits (*Parus major*). *Sci. Rep.* **2022**, *12*, 1–12, doi:10.1038/s41598-022-05059-4.

54. Touzot, M.; Lefebure, T.; Lengagne, T.; Secondi, J.; Dumet, A.; Konecny-Dupre, L.; Veber, P.; Navratil, V.; Duchamp, C.; Mondy, N. Transcriptome-Wide Deregulation of Gene Expression by Artificial Light at Night in Tadpoles of Common Toads. *Sci. Total Environ.* **2022**, *818*, 151734, doi:10.1016/j.scitotenv.2021.151734.

55. Chen, Y.; Wei, W.; Tzeng, D.T.W.; Owens, A.C.S.; Tang, H.; Wu, C.; Lin, S.; Zhong, S.; Yang, E. Effects of Artificial Light at Night (ALAN) on Gene Expression of *Aquatica ficta* Fire Fly Larvae. *Environ. Pollut.* **2021**, *281*, 1–8, doi:10.1016/j.envpol.2021.116944.

56. Uryu, O.; Tomioka, K. Circadian Oscillations Outside the Optic Lobe in the Cricket *Gryllus bimaculatus*. *J. Insect Physiol.* **2010**, *56*, 1284–1290, doi:10.1016/j.jinsphys.2010.04.010.

57. Tomioka, K.; Abdelsalam, S. Circadian Organization in Hemimetabolous Insects. *Zoolog. Sci.* **2004**, *21*, 1153–1162, doi:10.2108/zsj.21.1153.

58. Tomioka, K.; Chiba, Y. Photoperiodic Entrainment of Locomotor Activity in Crickets (*Gryllus bimaculatus*) Lacking the Optic Lobe Pacemaker. *J. Insect Physiol.* **1989**, *35*, 827–835, doi:10.1016/0022-1910(89)90098-X.

59. Hege, D.M.; Stanewsky, R.; Hall, J.C.; Giebultowicz, J.M. Rhythmic Expression of a PER-Reporter in the Malpighian Tubules of Decapitated *Drosophila*: Evidence for a Brain-Independent Circadian Clock. *J. Biol. Rhythms* **1997**, *12*, 300–308, doi:10.1177/074873049701200402.

60. Sadik, N.; Cruz, L.; Gurtner, A.; Rodosthenous, R.S.; Dusoswa, S.A.; Ziegler, O.; Van Solinge, T.S.; Wei, Z.; Salvador-Garicano, A.M.; Gyorgy, B.; et al. Extracellular RNAs: A New Awareness of Old Perspectives. In *Extracellular RNA. Methods in Molecular Biology*; Patel, T., Ed.; Springer: New York, 2018; Vol. 1740, pp. 1–15 ISBN 978-1-4939-7652-2.

61. Adewoye, A.B.; Kyriacou, C.P.; Tauber, E. Identification and Functional

Analysis of Early Gene Expression Induced by Circadian Light-Resetting in *Drosophila*. *BMC Genomics* **2015**, *16*, 1–9, doi:10.1186/s12864-015-1787-7.

62. Lee, C.; Parikh, V.; Itsukaichi, T.; Bae, K.; Edery, I. Resetting the *Drosophila* Clock by Photic Regulation of PER and a PER-TIM Complex. *Science (80-.).* **1996**, *271*, 1740–1744, doi:10.1126/science.271.5256.1740.

63. Yilmaz, A.; Lindenberg, A.; Albert, S.; Grübel, K.; Spaethe, J.; Rössler, W.; Groh, C. Age-Related and Light-Induced Plasticity in Opsin Gene Expression and in Primary and Secondary Visual Centers of the Nectar-Feeding Ant *Camponotus rufipes*. *Dev. Neurobiol.* **2016**, *76*, 1041–1057, doi:10.1002/dneu.22374.

64. Yan, S.; Zhu, J.; Zhu, W.; Zhang, X.; Li, Z.; Liu, X.; Zhang, Q. The Expression of Three Opsin Genes from the Compound Eye of *Helicoverpa armigera* (Lepidoptera: Noctuidae) Is Regulated by a Circadian Clock, Light Conditions and Nutritional Status. *PLoS One* **2014**, *9*, e111683, doi:10.1371/journal.pone.0111683.

65. Tomioka, K. Light and Serotonin Phase-Shift the Circadian Clock in the Cricket Optic Lobe in Vitro. *J. Comp. Physiol. - A Sensory, Neural, Behav. Physiol.* **1999**, *185*, 437–444, doi:10.1007/s003590050404.

66. Okada, Y.; Tomioka, K.; Chiba, Y. Circadian Phase-Response Curves for Light in Nymphal and Adult Crickets, *Gryllus bimaculatus*. *J. Insect Physiol.* **1991**, *37*, 583–590, doi:10.1016/0022-1910(91)90035-X.

67. Hölker, F.; Bolliger, J.; Davies, T.W.; Giavi, S.; Jechow, A.; Kalinkat, G.; Longcore, T.; Spoelstra, K.; Tidau, S.; Visser, M.E.; et al. 11 Pressing Research Questions on How Light Pollution Affects Biodiversity. *Front. Ecol. Evol.* **2021**, *9*, 1–13, doi:10.3389/fevo.2021.767177.

68. Grubisic, M.; van Grunsven, R.H. Artificial Light at Night Disrupts Species Interactions and Changes Insect Communities. *Curr. Opin. Insect Sci.* **2021**, *47*, 136–141, doi:10.1016/j.cois.2021.06.007.

69. Shortall, R.C.; Moore, A.; Smith, E.; Hall, J.M.; Woiwod, P.I.; Harrington, R. Long-Term Changes in the Abundance of Flying Insects. *Insect Conserv.*

Divers. **2009**, *2*, 251–260, doi:10.1111/j.1752-4598.2009.00062.x.

70. Hallmann, C.A.; Sorg, M.; Jongejans, E.; Siepel, H.; Hofland, N.; Schwan, H.; Stenmans, W.; Müller, A.; Sumser, H.; Hörren, T.; et al. More than 75 Percent Decline over 27 Years in Total Flying Insect Biomass in Protected Areas. *PLoS One* **2017**, *12*, doi:10.1371/journal.pone.0185809.

71. Altermatt, F.; Ebert, D. Reduced Flight-to-Light Behaviour of Moth Populations Exposed to Long-Term Urban Light Pollution. *Biol. Lett.* **2016**, *12*, 2016–2019, doi:10.1098/rsbl.2016.0111.

72. Taylor, S.C.; Nadeau, K.; Abbasi, M.; Lachance, C.; Nguyen, M.; Fenrich, J. The Ultimate QPCR Experiment: Producing Publication Quality, Reproducible Data the First Time. *Trends Biotechnol.* **2019**, *37*, 761–774, doi:10.1016/j.tibtech.2018.12.002.

73. R Core Team R: A Language and Environment for Statistical Computing. *R Found. Stat. Comput. Vienna, Austria.* **2022**, 2022.

74. Pinheiro, J.B. Nlme: Linear and Nonlinear Mixed Effects Models. *R Packag. version 3.1-157* **2022**, 2022.

FIGURES AND CAPTIONS

Figure 1. Transcriptional response of *G. bimaculatus* to dim ALAN. The relative gene expression (mean ± s.e.) following 30 min light pulse of 2 lx (dark grey), 5 lx (light grey) or 40 lx (yellow), and no-pulse control (black). Results are shown for the brain (A), optic lobe (B), Malpighian tubules (C), and haemolymph (D). *p < 0.05.

Figure 2: Normalized mean (± s.e.) of the relative gene expression of (A) *opLW*, (B) *cry2*, (C) *per*, (D) *c-fosB*, (E) *cry1,* and (F) Principal component analysis (PCA) of relative gene expression in the brain (blue), optic lobe (orange), Malpighian tubules (green), and haemolymph (yellow). Within-tissue normalization was performed relative to the mean of control. Different letters indicate significant differences. In the PCA, each point represents one individual sample and ellipses contain 95% of the group. PCA plot is illustrated on the two principal axes, explaining 78.98% of the variability.

Figure 3: Linear discriminant analysis of transcriptional response to ALAN. The analysis was carried out under four different light treatments: 0 lx (black), 2 lx (dark grey), 5 lx (light grey), 40 lx (yellow); sampled in the brain (A), optic lobe (B), Malpighian tubules (C), and haemolymph (D). Each point represents one individual sample while ellipses represent a 95% confidence level. Each plot is illustrated on the two principal axes of the LDA, explaining > 65% of the variability.

Table A1: Sample sizes of individual male *Gryllus bimaculatus* adult crickets used for each tissue and treatment of the experiment. Each column represents a different gene.

Tissue	Treatment	*Rpl18a*	*actin*	*opLW*	*cry2*	*per*	*c-fosB*	*cry1*
Brain	0lx	9	9	8	9	8	9	8
	2lx	8	8	8	8	8	8	8
	5lx	6	6	6	6	6	6	6
	40lx	5	5	5	5	3	5	3
Optic lobe	0lx	4	4	4	3	3	4	2
	2lx	6	6	6	6	6	6	6
	5lx	6	6	6	6	6	6	6
	40lx	6	6	6	5	6	6	6
Malpighian tubules	0lx	14	14	13	14	12	13	12
	2lx	9	9	8	9	9	9	9
	5lx	12	12	10	11	11	11	11
	40lx	12	12	12	12	11	12	12
Haemolymph	0lx	5	5	4	5	4	4	3
	2lx	5	5	5	5	4	5	4
	5lx	5	5	4	4	4	4	4
	40lx	7	7	5	6	6	6	6

Table A2: Sequences of primers used for qPCR of circadian gene expression in adult male *Gryllus bimaculatus* crickets exposed to a pulse of one of four different ALAN intensities.

Gene	Forward	Reverse
Rpl18a	GCTCCGGATTACATCGTTGC	GCCAAATGCCGAAGTTCTTG
actin	CCTGGCATTGCTGATAGGAT	CCTGCTTGGAGATCCACATT
opLW	CGCTCCTACATCCTCGTCTACTC	CGTTCATCTTCTTGGCTTGCT
cry2	TTCGACACGGAAGGTCTGCT	CGGCCAAATGATGCTACCC
per	GTGGAGGAGAACCGGTGTAA	CCAGGTTCCTCTTCAGCAAG
c-fosB	GTTGCCGTCTGAATGGAAAT	GGAGGTTTTGCACCACTTGT
cry1	GGGGCGAATACATCAAAAGA	CGACAATGAGGTGGAGGATT

Figure S1: Spectrogram of the light bulb used for acclimation and for the experimental pulses. The light spectrum was recorded using the Sekonic Sprectromaster C-700 (North White Plans, NY, USA).