Complex aneuploidy triggers autophagy and p53-mediated apoptosis and impairs the second lineage segregation in human preimplantation embryos

Marius Regin[1], Yingnan Lei[1], Edouard Couvreu De Deckersberg[1], Yves Guns[2], Pieter Verdyck[3], Greta Verheyen[2], Hilde Van de Velde[2,4], Karen Sermon[1,#], Claudia Spits[1,#,*]

[1]Research Group Reproduction and Genetics, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussel, Laarbeeklaan 103, 1090 Jette, Brussels, Belgium

[2]Brussels IVF, Universitair Ziekenhuis Brussel, Laarbeeklaan 101, 1090 Jette, Brussels, Belgium

[3]Vrije Universiteit Brussel (VUB), Universitair Ziekenhuis Brussel (UZ Brussel), Clinical Sciences, Research Group Reproduction and Genetics, Centre for Medical Genetics, Brussels, Belgium

[4]Research Group Reproduction and Immunology, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussels, Laarbeeklaan 103, 1090 Jette, Brussels, Belgium

[#]These authors contributed equally to this work.

*Corresponding author: Claudia.Spits@vub.be

ABSTRACT

Aneuploidy is present in about 80% of human preimplantation embryos and is thought to largely explain our relatively low fertility. However, the percentage of aneuploid cells decreases progressively during the early cleavages and chromosomally mosaic human embryos have been shown to result in healthy new-borns with normal karyotypes. These observations imply a selective loss of aneuploid cells, but the underlying mechanisms are not yet fully understood in human embryos. From studies in mouse embryos and human 2D gastruloids we know that aneuploid cells are eliminated from the epiblast by apoptosis and rather tolerated in the trophectoderm and that aneuploid cells show proteotoxic stress, autophagy and p53 activation. Here, we show that, complex aneuploid human blastocysts present proteotoxic stress, autophagy, deregulated p53 signalling and apoptosis in the trophectoderm and show impaired development of the epiblast and the primitive endoderm. Our findings explain why fully aneuploid embryos fail to further develop and shed light on the mechanisms by which aneuploid cells are removed from mosaic human embryos.

Introduction

Aneuploidy is common in human preimplantation embryos from both natural (Munné *et al.*, 2020a) and medically assisted reproduction cycles (Voullaire *et al.*, 2000; Wells and Delhanty, 2000; Vanneste *et al.*, 2009; Johnson *et al.*, 2010; Chavez *et al.*, 2012; Mertzanidou *et al.*, 2013b, 2013a; Chow *et al.*, 2014; Fragouli *et al.*, 2019; Popovic *et al.*, 2020; Starostik *et al.*, 2020; Capalbo *et al.*, 2021; Yang *et al.*, 2021). Depending on the study, up to 80% of human cleavage stage embryos are found to be aneuploid, of which 70% show chromosomal mosaicism, i.e. the presence of at least two cell lineages with different genomic content (Regin *et al.*, 2022). The reasons behind this high rate of chromosomal abnormalities of mitotic origin is still largely unknown (McCoy, 2017) but could be a combination of a weak spindle assembly checkpoint (SAC) that is uncoupled of apoptosis until the blastocyst stage (Jacobs *et al.*, 2017; Vázquez-Diez *et al.*, 2019), in combination with an abundancy of transcripts of anti-apoptotic genes in cleavage-stage embryos (Yan *et al.*, 2013). After embryonic genome activation, the rapidly increasing expression of pro-apoptotic genes along with the full functionality of the SAC establishes the control of mitotic errors.

It is by now well established that chromosomally mosaic embryos can result in healthy new-borns with normal karyotypes, implying a progressive and selective loss of aneuploid cells during development (Greco *et al.*, 2015; Fragouli *et al.*, 2017, 2019; Munné *et al.*, 2017, 2020b; Capalbo *et al.*, 2021; Viotti *et al.*, 2021; Yang *et al.*, 2021). In the preimplantation embryo, several studies have shown that the proportion of aneuploid cells within the embryo decreases three days post fertilization (3dpf) (van Echten-Arends *et al.*, 2011; Fragouli *et al.*, 2019; Yang *et al.*, 2021) with no apparent preferential allocation of aneuploid cells during the first lineage segregation

event to either trophectoderm (TE) or inner cell mass (ICM) (Capalbo *et al.*, 2013;

Popovic *et al.*, 2018; Starostik *et al.*, 2020; Ren *et al.*, 2022). However, aneuploid

cells can be excluded as cell debris (Orvieto *et al.*, 2020) or allocated to the

blastocoel cavity and to peripheral cells that do not participate in the formation of the

embryo (Griffin *et al.*, 2022). In some studies, (Starostik *et al.*, 2020; Griffin *et al.*,

2022) the TE contained a slightly higher number of aneuploid cells than the ICM,

without reaching statistical significance. In their landmark paper, (Bolton *et al.*, 2016)

found a quite similar but statistically significant 6% TE enrichment in mouse embryos

treated with the SAC inhibitor reversine to induce mosaic aneuploidy. This small

enrichment fell in the same confidence interval as the Starostik study, therefore TE

enrichment cannot be ruled out in the human embryo. After implantation, aneuploid

cells become progressively more frequent in the trophoblast lineage which is derived

from TE cells as compared to the epiblast (EPI) and primitive endoderm (PrE)

lineages which are derived from the ICM cells during the second lineage segregation

(Starostik *et al.*, 2020). This suggests that the trophoblast cells are more tolerant to

aneuploidy, in line with the rate of 1-2% of confined placental mosaicism observed in

ongoing pregnancies in the general population (Kalousek and Vekemans, 1996) and

with the presence of aneuploid cells in the placental lineages only (Zamani Esteki et

al., 2019). This is further supported by a recent study showing that genetic

mosaicism is a normal feature of the placenta and is due to genetic bottlenecks that

occur early in development (Coorens *et al.*, 2021). Taken together, strong evidence

supports the selective elimination of aneuploid cells from the embryonic lineage in

the pre- and post-implantation development, while the placenta appears to be more

tolerant to genetic imbalances.

From a mechanistic point of view, this selective clearance is thought to be mediated

by two intracellular stress responses that have been particularly well studied in

cancer aneuploidy and in yeast models (reviewed in Santaguida and Amon, 2015;

Zhu et al., 2018). First, aneuploidy results in gene-dosage defects, leading to an

unbalanced protein pool and proteotoxic stress (Oromendia *et al.*, 2012; Ohashi *et*

al., 2015). If this remains unresolved by activation of autophagy, CASP8/LC3B/p62

will interact to activate the apoptosis cascade (Pan *et al.*, 2011; Stingele *et al.*, 2012;

Dürrbaum *et al.*, 2014). Second, aneuploidy results in replication stress that can lead

to DNA-damage, resulting in *p53* activation and ultimately to apoptosis (Zhu *et al.*,

2018; Regin *et al.*, 2022). Bolton *et al.* (Bolton *et al.*, 2016) showed that aneuploid

cells are selectively cleared from chimeric mouse embryos generated from control

blastomeres mixed with aneuploid blastomeres obtained using a treatment with the

SAC inhibitor reversine. Later work by the same group showed that apoptosis of

these aneuploid cells requires autophagy and *p53* activation and that euploid cells in

the mouse embryo compensate this cell loss by increased proliferation (Singla *et al.*,

2020). The authors proposed that the proteotoxic stress in these aneuploid chimeric

mouse embryos activated the p53 pathway, which in turn activated autophagy and

apoptosis. In a recent study using human 2D gastruloids with reversine-induced

aneuploidy, apoptosis was only induced in differentiating but not in pluripotent cells,

although aneuploidy induced an increase in total nuclear p53 proteins (Yang *et al.*,

2021). The apoptotic response to aneuploidy also seems to be lineage-specific. In

the mouse, aneuploid cells in the ICM are eliminated by apoptosis while they are

rather tolerated in the TE (Bolton *et al.*, 2016). Furthermore, in a follow-up study,

(Singla *et al.*, 2020) demonstrated that during the peri-implantation period, aneuploid

cells were eliminated from the EPI and the PrE in similar rates, and that the EPI-

specific elimination of aneuploid cells by apoptosis is coupled to increased

proliferation rates of euploid cells. Similarly, in human gastruloids, aneuploid cells are

eliminated by apoptosis from the embryonic germ layers but are tolerated in the TE-

like cells (Yang *et al.*, 2021).

The type of aneuploidy and the percentage of aneuploid cells within a

euploid/aneuploid mosaic embryo determine its developmental capacity. The viability

of mouse embryos inversely correlates to the proportion of aneuploid cells, with 50%

of euploid cells being sufficient to sustain normal development (Bolton *et al.*, 2016).

This same threshold appears to apply to human embryos, where embryos with less

than 50% of aneuploid cells have higher developmental potential than those

containing more than 50%, albeit both performing worse than fully euploid embryos

(Viotti et al., 2021, Capalbo et al., 2021). Conversely, whole chromosome gains and

losses that are present in every cell of the human embryo are usually linked to

lethality with the notable exception of trisomies 13 (Patau syndrome), 18 (Edwards

syndrome) and 21 (Down syndrome), and sex chromosome abnormalities. Shahbazi

et al. recently described the consequences of carrying specific meiotic aneuploidies

on human embryo development using an *in vitro* implantation model (Shahbazi *et al.*,

2020). They showed that human embryos with trisomy 15, 16 and 21 and monosomy

21 readily all reached the blastocyst stage, and that monosomy 21 embryos arrested

after implantation while trisomy 15 and 21 embryos developed further. Interestingly,

trisomy 16 embryos showed a hypoproliferative trophoblast due to excessive *CDH1*

expression, a gene located on chromosome 16.

In this study, we investigated the cellular consequences of aneuploidy during human

preimplantation development, with a focus on stress response and lineage

segregation events. We studied embryos with complex meiotic aneuploidy rather

than mosaic embryos to ascertain that our observations were significant.

Methods

Ethical approval. All experiments have been approved by the local Commission of Medical Ethics of the UZ Brussel (B.U.N. 143201628722) and the Federal Committee for Medical and Scientific Research on Human Embryos *in vitro* (AdV069 and AdV091). Patients from Brussels IVF (UZ Brussel) donated their embryos for research after written informed consent.

Culture of human pre-implantation embryos. We warmed vitrified human blastocysts (5-6dpf) after PGT or at 3dpf using the Vitrification Thaw Kit (Vit Kit-Thaw; Irvine Scientific, USA) following manufacturer's instructions. Embryos were graded before vitrification according to Gardner and Schoolcraft criteria (Gardner and Schoolcraft, 1999). Post PGT-embryos were left to recover for 2h in 25 µL droplets of Origio Sequential Blast^{TM} medium (Origio) at 37°C with 5% O_2, 6% CO_2 and 89% N_2, after which they underwent a second TE biopsy (Fig. 1a). For the reversine experiment, we cultured the 3dpf embryos for 24h in the same medium supplemented with 0.5 µM reversine (Stem Cell Technologies) and after wash-out we let them develop until 5dpf and subsequently took a biopsy (Fig. 1b). The PGT embryos used for the second experiment series (Fig. 4a) were cultured for 16h after warming to allow them to progress to the second lineage. Prior to fixation we live-stained all the embryos with either Caspase-3/7 or Caspase-8 (Supplementary Table 30) for 30 min and subsequently removed the zona pellucida using Acidic Tyrode's Solution (Sigma-Aldrich) for 30 sec – 2 min and subsequently washed in PBS/2%BSA.

Biopsy procedure. The zona pellucida was opened (15-25µm) on 4dpf using a Saturn Laser (Cooper Surgical). For PGT cycles, expanded good quality embryos

were biopsied on 5dpf; early blastocysts were evaluated again for expansion on 6dpf.

Briefly, embryos were transferred to HEPES buffered medium and biopsied on a

Nikon inversed microscope equipped with a Saturn Laser. The herniated TE cells

were then aspirated and cut using manual laser shoots. In case of sticking cells this

was supported by mechanical cutting. After successful tubing, the blastocysts were

vitrified using CBS-VIT High-Security straws (CryoBioSystem, L'Aigle, France) with

dimethylsulphoxide-ethylene glycol-sucrose as the cryoprotectants (Irvine Scientific

Freeze Kit, Newtownmountkennedy, County Wicklow, Ireland) (Van Landuyt *et al.*,

2011). The TE biopsy for RNA-sequencing (Fig. 1a,b) was performed with the same

procedure, cells were biopsied form the same opening in the zona pellucida.

PGT. During PGT we analyzed a TE sample with multiple cells for each embryo.

Next Generation Sequencing: The chromosomal analysis was performed by WGA

(Sureplex, Illumina) followed by library preparation (KAPA HyperPlus, Roche),

sequencing on a NovaSeq (Illumina) and an *in-house* developed interpretation

pipeline. The analysis has an effective resolution of 5 Mb.

SNP-array: We relied on whole genome amplification (MDA, Repli-G) followed by

genome wide SNP array using Karyomapping (Vitrolife) with Bluefuse software

(Vitrolife). In addition, SNP array data were analyzed with an *in-house* developed

interpretation pipeline for aneuploidy detection.

RNA-sequencing. Multiple (5-10) TE cells were used as input to generate full length

cDNA with the SMART-Seq^{TM} v4 Ultra^{TM} Low Input RNA Kit (Clontech Laboratories,

Inc.) according to the manufacturer's instructions. The quality of cDNA was checked

using the AATI Fragment Analyzer (Advances Analytical). Library preparation was

performed using the KAPA HyperPlus Library Preparation Kit (KAPA Biosystems)

according to the manufacturer's instructions. During cDNA synthesis and library

preparation we used 17 and 11 PCR cycles, respectively. Sequencing was

performed on a NovaSeq 6000 (Illumina) with 25 million reads per sample.

Bioinformatics analysis. Reads were trimmed using cutadapt version 1.11 to

remove the "QuantSEQ FWD" adaptor sequence. We checked the quality of the

reads using the FastQC algorithm (Love *et al.*, 2014). Count tables were generated

using STAR (Dobin *et al.*, 2013) (version 2.5.3) through mapping against the

Genome Reference Consortium Human Build 38 patch release 10 (GRCh38.p10)

combined with a general transfer format (GTF) file, both downloaded from the

Ensembl database. The RNA-sequencing by Expectation Maximization (RSEM) (Li

and Dewey, 2011) software (version 1.3.0) was used to produce the count tables.

Differential gene expression analysis was performed using R-studio (Version

1.1.456) with the edgeR (Robinson *et al.*, 2009) and limma (Ritchie *et al.*, 2015)

packages. We included genes with a count per million greater than 1 in at least two

samples. The trimmed mean of M values (Robinson and Oshlack, 2010) algorithm

was used for normalization. The normalized counts were then transformed in a $\log_2$

fold-change ($\log_2$FC) table with their associated statistics. In each comparison, genes

with a $| \log_2$FC $| > 1$ and an p-value < 0.05 were considered as significantly

differentially expressed.

Gene set enrichment analysis (GSEA) was performed using the 'investigate gene

sets' function on the website of the Molecular Signature Database (http://www.gsea-

msigdb.org/gsea/msigdb/annotate.jsp). We used all differentially expressed genes

(Supplementary Tables 2, 3) as input and investigated Hallmark, C2 and C5 gene

sets with FDR-q values < 0.05. For the analysis of gene dosage effects, we could

only use the TOP500 differentially expressed genes by p-value since the Molecular

Signature Database changed the cut-off during the period of our data analysis (raw

data, Supplementary Tables 26, 27). In the figures 1f and g we used a supervised

approach and sorted the significantly deregulated gene sets by the total amount of

genes in overlap. Figure 2f was also supervised but the gene sets were sorted by

FDR-q value.

Ingenuity Pathway analysis (IPA, Qiagen Inc.) was used to investigate activated,

deregulated and inhibited upstream regulators and canonical pathways. As for the

GSEA we used all differentially expressed genes of our two test sets with their

respective log_2FC and p-value. For the upstream regulators IPA computes z-scores

with associated p-values that are considered significant when $p < 0.05$. If significant,

a z-score > 2 means activated, a z-score < 2 means inhibited and in between means

deregulated. For the canonical pathways IPA computes z-scores with associated p-

values that are considered significant when $p < 0.05$. If significant, we considered a

z-score > 1.5 as activated, a z-score < 1.5 as inhibited and in between as

deregulated. We generated all figures in a supervised approach.

Transcription factor analysis of all differentially expressed genes was performed

using Enrichr (https://maayanlab.cloud/Enrichr/). We selected the outputs from ChEA

2016 and ENCODE ChEA consensus and considered those transcription factors as

significant with a p-value < 0.05.

The copy number variation (CNV) analysis was inferred from our RNA-sequencing

data set using *inferCNV* R package (version 1.7.1) (Inferring copy number alterations

from   tumor   single   cell   RNA-Seq   data

(https://github.com/broadinstitute/inferCNV/wiki"). Our count table was used as input

and compared to the reference set which in our case was the expression data of the

euploid embryos. The analysis followed the standard workflow in inferCNV with the

following parameters: "denoise" mode, value of 1 was used for the "cutoff", prediction

of CNV states using default Hidden Markov Models (HMM) method,

"cluster_by_groups" was set to TRUE, and other values were set by default.

All figures were generated using R-studio (Version 1.1.456).

Gene dosage analysis. We calculated the total number of loci with imbalanced gene

expression based on the genetic abnormalities identified during the PGT. For this, we

used the number of coding loci per chromosomal region as listed in the ENSMBL

database (ensemble.org) and included both gains and losses, either mosaic or

homogeneously present, and both full chromosomal aneuploidy as well as segmental

abnormalities. The embryos were ranked based on the total number of imbalanced

loci and separated into a low- and a high-imbalanced gene dosage group. The low-

dosage group contained the embryos with the lowest 50^{th} percentile in number of

unbalanced loci, the high-dosage group the embryos with 50^{th} percentile or higher

(Supplementary Table 25, 29).

Immunocytochemistry. Embryos were fixed using 4% paraformaldehyde (Sigma-

Aldrich) for 10 min, washed three times in PBS/2%BSA for 5 min and then

permeabilized using 0.1% Triton X-100 (Sigma-Aldrich) for 20 min followed by

another washing cycle. We used 10% FBS supplemented with 2% BSA for 1h to

block non-specific protein binding. The embryos were then incubated overnight with

primary antibodies (Supplementary Table 30) diluted in blocking buffer at 4°C. The

next day, they underwent another washing cycle using PBS/2%BSA and were then

incubated with secondary antibodies (1:200, Supplementary Table 26) for 1h at room

temperature. After washing, nuclei were stained using Hoechst 33342 (5µg/mL, Life

Technologies) diluted in PBS/2%BSA for 15 min. For imaging we mounted the

embryos on poly-L-lysine coated glass slides (Sigma-Aldrich) in 3µL of PBS. To

avoid flattening of the embryos we used Secure-Seal^{TM} Spacers (9mm diameter,

0.12mm deep, Thermofisher) before putting the coverslips in place. For the

experiments that required re-staining (Fig. 4a) we imaged the embryos in 3µL

PBS/2%BSA in 18 well µ-Slides (Ibidi) and subsequently recuperated them. After

recuperation we photobleached the embryos stained for CASP3/7 by using maximum

laser power of the 488 nm channel and maximum size of the pinhole for 10 min. The

embryos were then re-stained using rat-anti-GATA4 primary antibody and the

matching Alexa Fluor 488 secondary antibody (Supplementary Table 30) (to replace

the CASP3/7 staining).

Imaging and image analysis. Confocal imaging was performed using the LSM800

(Zeiss). Z-stacks were taken with a LD C-Apochromat 40x/1.1 NA water immersion

objective in optical sections of 1µm. Nuclei were counted using Arivis Vision 4D

(3.4.0). All other measurements were performed using the Zen 2 (blue edition)

software based on the optical sections and/or orthogonal projections.

Statistics. The type of statistical test and p-values are mentioned in the figure

legends. For experiments that contained groups with small sample sizes we used

non-parametric tests (Mann-Whitney or Kruskal-Wallis test). Fisher-exact test was

used to determine the dependency of the ploidy status on the presence or absence

of primitive endoderm. Otherwise we used Student's t-test with Welch's correction

when we assumed normal distribution and unequal standard deviations. We used

GraphPad Prism 9.0.0 or R-studio for statistical testing. The trend analysis regarding

gene dosage effects was performed with SPSS using the Jonkheere-Terpstra test.

Results

Human embryos with meiotic complex aneuploidy show transcriptional signatures of a deregulated p53 pathway and apoptosis

We first tested the hypothesis that chromosomally unbalanced cells are cleared during human preimplantation development by apoptosis, mediated by both sustained p62-LC3B accumulation as a marker for autophagy and DNA-damage-mediated p53 activation. Fifty cryopreserved human blastocysts (5 or 6dpf) that had been previously diagnosed by Preimplantation Genetic Testing (PGT) as either euploid (n=28) or containing non-mosaic complex aneuploidy of at least two whole chromosomes and assumed of meiotic origin (except for one)(n=22, further referred to as aneuploid, Fig. 1a, Supplementary Table 1) underwent a second TE biopsy to collect cells for RNA sequencing. Eleven warmed human cleavage stage embryos that were treated with 0.5 µM reversine from 3dpf to 4dpf and further developed to the blastocyst stage (referred to as reversine-treated embryos, Fig. 1b) were also biopsied (on 5dpf or 6dpf) to serve as a comparison with the results obtained in the mouse. The concentration of 0.5 µM reversine was chosen based on previous experiments in mouse embryos (Bolton *et al.*, 2016; Singla *et al.*, 2020). Of these, 14 euploid, 20 aneuploid and 11 reversine-treated embryos (total=45) yielded good quality RNA-sequencing results.

Unsupervised hierarchical clustering and principal component analysis using all expressed genes revealed no clustering of the euploid, aneuploid or reversine-treated embryos in distinct groups (Fig. 1c, Supplementary Fig. 1a). We used InferCNV (https://github.com/broadinstitute/inferCNV/wiki") to bioinformatically infer unbalanced chromosomal copy numbers from gene expression data, based on

higher or lower expression of genes located on chromosomes with gains or losses

respectively (Fig. 1d, Supplementary Fig. 1b, Supplementary Table 1). We compared

the outcomes to the results obtained during PGT and found a match for 45/48 of the

full chromosome aneuploidies, while euploidy was correctly predicted in all cases

(Supplementary Fig. 1b). These results show that aneuploidy in human blastocysts

results in significant abnormal gene dosage effects. The exact karyotypes of the

reversine-treated embryos could not be determined since reversine induces different

aneuploidies in each cell, leading to mosaic TE biopsies. However, InferCNV allowed

us to determine a trisomy 12 meiotic chromosomal abnormality in one reversine-

treated embryo (Supplementary Figure 1c), confirming that the reversine treatment

causes random aneuploidy while naturally occurring meiotic aneuploidies are readily

determined.

Despite the abnormal gene dosage, and in line with the lack of clear clustering in the

PCA, the differential gene expression analysis yielded very few genes with a false

discovery rate (FDR-q-value) under 0.05. These findings are similar to those of other

studies on the transcriptome of aneuploid human embryos (Groff *et al.*, 2019;

Maxwell *et al.*, 2022). Therefore, we used the differential gene expression with a

cutoff value of |log$_2$ fold change|>1 and -log$_{10}$(p-value)<0.05, and primarily focused

on pathway analysis. We found 547 upregulated and 463 downregulated genes in

aneuploid versus euploid embryos (Fig. 1e, Supplementary Table 2). Amongst the

top upregulated genes was Growth Differentiation Factor 15 (*GDF15*), a gene that

has been found to be upregulated in human aneuploid cells (Dürrbaum *et al.*, 2014)

including aneuploid human embryos (Starostik *et al.*, 2020). When comparing

reversine-treated versus euploid embryos we observed 539 upregulated and 355

downregulated genes (Fig. 1f, Supplementary Table 3). While only 9.6% of the

differentially expressed genes are in common between the aneuploid and reversine-

treated embryos (Fig. 1g, h, Supplementary Table 4), pathway analysis revealed

several commonly deregulated pathways. We identified deregulation of the p53

pathway and apoptosis (FDR q-value < 0.05, Fig. 1i, Supplementary Tables 5, 6) and

deregulation of cell population proliferation, apoptotic process, regulation of cell

death, and cell cycle (Fig. 1j, Supplementary Tables 7, 8). The prediction of p53

activation is also supported by differential expression of several p53 target genes

(Jaiswal *et al.*, 2021), such as *MDM2* and *PPM1D* (feedback); *CDKN1A, BTG2,*

GADD45A (cell-cycle arrest); *TNFRSF10B* (apoptosis), *DDB2* and *RRM2B* (DNA-

repair); *FDXR* (metabolism) and *SESN1* (translation control) (Supplementary Table

2). Other pathways of interest related to DNA damage and p53, such as cellular

response to DNA damage stimulus and DNA damage response signal transduction

by p53 class mediator were specific to the aneuploid group. Cellular senescence

signatures in the aneuploid embryos and selective autophagy in the reversine-treated

group were also deregulated (Supplementary Tables 9, 10). Ingenuity pathway

analysis (IPA) showed that upstream regulators *CDK19* and *AURK* (cell cycle) and

GAS2L3 and *ANLN* (cell growth) were inhibited in the aneuploid group (Fig. 1k,

Supplementary Table 11). In line with the GSEA analysis, *p53* was the most

activated upstream regulator. Reversine-treated embryos activated *ATM* which is a

well-known sensor of DNA-damage that mediates target genes such as *p53* (Banin *et*

al., 1998) (Fig. 1l, Supplementary Table 12). Canonical pathway analysis by IPA

confirmed that p53 signalling, autophagy, cell-cycle pathways and senescence are

deregulated in aneuploid embryos (Fig. 1m, Supplementary Table 13). The TE of

reversine-treated embryos showed activated senescence, similar to the mouse

reversine-treated embryos (Bolton *et al.*, 2016) (Fig. 1n, Supplementary Table 14).

Transcription factor analysis identified significant *p53* expression in both aneuploid

and reversine-treated groups (Supplementary Tables 15, 16).

We next compared our set of differentially expressed genes to those identified by

other studies investigating the gene expression of euploid and aneuploid embryos,

namely: mosaic monosomies 4, 7, 10, 15, 19 and mosaic trisomies 15 and 19

(Maxwell *et al.*, 2022), trisomy 16 and monosomy 16 (Fuchs Weizman *et al.*, 2019),

monosomy 21 (Sanchez-Ribas *et al.*, 2019) and embryos that contained various

types of chromosomal abnormalities (Vera-Rodriguez *et al.*, 2015; Licciardi *et al.*,

2018; Starostik *et al.*, 2020) (detailed overview in Supplementary Table 17). Overall,

we found modest overlaps (Supplementary Figure 1d,e, Supplementary Tables 18-

20). Most commonalities were with the data of Starostik et al, where we found an

overlap in 87 genes (2.3%) (Starostik *et al.*, 2020). These included target genes of

the p53 pathway such as *MDM2* and *CDKN1A* (Supplementary Fig. 1d,

Supplementary Table 20). Interestingly, GSEA of these 87 genes identified an

enrichment of intrinsic apoptotic signalling pathway by p53 class mediator and the

apoptotic signalling pathway, in line with the findings of our dataset (Supplementary

Table 19).

High gene-dosage imbalances result in stronger cellular stress signatures in
human embryos

To test if there was a relationship between the level of stress signatures and the size

of chromosomal imbalances in embryos as previously described in cancer (Dürrbaum

and Storchová, 2016), we separated the aneuploid group into a low- and a high-

imbalanced gene dosage group, based on the total number of coding loci that were

affected by the chromosomal abnormalities. The low-dosage group contained the

embryos under the 50th percentile when ranking the embryos from the lowest to the

highest number of unbalanced loci, the high-dosage group the embryos with 50th

percentile or higher (Supplementary Table 21). Hierarchical clustering using the top

500 deregulated genes showed that all high-dosage embryos clustered together and

separated from the euploid embryos, while the low-dosage embryos clustered with

either group (Fig. 2a). Differential gene expression with |log$_2$ fold change|>1 and -

log$_{10}$(p-value)<0.05 resulted in 404 upregulated and 340 downregulated genes in

low-dosage versus euploid embryos and in 716 upregulated and 627 downregulated

genes in high-dosage versus euploid embryos with an overlap of 104 (10%) between

both aneuploid groups (Fig. 2b-e, Supplementary Tables 22,23). We identified

deregulation of TNFA signalling via NFkB, p53 pathway and apoptosis for both

groups (FDR q-value < 0.05, Fig. 2f, Supplementary Tables 24,25). High-dosage

embryos additionally showed an unfolded protein response, suggesting that this

signature may only be present in the embryos with more and/or larger abnormal

chromosomes. In the high-dosage group more of the genes of the p53 and apoptosis

pathways were deregulated, and with higher significance (Fig. 2g,h, Supplementary

Tables 26,27). The gene expression levels in the p53 and apoptosis pathways show

a stepwise distribution between the gene dosage groups (Fig. 2i).

Aneuploidy triggers proteotoxic stress, autophagy and apoptosis in human

embryos

To further investigate the cellular stress responses identified by the transcriptomic

analysis, we immunostained the same embryos that underwent TE biopsy for

transcriptome analysis. Thirty-seven (18 euploid, 19 aneuploid) of the 50 PGT

embryos and 8 of the 11 reversine-treated embryos were successfully stained for

markers of apoptosis, proteotoxic stress and autophagy (Fig. 3a,b). Aneuploid

embryos showed increased intensity levels of active CASP3/7 (13 euploid and 8

aneuploid embryos stained) and CASP8 (6 euploid, 9 aneuploid and 8 reversine-

treated embryos stained) (Fig. 3c-f). Both aneuploid and reversine-treated embryos

had lower cell numbers than euploid embryos (Fig. 3g). We investigated whether

proteotoxic stress and autophagy (Oromendia *et al.*, 2012; Stingele *et al.*, 2012;

Dürrbaum *et al.*, 2014; Ohashi *et al.*, 2015; Singla *et al.*, 2020) were also present in

aneuploid human embryos. The number of puncta per cell of autophagy markers

LC3B (19 euploid, 22 aneuploid and 8 reversine-treated embryos stained) and p62 (6

euploid and 9 aneuploid embryos stained) and the intensity levels of proteotoxic

stress marker HSP70 (13 euploid, 8 aneuploid and 8 reversine-treated embryos

stained) were significantly increased in both aneuploid and reversine-treated

embryos (Fig. 3h-k). The gene expression profiles suggest that the accumulation of

autophagic proteins in aneuploid embryos is caused by increased autophagic flux

due to differential expression of the *p53* target gene DNA Damage Regulated

Autophagy Modulator-1 (*DRAM1*), rather than by inhibition of autophagy

(Supplementary Table 2). DRAM1 is located in the lysosomal membrane and it

increases autophagic flux and apoptosis after stress-induced p53 activation in cancer

cells (Crighton *et al.*, 2006; Guan *et al.*, 2015).

Aneuploidy increases apoptosis in the trophectoderm and impairs the second

lineage segregation in the inner cell mass

We further investigated if these effects are cell-type specific, as shown in the mouse

(Bolton *et al.,* 2016; Singla *et al.*, 2020) and human 2D gastruloids (Yang *et al*.,

2021), and how they affect the first lineage segregation event to TE and ICM, and the

second lineage segregation event to EPI and PrE.

We co-stained a new batch of expanded blastocysts (5dpf or 6dpf) for OCT4 (10

euploid, 14 aneuploid, ICM/EPI) and CASP3/7 (10 euploid, 14 aneuploid), LC3B (4

euploid, 5 aneuploid) or Serine 15 p53 (3 euploid, Aneuploid embryos showed a

higher percentage of cells with activated p53 5 aneuploid) (Fig. 4a). All information

on embryo score and developmental day and stage can be found in Supplementary

Table 28. The latter marker was chosen to demonstrate that the transcriptomic

signature of p53 signalling is caused by DNA-damage-induced phosphorylation of

Serine-15 on p53 (Loughery *et al.*, 2014). Aneuploid embryos showed a higher

percentage of cells with activated p53 (Fig. 4b,c), as suggested by our transcriptomic

data (Figure 1), although these differences did not reach significance when

considering the ICM/EPI (OCT4-positive) cells or TE (OCT4-negative) separately

(Supplementary Fig. 2a-c). We then investigated cell type specificity of proliferative

defects and apoptosis. Both the aneuploid TE and OCT4-positive cells showed

decreased cell numbers (Fig. 4d-f) that led to an overall decrease of cell numbers in

whole aneuploid embryos (Fig. 4g), confirming our previous experiment (Fig. 3g).

The observed increased percentage of nuclear CASP3/7 positive cells in the TE of

aneuploid embryos (Fig. 4h,i) and increased CASP3/7 mean intensity per cell in

whole aneuploid embryos (Fig. 4h,j) points to apoptosis as the cause of this decrease

in cell numbers. However, no OCT4-positive cells of either euploid or aneuploid

group showed nuclear CASP3/7 signal (Fig. 4h). We also found CASP3/7-positive

micronuclei which were increased in number in aneuploid embryos (Fig. 4h,

Supplementary Fig. 2d,e).

The observation (figure 4f) that the ICM/EPI cell numbers of aneuploid embryos fell

into two groups with either low or high cell counts led us to further investigate the

impact of aneuploidies on the second lineage segregation event. All blastocysts were

re-stained for GATA4 (PrE marker) and imaged together with OCT4 (ICM/EPI

marker). We found that while all euploid embryos contained GATA4-positive cells,

7/14 aneuploid embryos had no GATA4-positive cells (Fig. 4k,l). Further, in those

seven aneuploid embryos with a GATA4-positive PrE cells, we observed a

significantly lower PrE/EPI ratio (Fig. 4m). We then investigated whether there are

different levels of LC3B as marker of autophagy between the aneuploid TE and ICM

in an additional 5 aneuploid and 4 euploid embryos. Aneuploid embryos displayed

increased levels of autophagy mainly attributable to the TE (Fig. 4n-q), while their

ICM/EPI cells showed a small increase in autophagy which did not reach significance

(Fig. 4q). The aneuploid embryos could be separated in two groups: a first group with

low levels of autophagy lacking GATA4-positive cells in 2 out of 3 embryos (Fig. 4n,

left and right panels, 2^{nd} row) and a high-level autophagy group in which both

embryos contained GATA4-positive cells (Fig. 4n, left and right panels 3^{rd} row).

Lastly, we separated the aneuploid embryos into a low- and a high-dosage group as

done for the gene-expression analysis (Supplementary Table 29). Trend analysis

showed that the higher the number of imbalanced loci, the lower the cell numbers in

the TE and PrE but not in the EPI (Fig. 4r).

Discussion

In this work, we studied the consequences of aneuploidy in human preimplantation

embryos that carried a uniformly present, presumed meiotic, complex aneuploidy.

We selected this type of genetic imbalances because we hypothesized that in the

embryo, these would result into stronger and potentially different cellular stress

responses than single meiotic aneuploidy or mosaic aneuploidy. In our study we find

that, while aneuploid embryos had lower cell numbers in both TE and ICM/EPI, the

effect of aneuploidy was cell-type dependent. Aneuploid TE presented transcriptomic

signatures of p53 signalling and apoptosis, and immunostaining of whole embryos

showed that aneuploidy results in increased proteotoxic stress, autophagy, DNA

damage-mediated activated p53 and subsequent apoptosis in the TE. In contrast,

aneuploidy impaired the second lineage differentiation, in particular PrE formation.

Overall, our gene-expression results are in line with previously published RNA-

sequencing studies on aneuploid human embryonic cells that identified

transcriptional signatures consistent with DNA damage (Vera-Rodriguez *et al.*, 2015;

Licciardi *et al.*, 2018; Fuchs Weizman *et al.*, 2019), alteration of cell proliferation

(Starostik *et al.*, 2020; Maxwell *et al.*, 2022), the cell cycle (Vera-Rodriguez *et al.*,

2015; Licciardi *et al.*, 2018; Fuchs Weizman *et al.*, 2019; Starostik *et al.*, 2020),

deregulation of autophagy (Licciardi *et al.*, 2018; Sanchez-Ribas *et al.*, 2019), p53

signalling (Licciardi *et al.*, 2018) and apoptosis (Licciardi *et al.*, 2018; Groff *et al.*,

2019; Sanchez-Ribas *et al.*, 2019; Maxwell *et al.*, 2022). However, when comparing

our dataset of differentially expressed genes to these studies, we only found modest

overlaps at the level of individual genes. This is probably because embryos of

different developmental stages were included in the studies, containing specific

aneuploidies, and of different sample types (overview in Supplementary Table 17).

Naturally acquired and induced aneuploidies only share a few differentially expressed

genes but show overall equal signatures after pathway analysis, suggesting that

reversine-based models of aneuploidy are very close but may not fully mimic

endogenous aneuploidy. We also observed aneuploidy-mediated gene dosage

effects in the TE cells, as extensively reported in human and mouse aneuploid

cancer cell lines (Dürrbaum and Storchová, 2016), and found that a higher dosage

leads to stronger p53 signalling and apoptosis responses.

We next tested the predictions made based on the gene-expression profiles with

immunostaining. First, we found evidence of proteotoxic stress and increased

expression of autophagy proteins, as well as activation of apoptosis. The most likely

series of events is that p62/LC3B/CASP8 interaction results in autophagy-mediated

apoptosis in aneuploid human embryos. Nevertheless, we cannot rule out indirect

activation of apoptosis mediated by autophagy due to the degradation of cell

organelles (Gump and Thorburn, 2011). This is supported by the observation in the

aneuploid mouse embryo model that autophagy is independent of the mTOR

pathway (Singla *et al.*, 2020). Other upstream regulators of autophagy could be

involved, such as the cGAS-STING pathway (Krivega *et al.*, 2021) or, as also

observed in our dataset, p53-mediated activation of DRAM1 (Crighton *et al.*, 2006)

that can induce apoptosis by increasing BAX protein levels (Guan *et al.*, 2015).

Second, we confirmed the presence of activated *p53* in aneuploid human embryos.

We specifically stained for p53 with phosphorylation on Serine 15, mediated by

kinases such as ATM and ATR as a direct result of DNA-damage (Loughery *et al.*,

2014). This indicates that aneuploidy also induces the DNA-damage response in the

embryo, as found in the mouse and 2D gastruloids (Singla *et al.*, 2020; Yang *et al.*,

2021). An interesting observation is that aneuploid embryos contained increased

numbers of CASP3/7 positive micronuclei. This is similar to findings in a non-human

primate model showing that abnormal chromosomes can be encapsulated in

micronuclei, which can subsequently be selectively eliminated from the embryo

(Daughtry *et al.*, 2019). In human cleavage stage embryos these micronuclei have

been shown to contain genetic material that originates from chromosome breakages

due to replication fork stalling and DNA-damage (Palmerola *et al.*, 2022), further

highlighting how aneuploidy is leading to DNA-damage in the human embryo. In sum,

our results in this part are in line with previous findings on mouse and 2D gastruloid

models, and suggest that at least two different stress responses are activated in fully

aneuploid embryos with complex karyotypes, both converging on apoptosis if not

resolved. First, proteotoxic stress resulting from gene-dosage imbalances mediates

the unfolded protein response and autophagy. Second, DNA-damage mediates p53

activation that likely occurs due to replication stress as occurring in cancer and yeast

(Santaguida and Amon, 2015; Zhu *et al.*, 2018).

In the second part of our work, we focused on studying the lineage-specific response

to complex aneuploidy, as the studies on reversine-treated mouse embryos (Bolton

et al., 2016; Singla *et al.*, 2020) and human 2D gastruloids (Yang *et al.*, 2021) also

identified a cell-type specific effect of aneuploidy. In the mouse, aneuploid cells are

eliminated by apoptosis in the EPI, while tolerated in the TE, and in human 2D

gastruloids, aneuploid cells are eliminated in the post-gastrulation embryonic germ

layers but tolerated in the TE.

In our work, we used 5-6dpf human expanded blastocysts to study the consequences

of aneuploidy on lineage segregation, identifying the ICM/EPI as staining positive for

OCT4 and the PrE as positive for GATA4. We confirmed that the embryos analysed

were expanded blastocysts which had undergone the first lineage segregations, as

OCT4 was restricted to the inner cells and absent in TE cells (Niakan and Eggan,

2013). The identity of human ICM and EPI cells is currently under vigorous debate

(Meistermann *et al.*, 2021) and exclusive markers of either cell type are untested,

therefore we cannot assign ICM or EPI identity to OCT4-positive cells. However, in

blastocysts with GATA4-positive cells the second lineage segregation had occurred

and PrE was established, and consequently we confidently assigned EPI identity to

OCT4-positive cells in these blastocysts. Conversely, in GATA4-negative embryos

OCT4-positive cells could be either ICM or EPI cells. Interestingly, we never found

GATA4-positive cells in the absence of OCT4-positive cells, supporting the

hypothesis that PrE cells differentiate from EPI cells (Meistermann *et al.*, 2021).

While we detected lower cell numbers in aneuploid OCT4-positive ICM/EPI cells,

they did not contain a single CASP3/7 positive cell. This difference with the mouse

results is likely due to species-specific properties of the pluripotent cells: the lack of

apoptosis in the human ICM/EPI could be due to an uncoupling between aneuploidy-

induced stress signals and apoptosis, as observed in human and mouse pluripotent

stem cells (Mantel *et al.*, 2007) . The lower cell numbers could be explained by two

other known effects of *p53* activation. First, *p53* activation decreases cell

proliferation, leading to lower cell numbers. Secondly, *p53* activation may

downregulate *OCT4* and *NANOG* expression, as seen in mouse embryonic stem

cells (Lin *et al.*, 2005), leading to loss of pluripotency and driving the cells to

differentiation and out of the ICM/EPI. It is worth noting that our results are in contrast

to findings of Victor et al, where both TE and ICM/EPI of mosaic and aneuploid

embryos showed increased apoptosis (Victor *et al.*, 2019). Further work will be

needed to understand if these discrepancies are due to technical or biological

factors.

A second difference we observed was that the aneuploid TE had increased levels of

autophagy markers, which did not reach statistical significance in the ICM/EPI.

Interestingly, the aneuploid OCT4-positive results separated into a high and a low-

level autophagy group, where the low-level group seemed to lack GATA4-positive

PrE cells. Although based on a few samples, this data suggests that an ICM in a

transcriptionally earlier state of pluripotency, i.e. GATA4-negative and prior to the

second lineage segregation event, might be less sensitive to the consequences of

aneuploidy, i.e. show less autophagy. We thus propose that aneuploid human

embryonic cells in the ICM become more sensitive from the second lineage

segregation onwards. We hypothesize that this effect may even become stronger

and more specific to the embryonic lineage after onset of gastrulation, as elegantly

shown in human 2D gastruloids (Yang *et al.*, 2021).

Finally, we found that in those embryos that had already undergone the second

lineage segregation event, the aneuploid embryos had a significantly lower ratio of

PrE to EPI cells than euploid embryos. Further, this effect was dosage-dependent,

with embryos with larger gene-dosage imbalances showing significantly lower cell

numbers in the TE and PrE than euploid embryos or embryos with lower gene-

dosage imbalances. This suggests that complex aneuploidy both impairs TE/ICM

proliferation and PrE formation or that the PrE is more sensitive to aneuploidy-

mediated apoptosis than the less differentiated epiblast. Whether this effect is also

driven by p53 activation remains to be confirmed.

Taken together, our results suggest that while the same pathways appear to be

activated in human and mouse embryos with complex aneuploidy, human embryos

have a different response when it comes to the downstream effect of these pathways

in the different lineages. While the mouse readily depletes aneuploid cells by

apoptosis specifically from the embryonic lineage before implantation (Bolton *et al.*,

2016; Singla *et al.*, 2020), in the preimplantation human embryo, apoptosis appears

to be more active in the TE. The results suggest that in the human embryo,

undifferentiated pluripotent cells have a different response to aneuploidy-induced

cellular stress than the neighbouring differentiated cells.

To conclude, our results on human embryos with complex meiotic aneuploidy provide

further insight into the mechanisms by which human embryos respond to gene-

dosage imbalances, and may contribute to understanding how aneuploid cells are

selectively eliminated in mosaic embryos. Further research, especially by using

human embryos with endogenous chromosomal mosaicism, will be needed to shed

light on the interactions between proteotoxic stress, the p53 pathway and

differentiation during early development, for which the most novel models of human

embryo implantation (Deglincerti *et al.*, 2016; Shahbazi *et al.*, 2016, 2020; Kagawa *et*

al., 2021) will prove to be of incalculable value.

References

Banin S, Moyal L, Shieh SY, Taya Y, Anderson CW, Chessa L, Smorodinsky NI, Prives C, Reiss Y, Shiloh Y, *et al.* Enhanced phosphorylation of p53 by ATM in response to DNA damage. *Science* [Internet] 1998;**281**:1674–1677. Science.

Bolton H, Graham SJL, Aa N van der, Kumar P, Theunis K, Fernandez Gallardo E, Voet T, Zernicka-Goetz M. Mouse model of chromosome mosaicism reveals lineage-specific depletion of aneuploid cells and normal developmental potential. *Nat Commun* [Internet] 2016;**7**:1–12. Nature Publishing Group.

Capalbo A, Poli M, Rienzi L, Girardi L, Patassini C, Fabiani M, Cimadomo D, Benini F, Farcomeni A, Cuzzi J, *et al.* Mosaic human preimplantation embryos and their developmental potential in a prospective , non-selection clinical trial. Am. J. Hum. Genet., 108 (2021), pp. 2238-2247.

Capalbo A, Wright G, Elliott T, Ubaldi FM, Rienzi L, Nagy ZP. FISH reanalysis of inner cell mass and trophectoderm samples of previously array-CGH screened blastocysts shows high accuracy of diagnosis and no major diagnostic impact of mosaicism at the blastocyst stage. *Human Reproduction* 2013;**28**:2298–2307. Oxford University Press.

Chavez SL, Loewke KE, Han J, Moussavi F, Colls P, Munne S, Behr B, Reijo Pera RA. Dynamic blastomere behaviour reflects human embryo ploidy by the four-cell stage. *Nat Commun* [Internet] 2012;**3**:. Nat Commun.

Chow JFC, Yeung WSB, Lau EYL, Lee VCY, Ng EHY, Ho PC. Array comparative genomic hybridization analyses of all blastomeres of a cohort of embryos from young IVF patients revealed significant contribution of mitotic errors to embryo mosaicism at the cleavage stage. *Reprod Biol Endocrinol* [Internet] 2014;**12**:. Reprod Biol Endocrinol.

Coorens THH, Oliver TRW, Sanghvi R, Sovio U, Cook E, Vento-Tormo R, Haniffa M, Young MD, Rahbari R, Sebire N, *et al.* Inherent mosaicism and extensive mutation of human placentas. *Nature* 2021;**592**:80–85. Nature Research.

Crighton D, Wilkinson S, O'Prey J, Syed N, Smith P, Harrison PR, Gasco M, Garrone O, Crook T, Ryan KM. DRAM, a p53-Induced Modulator of Autophagy, Is Critical for Apoptosis. *Cell* 2006;**126**:121–134. Elsevier B.V.

Daughtry BL, Rosenkrantz JL, Lazar NH, Fei SS, Redmayne N, Torkenczy KA, Adey A, Yan M, Gao L, Park B, *et al.* Single-cell sequencing of primate preimplantation embryos reveals chromosome elimination via cellular fragmentation and blastomere exclusion. *Genome Res* [Internet] 2019;**29**:367–382. Cold Spring Harbor Laboratory Press.

David K Gardner, William B Schoolcraft. Culture and transfer of human blastocysts. *Curr Opin Obstet Gynecol* 1999;**11**:307–311.

661 Deglincerti A, Croft GF, Pietila LN, Zernicka-Goetz M, Siggia ED, Brivanlou AH. Self-
662   organization of the in vitro attached human embryo. *Nature* [Internet]
663   2016;**533**:251–254.
664
665 Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson
666   M, Gingeras TR. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
667   2013;**29**:15–21.
668
669 Dürrbaum M, Kuznetsova A, Passerini V, Stingele S, Stoehr G, Storchová Z. Unique
670   features of the transcriptional response to model aneuploidy in human cells.
671   *BMC Genomics* [Internet] 2014;**15**:139. BioMed Central.
672
673 Dürrbaum M, Storchová Z. Effects of aneuploidy on gene expression: implications for
674   cancer. *FEBS J* [Internet] 2016;**283**:791–802. John Wiley & Sons, Ltd.
675
676 Echten-Arends J van, Mastenbroek S, Sikkema-Raddatz B, Korevaar JC, Heineman
677   MJ, Veen F van der, Repping S. Chromosomal mosaicism in human
678   preimplantation embryos: a systematic review. *Hum Reprod Update* [Internet]
679   2011;**17**:620–627. Oxford University Press.
680
681 Fragouli E, Alfarawati S, Spath K, Babariya D, Tarozzi N, Borini A, Wells D. Analysis
682   of implantation and ongoing pregnancy rates following the transfer of mosaic
683   diploid–aneuploid blastocysts. *Hum Genet* [Internet] 2017;**136**:805–819.
684   Springer Verlag.
685
686 Fragouli E, Munne S, Wells D. The cytogenetic constitution of human blastocysts:
687   insights from comprehensive chromosome screening strategies. *Hum Reprod*
688   *Update* [Internet] 2019;**25**:15–33. Oxford University Press.
689
690 Fuchs Weizman N, Wyse BA, Antes R, Ibarrientos Z, Sangaralingam M, Motamedi
691   G, Kuznyetsov V, Madjunkova S, Librach CL. Towards Improving Embryo
692   Prioritization: Parallel Next Generation Sequencing of DNA and RNA from a
693   Single Trophectoderm Biopsy. *Sci Rep* [Internet] 2019;**9**:1–11. Nature Publishing
694   Group.
695
696 Greco E, Minasi MG, Fiorentino F. Healthy Babies after Intrauterine Transfer of
697   Mosaic Aneuploid Blastocysts. *New England Journal of Medicine* [Internet]
698   2015;**373**:2089–2090. Massachusetts Medical Society.
699
700 Griffin DK, Brezina PR, Tobler K, Zhao Y, Silvestri G, Mccoy RC, Anchan R, Benner
701   A, Cutting GR, Kearns WG. The human embryonic genome is karyotypically
702   complex, with chromosomally abnormal cells preferentially located away from
703   the developing fetus. *Human Reproduction* [Internet] 2022; Oxford University
704   Press (OUP).
705
706 Groff AF, Resetkova N, DiDomenico F, Sakkas D, Penzias A, Rinn JL, Eggan K.
707   RNA-seq as a tool for evaluating human embryo competence. *Genome Res*
708   [Internet] 2019;**29**:1705–1718. Cold Spring Harbor Laboratory Press.
709

Guan JJ, Zhang XD, Sun W, Qi L, Wu JC, Qin ZH. DRAM1 regulates apoptosis through increasing protein levels and lysosomal localization of BAX. *Cell Death Dis* [Internet] 2015;**6**:e1624–e1624. Nature Publishing Group.

Gump JM, Thorburn A. Autophagy and apoptosis: what is the connection? *Trends Cell Biol* [Internet] 2011;**21**:387–392. Trends Cell Biol.

Jacobs K, Velde H van de, Paepe C de, Sermon K, Spits C. Mitotic spindle disruption in human preimplantation embryos activates the spindle assembly checkpoint but not apoptosis until Day 5 of development. *MHR: Basic science of reproductive medicine* [Internet] 2017;**23**:321–329. Oxford University Press.

Jaiswal SK, Raj S, DePamphilis ML. Developmental Acquisition of p53 Functions. *Genes (Basel)* [Internet] 2021;**12**:1675. MDPI.

Johnson DS, Cinnioglu C, Ross R, Filby A, Gemelos G, Hill M, Ryan A, Smotrich D, Rabinowitz M, Murray MJ. Comprehensive analysis of karyotypic mosaicism between trophectoderm and inner cell mass. *MHR: Basic science of reproductive medicine* [Internet] 2010;**16**:944–949. Oxford University Press.

Kagawa H, Javali A, Khoei HH, Sommer TM, Sestini G, Novatchkova M, Scholte op Reimer Y, Castel G, Bruneau A, Maenhoudt N, *et al.* Human blastoids model blastocyst development and implantation. *Nature* [Internet] 2021;1–9. Nature Publishing Group

Kalousek DK, Vekemans M. Confined placental mosaicism. *J Med Genet* [Internet] 1996;**33**:529–533. BMJ Publishing Group.

Krivega M, Stiefel CM, Karbassi S, Andersen LL, Chunduri NK, Donnelly N, Pichlmair A, Storchová Z. Genotoxic stress in constitutive trisomies induces autophagy and the innate immune response via the cGAS-STING pathway. *Communications Biology 2021 4:1* [Internet] 2021;**4**:1–16. Nature Publishing Group.

Landuyt L Van, Stoop D, Verheyen G, Verpoest W, Camus M, Velde H Van De, Devroey P, Abbeel E Van Den. Outcome of closed blastocyst vitrification in relation to blastocyst quality: Evaluation of 759 warming cycles in a single-embryo transfer policy. *Human Reproduction* 2011;**26**:527–534. Oxford University Press.

Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 2011;**12**:323.

Licciardi F, Lhakhang T, Kramer YG, Zhang Y, Heguy A, Tsirigos A. Human blastocysts of normal and abnormal karyotypes display distinct transcriptome profiles. *Sci Rep* [Internet] 2018;**8**:1–9. Nature Publishing Group.

Lin T, Chao C, Saito S, Mazur SJ, Murphy ME, Appella E, Xu Y. p53 induces differentiation of mouse embryonic stem cells by suppressing Nanog expression. *Nat Cell Biol* [Internet] 2005;**7**:165–171. Nat Cell Biol.

Loughery J, Cox M, Smith LM, Meek DW. Critical role for p53-serine 15 phosphorylation in stimulating transactivation at p53-responsive promoters. *Nucleic Acids Res* [Internet] 2014;**42**:7666–7680. Oxford University Press.

Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014;**15**:. BioMed Central Ltd.

Mantel C, Guo Y, Lee MR, Kim M-K, Han M-K, Shibayama H, Fukuda S, Yoder MC, Pelus LM, Kim K-S, *et al.* Checkpoint-apoptosis uncoupling in human and mouse embryonic stem cells: a source of karyotpic instability. *Blood* [Internet] 2007;**109**:4518–4527.

Maxwell SM, Lhakhang TC, Lin Z, Kramer YG, Zhang Y, Wang F, Heguy A, Tsirigos A, Grifo JA, Licciardi F. Investigation of Global Gene Expression of Human Blastocysts Diagnosed as Mosaic using Next-generation Sequencing. *Reproductive Sciences* [Internet] 2022;1–11.

McCoy RC. Mosaicism in Preimplantation Human Embryos: When Chromosomal Abnormalities Are the Norm. *Trends in Genetics* [Internet] 2017;**33**:448–463. Elsevier.

Meistermann D, Bruneau A, Loubersac S, Reignier A, Firmin J, François-Campion V, Kilens S, Lelièvre Y, Lammers J, Feyeux M, *et al.* Integrated pseudotime analysis of human pre-implantation embryo single-cell transcriptomes reveals the dynamics of lineage specification. *Cell Stem Cell* 2021;**28**:1625-1640.e6.

Mertzanidou A, Spits C, Nguyen HT, Velde H van de, Sermon K. Evolution of aneuploidy up to Day 4 of human preimplantation development. *Human Reproduction* [Internet] 2013a;**28**:1716–1724. Oxford University Press.

Mertzanidou A, Wilton L, Cheng J, Spits C, Vanneste E, Moreau Y, Vermeesch JR, Sermon K. Microarray analysis reveals abnormal chromosomal complements in over 70% of 14 normally developing human embryos. *Human Reproduction* [Internet] 2013b;**28**:256–264. Oxford University Press.

Munné S, Blazek J, Large M, Martinez-Ortiz PA, Nisson H, Liu E, Tarozzi N, Borini A, Becker A, Zhang J, *et al.* Detailed investigation into the cytogenetic constitution and pregnancy outcome of replacing mosaic blastocysts detected with the use of high-resolution next-generation sequencing. *Fertil Steril* [Internet] 2017;**108**:62-71.e8. Elsevier Inc.

Munné S, Nakajima ST, Najmabadi S, Sauer M v., Angle MJ, Rivas JL, Mendieta L v., MacAso TM, Sawarkar S, Nadal A, *et al.* First PGT-A using human in vivo blastocysts recovered by uterine lavage: Comparison with matched IVF embryo controls. *Human Reproduction* [Internet] 2020a;**35**:70–80. Oxford University Press.

Munné S, Spinella F, Grifo J, Zhang J, Beltran MP, Fragouli E, Fiorentino F. Clinical outcomes after the transfer of blastocysts characterized as mosaic by high resolution Next Generation Sequencing- further insights. *Eur J Med Genet* 2020b;**63**:103741. Elsevier Masson SAS.

810 Niakan KK, Eggan K. Analysis of human embryos from zygote to blastocyst reveals
811 distinct gene expression patterns relative to the mouse. *Dev Biol* [Internet]
812 2013;**375**:54–64. Dev Biol.
813
814
815 Ohashi A, Ohori M, Iwai K, Nakayama Y, Nambu T, Morishita D, Kawamoto T,
816 Miyamoto M, Hirayama T, Okaniwa M, *et al.* Aneuploidy generates proteotoxic
817 stress and DNA damage concurrently with p53-mediated post-mitotic apoptosis
818 in SAC-impaired cells. *Nat Commun* [Internet] 2015;**6**:1–16. Nature Publishing
819 Group.
820
821 Oromendia AB, Dodgson SE, Amon A. Aneuploidy causes proteotoxic stress in
822 yeast. *Genes Dev* 2012;**26**:2696–2708.
823
824 Orvieto R, Shimon C, Rienstein S, Jonish-Grossman A, Shani H, Aizer A. Do human
825 embryos have the ability of self-correction? *Reprod Biol Endocrinol* [Internet]
826 2020;**18**:. Reprod Biol Endocrinol.
827
828 Palmerola KL, Amrane S, Los Angeles A de, Xu S, Wang N, Pinho J de, Zuccaro M
829 v, Taglialatela A, Massey DJ, Turocy J, *et al.* Replication stress impairs
830 chromosome segregation and preimplantation development in human embryos.
831 *Cell* [Internet] 2022.
832
833 Pan J-A, Ullman E, Dou Z, Zong W-X. Inhibition of protein degradation induces
834 apoptosis through a microtubule-associated protein 1 light chain 3-mediated
835 activation of caspase-8 at intracellular membranes. *Mol Cell Biol* [Internet]
836 2011;**31**:3158–3170. American Society for Microbiology.
837
838 Popovic M, Dhaenens L, Boel A, Menten B, Heindryckx B. Chromosomal mosaicism
839 in human blastocysts: The ultimate diagnostic dilemma. *Hum Reprod Update*
840 [Internet] 2020;**26**:313–334. Oxford University Press.
841
842 Popovic M, Dheedene A, Christodoulou C, Taelman J, Dhaenens L, Nieuwerburgh F
843 Van, Deforce D, Abbeel E Van Den, Sutter P De, Menten B, *et al.* Chromosomal
844 mosaicism in human blastocysts: The ultimate challenge of preimplantation
845 genetic testing? *Human Reproduction* [Internet] 2018;**33**:1342–1354. Oxford
846 University Press.
847
848 Regin M, Spits C, Sermon K. On the origins and fate of chromosomal abnormalities
849 in human preimplantation embryos: an unsolved riddle. *Mol Hum Reprod*
850 [Internet] 2022; Mol Hum ReprodAvailable from:
851 https://academic.oup.com/molehr/advance-
852 article/doi/10.1093/molehr/gaac011/6566308.
853
854 Ren Y, Yan Z, Yang M, Keller L, Zhu X, Lian Y, Liu Q, Li R, Zhai F, Nie Y, *et al.*
855 Regional and developmental characteristics of human embryo mosaicism
856 revealed by single cell sequencing. *PLoS Genet* 2022;**18**:. Public Library of
857 Science.
858

859 Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. Limma powers
860   differential expression analyses for RNA-sequencing and microarray studies.
861   *Nucleic Acids Res* [Internet] 2015;**43**:e47. Oxford University Press.
862
863 Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for
864   differential expression analysis of digital gene expression data. *Bioinformatics*
865   2009;**26**:139–140. Oxford University Press.
866
867 Robinson MD, Oshlack A. *A scaling normalization method for differential expression*
868   *analysis of RNA-seq data* [Internet]. 2010.
869
870 Sanchez-Ribas I, Diaz-Gimeno P, Sebastián-León P, Mercader A, Quiñonero A,
871   Ballesteros A, Pellicer A, Domínguez F. Transcriptomic behavior of genes
872   associated with chromosome 21 aneuploidies in early embryo development.
873   *Fertil Steril* 2019;**111**:991-1001.e2. Elsevier Inc.
874
875 Santaguida S, Amon A. Short- and long-term effects of chromosome mis-segregation
876   and aneuploidy. *Nat Rev Mol Cell Biol* 2015;**16**:473–485.
877
878 Shahbazi MN, Jedrusik A, Vuoristo S, Recher G, Hupalowska A, Bolton V, Fogarty
879   NME, Campbell A, Devito LG, Ilic D, *et al.* Self-organization of the human
880   embryo in the absence of maternal tissues. *Nat Cell Biol* [Internet] 2016;**18**:700–
881   708. Nature Publishing Group.
882
883 Shahbazi MN, Wang T, Tao X, Weatherbee BAT, Sun L, Zhan Y, Keller L, Smith GD,
884   Pellicer A, Scott RT, *et al.* Developmental potential of aneuploid human embryos
885   cultured beyond implantation. *Nat Commun* [Internet] 2020;**11**:1–15. Nature
886   Research.
887
888 Singla S, Iwamoto-Stohl LK, Zhu M, Zernicka-Goetz M. Autophagy-mediated
889   apoptosis eliminates aneuploid cells in a mouse model of chromosome
890   mosaicism. *Nat Commun* [Internet] 2020;**11**:1–15. Nature Research.
891
892 Starostik MR, Sosin OA, McCoy RC. Single-cell analysis of human embryos reveals
893   diverse patterns of aneuploidy and mosaicism. *Genome Res* 2020;**30**:814–826.
894
895 Stingele S, Stoehr G, Peplowska K, Cox J, Mann M, Storchova Z. Global analysis of
896   genome, transcriptome and proteome reveals the response to aneuploidy in
897   human cells. *Mol Syst Biol* [Internet] 2012;**8**:608. EMBO Press.
898
899 Vanneste E, Voet T, Caignec C le, Ampe M, Konings P, Melotte C, Debrock S,
900   Amyere M, Vikkula M, Schuit F, *et al.* Chromosome instability is common in
901   human cleavage-stage embryos. *Nat Med* [Internet] 2009;**15**:577–583. Nature
902   Publishing Group.
903
904 Vázquez-Diez C, Paim LMG, FitzHarris G. Cell-Size-Independent Spindle Checkpoint
905   Failure Underlies Chromosome Segregation Error in Mouse Embryos. *Current*
906   *Biology* [Internet] 2019;**29**:865-873.e3. Cell Press.
907

908 Vera-Rodriguez M, Chavez SL, Rubio C, Pera RAR, Simon C. Prediction model for
909   aneuploidy in early human embryo development revealed by single-cell analysis.
910   *Nat Commun* [Internet] 2015;**6**:7601. Nature Publishing Group.
911
912
913
Victor AR, Tyndall JC, Brake AJ, Lepkowsky LT, Murphy AE, Griffin DK, McCoy RC,
Barnes FL, Zouves CG, Viotti M. One hundred mosaic embryos transferred
prospectively in a single clinic: exploring when and why they result in healthy
pregnancies. *Fertil Steril* [Internet] 2019;**111**:280–293. Elsevier Inc.
Viotti M, Victor AR, Barnes FL, Zouves CG, Besser AG, Grifo JA, Cheng EH, Lee
MS, Horcajadas JA, Corti L, *et al.* Using outcome data from one thousand
mosaic embryo transfers to formulate an embryo ranking system for clinical use.
Fertil Steril 2021;**115**:1212–1224. Elsevier Inc.
922
923 Voullaire L, Slater H, Williamson R, Wilton L. Chromosome analysis of blastomeres
924   from human embryos by using comparative genomic hybridization. *Hum Genet*
925   [Internet] 2000;**106**:210–217.
926
927 Wells D, Delhanty JD. Comprehensive chromosomal analysis of human
928   preimplantation embryos using whole genome amplification and single cell
929   comparative genomic hybridization. *Mol Hum Reprod* [Internet] 2000;**6**:1055–
930   1062.
931
932 Yan L, Yang M, Guo H, Yang L, Wu J, Li R, Liu P, Lian Y, Zheng X, Yan J, *et al.*
933   Single-cell RNA-Seq profiling of human preimplantation embryos and embryonic
934   stem cells. *Nat Struct Mol Biol* [Internet] 2013;**20**:1131–1139. Nature Publishing
935   Group.
936
937 Yang M, Rito T, Metzger J, Naftaly J, Soman R, Hu J, Albertini DF, Barad DH,
938   Brivanlou AH, Gleicher N. Depletion of aneuploid cells in human embryos and
939   gastruloids. *Nat Cell Biol* [Internet] 2021;**23**:314–321. Nature Research.
940
941 Zamani Esteki M, Viltrop T, Tšuiko O, Tiirats A, Koel M, Nõukas M, Žilina O, Teearu
942   K, Marjonen H, Kahila H, *et al.* In vitro fertilization does not increase the
943   incidence of de novo copy number alterations in fetal and placental lineages. *Nat*
944   *Med* 2019;
945
946 Zhu J, Tsai HJ, Gordon MR, Li R. Cellular Stress Associated with Aneuploidy. *Dev*
947   *Cell* [Internet] 2018;**44**:420–431. Cell Press.
948
949

Data availability

Data are available from the corresponding author upon reasonable request.

Acknowledgements

The authors would like to thank Marleen Carlé from the Center for Medical Genetics (UZ Brussel) for assisting during the tubing of trophectoderm samples. We thank Wilfried Cools from the Biostatistics and Medical Informatics Group (VUB) for the statistical advice and the members of the BRIGHTcore facility (UZ Brussel) for performing the RNA sequencing. We thank Prof. Rajiv McCoy for the scientific input and data exchange. This study was funded by the Fonds for Scientific Research in Flanders -G017218N- (Fonds Wetenschappelijk Onderzoek – Vlaanderen [FWO]). M.R. and E.C.D.D. are doctoral fellows at the FWO.

Author contributions

M.R. carried out all the experiments unless stated otherwise, interpreted and analyzed the results and co-wrote the manuscript. Y.L. and E.C.D.D. performed the bioinformatics analysis. Y.G. performed the trophectoderm biopsies. P.V. provided and supervised the diagnoses of the PGT embryos. G.V. and H.V.D.V. provided the embryos for this study. K.S. interpreted and supervised the experimental work and co-wrote the manuscript. C.S. designed, interpreted and supervised the experimental work and co-wrote the manuscript. All co-authors proofread the manuscript.

Competing interests

The authors declare no competing interests.

Fig. 1: RNA-sequencing of trophectoderm cells reveals deregulation of DNA damage, p53, cell-cycle, autophagy and apoptosis pathways in chromosomally abnormal human embryos.

a. Human blastocysts 5 or 6 days post-fertilization (dpf) underwent Preimplantation Genetic Testing (PGT) for monogenic diseases and/or aneuploidy screening, followed by cryopreservation. Embryos that were not transferred were donated for research after written informed content. After warming and 2hrs recuperation, euploid and fully aneuploid research embryos underwent a second TE biopsy for RNA-sequencing. The remainder of the embryos was live-stained with either CASP3/7 or CASP8 for 30min and fixed for immunostaining for proteins of interest (see figure 3). **b**. 3dpf human embryos were warmed and treated with 0.5μM reversine for 24hrs then transferred to normal culture medium until 5dpf to undergo TE biopsy for RNA-sequencing. After 2hrs recuperation the embryos were live-stained with CASP8 for 30min and fixed for immunostaining for proteins of interest (see figure 3). **c**. Unsupervised hierarchical clustering of all coding genes. Color coding: beige = euploid, red = aneuploid and blue = reversine-treated. **d**. Comparison between the diagnosis of aneuploid embryos obtained after PGT (list on the left) and the results obtained after using InferCNV after RNA-sequencing (plot on the right). **e**,**f**. Volcano plots after differential gene expression analysis with a cutoff value of |$\log_2$ fold change| > 1 and -$\log_{10}$(p-value) < 0.05 for aneuploid versus euploid (**e**) and reversine versus euploid (**f**). **g**,**h**. Venn diagrams comparing the upregulated (**g**) and downregulated (**h**) genes in aneuploid versus reversine treated embryos. **i**,**j**. Supervised plot of pathway analysis using the Molecular Signatures Database (FDR < 0.05) based on the list of all differentially expressed genes between aneuploid or reversine versus euploid embryos. Pathways are ranked based on the total number

of genes in overlap of both test sets, ranked from sets with the highest number of

genes in overlap to sets with the lowest. Outputs of the Hallmark library and C5

ontology gene sets are shown in (**i**) and (**j**), respectively. **k,l**. Supervised plots after

Ingenuity Pathway Analysis of upstream regulators with an activation z-score ≤ -2

and ≥ 2 and a p-value < 0.05 in the aneuploid (**k**) or reversine (**l**) versus euploid

embryos. **m,n**. Supervised plots after Ingenuity Pathway Analysis of canonical

pathways in aneuploid (**m**) or reversine (**n**) versus euploid embryos. Pathways are

ranked by p-value from the most to the least significant, those with a p-value < 0.05

and with an activation z-score ≤ -1.5 and ≥ 1.5 were considered as inhibited or

activated, respectively. Significant pathways with an activation z-score between -1.5

and 1.5 were considered as deregulated.

1008

Fig. 2: Human embryos with the highest number of genes with abnormal copy

number show stronger p53 pathway and apoptosis response.

a. Supervised hierarchical clustering of the TOP 500 deregulated genes sorted by p-

value. Color coding: blue = euploid , green = Low dosage and red = High dosage.

b,c. Volcano plots after differential gene expression analysis with a cutoff value of

|$\log_2$ fold change| > 1 and -$\log_{10}$(p-value) < 0.05 for low dosage versus euploid

embryos (**b**) and high dosage versus euploid embryos (**c**). **d,e**, Venn diagrams

comparing the upregulated (**d**) and downregulated (**e**) genes in low dosage versus

high dosage embryos. **f**. Supervised plot of pathway analysis using the Hallmark

library of the Molecular Signatures Database (FDR < 0.05) based on the list of the

TOP 500 expressed genes by p-value between low or high dosage versus euploid

embryos. Pathways are ranked based on FDR-q-values of both test sets, the most

significant deregulated pathway ranked from top to least significant pathway. **g**,**h**. Dot

plots of the genes in overlap with the Hallmark p53 pathway (**g**) and Hallmark

apoptosis pathway (**h**) of both low and high dosage embryos sorted by $\log_2$ Fold

Change from the highest to the lowest. Dots with an increased size and a darker blue

represent a higher value of $\log_2$ Fold Change and lower p-value, respectively. **i**, Violin

plots with box and whisker plots of the counts per million mapped reads of a

supervised set of 10 genes (from Supplementary Table 2) that are part of apoptosis

(*BAD, BCL2L1, TNFRSF10B*), p53 pathway (*DRAM1, MDM2, CDKN1A/B, PURPL*),

growth arrest (*GADD45A*), and DNA-damage (*DDB2*). *p = 0.027, **p = 0.010, ***p =

0.004, ****p < 0.001 using the Jonkheere-Terpstra test. Box and whisker plots show

median and minimum to maximum values.

Fig. 3: Immunostaining reveals increased apoptosis, autophagy, proteotoxic

stress and DNA-damage in chromosomally abnormal human embryos.

a,**b**. Same experimental set-up as described in **Fig. 1 a**,**b**, but focused on

immunostaining analysis. **c**, Orthogonal projections after immunostaining of euploid

and aneuploid embryos for DNA (white) and CASP3/7 (green). **d**, Orthogonal

projections after immunostaining of euploid, aneuploid and reversine-treated

embryos for DNA (white) and CASP8 (green). **e**, CASP3/7 mean intensity per cell.

Euploid n=13, Aneuploid n=8 embryos. Mann-Whitney test, *p = 0.0199. **f**, CASP8

mean intensity per cell. Euploid n=6, Aneuploid n=9, Reversine-treated n=8 embryos.

Kruskal-Wallis test, *p = 0.0394, **p = 0.0085. **g**, Number of nuclei per embryo.

Euploid n=19, Aneuploid n=18, Reversine-treated n=8 embryos. Kruskal-Wallis test,

****p < 0.0001, *p = 0.0305. **h**, Orthogonal projections after immunostaining of

euploid, aneuploid and reversine-treated embryos for DNA (white), LC3B (turquoise)

and HSP70 (magenta). **i**, LC3B puncta per cell. Euploid n=19, Aneuploid n=22

embryos, Reversine-treated n=8 embryos. Kruskal-Wallis test, *p = 0.0189, **p = 0.0074. **j**, HSP70 mean intensity per cell. Euploid n=13, Aneuploid n=8, Reversine-treated n=8 embryos. Kruskal-Wallis test, **p = 0.0077, ***p = 0.0008. **k**, Orthogonal projections after immunostaining of euploid, aneuploid for DNA (white) and p62 (turquoise). **l**, p62 puncta per cell. Euploid n=6, Aneuploid n=9 embryos. Mann-Whitney test, **p = 0.0076.

Brightfield pictures were obtained during confocal imaging. All scale bars are 20 μm. Box and whisker plots show median and whiskers show minimum to maximum values. Bar plots show mean ± s.d. For all plots each dot represents a single embryo and ns = not significant.

Fig. 4: Aneuploid human embryos show less cells in trophectoderm and OCT4-positive cells and impaired primarily primitive endoderm differentiation

a, 5 or 6dpf post-PGT-A euploid and fully aneuploid human blastocysts were warmed and kept for 16hrs in culture to allow for the second lineage segregation. The blastocysts were live-stained with CASP3/7 for 30min and fixed for immunostaining proteins of interest. In a first scan we analyzed OCT4 (ICM), CASP3/7 (apoptosis), LC3B (autophagy) and S15p53 (DNA damage) in both lineages. Embryos were then re-stained with GATA4 as a marker for PrE. **b,** Orthogonal projections after immunostaining of euploid, aneuploid for DNA (white) and Serine (S) 15 p53 (turquoise). **c**, Percentage (%) of Serine 15 p53 positive cells per embryo. Euploid n = 3, Aneuploid n = 5. Mann-Whitney test, *p = 0.0357. **d**, Orthogonal projections after immunostaining of euploid and aneuploid embryos for DNA (white) and OCT4 (magenta). The first aneuploid panel (Trisomy 14 and 16) shows a similar number of

ICM cells compared to the euploid embryo. The second aneuploid panel (Trisomy 14

and 22) shows an embryo without an ICM. **e,f**,**g**, Differences in number of nuclei per

embryo between euploid and aneuploid embryos in the OCT4-negative cells

(trophectoderm) (**e**) **p = 0.0092 , OCT4-positive cells (ICM/EPI), *p = 0.0412 (**f**) and

whole embryo **p = 0.0067 (**g**). Euploid n = 10, Aneuploid n = 14. Student's t-test

with Welch's correction**. h**, Orthogonal projections after immunostaining of euploid

and aneuploid embryos for DNA (white), CASP3/7 (green) and OCT4 (magenta). **i**,

Percentage (%) of CASP3/7 positive (+) cells in the trophectoderm lineage (OCT4-

negative cells). Euploid n = 9, Aneuploid n = 14. Student's t-test with Welch's

correction, *p = 0.0453. **j**, CASP3/7 mean intensity per cell of whole embryos.

Euploid n = 9, Aneuploid n = 14. Student's t-test with Welch's correction, *p = 0.0332.

k, Orthogonal projections after immunostaining of euploid and aneuploid embryos for

DNA (white), OCT4 (magenta) and GATA4 (green). The first aneuploid panel

(trisomy 13 and 15) shows an embryo with presence of a GATA4 positive cell (PrE).

The second aneuploid panel (Trisomy 7 and monosomies 5, 12 and 21) shows an

embryo that did not contain GATA4 positive cells. **l**, Differences in the number of

cells per embryo that were OCT4 positive and either GATA4 negative (red) or

positive (green) between euploid and aneuploid embryos. In case GATA4-positive

cells (PrE) were present we considered the GATA4-negative cells to be epiblast

(EPI). All euploid embryos contained GATA4-positive cells, 7/14 aneuploid embryos

had an ICM completely lacking GATA4-positive cells (Fisher-exact test, p = 0.0188).

m, Differences in the PrE/EPI ratio between euploid and aneuploid embryos that had

an ICM containing GATA4-positive cells. Euploid n = 9, Aneuploid n = 7. Note here

that one euploid embryo (#6, **l**) was lost during re-staining for GATA4. Mann-Whitney

test, **p = 0.0089. **n**, Orthogonal projections after immunostaining of euploid and

aneuploid embryos for DNA (white), LC3B (turquoise) and OCT4 (magenta). Yellow

circle indicates the ICM. The first aneuploid panel (Trisomy 1 and 20 and monosomy

18) shows OCT4-positive cells (ICM/EPI) with low levels of autophagy and, after re-

staining for GATA4, without PrE (right panel, ICM zoom). The second aneuploid

panel (Monosomy 14) shows an embryo with high levels of autophagy in the ICM that

contains PrE cells (right panel, ICM zoom). **o,p,q.** Differences in LC3B puncta per

cell between euploid and aneuploid embryos in the whole embryo,*p = 0.0317

(**o**),(OCT4-negative cells (trophectoderm),*p = 0.0317 (**p**) and OCT4-positive cells

(ICM/EPI) (**q**). Euploid n = 4, Aneuploid n = 5. Mann-Whitney test. **r**, Violin plots with

box plots of OCT4-positive-cells (ICM/EPI) per embryo (left, ns), OCT4-negative cells

(trophectoderm) per embryo (center, *p = 0.043) and GATA4-positive-cells (PrE) per

embryo (right, ****p < 0.001, Jonkheere-Terpstra test) depending on gene dosage.

Euploid (light blue), Low dosage (middle blue) and High dosage (darkest blue).

Yellow arrows indicate presence of the signal and yellow arrow heads indicate

absence of signal. Brightfield pictures were obtained during confocal imaging. All

scale bars are 20 μm. Box and whisker plots show median and minimum to

maximum values. Bar plots show mean ± s.d. For all plots each dot represents a

single embryo; ns = not significant. T = Trisomy, M = Monosomy

1113

Supplementary Fig. 1: Expanded RNA-sequencing analysis

a, Principal component analysis after RNA-sequencing of euploid (green) versus

aneuploid (red) and reversine (blue) embryos. **b**, InferCNV analysis of individual

euploid samples compared to the reference set. **c**, InferCNV analysis of individual

reversine samples compared to the reference set. **d**, Venn Diagram of overlap

between differentially expressed genes of this study with two studies that analyzed

cleavage stage embryos that contain euploid and diverse types of aneuploidy from cleavage stage embryos (Vera-Rodriguez et al. 2015) and single cells from 4 and 7 dpf human embryos from TE, ICM, EPI and PrE lineages (Starostik et al. 2020). **e**, Venn Diagram of overlap between differentially expressed genes of this study with four studies that analyzed specific aneuploidies (Licciardi et al. 2018, Fuchs-Weizman et al. 2019, Sánchez-Ribas et al. 2019 and Maxwell et al. 2022).

Supplementary Fig. 2: S15-p53 expression in TE vs OCT4-positive cells (ICM/EPI) and increased micronuclei in aneuploid embryos

a, Orthogonal projections after immunostaining of euploid and aneuploid embryos for DNA (white), S15p53 (turquoise) and OCT4 (magenta). **b**, Percentages of nuclei positive for S15p53 per embryo in euploid (n = 3) and aneuploid (n = 4) embryos in the ICM. **c**, Percentages of nuclei positive for S15p53 per embryo in euploid (n = 3) and aneuploid (n = 5) embryos in the TE. **d**, Left panel: Orthogonal projections after immunostaining of euploid and aneuploid embryos for DNA (white) and CASP3/7 (green) showing micronuclei. Yellow box indicates zoomed-in region in the right panel. Yellow arrows indicate presence of micronuclei and overlap between DNA and CASP3/7. **e**, Number of micronuclei per cell in euploid and aneuploid whole embryos. Euploid n = 9, Aneuploid n = 14. Student's t-test with Welch's correction, *p = 0.0115.

Brightfield pictures were obtained during confocal imaging. All scale bars are 20 μm. Box and plots show median and minimum to maximum values. Bar plots show mean ± s.d. For all plots each dot represents a single embryo; ns = not significant. ROI = Region of interest.

1143

1144

a

dpf5/6

PGT   RNA-sequencing

Euploid   Aneuploid

Immunostaining
CASP3/7, CASP8, LC3B
p62, HSP70

b

dpf3 → dpf4 → dpf5

0.5µM reversine   washout

RNA-sequencing   Immunostaining
CASP8, LC3B, HSP70

c

euploid   aneuploid   reversine

d

PGT   InferCNV   gain   loss

M14/21
M17
T5;M14/19
M14/19;T16
T9;M22
M14
T2/22;M13
T2/10
T18;M22
T7;M11
M14/21/22;T18
M13;T22
T19;M22
T17;M18/19/21
T10/22
T15/22;M18
T1/15/18/22
T2/22;M13
M5/21;T9
M2/19

Chromosomes
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

e   Aneuploid vs Euploid

-log₁₀ p-value
log₂ Fold change

f   Reversine vs Euploid

-log₁₀ p-value
log₂ Fold change

g   upregulated genes

547   118   539
Aneuploid   Reversine

h   downregulated genes

463   84   355
Aneuploid   Reversine

i   Hallmark library   aneuploid   reversine

p53 pathway
Hypoxia
Glycolysis
Estrogen response early
Apoptosis
Estrogen response late
G2M checkpoint
Mitotic spindle

Genes in overlap

j   C5 library   aneuploid   reversine

Cell population proliferation
Identical protein binding
Apoptotic process
Regulation of cell death
Cell cycle
Negative regulation of cell death
Cellular response to DNA damage stimulus
Intrinsic apoptotic in response to DNA damage
DNA damage response by p53 class mediator
Mitotic G1/S transition checkpoint

Genes in overlap

k   Aneuploid vs Euploid   activated   inhibited

CDK19
AURK
GAS2L3
FGF2
ERK
ANLN
TP73
TP53
TOB1
RB1
RASSF1
TNF
IL1B
IL15
IFNG
RELA
MAP3K8

Activation Z-score

l   Reversine vs Euploid   activated

FOXM1
ATM
IKBKB
FGF2
EGFR
DUX4
AREG
TAZ
TNF
CX3CL1
AKT

Activation Z-score

m   activated   deregulated   inhibited   Aneuploid vs Euploid

IL-15 production
p53-signalling
Estrogen-mediated S-phase entry
G1/S Checkpoint regulation
Autophagy
TGF-beta-signalling
STAT3-pathway
Estrogen biosynthesis
Senescence pathway
BRCA1 in DNA damage response
JAK/STAT signalling
Cell cycle regulation by BTG proteins

Activation Z-score

n   activated   deregulated   Reversine vs Euploid

IL-15 production
JAK/STAT signalling
STAT3-pathway
Senescence pathway

Activation Z-score

a dpf5/6

PGT
Euploid Aneuploid

RNA-sequencing

Immunostaining
CASP3/7, CASP8, LC3B
p62, HSP70

b dpf3 → dpf4 washout dpf5

0.5μM reversine

RNA-sequencing

Immunostaining
CASP8, LC3B, HSP70

c Brightfield | DNA | CASP3/7 | MERGE

Euploid

Aneuploid

e Caspase-3/7 mean intensity per cell
Euploid   Aneuploid
*

f Caspase-8 mean intensity per cell
Euploid   Aneuploid   Reversine
**   ns   *

g Nuclei per embryo
Euploid   Aneuploid   Reversine
****   ns   *

d Brightfield | DNA | CASP8 | MERGE

Euploid

Aneuploid

Reversine

h Brightfield | DNA | LC3B | HSP70 | MERGE

Euploid

Aneuploid

Reversine

i LC3B puncta/cell
Euploid   Aneuploid   Reversine
**   ns   ***

j HSP70 mean intensity/cell
Euploid   Aneuploid   Reversine
**   ns   *

k Brightfield | DNA | p62 | MERGE

Euploid

Aneuploid

l p62 puncta/cell
Euploid   Aneuploid
**

a PCA plot

b

c

d Differentially expressed genes of diverse types of aneuploidy

e Differentially expressed genes of specific aneuploidies

a

	DNA	S15p53	OCT4	MERGE
Euploid				
Aneuploid				

b Inner cell mass

% of S15p53 positive cells

ns

Euploid   Aneuploid

c Trophectoderm

% of S15p53 positive cells

ns

Euploid   Aneuploid

d

	DNA	CASP3/7	MERGE		DNA	CASP3/7	MERGE
Euploid				ROI			
Aneuploid							

e

micronuclei per cell

*

Euploid   Aneuploid