Ralstonia solanacearum core effector RipE1 interacts and cleaves the Arabidopsis exocyst component Exo70B1

Dimitra Tsakiri[1§], Konstantinos Kotsaridis[1§], Vassiliki A. Michalopoulou[1], Michael Kokkinidis[1,2], Panagiotis F. Sarris[1,2,3]*

[1] Department of Biology, University of Crete, 714 09 Heraklion, Crete, Greece,
[2] Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology-Hellas, Heraklion, Crete, Greece,
[3] Biosciences, University of Exeter, Exeter, United Kingdom.

*Corresponding Author:
Professor Panagiotis F. Sarris: p.sarris@imbb.forth.gr; p.sarris@uoc.gr

[§] Equal contribution

Keywords

Pathogen effector, T3E, Plant defense, Cysteine protease, T3SS, exocyst complex

Abstract

Ralstonia solanacearum depends on numerous virulence factors, also known as effectors, to promote disease in a wide range of economically important host-plants. Although some of these effectors have been characterized, none has been shown to target the host's secretion machinery so far. Here, a screening was performed, using an extended library of NLR plant immune receptors' integrated domains (IDs), to identify new effector targets. The results uncovered that the core effector RipE1, of the *R. solanacearum* species complex, among other targets, associates with Arabidopsis exocyst component Exo70B1. RipE1, in accordance with its predicted cysteine protease activity, cleaves Exo70B1 *in vitro* between its sequence and is able to promote Exo70B1 degradation *in planta*. RipE1 enzymatic activity additionally results in the activation of TN2-dependent cell death. TN2 is an atypical NLR that has been proposed to guard Exo70B1. Despite the fact that RipE1 has been previously reported to activate defense responses in model plant species, we present here a *Nicotiana* species, in which RipE1 expression does not activate cell death. Overall, this study uncovers a new RipE1 host target, while providing evidence and novel tools to advance *in-depth* studies of RipE1 and homologues effectors.

Introduction

Plants, unlike animals, solely rely on their innate immune system to activate defense responses and fight off pathogens (1,2). To counterattack these responses, pathogens have evolved multiple virulence components, also known as effectors, that manipulate eukaryotic cell host metabolism, including pathways involved in innate immunity (3,4).

Plant innate immunity involves two intertwined cross-talking pathways that discern signs of invasion (5,6). Plant pattern-recognition receptors (PRRs) sense extracellular danger signals, mostly pathogen-associated molecular patterns (PAMPs), and activate a series of defense responses that constitute PAMP-triggered immunity (PTI) (7,8). Pathogen effectors translocated inside the host cell to block PTI, are recognized by plant intracellular nucleotide-binding leucine-rich repeat receptors (NLRs) that subsequently trigger the effector-triggered immunity (ETI) responses (7,9). Effectors' recognition often leads to induction of a focused rapid programmed cell death known as the hypersensitive response (HR), that stops pathogen spread (10,11). NLRs can recognize effectors either by directly binding or indirectly by perceiving modifications caused by effectors' activity on the operational target proteins (guardee model) or by a second protein that mimics their cellular target (decoy model) (12,13). A specialized version of direct recognition by NLRs is described via NLR integrated protein domains (NLR-IDs) (12,14). This model refers to NLRs carrying non-canonical domain fusions (integrated domains – IDs) that resemble pathogen targets to attract effectors and facilitate recognition (14,15). The evolutionary arms race between plants and microbial pathogens continues with either effector modification or the rise of new effectors that interfere with NLR activation or other downstream ETI signaling events, resulting in effectors' triggered susceptibility (ETS) (16).

The exocyst is an octameric protein complex that mediates tethering of secretory vesicles to the plasma membrane, before their SNARE (soluble N-ethylmaleimide- sensitive factor attachment protein receptors) - dependent fusion (17,18). Although it was first identified in yeast, the exocyst complex is highly conserved among eukaryotes and is also present in plants (18–20). Among exocysts from different organisms that have been studied to date, the complex seems to be composed of the same eight subunits -Sec3,

Sec5, Sec6, Sec8, Sec10, Sec15, Exo70, and Exo84- with the main difference in plants being the multiple paralogues encoding some of the subunits (20,21). In plants, the exocyst complex is involved in many physiological processes and there is strong evidence of its contribution to both PTI and ETI responses (22–29). The exocyst's role in plant immunity is further validated through the fact that different pathogens have acquired effectors that target exocyst components or their interactors and compromise defense-related exocytosis (30–35). In plants, Exo70 paralogues seem to be the most abundant, with Arabidopsis having 23 different Exo70 genes (24). Several of these Exo70 paralogues have been implemented in plant immune responses and are required for defense against distinct pathogens (25, 36, 37). Specifically, Arabidopsis Exo70B1 role in plant defense is highlighted by not only its interaction with key regulators and immune receptors, but also, by its involvement in plant autophagy, an emerging player in plant immunity responses (38–42). Furthermore, a number of different pathogen effectors target Arabidopsis Exo70B1, aiming to disrupt host secretion of defense-related molecules (30, 34, 35).

Ralstonia solanacearum is a destructive plant pathogen and the causative agent of bacterial wilt disease, affecting more than 250 plant species, including agronomically important crops (43–45). As in the case for many Gram-negative bacterial pathogens of both plants or animals, its Type III Secretion System (T3SS) is considered as a key virulence mechanism and is necessary for successful colonization of plant tissues by *R. solanacearum* (44, 46). T3SS is indispensable for the pathogen, since it serves for the injection of more than 70 Type III Effectors (T3Es) inside the host cell to promote colonization (44,47,48).

RipE1 is a T3E present in most of the so-far sequenced strains of the *R. solanacearum* species complex and seems to be highly conserved across different phylotypes (47, 48). Recent data suggest the effector's contribution to virulence and although it seems to trigger plant immunity in *Nicotiana benthamiana*, *Nicotiana tabacum* and *Arabidopsis thaliana*, other *R. solanacearum* T3Es are potentially able to suppress these responses (49–53). RipE1 has been shown to promote degradation of JAZ (jasmonate-ZIM-domain) repressors through its cysteine protease activity, inducing JA signaling and consequently

suppressing SA responses, in accordance with the earlier characterized RipE1 homologue

HopX1 (formerly AvrPphE) from *Pseudomonas syringae* (49, 54). In addition, cysteine

protease activity of RipE1 is known to be required for triggering immune responses in

both *N. benthamiana* and *A. thaliana* (51). Apart from their catalytic triad,

HopX/AvrPphE homologues additionally share a conserved N-terminal domain, the so-

called A-box, which is of unknown function but apparently necessary for effector

function *in planta* (55). For RipE1, the A-box domain is also required for triggering

immune responses in *N. benthamiana* (51).

In this work we aimed to uncover new targets of the conserved *R. solanacearum* T3E

RipE1 and further investigate its contribution to the pathogen's virulence. We provide

evidence for multiple potential RipE1 targets and propose evidence for the *Arabidopsis*

thaliana exocyst component Exo70B1 as a subcellular target of the particular effector *in*

planta. We use AI-based effector structure and function predictions and demonstrate *in*

vitro cleavage and *in planta* degradation of Exo70B1 by cysteine protease RipE1.

Overall, this study grants novel insight into discovering plant pathogen effectors'

distinctive targets and is the first report of a *R. solanacearum* effector targeting a

component of the host secretion machinery.

Results and Discussion

RipE1 intracellular function based on structure prediction

The AlphaFold, an artificial intelligence (AI), developed by DeepMind, protein structure prediction tool and it has been proposed as a highly accurate method to predict protein structures, compared to other well established methods (56). The 3D structure of RipE1 (47.31 kDa) was predicted using the AlphaFold Colab (**Fig. 1A**). The resulted structure confirmed that RipE1 has a cysteine protease folding. Cysteine proteases (thiol proteases) are enzymes with the ability of protein degradation by deprotonation of a thiol in the enzyme's active site (57). Therefore, all enzymes in this category have a common catalytic triad or dyad involving a nucleophilic cysteine residue (57). The predicted RipE1 structure was compared to Protein Data Bank structures via the DALI online server (58). The highest similarity was with the *Legionella pneumophila* periplasmic protease LapG (59) (**Fig. 1C**). According to this comparison, the extending hydrophobic surface patch at the active site of RipE1 (**Fig. 1B**) may suggest a role as an interaction interface for substrate binding. Subsequently, the superimposition (**Fig. 1D**) of the domain of the active sites between the two proteins had a root-mean-square deviation (RMSD) of 1.544. The catalytic triad cysteine-histidine-aspartate (C-H-D) of RipE1, specifically C172-H203-D222 (**Fig. 1A**), appeared to be identical with the LapG C137-H172-D189 catalytic triad (**Fig. 1D**). RipE1 belongs to the HopX1/AvrPphE effector family, the members of which share a conserved N-terminal domain and a catalytic triad with predicted cysteine protease activity (54, 55, 60). RipE1 produced by the *R. solanacearum* RS1000 strain has been shown to possess this enzymatic activity and it seemingly depends on it to cleave JAZ repressors (49). Using AI-based predictions (**Figure 1**), we show that RipE1 produced by the GMI1000 strain is a cysteine protease, as well. This further supports the findings of Sang et al (51), who reported that GMI1000-RipE1 loses its immunity-inducing attribute in *Nicotiana benthamiana* and *Arabidopsis thaliana* plants, when the cysteine residue belonging to the catalytic triad, is substituted.

RipE1 interacts with six ID protein domains and with Arabidopsis Exo70B1 in yeast

To identify new putative subcellular targets of RipE1, we performed a yeast two-hybrid (Y2H) screening between RipE1 and 22 eukaryotic protein domains (**Table S1**) that are fused to plant NLRs as Integrated Domains (NLR-IDs). We uncovered a total of 6 interactions (**Fig. 2A**) between RipE1 and NLR-IDs containing conserved characteristic domains of large plant protein families. RipE1 interacted with a TRX-like, a WRKY-like, the Exo70-like, and the CG-like NLR-IDs of our list, while weak interactions were also detected between RipE1 and the TFIIS-like, HSF-like and Ubox-like domains (**Fig. 2A, Table S1**). The expression of either RipE1 or any of the six NLR-IDs in yeast were not able to activate the reporter gene in the absence of an interactor, shortening the possibility of false-positive results (**Fig. S1A**). The conserved eukaryotic domains that interacted with RipE1 are commonly found on proteins that are tightly involved in regulation and facilitation of plant innate immunity processes (23,61–64).

Pathogen effectors often interfere with multiple host pathways to promote virulence (4) and our results, along with the pre-existing literature, conjointly point out that RipE1 seems to be implicated in several T3E functions (49–51). Our findings additionally provide new putative RipE1 subcellular targets and we hope that they advance the discovery of more functions of this conserved *R. solanacearum* effector.

For the purposes of this study, we further investigated the interaction between RipE1 and Exo70-ID, aiming to uncover an effector target that belongs to the host secretory pathway. Based on our previously performed Blastp analysis using the Exo70-like ID as a query (34), we hypothesized that *At*Exo70B1, an Exo70-containing *Arabidopsis thaliana* protein and a component of the plant exocyst complex, is most likely to be targeted by RipE1. We proceeded by testing the interaction between Exo70B1 and RipE1 in yeast. As shown in **Figure 2B**, Exo70B1 not only interacted with RipE1, but also with two mutants of the effector, the RipE1 C172A, carrying a C172A substitution at the active site and the RipE1 ΔA lacking eight conserved amino acids (121-128) found in the N-termini of the HopX1 effector family, also known as the A-box (51, 55). The two mutations were designed according to the structural predictions presented in **Figure 1**, but also due to their previously reported inability to induce plant immunity responses in model plants (51).

RipE1 interacts with *At*Exo70B1 *in planta*

Although GMI1000-RipE1 has been shown to trigger a hypersensitive response in *Nicotiana benthamiana*, *Nicotiana tabacum* and induce defense responses in *Arabidopsis thaliana* (50, 51), the quite similar (~82%) in sequence RS1000-RipE1 did not trigger such responses when transiently expressed in *Nicotiana benthamiana* leaves (49). In this study, we screened various *Nicotiana* species, including various *Nicotiana sylvestris* ecotypes, to find a new plant model for our *in-planta* experiments. We discovered that GMI1000-RipE1 (hereafter RipE1) did not trigger an HR response when transiently expressed in leaves of the *Nicotiana sylvestris* ecotype NS2706 from our collection, via *Agrobacterium*-mediated infiltration (**Fig. S1B**) and thus, we used it as a novel tool and performed all our further experiments on this ecotype. To test the possible colocalization of RipE1 and Exo70B1 *in planta*, we transiently expressed both proteins fused to fluorescent tags, -YFP and -mCherry respectively, in *Nicotiana sylvestris* leaf tissue, observed at 48hpi via confocal microscopy. As shown in **Figure 3A**, RipE1 and Exo70B1 colocalized in the periphery of the plant cells, an observation supported by the *in-silico* colocalization analysis presented in **Figure 3B**. To further investigate the interaction between RipE1 and Exo70B1 *in planta*, we used Bimolecular Fluorescent Complementation (BiFC) assay, transiently co-expressing RipE1 tagged with C-terminal YFP (cCFP) and Exo70B1 tagged with N-terminal YFP (nVenus). YFP signal was detected in the cell periphery via confocal microscopy 48hpi, as shown in **Figure 3D**, indicating the association between the two proteins *in planta*. YFP signal was also detected when the two RipE1 mutants -RipE1 C172A and RipE1 ΔA- were used (**Fig. S2**). Deletion of the A-box, had previously been shown to abolish cell death induction by RipE1 in *N. benthamiana* (51). This domain, although conserved in the HopX1/AvrPphE effector family, is of unknown function and has been earlier proposed as a domain that facilitates effector-target interactions (54, 55). Our results show that deletion of this 8-amino acid domain does not alter interaction capacity between RipE1 and Exo70B1 (**Figure 2B, Figure S2**) and thus, its role in effector function remains to be elucidated. We additionally performed co-immunoprecipitation assays to validate this interaction.

Exo70B1-HF pull-down did precipitate RipE1-myc, after their *Agrobacterium*-mediated

transient co-expression (**Fig. 3C**), advocating the association between the two proteins in

the plant cells. These findings are derived from the Y2H screening presented and

discussed above (**Fig. 2**, **Table S1**). After further investigation of the interaction between

RipE1 and Exo70-ID, we established *Arabidopsis thaliana* exocyst component Exo70B1

as a subcellular target of RipE1, highlighting the benefits of employing NLR-ID domains

as the means to uncover new potential targets of pathogen effectors.

RipE1 cleaves *At*Exo70B1 *in vitro*

To expand our understanding of the interaction between Exo70B1 and RipE1, we

predicted and visualized the interaction *in silico* (**Fig. 4A**). Based on this, the Exo70B1

146-201 aa region appears to approach the catalytic triad of RipE1 cysteine protease. This

observation, along with the interaction between the two proteins (**Fig. 2B**, **Fig. 3**),

indicate that Exo70B1 is a substrate for RipE1. In order to test the proteolytic effect of

RipE1 on Exo70B1, *in vitro* cleavage assays were performed with the purified proteins

(**Fig.4B**). Purified 6xHis-SUMO3-Exo70B1 was incubated with RipE1-6xHis, with and

without the addition of cysteine protease inhibitor leupeptin. Each protein of interest and

the 6xHis-SUMO3 tag were all incubated separately, to serve as negative controls for the

assay. The reactions were analyzed via SDS-PAGE, followed by anti-his-tag western

blotting (**Fig. 4B**). RipE1 appeared to have autoproteolytic activity in all cases, including

in the presence of leupeptin. On the other hand, a band representing cleaved 6xHis-

SUMO3-Exo70B1 (see **Fig. 4B**, orange box) was detected only in absence of this

inhibitor and in presence of RipE1 (**Fig. 4B**).

The *in-vitro* cleavage assay was repeated, after His:SUMO3 tag was removed from

Exo70B1. The reaction was analyzed via SDS-PAGE, followed by Silver staining (**Fig**.

4C, left). Based on the cleaved 6xHis-SUMO3-Exo70B1 (**Fig. 4B**) band molecular

weight and with the His-SUMO3 tag having been removed, three zones were selected to

be further analyzed via mass spectrometry, marked by the arrows on **Figure 4C**. The

analysis of the MS-derived data identified peptides of Exo70B1 present in these zones, as

indicated by the grey areas (**Fig. 4C**, right). These findings confirmed that Exo70B1 is

cleaved by RipE1 *in vitro*, between the particular sequence.

RipE1 promotes the degradation of Exo70B1 *in planta*

To investigate the possible degradation of Exo70B1 by RipE1 *in planta*, we transiently

expressed the proteins of interest in *N. sylvestris* tissue via *Agrobacterium*-mediated

infiltration. Proteins were extracted and analyzed via SDS-PAGE and immunoblotting

(**Fig. 5A**) and a total of three replications were used for *in-silico* quantification of

proteins band intensity (**Fig. 5B**). In agreement with the observations regarding *in vitro*

cleavage (**Fig. 4**), expression of RipE1 seemed to decrease Exo70B1 protein levels after

their transient co-expression *in planta*, in a statistically significant manner (**Fig. 5B**).

Since we had used agrobacteria carrying empty vector (EV) to balance inoculum OD_{600},

we additionally compared RipE1 levels in these experiments (**Fig. S3A**). In the case of

RipE1 levels, no significant difference was detected (**Fig. S3A**), ruling out possible

reduction of Exo70B1 due to co-expression of two constructs at the same plant tissue.

There is strong evidence that Exo70B1 is guarded by the TN2 (TIR-NBS2) NLR in

Arabidopsis in a CPK5 (Calcium-dependent Protein Kinase 5)-dependent manner and

that Exo70B1 degradation induces NLR-mediated immunity responses (30, 39, 59). TN2

expression in *Nicotiana tabacum* triggers a strong cell death response, suppressed by co-

expression with Exo70B1 and subsequently rescued by the *Pseudomonas syringae*

effector AvrPtoB, which induces Exo70B1 degradation (30). Based on our findings so

far, we hypothesized that Exo70B1 degradation by RipE1 would be able to induce TN2-

dependent cell death response. We observed that TN2 also triggers a cell death response

in *Nicotiana sylvestris* that was successfully suppressed by co-expression of Exo70B1

(**Fig. 5C-5F**). In accordance with our hypothesis, co-expression of TN2 and Exo70B1

with RipE1 did rescue TN2-mediated HR in *N. sylvestris* (**Fig. 5C-5D**), suggesting that

RipE1 promotes Exo70B1 degradation *in planta*. When the RipE1 mutant C172A was

used to perform similar assays, RipE1 C172A failed to rescue TN2-mediated HR,

indicating that this rescue is dependent on the effector's cysteine protease activity (**Fig.**

5E-5F). This NLR activation could possibly cause the previously observed C172-

dependent induction of immune responses by RipE1 in Arabidopsis (51), however, this
hypothesis requires further investigation. These results and the so-far published data
showcase that RipE1 function as a virulence or avirulence factor varies among different
strains and hosts and could promote the discovery of one or more existing plant NLRs
that can recognize RipE1, thus contributing to pursuing resistance against *R.*
solanacearum. To supplement our data on Exo70B1 degradation by RipE1 *in planta*, we
transiently expressed theExo70B1 simultaneously fused to two fluorescent tags. N-
terminal YFP and C-terminal mCherry fluorescent epitopes were used as tags for
Exo70B1, respectively. FRET efficiency was measured between the two epitopes, after
transient co-expression of the described construct, along with RipE1 or GUS (**Fig. S3**).
Although the difference was not that obvious to the eye in the FRET efficiency images
(**Fig. S3A**), our *in-silico* quantification revealed a statistically significant decrease in
FRET efficiency between YFP and mCherry when RipE1 was present, compared to the
GUS control (**Fig. S3B**).

Conclusion

Overall, the evidence presented here heavily supports the fact that RipE1 targets and
cleaves Arabidopsis exocyst component Exo70B1, possibly adding the manipulation of
the host secretion machinery by RipE1 to the pathogen's arsenal of virulence means.
However, this plant protein is also involved in other eukaryotic pathways and given the
displayed Exo70B1 embroilment in plant autophagy (40), whether cleavage by RipE1
aims to interfere with vesicle trafficking or other host cell processes needs to be
elucidated.

Materials and Methods

In silico protein structure prediction and analysis

The 3D structure of the full length RipE1 protein was predicted using the AlphaFold Colab (66,67). Subsequently, the predicted structure was compared with those in the Protein Data Bank via the Dali (58) online server. The visualization of the protein structures was performed by PyMOL (68).

Plasmid construction

Genes of interest were amplified using Phusion® High-Fidelity DNA Polymerase (NEB) and PCR products were purified with the Nucleospin® Gel and PCR Clean-up kit MACHEREY-NAGEL). All constructs used in this study were generated via Golden Gate (GG) Cloning, using the following plasmids as backbone vectors: pGADT7:RFP, pGBKT7:RFP for yeast and pICH86988, pICH86966 for plant assays. RipE1 was amplified from *Ralstonia solanacearum* genomic DNA. *R. solanacearum* GMI1000 was kindly donated by Prof. Nemo Peeters (INRA-CNRS) and gDNA was extracted using the CTAB method (69). RipE1 mutations were designed according to structural predictions (**Figure 1**) and recent studies (49,51) and were generated through PCR followed by GG Cloning, using the primers listed in **Table S2**. Exo70B1 and NLR-ID containing constructs were generated as described earlier in a recent study from our lab (34). pMDC:spC7HPB for *in planta* transient expression of Chp7 was a kind donation of Prof. Jane Glazebrook (70).All constructs generated in this study were verified by Sanger sequencing.

Bacterial strains and growth conditions

Escherichia coli (Stellar or DH10b strains) were cultured in LB medium (1% tryptone, 0.5% yeast extract, 1% sodium chloride for liquid cultures and additionally 1,5% agar for plates) containing appropriate antibiotics: ampicillin 100μg/ml or kanamycin 50μg/ml. *E. coli* cells were grown either on LB plates or in liquid LB medium at 37°C for 16h. For

liquid cultures, shaking was applied at approximately 200 rpm. *Agrobacterium*
tumefaciens (AgL-1, C58C1 or GV3101 strains) were also cultured in LB medium
containing appropriate antibiotics: rifampicin 50μg/ml and ampicillin 100μg/ml,
kanamycin 50μg/ml or gentamycin 10μg/ml. *Ralstonia solanacearum* strain GMI1000
was cultured in NB/NA medium (0.5% tryptone, 0.3% yeast extract, 0.5% sodium
chloride for liquid cultures and additionally 1,5% agar for plates) and grown at 28° for 2
days. For liquid cultures, shaking was applied at approximately 200 rpm.

Y2H assay

All plant proteins were cloned as baits (in the pGBKT7-RFP vector) and bacterial
effectors were cloned as preys (in the pGADT7-RFP vector), or as otherwise stated.
Different pairs of constructs were co-transformed into *Saccharomyces cerevisiae* strain
PJ694a (71), which carries the following auxotrophic traits: gal4Δ, gal80Δ, trp1-901
leu2-3 ura3-52 his3-200, *LYS2::GAL1-HIS3*, *GAL2-ADE2*, *met2::GAL7-lacZ.* At least
500ng from each plasmid (bait and prey) was used to transform *S. cerevisiae* PJ694a
competent cells, freshly prepared with the lithium acetate method (72). Transformed cells
were selected on SD–Leu–Trp medium (SD– LW) agar plates. Two single clones were
transferred from the SD–LW medium and were re-cultured on SD–LW (for verification
of growth) and SD–Leu–Trp–Ade (–LWA, auxotrophic selection) medium. For the
dilutions, a single clone was transferred to liquid SD–LW medium, grown until saturation
and diluted as stated in the figures, after OD_{600} measurements. Yeast cells in this study
were cultured in YPDA rich medium or SD-LW and SD-LWA minimal media at 28-
30°C.

Plant material

The *Nicotiana sylvestris* plants from various ecotypes that were used in our study were
grown in greenhouse conditions. For *in planta* transient expression via *Agrobacterium*-
mediated infiltration, 5-week-old *N. sylvestris* NS2706 plants were used. Agrobacteria
carrying indicated constructs were grown in liquid cultures until saturation and cell

pellets were diluted in 10mM MgCl2 and 10mM MES. Agrobacteria carrying each

construct were then used at an OD_{600} score of 0.5 to infiltrate *N. sylvestris* leaf areas,

while agrobacteria carrying empty vector were used to equalize the total OD_{600} score in

all inoculums of each experiment.

Confocal microscopy

All confocal images were captured in a Leica SP8 laser scanning confocal microscope.

All leaf tissue samples were imaged with a 40x water objective, after *Agrobacterium-*

mediated transient expression at 48-72hpi. The confocal images were processed with

ImageJ. For colocalization, all indicated proteins were fused to YFP or mCherry

fluorescent epitopes. YFP was excited using a 514-nm argon laser and emission was

detected at 520–550 nm. mCherry was excited using a 561-nm laser and emission was

detected at 593–628nm. The colocalization analyses were performed with PSC using the

Coloc2 addon in Fiji (73,74). The threshold regression used was Costes, PSF 100 and

Costes randomizations 10 (75). A total of 15 regions of interest (ROIs) were chosen for

the PSC analysis. For BiFC, all indicated peoreins were fused to C-terminal nVENUS or

cCFP epitope tags and subsequently, YFP signal was detected. Chlorophyll was detected

between 653 and 676 nm and excited at 561 nm. Leica software LASX was used to add

scale bars to the images and to display the z-stack maximum projection. Details of the

FRET-SE method are described in the Confocal Application Notes Pdf prepared by

Myriam Gastard, Ph.D. Exton Support Group, Leica Microsystems, Inc. (76,77). For

calculation of FRET efficiency, Method 2 was used (76,77). For quantification of FRET-

SE signal, at least 300 counts were measured using FIJI v. 1.49 software

(https://imagej.nih.gov/) and the measurements were submitted to were submitted to t-test

followed by Welch's correction on GraphPad Prism version 8.0.0 for Windows,

GraphPad Software, San Diego, California USA, (www.graphpad.com). Differences with

p value <0.05 were considered statistically significant (*), while multiple asterisks were

used to indicate profoundly low p value (as stated in the corresponding figure legends).

Protein extraction from plant leaf tissue

Agroinfiltrated leaf tissue was ground at 48hpi in liquid nitrogen and proteins were extracted in GTEN buffer (10% glycerol, 25-mM Tris–Cl pH 7.5, 1-mM ethylenediaminetetraacetate, 150-mM NaCl), supplemented with fresh 5-mM DTT, protease inhibitor cocktail (Calbiochem) and 0.15% v/v NP-40, applying vigorous shaking. After a 10-minute centrifugation at high speed (14,000g) at 4°C, the supernatant was collected in a clean microcentrifuge tube and was used for further protein analysis (co-immunoprecipitation, SDS-PAGE and immunoblotting).

Co-Immunoprecipitation

Protein extract from leaf tissue was incubated for 2h at 4°C with an anti-flag mouse antibody at 1:4,000 dilution (Sigma, F1804), followed by a 1h-incubation at 4°C with PureProteomeProtein A/G Magnetic Beads Mix (Millipore). Beads were washed 3 times with GTEN buffer, before proceeding to SDS-PAGE and immunoblotting.

SDS PAGE and Immunoblotting

Protein samples from plant leaves were prepared for SDS-PAGE electrophoresis by adding 1x SDS loading dye and boiling the samples for 5 min. The samples were then separated via 8% or 12% SDS–PAGE (stated in images captions), electroblotted to a PVDF membrane (Millipore), then blocked for 1h with 5% w/v milk in Tris-buffered saline (TBS; 150-mM NaCl, 50- mM Tris–HCl, pH 7.6) and incubated for 1h with primary antibodies (1:4,000 dilution, anti-flag mouse -Sigma #F1804 or anti-myc mouse -Cell Signaling #2276). Membranes were then incubated for 1h with a secondary antibody (anti-mouse HRP conjugate at 1:10,000 dilution -Cell Signaling) and chemiluminescent substrate ultra (Millipore) was applied for 5 min before camera detection in Sapphire Biomolecular Imager (Azure Biosystems). For protein band intensity analysis, gel lanes (both CBB staining and immunoblot) from 3 separate experimental replications were selected and analyzed in FIJI v. 1.49 software (https://imagej.nih.gov/). Band intensity measurements were then submitted to were

submitted to t-test followed by Welch's correction on GraphPad Prism version 8.0.0 for

Windows, GraphPad Software, San Diego, California USA, (www.graphpad.com).

Differences with p value <0.05 were considered statistically significant (*), while

multiple asterisks were used to indicate profoundly low p value (as stated in the

corresponding figure legends).

Protein production and purification

RipE1 gene was cloned and inserted into the pET26b (ColE1 plasmids) vector carrying a

C-terminal 6-His tag and transformed into the *E. coli* strain BL21. A sufficient amount of

soluble protein was obtained after induction using the following conditions: Cells were

grown in LB medium containing 30μg/ml kanamycin until an OD_{600} value of 0.6-0.8 was

reached. The culture was induced with 0.5 mM IPTG and left to grow at 20°C overnight.

The cell paste was re-suspended in 100 ml lysis buffer containing 25 mM Tris pH 8.0,

300 mM NaCl, 5 mM imidazole and 15 mM β-mercaptoethanol, and homogenized. After

adding protease inhibitors (20 μg/ml leupeptin, 1 mM PMSF and 150 μg/ml

benzamidine), the cells were disrupted with sonication in ice, 10 sonication cycles of 30 s

each, with cooling intervals of 30 s. The precipitate was subsequently removed by

centrifugation at 12,000 rpm at 4°C for 45 min. Purification was carried out using His-tag

affinity chromatography at 4°C with a 8 ml Ni-NTA Qiagen column pre-equilibrated in

lysis buffer and initially washed stepwise with 10, 20 and 30 mM imidazole. With a

subsequent increase in imidazole concentration the protein started eluting at 100 mM

imidazole. Fractions containing the protein were dialyzed against the storage buffer

containing 25 mM Tris pH 8.0, 100 mM NaCl and 10 mM β-mercaptoethanol. In case of

EXO70B1, the full-length protein was cloned and inserted into the LIC 1.10 vector

carrying a N-terminal 6-His-SUMO3 tag, and transformed into the *E. coli* strain

Rosetta™(DE3)**.** The same protocol was followed and the fractions containing the protein

were dialyzed against the storage buffer (25 mM Tris pH 8.0, 100 mM NaCl, 10 mM β-

mercaptoethanol and 0.5 nM SENP-2 protease). Subsequently, a reverse purification

protocol was carried out using His-tag affinity chromatography. The elution fractions

containing the proteins, were polled and concentrated to a final volume of 2 ml. Size

exclusion chromatography was performed in 20°C ÄKTA purifier system (Amersham) and a prepacked Hi-Prep 16/60 Sephacryl S-200 high-resolution column (GE Healthcare). Flow rate was 0.5 ml/min, and elution was monitored at 280 nm. Using a 2ml-loop the proteins were loaded, and the 2ml fractions were collected and analyzed via a 10% SDS-polyacrylamide gel.

In vitro cleavage assay

Purified *A. thaliana* Exo70B1 was incubated with purified RipE1 in reaction buffer (25 mM Tris pH 8.0, 100 mM NaCl), for 3 h at room temperature. For the inhibitory assay, RipE1 was incubated with 100 μM leupeptin prior the cleavage assay, at room temperature for 1 h before adding the substrate. After the indicated times, the reaction was stopped by the addition of SDS sample buffer, and the samples were subjected to SDS-polyacrylamide gel electrophoresis and analyzed by Western blotting.

NanoLC-MS/MS Analysis

The nanoLC-MS/MS analysis was performed on an EASY-nLC II system (Thermo Scientific) coupled with an LTQ-Orbitrap XL ETD (Thermo Scientific) through a nanoES ion source (Thermo). Data were acquired with Xcalibur software (LTQ Tune 2.5.5 sp1, Thermo Scientific). Prior to the analysis, the mass spectrometer was calibrated with a standard ESI positive ion calibration solution of caffeine (Sigma), L-methionyl-arginyol-phenylalanylalanine acetate H2O (MRFA, Research Plus, Barnegat, NJ), and perfluoroalkyl triazine (Ultramark 1621, Alfa Aesar, Ward Hill, MA). Samples were reconstituted in 0.5% formic acid, and the tryptic peptide mixture was separated on a reversed-phase column (Reprosil Pur C18 AQ, particle size = 3 μm, pore size = 120 □ (Dr. Maisch, AnaLab, Athens, Greece), fused silica emitters 100 mm long with a 75 μm internal diameter (New Objective) packed in-house using a pressurized (35 to 40 bars of helium) packing bomb. The nanoLC flow rate was 300 nl min−1. The LC mobile phase consisted of 0.5% formic acid in water (A) and 0.5% formic acid in acetonitrile (B). A multi-step gradient was employed, from 5% to 30% B in 120 min, to 90% B in 10 min.

After the gradient had been held at 90% B for 5 min, the mobile phase was re-
equilibrated at initial gradient conditions. The MS was operated with a spray voltage of
2300 V, a capillary voltage of 35 V, a tube lens voltage of 140 V, and a capillary
temperature of 180 °C. A survey scan was acquired in the range of m/z 400–1800 with an
AGC MS target value of 106 (resolving power of 60,000 at m/z 400). The 10 most
intense precursor ions from each MS scan were subjected to collision-induced
dissociation in the ion trap.

Data Analysis of MS/MS-derived Data

The MS raw data were loaded in Proteome Discoverer 2.2.0 (Thermo Scientific) and run
using Mascot 2.3.02 (Matrix Science, London, UK) and Sequest HT (Thermo Scientific)
search algorithms against the EXO70B1 Sequence. A list of common contaminants was
included in the database. For protein identification, the following search parameters were
used: precursor error tolerance = 10 ppm, fragment ion tolerance = 0.8 Da, trypsin full
specificity, maximum number of missed cleavages = 3, and cysteine alkylation as a fixed
modification, Oxidation of Methionine, Acetylation of N-term, Phosphorylation (The
node ptmRS was used) of Serine, Threonine and Tyrosine as Variable Modifications. To
calculate the protein false discovery rate, a decoy database search was performed
simultaneously with strict criteria set to 0.01 and relaxed criteria to 0.05.

Statistical analysis and plotting of Hypersensitive Response (HR) assay results

Statistical analysis and plotting were performed using GraphPad Prism version 8.0.0 for
Windows, GraphPad Software, San Diego, California USA (www.graphpad.com). For
HR assays, a score 0-5 was assigned to each observed phenotype, representing cell death
intensity. HR scores from 12 replicates were submitted to t-test followed by Welch's
correction. Differences with p value <0.05 were considered statistically significant (*),
while multiple asterisks were used to indicate profoundly low p value (as stated in the
corresponding figure legends).

Author contributions

P.F.S. designed the research and supervised the project; D.T. carried out the experiments; K.K. purified recombinant proteins and performed in vitro protein assays and performed the AlphaFold analysis; V.A.M. supervised the FRET analysis and the confocal microscopy experiments. M.K. and V.A.M. contributed experimental materials. P.F.S., D.T. and K.K. wrote the manuscript. P.F.S. and M.K. edited the manuscript. All authors have read and approved the manuscript.

References

1. Duxbury Z, Ma Y, Furzer OJ, Huh SU, Cevik V, Jones JDG, et al. Pathogen perception by NLRs in plants and animals: Parallel worlds. BioEssays. 2016;38(8):769–81.

2. Mermigka G, Amprazi M, Mentzelopoulou A, Amartolou A, Sarris PF. Plant and Animal Innate Immunity Complexes: Fighting Different Enemies with Similar Weapons. Trends Plant Sci [Internet]. 2019;1–12. Available from: https://doi.org/10.1016/j.tplants.2019.09.008

3. Lal NK, Thanasuwat B, Huang P jui, Cavanaugh KA, Carter A, Michelmore RW, et al. Phytopathogen Effectors Use Multiple Mechanisms to Manipulate Plant Autophagy. Cell Host Microbe [Internet]. 2020;28(4):558-571.e6. Available from: https://doi.org/10.1016/j.chom.2020.07.010

4. Khan M, Seto D, Subramaniam R, Desveaux D. Oh, the places they'll go! A survey of phytopathogen effectors and their host targets. Plant J. 2018;93(4):651–63.

5. Yuan M, Jiang Z, Bi G, Nomura K, Liu M, He SY, et al. Pattern-recognition receptors are required for NLR-mediated plant immunity. Nature [Internet]. 2021;(April 2020). Available from: http://dx.doi.org/10.1038/s41586-021-03316-6

6. Yuan M, Ngou BPM, Ding P, Xin XF. PTI-ETI crosstalk: an integrative view of plant immunity. Curr Opin Plant Biol [Internet]. 2021;62:102030. Available from: https://doi.org/10.1016/j.pbi.2021.102030

7. Jones JDG, Dangl JL. The plant immune system. Nature. 2006;444:323–9.

8. Lee JH, Kim H, Chae WB, Oh MH. Pattern recognition receptors and their interactions with bacterial type III effectors in plants. Genes and Genomics [Internet]. 2019;41(5):499–506. Available from: http://dx.doi.org/10.1007/s13258-019-00801-1

9. Kapos P, Devendrakumar KT, Li X. Plant NLRs: From discovery to application. Plant Sci. 2019;279:3–18.

10. Li X, Kapos P, Zhang Y. NLRs in plants. Curr Opin Immunol [Internet]. 2015;32:114–21. Available from: http://dx.doi.org/10.1016/j.coi.2015.01.014

11. Camagna M, Takemoto D. Hypersensitive Response in Plants. eLS. 2018;1–7.

12. Cesari S. Multiple strategies for pathogen perception by plant immune receptors. New Phytol. 2018;219:17–24.

13. van der Hoorn RAL, Kamoun S. From Guard to Decoy: A New Model for Perception of Plant Pathogen Effectors. Plant Cell [Internet]. 2008;20(8):2009–17. Available from: http://www.plantcell.org/cgi/doi/10.1105/tpc.108.060194

14. Baggs E, Dagdas G, Krasileva K V. NLR diversity, helpers and integrated domains: making sense of the NLR IDentity. Curr Opin Plant Biol [Internet]. 2017;38(Figure 1):59–67. Available from: http://dx.doi.org/10.1016/j.pbi.2017.04.012

15. Sarris PF, Cevik V, Dagdas G, Jones JDG, Krasileva K V. Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens. BMC Biol [Internet]. 2016;14(1). Available from: http://dx.doi.org/10.1186/s12915-016-0228-7

16. Schreiber KJ, Chau-Ly IJ, Lewis JD. What the wild things do: Mechanisms of plant host

manipulation by bacterial type III-secreted effector proteins. Microorganisms.
2021;9(5):1–48.

17.   Saeed B, Brillada C, Trujillo M. Dissecting the plant exocyst. Curr Opin Plant Biol
[Internet]. 2019;52:69–76. Available from: https://doi.org/10.1016/j.pbi.2019.08.004

18.   Mei K, Guo W. The exocyst complex. Curr Biol [Internet]. 2018;28(17):R922–5.
Available from: http://dx.doi.org/10.1016/j.cub.2018.06.042

19.   TerBush DR, Maurice T, Roth D, Novick P. The Exocyst is a multiprotein complex
required for exocytosis in Saccharomyces cerevisiae. EMBO J. 1996;15(23):6483–94.

20.   Elias M, Drdova E, Ziak D, Bavlnka B, Hala M, Cvrckova F, et al. The exocyst complex
in plants. Cell Biol Int. 2003;27(3):199–201.

21.   Picco A, Irastorza-Azcarate I, Specht T, Böke D, Pazos I, Rivier-Cordey AS, et al. The In
Vivo Architecture of the Exocyst Provides Structural Basis for Exocytosis. Cell.
2017;168(3):400-412.e18.

22.   Gu Y, Zavaliev R, Dong X. Membrane Trafficking in Plant Immunity. Mol Plant
[Internet]. 2017;10(8):1026–34. Available from:
http://dx.doi.org/10.1016/j.molp.2017.07.001

23.   Martin-Urdiroz M, Deeks MJ, Horton CG, Dawe HR, Jourdain I. The exocyst complex in
health and disease. Front Cell Dev Biol. 2016;4(APR):1–22.

24.   Du Y, Overdijk EJR, Berg JA, Govers F, Bouwmeester K. Solanaceous exocyst subunits
are involved in immunity to diverse plant pathogens. J Exp Bot. 2018;69(3):655–66.

25.   Peenková T, Hála M, Kulich I, Kocourková D, Drdová E, Fendrych M, et al. The role for
the exocyst complex subunits Exo70B2 and Exo70H1 in the plant-pathogen interaction. J
Exp Bot. 2011;62(6):2107–16.

26.   Žárský V, Sekereš J, Kubátová Z, Pečenková T, Cvrčková F. Three subfamilies of exocyst
EXO70 family subunits in land plants: early divergence and ongoing functional
specialization. J Exp Bot. 2020;71(1):49–62.

27.   Stegmann M, Anderson RG, Westphal L, Rosahl S, McDowell JM, Trujillo M. The
exocyst subunit Exo70B1 is involved in the immune response of Arabidopsis thaliana to
different pathogens and cell death. Plant Signal Behav. 2014;8(12):1–5.

28.   Wang W, Liu N, Gao C, Cai H, Romeis T, Tang D. The Arabidopsis exocyst subunits
EXO70B1 and EXO70B2 regulate FLS2 homeostasis at the plasma membrane. New
Phytol. 2020;

29.   Brillada C, Teh O, Ditengou FA, Lee C, Klecker T, Furlan G, et al. Exocyst subunit
Exo70B2 is linked to immune signalling and autophagy. 2020;

30.   Wang W, Liu N, Gao C, Rui L, Tang D. The Pseudomonas Syringae Effector AvrPtoB
Associates With and Ubiquitinates Arabidopsis Exocyst Subunit EXO70B1. Front Plant
Sci [Internet]. 2019 Aug 29;10(August):1–16. Available from:
https://www.frontiersin.org/article/10.3389/fpls.2019.01027/full

31.   Du Y, Mpina MH, Birch PRJ, Bouwmeester K, Govers F. Phytophthora infestans RXLR
effector AVR1 interacts with exocyst component Sec5 to manipulate plant immunity.
Plant Physiol. 2015;169(3):1975–90.

32.   Petre B, Contreras MP, Bozkurt TO, Schattat MH, Sklenar J, Schornack S, et al. Host-
587   interactor screens of Phytophthora infestans RXLR proteins reveal vesicle trafficking as a
588   major effector-targeted process. Plant Cell. 2021;33(5):1447–71.

589   33.   Redditt TJ, Chung E-H, Zand Karimi H, Rodibaugh N, Zhang Y, Trinidad JC, et al.
590   AvrRpm1 Functions as an ADP-Ribosyl Transferase to Modify NOI-domain Containing
591   Proteins, Including Arabidopsis and Soybean RPM1-interacting Protein 4. Plant Cell.
592   2019;tpc.00020.2019.

593   34.   Michalopoulou VA, Mermigka G, Kotsaridis K, Mentzelopoulou A, Celie PHN, Moschou
594   PN, et al. The host exocyst complex is targeted by a conserved bacterial type III effector
595   protein that promotes virulence. Plant Cell [Internet]. 2022;00:1–25. Available from:
596   https://www.biorxiv.org/content/10.1101/2020.11.06.371260v1%0Ahttps://www.biorxiv.o
597   rg/content/10.1101/2020.11.06.371260v1.abstract

598   35.   Afzal AJ, da Cunha L, Mackey D. Separable fragments and membrane tethering of
599   Arabidopsis RIN4 regulate its suppression of PAMP-triggered immunity. Plant Cell.
600   2011;23(10):3798–811.

601   36.   Stegmann M, Anderson RG, Ichimura K, Pecenkova T, Reuter P, Žárský V, et al. The
602   ubiquitin ligase PUB22 targets a subunit of the exocyst complex required for PAMP-
603   triggered responses in arabidopsis c w. Plant Cell. 2012;24(11):4703–16.

604   37.   Ostertag M, Stammler J, Douchkov D, Eichmann R, Hückelhoven R. The conserved
605   oligomeric Golgi complex is involved in penetration resistance of barley to the barley
606   powdery mildew fungus. Mol Plant Pathol. 2013;14(3):230–40.

607   38.   Sabol P, Kulich I, Žárský V. RIN4 recruits the exocyst subunit EXO70B1 to the plasma
608   membrane. J Exp Bot. 2017;68(12):3253–65.

609   39.   Zhao T, Rui L, Li J, Nishimura MT, Vogel JP, Liu N, et al. A Truncated NLR Protein,
610   TIR-NBS2, Is Required for Activated Defense Responses in the exo70B1 Mutant. PLoS
611   Genet. 2015;11(1):1–28.

612   40.   Kulich I, Pečenková T, Sekereš J, Smetana O, Fendrych M, Foissner I, et al. Arabidopsis
613   Exocyst Subcomplex Containing Subunit EXO70B1 Is Involved in Autophagy-Related
614   Transport to the Vacuole. Traffic. 2013;14(11):1155–65.

615   41.   Hofius D, Li L, Hafrén A, Coll NS. Autophagy as an emerging arena for plant–pathogen
616   interactions. Curr Opin Plant Biol. 2017;38:117–23.

617   42.   Leary AY, Savage Z, Tumtas Y, Bozkurt TO. Contrasting and emerging roles of
618   autophagy in plant immunity. Curr Opin Plant Biol [Internet]. 2019;52:46–53. Available
619   from: https://doi.org/10.1016/j.pbi.2019.07.002

620   43.   Fegan M, Prior P. How complex is the Ralstonia solanacearum species complex. In: Allen
621   C, Hayward AC, Prior P, editors. Bacterial wilt disease and the Ralstonia solanacearum
622   species complex. Saint Paul, USA: APS Press; 2005. p. 449–61.

623   44.   Landry D, González-Fuente M, Deslandes L, Peeters N. The large, diverse, and robust
624   arsenal of Ralstonia solanacearum type III effectors and their in planta functions. Mol
625   Plant Pathol. 2020;21(10):1377–88.

626   45.   Mansfield J, Genin S, Magori S, Citovsky V, Sriariyanum M, Ronald P, et al. Top 10 plant
627   pathogenic bacteria in molecular plant pathology. Mol Plant Pathol. 2012;13(6):614–29.

46. Meng F. The Virulence Factors of the Bacterial Wilt Pathogen Ralstonia solanacearum. J
Plant Pathol Microbiol. 2013;04(03):3–5.

47. Sabbagh CRR, Carrere S, Lonjon F, Vailleau F, Macho AP, Genin S, et al. Pangenomic
type III effector database of the plant pathogenic Ralstonia spp. PeerJ. 2019;2019(8):1–21.

48. Peeters N, Carrère S, Anisimova M, Plener L, Cazalé A-C, Genin S. Repertoire, unified
nomenclature and evolution of the Type III effector gene set in the Ralstonia
solanacearum species complex. BMC Genomics [Internet]. 2013;14(1):859. Available
from: http://bmcgenomics.biomedcentral.com/articles/10.1186/1471-2164-14-859

49. Nakano M, Mukaihara T. Comprehensive Identification of PTI Suppressors in Type III
Effector Repertoire Reveals that Ralstonia solanacearum Activates Jasmonate Signaling at
Two Different Steps. Int J Mol Sci. 2019;20(5992).

50. Jeon H, Kim W, Kim B, Lee S, Jayaraman J, Jung G, et al. Ralstonia solanacearum type
III effectors with predicted nuclear localization signal localize to various cell
compartments and modulate immune responses in nicotiana spp. Plant Pathol J.
2020;36(1):43–53.

51. Sang Y, Yu W, Zhuang H, Wei Y, Derevnina L, Yu G, et al. Intra-strain Elicitation and
Suppression of Plant Immunity by Ralstonia solanacearum Type-III Effectors in Nicotiana
benthamiana. Plant Commun [Internet]. 2020;1(4):100025. Available from:
https://doi.org/10.1016/j.xplc.2020.100025

52. Yu G, Xian L, Xue H, Yu W, Rufian JS, Sang Y, et al. A bacterial effector protein
prevents mapk-mediated phosphorylation of sgt1 to suppress plant immunity. PLoS
Pathog [Internet]. 2020;16(9):1–30. Available from:
http://dx.doi.org/10.1371/journal.ppat.1008933

53. Nakano M, Ichinose Y, Mukaihara T. Ralstonia solanacearum Type III Effector RipAC
Targets SGT1 to Suppress Effector-Triggered Immunity . Plant Cell Physiol.
2021;61(12):2067–76.

54. Gimenez-Ibanez S, Boter M, Fernández-Barbero G, Chini A, Rathjen JP, Solano R. The
Bacterial Effector HopX1 Targets JAZ Transcriptional Repressors to Activate Jasmonate
Signaling and Promote Infection in Arabidopsis. PLoS Biol. 2014;12(2).

55. Nimchuk ZL, Fisher EJ, Desveaux D, Chang JH, Dangl JL. The HopX (AvrPphE) Family
of Pseudomonas syringae Type III Effectors Require a Catalytic Triad and a Novel N-
Terminal Domain for Function . Mol Plant-Microbe Interact. 2007;20(4):346–57.

56. Fowler NJ, Williamson MP. The accuracy of protein structures in solution determined by
AlphaFold and NMR. Structure [Internet]. 2022;30(7):925-933.e2. Available from:
https://doi.org/10.1016/j.str.2022.04.005

57. Rawat A, Roy M, Jyoti A, Kaushik S, Verma K, Srivastava VK. Cysteine proteases:
Battling pathogenic parasitic protozoans with omnipresent enzymes. Microbiol Res.
2021;249(April):126784.

58. Holm L. Using Dali for Protein Structure Comparison. Methods Mol Biol. 2020;2112:29–
42.

59. Chatterjee D, Boyd CD, O'Toole GA, Sondermann H. Structural characterization of a
conserved, calcium-dependent periplasmic protease from Legionella pneumophila. J
Bacteriol. 2012;194(16):4415–25.

60. Kotsaridis K, Tsakiri D, Sarris PF. Understanding enemy's weapons to an effective
prevention: common virulence effects across microbial phytopathogens kingdoms. Crit
Rev Microbiol [Internet]. 2022;0(0):1–15. Available from:
https://doi.org/10.1080/1040841X.2022.2083939

61. Frederickson Matika DE, Loake GJ. Redox regulation in plant immune function. Antioxid
Redox Signal. 2014;21(9):1373–88.

62. Wani SH, Anand S, Singh B, Bohra A, Joshi R. WRKY transcription factors and plant
defense responses: latest discoveries and future prospects. Plant Cell Rep [Internet].
2021;40(7):1071–85. Available from: https://doi.org/10.1007/s00299-021-02691-8

63. Cheval C, Aldon D, Galaud JP, Ranty B. Calcium/calmodulin-mediated regulation of
plant immunity. Biochim Biophys Acta - Mol Cell Res [Internet]. 2013;1833(7):1766–71.
Available from: http://dx.doi.org/10.1016/j.bbamcr.2013.01.031

64. Tsuda K, Somssich IE. Transcriptional networks in plant immunity. New Phytol.
2015;206:932–47.

65. Liu N, Hake K, Wang W, Zhao T, Romeis T, Tang D. Calcium-dependent protein kinase5
associates with the truncated NLR protein TIR-NBS2 to contribute to exo70B1-mediated
immunity. Plant Cell. 2017;29(4):746–59.

66. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold
Protein Structure Database: massively expanding the structural coverage of protein-
sequence space with high-accuracy models. Nucleic Acids Res. 2022;50(D1):D439–44.

67. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
protein structure prediction with AlphaFold. Nature. 2021;596(7873):583–9.

68. The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.

69. Minas K, Mcewan NR, Newbold CJ, Scott KP. Optimization of a high-throughput CTAB-
based protocol for the extraction of qPCR-grade DNA from rumen fluid, plant and
bacterial pure cultures. FEMS Microbiol Lett. 2011;325(2):162–9.

70. Lu Y, Hatsugai N, Katagiri F, Ishimaru CA, Glazebrook J. Putative serine protease
effectors of clavibacter michiganensis induce a hypersensitive response in the apoplast of
nicotiana species. Mol Plant-Microbe Interact. 2015;28(11):1216–26.

71. James P, Halladay J, Craig EA. Genomic Libraries and a Host Strain Designed for Highly
Efficient Two-Hybrid Selection in Yeast. Genetics. 1996;(144):1425–36.

72. Ito H, Murata K, Kimura A. Transformation of Intact Yeast Cells Treated with Alkali
Cations or Thiol Compounds. Agric Biol Chem. 1984;48(2):341–7.

73. French AP, Mills S, Swarup R, Bennett MJ, Pridmore TP. Colocalization of fluorescent
markers in confocal microscope images of plant cells. Nat Protoc. 2008;3(4):619–28.

74. Moschou PN, Smertenko AP, Minina EA, Fukada K, Savenkov EI, Robert S, et al. The
caspase-related protease separase (EXTRA SPINDLE POLES) regulates cell polarity and
cytokinesis in arabidopsis. Plant Cell. 2013;25(6):2171–86.

75. Costes S V., Daelemans D, Cho EH, Dobbin Z, Pavlakis G, Lockett S. Automatic and
quantitative measurement of protein-protein colocalization in live cells. Biophys J
[Internet]. 2004;86(6):3993–4003. Available from:

712   http://dx.doi.org/10.1529/biophysj.103.038422

76.   Wouters FS, Verveer PJ, Bastiaens PIH. Imaging biochemistry inside cells. Trends Cell
Biol. 2001;11(5):203–11.

77.   Van Rheenen J, Langeslag M, Jalink K. Correcting Confocal Acquisition to Optimize
Imaging of Fluorescence Resonance Energy Transfer by Sensitized Emission. Biophys J
[Internet]. 2004;86(4):2517–29. Available from: http://dx.doi.org/10.1016/S0006-
3495(04)74307-6

78.   Smirnov MN, Smirnov VN, Budowsky EI, Inge-Vechtomov SG, Serebrjakov NG. Red
pigment of adenine-deficient yeast Saccharomyces cerevisiae. Biochem Biophys Res
Commun. 1967;27(3):299–304.

79.   Weisman LS, Bacallao R, Wickner W. Multiple methods of visualizing the yeast vacuole
permit evaluation of its morphology and inheritance during the cell cycle. J Cell Biol.
1987;105(4):1539–47.

Figures legends

Figure 1. Presentation of *R. solanacearum* GMI1000 RipE1 3D structure prediction. (A) 3D structure prediction using AlphaFold Colab (N-terminal is shown as blue, C-terminal is shown as red). The first 76 residues are presented as unfolded. The conserved catalytic triad (Cys-His-Asp, C-H-D) and the hydrogen bond between H203 and D222is shown bottom left. **(B)** Surface hydrophobicity (Red: Hydrophobic surface). **(C)** Crystal structure of *L. pneumophila* LapG. **(D)** Superimposition of LapG (cyan) and RipE1 (purple) proteins active sites.

Figure 2. RipE1 interacts with Exo70-like ID and *At*Exo70B1 in yeast. (A) Interactions between RipE1 and six NLR-ID domains, uncovered by yeast two-hybrid screening (for more details see **Table S1**). Yeast cells carrying the indicated baits and preys were spotted on SDC-LW (plasmid selection) and SDC-LWA plates (interaction selection). Photographs were taken after 4 days of incubation. The red color in some of the colonies is a result of accumulation of a derivative of 5-aminoimidazole ribotide (78,79). **(B)** Interactions between wild-type RipE1, RipE1 C172A mutant, RipE1 ΔA mutant and *Arabidopsis thaliana* Exo70B1. Serial dilutions of cells carrying the indicated baits and preys were spotted on SDC-LW and SDC-LWA selective plates. Initial OD_{600} of cells was 1 unit and subsequent dilutions were made 0.1x, 0.01x and 0.001x. Photographs were taken after 4 days of incubation.

Figure 3. RipE1 interacts with *At*Exo70B1 *in planta*. (A) Colocalization between RipE1 and Exo70B1 *in planta*. YFP-tagged RipE1 and mCherry-tagged Exo70B1 were transiently expressed in *Nicotiana sylvestris* leaves and detected via confocal microscopy 48hpi. YFP and mCherry signals were overlapping at the cell periphery. **(B)** Colocalization analysis, supporting **Figure 1A**. 15 ROIs were analyzed using Coloc2 in ImageJ and the Pearson's R and Spearman's R parameters were plotted to visualize colocalization probability. **(C)** Co-immunoprecipitation of RipE1 and Exo70B1 from plant tissue. Myc-tagged RipE1 and HF (6xHis and 3xFLAG)-tagged Exo70B1 transiently expressed in *N. sylvestris* leaves and an IP-flag assay was performed 48hpi. RipE1-myc is 52 kDa and Exo70B1-HF is 77 kDa. **(D)** *In planta* validation of the RipE1/Exo70B1 interaction using a BiFC assay in *N. sylvestris* plants. RipE1 and Exo70B1 were fused at their C-termini with cCFP and nVenus epitope tags, respectively, and the YFP signal was detected via confocal microscopy 48hpi (middle). A maximum projection of the z-stack is also presented (left). As negative control, RipE1-cCFP was co-expressed with RRS1-nVenus and no YFP signal was observed (right). Bars = 80μm.

Figure 4. RipE1 cleaves Exo70B1 between its sequence *in vitro*. (A). *In-silico* prediction of the interaction between Exo70B1 (cyan) and RipE1 (magenda). The boxed area appears magnified on the right and shows the Exo70B1 region (cyan) that approaches the catalytic triad of RipE1 (blue). **(B)** *In-vitro* cleavage of Exo70B1 by RipE1 is shown in this immunoblot. Reactions were analyzed via SDS-PAGE, followed by immunoblotting with His antibody. **(C)** Purified Exo70B1, after removal of 5xHis-SUMO3 tag, was incubated with RipE1-6xHis and the sample was analyzed via SDS-PAGE followed by Silver staining (left). Three selected SDS-PAGE zones, as indicated by arrows, were further analyzed by mass spectrometry. Grey regions on the right show the MS-identified Exo70B1 peptides that derived after cleavage by RipE1.

Figure 5. RipE1 promotes degradation of Exo70B1 *in planta*. (A) RipE1 expression promotes decrease in Exo70B1 protein levels in planta. These immunoblots support the statistical analysis presented in (B) and are one of the 3 replications performed for this analysis. RipE1 fused to Myc epitope and Exo70B1 fused to HF (6xHis and 3xFlag) epitope were transiently expressed in *Nicotiana sylvestris* leaves. Extracted proteins (48hpi) were analyzed via SDS-PAGE and immunoblotting to compare protein levels between the tissue samples expressing the single proteins (empty vector was used to balance agrobacteria density) and those co-expressing the interacting partners RipE1/Exo70B1. The presented analyzed samples have originated from the same leaf, to achieve comparability. Coomassie Brilliant Blue (CBB) staining images accompany each immunoblot and were used to normalize protein levels according to loaded protein quantity. **(B)** Statistical analysis of relative immunoblot band intensity indicating Exo70B1 protein levels extracted from plant tissue. This analysis was performed using 3 total experimental replicates, where total protein was extracted from *N. sylvestris* leaves, after *Agrobacterium*-mediated transient expression (48hpi) of Exo70B1 and Exo70B1 along with RipE1. Empty vector (EV) was used in the samples expressing only Exo70B1 to balance agrobacteria density. Compared relative band intensities (%) were submitted to t-test followed by Welch's correction. *p value=0.0284 **(C)** *Nicotiana sylvestis* leaf representing one of the 12 replicates used for the statistical analysis of hypersensitive response (HR) assays presented in (D). The indicated proteins were transiently expressed via *Agrobacterium*-mediated infiltration at a total OD_{600} of 0.5. The photograph was captured 3dpi. **(D)** Statistical analysis of the 12 HR assays replicates testing wild type RipE1 ability to rescue TN2-induced cell death phenotype via Exo70B1 degradation. Compared phenotypes were submitted to t-test followed by Welch's correction. ***p value=0.0003, *p value=0.0125. **(E)** *N. sylvestis* leaf representing one of the 12 replicates used for the statistical analysis of HR assays presented in (F). The indicated proteins were transiently expressed via *Agrobacterium*-mediated infiltration at a total OD_{600} of 0.5. The photograph was captured 3dpi. (+) on the upper left infiltrated area represents an additional cell-death inducing construct that was tested as a putative positive control (effector Chp7 fused to PR1 secretion peptide) and was not included in the statistical analysis of HR scores. **(F)** Statistical analysis of the 12 HR assays replicates testing mutated RipE1 C172A ability to rescue TN2-induced cell death phenotype via Exo70B1 degradation. Compared phenotypes were submitted to t-test followed by Welch's correction. **p value=0.0085, ns: no significance p value>0.05.

Supplementary Tables and Figures legends

Table S1. NLR-IDs that were tested for interaction with RipE1. NLR-ID domains originating from both dicot (*Arabidopsis thaliana*, *Brassica napus*) and monocot (*Oryza sativa*, *Hordeum vulgare*) plant genomes were used in a yeast two-hybrid screening to test their possible interaction with effector RipE1. Each putative interaction was tested at least 3 times.

(-) no interaction was detected

(+) a relatively weak interaction was detected, *not* detectable in all replications

(++) a relatively weak interaction was detected, detectable in all replications

(+++) a relatively strong interaction was detected, detectable in all replications

Table S2. List of primers used in this study. The primers listed here were designed to clone effector wild type RipE1 and its derived mutants RipE1 C172A and RipE1 ΔA, in Golden Gate-compatible vectors for expression in yeast (yeast two-hybrid assay, Y2H) and *Agrobacterium*-mediated transient expression in plant tissue.

Figure S1. Representative images supplementing main Figures 2 and 3. (A) Neither RipE1 or any of the NLR-IDs were able to induce expression of the reporter gene in a yeast two-hybrid assay, in the absence of an interacting protein. Empty bait and prey vectors, respectively, were used to test possible auto-activities. The results support the interactions in yeast, that are presented in **Figure 2**. **(B)** Wild type RipE1 or mutants C172A and ΔA do not induce a hypersensitive response (HR) in *Nicotiana sylvestris* selected ecotype NS2706. Effector Chp7 – fused to PR1 secretion peptide for apoplastic secretion-was used as a positive HR-inducing control and GUS reporter gene was used as a negative control. This photograph was taken 4dpi. This experiment was repeated at least 6 times with the same exact results. **(C)** Leaf discs from **(B)** were observed via confocal microscopy to confirm expression of proteins of interest that did not induce HR. All indicated proteins were c-terminally fused to YFP fluorescent epitope. These images were captured 5dpi. **(B) and (C)** collectively support the further use of *Nicotiana sylvestris* NS2706 as a model for our *in-planta* experiments.

Figure S2. RipE1 mutants associate with *At*Exo70B1 *in planta*. *In planta* validation of the RipE1 C712A/Exo70B1 and RipE1 ΔA/Exo70B1 association using a BiFC assay in *Nicotiana sylvestris* plants. RipE1 mutated proteins and Exo70B1 were fused at their C-termini with cCFP and nVenus epitope tags, respectively, and the YFP signal was detected via confocal microscopy 48hpi.

Figure S3. Images supplementing main Figure 5. (A) Statistical analysis of relative
immunoblot band intensity indicating RipE1 protein levels extracted from plant tissue. This
analysis was performed using 3 total experimental replicates, where total protein was extracted
from *N. sylvestris* leaves, after *Agrobacterium*-mediated transient expression (48hpi) of RipE1
and RipE1 along with Exo70B1. A representative immunoblot is shown in **Figure 5A**. Empty
vector (EV) was used to balance agrobacteria density in the samples expressing only RipE1.
Compared relative band intensities (%) were submitted to t-test followed by Welch's correction.
ns: no significance p value>0.05. **(B)** RipE1 expression caused a decrease in YFP-mCherry FRET
efficiency, compared with GUS control. YFP and mCherry fluorescent epitopes were N-
terminally and C-terminally, respectively, fused to Exo70B1. Two *N. sylvestris* leaf discs
transiently expressing this construct, along with either GUS reporter gene or effector RipE1, were
sampled to measure FRET efficiency via confocal microscopy. This figure was used for the
statistical analysis presented in (C). **(C)** Statistical analysis of FRET efficiency between. YFP and
mCherry fluorescent epitopes. Pixel intensity from at least 300 points was measured in FiJi and
the results were submitted to t-test followed by Welch's correction. ****p value=0.0001.
854

Figure 1

A

N

C

1   76   425

RipE1 *R. solanacearum* GMI1000

D²²²

C¹⁷²

H²⁰³

conserved
C-H-D
triad

B

conserved
C-H-D
triad

C

D

C¹⁷²
C¹³⁷

H²⁰³
H¹⁷²

D²²²
D¹⁸⁹

Figure 2

A

RipE1 prey x
TRX-like ID bait

RipE1 prey x
WRKY-like ID bait

RipE1 prey x
TFIIS-like ID bait

RipE1 prey x
Exo70-like ID bait

RipE1 prey x
CG-like ID bait

RipE1 prey x
HSF-like ID bait

RipE1 prey x
Ubox-like ID bait

SD
-LW

SD
-LWA

B

SD -LW

SD -LWA

RipE1 prey **x**
AtExo70B1 bait

RipE1 C172A prey **x**
AtExo70B1 bait

RipE1 ΔA prey **x**
AtExo70B1 bait

Empty prey **x**
AtExo70B1 bait

RipE1 prey **x**
empty bait

RipE1 C172A prey **x**
empty bait

RipE1 ΔA prey **x**
empty bait

Cell density

Figure 3

Figure 4

A

−	+	+	RipE1:myc
+	−	+	Exo70B1:HF

CBB

75 kDa — a-flag (Exo70B1, 77kDa)

63 kDa — a-myc (RipE1, 51kDa)

CBB

B

Relative band intensity (%)

*

+EV   +RipE1

Exo70B1

C

RipE1

TN2

TN2
+ RipE1

Exo70B1

TN2
+ Exo70B1
+ RipE1

TN2
+ Exo70B1

D

HR Score

***

*

TN2   TN2 + Exo70B1   TN2 + Exo70B1 + RipE1   Exo70B1   RipE1   TN2 + RipE1

E

RipE1 C172A

+

TN2

TN2
+ RipE1 C172A

TN2
+ Exo70B1

TN2
+ Exo70B1
+ RipE1 C172A

Exo70B1

F

HR Score

**

ns

TN2   TN2 + Exo70B1   TN2 + Exo70B1 + RipE1 C172A   Exo70B1   RipE1 C172A   TN2 + RipE1 C172A