CRTC-1 Balances Histone Trimethylation and Acetylation to Promote Longevity

Carlos G. Silva-García[1], Laura I. Láscarez-Lagunas[2], Caroline Heintz[1], Katharina Papsdorf[3], Lourdes Pajuelo[1], Arpit Sharma[1], Monica P. Colaiácovo[2], Anne Brunet[3,4] and William B. Mair[1*]

[1]Department of Molecular Metabolism, Harvard T. H. Chan School of Public Health, Harvard University, Boston, Massachusetts 02115, USA.
[2]Department of Genetics, Blavatnik Institute, Harvard Medical School, Boston, Massachusetts 02115, USA.
[3]Department of Genetics, Stanford University, 200 Jane Stanford Way, Stanford, California 94305, USA.
[4]Glenn Laboratories for the Biology of Aging, Stanford University, Stanford, California 94305, USA.

*Corresponding author: wmair@hsph.harvard.edu (WBM)

SUMMARY

Loss of function during ageing is accompanied by transcriptional drift, altering gene expression and contributing to a variety of age-related diseases. CREB regulated transcriptional coactivators (CRTCs) have emerged as key regulators of gene expression that might be targeted to promote longevity. Here, we define the role of the *Caenorhabditis elegans* CRTC-1 in the epigenetic regulation of longevity. Endogenous CRTC-1 binds chromatin factors, including components of the COMPASS complex, which trimethylates lysine 4 on histone H3 (H3K4me3). CRISPR editing of endogenous CRTC-1 reveals that the CREB-binding domain in neurons is specifically required for H3K4me3-dependent longevity. However, this effect is independent of CREB but instead acts via the transcription factor AP-1. Strikingly, CRTC-1 also mediates global histone acetylation levels, and this acetylation is essential for H3K4me3-dependent longevity. Indeed, overexpression of an acetyltransferase enzyme is sufficient to promote longevity in wild-type worms. CRTCs, therefore, link energetics to longevity by critically fine-tuning histone acetylation and methylation to promote healthy ageing.

MAIN TEXT

Transcriptional regulators have emerged as evolutionarily conserved mechanisms in ageing and longevity[1]. Gene expression can be optimised for health and longevity by manipulating transcription factors and their regulators, making them attractive targets to treat or prevent age-related diseases. Recent studies in the nematode *C. elegans* have highlighted a family of cofactors, cAMP response element-binding protein (CREB)-regulated transcriptional coactivators (CRTCs), as novel modulators of ageing that link energy sensing to transcription[2,3]. CRTCs are multifunctional factors, and despite their importance in multiple human age-related pathologies[4], it is not yet clear how specific roles of CRTCs coordinate organismal longevity.

Mammals possess three CRTC family members. CRTCs lack DNA-binding activity and depend on transcription factor partners for recruitment to DNA to stimulate gene transcription. Thus, CRTCs contain a conserved N-terminal coiled-coil domain, the CREB binding domain, required to bind CREB and other bZIP transcription factors[5]. *C. elegans* possess a single, highly conserved CRTC family member, CRTC-1. We previously demonstrated that when AMP-activated protein kinase (AMPK) is constitutively active, it inactivates CRTC-1 directly by phosphorylation and promotes longevity. In addition, CRTC-1 acts through its canonical transcriptional partner CREB to block this longevity[2,3]. CRTCs are traditionally described as transcriptional coactivators, but recently additional functions of this family of proteins have been revealed, including ER-to-Golgi transport through COPII-mediated vesicle trafficking and splicing and chromatin regulation[6–9]. Here we identify a novel role of CRTC-1 in the epigenetic regulation of longevity. Furthermore, we demonstrate that CRTC-1 promotes longevity through a single defined domain by coupling two different modes of action: transcriptional activation and histone acetylation.

CRTC-1-binding proteins

To identify CRTC-1 protein interactions, we tagged endogenous CRTC-1 via CRISPR Cas-9 editing to generated a *crtc-1::3xFLAG* strain and performed immunoprecipitation (IP) followed by liquid chromatography coupled to mass spectrometry (LC-MS). We found 137 potential CRTC-1 interacting proteins (**Fig. 1a and Supplementary Table 1**), including some known mammalian CRTCs interactors such as SEC-31/SEC31, TAX-6/calcineurin, and to a lesser extent, AAK-2/AMPK alpha 2[2,8,10] (**Fig. 1a**). Using STRING *in silico* analysis[11], we found CRTC-1 binds proteins with WD40 repeats, including WDR-5.1/WDR5 (**Fig. 1b**). WDR-5.1, along with methyltransferase SET-2/SET1 and ASH-2/ASH2L (components of the COMPASS complex), catalyses H3K4me3 and regulates lifespan in *C. elegans*[12,13]. Thus, H3K4me3-deficient animals show an activation of the SBP-1/SREBP1 transcription factor, which in turn drives a transcriptional network to promote fat accumulation with specific enrichment of monounsaturated fatty acids (MUFAs), which are necessary for the lifespan extension[13]. Since CRTCs regulate global energy homeostasis, and we previously showed that CRTC-1 modulates metabolism during AMPK-dependent longevity[2,3], we reasoned that CRTC-1 could play a novel role in COMPASS-dependent longevity due to its interaction with WDR-5.1 (**Fig. 1a and 1b**).

Fig. 1. The co-transcriptional function of CRTC-1 is required for H3K4me3-mediated longevity. a, Volcano plot showing proteins identified during immunoprecipitation analysis of endogenous CRTC-1::3xFLAG tagged strain relative to wild-type non-tagged strain (N2). The dotted line indicates a one-fold change. Four biological replicates. P-values were calculated by two sample t-test. See Supplementary Table 1 for IP statistics. **b**, STRING protein-protein interaction network of CRTC-1-binding proteins identified during immunoprecipitation analysis. The network nodes are proteins. The edges represent the predicted functional associations. Redline (presence of fusion evidence), green line (neighbourhood evidence), blue line (co-occurrence evidence), purple line (experimental evidence), yellow line (text mining evidence), light blue line (database evidence), and blackline (co-expression evidence). **c**, Schematic of endoplasmic reticulum (ER)–Golgi traffic regulated by the SEC31-binding site of CRTC (top). Amino acid sequence localisation of the SEC31-binding site and CRISPR edit (bottom). **d**, Schematic of co-transcriptional regulation by the CREB-binding domain of CRTC (top). Amino acid sequence localisation of the CREB-binding domain site and CRISPR edits (bottom). **e**, *crtc-1*$^{Sec31-/-}$ specifically regulates SREBP1/SBP-1 activation. Intestinal images of GFP::SBP-1 reporter in wild type, *crtc-1*$^{Sec31-/-}$, and *crtc-1*$^{CBD-/-}$ animals. Quantification of GFP::SBP-1 nuclear accumulation; mean ± SEM of n = 123-133 nuclei, pooled from at least three independent experiments, ****p<0.0001 by Mann-Whitney test. **f**, *crtc-1*$^{CBD-/-}$ specifically regulates CRE promoter transcriptional activation. Images of CREp::GFP reporter in neurons of the head in wild type, *crtc-1*$^{Sec31-/-}$, and *crtc-1*$^{CBD-/-}$ animals. Quantification of CREp::GFP expression; mean ± SEM of n = 36-44 heads, pooled from at least three independent experiments, ****p<0.0001 by Mann-Whitney test. **g-h**, Survival curves showing that *set-2(ok952)* mutants live longer than wild type (**g**), and *crtc-1*$^{Sec31-/-}$ does not suppress this longevity phenotype (**h**). **i-j**, Survival curves demonstrating that *set-2(ok952)* mutants live longer than wild type (**i**), and *crtc-1*$^{CBD-/-}$ completely suppresses this longevity phenotype (**j**). See Supplementary Table 2 for lifespan statistics.

CRTC-1 specifically regulates lifespan extension under H3K4me3 deficiency

We generated a novel *C. elegans crtc-1* null mutant strain (*crtc-1*^{null}) to test its role in COMPASS-dependent longevity and other longevity pathways. The absence of CRTC-1 did not affect *crtc-1*^{null} mutant lifespan compared to wild-type animals (**Extended Data Fig. 1a**). As previously reported[12], RNA interference (RNAi) of H3K4me3 modifiers *set-2, ash-2,* and *wdr-5.1* extended lifespan in wild-type worms (**Extended Data Fig. 1b, d, and f**). In *crtc-1*^{null} mutants, we still observed a lifespan extension with *wdr-5.1* RNAi (**Extended Data Fig. 1c**). However, lifespan extension by *ash-2* RNAi was significantly reduced in *crtc-1*^{null} mutants and *crtc-1*^{null} entirely suppressed *set-2* RNAi longevity (**Extended Data Fig. 1e and g**). To further explore the role of CRTC-1 in SET-2-dependent longevity, we crossed the *crtc-1*^{null} strain into the long-lived *set-2(ok952)* strain[12]. We found that the absence of CRTC-1 fully suppressed lifespan extension in *set-2* mutants (**Extended Data Fig. 1h-i**).

We next examined whether CRTC-1 is a global modulator of other longevity pathways. We crossed *crtc-1*^{null} animals into *raga-1(ok386)* (mTORC1 pathway) and *clk-1(qm30)* (electron transport chain) mutants and reduced insulin/insulin-like growth factor-1 signalling via *daf-2 RNAi*[14–17]. In all conditions, we observed a significant increase of lifespan in both wild-type and *crtc-1*^{null} backgrounds (**Extended Data Fig. 1j-o**). Overall, our data show that, instead of being required for all longevity pathways, CRTC-1 activity is specifically required for H3K4me3-dependent longevity. In addition, given our previous finding that activation of CRTC-1 suppresses AMPK mediated longevity[2,3], these data highlight a complex and contextual role for how CRTC-1 activity state can impact healthy ageing.

Targeting CRTC-1 functions in longevity

Since CRTCs have multiple functions related to different cellular processes and age-related diseases[4], we hypothesised that a specific function of CRTC-1 might be required to promote longevity under conditions of H3K4me3 deficiency. Mammalian CRTC2 regulates SREBP1 activation and integrates it to TORC1 activity through COPII vesicle formation and competitive sequestration of Sec31A[8]. In *C. elegans*, SBP-1/SREBP1 activation is required for MUFAs synthesis and longevity in H3K4me3-deficient animals[13]. Supporting these data, we found in our immunoprecipitation experiments that CRTC-1 interacts with the *C. elegans* counterpart of

Fig. 2. CRTC-1^{CBD} mediates specific longevity pathways and acts through AP-1 to extend lifespan in H3K4me3-deficient animals. **a**, GC-MS quantification of fatty acids. Mean ± SEM, *p<0.05, **p<0.01, and ***p<0.001 by unpaired two-way ANOVA. **b-c**, Cluster analysis identified 770 genes that show increased expression in *set-2(ok952)* that is reversed in the double *crtc-1$^{CBD-/-}$::set-2(ok952)* mutant (**a**) and 1704 genes that show decreased expression in *set-2(ok952)* that is reversed in the absence of CRTC-1^{CBD} (**b**). **d**, Over-representation analysis of the Gene Ontology biological process terms for genes comprising each cluster shown in panels (**a-b**). **e-f**, Single site analyses to detect over-represented conserved transcription factor binding sites in up > down (**e**) and down > up (**f**) groups. Analysis was done using oPOSSUM version 3.0. **g-j**, Survival curves showing that the lifespan extension in *set-2(ok952)* (**g**) mutants does not require the transcriptional factors *crh-1(n3315)* (**h**), *crh-2(gk3293)* (**i**), and *atf-6(ok551)* (**j**). **k-n**, AP-1 components, *fos-1* and *jun-1*, do not suppress *set-2(ok952)* longevity separately (**k-m**), but the absence of the entire complex (*jun-1(gk557);fos-1RNAi*) fully suppresses *set-2(ok952)* longevity (**n**). See Supplementary Table 2 for lifespan statistics.

Sec31A, SEC-31 (**Fig. 1a-b**). In addition, we previously showed that a constitutively activated nuclear form of CRTC-1 suppresses AMPK-dependent longevity by promoting CRH-1/CREB-dependent transcription[2,3]. Based on our data and published work, we sought to uncouple these two distinct CRTC-1 functions, 1) ER-to-Golgi transport and 2) transcriptional regulation, and assess their role in H3K4me3-dependent longevity.

We leveraged CRISPR/Cas9 to edit functional domains of endogenous *crtc-1* specifically, mutating the amino acids in CRTC-1 that regulate Sec31A-binding[8] and CREB-binding[18] independently (**Fig. 1c-d**). To define whether these edits, Sec31 and CBD, indeed uncouple the specific functions of CRTC-1, we used *in vivo* readouts for SBP-1 processing and transcription

activation[19]. Using a GFP::SBP-1 reporter, we quantified its nuclear accumulation in intestinal cells as an indication of its activation[13]. We confirmed that the specific *crtc-1$^{Sec31-/-}$* mutant shows increased SBP-1 activation (**Fig. 1e**). To assess CRTC-1 transcriptional effects, we used a CREB transcriptional reporter, a cAMP response element sequence fused with GFP (CREp::GFP)[2]. The CREp::GFP reporter is mainly expressed in head neurons, and its expression is specifically reduced in the *crtc-1$^{CBD-/-}$* mutant (**Fig. 1f**). Critically, these two CRTC-1 functions can be specifically uncoupled since *crtc-1$^{CBD-/-}$* mutants do not increase SBP-1 activation, and *crtc-1$^{Sec31-/-}$* mutants do not reduce CREp::GFP expression (**Fig. 1e-f**). Our data confirm that the roles of CRTC-1 in transcriptional regulation and ER-to-Golgi trafficking are conserved in *C. elegans*, and importantly that we can selectively and specifically uncouple these functions in endogenous CRTC-1.

To identify the specific CRTC-1 function that regulates H3K4me3-dependent longevity, we crossed the *crtc-1$^{CBD-/-}$* and *crtc-1$^{Sec31-/-}$* strains into *set-2* long-lived mutants and analysed their role in lifespan. Strikingly, although H3K4me3-deficient animals promote MUFAs accumulation through SBP-1 activation[13], the Sec31 point mutation did not suppress longevity in these animals (**Fig. 1g-h**). In contrast, the CBD point mutation fully suppressed lifespan extension in *set-2* worms (**Fig. 1i-j**). These data indicate that the co-transcriptional role of CRTC-1 through its CBD domain (hereafter named CRTC-1^{CBD}, referring to the wild-type form) regulates H3K4me3-dependent longevity. Contrary to AMPK longevity, where the CRTC-1-mediated transcriptional activation suppresses longevity[2,3], H3K4me3 deficiency (where overall transcription levels are decreased) requires transcriptional activation driven by CRTC-1^{CBD} to extend lifespan. Therefore, our data suggest that transcriptional activation is necessary to promote longevity in H3K4me3-deficient animals.

CRTC-1^{CBD} regulates specific longevity-related pathways

H3K4me3-deficient animals accumulate MUFAs, and these types of fatty acids are necessary for lifespan extension[13]. Therefore, we sought to define if CRTC-1^{CBD} regulates MUFAs accumulation in *set-2* mutants. To further assess potential changes in specific fatty acid synthesis, we performed gas chromatography coupled to

a up > down group

PVD neurons (289 genes)
Outer labial sensilium (292 genes)
Head mesodermal cell (301 genes)
Intestine (526 genes)
ASE neurons (158 genes)

0.0 0.5 1.0 1.5 2.0
-log$_{10}$ (P value)

b down > up group

Germ line (1650 genes)
Reproductive system (1684 genes)
Gonadal primordium (538 genes)
Embryonic founder cell (82 genes)
Germ line precursor cell (25 genes)

0.0 0.5 1.0 1.5 2.0
-log$_{10}$ (P value)

c AP-1 targets

PVD neurons (233 genes)
Outer labial sensilium (236 genes)
Head mesodermal cell (224 genes)
Cephalic sensillum (44 genes)
ASE neurons (127 genes)

0.0 0.5 1.0 1.5 2.0
-log10 (P value)

d

Muscles | Intestine | Neurons | Gonad

CRTC-1::GFP

Brightfield

Head

e (Survivorship vs Age (Days))

Wild type
set-2
crtc-1$^{CBD-/-}$
crtc-1$^{CBD-/-}$;set-2
crtc-1$^{CBD-/-}$;set-2;
rab-3p::crtc-1$^{CBD+/+}$

*P = <0.0001

f

Wild type
set-2
crtc-1$^{CBD-/-}$
crtc-1$^{CBD-/-}$;set-2
crtc-1$^{CBD-/-}$;set-2;
myo-3p::crtc-1$^{CBD+/+}$

P = <0.0001
*P = 0.6146

g

Wild type
set-2
crtc-1$^{CBD-/-}$
crtc-1$^{CBD-/-}$;set-2
crtc-1$^{CBD-/-}$;set-2;
nep-17p:::::crtc-1$^{CBD+/+}$

P = <0.0001
*P = 0.8685

Fig. 3. CRTC-1^{CBD} regulates longevity cell nonautonomously. a-b, Tissue enrichment analyses for genes comprising up > down (**a**) and down > up (**b**) groups. Analysis was done using WormBase version WS280. **c**, Tissue enrichment analysis of AP-1 target genes from the up > down group using WormBase version WS280. **d**, Confocal images showing that endogenous CRTC-1::GFP is expressed in muscles, intestine, and neurons but not in the germline. Arrows point CRTC-1::GFP expression in nuclei from muscles and intestine. **e-g**, Survival analysis demonstrating that lifespan extension in *set-2(ok952)* mutants is suppressed by *crtc-1$^{CBD-/-}$* and significantly rescued by a single copy of neuronal *crtc-1$^{CBD+/+}$* (**e**), but not when *crtc-1$^{CBD+/+}$* is expressed in muscles (**f**) or intestine (**g**). See Supplementary Table 2 for lifespan statistics.

mass spectrometry (GC-MS) (**Fig. 2a**). We found that double *crtc-1$^{CBD-/-}$;set-2* mutants did not show significantly reduced levels of MUFAs compared to single *set-2* mutants (**Fig. 2a**), suggesting that CRTC-1^{CBD} does not regulate *set-2* longevity via altering MUFAs accumulation.

Since our data indicate that the co-transcriptional role of CRTC-1 is necessary to extend lifespan in H3K4me3-deficient animals (**Fig. 1i-j**), we reasoned that CRTC-1^{CBD}-dependent transcriptional changes in *set-2* mutants might reveal mechanistic insight into how CRTC regulates COMPASS-mediated longevity. Via RNA sequencing, we found that CRTC-1^{CBD}-dependent genes in *set-2* animals cluster in two main groups (**Fig. 2b-c and Supplementary Table 3**). Interestingly, functional analysis within these two clusters revealed that Gene Ontology (GO) terms are clearly different in each group (**Fig. 2d and Extended Data Fig. 2a-b, and Supplementary Table 3**). In the up > down group, we observed an enrichment of processes involved in activation of the MAPK cascade, including immune and stress responses, whereas GO terms related to translation such as ribosome biogenesis, ribonucleoprotein assembly, and RNA processing are represented in the down > up group (**Fig. 2d**). Importantly, both activation of MAPK cascade and reduction of translation are defined signatures of longevity[20–22], and it is to be noted that

CRTC-1^{CBD} specifically regulates expression of genes involved in these processes. Furthermore, in agreement with the fact that CRTC-1^{CBD} does not alter MUFA accumulation (**Fig. 2a**), we did not observe any CRTC-1^{CBD}-dependent transcriptional changes that directly regulate lipid metabolism during our functional analysis in both groups (**Fig. 2d**). Together our data suggest that instead of a generalised dysregulation of transcription by the absence of CRTC-1^{CBD} in H3K4me3-deficient animals, CRTC-1^{CBD}-dependent transcriptional regulation of genes in specific pathways is coupled to longevity.

AP-1 regulates longevity in H3K4me3-deficient animals
The CBD domain of CRTC-1 is critical to associate with bZIP factors[18], including CREB, the heterodimer AP-1, and the activating transcription factor (ATF)6[5,23–25]. Using our transcriptional data (**Fig. 1b-c**), we performed a Single Site Analysis to detect over-represented conserved transcription factor binding sites[26–28]. We observed that CREB1 and AP-1 transcription factors have gene targets in both groups (**Fig. 2e-f and Supplementary Table 4**). However, only AP-1 was above the recommended threshold[26] in the up > down group (mean + 1* standard deviation, **Fig. 2a**). This means that those genes that were upregulated in H3K4me3-deficient animals and then downregulated by the absence of CRTC-1^{CBD} (**Fig. 2b**) are significant targets of AP-1 (**Fig. 2e and Supplementary Table 4**).

To test the causal role of CRTC-1 transcription factor partners in H3K4me3-dependent longevity, we used a combination of mutants and RNAi for their worm counterparts: CRH-1/CREB1, CRH-2/CREB3L1, ATF-6/ATF6, and FOS-1:JUN-1/AP-1[5,23–25]. Although we have previously shown that CRTC-1 governs AMPK-mediated longevity and metabolism via the CREB transcription factor[3], we found that CRH-1 and CRH-2 are dispensable for longevity in *set-2* mutants (**Fig. 2g-i**). The *atf-6(ok551)* mutant by itself is long-lived[29]. *set-2* deletion significantly increases lifespan in *atf-6* mutants, although to a lesser extent than in WT (**Fig. 2j**). Separately, inactivation of AP-1 components, JUN-1 and FOS-1 alone did not suppress *set-2* longevity (**Fig. 2k-m**). Strikingly, however, the absence of the entire AP-1 complex (*jun-1(gk557);fos-1RNAi*) fully suppresses *set-2* longevity (**Fig. 2n**). These data indicate that the complete heterodimer AP-1 transcription factor is necessary to extend longevity and point to the CRTC-1^{CBD}:AP-1 complex as the critical transcriptional core that promotes longevity in H3K4me3-deficient animals.

CRTC-1^{CBD} regulates lifespan cell nonautonomously
H3K4me3 modifiers function at least in part in the germline to promote longevity[12,13]. We demonstrate here that CRTC-1^{CBD} is required to extend lifespan of *C. elegans* with an H3K4me3 deficiency (**Fig. 1 and Extended Data Fig. 1s**), and CRTC-1 is known to mediate longevity from neurons during AMPK activation[3]. These data suggest a potential inter-tissue interaction in the H3K4me3 > CRTC-1^{CBD} pathway to coordinate longevity. We performed tissue enrichment analysis using the genes present

Fig. 4. CRTC-1^{CBD} maintains histone acetylation levels and these are necessary for lifespan extension. a, H3K9Ac, H3K18Ac, and H3K27Ac marks are reduced in a CRTC-1^{CBD}-dependent manner under H3K4me3 deficiency induced by a loss of function of *set-2*. Western blots of wild-type, *set-2(ok952)*, *crtc-1^{CBD-/-}*, and *crtc-1^{CBD-/-};set-2(ok952)* worms. H3 levels were used as loading controls (representative immunoblot of three independent blots). See Extended Data Fig. 3 for Western blot quantifications. **b**, Fluorescence images of the distal region of *C. elegans* gonad from wild-type, *set-2(ok952)*, *crtc-1^{CBD-/-}*, *crtc-1^{CBD-/-};set-2(ok952)* and *crtc-1^{CBD-/-};set-2(ok952);rab-3p::crtc-1^{CBD+/+}* worms co-immunostained for H3K4me3 (gray), H3K9Ac (green), and DAPI (magenta). **c**, Quantification of fluorescence intensity detected for H3K4me3 (**a**). *set-2* and *crtc-1^{CBD-/-};set-2* worms showed a significant reduction of the H3K4me3 mark, which is not rescued in the *crtc-1^{CBD-/-};set-2(ok952);rab-3p::crtc-1^{CBD+/+}* strain. Mean ± SEM of n = 10-14 gonads, pooled from three independent experiments, *p<0.05, **p<0.01, and ***p<0.001 by unpaired t test. ns = not significant. **d**, Quantification of fluorescence intensity detected for H3K9Ac (**a**). *crtc-1^{CBD-/-}* and *crtc-1^{CBD-/-};set-2(ok952)* worms showed a significant reduction of the H3K9Ac mark, which is rescued in the *crtc-1^{CBD-/-};set-2(ok952);rab-3p::crtc-1^{CBD+/+}* strain. Mean ± SEM of n = 10-14 gonads, pooled from three independent experiments, ***p<0.001 and ****p<0.0001 by unpaired t test. ns = not significant. **e-f**, Survival curves demonstrating that lifespan extension in *set-2(ok952)* mutants is suppressed by *cbp-1* (**e**) and *pcaf-1* (**f**) RNAi. (**g**) Survival curves demonstrating that overexpression (OE) of PCAF-1 increases wild-type lifespan in three different lines. See Supplementary Table 2 for lifespan statistics.

in the two main groups obtained from our RNA-Seq data (**Fig. 2b-c**). In the up > down group, we found a significant enrichment of genes expressed in neurons (**Fig. 3a**). In contrast, those genes that belong to the down > up group are enriched in the gonad (**Fig. 3b**). Additionally, we performed a tissue enrichment analysis specifically for AP-1 targets from the up > down group (**Fig. 2e**). Similar to CRTC-1, we found a significant enrichment of AP-1 gene targets expressed in the nervous system (**Fig. 3c**).

To link the tissue enrichment analysis to CRTC-1 tissue localisation, we characterised the endogenous expression of CRTC-1. CRTC-1::GFP is particularly high in neurons, while also present in muscles and intestine but not in the germline (**Fig. 3d**). Furthermore, our data demonstrate that the co-transcriptional role of CRTC-1 regulates H3K4me3-dependent longevity (**Fig. 1i-j**), which correlates with transcriptional gene activation from the neuron-enriched up > down group (**Fig. 2b and 3a**). Together,

these findings lead us to hypothesise that CRTC-1 modulates H3K4me3-dependent longevity cell nonautonomously, potentially from the nervous system.

To directly examine the role of neuronal CRTC-1^{CBD} activity on H3K4me3-dependent longevity, we rescued CRTC-1^{CBD} activity in double *crtc-1^{CBD};set-2* mutants. Using our SKI LODGE system[30], we inserted a single copy of *crtc-1* cDNA wild-type sequence into an intergenic cassette containing the *rab-3* promoter and *rab-3 3' UTR* sequences to drive gene expression specifically in neurons. We found that expressing CRTC-1 from the *rab-3* promoter fully rescued the lifespan extension in *crtc-1^{CBD-/-};set-2* mutants (**Fig. 3e**). In contrast, expressing CRTC-1 directly in muscles or intestine by single copy transgene using the muscle-specific *myo-3* or intestine-specific *nep-17* promoters, respectively, did not rescue the lifespan extension in the double mutants (**Fig. 3f-g**). Taken together, these results indicate that CRTC-1^{CBD} activity specifically in neurons modulates longevity in animals with H3K4me3 deficiency.

Overall, our data indicate that rather than modulating global transcriptional changes, CRTC-1^{CBD}:AP-1 activity controls specific pathways necessary for H3K4me3 longevity. These specific changes in transcriptional regulation are associated with active transcription since we demonstrate that the coactivating transcriptional role of CRTC-1 is required for longevity (**Fig. 1, 2, and 3**). Furthermore, this CRTC-1^{CBD}-dependent active transcription occurs when global transcriptional levels are reduced in H3K4me3-deficient animals to induce transcription and hence longevity. Since mammalian CRTCs are involved in chromatin regulation to activate transcription[6], we aimed to analyse if chromatin modifications could link activation of gene expression to CRTC-1^{CBD} function.

Histone acetylation promotes longevity

Mammalian CRTC2 associates with the KAT2B acetyltransferase, which in turn increases the H3K9Ac at promoters of gluconeogenic genes, upregulating transcription[6]. CRTC2 also interacts with p300 and CBP acetyltransferases[7]. The *C. elegans* ortholog of the mammalian acetyltransferases p300 and CBP, CBP-1, was recently shown to regulate H3K18Ac and H3K27Ac levels[31]. Therefore, we sought to explore whether CRTC-1^{CBD} mediates H3K9Ac, H3K18Ac, and H3K27Ac marks in *set-2* mutants. We found that in *set-2* mutants, all three

acetylation marks were increased: H3K27Ac (+5%), H3K9Ac (+15%), and H3K18Ac (+25%). The same histone marks were then reduced in single *crtc-1*[CBD-/-] mutants: H3K9Ac (-16%), H3K18Ac (-14%), and H3K27Ac (-5%). Strikingly, however, we observed a dramatic reduction of acetylation in the double *crtc-1*[CBD-/-]*;set-2* mutants: H3K9Ac (-36%), H3K18Ac (-34%), and H3K27Ac (-45%) (**Fig. 4a and Extended Data Fig. 3b-c**). In contrast, *crtc-1*[CBD-/-] did not change global H3K4me3 levels (**Fig. 4a and Extended Data Fig. 3a**). These results indicate that the CRTC-1[CBD] regulates H3K9Ac, H3K18Ac, and H3K27Ac marks, particularly in H3K4me3-deficient animals.

Given that H3K4me3 modifiers promote longevity from germline[12,13] and CRTC-1[CBD] modulates H3K4me3 longevity from neurons (**Fig. 3**), we asked whether CRTC-1[CBD] regulates histone acetylation cell nonautonomously. Similar to our Western blot analyses (**Fig. 4a**), *crtc-1*[CBD-/-] did not change H3K4me3 levels in the gonad (**Fig. 4b-c**), but H3K9Ac levels were significantly reduced in both *crtc-1*[CBD-/-] and *crtc-1*[CBD-/-]*;set-2* animals (**Fig. 4b and d**). Interestingly, restoring *crtc-1*[CBD+/+] expression only in neurons fully rescued H3K9Ac germline levels in *crtc-1*[CBD-/-]*;set-2* mutants. (**Fig. 4b and d**). In line with our previous data, the neuronal rescue of CRTC-1[CBD] did not affect reduced levels of H3K4me3 in the gonad (**Fig. 4b-c**). Therefore, similarly to lifespan, CRTC-1[CBD] regulates H3K9Ac cell nonautonomously from neurons. Overall, our data indicate that CRTC-1[CBD] regulates histone acetylation, notably when H3K4me3 levels are reduced, suggesting that histone acetylation contributes to lifespan extension in H3K4me3-deficient animals.

To test the hypothesis that histone acetylation is required for longevity during H3K4me3 deficiency, we knocked down the *C. elegans* orthologs of the histone acetyltransferases associated with CRTCs in mammals that also regulate the acetylation marks mentioned above, PCAF-1/KAT2B and CBP-1/p300/CBP[6,7]. Supporting our previous findings, we found that both *cbp-1* and *pcaf-1* knockdowns entirely suppressed longevity in *set-2* mutants (**Fig. 4e-f**). Remarkably, *cbp-1* knockdown had a dramatic effect on *set-2* mutants. The lifespan in *set-2;cbp-1 RNAi* was shorter than *wild type;cbp-1 RNAi* lifespan (**Fig. 4e**), indicating that CBP-1 plays a more essential role in H3K4me3-deficient animals than wild-type animals, wherein H3K4me3 levels are normal.

These data suggest histone acetylation is required to extend lifespan in the context of H3K4me3 deficiency. Finally, we reasoned that if histone acetylation positively affects ageing, the lifespan would be increased by overexpressing histone acetyltransferases associated with CRTCs. Indeed, overexpression of PCAF-1 was sufficient to extend lifespan in the wild-type background by 40% (**Fig. 4g**). Collectively, these data suggest that CRTC-1[CBD] mediates longevity during H3K4me3 deficiency via remodelling global histone acetylation levels, and remarkably, that this acetylation is sufficient to promote longevity in wild-type worms, identifying histone acetylation as a novel marker for longevity.

DISCUSSION

CRTCs have several functions and have been implicated in many age-related diseases such as cancer, neurodegeneration, diabetes, and other metabolic disorders[4,32]. Thus, CRTCs are key integrators that regulate cellular signalling, and their variety of functions is context dependent. Linking CRTC functions to specific physiological processes will, therefore, offer novel opportunities for clinical therapies. We previously demonstrated that an overexpressed active form of CRTC-1 in neurons suppresses lifespan extension when AMPK is constitutively activated in the entire worm[3]. Surprisingly, in this study, we found a positive role for CRTC-1 in regulating longevity due to H3K4me3 deficiency. Several positive and negative roles have been described for CRTCs in cancer, diabetes, and autoimmune disease[4,32]. Now, we discover a dual role of CRTC-1 in ageing, supporting the idea that CRTC's roles in maintaining homeostasis are context-dependent.

We propose a model in which lifespan extension induced by H3K4me3 methyltransferase deficiency requires a specific function (co-transcriptional regulation) of CRTC-1 driven by its CBD domain, which in turn triggers a neuron-to-periphery signal via the AP-1 transcription factor, leading to activation of gene expression and maintenance of histone acetylation levels. Surprisingly, although H3K4me3 deficiency couples lipid metabolism to longevity through the SBP-1/SREBP1 transcription factor in worms[13] and mammalian CRTC2 regulates lipid metabolism by inducing COPII trafficking-dependent activation of SREBP1[8], we demonstrate that this CRTC-1 function is not required for H3K4me3 longevity. Although we have demonstrated that a deficiency in H3K4me3 promotes longevity[12,13,33], recent work suggests that this deficiency shortens lifespan[34]. However, the food source differs from our conditions, and in addition, RNAi in *set-2* extends lifespan in wild-type worms[12,13,33,35].

It is as yet unclear how CRTC-1[CBD] regulates histone acetylation. One of the primary sources of an acetyl group for histones is the metabolite acetyl-CoA, and it has been shown that cellular metabolism is intertwined with chromatin dynamics since many metabolites constitute histone and DNA modifications[36]. Since CRTCs have been extensively associated with metabolic pathways[4], it is possible that CRTC-1[CBD] regulates histone acetylation by controlling metabolism and metabolic intermediaries. Rather than just a global increase in histone acetylation or a global reduction in histone trimethylation, the essential factor is which genes are targeted by these marks in specific contexts and tissues in order to promote longevity. Therefore, it is critical to identify components that integrate histone marks with gene expression, such as CRTC-1. Finally, if promoting histone acetylation induces longevity, as our data indicate, selectively targeting histone acetylation modifiers in specific tissues may be sufficient to generate an organismal response that promotes healthier ageing.

ACKNOWLEDGMENTS

We thank the Caenorhabditis Genetics Center for providing worm strains. We thank Zon Weng Lai from the Harvard Chan Advanced Multiomics Platform (ChAMP) for the mass spectrometry analysis for immunoprecipitation. We thank Jihe Liu from the Harvard Chan Bioinformatics Core for the RNA-seq analysis. We are also grateful to members of the Mair laboratory for helpful discussions on the project and the manuscript.

FUNDING

WBM is funded by the National Institutes of Health, NIA/NIH R01AG051954, NIA/NIH R01AG059595, NIA/NIH R01AG067106, NIA/NIH R01AG054201, and NIA/NIH R01AG044346. This work was also supported by the American Federation for Aging Research/
10 Glenn Foundation for Medical Research Breakthroughs in Gerontology Award. Work in MPC's lab is funded by NIGMS/NIH R01GM072551. CGSG is funded by NIA/NIH K99AG065508.

AUTHOR CONTRIBUTIONS

CGSG and WBM designed the study. CGSG performed most experiments and analysed results. LILL performed immunostainings and image analysis in MPC lab. KP performed gas chromatography/mass spectrometry analysis of fatty acid profiles in AB lab. CH, LP and AS performed lifespan repeats. CGSG and WBM wrote the manuscript, implemented comments and edits from all authors.

DECLARATION OF INTERESTS

The authors declare no competing interests.

METHODS

Materials & Correspondence

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, William Mair (wmair@hsph.harvard.edu). All unique and stable reagents generated in this study are available from public repositories (e.g., Caenorhabditis Genetics Center for *C. elegans* strains) or from the Lead Contact with a completed Materials Transfer Agreement. Source transcriptomic data will be available, and other original data will be available upon request after publication.

C. elegans strains and husbandry

N2 wild-wild type, MQ130 [*clk-1(qm30)*], VC222 [*raga-1(ok952)*], RB772 [*atf-6(ok551)*], VC3149 [*crh-2(gk3293)*], and VC1200 [*jun-1(gk557)*] *C. elegans* strains were obtained from the Caenorhabditis Genetic Center, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). The *crh-1(n3315)* mutant was provided by Mark Alkema. Worms were maintained on standard nematode growth media (NGM) seeded with *E. coli* OP50-1 and maintained at 20 ºC.

Microbe strains

OP50-1 bacteria were cultured overnight in LB at 37 ºC, after which 100 ml of liquid culture was seeded on plates to grow for two days at room temperature. Unless otherwise noted, worms were grown at 20 ºC on the strain *E. coli* HT115 (empty vector, EV) for all experiments. RNAi experiments employed HT115 bacteria from the Ahringer library (Source Bioscience) expressing dsRNA against the gene noted or empty vector control. HT115 bacteria were cultured overnight in LB containing 100 mg ml^{-1} carbenicillin and 12.5 ml ml^{-1} tetracycline at 37 C, after which 100 mL of LB was seeded on NGM plates containing 100 mg ml^{-1} carbenicillin to grow for two days at room temperature. dsRNA expression was induced by adding 100 ml IPTG (100 mM) at least 2 hours before worms were introduced to the plates. RNAi experiments for *ash-2*, *set-2*, and *wdr-5.1* were performed as previously described[13].

Microinjection and CRISPR/Cas9-triggered homologous recombination

All CRISPR edits and insertions required to generate the strains were performed using the previously described CRISPR protocol[30,37]. Briefly, homology repair templates were amplified by PCR, using primers that introduced a minimum stretch of 35 bp homology at both ends. Single-stranded oligo donors (ssODN) were also used as repair templates. CRISPR injection mix reagents were added in the following order: 0.375 µl Hepes pH 7.4 (200 mM), 0.25 µl KCl (1 M), 2.5 µl tracrRNA (4 µg/µl), 0.6 µl *dpy-10* crRNA (2.6 µg/µl), 0.25 µl *dpy-10* ssODN (500 ng/µl), and PCR or ssODN repair template(s) up to 500 ng/µl final in the mix. Water was added to reach a final volume of 8 µl. 2 µl purified Cas9 (12 µg/µl) added at the end, mixed by pipetting, spun for 2 min at 13000 rpm and incubated at 37 ºC for 10 min. Mixes were microinjected into the germline of day 1 adult hermaphrodite worms using standard methods[38].

Single-Copy Transgene Construction

crtc-1 single-copy neuronal rescue strain was generated by CRISPR according to[30]. Specifically, a homology repair template (HR) containing *crtc-1* cDNA sequence was amplified from plasmid pIM41. A CRISPR mix containing the *crtc-1* HR template was prepared according to[30] and injected into the WBM1144 strain. The resulting CRISPR-edited allele wbmIs92 was outcrossed 6 times to N2 before use for experiments.

Survival Analyses

Lifespan experiments were performed as described previously[3,39]. Unless otherwise noted, all experiments were performed on standard NGM containing 100 mg ml^{-1} carbenicillin at 20 ºC and on HT115 (empty vector, EV). RNAi experiments were performed from hatching in all cases except for *cbp-1 RNAi*, which started on day 1 of adulthood, on standard NGM containing carbenicillin (100 mg ml^{-1}). Expression of dsRNA was induced by

pipetting 100 ml IPTG solution (100 mM, containing 100 mg ml^{-1} carbenicillin and 12.5 ml ml^{-1} tetracycline) onto HT115 lawns prior to place worms. Worms were synchronised by bleaching using gravid adults. After bleaching, embryos were placed on plates. When the progeny reached adulthood (~72 h), 100 worms were transferred to fresh plates with 20 worms per plate. Worms were transferred to fresh plates every other day until reproduction had ceased (day 9-12). Survival was scored every 1–2 days, and a worm was deemed dead when unresponsive to 3 taps on the head and tail. Worms were censored due to contamination on the plate, leaving the NGM, eggs hatching inside the adult, or loss of vulval integrity during reproduction. Lifespan analysis was performed using GraphPad Prism.

RNA Sequencing

More than 2000 day-one adults were used for each sample. Four biological replicate samples were collected for each genotype. Worms were collected in M9 buffer and washed three times in M9 buffer to remove bacteria. Liquids were removed after last centrifugation, and QIAzol lysis reagent (Qiagen, 79306) was added to each sample before snap-freezing in liquid nitrogen. All samples were stored in a -80℃ freezer until RNA extraction. To break the worm cuticle and improve RNA yield, all samples underwent five freeze-thaw cycles. In each cycle, samples were thawed at 37 ºC and then snap-frozen in liquid nitrogen. RNA extraction was performed using QIAGEN RNeasy Mini Kit (QIAGEN, 74104) following manufacturer's instructions. RNA quality was confirmed using TapeStation system (Agilent Technologies). mRNA libraries were prepared using KAPA mRNA HyperPrep, poly-A selection (Roche) following manufacturer's instructions. Library quality was checked using TapeStation system (Agilent Technologies). Libraries were then pooled and sequenced with Illumina NovaSeq SP Single Lane using 50 read length, pair-end settings.

Gene Expression and Functional Analysis

All samples were processed using an RNA-seq pipeline implemented in the bcbio-nextgen project (https://bcbio-nextgen.readthedocs.org/en/latest/). Raw reads were examined for quality issues using FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) to ensure library generation and sequencing data were suitable for further analysis. Reads were aligned to the Ensembl94 build of the *C. elegans* genome using STAR[40]. Quality of alignments was assessed by checking for evenness of coverage, rRNA content, genomic context of alignments, complexity, and other quality checks. Using Salmon[41], expression quantification was performed to identify transcript-level abundance estimates and then collapsed down to the gene level using the R Bioconductor package tximport[42]. Principal components analysis (PCA) and hierarchical clustering methods validated the clustering of samples from the same sample group. Differential expression was performed at the gene level using the R Bioconductor package

DESeq2[43]. Differentially expressed genes were identified using the Wald test, and significant genes were obtained using an FDR threshold of 0.05. Significant genes were separated into clusters based on similar expression profiles across the defined sample groups. Gene lists for each cluster were used as input to the R Bioconductor package cluster Profiler[44] to perform an over-representation analysis of Gene Ontology (GO) biological process terms.

Microscopy of mounted worms and image analysis

For imaging, one-day-old adult animals were mounted on 2% agarose pads and anaesthetised with 20 mM tetramisole in M9. For visualisation of intestinal nuclear SBP-1 expression, a transgenic line expressing a translational fusion between GFP and SBP-1 was used. For visualisation of CRE reporter expression, a transgenic strain expressing a transcriptional reporter was used, CREp::GFP. In both cases, images were acquired using a Zeiss Axio Imager.M2 microscope equipped with an ApoTome.2 system and an AxioCam MRc camera using identical exposure settings for all experiments. The mean fluorescence intensity in arbitrary units (a.u.) of each intestinal nucleus, for the GFP::SBP-1 reporter, was quantified using the oval brush tool in Fiji ImageJ. The mean fluorescence intensity in arbitrary units of each head, for the CREp::GFP reporter, was quantified using the polygon selection tool in Fiji ImageJ. Then, mean intensity values in arbitrary units were graphed using Graphpad Prism, and statistical significance was determined using a two-tailed Mann-Whitney test. For imaging of endogenous CRTC-1::GFP, images were taken in the Sabri Ulker imaging lab using a Yokogawa CSU-X1 spinning confocal disk system (Andor Technology, South Windsor, CT, USA) combined with a Nikon Ti-E inverted microscope (Nikon Instruments, Melville, NY, USA). Images were taken using a 40x objective lens, Zyla cMOS (Zyla 4.2 Plus USB3) camera, and 488 nm Laser for GFP.

Immunostainings and image analysis

Whole-mount preparations of dissected gonads and immunostainings were performed as in[45]. Briefly, gonads from 24 h post-L4 hermaphrodites were dissected and fixed with 1% formaldehyde for 5 minutes, freeze-cracked, and post-fixed in ice-cold 100% methanol 1 minute, followed by blocking with 1% BSA for 1 hour. The following primary antibodies were used at the indicated dilutions: mouse α-H3K4me3 (1:500, MAB Institute Inc Wako 305-348191) and rabbit α-H3K9Ac (1:1000, Abcam ab10812). The secondary antibodies from Jackson ImmunoResearch Laboratories (West Grove, PA) were used at the following dilutions: α-mouse Cy-3 (1:200) and α-rabbit Alexa 488 (1:500). DAPI was used to counterstain DNA. Vectashield from Vector Laboratories (Burlingame, CA) was used as a mounting media and anti-fading agent. Imaging was performed using an IX-70 microscope (Olympus) with a cooled CCD camera (model CH350; Roper Scientific) controlled by the DeltaVision system (Applied Precision). Images were collected using a 40x objective,

and Z-stacks were set at 0.2 μm thickness intervals using identical exposure settings for each antibody. Image deconvolution was done using the SoftWoRX 3.3.6 program (Applied Precision) and processed with Fiji ImageJ. Gonads were imaged by dividing the gonad into three sections from distal to loop regions. To quantify H3K4me3 and H3K9Ac expressions, intensity in arbitrary units of each section was quantified using the polygon selection tool in Fiji. Then, all values from each section of a single gonad were pulled together. Mean intensity values in arbitrary units were graphed using Graphpad Prism, and statistical significance was determined by unpaired t-test.

Co-immunoprecipitation

Pulled-down proteins were obtained from CRTC-1::3xFLAG and N2 one-day-old animals. Animals were grown at 20 °C, collected in M9 medium, last two washes were done with lysis buffer with glycerol (100 mM HEPES pH 7.5, 2 mM $MgCl_2$, 300 mM, 5 mM 2-Mercaptoethanol, 0.05% NP-40, and 10% glycerol) and protease (Sigma-Aldrich, 8340) and phosphatase (Roche, 04906837001) inhibitors. Liquids were removed after the last centrifugation, and then samples were frozen in liquid nitrogen. Liquid nitrogen-frozen pellets equivalent to 1 g of animals were homogenised in 3 ml of lysis buffer (Sigma-Aldrich, L3412; 50 mM Tris HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA, and 1% TRITON X-100) with protease and phosphatase inhibitors. Worms were lysed via sonication (Qsonica, Q700). Protein concentration was measured using Pierce BCA protein assay kit (Thermo Fisher Scientific, PI23227) following manufacturer's instructions. Anti-FLAG M2 magnetic beads (Sigma-Aldrich, M8823) were used to immunoprecipitate CRTC-1::3xFLAG. Beads were prepared by washing 3 times with 1x wash buffer (Sigma-Aldrich, W0390; 0.5 M Tris HCl, pH 7.4, and 1.5 M NaCl). Each supernatant from sonication was incubated with pre-washed beads overnight with end-over-end rotation at 4 °C. After incubation, beads were washed 3 times 5 minutes with lysis buffer (Sigma-Aldrich, L3412) on a rotator and protease and phosphatase inhibitors. Next, beads were rewashed 3 times 5 minutes with 1x wash buffer (Sigma-Aldrich, W0390) on a rotator and with protease and phosphatase inhibitors. Then, samples were ready to process for mass spectrometry.

Mass spectrometry for immunoprecipitation

Pulled-down proteins from CRTC-1::3xFLAG and N2 immunoprecipitations were brought to pH 7.5 with 200 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid). Proteins were reduced using 5 mM dithiothreitol (Sigma-Aldrich) at 37 °C for 1 h, followed by alkylation of cysteine residues using 15 mM iodoacetamide (Sigma-Aldrich) in the dark at room temperature for 1 h. Excessive iodoacetamide was quenched using 10 mM dithiotheritol. Protein mixtures were diluted in 1:6 ratio (v/v) using ultrapure water prior to digestion using sequencing grade trypsin (Promega) at 37 °C for 16 h. Subsequently, digested peptides were desalted using self-packed C18 STAGE tips (3 M EmporeTM)[46] for LC-MS/MS analysis. Desalted peptides were resuspended in 0.1% (v/v) formic acid and loaded onto HPLC-MS/MS system for analysis on an Orbitrap Q-Exactive Exploris 480 (Thermo Fisher Scientific) mass spectrometer coupled to an Easy nanoLC 1000 (Thermo Fisher Scientific) with a flow rate of 250 nl/min. The stationary phase buffer was 0.5% formic acid, and mobile phase buffer was 0.5% (v/v) formic acid in acetonitrile. Chromatography for peptide separation was performed using increasing organic proportion of acetonitrile (5-40% (v/v)) over a 120 min gradient) on a self-packed analytical column using PicoTipTM emitter (New Objective, Woburn, MA) using Reprosil Gold 120 C-18, 1.9 μm particle size resin (Dr Maisch, Ammerbuch-Entringen, Germany). The mass spectrometry analyser operated in data-dependent acquisition mode with a top ten method at a 300-2000 Da mass range.

Mass spectrometry data were processed by MaxQuant software version 1.5.2.8 [47] using the following setting: oxidised methionine residues and protein N-terminal acetylation as variable modification, cysteine carbamidomethylation as fixed modification, first search peptide tolerance 20 ppm, main search peptide tolerance 4.5 ppm. Protease specificity was set to trypsin with up to 2 missed cleavages were allowed. Only peptides longer than five amino acids were analysed, and the minimal ratio count to quantify a protein is 2 (proteome only). The false discovery rate (FDR) was set to 1% for peptide and protein identifications. Database searches were performed using the Andromeda search engine integrated into the MaxQuant environment [48] against the UniProt *Caenorhabditis elegans* database containing 27,390 entries (November 2020). "Matching between runs" algorithm with a time window of 0.7 min was employed to transfer identifications between samples processed using the same nanospray conditions. Protein tables were filtered to eliminate identifications from the reverse database and common contaminants.

Gas chromatography/mass spectrometry analysis of fatty acid profiles

Quantification of long-chain fatty acids was performed as described[13]. Briefly, for each condition, approximately 500 age-synchronised adults on day 1 were collected in M9 buffer and washed 3 times to remove residual bacteria in the worm pellets. Worm pellets were lyzed by sonication, and the protein concentration of the lysate was determined using the Pierce BCA Protein Assay Kit (Thermo- Scientific). The fatty acid C13:0 (NuChek Prep, dissolved in methanol) was added to each sample to serve as the internal reference control for variations introduced during derivatisation and extraction steps. Fatty acids were derivatised into their respective fatty acid methyl ester (FAME) by incubation in 2% H_2SO_4 (Sigma Aldrich) in methanol (Fisher) at 55°C overnight. The reaction was stopped by the addition of 1.5 ml water (Fisher, MS grade). FAMEs were extracted in a 300 μl hexane (Sigma Aldrich) by vigorous vortexing and centrifuging at 1000 rpm for 1 minute. The hexane layer

containing the FAMEs was transferred into an amber GC vial (Agilent). FAME analysis was performed using an Agilent 7890A gas chromatograph equipped with an HP-5MS column. Each FAME peak was identified based on its retention time and unique fragmentation ions and quantified using a serial dilution standard curve. FAME abundance measured by GC-MS was normalised to the internal C13:0 reference control of each sample. For each sample, FAMEs concentration (μg/ml) was normalised to protein concentration (mg/ml) as microgram of fatty acid detected per milligram of protein (μg/mg). The fatty acid concentration (μg/mg) for each mutant was normalised to the fatty acid concentration (μg/mg) of N2 control. The final ratio is expressed as relative fatty acid levels in the graph. One experiment was carried out with three replicates. Relative fatty acid abundances were plotted in Prism 8. Statistically, significant differences between samples were assessed using the unpaired, non-parametric Mann Whitney test with Benjamini and Hochberg test for multiple hypothesis correction.

Western blots

More than 2000 day-one adults were used for each sample. Worms were collected in M9 buffer and washed three times in M9. Liquids were removed after centrifugation, and samples were frozen in liquid nitrogen. For worm lysis, RIPA buffer containing protease inhibitors (Sigma-Aldrich, 8340) and phosphatase inhibitors (Roche, 04906845001) was added to each sample at the same volume as the worm pellet. Worms were lysed via sonication (Qsonica, Q700). Protein concentration was measured using Pierce BCA protein assay kit (Thermo Fisher Scientific, PI23227) following manufacturer's instructions. 4x Leammli sample buffer (Bio-Rad, 1610747) was added to denature the proteins, and samples were heated to 95 °C for 5 min. Samples containing 30-40 mg protein were loaded to 10-20% Tris-Glycine gels (Thermo Fisher Scientific XP10205BOX) for SDS-PAGE. Proteins were transferred to PVDF membranes (Thermo Fisher Scientific, LC2005) and blocked with 5% non-fat milk powder in TBST. The following primary antibodies were used at the indicated dilutions: rabbit α-H3K4me3 (1:1000, Abcam ab8580), rabbit α-H3K9Ac (1:1000, Abcam ab10812), rabbit α-H3K18Ac (1:1000, Sigma-Aldrich 07-354), rabbit α-H3K27Ac (1:1000, Abcam ab4729), and rabbit α-H3 (1:1000, Cell Signaling 9715). The secondary antibody was α-rabbit IgG HRP-linked (1:1000, genes were identified using the Wald test. For all experiments, p values < 0.05 were considered significant.

Cell Signaling 7074). Antibody signals were developed using SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo Fisher Scientific, 34577), and bands were quantified with Fiji ImageJ.

Single site analyses

Transcription factor binding sites (TFBS) analysis was done using oPOSSUM program, version 3.0 [28]. The parameters used in the oPOSSUM were as follows: conservation cut-off = 0.4, matrix score threshold = 85%, analyzed sequences = 1 kb upstream of transcription start sites, Z-score cut-off = 10, Fisher score cut-off = 7.

In silico analysis for protein-protein interaction networks

Immunoprecipitation data was used to identify protein-protein interactions for CRTC-1. The analysis was done using STRING, version 11.0 (http://www.string-db.org). The parameters used in the STRING were as follows: network type = full network, meaning of network edges = evidence, active interaction sources = all options (textmining, experiments, databases, co-expression, neighbourhood, gene fusion, and co-occurrence), minimum required interaction score = 0.400, maximum number of interactions to show = no more than 5 interactions (1st shell).

Tissue-enrichment analysis

Tissue-enrichment analysis was performed using the Enrichment Analysis tool [49] from WormBase version WS280. q value threshold = 0.1 was used as parameter.

Quantification and statistical analysis

Fiji ImageJ was used for image quantifications. GraphPad Prism 9 was used for statistical analyses. Statistical details of experiments can be found in the Figures legends. The Log-rank (Mantel-Cox) method was used to compare survival curves. The Mann-Whitney test was used to analyse GFP::SBP-1 and CREp::GFP expressions. Two sample t-test was used to compare IP samples in volcano plot using Perseus software 1.5.1.6. Mann Whitney test with Benjamini and Hochberg test for multiple hypothesis correction was used to compare fatty acid levels. The unpaired t-test was used to compare the expression of antibody signals in immunofluorescence samples. For RNA-Seq analysis, differentially expressed

REFERENCES

1. Corrales, G. M. & Alic, N. Evolutionary Conservation of Transcription Factors Affecting Longevity. *Trends Genet* 36, 373–382 (2020).
2. Mair, W. *et al.* Lifespan extension induced by AMPK and calcineurin is mediated by CRTC-1 and CREB. *Nature* 470, 404–408 (2011).
3. Burkewitz, K. *et al.* Neuronal CRTC-1 Governs Systemic Mitochondrial Metabolism and Lifespan via a Catecholamine Signal. *Cell* 160, 842–855 (2015).
4. Escoubas, C. C., Silva-García, C. G. & Mair, W. B. Deregulation of CRTCs in Aging and Age-Related Disease Risk. *Trends in Genetics* 33, 303–321 (2017).
5. Conkright, M. D. *et al.* TORCs: transducers of regulated CREB activity. *Molecular cell* 12, 413–23 (2003).
6. Ravnskjaer, K. *et al.* Glucagon regulates gluconeogenesis through KAT2B- and WDR5-mediated epigenetic effects. *The Journal of clinical investigation* 123, 4318–28 (2013).
7. Liu, Y. *et al.* A fasting inducible switch modulates gluconeogenesis via activator/coactivator exchange. *Nature* 456, 269–273 (2008).
8. Han, J. *et al.* The CREB coactivator CRTC2 controls hepatic lipid metabolism by regulating SREBP1. *Nature* 524, 243–246 (2015).
9. Amelio, A. L., Caputi, M. & Conkright, M. D. Bipartite functions of the CREB co-activators selectively direct alternative splicing or transcriptional activation. *The EMBO Journal* 28, 2733–2747 (2009).
10. Screaton, R. A. *et al.* The CREB coactivator TORC2 functions as a calcium- and cAMP-sensitive coincidence detector. *Cell* 119, 61–74 (2004).
11. Szklarczyk, D. *et al.* STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* 47, gky1131 (2018).
12. Greer, E. L. *et al.* Members of the H3K4 trimethylation complex regulate lifespan in a germline-dependent manner in C. elegans. *Nature* 466, 383 (2010).
13. Han, S. *et al.* Mono-unsaturated fatty acids link H3K4me3 modifiers to C. elegans lifespan. *Nature* 544, 185 (2017).
14. Zhang, Y. *et al.* Neuronal TORC1 modulates longevity via AMPK and cell nonautonomous regulation of mitochondrial dynamics in C. elegans. *Elife* 8, e49158 (2019).
15. Kenyon, C., Chang, J., Gensch, E., Rudner, A. & Tabtiang, R. A C. elegans mutant that lives twice as long as wild type. *Nature* 366, 461–464 (1993).
16. Liu, G. & Sabatini, D. mTOR at the nexus of nutrition, growth, ageing and disease. *Nat Rev Mol Cell Bio* 21, 183–203 (2020).
17. Dillin, A. *et al.* Rates of Behavior and Aging Specified by Mitochondrial Function During Development. *Science* 298, 2398–2401 (2002).
18. Luo, Q. *et al.* Mechanism of CREB recognition and coactivation by the CREB-regulated transcriptional coactivator CRTC2. *Proceedings of the National Academy of Sciences* 109, 20865–20870 (2012).
19. Yang, F. *et al.* An ARC/Mediator subunit required for SREBP control of cholesterol and lipid homeostasis. *Nature* 442, 700–704 (2006).
20. Hansen, M. *et al.* Lifespan extension by conditions that inhibit translation in Caenorhabditis elegans. *Aging Cell* 6, 95–110 (2007).
21. Pan, K. Z. *et al.* Inhibition of mRNA translation extends lifespan in Caenorhabditis elegans. *Aging Cell* 6, 111–119 (2007).
22. Troemel, E. R. *et al.* p38 MAPK Regulates Expression of Immune Response Genes and Contributes to Longevity in C. elegans. *Plos Genet* 2, e183 (2006).
23. Iourgenko, V. *et al.* Identification of a family of cAMP response element-binding protein coactivators by genome-scale functional analysis in mammalian cells. *Proceedings of the National Academy of Sciences of the United States of America* 100, 12147–52 (2003).
24. Canettieri, G. *et al.* The coactivator CRTC1 promotes cell proliferation and transformation via AP-1. *Proceedings of the National Academy of Sciences* 106, 1445–1450 (2009).
25. Wang, Y., Vera, L., Fischer, W. H. & Montminy, M. The CREB coactivator CRTC2 links hepatic ER stress and fasting gluconeogenesis. *Nature* 460, 534–537 (2009).
26. Kwon, A. T., Arenillas, D. J., Hunt, R. W. & Wasserman, W. W. oPOSSUM-3: Advanced Analysis of Regulatory Motif Over-Representation Across Genes or ChIP-Seq Datasets. *G3 Genes Genomes Genetics* 2, 987–1002 (2012).
27. Sui, S. J. H., Fulton, D. L., Arenillas, D. J., Kwon, A. T. & Wasserman, W. W. oPOSSUM: integrated tools for analysis of regulatory motif over-representation. *Nucleic Acids Res* 35, W245–W252 (2007).
28. Sui, S. J. H. *et al.* oPOSSUM: identification of over-represented transcription factor binding sites in co-expressed genes. *Nucleic Acids Res* 33, 3154–3164 (2005).
29. Burkewitz, K. *et al.* Atf-6 Regulates Lifespan through ER-Mitochondrial Calcium Homeostasis. *Cell Reports* 32, 108125 (2020).
30. Silva-García, C. G. *et al.* Single-Copy Knock-In Loci for Defined Gene Expression in Caenorhabditis elegans. *G3: Genes, Genomes, Genetics* 9, g3.400314.2019 (2019).
31. Li, T. Y. *et al.* The transcriptional coactivator CBP/p300 is an evolutionarily conserved node that promotes longevity in response to mitochondrial stress. *Nat Aging* 1, 165–178 (2021).
32. Tasoulas, J., Rodon, L., Kaye, F. J., Montminy, M. & Amelio, A. L. Adaptive Transcriptional Responses by CRTC Coactivators in Cancer. *Trends Cancer* 5, 111–127 (2019).
33. Greer, E. L. *et al.* Transgenerational epigenetic inheritance of longevity in Caenorhabditis elegans. *Nature* 479, 365 (2011).
34. Caron, M. *et al.* Loss of SET1/COMPASS methyltransferase activity reduces lifespan and fertility in Caenorhabditis elegans. (2021) doi:10.1101/2021.06.07.447374.
35. Silva-García, C. G. & Mair, W. B. Confirming the Pro-longevity Effects of H3K4me3-deficient set-2 Mutants in Extending Lifespan in C. elegans. *Biorxiv*

2022.08.02.502497 (2022) doi:10.1101/2022.08.02.502497.

36. Boon, R., Silveira, G. G. & Mostoslavsky, R. Nuclear metabolism and the regulation of the epigenome. *Nat Metabolism* 1–14 (2020) doi:10.1038/s42255-020-00285-4.

37. Paix, A., Folkmann, A., Rasoloson, D. & Seydoux, G. High Efficiency, Homology-Directed Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein Complexes. *Genetics* 201, 47–54 (2015).

38. Evans, T. Transformation and microinjection. *WormBook* (2006) doi:10.1895/wormbook.1.108.1.

39. Weir, H. J. *et al.* Dietary Restriction and AMPK Increase Lifespan via Mitochondrial Network and Peroxisome Remodeling. *Cell Metab* 26, 884-896.e5 (2017).

40. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21 (2013).

41. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* 14, 417–419 (2017).

42. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. *F1000research* 4, 1521 (2015).

43. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15, 550 (2014).

44. Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. *Omics J Integr Biology* 16, 284–287 (2012).

45. Lascarez-Lagunas, L. I., Herruzo, E., Grishok, A., San-Segundo, P. A. & Colaiácovo, M. P. DOT-1.1-dependent H3K79 methylation promotes normal meiotic progression and meiotic checkpoint function in C. elegans. *Plos Genet* 16, e1009171 (2020).

46. Rappsilber, J., Ishihama, Y. & Mann, M. Stop and Go Extraction Tips for Matrix-Assisted Laser Desorption/Ionization, Nanoelectrospray, and LC/MS Sample Pretreatment in Proteomics. *Anal Chem* 75, 663–70 (2003).

47. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* 26, 1367–1372 (2008).

48. Cox, J. *et al.* Andromeda: A Peptide Search Engine Integrated into the MaxQuant Environment. *J Proteome Res* 10, 1794–1805 (2011).

49. Angeles-Albores, D., Lee, R. Y. N., Chan, J. & Sternberg, P. W. Tissue enrichment analysis for C. elegans genomics. *Bmc Bioinformatics* 17, 366 (2016).

EXTENDED DATA AND SUPPLEMENTARY TABLES TITLES

Extended Data Fig. 1. CRTC-1 specifically regulates H3K4me3-mediated longevity.

a, Lifespan curve showing that *crtc-1^null* mutant has a similar lifespan to wild-type animals. **b-i**, *crtc-1^null* mutant has no effect on lifespan extension by *wdr-5.1 RNAi* (**b-c**), reduces *ash-2 RNAi*-mediated longevity (**d-e**), and fully suppresses lifespan extension in *set-2 RNAi* (**f-g**) animals and *set-2(ok952)* mutants (**h-i**). **j-o**, Lifespan curves showing that CRTC-1 is not required to extend lifespan in *raga-1(ok386)* (**j-k**), *clk-1(qm30)* (**l-m**), and *daf-2 RNAi* (**n-o**) worms. See Supplementary Table 2 for lifespan statistics.

Extended Data Fig. 2. CRTC-1^CBD drives a transcriptional shift in H3K4me3-deficient animals.
a, Principal component analysis of the RNA-Seq data showing similarity between samples. **b**, Over-representation analysis of the Gene Ontology biological process terms for genes comprising all comparisons from the RNA-Seq samples: wild-type, *set-2*, *crtc-1^CBD-/-*, and *crtc-1^CBD-/-;set-2* strains.

Extended Data Fig. 3. Global levels of trimethylation and acetylation in the SET-2-CRTC-1^CBD pathway.
a-d, Quantification of Western blots of wild-type, *set-2*, *crtc-1^CBD-/-*, and *crtc-1^CBD-/-;set-2* worms for H3K4me3 (**a**), H3K9Ac (**b**), H3K18Ac (**c**), and H3K27Ac (**d**) marks. H3 levels were used as loading controls. n = 3 independent blots.

Supplementary Table 1. List of proteins that interact with CRTC-1.

Supplementary Table 2. Lifespan statistics.

Supplementary Table 3. List of differential gene expression (DGE), gene ontology (GO), and tissue enrichment analyses from RNA-seq.

Supplementary Table 4. List of AP-1 targets in up > down group.

Supplementary tables and other reagents generated in this study are available upon request to the lead contact.

C. elegans strains	Source
N2: wild isolate	Caenorhabditis Genetics Center
WBM1241: *crtc-1(wbm54)* I [CRTC-1::3xFLAG]	This paper
WBM1254: *crtc-1(wbm57)* I [*crtc-1^{null}*]	This paper
RB1025: *set-2(ok952)* III [WBM1272 6x outcrossed]	Laboratory of Anne Brunet
VC222: *raga-1(ok386)* II [WBM499 6x outcrossed]	Caenorhabditis Genetics Center
MQ130: *clk-1(qm30)* III	Caenorhabditis Genetics Center
WBM1285: *set-2(ok952)* III;*crtc-1(wbm57)* I	This paper
WBM1337: *crtc-1(wbm57)* I;*raga-1(ok386)* II	This paper
WBM1338: *crtc-1(wbm57)* I;*clk-1(qm30)* III	This paper
WBM935: *crtc-1(wbm5)* I [*crtc-1$^{Sec31-/-}$*]	This paper
WBM1041: *crtc-1(wbm18)* I [*crtc-1$^{CBD-/-}$*]	This paper
WBM1286: *set-2(ok952)* III;*crtc-1(wbm18)* I	This paper
WBM1035: *set-2(ok952)* III;*crtc-1(wbm5)* I	This paper
ABR13: pEx141[*sbp-1p*::GFP::SBP-1 + *rol-6*(su1006)]	Laboratory of Anne Brunet
WBM945: *crtc-1(wbm5)* I;pEx141[*sbp-1p*::GFP::SBP-1 + *rol-6(su1006)*]	This paper
WBM1113: *crtc-1(wbm18)* I;pEx141[*sbp-1p*::GFP::SBP-1 + *rol-6(su1006)*]	This paper
AGD545: N2; UthEx540[pCRE4xTATA::GFP + pRF4]	Laboratory of Andrew Dillin
WBM1062: *crtc-1(wbm18)* I;UthEx540[pCRE4xTATA::GFP + pRF4]	This paper
WBM1114: *crtc-1(wbm5)* I;UthEx540[pCRE4xTATA::GFP + pRF4]	This paper
WBM936: *crtc-1(wbm8)* [CRTC-1::GFP]	This paper
WBM1207: N2, wbmIs92[*rab-3p*::3XFLAG::*crtc-1* cDNA::*rab-3* 3'UTR, *wbmIs66] VI	This paper
WBM1291: *set-2(ok952)* III;*crtc-1(wbm18)* I;wbmIs92[*rab-3p*::3XFLAG::*crtc-1* cDNA::*rab-3 3'UTR*, *wbmIs66] IV	This paper
WBM1196: N2, wbmIs85[*myo-3p*::3XFLAG::*crtc-1* cDNA::*unc-54* 3'UTR, *wbmIs61] I	This paper
WBM1294: *set-2(ok952);crtc-1(wbm18)*;wbmIs85[*myo-3p*::3XFLAG::*crtc-1* cDNA::*unc-54* 3'UTR, *wbmIs61] I	This paper
WBM1600: N2, wbmIs147[*nep-17p*::3XFLAG::*crtc-1* cDNA::*unc-54* 3'UTR, *wbmIs136] V	This paper
WBM1608: *set-2(ok952);crtc-1(wbm18)*;wbmIs147[*nep-17p*::3XFLAG::*crtc-1* cDNA::*unc-54* 3'UTR, *wbmIs136] V	This paper
crh-1(n3315) III	Laboratory of Mark Alkema
WBM1037: *set-2(ok952)* III;*crh-1(n3315)* III	This paper
VC3149: *crh-2(gk3293)* II	Caenorhabditis Genetics Center
WBM1132: *set-2(ok952)* III;*crh-2(gk3293)* II	This paper
RB772: *atf-6(ok551)* X [WBM841: 8x outcrossed]	Caenorhabditis Genetics Center
WBM1130: *set-2(ok952)* III;*atf-6(ok551)* X	This paper
VC1200: *jun-1(gk557)* II	Caenorhabditis Genetics Center
WBM1131: *set-2(ok952)* III;*jun-1(gk557)* II	This paper
WBM1432: N2; wbmEx415[pCS3(*eft-3p::pcaf-1* cDNA::*unc-54* 3'UTR) + pCFJ104(*myo-3p*::mCherry::*unc-54* 3'UTR)]	This paper
WBM1433: N2; wbmEx416[pCS3(*eft-3p::pcaf-1* cDNA::*unc-54* 3'UTR) + pCFJ104(*myo-3p*::mCherry::*unc-54* 3'UTR)]	This paper
WBM1434: N2; wbmEx417[pCS3(*eft-3p::pcaf-1* cDNA::*unc-54* 3'UTR) + pCFJ104(*myo-3p*::mCherry::*unc-54* 3'UTR)]	This paper

Oligonucleotoides
CGSG79: agcgcaattgcatcacacatct, *crtc-1::3xFLAG*
CGSG245: agagatagagacatggtgccagg, *crtc-1::3xFLAG*
CGSG94: tcttttcccgtctctctctccc, *crtc-1^{null}*
CGSG296: gtcgttctggtcttcctcgttg, *crtc-1^{null}*
CGSG80: atcgacacaagcggctactgta, *crtc-1^{null}*
CGSG293: tcgcgcgaacatttgaatttgtc, *set-2(ok952)*
CGSG294: ctcgtatggagtttgccggaag, *set-2(ok952)*
CGSG295: gaaggcatcagtttcgacccc, *set-2(ok952)*
raga-1(ok386) ExtF: tgctcatgtcaagtaagggatcaattc
raga-1(ok386) IntF: ttcaagtccgaaacagtcaattctc
raga-1(ok386) R: ggaactgaagcgatcacaccgac
clk-1(qm30) F: ctagaactccttgccgacgatcctg
clk-1(qm30) R: gaaatcaataattttttggggggaatcgcattac
CGSG94: tcttttcccgtctctctctccc, *crtc-1$^{CBD-/-}$*
CGSG95: acgcatcctatgaatgttcctacatc, *crtc-1$^{CBD-/-}$*

CGSG70: caatgcccattccgcaacag, *crtc-1*^{Sec31}
CGSG71: gggcgatcgtgataatccattcg, *crtc-1*^{Sec31}
CGSG104: gaactttgcagtttggttgtagtg, wbmIs92
CGSG106: tcaatccgttcatttgagccc, wbmIs92
CGSG156: acagcgtgttatgtaacggagc, wbmIs92, wbmIs85, wbmIs147
CGSG79: agcgcaattgcatcacacatct, *crtc-1::GFP*
CGSG80: atcgacacaagcggctactgta, *crtc-1::GFP*
CGSG81: atcaccttcaccctctccactg, *crtc-1::GFP*
CGSG143: taagcttctcacacctttctctcg, wbmIs85
CGSG124: cctcccctcatctcaattatcccg, wbmIs85
CGSG205: accttcgacctcactttcccctc, wbmIs147
CGSG206: ccgtcctgaagtatacccagatcc, wbmIs147
crh-1 upper: cttgaaggcacatctggg
crh-1 lower 1: cattagcgtgattactctg
crh-1 lower 2: ccgaaacgcgattttccagac
crh-2 F: tgagggagaggcggaggacac
crh-2 R: atggctattcgacaacatctaatcactg
crh-2 Int: ttcggcgagattcttgtgcag
CGSG185: catgcgcattccttctatcaacac, *jun-1(gk557)*
CGSG186: tctccgacgaccagaccatc, *jun-1(gk557)*
CGSG187: cacgaatcgaattgttcgggc, *jun-1(gk557)*
CGSG152: taaagatcatgacatcgattacaaggatgacgatgacaagtcaaattcgaatacaccacg, *myo-3/rab3-p::crtc-1* SKI LODGE cloning
CGSG233: tgagtttttatagatagtataatagaacgtagaatttctaattttggagcatttgtgtac, *rab3-p::crtc-1* SKI LODGE cloning
CGSG232: aattggacttagaagtcagaggcacgggcgcgagatgctaattttggagcatttgtgtac, *myo3-p::crtc-1* SKI LODGE cloning
CGSG346: cccgggatgaaacaacgcgactctctgg, *pcaf-1* OE cloning
CGSG347: gcggccgcttaacatagaggtttttgctcggggg, *pcaf-1* OE cloning
pCS3: *eft-3p::pcaf-1 cDNA::unc-54 3'UTR*
pCFJ104: *myo-3p::mCherry::unc-54 3'UTR*