1 **Comparative genomic and metagenomic investigations of the *Corynebacterium***
2 ***tuberculostearicum* species complex reveals potential mechanisms underlying**
3 **associations to skin health and disease**
4

5 Rauf Salamzade[1], Mary Hannah Swaney[1], Lindsay R. Kalan[1,2]
6 [1]Department of Medical Microbiology and Immunology, School of Medicine and Public Health,
7 University of Wisconsin, Madison, WI, USA
8 [2]Department of Medicine, School of Medicine and Public Health, University of Wisconsin,
9 Madison, WI, USA
10
11 Address for Correspondence: Lindsay Kalan, 1550 Linden Dr, 6155 Microbial Sciences Building,
12 Madison, WI, USA, 53706. lkalan@wisc.edu

13
14 **ABSTRACT**
15

Corynebacterium are a diverse genus and dominant member of the human skin
microbiome. Recently, we reported that the most prevalent *Corynebacterium* species found on
skin – including *Corynebacterium tuberculostearicum* and *Corynebacterium kefirresidentii* –
comprise a narrow species complex despite the diversity of the genus. Here, we apply high-
resolution phylogenomics and comparative genomics to describe the structure of the *C.
tuberculostearicum* species complex. We find this species complex is missing a fatty acid
biosynthesis gene family which is often found in multi-copy in approximately 99% of other
Corynebacterium species. Conversely, this species complex is enriched for multiple genetic
traits, including a gene encoding for a collagen-like peptide. Further, through metagenomic
investigations, we find that one species within the complex, *C. kefirresidentii*, increases in
relative abundance during atopic dermatitis flares and show that most members of this species
possess a colocalized set of putative virulence genes.

INTRODUCTION

Several species of *Corynebacterium* have been isolated from human skin, often associated with diseases such as diphtheria and regarded as opportunistic pathogens[1,2]. Only recently, with the application of metagenomics to comprehensively survey the skin microbiome, have we begun to develop a better understanding of this understudied genus and assess mechanisms underlying their basis as core constituents of skin microbiomes[3–5].

Regarded as a lipophilic and disease-associated species, *C. tuberculostearicum* was first named and associated with cases of diphtheria in 1984[1], later found to be associated with rhinosinusitis[6], and more recently shown to elicit specific inflammatory signaling pathways of skin cells[5]. While it was noted that *C. tuberculostearicum* could be closely related to other *Corynebacterium* species when it was first validated as a species in 2004[7], this analysis was based on 16S sequences which provide limited resolution for such investigations. Later work suggested that misclassification of *C. tuberculostearicum* is likely prevalent by clinical biochemical assays and that the species is potentially underreported[8].

We recently found that the three most prevalent species of *Corynebacterium* in healthy skin microbiomes from a metagenomic survey in our lab[9], currently classified as *Corynebacterium tuberculostearicum*, *Corynebacterium kefirresidentii*, and *Corynebacterium aurimucosum* type E in GTDB[10], belong to a narrow clade, which includes at least one additional species[11]. Because the average nucleotide identity (ANI) between genomes in this clade is relatively high and the lowest value observed was at 88%, we refer to the clade as the *C. tuberculostearicum* species complex[10,11]. While *C. aurimucosum* and *C. tuberculostearicum* have been validated as species[7,12], reassessment of genomes classified as these species in NCBI's GenBank database by GTDB suggests that some are mislabelled[10]. *C. kefirresidentii* was only recently proposed as a new species and has not yet been validated by the International Committee on Systematics of Prokaryotes[13].

Isolates of *C. kefirresidentii* sampled from healthy human skin are lipophilic similar to *C. tuberculostearicum* and also grow better in the presence of compounds commonly found in sweat[9]. This finding is in accordance with our earlier observation that members of the *C. tuberculostearicum* species complex are often found at body sites regarded as moist[11]. In this study, we apply comparative genomics and metagenomics analytics to develop a better understanding of the genetic factors underlying the prevalence of the *C. tuberculostearicum* species complex on healthy skin and determine they lack key metabolic enzymes likely rendering them lipid-dependent. In addition, we assess the biogeographical distribution of individual species within the complex across different skin body sites and whether they associate with atopic dermatitis flares.

RESULTS

A high-resolution phylogeny of the *C. tuberculostearicum* species complex

A set of 26 genomes belonging to the *C. tuberculostearicum* species complex were used for phylogenomic investigation of the complex (Table S1). This set of genomes includes 22 genomes we had previously described as part of the complex[11]. In addition, we added three

recently reported metagenome-assembled genomes (MAGs) from skin predicted to correspond
to novel *Corynebacterium* species[14] and the genome of a *C. kefirresidentii* we recently isolated
from human skin[9]. All 26 genomes, including a total of 8 MAGs, were regarded as high
completion and showed low rates of contamination by CheckM analysis[15] (Table S1).
A set of 1,250 single-copy-core (SCC) orthologs was identified amongst the 26 genomes
belonging to the species complex and used to construct a high-resolution phylogeny following
filtering of sites predicted to be affected by recent or ancestral recombination[16,17] (Figure 1). This
phylogeny was found to be highly concordant with species classifications for the genomes by
GTDB-tk[18] run with GTDB release 207[10], but not their taxonomic designations on GenBank
(Table S1).
ANI between genomes belonging to different species ranged from a lower limit of
~88%[11] to nearly 95% (Figure S1). This suggests recent speciation occurring within the species
complex, in particular within the subclade containing the species *Corynebacterium*
sp900539985, *C. aurimucosum* type E*,* and *C. tuberculostearicum*, where genomes from
different species can exhibit ANI greater than 94% to each other, which is slightly under the
95% ANI threshold commonly used to delineate species[19,20]. In addition, the ANI between
genomes *C. spp.* SMGC_38 and *C. spp.* SMGC_337 was estimated as being between 93% and
95%, suggesting recent speciation between them. These two newly reported species[14],
exhibited greater than 90% ANI to *C. kefirresidentii* genomes (Figure S1). The MAG proposed to
correspond to a novel *Corynebacterium* species[14], *Corynebacterium spp.* SMGC_92, was found
to exhibit high ANI to an established but currently unnamed species in GTDB, *C.* sp900539985,
and designated as belonging to this species. Due to close phylogenetic placement and ANI
estimates exceeding 95%, we further propose that GTDB species *C. tuberculostearicum* and *C.*
tuberculostearicum type C should simply be regarded as one species (*C. tuberculostearicum*).
Overall, the phylogeny constructed using the core-genome with sites predicted to be
affected by recombination removed, was well supported as indicated by high bootstrap values at
ancestral nodes to species and sets of species (Figure 1). However, we were unable to
confidently resolve whether *C. sp900539985* are more related to *C. tuberculostearicum*, as ANI
analysis suggests, or *C. aurimucosum* (Figure 1, S1).
101
102   **Unique genes and the biogeographic distribution of the *C. tuberculostearicum* species**
103   **complex**
104
To address potential biases predicting prevalence of the *C. tuberculostearicum* species
complex associated with the metagenomic survey performed in our laboratory[4], we used
StrainGE[21] to profile the presence of representative *Corynebacterium* genomes amongst
published metagenomic datasets profiling the healthy skin microbiome[3] including a study
profiling the skin microbiome in atopic dermatitis, a common pediatric allergic disease[22]. In
concordance with our previous findings[11], we observed that all three species from the complex
represented in the StrainGE database are highly prevalent in these additional metagenomic
datasets relative to other *Corynebacterium* species (Figure 2A; Table S2). We further observed
that *C. tuberculostearicum* and *C. kefirresidentii* were differentially abundant at distinct body
sites. While *C. tuberculostearicum* were prevalent and found in high abundances at foot

associated body sites, in particular the toe web space, *C. kefirresidentii* were more prevalent in
nasal and surrounding alar crease sites (Figure 2B).

To understand the basis of the enrichment of the *C. tuberculostearicum* species
complex, we performed comparative genomics to determine if particular homolog groups were
found more commonly or less commonly than expected within the complex relative to the rest of
the genus (Table S3). A representative set of 187 *Corynebacterium* genomes was selected
based on dereplication[23] of those designated as belonging to the genus in GTDB at 95% ANI, to
approximate selection of a single representative genome per species.

We identified only two homolog groups which were statistically significant for being
absent in the species complex but highly conserved throughout the genus. One of these was
annotated as a short-chain dehydrogenase reductase and was absent in all four representative
genomes of the *C. tuberculostearicum* species complex but found at a median copy count of
three in 98.9% other *Corynebacterium*. Searching for highly conserved domains on the
consensus sequence of the homolog group revealed a more precise annotation of 3-oxoacyl-
ACP reductase FabG[24], thus suggesting a role in fatty acid biosynthesis[25].

A larger set of proteins were identified as being statistically enriched in the *C.*
tuberculostearicum species complex in relation to other *Corynebacterium*. We filtered for 26
homolog groups which were found to be nearly core to the species complex (found in $\geq$95% of
the 26 *C. tuberculostearicum* genomes). While the majority of these homolog groups lacked
reliable functional annotations, we identified three to putatively correspond to acetyltransferases
and an additional three to be factors in cellular regulation or translation. The largest homolog
group, found in only 3.8% of *Corynebacterium* outside of the species complex, was predicted to
encode a collagen-like peptide. Collagen-like proteins have previously been suggested to
function as adhesins of pathogens to host cells[26–29] and could correspond to an analogous
mechanism in the *C. tuberculostearicum* species complex for attaching to skin cells.

Species specific partitioning into distinct moist body sites and association with atopic
dermatitis

Because prior taxonomic classifications of genomes were incongruent with their
phylogenomic placement, it was previously impossible to accurately associate individual species
within the *C. tuberculostearicum* species complex with skin-related diseases. With our resolved
view of the species complex, we thus reinvestigated the metagenomic dataset from Byrd *et al.*
to assess the relative abundance of *C. tuberculostearicum*, *C. kefirresidentii*, and type E *C.*
aurimucosum at different stages of atopic dermatitis for a pediatric cohort. We found that *C.*
kefirresidentii was more frequently detected during flares, in comparison to baseline (*p*=1.5e-4)
and healthy control microbiomes (*p*=4.0e-4). Furthermore, in certain metagenomes sampled
during a flare, *C. kefirresidentii* exhibited a substantial increase in relative abundance (Figure
3A; S2).

To further affirm the association of *C. kefirresidentii* with atopic dermatitis flares, we also
applied an alignment-based approach in which we aligned metagenomic reads to the single-
copy core genes of the complex and measured coverage of species specific alleles. *C.*
kefirresidentii was more evidently present at higher abundances primarily in metagenomes from
flares and post-flare samplings (Figure 3B), which also tended to have the lowest viable

sequencing data after filtering metagenomes for quality, adapter sequences, and human
contamination (Figure S3). We further developed a ratio statistic to assess the abundance of *C.*
kefirresidentii to *C. tuberculostearicum* and type E *C. aurimucosum* individually within each
metagenome. *C. kefirresidentii* were more abundant than type E *C. aurimucosum* and *C.*
tuberculostearicum combined in 52.7% of metagenomes survey from flares, which was
significantly more than for baseline (25.7%; *p*=2.3e-4) and healthy control (21.1%; *p*=3.8e-5)
metagenomes (Figure 3C).
To understand what potential genetic factors might be contributing to the unique
association of *C. kefirresidentii* with atopic dermatitis flares, we performed comparative
genomics to identify traits which were lost or gained in their genomes relation to the closely
related *C. tubercuostearicum* and type E *C. aurimucosum* species. We found a total of 97
homolog groups were either enriched or depleted in *C. kefirresidentii* relative to *C.*
tubercuostearicum and type E *C. aurimucosum* (Table S4). *C. tubercuostearicum* and type E *C.*
aurimucosum feature a core thermonuclease that is homologous to the virulence factor *nuc*[30]
from *Staphylococcus aureus*, but missing in all *C. kefirresidentii.* In contrast, two distinct co-
located sets of genes, one predicted to encode for thiamine biosynthesis and the other encoding
molybdopterin-binding enzymes, were enriched in *C. kefirresidenttii* but missing or significantly
lacking in *C. tubercuostearicum* and type E *C. aurimucosum* genomes. Of note, molybdenum
enzymes have recently been discussed as important virulence factors for infection in a variety of
diverse pathogens[31,32].
179
180   **DISCUSSION**
181
The *C. tuberculostearicum* species complex features the three major species of
Corynebacterium found on human skin[11]. Despite this, fewer than 30 representative genomes
across all species belonging to the complex are present in the most recent GTDB release and
eight of the 26 genomes (30.8%) used in this study were MAGs. The lack of genomes for the
species complex, despite its prevalence on healthy skin, might be in part due to difficulties with
culturing. For instance, we recently found that a single *C. kefirresidentii* isolate grows best when
high concentrations of compounds found in both sebum and sweat are available[9].
Here we determine that the *C. tuberculostearicum* species complex appears to be
missing a fatty acid biosynthesis gene family that is often found in multi-copy in other
Corynebacterium, potentially explaining their lipophilic nature[7,8]. However, other
Corynebacterium species outside of this species complex
have also been characterized as lipophilic[9,33], suggesting convergent evolution with different
genetic factors underlying the shared trait in diverged species across the genus. Comparative
genomics further highlighted several genes encoding proteins that are specific to the species
complex, though the majority of these lack any functional annotation and future molecular
characterization will be essential to further our functional understanding of them. Interestingly,
one of these genes, core to the species yet present in less than 4% of other *Corynebacterium*,
encodes for a collagen-like peptide and could be involved in adherence to skin.
We further identify an increase in relative abundance of *C. kefirresidentii* in
metagenomes sampled during atopic dermatitis flares. Comparative genomics between *C.*
kefirresidentii with *C. tuberculostearicum* and *C. aurimucosum* type E genomes found that

multiple genes differentiate *C. kefirresidentii* from its two neighboring species. Most striking are
a set of five colocated molybdopterin-associated enzymes, which are commonly associated with
virulence[31,32].
Two ATCC isolates (ATCC-35693 / 2628 LB and ATCC-35694 / FPSA)[34], designated as
C. tuberculostearicum and originating from the original study first describing the species taken
from leprosy patients[1], have more recently been used in studies to associate the species to
rhinosinusitis[6] and elicitation of a specific host inflammatory signaling pathway[5]. To our
knowledge, these isolates lack genomes however and might currently be misclassified since
other species of *Corynebacterium* have been reported to similarly produce tuberculostearic acid,
the trait initially used to differentiate the species from others in the genus[7,12,35]. Identifying
genetic markers to distinguish species within the *C. tuberculostearicum* species complex and
continued efforts in whole genome and metagenomic sequencing will thus be critical to correctly
associate the contributions of particular species within the complex to diseases.
216
217
218 **ACKNOWLEDGMENTS**
219

220 This work was supported by grants from the National Institutes of Health (NIAID U19AI142720
221 and NIGMS R35GM137828 [L.R.K]). The content is solely the responsibility of the authors and
222 does not necessarily represent the official views of the National Institutes of Health. The authors
223 gratefully acknowledge J.Z. Alex Cheong, Dr. Nicole Lane Starr, Lucas van Dijk, Dr. Abigail
224 Manson, and Dr. Ashlee Earl for helpful discussions.

225
226 **METHODS**
227
Genome selection and taxonomic reclassification
We had previously identified a set of 22 genomes through an interactive process in
which we first identified the clade using phylogenomics as consisting of the five GTDB defined
species: *C. tuberculostearicum*, *C. tuberculostearicum*_C, *C. aurimucosum*_E, and *C.
kefirresidentii* (using release R202)[11]. Five additional genomes belonging to the clade were
identified in NCBI which were missing in GTDB release R202 and validated to belong to one of
the five species through GTDB-tk classification[18]. In our earlier study, we also found that three
recently published MAGs from skin microbiomes, proposed novel species of *Corynebacterium*
(based on GTDB release 88)[14], had >88% ANI to one of the prior defined 22 genomes. We thus
included these three genomes, in addition to the genome for a *C. kefirresidentii* isolate, LK1134,
we had isolated from healthy skin and profiled the growth dynamics off across different
concentrations of artificially synthesized sebum and sweat[9].
Systematic reclassification on the total set of 26 genomes proposed to belong to the *C.
tuberculostearicum* species complex was performed using GTDB-tk2 with the latest GTDB
release R207. All genomes were assigned to one of the five previously mentioned GTDB
species, except for the MAGs *C. spp.* SMGC_38 and *C. spp.* SMGC_337[14], which appear to still
correspond to novel species of close relation to *C. kefirresidentii*. Because MAGs were included,
the contamination and completeness of genomes was assessed using CheckM (v1.2.1) (Table
S1).

For comparative genomics of the *C. tuberculostearicum* species complex against other
species from the *Corynebacterium* genus, all 1,296 genomes in the genus from GTDB R207
were downloaded[10]. Genome dereplication was performed using dRep[23] with a secondary ANI
filter of 95% using fastANI[19]. This retained a set of 187 distinct genomes, including four
representatives of the *C. tuberculostearicum* species complex.
252
Homolog group inference and functional annotation
Gene calling was performed on genomes using prodigal[36] and OrthoFinder (v2.5.4)[16]
was subsequently used to perform homolog group detection independently amongst the set of
26 *C. tuberculostearicum* genomes and the genus-wide set of 187 representative
Corynebacterium genomes. For both analyses, the course set of initial orthogroups following
MCL were used instead of phylogenetic hierarchical orthogroups.
Functional annotation was only performed for homolog groups which comparative
genomics investigations suggested were significantly more prevalent or lacking within certain
phylogenetic clades. Annotation was primarily performed using individual protein sequences
from such homolog groups via the EggNOG mapper webserver run with default settings[37].
Consolidation of annotations was performed across proteins belonging to each homolog group.
For homolog groups predicted to be enriched or depleted in the *C. tuberculostearicum* species
complex relative to other *Corynebacterium*, we further attempted annotation by generating
consensus amino acid sequences for each homolog group and using NCBI's conserved domain
search[24] as well as Phyre2 to assess structural similarity to structurally characterized proteins[38].
268
Phylogenomics of the *C. tuberculostearicum* species complex and the representative
selection of *Corynebacterium*
We identified 1,250 homolog groups as single-copy orthologs between the 26 *C.*
tuberculostearicum species complex genomes. Protein alignments were constructed using
MUSCLE (v5)[39] and converted to codon alignments using PAL2NAL[40]. RAxML (v8.2.12) was
then used to construct an initial maximum likelihood phylogeny on a concatenation of codon
alignments (1,239,366 bp) with the GTRCAT model and 1,000 bootstraps. Core codon
alignments were projected onto and ordered according to the reference genome *C.*
kefirresidentii FDAARGOS_1055 and, together with the RAxML phylogeny, used as input for
ClonalFrameML (v1.12)[17] to infer sites affected by recombination by first using a simple model,
with parameter emsim set to 100 and kappa (the relative rate of transitions to transversions) set
to 5.09664. For reference projection, gaps between single-copy orthologs were set to be the
length of distances in between corresponding genes in the reference genome and codon
alignments were reverse complemented if the gene was predicted to occur on the antistrand.
Non-core sites were specified as the gaps placed in between single copy core orthologs so as
to appropriately model the strength of linkage between them. Results from running
ClonalFrameML with a simple model were used to initialize values for R/theta, 1/delta, and nu
(set to 0.126997, 0.0027686, and 0.0583416, respectively) and run ClonalFrameML with a per-
branch model with the embranch_dispersion parameter set 0.1, to allow for greater dispersion in
parameters amongst branches. The initial alignment was filtered for sites predicted to be
affected by recombination and subsequently subset for 15,936 sites which lacked ambiguity for
any of the 26 genomes. This alignment was finally used to construct a high-resolution,

maximum likelihood phylogeny of the species complex using RAxML with the same modeling
strategy as applied initially.
For phylogeny construction of the full genus, we used GToTree[41] (v1.6.36) with the set
of single-copy gene sets specific to Actinomycetota (formerly Actinobaceria) and FastTree2
phylogeny inference[42].
296
Statistical testing for enrichment or depletion of homolog groups in clades for
comparative genomics analyses
Assessment of enrichment or depletion of homolog groups for two sets of genomes was
carried out using a two-sided permutation test based on mean copy-count differences with
100,000 resamples. This allowed us to detect homolog groups which were not merely
differentially present, but also those which had greater copy-counts in one set of genomes
relative to the other. A generalizable program for this testing is provided in the Github repository:
https://github.com/Kalan-Lab/Salamzade_etal_CtuberCompGen. Multiple testing correction was
performed for both such comparative genomics analyses using the Benjamini-Hochberg
procedure for P-value adjustment to control the false discovery rate. For homolog groups found
to be enriched in the four *C. tuberculostearicum* species complex representative genomes
relative to the complementary set of 183 *Corynebacterium* genomes, we further required that
they map to homolog groups deemed to be $\geq$95% near-core amongst the 26 *C.*
tuberculostearicum species complex genomes (there was a total of 1,871 near-core homolog
groups).
312
Metagenomic detection of *Corynebacterium* strains and species
Three independent sets of metagenomes profiling the microbiome of healthy skin and
skin affected by atopic-dermatitis were processed using a pipeline previously described[4]. Briefly,
metagenomes were processed for quality and human contamination using fastp[43] and
KneadData (https://github.com/biobakery/kneaddata). Mock and negative control metagenomes
from the Swaney *et al.* 2022 study were excluded[4] as were 22 metagenomes from the Oh *et al.*
2016 dataset marked as "human depletion" or "whole genome amplification" under "Notes
(excluded for comparisons)" in their spreadsheet describing all samples[3]. Deeply sequenced
metagenomes from Oh *et al.* 2016 were randomly downsampled to 25 million read pairs using
seqtk (https://github.com/lh3/seqtk). In addition, as described previously[4], five metagenomes
from the Byrd *et al.* 2017 dataset[22] (3 flare and 2 post-flare) were excluded due to insufficient
sequencing data following processing. All symmetric metagenomes (e.g. metagenomes
corresponding to sampling the left and right sides of a body site) from the Oh *et al.* 2016 and
Byrd *et al.* 2017 metagenomes were individually considered.
Detection of strains or representative genomes was performed using StrainGST[21] with a
custom database constructed from complete *Corynebacterium* genomes from NCBI, as
described previously[11]. Briefly, this database featured 164 distinct representative genomes
following dereplication using programs for StrainGST (0.1+191.g7fcfbcd.dirty) database
creation[21]. These genomes were run through GTDB-tk with database release R207[10,18] to
generate consistent taxonomic classifications. One genome regarded as *Corynebacterium* in
RefSeq[44] at the time of database creation was classified as not part of the genus by GTDB-tk
and not observed in any of the metagenomic datasets. Another genome was classified as a

novel *Corynebacterium spp.* not yet given an identifier in GTDB R207. We counted the number
of distinct species detected in metagenomes from the three datasets and illustrated them as a
multi-bar graph on the phylogeny of 187 *Corynebacterium* by species name. Because *C.*
tuberculostearicum and *C. tuberculostearicum* type C classified genomes were found in this
study to phylogenetically group as one, we considered them as a single species and mapping of
detection counts for the *C. tuberculostearicum* representative genome in StrainGST was
allowed to the *C. tuberculostearicum* type C representative genome in the set of 187 genomes
used for phylogeny construction and comparative genomics. The species *C. sp900539985* from
the *C. tuberculostearicum* species complex, which was represented in the phylogeny of 187
genomes, was not represented in the StrainGST database as it lacks a complete genome
currently. Further, there was a single representative strain / genome for each of the three
species from the complex represented in the StrainGST database: *C. tuberculostearicum*, *C.*
aurimucosum_E, and *C. kefirresidentii*. The relative abundance estimation for these three
species was thus computed based on the relative abundance computed for the corresponding
representative strains / genomes by StrainGST (*rapct*) after updating to version 1.3.3, to
incorporate updates to how relative abundance was being computed relative to earlier versions.
Confirmation of StrainGST results using an independent approach: StrainGST is a k-mer
based approach to assess strain presence in metagenomes and to complement it we used an
independent and complementary alignment based approach to gauge whether *C. kefirresidentii*
were enriched during atopic dermatitis flares in the Byrd *et al.* 2017 metagenomes. For this
approach, we aligned metagenomic reads individually to 4,514 representative genes from 367
homolog groups which comparative genomics had revealed were single-copy orthologs and
featured at least 20 sites where *C. kefirresidentii* had a core allele that was not observed in the
complementary set of genomes in the *C. tuberculostearicum* species complex. Reads were
aligned as unpaired using Bowtie 2[45] with parameters "--very-sensitive-local --no-unal -a -x".
Reads were aligned as unpaired because gene sequences are short and pairs can thus align
poorly. Additionally, all alignments for each read are requested because multiple representative
genes were considered per homolog group. We then processed alignments to each
representative gene where coverage at 1X was not observed for 90% of the sites. Following this
preliminary scan for checking that genes were loosely covered, we assessed alignments to the
gene met either of the following two criteria: (i) $\geq$99% identity with length $\geq$60 bp or (2) $\geq$95%
identity with length $\geq$100 bp for the core alignment region (excluding flanking regions to account
for alignments hanging off representative gene edges). We also allowed a maximum of 5 indelic
sites within the core alignment region for the alignment to be considered. Alignments which met
these criteria were then further processed and retained if they had the best mapping score
observed for a particular read (in the case of ties, all alignments with the best Bowtie 2 mapping
score were retained). Finally, allele counts were tallied for each metagenome at each site in
homolog group codon alignments based on mapping of the query base to the reference /
representative gene position in the codon alignments. Only positions in reads which had greater
than 30 base quality were considered.
Results from determining the presence of alleles at sites along homolog group codon
alignments for each metagenome were processed to identify the proportion of *C. kefirresidentii*
specific alleles covered by at least three reads. The mean coverage of these well-covered *C.*
kefirresidentii specific alleles were computed and normalized for sequencing depth through

dividing by a metagenome specific scalar value (Figure S3). This scalar value corresponded to
the ratio of the number of bases the metagenome featured in relation to the metagenome with
the fewest number of bases in the final forward read files following processing.
We additionally identified *C. tuberculostearicum* and type E *C. aurimucosum* specific
alleles along the same set of 367 core homolog groups and quantified the mean coverage of
species specific alleles within each metagenome (inclusive of coverage at alleles which were
uncovered). This allowed us to compute a ratio statistic measuring the abundance of *C.*
kefirresidentii to *C. tuberculostearicum* and type E. *C. aureimucosum* with a pseudocount of 1
added to both the numerator and denominator to avoid dividing by zero. A value greater than 1
for this statistic thus indicates that *C. kefirresidentii* are more prevalent than *C.*
tuberculostearicum and type E. *C. aureimucosum* within an individual metagenome; whereas, a
value less than 1 supports the opposite. Our code for metagenomic alignment and calculation of
the ratio statistic is provided in the Github repository: https://github.com/Kalan-
Lab/Salamzade_etal_CtuberCompGen.
Statistical testing: To test whether StrainGST-based detection of *C. kefirresidentii* was
significantly more prevalent within metagenomes in atopic dermatitis flare metagenomes relative
to baseline and healthy control metagenomes, we used the two-sided Fisher's exact test. The
two-sided Fisher's exact test was also used to statistically assess whether the ratio statistic was
larger than 1 in metagenomes sampled during atopic dermatitis flares relative to baseline and
healthy control metagenomes.
399
400 **Visualizations**
401 　　　　iTol was used for illustrating phylogenetic trees and associated features[46]. The R library
402 ggplot2[47] was used to visualize heatmaps, scatterplots, and bar graphs. Circleator[48] was used to
403 generate an overview of conservation and showcase genes highlighted by comparative
404 genomics along the chromosome of *C. kefirresidentii* FDAARGOS_1055.
405
REFERENCES

1. Brown, S., Lanéelle, M. A., Asselineau, J. & Barksdale, L. Description of Corynebacterium
tuberculostearicum sp. nov., a leprosy-derived Corynebacterium. *Ann. Microbiol.* **135B**,
251–267 (1984).
2. Bernard, K. The genus corynebacterium and other medically relevant coryneform-like
bacteria. *J. Clin. Microbiol.* **50**, 3152–3158 (2012).
3. Oh, J. *et al.* Temporal Stability of the Human Skin Microbiome. *Cell* **165**, 854–866 (2016).
4. Swaney, M. H., Sandstrom, S. & Kalan, L. R. Cobamide Sharing Is Predicted in the Human
Skin Microbiome. *mSystems* e0067722 (2022).
5. Altonsy, M. O. *et al.* Corynebacterium tuberculostearicum, a human skin colonizer, induces
the canonical nuclear factor-κB inflammatory signaling pathway in human skin cells.
Immun. Inflamm. Dis. **8**, 62–79 (2020).
6. Abreu, N. A. *et al.* Sinus microbiome diversity depletion and Corynebacterium
tuberculostearicum enrichment mediates rhinosinusitis. *Sci. Transl. Med.* **4**, 151ra124
(2012).
7. Feurer, C. *et al.* Taxonomic characterization of nine strains isolated from clinical and
environmental specimens, and proposal of Corynebacterium tuberculostearicum sp. nov.
Int. J. Syst. Evol. Microbiol. **54**, 1055–1061 (2004).
8. Hinic, V. *et al.* Corynebacterium tuberculostearicum: a potentially misidentified and

multiresistant Corynebacterium species isolated from clinical specimens. *J. Clin. Microbiol.*
50, 2561–2567 (2012).

9.   Swaney, M. H., Nelsen, A., Sandstrom, S. & Kalan, L. R. Sweat and sebum preferences of
the human skin microbiota. doi:10.1101/2022.08.13.503869.

10. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy.
Nucleic Acids Res. (2021) doi:10.1093/nar/gkab776.

11. Salamzade, R. *et al.* lsaBGC provides a comprehensive framework for evolutionary
analysis of biosynthetic gene clusters within focal taxa. *bioRxiv* 2022.04.20.488953 (2022)
doi:10.1101/2022.04.20.488953.

12. Yassin, A. F., Steiner, U. & Ludwig, W. Corynebacterium aurimucosum sp. nov. and
emended description of Corynebacterium minutissimum Collins and Jones (1983). *Int. J.*
Syst. Evol. Microbiol. **52**, 1001–1005 (2002).

13. Blasche, S., Kim, Y. & Patil, K. R. Draft Genome Sequence of Corynebacterium
kefirresidentii SB, Isolated from Kefir. *Genome Announc.* **5**, (2017).

14. Saheb Kashaf, S. *et al.* Integrating cultivation and metagenomics for a multi-kingdom view
of skin microbiome diversity and functions. *Nat Microbiol* **7**, 169–179 (2022).

15. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

16. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238 (2019).

17. Didelot, X. & Wilson, D. J. ClonalFrameML: efficient inference of recombination in whole
bacterial genomes. *PLoS Comput. Biol.* **11**, e1004041 (2015).

18. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2019).

19. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.
Nat. Commun. **9**, 1–8 (2018).

20. Olm, M. R. *et al.* Consistent Metagenome-Derived Metrics Verify and Delineate Bacterial
Species Boundaries. *mSystems* **5**, (2020).

21. van Dijk, L. R. *et al.* StrainGE: a toolkit to track and characterize low-abundance strains in
complex microbial communities. *Genome Biol.* **23**, 74 (2022).

22. Byrd, A. L. *et al.* Staphylococcus aureus and Staphylococcus epidermidis strain diversity
underlying pediatric atopic dermatitis. *Sci. Transl. Med.* **9**, (2017).

23. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate
genomic comparisons that enables improved genome recovery from metagenomes through
de-replication. *ISME J.* **11**, 2864–2868 (2017).

24. Lu, S. *et al.* CDD/SPARCLE: the conserved domain database in 2020. *Nucleic Acids Res.*
48, D265–D268 (2020).

25. Gago, G., Diacovich, L., Arabolaza, A., Tsai, S.-C. & Gramajo, H. Fatty acid biosynthesis in
actinomycetes. *FEMS Microbiol. Rev.* **35**, 475–497 (2011).

26. Chen, S.-M. *et al.* Streptococcal collagen-like surface protein 1 promotes adhesion to the
respiratory epithelial cell. *BMC Microbiol.* **10**, 320 (2010).

27. Yu, Z., An, B., Ramshaw, J. A. M. & Brodsky, B. Bacterial collagen-like proteins that form
triple-helical structures. *J. Struct. Biol.* **186**, 451–461 (2014).

28. Bachert, B. A. *et al.* A Unique Set of the Burkholderia Collagen-Like Proteins Provides
Insight into Pathogenesis, Genome Evolution and Niche Adaptation, and Infection
Detection. *PLoS One* **10**, e0137578 (2015).

29. Qiu, Y., Zhai, C., Chen, L., Liu, X. & Yeo, J. Current Insights on the Diverse Structures and
Functions in Bacterial Collagen-like Proteins. *ACS Biomater Sci Eng* (2021)

doi:10.1021/acsbiomaterials.1c00018.

30. Berends, E. T. M. *et al.* Nuclease expression by Staphylococcus aureus facilitates escape
from neutrophil extracellular traps. *J. Innate Immun.* **2**, 576–586 (2010).

31. Zhong, Q., Kobe, B. & Kappler, U. Molybdenum Enzymes and How They Support Virulence
in Pathogenic Bacteria. *Front. Microbiol.* **11**, 615860 (2020).

32. Williams, M. J. *et al.* bis-Molybdopterin guanine dinucleotide is required for persistence of
Mycobacterium tuberculosis in guinea pigs. *Infect. Immun.* **83**, 544–550 (2015).

33. Tauch, A. *et al.* Complete genome sequence and analysis of the multiresistant nosocomial
pathogen Corynebacterium jeikeium K411, a lipid-requiring bacterium of the human skin
flora. *J. Bacteriol.* **187**, 4671–4682 (2005).

34. Institute of Medicine (US) Committee on Resource Sharing, Berns, K. I., Bond, E. C. &
Manning, F. J. *The American Type Culture Collection*. (National Academies Press (US),
1996).

35. Funke, G., Osorio, C. R., Frei, R., Riegel, P. & Collins, M. D. Corynebacterium confusum
sp. nov., isolated from human clinical specimens. *Int. J. Syst. Bacteriol.* **48 Pt 4**, 1291–1296
(1998).

36. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11**, 119 (2010).

37. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

38. Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. E. The Phyre2 web
portal for protein modeling, prediction and analysis. *Nat. Protoc.* **10**, 845–858 (2015).

39. Edgar, R. C. MUSCLE v5 enables improved estimates of phylogenetic tree confidence by
ensemble bootstrapping. *bioRxiv* 2021.06.20.449169 (2021)
doi:10.1101/2021.06.20.449169.

40. Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34**, W609–12
(2006).

41. Lee, M. D. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* **35**, 4162–
4164 (2019).

42. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood
trees for large alignments. *PLoS One* **5**, e9490 (2010).

43. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

44. Li, W. *et al.* RefSeq: expanding the Prokaryotic Genome Annotation Pipeline reach with
protein family model curation. *Nucleic Acids Res.* **49**, D1020–D1028 (2021).

45. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9, 357–359 (2012).

46. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Res.* **47**, W256–W259 (2019).

47. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer, New York, NY, 2009).

48. Crabtree, J. *et al.* Circleator: flexible circular visualization of genome-associated data with
BioPerl and SVG. *Bioinformatics* **30**, 3125–3127 (2014).

Figure 1: A high-resolution phylogenomic view of the _C. tuberculostearicum_ species complex. A maximum-likelihood phylogeny of the _C. tuberculostearicum_ species complex, constructed from 15,936 sites predicted to not be impacted by recombination and not exhibit ambiguity in any of the 26 genomes, is shown after midpoint rooting. Bootstrap values are indicated at inner-nodes.

Figure 2: The *C. tuberculostearicum* species complex across skin metagenomic datasets and body sites. A) A phylogeny of the 187 *Corynebacterium* is shown with the *C. tuberculostearicum* species complex highlighted in gold. A single track is shown for species which were represented in the StrainGST database which we used to detect strains/species within metagenomes. Bars correspond to the number of skin-related metagenomic samples which were found to feature a species, with colors representing the study the metagenomic. **B**) The estimated relative abundance of *C. kefirresidentii* (*Ck*), *C. tuberculostearicum* (*Ct*), and *C. aurimucosum* type E (*Ca*E) is shown across Oh *et al.* 2016 and Swaney *et al.* 2022 metagenomes from select body sites in adults. Metagenomes without any of the three species detected are not shown.

Figure 3: Increase in *C. kefirresidentii* relative abundance in metagenomes sampled during atopic dermatitis flares. A) The estimated relative abundance of *Corynebacterium* species in the species complex is shown across all metagenomes from Byrd *et al.* 2017 with at least one species from the complex detected. **B)** Average coverage of *C. kefirresidentii* specific alleles along 367 single-copy-orthologs of the *C. tuberculostearicum* species is shown. Only sites/alleles which were covered by at least three reads are regarded. **C)** The ratio of the total average coverage for *C. kefirresidentii* specific alleles (inclusive of uncovered alleles) to the sum of total average coverage for *C. tuberculostearicum* specific alleles and *C. aurimucosum* type E specific alleles is shown. A pseudo-count of 1 was added to both the numerator and denominator. **D)** Genomic location of homolog groups identified as enriched in *C. kefirresidentii* genomes relative to *C. tuberculostearicum* and *C. aurimucosum* type E genomes along the chromosome of *C. kefirresidentii* FDAARGOS_1055 is shown as the outermost track in gold. The most inner two rings in black show the location of coding genes on the reverse and forward strands. The following rings (from inner to outer) show the conservation of homologs of the gene in *C. kefirresidentii* (orange), *C. tuberculostearicum* (purple), and *C. aurimucosum* type E (blue) genomes. The co-located set of genes found to be conserved in 80% of *C. kefirresidentii* featuring five molybdopterin associated enzymes is highlighted in grey and zoomed in at 10X.