Title: Planarian dorsoventral Netrins control a muscle midline signaling center and regulate blastema

formation

Authors and Affiliations:

Erik G. Schad[1,3]

Christian P. Petersen*[1,2]

1-Department of Molecular Biosciences, Northwestern University, Evanston, IL 60208

2-Robert Lurie Comprehensive Cancer Center, Northwestern University, Evanston IL 60208

3-present address: Genective, Weldon, IL 61882

*-corresponding author, christian-p-petersen@northwestern.edu

Abstract

Integration of positional information across body axes is likely critical for whole-body regeneration to define the territories of missing tissue in three dimensions with fidelity. The body-wall musculature in planarians expresses patterning factors regulating the anteroposterior, dorsoventral, and mediolateral axes, but how this information coordinates is not fully understood. We identify a previously described factor specifically expressed in dorsal midline muscle as a BMP/Activin decoy receptor *bambi-2*. Analysis of scRNAseq indicates *bambi-2+* cells coexpress midline-specifying transcription factor *pitx* and longitudinal muscle-specifying factor *myoD*, and production of *bambi-2+* cells requires these factors. In laterally amputated animals regenerating an entirely new midline, *bambi-2+* cells are initially formed at the wound site, then dynamically spread, and ultimately reset to restore bilateral symmetry. We further identify a system of dorsoventral Netrin and Netrin receptor signals expressed from body-wall muscle that control midline identity and blastema morphology. Ventral and laterally expressed *netrins -1, -4,* and *-5* signal via dorsally-enriched netrin repulsion receptors *unc5-C, unc5-E,* and *dcc-2*, which together limit mediolateral spread of *bambi-2+* dorsal midline muscle and influence the architecture of the muscle system. Our results suggest a model in which ventral determinants dictate mediolateral information important for blastema morphology.

Introduction

Whole body regeneration involves tissue restoration from a variety of possible truncations to axis content after amputation. Species of planarians, acoels, and cnidarians exemplify these abilities, as they are able to regenerate essentially entirely new animals after even severe injuries such as decapitation (Reddien, 2018; Vogg et al., 2019; Srivastava, 2022). The nature of robust positional information is therefore important to understand how extreme regenerative ability is possible. The information systems controlling regenerative ability across each body axis have often been considered separately, but surgical injuries also frequently damage more than one orthogonal body axis. How overall patterning across axes is accomplished

is incompletely understood and may be important for understanding how regenerative organisms correctly

match blastema specification to the identity and content of pre-existing tissue

Flatworm planarians can regenerate from nearly any type of surgical injury to restore axis content

along their anteroposterior (AP), dorsoventral (DV), and mediolateral (ML) axes. A system of posterior

Wnts and anterior Wnt inhibitors are expressed in gradients along the AP axis, and inhibition of canonical

Wnt signaling results in ectopic head regeneration while over-activation due to inhibition of anti-Wnt

factors results in ectopic tail regeneration (Gurley et al., 2008; Iglesias et al., 2008; Petersen and Reddien,

2008; Petersen and Reddien, 2009; Gurley et al., 2010; Petersen and Reddien, 2011; Owen et al., 2015;

Reuter et al., 2015; Stuckemann et al., 2017). In addition, two Wnt/FGFRL systems control head and trunk

regionalization in conjunction with Disheveled and Src signals (Cebria et al., 2002; Almuedo-Castillo et

al., 2011; Lander and Petersen, 2016; Scimone et al., 2016; Bonar et al., 2022). By contrast, the dorso-

ventral (DV) axis is controlled through dorsal *bmp4* signals and ventral *admp*, which signal in a negative

feedback loop (Molina et al., 2007; Orii and Watanabe, 2007; Reddien et al., 2007; Gavino and Reddien,

2011). Finally, medio-lateral axis patterning is regulated by medially expressed *slit* which opposes laterally

expressed *wnt5* (Cebria et al., 2007; Gurley et al., 2010).

The major signals specifying AP, DV, and ML pattern in planarians are expressed from

subepidermally located muscle body-wall cells that surround the inner parenchymal region containing the

neoblast planarian stem cells (Witchley et al., 2013). Neoblasts are pluripotent and generate all planarian

adult differentiated cells, thus underlying the animal's ability to replace diverse missing tissues and undergo

perpetual tissue homeostasis in the absence of injry (Wagner et al., 2011; Reddien, 2018; Zeng et al., 2018).

Planarian muscle cells are mononucleated and project contractile fibers across several orientations

(longitudinal, diagonal, circular, and dorsoventral orientations) to form a meshwork plexus (Witchley et al.,

2013; Cebria, 2016). The majority of patterning factors controlling regionalization are expressed from

multiple muscle subtypes (Scimone et al., 2017). However, analysis of cell fate transcription factors

required for producing individual muscle subtypes found that some patterning genes have expression

restricted to muscle oriented in particular directions. For example, injury triggers expression of the Wnt

inhibitor *notum* and the Activin pathway modulator *follistatin*, important for wound polarization and

outgrowth (Petersen and Reddien, 2011; Gavino et al., 2013; Roberts-Galbraith and Newmark, 2013;

Tewari et al., 2018), and inhibition of the longitudinal muscle fate transcription factor *myoD* prevents their

expression and leads to failed regeneration (Scimone et al., 2017). Likewise, circular muscle requires *nkx1.1*

for its production, selectively expresses *activin-2* involved in polarity determination (Cloutier et al., 2021),

and inhibition of *nkx1.1* leads to splitting the anterior pole and formation of split-axis animals (Scimone et

al., 2017). In addition, medial dorsoventral muscle controlled by *foxF1* and *gata4/5/6-2* factors is a source

of *slit* (Scimone et al., 2018). However, little is known about how the muscle system becomes organized

and how axis patterning systems communicate with each other to result in a restoration of form following

injury.

Prior studies have suggested the existence of communication across the body axes. The master

regulator of posterior identity *wnt1* is required for posterior regeneration and is expressed selectively in the

posterior along the dorsal midline (Petersen and Reddien, 2009; Gurley et al., 2010). *wnt1* is co-expressed

in a posterior subpopulation of dorsal midline (DM) muscle cells which express the previously unannotated

gene *dd23400* and extend across the majority of the AP axis (Fincher et al., 2018; Schad and Petersen,

2020). Overactivation of *wnt1* in *dd23400+* DM muscle cells through inhibition of *mob4* or *striatin*

components of the STRIPAK complex results in the expansion of posterior tissue in homeostasis and

regeneration, suggesting that regulation of DM muscle cells is important for body patterning (Schad and

Petersen, 2020). Additionally, the correct placement of the *notum+* anterior pole at the mediolateral and

dorsoventral midpoint is dependent on mediolateral components *wnt5* and *slit*, as well as the dorsoventral

regulator *bmp4* (Oderberg et al., 2017). These observations suggest that patterning information may

communicate across orthogonal axes and also the importance of the dorsal midline in regeneration.

However, the roles for dorsal midline muscle cells in regeneration are not fully understood.

In order to understand how midline cells contribute to regeneration, we analyzed single-cell

RNAseq data to identify factors expressed in midline muscle. This approach identified a set of transcription

factors required for DM muscle identity. We found that factors required for DM formation were also

required for lateral regeneration, suggesting the importance of these cells in facilitating outgrowth during

formation of a new mediolateral axis through regeneration. Furthermore, by tracking expression of midline

markers, we find that the re-establishment of dorsal midline during lateral regeneration proceeds from the

wound edge rather than the mediolateral center of the fragment. Using candidate approaches, analysis of

scRNAseq data, and functional analysis, we furthermore implicate Netrin related signals as regulators

restricting the domain of DM cells. Netrins are a class of secreted proteins known for their role in axonal

guidance (Hedgecock et al., 1990; Ishii et al., 1992; Kennedy et al., 1994; Serafini et al., 1994), and play a

role in axon growth and regeneration in planarians (Cebria and Newmark, 2005). Other known roles for

Netrins include cell survival (Mehlen et al., 1998; Llambi et al., 2001), organ morphogenesis (Dalvin et al.,

2003; De Breuck et al., 2003; Srinivasan et al., 2003; Yebra et al., 2003; Liu et al., 2004), and myotube

formation (Kang et al., 2004). In order to transduce their signals, Netrins bind a variety of targets including

UNC-5 (Hedgecock et al., 1990) and DCC (Chan et al., 1996; Keino-Masu et al., 1996; Kolodziej et al.,

1996; Fazeli et al., 1997). There are five genes encoding Netrins in planarians, all of which have prominent

expression in muscle cells and are regionally expressed across the ML and DV axes (Cebria and Newmark,

2005; Scimone et al., 2017; Fincher et al., 2018). In planarians, Netrin receptors (five UNC-5s, two DCCs)

are expressed widely, including in muscle cells (Fincher et al., 2018). Because Netrins and their receptors

are expressed in muscle cells regionally across the ML or DV axes like other genes controlling patterning

in planarians, we hypothesized that Netrins organize key features across these body axes, in particular, DM

muscle cells.

Results

dd23400/bambi-2+ is expressed in dorsal midline muscle

To identify factors important for DM muscle, we sought genes expressed in this cell population. Prior

single-cell RNA seq analysis identified a gene _dd23400_ expressed exclusively on the dorsal midline from

muscle cells. This gene was originally identified without discernable sequence similarity to other proteins

(Fincher et al., 2018). However, we noticed that PSI-blast of _dd23400_ recovered several homologs of a

family of Bmp/Activin membrane-bound inhibitors (BAMBI) (Figure S1) whose members possess

extracellular ActR domains, lack intracellular activation domains, and may therefore function as decoy

receptors (Onichtchouk et al., 1999). SMART domain analysis weakly identified an ActR domain in

dd23400 and also a transmembrane domain. Using maximum-likelihood phylogenies, *dd23400* placed with

strong support within a clade of bilaterian BAMBI homologs separate from BMPR1 receptors, and so we

named the gene *bambi-2*.

To examine the requirements for DM muscle cell production, we re-examined a prior whole-animal

single-cell RNAseq atlas generated by DropSeq to search for factors expressed in *bambi-2+* muscle cells

(Fincher et al., 2018)*.* Because *bambi-2* has relatively weak expression, we searched for cells co-expressing

bambi-2 and the midline factor *slit,* and compared these to bulk muscle cells using the scde package for

differential gene expression profiling from single-cell RNAseq data (Kharchenko et al., 2014). The

resulting analysis identified several genes putatively enriched for expression in *bambi2+* midline muscle

cells, including *pitx, bmp, msx, myoD, fgfr4, delta-like, TM211, nephrin, unc5-C,* and *fgfr4* (Figure 1A,

Table S1). A prior analysis using Smart-Seq to find transcriptome information for *wnt1+* cells identified

enriched expression (log fold-change > 2) in these cells for *pitx, wif, nephrin, msx, fgfr4, bmp, myoD, unc5-*

C, TM211, delta-like, and *bambi-2* (Li et al., 2019). As a further test of midline enrichment, we performed

colorimetric whole-mount in situ hybridizations on several candidates to examine whether which had

midline expression. We found midline-related expression for 11 out of 15 factors tested, with the remaining

4 lacking detection in the assay (Figure 1A, Figure S1). The transcription factor *pitx* was identified

previously as important for expression of *wnt1* (Currie and Pearson, 2013; März et al., 2013)*,* a master

regulator of posterior formation expressed from the posterior dorsal midline (Adell et al., 2009; Petersen

and Reddien, 2009; Gurley et al., 2010) and also for new expression of the midline determinant *slit* in

regeneration blastemas (Currie and Pearson, 2013; März et al., 2013). Posterior *wnt1* was subsequently

shown to be co-expressed in a posterior subset of *23400/bambi-2+* dorsal midline muscle cells (Schad and

Petersen, 2020). We used FISH to verify that *pitx* has expression along the dorsal midline (Figure 1B). We

further reasoned that other factors required for *wnt1* expression might also have expression on the dorsal

midline. The transcription factor *islet* is required for expression of *wnt1* (Hayashi et al., 2011) and although

our analysis of a DropSeq atlas did not identify enrichment for this gene in midline muscle, a prior analysis

using SmartSeq found enrichment for *islet* expression in *wnt1+* posterior midline cells (Li et al., 2019).

Using FISH, we found *islet* is expressed more broadly than *pitx,* but including dorsal midline cells in the

anterior and posterior of the animal (Figure 1B).

We observed that in addition to expression on the dorsal midline in the posterior, *islet* and *pitx* are

also expressed on the DM in the anterior (Figure 1B), so we reasoned that they could be required for all

bambi-2+ cells. We performed RNAi of *islet* and *pitx,* and found a nearly complete loss of *bambi-2*

expression (Figure 1C). *bambi-2+* DM muscle cells may additionally be specified by transcription factors

required for other muscle cell subsets. The *myoD* transcription factor is required for longitudinal muscle

throughout the body (Scimone et al., 2017), was recovered by scRNAseq analysis as enriched in *bambi-2+*

midline cells, and its expression domain includes a midline-enriched region expression pattern (Figure 1A).

Therefore, we tested whether *myoD* inhibition would impact *bambi-2* dorsal muscle expression. RNAi of

myoD, but not the *nkx1.1* transcription factor required for circular muscle, caused a loss of *bambi-2*

expression, suggesting that *bambi-2+* DM muscle cells are likely a subset of longitudinal muscle (Figure

1B). Because *pitx* and *islet* are known to be required for expression of *slit* (Currie and Pearson, 2013; März

et al., 2013), these results argue these factors likely control midline muscle cell identity in general. We

tested other markers of midline to confirm these findings. *dachsous* has midline expression (Vu et al.,

2019) and was enriched for expression in midline muscle by the scde analysis (Table S1, log2 upper-bound

enrichment of 3.10). *pitx* or *islet* RNAi resulted in a loss of *dachsous* expression (Figure 1D). Furthermore,

a homolog of *Wnt inhibitory factor (wif)* (Hsieh et al., 1999), itself a homolog of Drosophila *shifted* (Glise

et al., 2005)*,* was recovered as strongly enriched in *bambi-2+* muscle cells from the single-cell RNAseq

analysis (Figure 1A, log2 upper-bound enrichment of 10.92). FISH analysis revealed *wif* to be expressed

weakly along the dorsal midline, and inhibition of *pitx* or *islet* similarly reduced *wif* expression. Therefore,

pitx and *islet* are important for expression of several markers of dorsal midline muscle. We also used the

additional markers of midline muscle to examine the consequences of *bambi-2* inhibition on these cells.

bambi-2(RNAi) animals appeared phenotypically normal and did not fail to express midline *dachsous* or

wif (Figure 1C-D). Additionally, co-inhibition of another previously reported BAMBI homolog, *bambi ,*

also resulted in animals with normal appearance. Together, *bambi-2* is a marker of dorsal midline but likely

not critical for its formation.

The high specificity of expression of *bambi-2* for the dorsal midline enabled the opportunity to

examine the process of midline re-establishment during lateral regeneration. The expression of *bmp4* has

been used to examine midline formation during lateral regeneration previously. In *Schmidtea mediterranea*,

bmp4 is expressed both on the dorsal midline as well as laterally on the dorsal side in a graded manner

(Reddien et al., 2007), and in *Dugesia japonica, bmp* expression is prominent on the midline and also

detectable in a graded fasion from the dorsal midline (Orii, 1998). In *Dugesia japonica* animals undergoing

lateral regeneration, *bmp4* expression punctate and broad at 1 day post amputation, which could either

reflect pre-existing or new expression of *bmp4* (Orii, 1998). By 3 days of regeneration, expression primarily

appeared as a line running anteroposteriorly and located asymmetrically on the fragment and closer to side

forming a regeneration blastema. As regeneration proceeded, the most abundant *bmp4* signal still appeared

to be asymmetric in the animal but located closer to the left-right midpoint of the fragment. Similar

asymmetry in *bmp4* expression early in regeneration has been reported in *Schmidtea mediterranea* with

some differences. In lateral regeneration in that organism, new expression of *bmp4* was detected at 12 hours

in a punctate pattern close to the wound site (Reddien et al., 2007). By 48 hours, expression was stronger

on the blastema side of the fragment but without yet a discernable midline component, and midline

expression could be detected by 14 days. Other tests of *bmp4* expression in lateral regeneration have

detected greater expression early in regeneration (2-4 days) on the blastema side of fragments (Gavino and

Reddien, 2011). Together, these prior observations suggest midline formation is unlikely to proceed through

a process that places new midline at the geometric midpoint. However, because *bmp* expression is graded

and includes non-medial dorsal cells, the exact nature of midline regeneration is still not fully understood.

We tested the expression of *bambi-2* during a timecourse of lateral regeneration in asymmetrically

laterally amputated fragments that unambiguously initially lacked midline expression. *bambi-2* expression

appeared by 24 hours at the wound edge at a location furthest from the existing lateral-most domain. *Bambi-*

2 expression retained dorsal specificity throughout the timecourse, so this position would effectively mark

the dorsal-most position in the fragment remaining following the amputation (Figure 2). Over time, this

expression domain expanded while retaining dorsal specificity (100% of animals only had expression on

one DV side of the animal, N at least 8 animals per timepoint), reached a point of largest mediolateral

spread by 72 hours, followed by restriction to a location between the pre-existing tissue and growing

blastema. Consistent with prior studies, midline expression had not become fully restored to the middle of

the fragment by 7 days. These results support that dorsal midline regeneration proceeds initially in close

association with the wound site, followed by subsequent re-establishment.

Netrins from ventral/medial muscle domains coordinate to instruct blastema outgrowth shape

We hypothesized from these results that dorsoventral signals could be important for establishing

the *bambi-2+* expression domain. We noticed that one of the genes enriched in midline muscle expression

from the scRNAseq analysis (*unc5*-C) belonged to the Netrin signaling pathway, suggesting a possible role

for this pathway in DM muscle function. Netrins are secreted proteins that provide guidance cues for

migratory cells (Hedgecock et al., 1990; Ishii et al., 1992) and can signal to receiving cells across long

ranges (Colavita and Culotti, 1998; Hong et al., 1999; Keleman and Dickson, 2001; Finci et al., 2014; Boyer

and Gupton, 2018). The planarian genome contains five genes encoding Netrin proteins, each expressed in

muscle cells (Figure 3A-C, Figure S2) (Fincher et al., 2018), while *netrin-1* and *netrin-2* are also expressed

in neurons (Cebria and Newmark, 2005; Scimone et al., 2017). Planarian Netrins are expressed in a series

of ML and DV domains (Figure 3A). *netrin-1* and *netrin-3* are coexpressed in *collagen+* muscle cells on

the lateral edge of the animal, while *netrin-2* and *netrin-5* are expressed in medial muscle cells on the ventral

side (Figure 3A). *netrin-4* is expressed across the ventral side of the animal (Figure 3A). Sections of animals

co-stained with each Netrin and the muscle marker *collagen* showed that Netrins are expressed in several

ventral/medial domains within muscle. Because Netrins function in cell guidance and have graded

expression in muscle cells on the ML axis, we hypothesized that they play a role in organizing the DM

muscle domain. The planarian genome encodes several Netrin receptors: five putative UNC-5 receptors,

and two DCC receptors (Scimone et al., 2020). Each of these receptors is expressed broadly, but also has

graded expression in muscle cells (Fincher et al., 2018) (Figure 3B-C, Figure S2). *unc5-A,-B,-D* and *dcc-*

1/net-R are expressed on the lateral edge of the animal, while *unc5-C,-E* and *dcc-2* have prominent

expression in dorsal medial *collagen+* muscle cells (Figure 3B-C). In summary, Netrins are expressed in

either lateral (*netrin-1,-3),* ventral-medial (*netrin 2,-5*), or broad ventral (*netrin-4*) domains, while putative

receptors are expressed in lateral (*unc5-A,-B,-D* and *dcc-1/net-R*) or dorso-medial (*unc5-C,-E* and *dcc-2*)

domains (Figure 3D).

To investigate the role of Netrins during regeneration we inhibited each Netrin using RNAi. We

first inhibited each Netrin individually, resulting in no overt phenotypes of dysfunction in either anterior or

posterior regeneration (Figure 4A). We reasoned that Netrins could function redundantly, and so we

inhibited all five planarian Netrins through combined RNAi, and verified by qPCR for effective knockdown

(Figure S3). Inhibition of all Netrins simultaneously resulted in regeneration with midline defects of a

pointy blastema in the anterior (32/36 regenerating trunk fragments), and of indented blastemas in the

posterior (7/36) (Figure 4B). In addition, at a low frequency, inhibition of the five Netrins resulted in failed

anterior regeneration in amputated tail fragments (6/36) or failed posterior regeneration from amputated

head fragments (5/36 animals). We reasoned that these phenotypes were either due to inhibition of all

Netrins together or instead due to inhibition of subsets of Netrins. To test this hypothesis, we inhibited each

possible combination of two, three, or four Netrins, and scored animals for the penetrance and expressivity

of pointy blastema in anterior blastemas or indentations in posterior blastemas (Figure S4A-B). We found

that inhibition of all combinations of Netrins resulted in anterior pointy blastemas to varying degrees.

Inhibition conditions including RNAi of *netrin-4,-5* yielded the strongest effects on pointy blastemas,

followed by those including *netrin-1* and *netrin-2*. The laterally expressed *netrin-3* factor likely did not

contribute, because *netrin-1;-2;-3;-4;-5* RNAi yielded the same outcome as *netrin-1;-2;-4;-5* RNAi (Figure

S4A). Therefore, *netrin-1, -2 ,-4,* and *-5* contribute to normal anterior regeneration morphology.

A similar analysis of the indented blastema phenotype from posterior-facing wounds found a

somewhat distinct outcome. Whereas inhibition of all five Netrins simultaneously resulted in an indented

blastema, inhibition of most combinations of Netrins resulted in normal posterior regeneration. Inhibition

of *netrin-1, -4,* and *-5* increased the penetrance and expressivity of indentation over that from inhibition of

all netrins, indicating a particular importance for these three genes in enabling midline regeneration in the

posterior (Figure S4B). Consistent with this model, RNAi combinations excluding inhibition of any of these

three Netrins resulted in normal posterior regeneration, and co-inhibitions with *netrin-2* or *netrin-3* did not

strongly affect the outcome. Therefore, *netrin-1*, *netrin-4*, and *netrin-5* act together to ensure midline

regeneration in blastemas generating posterior tissue.

We also examined whether the low frequency of failed regeneration after inhibition of all Netrins

simultaneously could be explained by lack of injury-induced gene expression from muscle cells. However,

netrin-1;-2;-3;-4;-5(RNAi) animals had normal injury-induced expression of *notum* and *follistatin* (Figure

S5). We suggest that the failed regeneration phenotype could either reflect a distinct role for Netrins or a

more extreme version of the defects causing impaired blastema morphology. Nonetheless, these results

suggest that Netrin inhibition does not result in failed responsiveness of longitudinal muscle cells to injury

(Scimone et al., 2017). These results also suggest that the indented morphology phenotype is not likely due

to a problem with early expression of injury-induced *notum* or *wnt1*.

To further investigate the role of Netrin signaling in regeneration, we conducted RNAi experiments

to test for functions of the Netrin receptors. Inhibition of the receptors expressed in lateral domains (*unc5-*

A,-B,-D and *dcc-1/net-R*) appeared to result in normal regeneration, including when inhibited in

combination (Figure 4C). However, inhibition of either *unc5-C, unc5-E*, or *dcc-2* caused regeneration with

indentations to posterior blastemas in all cases, and indentations to anterior regeneration after *dcc-2* RNAi.

Dual inhibition of *unc5-E* and *unc5-C* or triple inhibition of *unc5-E;unc5-C;dcc-2* resulted in more strongly

expressive anterior and posterior blastema indentations. The indented posterior blastemas were reminiscent

of the posterior blastema morphology phenotypes after RNAi of *netrins-1, -4,* and *-5*, suggesting these

Netrins may signal through *unc5-E/unc5-C/dcc-2* receptors. Furthermore, while *netrin-1, netrin-4* and

netrin-5 have expression in ventral and lateral muscle domains, *unc5-E* and *unc5-C* have enrichment in a

dorsal domain (Figure 3). These observations implicate a system of Netrin ligands and receptors signaling

across muscle domains in to control midline blastema morphology.

<u>Netrins act as repulsive signals to restrict the midline domain</u>

Based on the overt mediolateral defects in blastema formation after Netrin RNAi, we tested the

expression of ML and DV patterning factors in these animals. *netrin-1;-2;-3;-4;-5(RNAi)* animals displayed

an increase in *bambi-2* domain width, especially in the anterior (Figure 5A). Similarly, midline expression

of *slit* was also expanded in *netrin-1;-2;-3;-4;-5(RNAi)* animals, while laterally expressed *wnt5* was

unchanged (Figure 5A-C). Furthermore, *netrin-1;-2;-3;-4;-5(RNAi)* animals retained expression of dorsal

bmp4 and ventral expression of *admp*, suggesting these animals did not undergo ventralization or

dorsalization (Figure 5A). Finally, *netrin-1;-2;-3;-4;-5(RNAi)* animals succeeded in expressing *wnt1+* at

the posterior pole and *notum* at the anterior pole, consistent with animals regenerating with normal head

and tail identity in spite of the blastema morphology defects (Figure 5A). We noted that although *notum*

expression was correctly located at the anterior pole, its expression was expanded over the Netrin(RNAi)

pointy blastema compared to controls. Prior studies suggest that a preexisting midline can determine the

location of a newly regenerated anterior pole (Oderberg et al., 2017), so perhaps the expanded midline

domain in *netrin-1;-2;-3;-4;-5(RNAi)* animals may cause formation of an expanded anterior pole. Together,

these findings implicate Netrins specifically in restricting dorsal midline identity. Based on these findings,

we specifically tested for any midline defects after inhibition of the Netrin receptors involved in midline

formation. Simultaneous RNAi of *unc5-C, unc5-E,* and *dcc-2* resulted in a similar expansion of the *bambi-*

2+ cell domain (Figure 3B-C). Therefore, the phenotypes of dorsal midline expansion correlate with

indented blastemas. Reception of Netrins through receptor heterodimers of Unc5 and DCC family members

are thought to signal repulsion during axon guidance and cell migration (Lai Wing Sun et al., 2011).

Furthermore, the *bambi-2+* dorsal midline cells are located at the point furthest away from the sources of

ventral and medial *netrin-1, netrin -4* and *netrin-5* but within the dorsal domains of *unc5-E* and *unc5-C*

expression. Therefore, we suggest that the *bambi-2+* cells are restricted dorsally because of repulsive

signaling between ventral/lateral *netrin-1,-4,-5* to dorsally expressed Unc5 receptors *unc5-E* and *unc5-C*

signaling through *dcc-2*.

The expression of Netrins within muscle cells led us to hypothesize that Netrins might control some

aspect of organization of the muscle fiber network important for blastema formation in regeneration. To

test this hypothesis, we inhibited all Netrins or Netrin receptors *unc5-E, unc5-C,* and *dcc-2*, and then used

immunofluorescence to label animals with a monoclonal antibody (6G10) that labels muscle cell fibers

(Ross et al., 2015) (Figure 5D). In *netrin-1,-2,-3,-4,-5(RNAi)* animals, muscle cell fibers were present in the

pointed anterior blastemas and appeared normal, although they appeared to track with the overall pointed

morphology. In *netrin-1,-2,-3,-4,-5(RNAi)* animals, some circular muscle fibers near the indented posterior

regenerated tissue inappropriately projected anteroposteriorly after crossing the midline. *unc5-E;unc5-*

C;dcc-2 RNAi led to a more severe defect of indentation, and we also noticed such animals have a zone of

muscle disorganization immediately anterior to the indentation along the midline, which could reflect a

muscle remodeling process linked to indented tissue. We conclude that a system of ventrally expressed

Netrins and dorsally expressed Netrin Receptors are likely not required for muscle fiber growth altogether,

but impact the architecture of the muscle system.

Discussion

Whole-body regeneration requires the re-establishment of patterning across all major body axes.

Here, we identify processes important for generating, restricting, and using *bambi-2+* dorsal midline muscle

in planarian regeneration. Inhibition of factors *pitx* and *islet* caused loss of midline expression of *bambi-2,*

dachsous, and *wif* and also prevented blastema outgrowth during lateral regeneration, suggesting the

importance of these cells in restoring mediolateral tissue distribution. We furthermore used *bambi-2*

expression as a probe to uncover the overall process of midline regeneration from fragments lacking a

midline. *bambi-2* expression initiates at the lateral wound edge. Subsequently, *bambi-2* expression expands

then appears to restrict to a region between new and old tissue, and differential growth over several weeks

ultimately restores bilateral symmetry (Figure 6A).

In addition, we describe the essential role of Netrin signaling in ML patterning during regeneration

in planarians (Figure 6B). *netrin-1,-2,-3,-4,-5* inhibition lead to an expansion of the *bambi-2+* cell domain,

but did not alter overall DV or AP identities. Netrins are known for their role in axonal growth (Hedgecock

et al., 1990; Ishii et al., 1992; Hamelin et al., 1993; Colavita and Culotti, 1998), including in planarians

(Cebria and Newmark, 2005), but Netrins can also guide myotube formation and other migrating cells in

murine development (Dalvin et al., 2003; De Breuck et al., 2003; Srinivasan et al., 2003; Yebra et al., 2003;

Kang et al., 2004; Liu et al., 2004), so it is plausible that Netrins may function to guide muscle cell growth

in planarians. Three putative Netrin receptors (*unc5-C, -E, dcc-2*) are expressed in dorso-medial muscle

cells, and their inhibition leads to an expansion of the *bambi-2+* cell domain. These results are consistent

with a model in which ventral Netrins act as repellants by binding to these receptors (Figure 6B). When

netrin-1,-2,-3,-4,-5 are inhibited together, anterior blastemas form with a pointed morphology, with

expanded *notum+* anterior pole cells. Prior work determined that the midline instructs where the anterior

pole is placed in the anterior blastema (Oderberg et al., 2017), suggesting that the expanded dorsal muscle

domain from Netrin RNAi conditions may cause the regeneration of an excessively wide anterior pole, thus

generating pointed blastemas. Alternatively, it is possible that the pointy blastema phenotype represents a

loss of other outputs for Netrin activity. For example, Netrins can function as attractants in axon guidance

or migration, and this function is thought to occur via DCC/DCC receptor dimers rather than DCC/Unc5

dimers (Lai Wing Sun et al., 2011). In our survey of Netrin receptor RNAi phenotypes, no condition

phenocopied the pointy blastema defect observed from Netrin RNAi. One possible explanation could be

that due to the shared use of DCC in both repulsion and attraction, it is not possible to use RNAi of these

factors to selectively disrupt the process of attraction. Alternatively, perhaps other receptors are the targets

of Netrin signaling disrupted during the acquisition of the pointy blastema phenotype. Along these lines,

we also did not find blastema failure phenotypes through inhibition of Netrin receptors. However, our

phenotypic analysis identified shared functions in controlling the regeneration of midline tissue in the

posterior as well as in muscle targeting, for a subset of Netrins and Netrin receptors. RNAi of *netrin-1;-4;-*

5 or *unc5-C;unc5-E;dcc-2* impacted the organization of the planarian musculature system, causing the

newly made circular muscle in regenerating blastemas to mistarget over the midline and instead forming

separated domains that grow into indented blastemas.

Whole body regeneration in planarians depends on axis information systems encoded in muscle

cells. Our study finds muscle-expressed ligands signaling through muscle-expressed receptors to control

regionalization of muscle identity. These observations suggest that regionalization signaling could take

place directly between muscle cells. Alternatively, given that *dcc-2* and other netrin receptors are expressed

more broadly, an alternative possible mechanism for pattern formation could involve the use of positional

cues from muscle to influence regionalization through control of stem cell differentiation into distinct

muscle domains. Netrins have dorsoventral expression domains opposing BMP in multiple organisms,

including in planarians (Cebria et al., 2007; Reddien et al., 2007), hemichordates (Lowe et al., 2006), acoels

(Srivastava et al., 2014), Drosophila (Harris et al., 1996; Mitchell et al., 1996), and within the vertebrate

neural tube (Serafini et al., 1994; Lai Wing Sun et al., 2011). In addition, Netrins have expression in body

wall muscle in planarians (Witchley et al., 2013), acoels (Raz et al., 2017), *Drosophila*, and *C. elegans*

(Ishii et al., 1992). Our observations point to an ancient and conserved use of Netrins for linking

dorsoventral and midline identity through muscle. The involvement of multiple Netrins and receptors for

controlling dorsal midline muscle and blastema morphology suggests the possibility of a dorsoventral code

of these factors that guide patterning of the blastema in order to restore bilateral symmetry during whole-

body regeneration.

Acknowledgements

We thank members of the Petersen lab for critical comments. CP acknowledges funding from NIGMS

R01GM129339, R01GM130835, and pilot project funding from the NSF-Simons Center for Quantitative

Biology at Northwestern University, an NSF (1764421)-Simons/SFARI (597491-RWC) MathBioSys

Research Center.

Author contributions: CP conceived of the study and acquired funding, ES designed and conducted experiments, CP and ES wrote the manuscript.

Declaration of Interests: There are no competing interests.

Figure Legends

Figure 1. *pitx, islet,* and *myoD* are required for expression of *bambi-2*.

(A) Analysis of prior scRNAseq data (Fincher et al 2018) comparing transcriptomes of dorsal muscle midline cells co-expressing *bambi-2(dd23400)* and *slit* versus *collagen+* muscle lacking *bambi-2* and *slit* expression. Table shows log2-fold change (upper bound, ub) from scde analysis. Right panels show WISH of candidates with expression on the posterior midline. Images represent 3-4 animals per condition. **(B)** Animals were fixed and FISH was performed to detect *islet* or *pitx* expression on the dorsal midline in the anterior (left) or posterior (right). Arrows indicate expression on the dorsal midline. **(C)** Homeostatic RNAi (8 feedings dsRNA) to inhibit transcription factors specific for midline cells (*islet, pitx*), L-muscle (*myoD*), and C-muscle (*nkx1-1*) followed by FISH to detect *bambi-2*. **(D)** FISH to detect expression of *dachsous* and *wif* midline markers after *pitx*, *islet* or *bambi-2* RNAi**.** Bars, 100 (B) or 200 microns (A,C).

Figure 2. Lateral regeneration progressively restores *bambi-2* midline expression through injury-induced expression and growth.

FISH of *bambi-2* expression during a timeseries of the regeneration of asymmetric lateral fragments that must regenerate the dorsal midline. Bottom, quantifications of the relative and absolute locations of the *bambi-2+* cells with respect to the wound edge (left), the relative size of the *bambi-2+* domain and size of pre-existing tissue domain lateral to the *bambi-2* domain (center), and absolute width of the regenerating fragments. *bambi-2* expression first arises at the lateral wound edge by 24 hours, followed by a progressive relative movement toward the left/right center of the fragment, rather than a mechanism that directly places these cells centrally. n=12-15 animals. * p<0.05, **p<0.01, ***p<0.001 by two-tailed unpaired t-tests.

Figure 3. Netrins and Netrin receptors are expressed in dorsoventral domains in muscle cells.

(A) FISH images showing *netrins* are expressed in along either ventromedial (*netrin-2, netrin-4, netrin-5)* or lateral (*netrin-1, netrin-3)* domains. Each image shows a ventral view. N=5-6 animals. **(B)** FISH images showing Netrin receptors are expressed in along either dorsomedial (*unc5-C, unc5-E, dcc-2)* or lateral (*unc5-A, unc5-B, unc5-D, dcc-1)* domains (4-7 animals probed per condition). (C) Images of cross-sections, dorsal side top. FISH images showing *netrins* are expressed in along either ventromedial (*netrin-2, netrin-4, netrin-5)* or lateral (*netrin-1, netrin-3)* domains (4-6 animals probed per condition) FISH images showing Netrin receptors are expressed in along either dorsomedial (*unc5-C, unc5-E, dcc-2)* or lateral (*unc5-A, unc5-B, unc5-D, dcc-1)* domains. Bars, 100 (A,B) or 50 microns (C). Arrows indicate expression of each tested gene coincident with *collagen+* muscle. (D) Cartoon of planarian cross-section displaying the approximate expression domains for each Netrin and Netrin receptor.

Figure 4. Netrins act cooperatively to control blastema morphology and regeneration.

(A) Live images showing RNAi of individual *netrins* results in normal regeneration. (10 animals per condition). **(B)** Simultaneous inhibition of all five Netrins causes regeneration of blastemas with an indentation in the posterior and pointed morphology in the anterior, as well as weakly penetrant

regeneration failure in head and tail fragments (arrows). **(C)** RNAi of *unc5-C, unc5-E* and *dcc-2*

individually or in combination results in indented blastemas (arrows). Bars, 200 microns.

Figure 5. Netrins control midline muscle cell organization and proper muscle morphology.

(A) FISH showing expression of patterning factors following inhibition of each Netrin. Midline (*bambi-2,*

slit) and anterior pole cells (*notum)* are expanded, while lateral (*wnt5*), posterior (*wnt1*), and dorsoventral

markers (*bmp-4, admp-1*) appear normal. *admp-* and *notum*-stained animals are from regenerating tail

fragments, all other panels show regenerating trunk fragments. Insets show enlargements of *bambi-2* and

slit expression **(B)** FISH showing expanded *bambi-2* expression domain following *unc5-C,-E,dcc-2* RNAi.

Insets show enlargements of *bambi-2* expression. **(C)** Quantification of midline expansion normalized to

body width in each condition. Bars show standard error of 5-10 replicates, *** $p < 0.001$ by 2-tailed unpaired

t-test. **(D)** Immunostaining with 6G10 antibody to visualize muscle fiber morphology after *netrin-1;-2;-3;-*

4;-5 RNAi and *unc5-C;unc5-E;dcc-2* RNAi in anterior (top) or posterior (bottom) blastemas. Muscle

morphology was disrupted at locations of the indented blastema. Animals were fixed after 8 dsRNA

feedings and 14 days of regeneration in all panels. Bars, 200 (A-B) or 100 microns (D).

Figure 6. Model

(A) In regeneration of lateral fragments, the *bambi-2+* dorsal midline muscle is re-established near the

wound site and not the geometric center of the fragment. *pitx* and *islet* are necessary both for *bambi-2*

expression and blastema outgrowth that restores bilateral symmetry (B) Three ventral netrins likely act

through dorsal netrin receptors to restrict the location of the *bambi-2+* cells.

464   **Supplemental Figures**

Figure S1. dd23400/*bambi-2* encodes a BAMBI decoy receptor

(A) Sequence alignment of dd23400 with BAMBI factors with conserved residues highlighted in red. Green bar indicates location of ActR domain. dd23400 has a weakly detected ActR domain (aa4-84, evalue 0.14), possesses a transmembrane domain (aa158-180, TMHMMv2.0) and like other BAMBI homologs lacks an intracellular kinase activation domain. DELTA Blast of dd23400 followed by PSI-blast (3 rounds, 500 iterations) recovers BAMBI orthologs from several species, including BAMBI from Osmia bicornis (evalue=1e-22), Megachile rotundata (evalue=3e-22) and Apis mellifera (evalue=3e-20).**(B)** Bayesian inference phylogenetic tree generated from alignment with MUSCLE and calculated with MrBayes v3.2.3 using a GTR substitution model with gamma-distributed rate variation across sites and a proportion of invariable sites. Runs of 50000 generations converged with an average standard deviation of split frequencies <0.01 and 25% of trees were discarded as burn-in. Tree displayed using FigTree v1.4.4 with branch labels indicating posterior probabilities. BAMBI sequences from Owenia fusiformis (Of, CAC9588793.1), Octopus bimaculoides (Ob, XP_014790456.1), Argiope bruennichi (Ab, KAF8794778.1), Araneus ventricosus (Av, GBL98371.1), Stegodyphus mimosarum (Sm, KFM81221.1), Xenopus laevis (Xl, CAB56858.1), Mus musculus (Mm, NP_036474.1), and Branchiostoma floridae (Bf, XP_035688850.1).Tree is rooted using BMPR1 receptors from Brachiostoma floridae (Bf), Danio rerio (Dr), and Mus musculus (Mm).

Figure S2. Expression of netrins and netrin receptors from cell atlas.

Plots of Netrin and Netrin receptor gene expression from a planarian cell atlas (Fincher et al., 2018 ; digiworm.wi.mit.edu). Top cartoon shows cluster identities. Netrins and receptors have expression in muscle cells and also in other cell types.

Figure S3. qPCR validating knockdown.

dsRNAs targeting netrins -1, -2, -3, -4, and -5 were mixed together and delivered to animals by feeding 6 times over 2 weeks ("pan-net") and compared to animals treated with control dsRNA. RNA was harvested

and qPCR conducted to measure knockdown of each Netrin factor as indicated, using *clathrin* transcript as

a normalizing control. Bars show standard error of 4 replicates, ***P<0.001 by 2-tailed unpaired t-test.

Figure S4. Netrins -1, -2, -4, and -5 promote proper anterior blastema morphology while *net-2, -4,*

***and -5* promote posterior blastema morphology.**

(A) Graph showing regeneration score of anterior blastemas for all combinations of Netrin RNAi. n=7-10

animals. Animals were qualitatively scored based on the degree of pointedness from P0 (normal), P1

(subtle), P2 (pointy), to P3 (highly pointed). In order to compare different conditions with varying

penetrance and expressivity, a weighted average phenotype strength score was computed for each treatment

by summing the numbers of animals scored in each category multiplied by a scalar of increasing size for

higher expressivity (i.e., score = [(# animals in P1) + 2 x (# animals in P2) +3 x (# animals in P3)]/(total

number scored)). Average phenotype strength scores were normalized to the value from inhibition of all

five Netrins for comparison. Right- Representative images of degrees of blastema pointedness in different

combinations of Netrin inhibitions. (B) Graph showing percentage of indented posterior blastemas for all

combinations of Netrin RNAi. n=7-10 animals. Right- Representative images of degrees of indented-ness

in different combinations of Netrin inhibitions. (C) Live images showing *netrin-1;-2;-3;-4;-5* are not

required for blastema formation in regenerating fragments lacking midline.

Figure S5. *netrins* are not required for wound-induced expression of *notum* and *follistatin.*

(A) Animals were fixed following 8 dsRNA feedings and fixed either 6 or 24 hours after injury. FISH

showing *notum* or *fst* expression is unchanged in All Netrin RNAi. n=>10 animals. In panel A bars are 200

microns.

Materials and Methods

Experimental Model

Asexual *Schmidtea mediterranea* animals (CIW4 strain) were maintained in 1x Montjuic salts

between 18–20°C. Animals were fed a puree of beef liver and starved for at least 7 days before experiments.

In situ hybridizations

Animals were fixed and bleached as described previously (Pearson et al., 2009; King and Newmark, 2013).

Riboprobes (digoxigenin- or fluorescein-labeled) were synthesized by in vitro transcription (Pearson et al.,

2009; King and Newmark, 2013). Antibodies were used in TNTx/10% horse serum

(Roche, Basel Switzerland) at a concentration of 1:2000 for anti-DIG-POD (Roche, Basel Switzerland),

and 1:2000 for anti-FL- (Roche), or 1:4000 for anti-dig-alkaline phosphatase used in colorimetric in situ

hybridizations (Roche, Basel Switzerland). For multiplex FISH, peroxidase conjugated enzyme activity

was quenched between tyramide reactions by sodium azide treatment (100 mM in 1xTNTx) for 45 min at

room temperature. Nuclear counterstaining was performed using Hoechst 33342 (Invitrogen, 1:1000 in

1xPBSTx for 45 minutes). Colorimetric in situ hybridizations were developed using NBT/BCIP as

described previously (Lander and Petersen, 2016). Primer sequences used for cloning to generate riboprobes

are listed in Table S2.

Immunostaining

Animals were fixed the same as in FISH (King and Newmark, 2013). Bleaching was performed in

80% methanol, 20% hydrogen peroxide on a lightbox overnight. Following rehydration washes, animals

were washed and rocked in PBSTx/10% horse serum (Roche, Basel Switzerland) 8 times for 6 hours.

Animals were incubated in the anti-muscle mouse monoclonal antibody 6G10-2C7 (Iowa Hybridoma Bank)

at 1:1000 overnight, then washed in PBSTx/10% horse serum 8 times for 6 hours. Animals were incubated

overnight in an anti-mouse Dig conjugated antibody in PBSTx/10% horse serum. After 8 PBSTx washes,

tyramide amplification was performed using commercial tyramide kit. Finally, animals were washed in

PBSTx 8 times, 20 minutes each.

RNAi administration

RNAi by feeding was performed in vitro transcribed dsRNA mixed with liver paste and 5% red

food dye (Rouhana et al., 2013). For single RNAi experiments, dsRNA at a concentration of 2-4 ug/ul was

mixed with liver in a 1:5 ratio. In RNAi experiments with multiple dsRNAs, each dsRNA was mixed with

liver for a total concentration of 1:10 for each dsRNA. In all RNAi experiments animals were fed RNAi

food every 2-4 days for the length of experiment indicated. For homeostatic experiments, animals were

starved for at least 5 days before fixing. In regeneration experiments, animals were amputated the same day

of the last feeding, or in Figure 1C, 1D, Figure S4 at 5 days after the final RNAi feeding. dsRNA targeting

the *Photinus pyralis* luciferase (not present in the planarian genome) was used as a negative control RNAi

condition. Primers sequences used for RNAi are listed in Table S2.

Image Acquisition and Quantification

Live images of animals were performed with a Leica M210F dissecting scope with a Leica DFC295 camera.

FISH images were performed a Leica DM5500B compound microscope with optical sectioning by Optigrid

structured illumination or a Leica Stellaris or Leica TCS SPE confocal compound microscopes. Fluorescent

images collected by compound microscopy are maximum projections of a z-stack and adjusted for

brightness and contrast using Metamorph, Adobe Photoshop, ImageJ, or Leica LAS.

qPCR

RNA was harvested from 4 biological replicates of 4 animals per condition using a Turrex tissue

homogenizer. cDNAs were synthesized using oligo-dT and Superscript III reverse transcriptase (Life

Technologies) for 1 hour at 50°C. EvaGreen qPCR mastermix (Midwest Scientific BEQPCR-S) was used

for amplification and detection. Per-sample log2-fold change expression were determined by the delta-

delta Ct method, normalizing to expression of housekeeping gene clathrin probe set control and to RNAi

negative controls targeting *Photinus pyralis* luciferase (not present in planarian genome). Statistical

significance was determined through unpaired two-tailed t-tests comparing ddCt values between control

RNAi and target RNAi conditions. Primers for qPCR: *netrin-1* (5'-taaagagcatcagggtctatgtga -3', 5'-

ataatatcgacgcttcaatttcca-3'); *netrin-2* (5'-gaattggtatgacaagacaactgc-3', 5'-gttttgcgtcgcatatactacaat-3');

netrin-3 (5'-cgaaaccctatgtaaaaagaagga-3', 5'-cccttggagattttataggacaga-3') ; *netrin-4* (5'-

cctgttaaattgaaatcgctctct-3', 5'-aataaaataatttggggatgctga-3') ; *netrin-5* (5'-tgcacaaaatgctacatttgaagt-3', 5'-

atcctttctgaccgcatgtact-3'); *clathrin* (5'-agccgcatgtagttggttagtg-3', 5'-agtctacagaaaaccatcaattgtttcaaa-3').

scRNAseq differential gene expression testing

We mined a prior single-cell RNAseq atlas generated by DropSeq (Fincher et al., 2018) and recovered 15

dd23400+slit+ cells (at least 1 read for *bambi-2/dd23400* and at least 2 reads for *slit*) and compared these

to 143 randomly selected muscle cells (at least 7 reads for muscle marker *collagen* dd840 and no reads for

dd23400 or *slit*) and analyzed for differential gene expression using the Single Cell Differential Gene

Expression (scde) method (Kharchenko et al., 2014). Table S1 shows results of differential testing along

with identity of cells selected for analysis.

Phylogenetic analysis

Bayesian inference phylogenetic tree was generated from alignment with MUSCLE and calculated with

MrBayes v3.2.3 using a GTR substitution model with gamma-distributed rate variation across sites and a

proportion of invariable sites. Runs of 50000 generations converged with an average standard deviation of

split frequencies <0.01 and 25% of trees were discarded as burn-in.

591   **References**
592

ADELL, T., SALÒ, E., BOUTROS, M. & BARTSCHERER, K. 2009. Smed-Evi/Wntless is required for β-catenin-dependent and-independent processes during planarian regeneration. *Development,* 136**,** 905-910.

ALMUEDO-CASTILLO, M., SALÓ, E. & ADELL, T. 2011. Dishevelled is essential for neural connectivity and planar cell polarity in planarians. *Proceedings of the National Academy of Sciences,* 108**,** 2813-2818.

BONAR, N. A., GITTIN, D. I. & PETERSEN, C. P. 2022. Src acts with WNT/FGFRL signaling to pattern the planarian anteroposterior axis. *Development,* 149.

BOYER, N. P. & GUPTON, S. L. 2018. Revisiting Netrin-1: One Who Guides (Axons). *Front Cell Neurosci,* 12**,** 221.

CEBRIA, F. 2016. Planarian Body-Wall Muscle: Regeneration and Function beyond a Simple Skeletal Support. *Front Cell Dev Biol,* 4**,** 8.

CEBRIA, F., GUO, T., JOPEK, J. & NEWMARK, P. A. 2007. Regeneration and maintenance of the planarian midline is regulated by a slit orthologue. *Dev Biol,* 307**,** 394-406.

CEBRIA, F., KOBAYASHI, C., UMESONO, Y., NAKAZAWA, M., MINETA, K., IKEO, K., GOJOBORI, T., ITOH, M., TAIRA, M., SANCHEZ ALVARADO, A. & AGATA, K. 2002. FGFR-related gene nou-darake restricts brain tissues to the head region of planarians. *Nature,* 419**,** 620-4.

CEBRIA, F. & NEWMARK, P. A. 2005. Planarian homologs of netrin and netrin receptor are required for proper regeneration of the central nervous system and the maintenance of nervous system architecture. *Development,* 132**,** 3691-703.

CHAN, S. S., ZHENG, H., SU, M. W., WILK, R., KILLEEN, M. T., HEDGECOCK, E. M. & CULOTTI, J. G. 1996. UNC-40, a C. elegans homolog of DCC (Deleted in Colorectal Cancer), is required in motile cells responding to UNC-6 netrin cues. *Cell,* 87**,** 187-95.

CLOUTIER, J. K., MCMANN, C. L., ODERBERG, I. M. & REDDIEN, P. W. 2021. activin-2 is required for regeneration of polarity on the planarian anterior-posterior axis. *PLoS Genet,* 17**,** e1009466.

COLAVITA, A. & CULOTTI, J. G. 1998. Suppressors of ectopic UNC-5 growth cone steering identify eight genes involved in axon guidance in Caenorhabditis elegans. *Dev Biol,* 194**,** 72-85.

CURRIE, K. W. & PEARSON, B. J. 2013. Transcription factors lhx1/5-1 and pitx are required for the maintenance and regeneration of serotonergic neurons in planarians. *Development,* 140**,** 3577-88.

DALVIN, S., ANSELMO, M. A., PRODHAN, P., KOMATSUZAKI, K., SCHNITZER, J. J. & KINANE, T. B. 2003. Expression of Netrin-1 and its two receptors DCC and UNC5H2 in the developing mouse lung. *Gene Expr Patterns,* 3**,** 279-83.

DE BREUCK, S., LARDON, J., ROOMAN, I. & BOUWENS, L. 2003. Netrin-1 expression in fetal and regenerating rat pancreas and its effect on the migration of human pancreatic duct and porcine islet precursor cells. *Diabetologia,* 46**,** 926-33.

FAZELI, A., DICKINSON, S. L., HERMISTON, M. L., TIGHE, R. V., STEEN, R. G., SMALL, C. G., STOECKLI, E. T., KEINO-MASU, K., MASU, M., RAYBURN, H., SIMONS, J., BRONSON, R. T., GORDON, J. I., TESSIER-LAVIGNE, M. & WEINBERG, R. A. 1997. Phenotype of mice lacking functional Deleted in colorectal cancer (Dcc) gene. *Nature,* 386**,** 796-804.

FINCHER, C. T., WURTZEL, O., DE HOOG, T., KRAVARIK, K. M. & REDDIEN, P. W. 2018. Cell type transcriptome atlas for the planarian Schmidtea mediterranea. *Science,* 360.

FINCI, L. I., KRUGER, N., SUN, X., ZHANG, J., CHEGKAZI, M., WU, Y., SCHENK, G., MERTENS, H. D. T., SVERGUN, D. I., ZHANG, Y., WANG, J. H. & MEIJERS, R. 2014. The crystal structure of netrin-1 in complex with DCC reveals the bifunctionality of netrin-1 as a guidance cue. *Neuron,* 83**,** 839-849.

GAVINO, M. A. & REDDIEN, P. W. 2011. A Bmp/Admp regulatory circuit controls maintenance and regeneration of dorsal-ventral polarity in planarians. *Curr Biol,* 21**,** 294-9.

GAVINO, M. A., WENEMOSER, D., WANG, I. E. & REDDIEN, P. W. 2013. Tissue absence initiates regeneration through follistatin-mediated inhibition of activin signaling. *Elife,* 2**,** e00247.

GLISE, B., MILLER, C. A., CROZATIER, M., HALBISEN, M. A., WISE, S., OLSON, D. J., VINCENT, A. & BLAIR, S. S. 2005. Shifted, the Drosophila ortholog of Wnt inhibitory factor-1, controls the distribution and movement of Hedgehog. *Dev Cell,* 8**,** 255-66.

GURLEY, K. A., ELLIOTT, S. A., SIMAKOV, O., SCHMIDT, H. A., HOLSTEIN, T. W. & SANCHEZ ALVARADO, A. 2010. Expression of secreted Wnt pathway components reveals unexpected complexity of the planarian amputation response. *Dev Biol,* 347**,** 24-39.

GURLEY, K. A., RINK, J. C. & SANCHEZ ALVARADO, A. 2008. Beta-catenin defines head versus tail identity during planarian regeneration and homeostasis. *Science,* 319**,** 323-7.

HAMELIN, M., ZHOU, Y., SU, M. W., SCOTT, I. M. & CULOTTI, J. G. 1993. Expression of the UNC-5 guidance receptor in the touch neurons of C. elegans steers their axons dorsally. *Nature,* 364**,** 327-30.

HARRIS, R., SABATELLI, L. M. & SEEGER, M. A. 1996. Guidance cues at the Drosophila CNS midline: identification and characterization of two Drosophila Netrin/UNC-6 homologs. *Neuron,* 17**,** 217-28.

HAYASHI, T., MOTOISHI, M., YAZAWA, S., ITOMI, K., TANEGASHIMA, C., NISHIMURA, O., AGATA, K. & TARUI, H. 2011. A LIM-homeobox gene is required for differentiation of Wnt-expressing cells at the posterior end of the planarian body. *Development,* 138**,** 3679-88.

HEDGECOCK, E. M., CULOTTI, J. G. & HALL, D. H. 1990. The unc-5, unc-6, and unc-40 genes guide circumferential migrations of pioneer axons and mesodermal cells on the epidermis in C. elegans. *Neuron,* 4**,** 61-85.

HONG, K., HINCK, L., NISHIYAMA, M., POO, M. M., TESSIER-LAVIGNE, M. & STEIN, E. 1999. A ligand-gated association between cytoplasmic domains of UNC5 and DCC family receptors converts netrin-induced growth cone attraction to repulsion. *Cell,* 97**,** 927-41.

HSIEH, J. C., KODJABACHIAN, L., REBBERT, M. L., RATTNER, A., SMALLWOOD, P. M., SAMOS, C. H., NUSSE, R., DAWID, I. B. & NATHANS, J. 1999. A new secreted protein that binds to Wnt proteins and inhibits their activities. *Nature,* 398**,** 431-6.

IGLESIAS, M., GOMEZ-SKARMETA, J. L., SALO, E. & ADELL, T. 2008. Silencing of Smed-betacatenin1 generates radial-like hypercephalized planarians. *Development,* 135**,** 1215-21.

ISHII, N., WADSWORTH, W. G., STERN, B. D., CULOTTI, J. G. & HEDGECOCK, E. M. 1992. UNC-6, a laminin-related protein, guides cell and pioneer axon migrations in C. elegans. *Neuron,* 9**,** 873-81.

KANG, J. S., YI, M. J., ZHANG, W., FEINLEIB, J. L., COLE, F. & KRAUSS, R. S. 2004. Netrins and neogenin promote myotube formation. *J Cell Biol,* 167**,** 493-504.

KEINO-MASU, K., MASU, M., HINCK, L., LEONARDO, E. D., CHAN, S. S., CULOTTI, J. G. & TESSIER-LAVIGNE, M. 1996. Deleted in Colorectal Cancer (DCC) encodes a netrin receptor. *Cell,* 87**,** 175-85.

KELEMAN, K. & DICKSON, B. J. 2001. Short- and long-range repulsion by the Drosophila Unc5 netrin receptor. *Neuron,* 32**,** 605-17.

KENNEDY, T. E., SERAFINI, T., DE LA TORRE, J. R. & TESSIER-LAVIGNE, M. 1994. Netrins are diffusible chemotropic factors for commissural axons in the embryonic spinal cord. *Cell,* 78**,** 425-35.

KHARCHENKO, P. V., SILBERSTEIN, L. & SCADDEN, D. T. 2014. Bayesian approach to single-cell differential expression analysis. *Nat Methods,* 11**,** 740-2.

KING, R. S. & NEWMARK, P. A. 2013. In situ hybridization protocol for enhanced detection of gene expression in the planarian Schmidtea mediterranea. *BMC Dev Biol,* 13**,** 8.

KOLODZIEJ, P. A., TIMPE, L. C., MITCHELL, K. J., FRIED, S. R., GOODMAN, C. S., JAN, L. Y. & JAN, Y. N. 1996. frazzled encodes a Drosophila member of the DCC immunoglobulin subfamily and is required for CNS and motor axon guidance. *Cell,* 87**,** 197-204.

LAI WING SUN, K., CORREIA, J. P. & KENNEDY, T. E. 2011. Netrins: versatile extracellular cues with diverse functions. *Development,* 138**,** 2153-69.

LANDER, R. & PETERSEN, C. P. 2016. Wnt, Ptk7, and FGFRL expression gradients control trunk positional identity in planarian regeneration. *Elife,* 5.

LI, D. J., MCMANN, C. L. & REDDIEN, P. W. 2019. Nuclear receptor NR4A is required for patterning at the ends of the planarian anterior-posterior axis. *Elife,* 8.

LIU, Y., STEIN, E., OLIVER, T., LI, Y., BRUNKEN, W. J., KOCH, M., TESSIER-LAVIGNE, M. & HOGAN, B. L. 2004. Novel role for Netrins in regulating epithelial behavior during lung branching morphogenesis. *Curr Biol,* 14**,** 897-905.

LLAMBI, F., CAUSERET, F., BLOCH-GALLEGO, E. & MEHLEN, P. 2001. Netrin-1 acts as a survival factor via its receptors UNC5H and DCC. *EMBO J,* 20**,** 2715-22.

LOWE, C. J., TERASAKI, M., WU, M., FREEMAN, R. M., JR., RUNFT, L., KWAN, K., HAIGO, S., ARONOWICZ, J., LANDER, E., GRUBER, C., SMITH, M., KIRSCHNER, M. & GERHART, J. 2006. Dorsoventral patterning in hemichordates: insights into early chordate evolution. *PLoS Biol,* 4**,** e291.

MÄRZ, M., SEEBECK, F. & BARTSCHERER, K. 2013. A Pitx transcription factor controls the establishment and maintenance of the serotonergic lineage in planarians. *Development,* 140**,** 4499-4509.

MEHLEN, P., RABIZADEH, S., SNIPAS, S. J., ASSA-MUNT, N., SALVESEN, G. S. & BREDESEN, D. E. 1998. The DCC gene product induces apoptosis by a mechanism requiring receptor proteolysis. *Nature,* 395**,** 801-4.

MITCHELL, K. J., DOYLE, J. L., SERAFINI, T., KENNEDY, T. E., TESSIER-LAVIGNE, M., GOODMAN, C. S. & DICKSON, B. J. 1996. Genetic analysis of Netrin genes in Drosophila: Netrins guide CNS commissural axons and peripheral motor axons. *Neuron,* 17**,** 203-15.

MOLINA, M. D., SALO, E. & CEBRIA, F. 2007. The BMP pathway is essential for re-specification and maintenance of the dorsoventral axis in regenerating and intact planarians. *Dev Biol,* 311**,** 79-94.

ODERBERG, I. M., LI, D. J., SCIMONE, M. L., GAVINO, M. A. & REDDIEN, P. W. 2017. Landmarks in Existing Tissue at Wounds Are Utilized to Generate Pattern in Regenerating Tissue. *Curr Biol,* 27**,** 733-742.

ONICHTCHOUK, D., CHEN, Y. G., DOSCH, R., GAWANTKA, V., DELIUS, H., MASSAGUE, J. & NIEHRS, C. 1999. Silencing of TGF-beta signalling by the pseudoreceptor BAMBI. *Nature,* 401**,** 480-5.

ORII, H. & WATANABE, K. 2007. Bone morphogenetic protein is required for dorso-ventral patterning in the planarian Dugesia japonica. *Dev Growth Differ,* 49**,** 345-9.

ORII, H. K., K; AGATA, K; WATANABE, K 1998. Molecular Cloning of the Bone Morphogenetic Protein (BMP) Gene from the Planarian Dugesia japonica. *Zoolog Sci,* 15**,** 6.

OWEN, J. H., WAGNER, D. E., CHEN, C. C., PETERSEN, C. P. & REDDIEN, P. W. 2015. teashirt is required for head-versus-tail regeneration polarity in planarians. *Development,* 142**,** 1062-72.

PEARSON, B. J., EISENHOFFER, G. T., GURLEY, K. A., RINK, J. C., MILLER, D. E. & SANCHEZ ALVARADO, A. 2009. Formaldehyde-based whole-mount in situ hybridization method for planarians. *Dev Dyn,* 238**,** 443-50.

PETERSEN, C. P. & REDDIEN, P. W. 2008. Smed-betacatenin-1 is required for anteroposterior blastema polarity in planarian regeneration. *Science,* 319**,** 327-30.

PETERSEN, C. P. & REDDIEN, P. W. 2009. A wound-induced Wnt expression program controls planarian regeneration polarity. *Proc Natl Acad Sci U S A,* 106**,** 17061-6.

PETERSEN, C. P. & REDDIEN, P. W. 2011. Polarized notum activation at wounds inhibits Wnt function to promote planarian head regeneration. *Science,* 332**,** 852-5.

RAZ, A. A., SRIVASTAVA, M., SALVAMOSER, R. & REDDIEN, P. W. 2017. Acoel regeneration mechanisms indicate an ancient role for muscle in regenerative patterning. *Nat Commun,* 8**,** 1260.

REDDIEN, P. W. 2018. The Cellular and Molecular Basis for Planarian Regeneration. *Cell,* 175**,** 327-345.

REDDIEN, P. W., BERMANGE, A. L., KICZA, A. M. & SANCHEZ ALVARADO, A. 2007. BMP signaling regulates the dorsal planarian midline and is needed for asymmetric regeneration. *Development,* 134**,** 4043-51.

REUTER, H., MARZ, M., VOGG, M. C., ECCLES, D., GRIFOL-BOLDU, L., WEHNER, D., OWLARN, S., ADELL, T., WEIDINGER, G. & BARTSCHERER, K. 2015. Beta-catenin-dependent control of positional information along the AP body axis in planarians involves a teashirt family member. *Cell Rep,* 10**,** 253-65.

ROBERTS-GALBRAITH, R. H. & NEWMARK, P. A. 2013. Follistatin antagonizes activin signaling and acts with notum to direct planarian head regeneration. *Proc Natl Acad Sci U S A,* 110**,** 1363-8.

ROSS, K. G., OMURO, K. C., TAYLOR, M. R., MUNDAY, R. K., HUBERT, A., KING, R. S. & ZAYAS, R. M. 2015. Novel monoclonal antibodies to study tissue regeneration in planarians. *BMC Dev Biol,* 15**,** 2.

ROUHANA, L., WEISS, J. A., FORSTHOEFEL, D. J., LEE, H., KING, R. S., INOUE, T., SHIBATA, N., AGATA, K. & NEWMARK, P. A. 2013. RNA interference by feeding in

vitro-synthesized double-stranded RNA to planarians: methodology and dynamics. *Dev Dyn,* 242**,** 718-30.

SCHAD, E. G. & PETERSEN, C. P. 2020. STRIPAK Limits Stem Cell Differentiation of a WNT Signaling Center to Control Planarian Axis Scaling. *Curr Biol,* 30**,** 254-263 e2.

SCIMONE, M. L., ATABAY, K. D., FINCHER, C. T., BONNEAU, A. R., LI, D. J. & REDDIEN, P. W. 2020. Muscle and neuronal guidepost-like cells facilitate planarian visual system regeneration. *Science,* 368.

SCIMONE, M. L., COTE, L. E. & REDDIEN, P. W. 2017. Orthogonal muscle fibres have different instructive roles in planarian regeneration. *Nature,* 551**,** 623-628.

SCIMONE, M. L., COTE, L. E., ROGERS, T. & REDDIEN, P. W. 2016. Two FGFRL-Wnt circuits organize the planarian anteroposterior axis. *Elife,* 5.

SCIMONE, M. L., WURTZEL, O., MALECEK, K., FINCHER, C. T., ODERBERG, I. M., KRAVARIK, K. M. & REDDIEN, P. W. 2018. foxF-1 Controls Specification of Non-body Wall Muscle and Phagocytic Cells in Planarians. *Curr Biol,* 28**,** 3787-3801 e6.

SERAFINI, T., KENNEDY, T. E., GALKO, M. J., MIRZAYAN, C., JESSELL, T. M. & TESSIER-LAVIGNE, M. 1994. The netrins define a family of axon outgrowth-promoting proteins homologous to C. elegans UNC-6. *Cell,* 78**,** 409-24.

SRINIVASAN, K., STRICKLAND, P., VALDES, A., SHIN, G. C. & HINCK, L. 2003. Netrin-1/neogenin interaction stabilizes multipotent progenitor cap cells during mammary gland morphogenesis. *Dev Cell,* 4**,** 371-82.

SRIVASTAVA, M. 2022. Studying development, regeneration, stem cells, and more in the acoel Hofstenia miamia. *Curr Top Dev Biol,* 147**,** 153-172.

SRIVASTAVA, M., MAZZA-CURLL, K. L., VAN WOLFSWINKEL, J. C. & REDDIEN, P. W. 2014. Whole-body acoel regeneration is controlled by Wnt and Bmp-Admp signaling. *Curr Biol,* 24**,** 1107-13.

STUCKEMANN, T., CLELAND, J. P., WERNER, S., THI-KIM VU, H., BAYERSDORF, R., LIU, S. Y., FRIEDRICH, B., JULICHER, F. & RINK, J. C. 2017. Antagonistic Self-Organizing Patterning Systems Control Maintenance and Regeneration of the Anteroposterior Axis in Planarians. *Dev Cell,* 40**,** 248-263 e4.

TEWARI, A. G., STERN, S. R., ODERBERG, I. M. & REDDIEN, P. W. 2018. Cellular and Molecular Responses Unique to Major Injury Are Dispensable for Planarian Regeneration. *Cell Rep,* 25**,** 2577-2590 e3.

VOGG, M. C., GALLIOT, B. & TSIAIRIS, C. D. 2019. Model systems for regeneration: Hydra. *Development,* 146.

VU, H. T., MANSOUR, S., KUCKEN, M., BLASSE, C., BASQUIN, C., AZIMZADEH, J., MYERS, E. W., BRUSCH, L. & RINK, J. C. 2019. Dynamic Polarization of the Multiciliated Planarian Epidermis between Body Plan Landmarks. *Dev Cell,* 51**,** 526-542 e6.

WAGNER, D. E., WANG, I. E. & REDDIEN, P. W. 2011. Clonogenic neoblasts are pluripotent adult stem cells that underlie planarian regeneration. *Science,* 332**,** 811-6.

WITCHLEY, J. N., MAYER, M., WAGNER, D. E., OWEN, J. H. & REDDIEN, P. W. 2013. Muscle cells provide instructions for planarian regeneration. *Cell Rep,* 4**,** 633-41.

YEBRA, M., MONTGOMERY, A. M., DIAFERIA, G. R., KAIDO, T., SILLETTI, S., PEREZ, B., JUST, M. L., HILDBRAND, S., HURFORD, R., FLORKIEWICZ, E., TESSIER-LAVIGNE, M. & CIRULLI, V. 2003. Recognition of the neural chemoattractant Netrin-1

by integrins alpha6beta4 and alpha3beta1 regulates epithelial cell adhesion and migration. *Dev Cell,* 5**,** 695-707.

ZENG, A., LI, H., GUO, L., GAO, X., MCKINNEY, S., WANG, Y., YU, Z., PARK, J., SEMERAD, C., ROSS, E., CHENG, L. C., DAVIES, E., LEI, K., WANG, W., PERERA, A., HALL, K., PEAK, A., BOX, A. & SANCHEZ ALVARADO, A. 2018. Prospectively Isolated Tetraspanin(+) Neoblasts Are Adult Pluripotent Stem Cells Underlying Planaria Regeneration. *Cell,* 173**,** 1593-1608 e20.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

| *bambi-2* | *slit* | *wnt5* | *bmp4* | *bmp4* | *admp* | *admp* | *notum* | *wnt1* |

Control RNAi

8xF
14dR

*netrin-1,-2,
-3,-4,-5* RNAi

B

Control RNAi   *unc5-C;-E;dcc-2* RNAi

8xF
14dR

bambi-2

C

relative midline size

Control   *netrin-1;-2;
-3;-4;-5*   *unc5-C;-E;
dcc-2*
RNAi

D

Control RNAi   *netrin-1,-2,
-3,-4,-5* RNAi   *unc5-C;-E;dcc-2*
RNAi

8xF
14dR

Muscle fibers (6G10)

Figure 6

A

bambi-2

L ——————— M ——————— L

↓ **amputation**

↓ pitx, islet

↓ **midline signals outgrowth**

L ——— M ——— L

B

Dorsal

unc5-C
unc5-E
dcc-2

bambi-2+ **midline muscle**

net-1   *net-4*   *net-1*

net-5

Ventral

Figure S1

Figure S2

Figure S3

Figure S4

A

Score for pointy blastema

Normalized score of blastema pointiness

RNAi condition

netrin-1	-	+	+	+	+	-	-	-	-	-	-	+	+	+	+	+	+	-	-	-	-	-	+	+	+	+	+	-	+
netrin-2	-	+	-	-	-	+	+	+	-	-	-	+	+	+	-	-	-	+	+	+	-	+	+	+	+	-	+	+	+
netrin-3	-	-	+	-	-	+	-	-	+	+	-	+	-	-	+	+	+	-	+	+	-	+	+	+	-	+	+	+	+
netrin-4	-	-	-	+	-	-	+	-	+	-	+	-	+	-	+	-	+	+	-	+	+	+	-	+	+	+	+	+	+
netrin-5	-	-	-	-	+	-	-	+	-	+	+	-	+	-	+	-	+	+	+	+	+	-	+	+	+	+	+	+	+

Control RNAi — 1/11

net-1;-2;-3 RNAi — 10/11

net-4;-5 RNAi — 2/11

net-1;-2;-4;-5 RNAi — 6/10

net-1;-3;-4;-5 RNAi — 4/10

net-1;-2;-3;-4;-5 RNAi — 10/10

8xF 14dR

B

Score for indented blastema

Percentage of indented posterior blastemas

RNAi condition

netrin-1	-	+	+	+	+	-	-	-	-	-	-	+	+	+	+	+	+	-	-	-	-	-	+	+	+	+	+	-	+
netrin-2	-	+	-	-	-	+	+	+	-	-	-	+	+	+	-	-	-	+	+	+	-	+	+	+	+	-	+	+	+
netrin-3	-	-	+	-	-	+	-	-	+	+	-	+	-	-	+	+	+	-	+	+	-	+	+	+	-	+	+	+	+
netrin-4	-	-	-	+	-	-	+	-	+	-	+	-	+	-	+	-	+	+	-	+	+	+	-	+	+	+	+	+	+
netrin-5	-	-	-	-	+	-	-	+	-	+	+	-	+	-	+	-	+	+	+	+	+	-	+	+	+	+	+	+	+

8xF 14dR

Control RNAi — 11/11

net-1;-2;-4;-5 RNAi — 3/10

net-1;-4;-5 RNAi — 5/11

Figure S5