1

4 **Title:** Nonlinear computation by a habenula-driven recurrent inhibitory network in the raphe

6
7 **Short title:** Facilitating serotonin release in a recurrent network
8
9

10 **Authors:**
11 Michael B. Lynn[1], Sean Geddes[1], Mohamad Chahrour[1], Sébastien Maillé[1], Léa Caya-
12 Bissonnette[1], Emerson Harkin[1], Érik Harvey-Girard[1], Samir Haj-Dahmane[5], Richard Naud[1,2,3,4],
13 Jean-Claude Béïque[1,2,3]*
14

Affiliations:
[1]Department of Cellular and Molecular Medicine, University of Ottawa, Ottawa, ON K1H 8M5,
Canada
[2]Centre for Neural Dynamics and AI, University of Ottawa, Ottawa, ON K1H 8M5, Canada
[3]Brain and Mind Research Institute, University of Ottawa, ON K1H 8M5, Canada
[4]Department of Physics, STEM Complex, Room 336, 150 Louis Pasteur Private, University of
Ottawa, Ottawa, ON K1N 6N5, Canada
[5]Department of Pharmacology and Toxicology, University at Buffalo, Buffalo, NY 14203,
U.S.A.
24

25 **Corresponding Authors**:

26 * Correspondence: jbeique@uottawa.ca (J.-C. B.).

Abstract

Serotonin (5-HT) neurons in the dorsal raphe nucleus (DRN) receive a constellation of long-range inputs, yet guiding principles of local circuit organization and underlying computations in this nucleus are largely unknown. Using inputs from the lateral habenula (LHb) to interrogate the processing features of the DRN, we uncovered 5-HT1A receptor-mediated recurrent connections between 5-HT neurons, refuting classical theories of autoinhibition. Cellular electrophysiology and imaging of a genetically encoded 5-HT sensor revealed that these recurrent inhibitory connections spanned the raphe, were slow, stochastic, strongly facilitating, and gated spike output. These features collectively conveyed highly non-linear dynamics to this network, generating excitation-driven inhibition and winner-take-all computations. *In vivo* optogenetic activation of LHb inputs to DRN, at frequencies where these computations are predicted to ignite, transiently disrupted expression of a reward-conditioned response in an auditory conditioning task. Together, these data identify a core computation supported by an unsuspected slow serotonergic recurrent inhibitory network.

Keywords: Serotonin; dorsal raphe nucleus; habenula; recurrent dynamics; short-term plasticity; winner-take-all

Introduction

The serotonergic dorsal raphe nucleus (DRN) is a heterogeneous midbrain nucleus which regulates mood and behavior through its widespread efferent innervation (Dorocic et al., 2014; Weissbourd et al., 2014). The coding features of DRN 5-HT neurons still remain largely elusive, although recent studies have shown that they respond to environmental variables such as reward and threat (Cohen et al., 2015; Li et al., 2016; Zhong et al., 2017; Seo et al., 2019), making them suited to supervise behavioral policies based on internal representations of environmental contexts. Moreover, the DRN has been linked to a diverse range of mood-related states such as aggression (Takahashi et al., 2015), patience (Miyazaki et al., 2014; Miyazaki et al., 2018; Miyazaki et al., 2020), and depressive and anxiety disorders (Graeff et al., 1996; Warden et al., 2012). DRN may additionally support cognitive flexibility or modify learning rate (Matias et al., 2017; Grossman et al., 2022). DRN microcircuits are composed of multiple intermingled cell types, including dopaminergic, glutamatergic, serotonergic and GABAergic neurons, organized into functionally distinct modules (Ren et al., 2018). However, the precise rules by which these circuits process incoming information to ultimately provide instructive cues to regulate behavior are largely unknown.

One major source of input to the DRN is from the lateral habenula (LHb), which provides monosynaptic excitatory drive to 5-HT neurons (Zhou et al., 2017). LHb neurons fire in response to punishments, reward omissions and graded threat information by phasic, high-frequency discharges (Matsumoto & Hikosaka, 2009; Proulx et al., 2014; Bromberg-Martin & Hikosaka, 2011; Lecca et al., 2020; Amo et al., 2014; Yang et al., 2018). Recent work has suggested that both the raphe and habenula regulate behavioral policy over long timescales by modifying internal representations (Marques et al., 2020). How incoming synaptic inputs from LHb are processed by DRN microcircuits into an internal state representation that persists over longer timescales to influence behavioral policy is currently unknown.

Here, to elucidate key computations performed by local DRN microcircuits on synaptic inputs, we combined optogenetic, electrophysiological, two-photon imaging, computational and behavioral approaches. Using long-range inputs from LHb to interrogate DRN functional circuitry, we uncovered an unexpected slow inhibitory conductance activated by local recurrent connections between 5-HT neurons. These recurrent connections, mediated by inhibitory 5-HT$_{1A}$ receptors (5-HT$_{1A}$Rs), were slow, highly facilitating, and strongly modulated spiking output of 5-HT neurons. We developed an experimentally-grounded model of the complex short-term dynamics of these recurrent connections, which predicts a sharp transition from net excitation to excitation-driven inhibition and winner-take-all dynamics between long-range inputs to ensembles in DRN. Key predictions of this model were validated in a simple auditory conditioning task. Together, these findings outline highly nonlinear computations supported by an unsuspected recurrent inhibitory network between 5-HT neurons in the DRN.

Results

Excitatory monosynaptic LHb inputs to DRN trigger a frequency-dependent indirect inhibition

As a first step towards uncovering organizing principles of microcircuitry within DRN, we activated LHb, a strong long-range input to DRN, and probed neural responses. We utilized a SERT-Cre::tdTom mouse line for whole-cell electrophysiological recordings from genetically identified 5-HT neurons, validating this approach with immunohistochemical staining (Fig. S1A-D; 98.4% overlap between TPH2 and tdTomato expression). Upon bilateral injection of AAV-GFP in LHb, we observed high LHb axon density in DRN (Fig. S2A) consistent with recent work (Zhou et al., 2017). Next, we injected AAV-ChR2(H134R)-mCherry bilaterally into LHb of SERT-Cre::tdTom mice (Fig. 1A and Fig. S1E, see Methods). Photostimulation induced strong and reliable action potentials in LHb neurons which were qualitatively similar to those following current step injections (Fig. S2B-C). Within the DRN, we targeted 5-HT neurons predominantly within the ventromedial DRN, an area exhibiting high LHb axonal density (Fig. S2A; Fig. S3; Zhou et al., 2017). Recording from identified DRN 5-HT neurons from this subregion in the voltage-clamp configuration (Fig. 1C-D), we photostimulated LHb axons with brief (1-3ms) pulses of blue light, triggering synaptic release of glutamate that activated AMPARs (Fig. 1E; V_m=-70 mV) and NMDARs (Fig. 1F; V_m=+40mV). LED-stimulated glutamate release was blocked by TTX and rescued by the K^+ channel blocker 4-AP (Fig. 1G), indicating that LHb makes monosynaptic contacts with DRN 5-HT neurons. These excitatory inputs from LHb were capable of driving 5-HT neurons to spike (Fig. 1H). Thus, LHb afferents directly excite 5-HT neurons, consistent with previous reports (Zhou et al., 2017). While habenula-driven feedforward inhibition onto 5-HT neurons has been previously reported (Zhou et al., 2017; but see Dorocic et al., 2014), we observed little evidence of optically triggered delayed IPSCs in our recording conditions (only 1.7% of 5-HT neurons exhibited IPSCs, and this only upon 20 Hz LHb train photostimulation, n=116). Activation of ChR2-expressing local DRN GABAergic neurons (using a SOM-Cre:: tdTom mouse line) triggered IPSCs in GABAergic neurons (Fig. S4A-G), in agreement with previous work showing local feedforward circuitry in the raphe (Geddes et al., 2016; Zhou et al., 2017). However, optogenetic activation of LHb inputs produced primarily subthreshold excitation of GABAergic neurons (peak depolarization of ~7mV; Fig. S4H-O), presumably limiting the ability of these long-range inputs to trigger disynaptic inhibition onto 5-HT neurons, at least in our recording conditions and locations. Thus, DRN 5-HT neurons within the ventromedial subdivision of DRN principally receive monosynaptic long-range excitatory input from LHb.

Populations of LHb neurons can discharge at high rates following negative reward prediction errors, depression or stress, or graded threat information (Matsumoto & Hikosaka, 2009, Yang et al., 2018, Dolzani et al., 2016, Amo et al., 2014; Groos & Helmchen, 2024; Andalman et al., 2019). To explore how such input patterns are parsed by DRN microcircuits, we delivered trains of optical stimuli at a range of frequencies consistent with previously reported *in vivo* phasic habenular responses (Amo et al., 2014; Groos & Helmchen, 2024). Train stimulation at 10Hz or 20Hz, surprisingly, triggered the appearance of a delayed outward current lasting hundreds of milliseconds, following the initial monosynaptic EPSCs (Fig. 1I). The overall kinetics of this protracted current, as well as its underlying rectifying I-V curve displaying a reversal potential at ~-77mV (Fig. 1J), suggest the involvement of a G-protein inwardly rectifying potassium (GIRK) channel (Newberry & Nicoll, 1984). While several G-protein coupled receptor subtypes couple to GIRK channels, the inhibitory 5-HT receptor of the 5-HT$_{1A}$ subtype are expressed at high level in DRN 5-HT neurons (Pazos & Palacios, 1985) and were

Figure 1: Habenula input to raphe triggers direct excitation and heterosynaptic inhibition in 5-HT neurons. **(A-B)** Schematic of experimental approach. **(C)** Epifluorescent image of tdTomato-positive 5-HT neurons. **(D)** Application of 5-CT (100nM) triggered a transient increase in holding potential. **(E)** Single pulses of light at Vm=-70mV triggered EPSCs (left) which were abolished by application of NBQX (10uM, right). **(F)** At Vm=+40mV, single pulses of light triggered EPSCs (left) which were abolished by application of APV (50uM, right). **(G)** Application of TTX (1uM) blocked EPSCs at Vm=-70mV, while subsequent application of 4-AP (1mM) recovered the current. **(H)** Voltage-clamp recordings of ChR2-EPSCs (Vm = -70 mV) and spiking behavior in response to presynaptic activation. **(I)** Representative 5-HT neuron responses to single pulses of light (left) and pulse trains at 10Hz (right). **(J)** Representative traces of the current-voltage (I-V) relationship of the delayed outward current (left) and group data (right, n=3 neurons, mean +/- SE) **(K)** Representative timecourse of peak amplitude stability over time. **(L)** Representative traces and timecourse showing the blockade of the outward current by WAY 100-635 (400 nM; left). Population data presented in a scatter-line plot (right, Pre-WAY, 7.47 +/- 0.8; Post-WAY, 0.28 +/- 0.4; n = 4 neurons; p = 0.001; paired two-sided t-test). All recordings performed at room temperature; see Fig. S10H-K for replication at 30-32C.

thus reasonable candidates. Bath application of the 5-HT$_{1A}$R antagonist WAY100-635 abolished
the outward current triggered by burst activation of LHb axons in the DRN (Fig. 1K-L) without
influencing the light-induced EPSCs (pre-WAY amplitude, 25.2 +- 12.5pA; post-WAY
amplitude, 17.9 +- 6.9pA; n=3 neurons, p=0.48, paired t-test). Across all recordings from
genetically identified 5-HT neurons, 33.9% showed no response to optogenetic LHb stimulation,
50.8% exhibited direct excitation only, 12.7% exhibited direct excitation and indirect 5-HT$_{1A}$
inhibition, and 0.84% exhibited only indirect 5-HT$_{1A}$ inhibition. Together, these results
demonstrate that trains of LHb input to DRN 5-HT neurons trigger rapid monosynaptic
glutamatergic-mediated excitation followed by a slower, 5-HT$_{1A}$R-mediated inhibition.
181
DRN 5-HT neurons are organized in a recurrent inhibitory network
We next sought to identify the cellular and network underpinnings of this non-canonical
mixed excitation/inhibition response, initially focusing on the protracted 5-HT$_{1A}$R-mediated
inhibition. A dominant and well-accepted model posits that 5-HT1ARs are autoreceptors. In this
model, individual 5-HT neurons, upon firing, limit their own excitability through the dendritic
release of 5-HT (Maxwell et al., 1983; Ridet et al., 1994; Bunin & Wightmann, 1998), which
activates these so-called inhibitory somatodendritic 5-HT$_{1A}$ autoreceptors in an autocrine fashion
(Gozlan et al., 1983; Weissmann-Nanopoulos et al., 1985; Pineyro & Blier, 1999). This
canonical autoreceptor model is nonetheless difficult to reconcile with our observations that train
activation of LHb inputs triggered 5-HT$_{1A}$R-mediated outward currents in nonspiking, voltage-
clamped 5-HT neurons (Fig. 1I,J), or in 5-HT neurons receiving no functional excitatory LHb
inputs (Fig. S2D-E). We therefore next sought to experimentally disambiguate the autoreceptor
hypothesis from the alternative hypothesis that DRN 5-HT neurons are organized in a recurrent
inhibitory network architecture (Fig. 2A).
We first reasoned that if 5-HT$_{1A}$Rs were indeed autocrine inhibitory autoreceptors,
evoked spikes from a single 5-HT neuron should elicit a 5-HT1AR-mediated inhibition in the
same isolated neuron, contributing to the post-spike afterhyperpolarization (AHP) commonly
observed in 5-HT neurons (Vandermaelen & Aghajanian, 1983). Whole-cell recordings,
however, revealed that the prominent post-spiking AHP triggered by current injection was
insensitive to application of the 5-HT$_{1A}$R antagonist WAY100-635 (Fig. 2B-D), yet fully
abolished by application of the SK channel blocker apamin (Fig. 2E-H), demonstrating that the
slow AHP in 5-HT neurons is mediated by SK channels, and not 5-HT$_{1A}$Rs. The absence of
autocrine effects on the AHP could be explained by the small number of action potentials
emitted in current-clamp experiments. Voltage-clamp experiments further showed that the robust
outward current that followed long, constant, depolarizing voltage steps in 5-HT neurons was
also sensitive to apamin, but not to WAY100-635 (Fig. S5). Together, these experiments failed
to support an autocrine function for 5-HT$_{1A}$Rs on 5-HT neurons.
To begin exploring the alternative possibility of a recurrent network between 5-HT
neurons in the DRN, we designed a series of targeted optogenetic strategies based on viral
transduction of 5-HT neurons with a Cre-dependent AAV ChETA-YFP in SERT-Cre::tdTom
mice (Fig. S5A, see Methods). *Ex vivo* slice recordings from DRN 5-HT neurons co-expressing
tdTomato and YFP-tagged ChETA opsin (Fig. S7B-C) showed that LED stimulation induced
robust inward photocurrents and depolarizations that induced spiking reliably up to ~20 Hz (Fig.
S7D-E). Train photostimulation of all transduced 5-HT neurons triggered a robust inhibitory 5-
HT$_{1A}$R conductance (Fig. S7F-G). We then reasoned that neuronal spiking induced by
photostimulating ChETA-expressing 5-HT neurons should reveal an additive 5-HT$_{1A}$R mediated

Figure 2: 5-HT neurons in raphe are organized in a recurrent inhibitory network. **(A)** Two hypothesized mechanisms of 5-HT1AR-based signaling in raphe. **(B)** Representative traces of spiking in response to direct current injection before and after WAY 100-635 application. **(C)** Direct comparison of AHP before and after application of WAY 100-635. **(D)** Representative scatter plot showing no change in the AHP amplitude with bath application of WAY 100-635. **(E)** Representative traces of spiking in response to direct current injection before and after APAMIN application. **(F)** Relationship between current injection strength and evoked spikes, in the presence or absence of APAMIN (n=3 neurons, mean +/- SE) **(G)** Direct comparison of AHP before and after application of APAMIN. **(H)** Relationship between spike number and AHP amplitude, in the presence or absence of APAMIN (n=3 neurons, mean +/- SE). **(I)** Schematic of local optogenetic experimental approach. **(J)** Representative traces from CHETA-expressing 5-HT neurons showing current step-driven spiking before (green) and after (black) bath application of WAY100-635. Grey lines represent single sweeps from the same neuron. **(K)** Quantification of AHP statistics for control and WAY100-635 conditions (AHP amplitude, p=0.96, n=4 neurons; AHP area, p=0.85, n=4 neurons; AHP tau, p=0.09, n=3 neurons; paired

two-sided t-tests). Grey lines represent single sweeps from the same neuron. **(L)** Representative traces from CHETA-expressing 5-HT neurons showing photostimulation-driven spiking before (green) and after (black) bath application of WAY100-635. **(M)** Quantification of AHP statistics for control and WAY100-635 conditions (AHP amplitude, p=0.03, n=5 neurons; AHP area, p=0.03, n=5 neurons; AHP tau, p=0.02, n=3 neurons; paired two-sided t-tests). **(N)** Schematic and confocal images of sparse local optogenetic approach (AAV-DIO-CHETA-YFP diluted in mannitol 2:1). **(O)** While recording from an opsin-expressing neuron, brief optical stimulation could reliably elicit spiking behavior. **(P)** While recording from a non-opsin-expressing serotonergic neuron, photostimulation at 20Hz produced an inhibitory current; note the absence of direct photocurrents. **(Q)** Representative traces of the current-voltage (I-V) relationship of the inhibitory current (left) and group data (right, n=5 neurons, mean +/- SE) **(R)** Representative traces and timecourse of TTX (1uM) bath application (left) and group data (right; n=3 neurons; grey average trace: mean +/- SE). **(S)** Representative traces and timecourse of WAY100-635 (400nM) bath application (left) and group data (right; n=6 neurons; grey average trace: mean +/- SE). All recordings performed at room temperature.

component of the resultant AHP in the recorded neuron if DRN 5-HT neurons were indeed
interconnected in a recurrent inhibitory network. To test this, we cross-calibrated single-cell
direct current injections and LED intensity such that both methods independently triggered
spiking of similar frequency (Fig. 2J and 2L). While the spiking triggered by direct current
injections induced a strong and highly reliable AHP in these neurons (*i.e.,* tdTomato and YFP-
tagged ChETA-positive) that was not affected by the 5-HT$_{1A}$R antagonist WAY100-635 (Fig.
2K-L), the same spiking triggered by LED photostimulation of the entire DRN network induced
a substantially larger AHP, and the additional component was blocked by WAY100-635 (Fig.
2M). These results fail to support the autocrine hypothesis of 5-HT$_{1A}$R function, and rather
suggest that 5-HT neurons are rather organized in a recurrent inhibitory network.
To more directly test the presence of recurrent connectivity in the raphe, we next
developed an AAV-ChETA-YFP transduction strategy in the DRN of SERT-Cre::tdTom mice
(Fig. 2N) to achieve sparse, mosaic-like, expression of ChETA-YFP in 5-HT neurons (see
Methods). This allowed us to obtain visually targeted recordings either from ChETA-expressing
5-HT neurons (*i.e.,* YFP and tdTomato expressing neurons that exhibited reliable optically
driven spiking; Fig. 2O) or from non ChETA-expressing 5-HT neurons (*i.e.,* YFP-negative,
tdTomato-positive 5-HT neurons). Targeted recordings from non ChETA-expressing 5-HT
neurons revealed that high-frequency LED photostimulation (which directly drives the
surrounding network but not the recorded neuron) induced delayed, strongly rectifying, outward
currents (Fig. 2P-Q) which were blocked by TTX and by WAY 100-635 (Fig. 2R-S). Taken
together, these optogenetic data strongly support the existence of an unsuspected recurrent
inhibitory network in DRN, consisting of 5-HT neurons communicating via 5-HT$_{1A}$Rs.
To further explore connectivity features between 5-HT neurons in the DRN, we
performed small, visually guided injections of fluorescent dextran retrograde tracer into either
dorsal DRN or ventral DRN in acute slices from SERT-Cre::tdTom mice (dorsal DRN injection
diameter: 105.6 +-37.7um, ventral DRN injection diameter: 91.0 +- 22.4um; p=0.69,
independent t-test, n=3 injections each). We quantified the locations of neurons expressing both
fluorescent dextran and tdTomato signals (Fig. S7A; see Methods; Trinh et al., 2016), revealing
dual-labeled neurons several hundred microns outside of the injection site (Fig S7B). The
locations of both dual-labeled neurons and single-labeled tdTomato-expressing 5-HT neurons,
were extracted for each individual injection (Fig. S7C). Pooling across all injections revealed a
spatial organization of labeled neurons relative to the injection center that was distinct for dorsal
and ventral injections (Fig. S7D). Labeled neurons were found at distances up to ~500 μm from
the injection center, and the distribution of distances from the injection site was significantly
different between dorsal and ventral injections (Fig. S7E; p=1.29*10^-3; Kolmogorov-Smirnov
test), suggesting submodule-specific differences in spatial connectivity profiles. Furthermore,
dorsal injections were most similar to unstructured connectivity (random sampling from all 5-HT
neurons), while ventral injections had more local organization of connectivity (Fig. S7F; see
Methods). As a final step, we computed a polar plot of the angle of labeled neurons relative to
the injection center, which for dorsal injections revealed an enrichment in ventrally located
labeled 5-HT neurons, and for ventral injections led to a more homogeneous distribution of
labeled neurons in polar coordinates (Fig. S7G-H). Together, these experiments outline a set of
spatial and anatomical features of the recurrent inhibitory network within DRN.
261
A quantitative description of release dynamics at 5-HT$_{1A}$R-mediated recurrent connections

We next parameterized key functional features of the serotonergic recurrent inhibitory
network. Train optogenetic stimulation (20Hz, 8 pulses) of 5-HT neurons triggered a 5-HT$_{1A}$R
conductance that peaked approximately 400 ms following the end of the train (392.6 ± 106.9ms)
and that slowly decayed (2.1 ± 0.46s). Varying the number of light pulses delivered at 20Hz
revealed a relationship between pulse number and postsynaptic conductance that was almost
linear, indicating that 5-HT$_{1A}$R coupling to GIRK channels can follow multiple successive
activation by endogenous 5-HT with limited desensitization (Fig. S8A-C). To infer presynaptic
release dynamics, we next calculated the response variance as a function of pulse number,
finding virtually no change in variance despite changes in amplitude (Fig. S8D). While such
dynamics are in principle consistent with deterministic neurotransmitter release (*i.e.,* P$_{rel}$ ~1; see
Methods for full details), artefactual inflation of presynaptic release probability has previously
been reported for optogenetic synaptic terminal activation (Zhang & Oertner, 2007) (see
Methods). These results nonetheless outline the timecourse of a GIRK- and 5-HT$_{1A}$R -mediated
synapse that quasi-linearly reports presynaptic 5-HT release over a physiological range (see
below).
To monitor 5-HT presynaptic release dynamics in the DRN circumventing the confound
brought about by optogenetic stimulation, we next recorded DRN 5-HT neurons while
electrically stimulating nearby serotonergic fibers with a bipolar concentric electrode (Fig. 3A).
A cocktail of synaptic blockers was bath-applied to isolate serotonergic transmission (Fig. 3A,
see Methods). High-frequency electrical stimulation readily triggered a 5-HT$_{1A}$R -mediated
outward current with a similar time course to that induced by LED stimulation (Fig. 3B). The
subset of 5-HT neurons displaying an electrically evoked outward current did not exhibit clear
anatomical organizing principles within the ventral DRN (Fig. S3B; circles filled with red). We
next delivered isolated electrical pulses under a low-intensity minimal stimulation regime (*i.e.,*
activating small numbers of individual 5-HT synapses) while monitoring 5-HT$_{1A}$R -mediated
currents. These recordings showed instances of clear successes (~5 pA) and failures (Fig. 3C) of
5-HT release, consistent with small numbers of 5-HT synapses each exhibiting a P$_{rel}$ < 1 (Lynn et
al., 2020). Fig. 3D depicts the distribution of event charges plotted alongside neuron-specific
control distributions (obtained from integrating an equivalent duration of noise; see Methods).
The bimodal event distribution, exhibiting a left peak overlapping with the noise distribution, is
consistent with probabilistic 5-HT transmission.
We next examined the short-term dynamics of these 5-HT synapses by delivering
electrical stimulation trains with a fixed pulse number (8) and variable frequency (Fig. 3E). A
deterministic synapse should exhibit identical responses over a range of input frequencies if
pulse number is fixed. While 1 Hz stimulation produced a notably weak response, higher
frequencies of 10 Hz and 20 Hz elicited progressively larger inhibitory currents (Fig. 3E-F). To
directly compare these protracted responses, which displayed peak amplitudes that were highly
dependent on stimulus parameters, we employed a z-score based metric of total charge. Briefly,
we computed the total charge evoked by each stimulus train, and compared this with a surrogate
noise distribution constructed by performing the same integration on an equivalent duration of
baseline recording (example neuron shown in Fig. 3G; two-sided t-test on trial distribution of
charges for events and noise; See Methods). While low-frequency (1 Hz) stimuli did not produce
responses significantly different from the surrogate noise distribution (Fig. 3H; 6/10 neurons
with p>0.05 for event *vs* noise charges, two-sided t-test), high-frequency (20 Hz) stimuli
produced progressively larger responses (Fig. 3H), consistent with short-term facilitation.

Figure 3: Serotonin release is probabilistic and strongly facilitating at inhibitory recurrent connections. **(A)** Experimental schematic. **(B)** Electrical stimulation at 20Hz evoked an outward current that is blocked by WAY100-635. **(C)** Representative traces depicting responses to low-amplitude electrical stimulation, showing sorted successes and failures from the same neuron. **(D)** Histograms of charge z-score, calculated for post-stimulus events (blue) and an equivalent period of non stimulus-driven noise from the same neuron (green) (n=424 events from 4 neurons). **(E)** Experimental setup for frequency-varying electrical stimulation experiments (left). Representative averaged traces from a single neuron depicting responses to electrical stimulation at various frequencies (right). **(F)** Expanded traces showing responses to 1Hz and 20Hz stimuli. **(G)** Charge histograms for stimulus-driven events and noise (one example neuron; 1Hz: p>0.05, 20Hz: p<0.01, two-sided t-tests). **(H)** Charge z-score as a function of frequency, with 95% noise C.I shown in green (n=10 neurons, two-sided t-tests for each neuron as in G). **(I)** Paired-pulse electrical stimulation paradigm. **(J)** Representative averaged traces from a single neuron depicting responses to electrical stimulation at various paired-pulse intervals. **(K)** Charge histograms for single-pulse responses and paired-pulse responses at 50ms for one example neuron. **(L)** Post-stimulus charge as a function of paired-pulse time interval (n=7 neurons). Green line depicts linear summation. **(M)** Pulse-chase electrical stimulation paradigm. **(N)** Representative averaged traces from a single neuron depicting responses to pulse-chase electrical stimulation protocol at various temporal intervals. **(O)** Post-stimulus charge as a function of pulse-chase time interval (n=6 neurons). Green line depicts linear summation. All recordings performed at 30-32C and [Ca2+]=2.5mM; see Fig. S10A-G for replication at lower external calcium concentrations.

Altogether, we conclude that the protracted 5-HT1AR-mediated inhibition reflects the
probabilistic release of 5-HT that exhibits substantial short-term facilitation.
To further quantify the timescale of these short-term synaptic dynamics, we delivered
patterned electrical stimulation, starting with classic paired-pulse stimulation experiments. When
paired-pulse stimuli were closely spaced (50-200 ms) we observed supralinear summation
compared to isolated single pulses; in contrast, paired-pulse intervals of 500 ms and longer
produced depressing responses (Fig. 3J-K). This short-term depression observed at timescales of
seconds is consistent with previous reports on $5\text{-}HT_{1A}R$-mediated inhibition (Courtney & Ford,
2016). By comparing the actual evoked charge with the predicted linear summation of isolated
single-pulse responses, we quantified the dynamics of these recurrent synapses across multiple
timescales (Fig. 3L). Together, these results indicate that the short-term dynamics of these 5-HT
synapses are dominated by strong short-term facilitation at short intervals, transitioning to mild
depression at timescales > 500 ms.
In the experiments above, we aimed to approach conditions of minimal stimulation (i.e.,
successes and failures). We additionally repeated these experiments with stronger electrical
stimulation (Fig. S9) and under low extracellular calcium conditions (Fig. S10A-G; $[Ca^{2+}]$=1.5
mM), and found that 5-HT release dynamics still exhibited strong short-term facilitation,
especially for short inter-pulse intervals in these conditions. Likewise, the recruitment of 5-
$HT_{1A}R$-mediated inhibition by high frequency optogenetic activation of LHb inputs to the DRN
(Fig. 1I) still remained in these conditions ($[Ca^{2+}]$=1.5 mM; 30-32 °C; Fig. S10H-K).
To examine meta-dynamic features of these 5-HT synapses, we delivered a second set of
patterned electrical stimuli consisting of a paired-pulse stimulation (20 Hz; ISI of 50 ms)
separated from another paired-pulse stimulation by some varying time interval (pulse-chase; Fig.
3M). These experiments revealed, unexpectedly, that the paired-pulse facilitation at 20 Hz is
highly dependent on stimulus history over a timescale of seconds. When paired-pulse sets were
separated by short intervals, facilitation was present for both sets, yielding a supralinear response
to the patterned input (Fig. 3N). However, when separated by intervals of >500 ms, the second
paired-pulse stimulus elicited notably smaller responses (less facilitation), indicating release
meta-dynamics, or a history-dependent change in facilitation (Fig. 3N). Charge quantification
metrics, similar to above (see Methods), revealed that this dependence on stimulation history
lasted up to 10 seconds (Fig. 3O).
Any effort to identify the network computations sustained by this recurrent inhibitory
network requires, as an obligatory intermediate step, an accurate quantitative model of the
synaptic dynamics described above. We therefore next developed a mathematical model
capturing these ground-truth short-term dynamics of 5-HT synapses. We first fitted the
experimentally-derived $5\text{-}HT_{1A}R$ -mediated inhibitory GIRK current with a simple summed
exponential model (Fig. 4A). Next, to adequately capture the short-term dynamics of 5-HT
synapses, we fitted our experimental data using a previously described spike-response plasticity
model (SRP; Rossbroich et al., 2021; see Methods). Briefly, a presynaptic spiketrain is
convolved with a kernel indicating the sign and timescale of synaptic dynamics, passed through a
sigmoidal readout, and convolved at each presynaptic spike-time with a postsynaptic event
kernel (here, describing a GIRK-mediated current). Using the paired-pulse electrical stimulation
data as input, we used a least-squares approach to find a best fit, yielding an optimized short-
term plasticity (STP) kernel that showed a short-timescale facilitation and long-timescale
depression components (Fig. 4B). The resultant model effectively captured the dynamics of the
paired-pulse experimental data across all timescales tested (Fig. 4C). Remarkably, despite not

Figure 4: A quantitative model of short-term release dynamics captures patterned inputs. **(A)** Representative trace of kinetics fitted to an averaged electrically evoked GIRK current, and parameter distribution across the population (n=6 neurons). **(B)** Schematic depicting the linear-nonlinear short-term plasticity model and the plasticity kernel fitted using a least-squares approach to paired-pulse data. **(C)** Post-stimulus charge as a function of paired-pulse time interval, depicting experimental mean (black) and model predictions with fitted kernel and nonlinearity (blue). **(D)** Post-stimulus charge as a function of pulse-chase time interval, depicting experimental mean (black) and model predictions (blue). Note that model was fitted to paired-pulse, not pulse-chase dataset. **(E)** Simulations showing responses to 1Hz (black) and 20Hz (blue) presynaptic input, the model variables, and the resulting GIRK conductance.

being fitted on this data, the model described the meta-facilitation dynamics observed in the
pulse-chase experiments – in effect acting as a test set for the model (Fig. 4D). This likely
reflects the fact that the kernel has a short window of facilitation, which afterwards tends towards
depression if subsequent stimuli arrive too late (such as in the pulse-chase experiments with
extended time intervals; see Methods). To provide further intuition about this model, we
conducted simple simulations where presynaptic trains were delivered to single neurons, and the
model returned the kinetics and response magnitude of our earlier electrical stimulation
experiments (Fig. 4E). Together, these results show that the complex dynamics of 5-HT release
in the DRN can be faithfully captured by a SRP model.
363
Synaptic facilitation dynamics and the spatial structure of 5-HT release in DRN
We next explored spatial features of 5-HT release in e*x vivo* slices of the DRN using two-
photon imaging of the genetically encoded 5-HT sensor GRAB5-HT2M expressed selectively in
5-HT neurons (Fig. 5A). Bath application of 5-HT (50 µM) generated a large increase in $GRAB_{5-HT}$
fluorescence in both neurites and cell bodies (Fig. 5B), acting as a positive control for $GRAB_{5-HT}$
expression and overall function. We next electrically stimulated 5-HT connections in DRN
slices at various stimulation frequencies (5-20Hz; while fixing pulse number at 8, similar to
above; see Fig. 3 and Methods). We manually annotated responding regions and observed a
strikingly nonlinear increase in fluorescence response as electrical stimulation frequency
increased (Fig. 5C-D). $GRAB_{5-HT}$-mediated fluorescence was sensitive to TTX application,
demonstrating that electrically induced release events were action potential-driven (Fig. S11). By
imaging 5-HT release directly, these data present a separate line of evidence showing facilitation
dynamics of 5-HT release as revealed earlier by whole-cell recordings (Fig. 3), and additionally
support a presynaptic locus for this phenomenon.
As stimulation frequency increased, $GRAB_{5-HT}$ -mediated fluorescence increased in
absolute intensity and became progressively more localized to hotspots within the imaged frame
(Fig. 5E, left; Fig. 5G). To assess the reliability of the fluorescence signal, we computed the
pairwise pixel correlation across repeated trials, and found that the response became more
spatially reliable with higher stimulation frequencies (Fig. 5H), as expected for a probabilistic
and facilitating population of synapses. $GRAB_{5-HT}$ fluorescence displayed a weak overall
correlation with tdTomato signal, likely reflecting a lack of strong somatic signal for the
biosensor (Fig. 5I).
Our whole-frame analyses (Fig. 5E) suggested that $GRAB_{5-HT}$ fluorescence responses
were spatially restricted (*i.e.,* hotspots). To quantify the spatial features of these hotspots in a
principled manner, we next computed the spatial autocorrelation of the $GRAB_{5-HT}$ whole-frame
response for each stimulation frequency (Fig. 5F, left; see Methods). As frequency increased, the
spatial autocorrelation function became broader, providing a metric of hotspot size (Fig. 5F). To
illustrate the expected outcome of the autocorrelation function should no spatial organization of
the $GRAB_{5-HT}$ response be present, we performed spatial shuffling of the sensor response pixels
at 20Hz (Fig. 5E, right). This produced an autocorrelation function that sharply peaked in the
center (correlation when images are overlaid) (Fig. 5F, right). We constructed a null distribution
of the autocorrelation values arising from these shuffled 5-HT sensor response pixels for each
condition (Fig. S12A-E; see Methods). By comparing this shuffled distribution with the real
distribution of stimulation-evoked autocorrelation values (Fig. S12E), we were able to delineate
a mask of statistically significant autocorrelation values for each region and stimulation
condition (Fig. S12F). These masks, representing a metric of hotspot size of the $GRAB_{5-HT}$

Figure 5: Synaptic facilitation dynamics and spatial structure of release revealed by GRAB5-HT serotonin sensor imaging in DRN. **(A)** Experimental schematic demonstrating 2P GRAB5-HT imaging approach. **(B)** Bath application of 5-HT generates an increase in GRAB signal. *Left*: fluorescence max-projection images from an example field of view demonstrating tdTomato signal, baseline GRAB signal before 5-HT application, and GRAB signal during 5-HT application. *Right*: quantification of whole-frame GRAB fluorescence during timecourse of 5-HT washon. **(C)** GRAB fluorescence responses are frequency-dependent. *Left*: Example region showing max-projection tdTomato and GRAB fluorescence images, as well as manually identified responsive ROIs (dotted ovals and inset). *Right*: GRAB sensor responses to electrical stimulation at 5, 10 and 20Hz (shaded regions) for the ROIs depicted on the left. Thin traces represent individual trials and thick traces represent means across all trials. **(D)** Frequency dependence of GRAB responses across ROIs (n=39), calculated as response integral (left) or response maximum (right; paired t-test between 5 and 20Hz data). **(E)** Example whole-frame tdTomato fluorescence (left) and binned GRAB responses to 5, 10 and 20Hz electrical stimulation (8 pulses), including 20Hz shuffled pixel control (right). **(F)** Spatial autocorrelation plots for the frames shown in E. Autocorrelation values are normalized from 0 to 1 for presentation. **(G)** Left: Pixelwise cumulative distribution of GRAB responses at 5, 10 and 20Hz for example region shown in E (pixel response index is expressed as a z-score vs 5Hz data). Right: Same as left, but across all regions (n=11; Kolmogorov-Smirnov test for 5Hz vs 20Hz) **(H)** Mean pixelwise Pearson R correlation of GRAB response across trials of repeated electrical stimulation at 5, 10 and 20Hz (n=11 regions; paired T-test, 5Hz vs 20Hz). **(I)** Mean pixelwise Pearson R correlation of GRAB response during stimulation, and tdTomato fluorescence (n=11 regions; paired T-test, 5Hz vs 20Hz). **(J)** Spatial clustering of GRAB5-HT signal is frequency-dependent. Shown are spatial masks where autocorrelation of GRAB signal significantly exceeds autocorrelation on shuffled frames (p=0.0001), with all regions overlaid (n=11 regions). **(K)** The total area of spatial masks in J at 5, 10 and 20Hz (n=11 regions; paired T-test, 5Hz vs 20Hz).

400   fluorescence signal, showed greater area as the stimulation frequency increased (Fig. 5J-K).
401   Together, these data indicate that GRAB$_{5-HT}$ fluorescence exhibits spatially localized hotspots
402   that scale in both area and intensity with stimulation frequency, consistent with facilitating 5-HT
403   release.
404
405   *Computations performed by a recurrent inhibitory network under excitatory drive*
406   To identify the network computations in DRN that arise from recurrent connections
between 5-HT neurons, dominated by short-term facilitation, we constructed a network model
that combined our mathematical description of the short-term plasticity of 5-HT synapses (Fig.
4) with previously described excitability parameters for 5-HT neurons (Fig. 6A; Harkin et al.,
2023; see Methods). This model consisted of a population of leaky-integrate-and-fire 5-HT
neurons, with sparse recurrent connections imbued with the described short-term recurrent
plasticity rules. 5-HT neurons received background depolarizing Poisson input, and a subset of
5-HT neurons received input from long-range targets (see Methods).
414   First, we considered how recurrence shaped the basic input-output properties of the
network. Responses to sustained 5 Hz or 20 Hz LHb activity yielded three periods of the DRN
population response (Fig. 6B-C): an early phasic period (0-200 ms), followed by a relaxation to a
steady-state response (200-400 ms), and finally a post-stimulus period (up to 1000ms after the
end of the stimulus). The early period resulted from 5-HT neurons responding to direct LHb
excitation before 5-HT$_{1A}$R -mediated feedback inhibition could modulate the response.
Consequently, the early period output rate was a rectified linear function of the input frequency
(Fig. 6D). In contrast, the steady-state response was non-monotonic, displaying an inverted-U
shape which peaked at input frequencies around 5 Hz and progressively decreased with higher
LHb input frequencies (Fig. 6E). This peculiar non-monotonic function, essentially representing
excitation-driven net inhibition, critically depended on the potent short-term facilitation at 5-HT
synapses, as *in silico* removal of these short-term dynamics returned a linear, monotonic, input-
output function (Fig. 6E). In contrast, the early or post-stimulus period were not appreciably
affected by the removal of short-term recurrent dynamics (Fig. 6D-F). This unusual non-
monotonic relationship held across a wide range of network connectivity parameters, including
changes to recurrent connection probability between 5-HT neurons (Fig S13), and changes to the
fraction of 5-HT neurons innervated by long-range input (Fig. S14). The latter manipulation,
inspired by previous work describing anatomical heterogeneity within DRN (Zhou et al., 2017;
Ren et al., 2018; Dorocic et al., 2014, Weissbourd et al., 2014), unexpectedly demonstrated that
the shape and inversion point of this unique input-output relationship was preserved across a
wide range of input fractions, from 10% to 50% of 5-HT neurons (Fig. S14A-C). The recurrent
drive of the network scaled with the frequency of inputs, but not their precise innervation
fraction (Fig. S14D-E). Removing the short-term facilitation dynamics resulted in recurrent drive
tracking a combination of input frequency and input innervation (Fig. S14F-J). Together, these
simulations suggest that short-term facilitation dynamics uniquely bestows on 5-HT output the
ability to track input frequency while normalizing for differences in fine connectivity features.
Because the core of this computation emerged largely independently of fine details of
connectivity features, it may generalize to other long-range inputs to the DRN.
442   Next, we considered whether recurrent inhibition, operating over behavioral timescales,
443   could gate interaction between functionally separate submodules or cellular assemblies
444   previously reported within DRN (Ren et al., 2018; Huang et al., 2019; Paquelet et al., 2022). We
445   extended the formalism developed above by implementing two segregated submodules that

Figure 6: Habenula input to raphe generates a nonmonotonic network response through recurrent short-term plasticity. **(A)** Schematic of the LIF network model depicting direct excitation from LHb (contacting 25% of neurons) and recurrent inhibition between 5-HT neurons. **(B)** Results of a simulation where a 5Hz step input from LHb was delivered to a subset of 5-HT neurons. From top, depicted are the LHb input frequency; the mean firing rates of all 5-HT neurons; a raster plot depicting firing of individual 5-HT neurons over time (grey neurons represent those receiving direct LHb input; black neurons represent those not receiving direct LHb input; raster is limited to the first 2000 of 5000 5-HT neurons for clarity); representative simulated voltage traces of 10 5-HT neurons, each in a distinct color; and representative simulated GIRK conductances from recurrent inhibition of the same 5-HT neurons, each in a distinct color. **(C)** Results of a simulation where a 20Hz step input from LHb was delivered to a subset of 5-HT neurons. Same panels as described in B. **(D)** Input-output transformation for the transient (0-200ms) response phase of all 5-HT neurons in the network, with (black) and without (blue) STP rules at recurrent serotonergic connections. **(E)** Input-output transformation for the stationary (200-400ms) response phase of all 5-HT neurons, with (black) and without (blue) STP rules at recurrent serotonergic connections. **(F)** Input-output transformation for the post-response (0-1s after stimulus) phase of all 5-HT neurons, with (black) and without (blue) STP rules at recurrent serotonergic connections.

received excitatory inputs from either LHb (Module 1) or a second generic area termed A
(Module 2; Fig. 7A), and asked how input from LHb would disrupt ongoing A-driven activity.
Low-frequency (5Hz) LHb input did not appreciably modulate the activity of 5-HT neurons
receiving input from area A, as expected by the functional segregation of these submodules (Fig.
7B-C, left). However, high-intensity LHb drive (20 Hz) sharply reduced the activity of A-
recipient 5-HT neurons (Fig. 7C-D, right) despite these LHb inputs not directly contacting A-
recipient 5-HT neurons. These interactions between submodules again critically depended on the
short-term dynamics between 5-HT neurons, as demonstrated by *in silico* removal of these
dynamics (Fig. 7E-G). Thus, strong LHb inputs to DRN have the ability to suppress anatomically
separate submodules through recurrent inhibition, in essence implementing a winner-take-all
computation.
Further simulations revealed that this winner-take-all computation was not confined to a
narrow range of network connectivity parameters. Decreasing the recurrent connection
probability tenfold yielded qualitatively similar winner-take-all interactions (Fig. S15). We
additionally explored changes in the spatial profile of recurrent connectivity (local vs global),
revealing that winner-take-all dynamics were remarkably consistent across different spatial
connectivity profiles (Fig. S16). In additional simulations, we explored how spatial overlap of
LHb and A inputs within DRN affected winner-take-all dynamics (Fig. S17), demonstrating that
strong winner-take-all dynamics emerged even in cases with high input overlap, enhanced by the
presence of short-term plasticity at recurrent connections. Finally, to better understand how
competing, graded long-range inputs are processed, we quantified the strength of the winner-
take-all effect as the input frequency of both LHb and A was varied systematically. This
mapping indicated that the winner-take-all effect scaled with the magnitude of both LHb and A
over a variety of connectivity parameter choices (Fig. S16E; Fig. S17F), and this scaling
depended on the presence of short-term recurrent facilitation (Fig. S16G; Fig. S17I). In
summary, the winner-take-all computation between subregions of DRN is present independent of
fine connectivity details, and is aided by short-term facilitation of 5-HT release in the DRN.
We next explored the biological plausibility of these winner-take-all rules through
additional slice physiology experiments, where either habenula input or local recurrent
connections were optogenetically stimulated to gate spiking output. First, a sparse ChETA
expression paradigm was employed in DRN as described earlier (Fig. 2; Fig. S18A). In ChETA-
negative 5-HT neurons, optical stimulation elicited outward 5-HT$_{1A}$R currents (Fig. S18B). To
directly test whether recurrent connections could gate spiking output of (ChETA-negative) 5-HT
neurons, we examined the ability of high-frequency (20 Hz) optical stimulation of DRN ChETA-
expressing neurons to perturb the persistent spiking induced by direct current injection of 5-HT
neurons (Fig. S18C-7D). Consistent with predictions from our computational model,
photostimulation triggered a significant reduction in spiking (Fig. S18D-E) which was mediated
by 5-HT$_{1A}$Rs (Fig. S18F-L). These data indicate that within DRN, recurrent inhibition can allow
populations of 5-HT neurons to significantly shape spiking output of separate 5-HT populations.
To ensure that the plausibility of these results extend to activation of long-range synaptic input,
we found that high-frequency optical stimulation of LHb afferents in slice reliably inhibited
spiking output of 5-HT neurons through 5-HT1ARs (Fig. S19).
Together, these results support a model wherein short-term facilitation at recurrent 5-HT
connections generates highly nonlinear network computations. Long-range inputs lead to either
excitation at low frequencies, or protracted network suppression at high frequencies; winner-
take-all dynamics additionally dominate between functionally distinct submodules at high

Figure 7: Winner-take-all dynamics generated by recurrent short-term plasticity. **(A)** Simulations with LHb inputs of either 5Hz (left) and 20Hz (middle) were delivered to 5-HT neurons during ramp input from area A. Schematic of LIF network model with two inputs (right). **(B)** Mean responses of 5-HT neurons to 5Hz (left) and 20Hz (right) input with STP rules intact. **(C)** Raster plot showing 5-HT neuron responses to 5Hz (left) and 20Hz (right) input with STP rules intact. **(D)** Input-output function, for input A, with STP rules intact and coincident LHb input at 0, 5 or 20Hz. **(E)** Mean responses of 5-HT neurons to 5Hz (left) and 20Hz (right) input with STP rules removed. **(F)** Raster plot showing 5-HT neuron responses to 5Hz (left) and 20Hz (right) input with STP rules removed. **(G)** Input-output function, for input A, with STP rules removed and coincident LHb input at 0, 5 or 20Hz. **(H)** Schematic of winner-take-all computation with recurrently connected submodules. When one input crosses an STP threshold, output from that submodule dominates while other submodules are suppressed through recurrent inhibition (left and centre). When neither input crosses an STP threshold, outputs from both submodules persist (right).

492   frequencies. *In silico* manipulations revealed that the short-term dynamics at these recurrent
493   inhibitory connections does not only critically contribute to this core computation, but also to the
494   robustness of these computations to meta-parameter changes in network architecture - possibly
495   allowing these computations to be stable in the face of the heterogeneous connectivity
496   parameters across DRN (described in Fig. S7; Zhou et al., 2017).
497
Learned cue-reward associations are conditionally modulated by habenular drive to the DRN.
As a final step, we set out to identify a behavioral manifestation of the non-linear
computations triggered in the DRN by activating the habenulo-raphe pathway at distinct input
frequencies. An internal state of reward anticipation can lead to measurable motor outputs, such
as post-cue anticipatory licking in a classical condition task; these licking dynamics are known to
be modulated by 5-HT activity (Matias et al., 2017). We thus developed a head-fixed classical
conditioning task where we could optogenetically manipulate LHb inputs to DRN while
monitoring licking behavior (Fig. 8A-B, see Methods). Mice underwent water deprivation before
being trained on an auditory classical conditioning paradigm with three possible reward
outcomes (Fig. 8B; 0uL, 5uL, 10uL). Trained mice displayed robust cue-driven anticipatory
licking, a conditioned response which indicates expression of a learned stimulus-reward
association (Fig. 8C; Cohen et al., 2012; see Methods for training criteria).
We optogenetically stimulated the habenulo-raphe pathway during the post-cue period of
interleaved trials, at variable frequencies where physiology and modeling work strongly
indicated that direct excitation (low frequency stimulation) or recurrent inhibition (triggered by
high frequency stimulation) would dominate within DRN (see Methods). During trials with a
large reward outcome, low-frequency (5Hz) optogenetic stimulation had no effect on
anticipatory licking, while high-frequency (20Hz) photostimulation induced a striking and rapid
decrease in anticipatory licking which recovered rapidly after stimulation ceased (Fig. 8E-G). In
contrast, during trials with a small reward outcome, neither 20Hz or 5Hz photostimulation
modulated anticipatory licking (Fig. 8J; Fig. S20B-C); during trials with no reward outcome,
photostimulating at 5Hz after the auditory tone triggered a slight, but at times clearly noticeable,
increase in licking which was not seen during 20Hz photostimulation (Fig. 8K; Fig. S20D-E). As
controls for visual artefacts and viral expression, we photostimulated with yellow light in the
same cohort (Fig. 8G, yellow; Miyazaki et al., 2018; see Methods), or photostimulated with blue
light in a separate cohort with a control viral vector (Fig. 8G; green; see Methods), producing no
changes in reward anticipation in either case.
As serotonergic circuits can alter conditioned responses or reward consumption over
longer timescales (Matias et al., 2017; Zhong et al., 2017), we asked whether our post-cue
optogenetic stimulation had an effect during the subsequent reward period, finding that licking
during this post-reward period was not significantly modulated (Fig. 8H). We additionally asked
whether post-cue photostimulation drove longer-timescale changes in licking behavior occurring
throughout the session. Our analysis, consisting of comparing early trials with nonstimulated late
trials occurring during the optogenetic stimulation block, failed to reveal significant long-term
alterations in reward anticipation (Fig. 8I; see Methods) consistent with a lack of modulation of
the underlying reward association itself. There was a positive correlation between baseline
anticipatory licking and optogenetic effect size for high-frequency photostimulation (Fig. S20H),
suggesting a divisive rather than subtractive inhibitory effect of pathway activation on reward
anticipation. Finally, the effect size of 20Hz and 5Hz stimulation on anticipatory licking showed
no relationship (Fig. S20I). To rule out nonspecific stimulation-induced motor effects, we

Figure 8: Habenula input to raphe disrupts the expression of learned reward associations in a value- and frequency-dependent manner. **(A)** Schematic illustrating experimental setup (left) and confocal image of LHb injection site (right). **(B)** Schematic of head-fixed classical conditioning task. **(C)** Mice learned to lick in response to the small reward-predicting and large reward-predicting cues (left), and displayed elevated licking after reward delivery (right) (n=9 mice). **(D)** Schematic of trial structure for optogenetic manipulations. **(E)** Mean licking across all large reward trials for a single mouse depicting control trials (black) or 5Hz optogenetic stimulation trials (blue). Inset shows cumulative distribution of licking for all large-reward trials. **(F)** Mean licking across all large reward trials for a single mouse depicting control trials (black) or 20Hz optogenetic stimulation trials (blue). Inset shows cumulative distribution of licking for all large-reward trials. **(G-I)** Summary statistics for optogenetic manipulation during post-tone period of large-reward trials. **(G)** Change in conditioned licking response during tone presentation, between control and optogenetic stimulation trials during opto block (paired two-sided t-tests). **(H)** Mean reward licking change between control and optogenetic stimulation trials (paired two-sided t-tests). **(I)** Change in post-tone licking for nonstimulated trials between baseline block and optogenetic stimulation block. **(J)** Change in post-tone licking between control and optogenetic stimulation trials (small reward trials). **(K)** Change in post-tone licking between control and optogenetic stimulation trials (no reward trials).

photostimulated at high frequencies during the reward consumption phase instead of in the post-
cue period, observing no significant modulation of licking in these conditions (Fig. S20F-G).
Taken together, these results support a model whereby habenular inputs to the raphe can
contextually modulate the expression of a conditioned reward response in a temporally precise
manner, providing a putative behavioral manifestation of recurrent inhibitory dynamics in this
circuit.
544
Discussion
Here, using a combination of optogenetic, electrophysiological, computational and
behavioral approaches, we provide evidence that DRN is organized in an unsuspected recurrent
inhibitory network, characterized by a broad connectivity within this nucleus, and strongly
facilitating release. We outlined a core set of computations supported by this connectivity motif,
including excitation-driven inhibition of network response at high input frequencies, and a robust
winner-take-all computation where anatomically independent raphe subsystems can be
antagonistically coupled via recurrent inhibition. These results add to the reported role of 5-HT
in modulating cognitive processing, not direct appetitive or aversive proximal behaviors
(Fonseca et al., 2015; Miyazaki et al., 2018) and links a local circuit's network architecture, and
associated synaptic dynamics, to a core set of ethologically relevant computations.
We present several lines of experimental evidence supporting a connectivity model
whereby 5-HT neurons in the DRN are organized in a local recurrent inhibitory network. Our use
of sparse optogenetic stimulation experiments represents in effect an optogenetic equivalent of
paired recordings from synaptically-connected neurons, which have provided decisive functional
validation of the recurrent connectivity nature of several networks in the brain (Miles & Wong,
1986; Mason et al., 1991; Markram et al., 1997; Debanne et al., 2008). These experiments,
unambiguously showing functional connections between 5-HT neurons, supports a model which
is largely inconsistent with the widely and long-held idea that 5-HT neurons are regulated in an
autocrine fashion by dendritically-released 5-HT acting on so-called $5\text{-HT}_{1A}\text{R}$ somatodendritic
autoreceptors (Gozlan et al., 1983; Weissmann-Nanopoulos et al., 1985; Pineyro & Blier, 1999).
This autoreceptor hypothesis makes several experimental predictions which we directly tested
and failed to find support for. While our results do not formally allow to distinguish whether
interconnected 5-HT neurons communicate through dendro-dendritic or canonical axo-dendritic
functional contacts, or whether release of 5-HT is spatially confined or consists of more spatially
diffuse volume transmission (Maxwell et al., 1983; Ridet et al., 1994; Bunin & Wightmann,
1998), we nonetheless note that classic axonal release provides the most parsimonious and
credible model accounting for our results. While a subset of 5-HT dendrites contains 5-HT
containing vesicles (Descarries et al., 1982; Chazal and Ralston, 1987), 5-HT containing axon
terminals are nonetheless widespread in the DRN and are structurally indistinguishable from
those found elsewhere in the brain (Chazal and Ralston, 1987). Importantly, this view does not
preclude the presence of NMDAR-dependent dendritic release of 5-HT that may operate in
distinct physiological conditions (Colgan et al., 2012). Furthermore, the facilitation of the short-
term 5-HT release stands in stark contrast to the robustly depressing dopamine release dynamics
recently described in striatum (Liu et al., 2018, Liu et al., 2021).
Previous experimental and theoretical work has identified a number of possible
computational functions of network recurrence, including winner-take-all competition (Coultrip
et al., 1992), signal amplification and selectivity (Douglas et al., 1995) and probabilistic
inference (Echeveste et al., 2020). Our results show that the recurrent inhibitory dynamics lead to

a non-monotonic input-output relationship in the DRN by supporting a strong excitation-driven inhibition, where larger excitatory inputs generate greater network suppression. A natural question is how this computation compares with gain control associated with classical feedforward inhibition motif. A range of distinct gain control regimes have been described with feedforward inhibition such as subtractive and divisive inhibition (Ayaz & Chance, 2009) and even non-monotonic inhibition (Mejias et al., 2014; Lewis et al., 2007). However, the recurrent inhibition motif described here is unique in that the phasic network response is dominated by direct excitation and linearly reports input intensity, while the delayed steady-state response exhibits non-monotonic behavior with increasing excitation. In principle, these temporally distinguishable epochs of 5-HT output dynamics could support multiplexed codes such as valence encoding over multiple timescales (Cohen et al., 2015). Our network model additionally uncovered an emergent winner-take-all-like computation in the DRN, where strong activation of connected modules of 5-HT neurons can inhibit activity of others. This builds on previous work describing anatomically segregated DRN subdomains (Ren et al., 2018), and implies that a winner-take-all computation governs state selection between submodules. While these simulations uncovered general algorithmic principles which were tested with a range of connectivity profiles in our network models, more precise quantitative information on the recurrent connectivity between submodules (Ren et al., 2018) or on the additional recruitment of feedforward GABAergic inhibition by some inputs (Zhou et al., 2017; Geddes et al., 2016) should bring about refinement of these general principles.

More broadly, our behavioral results provide evidence that LHb inputs to the DRN can rapidly and reversibly alter behavioral state, which is consistent with previous findings on the habenulo-raphe pathway in other model organisms (Amo et al., 2014; Marquez et al., 2020) and may represent an important role for this pathway in dynamic state selection. Indeed, cue-reward learning has been shown to lead to a gradual increase in the population activity of 5-HT neurons during reward anticipation, peaking around reward delivery (Zhong et al., 2017), and current theories postulate that 5-HT neurons represent valence information during associative learning (Liu et al., 2017). Thus, the DRN microcircuit has many of the core ingredients necessary to perform such contextual behavioral state transitions.

Animals often exhibit sharp thresholds when transitioning between ethologically relevant behavioral states such as exploitation and exploration (Marques et al., 2020). These state transitions require brain circuits which continually evaluate graded representations of either internal or external cues, enacting thresholding operations on behavioral timescales. Building on recent theories (Liu et al., 2020; Luo et al., 2016; Cohen et al., 2015; Marques et al., 2020), the winner-take-all computation between ensembles of recurrently connected 5-HT neurons, operating over protracted timescales, satisfies many of the conditions required for the largely exclusive, non-overlapping implementation of distinct naturalistic behaviors (Ren et al., 2018).

Acknowledgements
This work was supported by grants from the Canadian Institute for Health Research (JCB; RN), the uOttawa Brain and Mind Institute and the Canadian Heart and Stroke Foundation (JCB), Brain Canada (JCB), the Krembil Foundation (JCB) and the Canadian Foundation for Innovation (JCB). We thank members of the Béïque, Naud and Maler lab for helpful discussions. We would also like to thank Drs Salah El Mestikawy and Sylvain Williams for providing us with the SERT-Cre mouse line.

Author contributions

Conceptualization, M.B.L., S.G., S.H-D., R.N., and J.-C.B.; Methodology, M.B.L., S.G. ; Software, M.B.L.; Formal Analysis, M.B.L.; Investigation, M.B.L., S.G., M.C., E.H., S.M., E.H.-G., ; Resources, M.B.L., E.H. and R.N. ; Writing – Original Draft, M.B.L.; Writing – Review & Editing, M.B.L., R.N., and J.-C.B.; Supervision, J.-C.B.; Funding Acquisition, J.-C.B.

Declaration of interests
The authors declare no competing interests.

Data availability statement
The datasets and code generated during this study are available from the corresponding author upon reasonable request.

References

Abbot, L.F. & Regehr, W.G. (2004). Synaptic computation. Nature 431:796-803.

Amo, R., Fredes, F., Kinoshita, M., Aoki, R., Aizawa, H., Agetsuma, M., Aoki, T., Shiraki, T., Kakinuma, H., Matsuda, M., Yamazaki, M., Takahoko, M., Tsuboi, T., Higashijima, S.-I., Miyasaka, N., Koide, T., Yabuki, Y., Yoshihara, Y., Fukai, T. & Okamoto, H. (2014) The habenulo-raphe serotonergic circuit encodes an aversive expectation value essential for adaptive active avoidance of danger. Neuron 84:1034-1048.

Andalman, A.S., Burns, V.M., Lovett-Baron, M., Broxton, M., Poole, B., Yang, S.J., Grosenick, L., Lerner, T.N., Chen, R., Benster, T., Mourrain, P., Levoy, M., Rajan, K. & Deisseroth, K. (2019). Neuronal Dynamics Regulating Brain and Behavioral State Transitions. Cell 177:970-985.

Andrade, R., Malenka, R.C. & Nicoll, R.A. (1986). A G protein couples 5-HT and GABAB receptors to the same channels in hippocampus. Science 234:1261-1265.

Ayaz, A. & Chance, F.S. (2009). Gain modulation of neuronal responses by subtractive and divisive mechanisms of inhibition. J Physiol 101:958-968.

Broadhurst, P.L. (1957). Emotionality and the Yerkes-Dodson Law. J Exp Psychol 54:345-352.

Bromberg-Martin, E.S. & Hikosaka, O. (2011). Lateral habenula neurons signal errors in the prediction of reward information. Nat Neurosci 14:1209-1216.

Bunin, M.A. & Wightman, R.M. (1998). Quantitative evaluation of 5-hydroxytryptamine (5-HT) neuronal release and reuptake: an investigation of extrasynaptic transmission. J Neurosci 18:4854-4860.

Campagnola, L., Seeman, S.C., Chartrand, T., Kim, L., Hoggarth, A., Gamlin, C., Ito, S., Trinh, J., Davoudian, P., Radaelli, C., et al., (2022). Local connectivity and synaptic dynamics in mouse and human neocortex. Science 375: eabj5861. DOI: 10.1126/science.abj5861.

676 Celada, P., Puig, V.M., Casanovas, J.M., Guillazo, G. & Artigas, F. (2001). Control of dorsal
677 raphe serotonergic neurons by the medial prefrontal cortex: Involvement of serotonin-1A,
678 GABAA, and glutamate receptors. J Neurosci 21:9917-9929.
679
680 Chazal, G. & Ralston, H.J. (1987). Serotonin-containing structures in the nucleus raphe dorsalis
681 of the cat: An ultrastructural analysis of dendrites, presynaptic dendrites, and axon terminals. J
682 Comp Neurol 259:317-329.
683
684 Cohen, J.Y., Amoroso, M.W. & Uchida, N. (2015). Serotonergic neurons signal reward and
685 punishment on multiple timescales. eLife 2015;4:e06346.
686
687 Cohen, J.Y., Haesler, S., Vong, L., Lowell, B.B. & Uchida, N. (2012). Neuron-type-specific
688 signals for reward and punishment in the ventral tegmental area. Nature 482:85-88.
689
Coffey, K.R., Marx, R.G., Vo, E.K., Nair, S.G. & Neumaier, J.F. (2020). Chemogenetic
inhibition of lateral habenula projections to the dorsal raphe nucleus reduces passive coping and
perseverative reward seeking in rats. Neuropsychopharm 45:1115-1124.
Colgan, L.A., Cavolo, S.L., Commons, K.G. & Levitan, E.S. (2012). Action potential-
independent and pharmacologically unique vesicular serotonin release from dendrites. J Neurosci
32:15737-15746.
696
697 Coultrip, R., Granger, R. & Lynch, G. (1992). A cortical model of winner-take-all competition
698 via lateral inhibition. Neural Networks 5:47-54.
699
700 Courtney, N.A. & Ford, C.P. (2016). Mechanisms of 5-HT$_{1A}$ receptor-mediated transmission in
701 dorsal raphe 5-HT neurons. J Physiol 594:953-965.
702
703 Cragg, S. (2003). Variable dopamine release probability and short-term plasticity between
704 functional domains of the primate striatum. J Neurosci 23:4378-4385.
705
706 Debanne, D., Boudkkaszi, S., Campanac, E., Cudmore, R.H., Giraud, P., Fronzaroli-Molinieres,
707 L., Carlier, E. & Caillard, O. (2008). Paired-recordings from synaptically coupled cortical and
708 hippocampal neurons in acute and cultured brain slices. Nat Protocols 3:1559-1568.
709
710 de Hoz, L. & Nelken, I. (2014). Frequency tuning in the behaving mouse: Different bandwidths
711 for discrimination and generalization. PLOS One 9:e91676.
712
713 Denève. S. & Machens, C.K. (2016). Efficient codes and balanced networks. Nat Neurosci
714 19:375-382.
715
716 Descarries, L., Watkins, K.C., Garcia, S. & Beaudet, A. (1982). The serotonin neurons in nucleus
717 raphe dorsalis of adult rat: A light and electron microscope radioautographic study. J Comp
718 Neurol 207:239-254.
719

Dolzani, S.D., Baratta, M.V., Amat, J., Agster, K.L., Saddoris, M.P., Watkins, L.R. & Maier, S.F. (2016). Activation of a habenulo-raphe circuit is critical for the behavioral and neurochemical consequences of uncontrollable stress in the male rat. eNeuro 3:e0229-16.2016.

Dorocic, I.P., Futh, D., Xuan, Y., Johansson, Y., Pozzi, L., Silberberg, G., Carlen, M. & Meletis, K. (2014). A Whole-brain atlas of inputs to serotonergic neurons of the dorsal and median raphe nuclei. Neuron 83:663-678.

Douglas, R.J., Koch, C., Mahowald, M., Martin, K.A.C. & Suarez, H.H. (1995). Recurrent excitation in neocortical circuits. Science 269:981-985.

Echeveste, R., Aitchison, L., Hennequin, G. & Lengyel, M. (2020). Cortical-like dynamics in recurrent circuits optimized for sampling-based probabilistic inference. Nat Neurosci 23:1138-1149.

Eriksson, A., Raczkowska, M., Navawongse, R., Choudhury, R., Stewart, J.C., Tang, Y.L., Wang, Z., Claridge-Chang, A. (2017). Neuromodulatory circuit effects on Drosophila feeding behaviour and metabolism. Sci Rep 7:8839.

Geddes, S.D., Assadzada, S., Lemelin, D., Sokolovski, A., Bergeron, R., Haj-Dahmane, S. & Beique, J.-C. (2016). Target-specific modulation of the descending prefrontal cortex inputs to the dorsal raphe nucleus by cannabinoids. PNAS 113:5429-5434.

Gozlan, H., El Mestikawy, S., Pichat, L., Glowinski, J. & Hamon, M. (1983). Identification of presynaptic 5-HT autoreceptors using a new ligand: 3H-PAT. Nature 305:140-142.

Graeff, F.G., Guimaraes, F.S., de Andrade, T.G.C.S. & Deakin, J.F.W. (1996). Role of 5-HT in Stress, Anxiety and Depression. Pharmacol Biochem Behav 54:129-141.

Groos, D. & Helmchen, F. (2024). The lateral habenula: a hub for value-guided behavior. Cell Rep 43:113968.

Grossman, C.D., Bari, B.A. & Cohen, J.Y. (2022). Serotonin neurons modulate learning rate through uncertainty. Curr Biol 32:1-14.

Harkin, E.F., Lynn, M.B., Payeur, A., Boucher, J.-F., Caya-Bissonnette, L., Cyr, D., Stewart, C., Longtin, A., Naud, R. & Béïque, J.-C. (2023). Temporal derivative computation in the dorsal raphe network revealed by an experimentally driven augmented integrate-and-fire modeling framework. eLife 12:e72951.

Huang, K.W., Ochandarena, N.E., Philson, A.C., Hyun, M., Birnbaum, J.E., Cicconet, M. & Sabatini, B.L. (2019). Molecular and anatomical organization of the dorsal raphe nucleus. eLife 8:e46464.

Jackman, S.L. & Regehr, W.G. (2017). The mechanisms and functions of synaptic facilitation. Neuron 94:447-464.

Jones, E., Oliphant, E., Peterson, P., et al. (2001). SciPy: open source scientific tools for Python.
http://www.scipy.org.

Lammel, S., Lim, B.K., Ran, C., Huang, K.W., Betley, M.J., Tye, K.M., Deisseroth, K. &
Malenka, R.C. (2012). Input-specific control of reward and aversion in the ventral tegmental
area. Nature 491:212-217.

Lecca, S., Namboodiri, V.M.K., Restivo, L., Gervasi, N., Pillola, G., Stuber, G.D. & Mameli, M.
(2020). Heterogeneous habenular neuronal ensembles during selection of defensive behaviors.
Cell Rep 31:107752.

Lewis, J.E., Lindner, B., Laliberté, B. & Groothuis, S. (2007). Control of neuronal firing by
dynamic parallel fiber feedback: Implications for electrosensory reafference suppression. J Exp
Biol 210:4437-4447.

Li, Y., Zhong, W., Wang, D., Feng, Q., Li, Z., Zhou, J., Jia, C., Hu, F., Zeng, J., Guo, Q., Fu, L.
& Luo, Minmin. (2016). 5-HT neurons in the dorsal raphe nucleus encode reward signals. Nat
Comms 7:10503.

Liu, C., Goel, P. & Kaeser, P.S. (2021). Spatial and temporal scales of dopamine transmission.
Nat Neurosci 22:345-358.

Liu, C., Kershberg, L., Wang, J., Schneeberger, S. & Kaeser, P. (2018). Dopamine secretion is
mediated by sparse active zone-like release sites. Cell 172:706-718.

Liu, Z., Lin, R. & Luo, M. (2020). Reward contributions to serotonergic function. Annu Rev
Neurosci 43:141-162.

Luo, M., Li, Y. & Zhong, W. (2016). Do dorsal raphe 5-HT neurons encode "beneficialness"?
Neurobiol Learn Mem 135:40-49.

Lynn, M.B., Lee, K.F.H., Soares, C., Naud, R. & Béïque, J.-C. (2020). A synthetic likelihood
solution to the silent synapse estimation problem. Cell Rep 32:107916.

Markram, H., Lubke, J., Frotscher, M. & Sakmann, B. Regulation of synaptic efficacy by
coincidence of postsynaptic APs and EPSPs. (1997). Science 275:213-215.

Marquez, J.C., Li, M., Schaak, D., Robson, D.N. & Li, J.M. (2020). Internal state dynamics
shape brainwide activity and foraging behavior. Nature 577:239-243.

Mason, A., Nicoll, A. & Stratford, K. (1991). Synaptic transmission between individual
pyramidal neurons of the rat visual cortex *in vitro*. J Neurosci 11:72:84.

Matias S., Lottem, E., Dugué, G.P. & Mainen, Z.F. (2017). Activity patterns of serotonin
neurons underlying cognitive flexibility. eLife 6:e20552.

Matsumoto, M. & Hikosaka, O. (2009). Representation of negative motivational value in the
primate lateral habenula. Nat Neurosci 12:77-84.

Maxwell, D.J., Leranth, C. & Verofstad, A.A.J. (1983). Fine structure of 5-HT-containing axons
in the marginal zone of the rat spinal cord. Brain Res 266:253-259.

Mejias, J.F., Payeur, A., Selin, E., Maler, L. & Longtin, A. (2014). Subtractive, divisive and non-
monotonic gain control in feedforward nets linearized by noise and delays. Front Comp
Neurosci, doi: https://doi.org/10.3389/fncom.2014.00019.

Miles, R. & Wong, R.K.S. (1986). Excitatory synaptic interactions between CA3 neurones in the
guinea-pig hippocampus. J Physiol 373:397-418.

Miyazaki, K., Miyazaki, K.W., Sivori, G., Yamanaka, A., Tanaka, K.F. & Doya, K. (2020).
Serotonergic projections to the orbitofrontal and medial prefrontal cortices differentially
modulate waiting for future rewards. Sci Adv 6:eabc7246.

Miyazaki, K., Miyazaki, K.W., Yamanaka, A., Tokuda, T., Tanaka, K.F. & Doya, K. (2018).
Reward probability and timing uncertainty alter the effect of dorsal raphe 5-HT neurons on
patience. Nat Comms 9:2048.

Miyazaki, K.W., Miyazaki, K., Tanaka, K.F., Yamanaka, A., Takahashi, A., Tabuchi, S. &
Doya, K. (2014). Optogenetic activation of dorsal raphe 5-HT neurons enhances patience for
future rewards. Curr Biol 24:2033-2040.

Newberry, N.R. & Nicoll, R.A.A. (1984). A bicuculline-resistant inhibitory post-synaptic
potential in rat hippocampal pyramidal cells in vitro. J Physiol 348:239-254.

Paquelet, G.E., Carrion, K., Lacefield, C.O., Zhou, P., Hen, R. & Miller, B.R. (2022). Single-cell
activity and network properties of dorsal raphe nucleus serotonin neurons during emotionally
salient behaviors. Neuron, doi: 10.1016/j.neuron.2022.05.015.

Pazos, A. & Palacios, J.M. (1985). Quantitative autoradiographic mapping of serotonin receptors
in the rat brain. I. Serotonin-1 receptors. Brain Res 346:205-230.

Pineyro, G. & Blier, P. (1999). Autoregulation of 5-HT neurons: role in antidepressant drug
action. Pharmacol Rev 51:533-591.

Pozzorini, C., Mensi, S., Hagens, O., Naud, R., Koch, C. & Gerstner, W. (2015). Automated
high-throughput characterization of single neurons by means of simplified spiking neurons.
PLAS Comp Biol 11:e1004275.

Proulx C.D., Hikosaka, O. & Malinow R. (2014). Reward processing by the lateral habenula in
normal and depressive behaviors. Nat Rev Neurosci 17:1146-1152.

Ren, J., Friedmann, D., Xiong, J., Huguenard, J., Horowitz, M.A. & Luo, L. (2018).
Anatomically defined and functionally distinct dorsal raphe 5-HT sub-systems. Cell 175:472-
487.
861
862 Ridet, J.-L., Tamir, H. & Privat, A. (1994). Direct immunocytochemical localization of 5-
863 hydroxytryptamine receptors in the adult rat spinal cord: A light and electron microscopic study
864 using anti-idiotypic antiserum. J Neurosci Res 38:109-121.
865
866 Rossbroich, J., Trotter, D., Beninger, J., Toth, K. & Naud, R. (2021). Linear-nonlinear cascades
867 capture synaptic dynamics. PLOS Comp Bio 17(3): e1008013.
868
869 Salehi, B., Cordero, M.I. & Sandi, C. (2010). Learning under stress: the inverted-U-shape
870 function revisited. Learn Memory 17:522-530.
871
872 Seo, Changwoo., Guru, A., Jin, M., Ito, B., Sleezer, B.J., Ho, Y.-Y., Wang, E., Boada, C., Krupa,
873 N.A., Kullakanda, D.S., Shen, X.C., Warden, M.R. (2019). Intense threat switches dorsal raphe
874 5-HT neurons to a paradoxical operation mode. Science 363:538-542.
875
876 Stephens, D.W. & Krebs, J.R. (1986). Foraging Theory. Princeton: Princeton University Press.
877
878 Stimberg, M., Brette, R. & Goodman, D.F.M. (2019). Brian 2, an intuitive and efficient neural
879 simulator. eLife 2019;8:e47314.
880
881 Takahashi, A., Lee, R.X., Iwasato, T., Itohara, S., Arima, H., Bettler, B., Miczek, K.A. & Koide,
882 T. (2015). Glutamate input in the dorsal raphe nucleus as a determinant of escalated aggression
883 in male mice. J Neurosci 35:6452-6436.
884
885 Trinh, A.-T., Harvey-Girard, E., Teixeira, F. & Maler, L. (2016). Cryptic laminar and columnar
886 organization in the dorsolateral pallium of a weakly electric fish. J Comp Neurol 524:408–428.
887
888 Vandermaelen, C.P. & Aghajanian, G.K. (1983). Electrophysiological and pharmacological
889 characterization of serotonergic dorsal raphe neurons recorded extracellularly and intracellularly
890 in rat brain slices. Brain Res 289:109-119.
891
892 van der Walt, S., Colbert, S.C., and Varoquaux, G. (2011). The NumPy array: a structure for
893 efficient numerical computation. Comput Sci Eng 13, 22–30.
894
895 Warden, M.R., Selimbeyoglu, A., Mirzabekov, J.J., Lo, M., Thompson, K.R., Kim, S.-Y.,
896 Adhikari, A., Tye, K.M., Frank, L.M. & Deisseroth, K.D. (2012). A prefrontal cortex-brainstem
897 neuronal projection that controls response to behavioral challenge. Nature 492:428-432.
898
899 Weissbourd, B., Ren, J., DeLoach, K.E., Guenthner, C.J., Miyamichi, K. & Luo, L. (2014).
900 Presynaptic partners of dorsal raphe serotonergic and GABAergic neurons. Neuron 83:645-662.
901

Weissmann-Nanopoulos, D., Mach, E., Magre, J., Demassey, Y. & Pujol, J. F. (1985). Evidence
for the localization of 5HT1A binding sites on 5-HT containing neurons in the raphe dorsalis and
raphe centralis nuclei of the rat brain. Neurochem Int 7:1061-1072.

Yang, Y., Cui, Y., Sang, K., Dong, Y., Ni, Z., Ma, S. & Hu, H. (2018). Ketamine blocks bursting
in the lateral habenula to rapidly relieve depression. Nature 554:317-322.

Zhang, Y.-P. & Oertner, T.G. (2007). Optical induction of synaptic plasticity using a light-
sensitive channel. Nat Methods 4:139-141.

Zhong, W., Li, Y., Feng, Q., Luo, M. (2017). Learning and stress shape the reward response
patterns of 5-HT neurons. J Neurosci 37:8863-8875.

Zhou, L., Liu, M.-Z., Deng, J., Mu, D. & Sun, Y.-G. (2017). Organization of functional long-
range circuits controlling the activity of serotonergic neurons in the dorsal raphe nucleus. Cell
Rep 18:3018-3032.

Methods

Viral Constructs and Retrograde Tracer

Construct	Source	Titre (IU/µl)
AAV9.CAG.hChR2(H134R)-mCherry.WPRE.SV40	*(Penn Vector Core, AV-9-20938M)*	*~10^12*
AAV9.Ef1a.DIO.ChETA(E123T/H134R)-eYFP.WPRE.hGH	*(Penn Vector Core, AV-9-26968P)*	*~10^12*
rAAV2-retro-hSyn1-cre	*(Janelia Farm Research Campus)*	*~10^13*
AAV2/9.emCBA.eGFP	*Canadian Neuorphotonics Platform, Université Laval*	*~10^13*
AAV-EF1a-DIO-5-HT2M (Serotype: 2/9)	*Canadian Neuorphotonics Platform, Université Laval*	*~10^12*

Animal use

All experiments and procedures were performed in accordance with approved procedures and guidelines set forth by the University of Ottawa Animal Care and Veterinary Services. For this study, SERT-Cre mice (B6.Cg-Tg(Slc6a4- cre)ET33Gsat/Mmcd) were crossed with the R26R line (Rosa26R-tdTomato fluorescent protein; tdTom) to generate a SERT-Cre::Rosa-tdTomato mouse line (referred to in the manuscript as SERT-Cre::tdTom), and SOM-Cre mice tm2.1(cre)Zjh/J) were crossed with the R26R line to generate a SOM-Cre::Rosa-tdTom mouse line (referred to in the manuscript as SOM-Cre::tdTom). Mice were kept on a 12:12 hour light/dark cycle, with access to food and water ad libitum unless otherwise noted. A mix of male and female SERT-Cre::tdTom and SOM-Cre::tdTom mice (1-3 months for slice physiology experiments, 2-6 months for behavior experiments) were used. A mix of male and female SOM-Cre::Rosa-tdTom mice were used (1-3 months for slice physiology experiments).

For the behavior portion of the study, mice with a C57BL/6 background were utilized, either originating as mice with a wild-type genotype originating from the SERT-Cre:: tdTomato line, or as mice purchased from Charles River Labs. Mice were kept on a 12:12 hour reverse light/dark cycle and were water restricted to 85-90% of their initial weight during behavioral testing.

Slice physiology and anatomy

Stereotaxic Injections

All experiments and procedures were performed in accordance with approved procedures and guidelines set forth by the University of Ottawa Animal Care and Veterinary Services. For stereotaxic injections, mice were injected with 0.05 mg/kg buprenorphine and anesthetized by inhalation of isoflurane. Injections were performed using a 10 µl Hamilton syringe with a 33 gauge needle, or with a Nanoject III programmable injector with a pulled glass micropipette. 0.4 - 0.5 µl was injected per coordinate, injected at a rate of 0.1 – 0.2 ul per minute.

Stereotaxic co-ordinates are as follows: LHb (from Bregma skull landmark, in mm: AP -1.7, ML +/- 0.4, DV -3.0) and DRN (from Lambda skull landmark, in mm, 20 degree angle: AP -0.5, ML +1.0, DV -3.0; or 0 degree angle: AP -0.5, ML +0, DV-3.2, -3, -2.8, sequential injections to tile DV extent of DRN with total volume split between all 3 coordinates. Both injection types were utilized). For retrograde tracing, 0.4 µl of rAAV-retro-TdTomato was injected. For ChR2 (H134R) expression in LHb, 0.4 µl of AAV9.CAG.hChR2(H134R)-mCherry.WPRE.SV40 (titre ~10^{12} GC/mL) was injected bilaterally and animals were left to recover for 3 to 4 weeks prior to ex-vivo electrophysiological recordings. In anterograde tracing, AAV-GFP was injected in the LHb and animals were perfused 2-3 weeks post-injections.

For experiments requiring sparsely labeled cells in the DRN with ChETA-YFP, we further diluted virus in a 20 % mannitol/saline solution at a ratio of 2:1 (mannitol solution: virus solution). In these experiments, the virus solution was AAV9.Ef1a.DIO.ChETA(E123T/H134R)-eYFP.WPRE.hGH. The mixed mannitol and virus solution was then injected in the DRN at the coordinates above.

For experiments involving expression of ChETA in SOM interneurons in DRN, utilizing the SOM-Cre::Rosa-tdTom mouse line, we injected a total of 0.5uL of AAV9.Ef1a.DIO.ChETA(E123T/H134R)-eYFP.WPRE.hGH at the DRN coordinates above.

Slice preparation
LHb and DRN containing brain slices were prepared from 8 to 10 week old mice (3 to 4 weeks post-stereotaxic injections). For electrical stimulation experiments, brainstem slices were prepared from P21-P30 mice. Acute slices were prepared similarly to previously described methods (Geddes et al., 2016). In brief, mice were anesthetized by isoflurane inhalation (Baxter Corporation, Canada) and sacrificed by decapitation. The brain was removed, and brain slices were sectioned while immersed in ice-cold choline chloride-based cutting solution of the following composition (in mM): 119 choline-Cl, 2.5 KCl, 1 $CaCl_2$, 4.3 $MgSO_4$-$7H_2O$, 1 NaH_2PO_4, 1.3 sodium L-ascorbate, 26.2 $NaHCO_3$, and 11 glucose, and equilibrated with 95% O_2, 5% CO_2. Slices were recovered in a chamber containing standard Ringer's solution of the following composition (in mM): 119 NaCl, 2.5 $CaCl_2$, 1.3 $MgSO_4$-$7H_2O$, 1 NaH_2PO_4, 26.2 $NaHCO_3$, and 11 glucose, at a temperature of 37°C, continuously bubbled with a mixture of 95% O_2, 5% CO_2. Slices were left to recover for 1 hour in a recovery chamber and equilibrate to a temperature of approximately 21°C until recordings were performed.

Whole-Cell Electrophysiology
DRN neurons were visualized using an upright microscope: (1) Examiner D1 equipped with Dodt-contrast or differential-interference contrast (DIC) (Zeiss, Oberkochen, Germany; 40x/0.75NA objective) or, (2) BX51W1 equipped with DIC optics (Olympus, Center Valley, PA; 40x/0.80NA objective). 5-HT neurons were visually identified by TdTomato signal driven by the SERT promoter under EPI-fluorescence illumination (xCite Series 120pc). Whole-cell recordings were carried out using an Axon Multiclamp 700B amplifier and electrical signals were low-pass filtered at 2 kHz and sampled at 10 kHz with an Axon Digidata 1440A (or 1550) digitizer. Whole-cell recordings were performed using borosilicate glass patch electrodes (3-6 MΩ; World Precision Instruments, Florida) pulled on a Narishige PC-10 pipette puller (Narishige, Japan). Slices were constantly bathed in Ringer solution containing (in mM): 119 NaCl, 2.5 KCl, 1.3 $MgSO_4$, 2.5 $CaCl_2$, 1.0 NaH_2PO_4, 11 glucose, 26.2 $NaHCO_3$, and 0.1 L-Tryptophan saturated with 95% O_2 and 5% CO_2 (pH = 7.3; 295-310 mOsm/L). In the

experiments described in Fig. S4 (exploration of feedforward inhibition) and Fig. S10 (electrical stimulation of 5-HT afferents, and optogenetic stimulation of long-range LHB input to 5-HT neurons), the Ringer solution was identical to that described above except that 1.5mM $CaCl_2$ was used in place of 2.5mM $CaCl_2$ (as described in the caption).

Three intracellular recording solutions were used. For voltage-clamp experiments involving solely excitatory post-synaptic currents (EPSCs) recordings, we used an intracellular solution of the following composition: 115 mM cesium methane-sulfonate, 5 mM tetraethylammonium-Cl, 10 mM sodium phosphocreatine, 20 mM HEPES, 2.8 mM NaCl, 5 mM QX-314, 0.4 mM EGTA, 3 mM ATP(Mg2+), and 0.5 mM GTP, pH 7.25 (adjusted with CsOH; osmolarity, 280–290 mOsmol/L). For recordings involving 5HT1AR-mediated current or potential, we used the either of the following internal solutions: 1) 115 mM potassium gluconate, 20 mM KCl, 10 mM sodium phosphocreatine, 10 mM HEPES, 4 mM ATP(Mg2+), and 0.5 mM GTP, pH 7.25 (adjusted with KOH; osmolarity, 280–290 mOsmol/L): or 2) 115 mM KMeSO4; 5 mM KCl; 5 mM NaCl; 1 mM MgCl2; 0.020 mM EGTA; 10mM HEPES; 4 mM ATP; 0.4 mM GTP; and 10mM Na2phoshocreatine, pH7.4 (adjusted with KOH; osmolarity 280-290   mOsmol/L). For all voltage clamp recordings, access resistance was continuously monitored using a 125 ms, 2 mV hyperpolarizing pulse, at least 245 ms prior to electrical or optogenetic stimulation. Recordings were discarded when the access resistance changed by >30%. Liquid junction potential was not compensated for.

The slice physiology experiments described in Figures 1 and 2 were conducted at room temperature (20-24C). The electrical stimulation experiments described in Figures 3 and 4, as well as Figures S4 (exploration of feedforward inhibition) and S10 (replication of main results of Figure 3), were conducted nominally at 30-32°C through the use of an in-line bath heater and stage heater.

LED Photostimulation
LED photostimulation was delivered using a PlexBright LED module and controller (465 nm; Plexon, Texas). Light was delivered through a 200 µm patch fiber cable with a bare fiber tip with ~ 1 cm of glass exposed. This fiber was held at a ~45 degree angle to the slice, with the tip at a distance of approximately 1cm from the surface of the acute brain slice. The tip of the fiber emitted light in a narrow, approximately conical shape with a measured irradiance of 792 mW/mm^2 at the slice. The fiber's position was manually calibrated such that all of the emitted light was hitting the slice, focused on the portion of the slice where the physiologically probed neuron was. For EPSC/EPSP stimulation, unless otherwise stated, optically evoked-PSCs were stimulated at a frequency of 0.1 or 0.067 Hz with light pulses of 1-3 ms in length and an irradiance of 792 mW/mm2 measured. In driving 5-HT1AR-mediated currents, optical stimulation was accomplished using 10 ms pulses at 20 Hz for 400 ms.

Electrical stimulation of 5-HT afferents
Electrical stimuli were delivered through a bipolar concentric stimulating electrode located >100 um from the soma of the recorded neuron. Stimuli were 0.1 ms in duration, and consisted of frequencies provided in the text. For all electrical stimulation experiments, we bath-applied a cocktail of synaptic blockers to isolate the 5-HT1AR-mediated responses (i.e., 10 µM NBQX, 100 µM picrotoxin, 200 nM CGP 55845). All 5-HT electrical stimulation experiments were performed at near-physiological temperature (30-32 °C).

Immunohistochemistry

Mice were perfused using a minimally invasive method developed by Kenneth D. Eichenbaum and colleague in 2005. Briefly, mice were put under general anaesthesia with 1.5% isoflurane. The animal was then placed on its back for injection (1 mL PFA4% in 0.1 M PBS). The injection was done with a 1 mL syringes equipped with a 26G needle over a 30 seconds course. The correct position of the needle was confirmed by the appearance of blood in the needle hub.   Brains were then put in 10 mL of PFA4% overnight and then transferred in a 15 mL tube with 10 mL of PBS (0.1 M PBS, 0.1% Sodium azide). The fixed brains were then cut in 50 um thick slices with a vibrotome (Vibrotome,1500 sectioning system).

Free-floating sections were blocked in 1% BSA + 0.03% Triton X-100 (TX-100) for 45-60 min, and incubated in primary antibody in blocking solution at 4 °C. On the following morning, slices were washed three times, for 5 min each time, with 0.1% Tx-100 before incubation in the secondary antibody goat anti-rabbit–conjugated Alexa-647 (Life technologies, A21245) for 2 h at room temperature. Slices were washed with 0.1% Tx-100 three times, for 5 min each time, before mounting onto slides with Fluoromount-G (Southern Biothech, 0100-01). The primary antibody used for this paper were Tryptophan hydroxylase 2 (1:1,000; Millipore, ABN60), NeuN (1:1000; EnCor Biotechnology, RPCA-Fox3-AP) and Nissl (1:1000; NeuroTrace® 435/455 Blue Fluorescent Nissl Stain, Life technologies, N21479).

Fluorescent tracer injections in DRN

To examine the local connectivity and network structure in DRN, we performed visually guided injections of fluorescent tracer in DRN in acute slices (described fully in Trinh et al., 2015). Briefly, in these experiments, SERT-CRE-tdTomato animals (P30-P40) were sacrificed and acute slices prepared as described above (*slice preparation*). We utilized the following fluorescent dextrans interchangeably, observing similar expression and transport between them: Cascade Blue; Alexa Fluor 488; Alexa Fluor 647 (all suspended at 10 mg/mL in 0.1M phosphate-buffered saline (PBS) pH 7.2). At 5x-10x magnification with a Zeiss upright microscope with an epifluorescent lamp, fluorescent dextran (described above) was injected in either dorsal DRN or ventral DRN (all locations given in Fig. S7D, inset) by iontophoresis (five to ten 400 ms pulses of ~ –100 V via a 7 mOhm electrode). After a fluorescent sphere of ~100 μm diameter was observed, injection was stopped. Slices were incubated overnight in oxygenated ACSF to allow anterograde and retrograde transport, fixed in 4% paraformaldehyde and rendered transparent using the SeeDB procedure (see Trinh et al., 2015 for details). Slices were imaged with a Zeiss confocal microscope (LSM800 AxioObserver Z1 microscope; Zen software) with the appropriate filters for resolving the utilized fluorescent dextrans.

Data Analysis

All electrophysiological recordings were analyzed offline using Clampfit 10.4 (Molecular Devices), Origin 8.5 software (OriginLab) and the Numpy and Scipy technical computing libraries within Python. All data are presented as means +/- SEM. n refers to the number of cells recorded. Unless otherwise stated, statistical significances were derived using a standard Student's two-sided t-test ($p < 0.05$). P values of less than 0.05 were considered statistically significant and are indicated with an asterisk(*). Unless otherwise stated, all measurements were taken from distinct neurons.

All anatomy experiments (fluorescent tracer injections, described above) were analyzed using a combination of IMARIS and custom Python scripts. After acquiring images (described above in

Fluorescent tracer injections in DRN), we performed semi-automated cell identification using the IMARIS image analysis software package (Oxford Instruments), employing the spots function. This allowed us to identify neurons dual-labeled with tdTomato and the fluorescent dextran dye, and extract their coordinates in three dimensions (X,Y,Z) relative to the centre of the injection. Only neurons that were clearly outside of the extent of the dextran injection sphere were considered. We additionally extracted the locations of all neurons that were tdTomato positive (using Imaris) to act as a null distribution to compare with the co-expressing neurons. The validity of all identified neurons was manually checked against the raw fluorescence images. Together, this analysis yielded, for each individual dextran injection, a dataset of 3-dimensional coordinates for dual-labeled (tdTomato and fluorescence dextran) 5-HT neurons and single-labeled (tdTomato) 5-HT neurons.

Custom Python scripts employing the Numpy and Scipy technical computing libraries were utilized to perform analysis on this dataset. To compare the observed probability distribution over distance of dual-labeled neurons with the null model of unstructured connectivity, we first constructed a probability distribution of all single-labeled (tdTomato) 5-HT neurons over distance. We computed the difference, at each distance bin, between the probability distribution for dual-labeled neurons and the probability distribution for single-labeled neurons, yielding a delta probability distribution over distance. To construct a polar plot, we computed the angle of each cell relative to the injection centre and computed a probability distribution over these angles, calculating a delta probability distribution over angle similar to above.

Drugs and Chemicals
Picrotoxin, 2,3-Dioxo-6-nitro-1,2,3,4-tetrahydrobenzo[f]quinoxaline-7-sulfonamide di-sodium salt (NBQX), DL-2-Amino-5-phosphonopentanoic acid (D,L-APV), picrotoxin, CGP 55845 and WAY 100-635 were purchased from Abcam (Cambridge, MA). Tetrodotoxin (TTX) was purchased from Tocris Bioscience (Ellisville, MO). 4-Aminopuridine (4-AP) was purchased from Sigma Aldrich (St. Louis, MO).

Two-photon imaging of a serotonin biosensor
Stereotaxic Injections
Mice were injected with 0.05 mg/kg buprenorphine and anesthetized by inhalation of isoflurane. Injections were performed using a Nanoject III programmable injector with a pulled glass micropipette. 0.4µl was injected at a rate of 0.1µl per minute. Stereotaxic co-ordinates are as follows: DRN (from Lambda skull landmark, in mm, 0 degree needle angle: AP +0.4, ML +0, DV-3.2, -3, -2.8, sequential injections to tile DV extent of DRN with total volume split between all 3 coordinates.) 0.4 µl of AAV-EF1a-DIO-5-HT2M (Serotype:2/9; titre ~$4.3*10^{12}$ GC/mL) was injected and animals were left to recover for 3 to 4 weeks prior to 2P imaging experiments.

Two-photon imaging
Brain slices were prepared using the method described above in *Slice physiology and anatomy,* subsection *Slice preparation.* Slices were positioned in a recording chamber for imaging, and received constant bath application of Ringer solution containing (in mM): 119 NaCl, 2.5 KCl, 1.3 MgSO4, 2.0 CaCl2, 1.0 NaH2PO4, 11 glucose, 26.2 NaHCO3, and 0.1 L-Tryptophan saturated with 95% O2 and 5% CO2 (pH = 7.3; 295-310 mOsm/L). All imaging was conducted at a measured temperature of 30-32C through the use of an inline heater and stage heater. Two-photon imaging was performed using an ultrafast Ti:Sapphire pulsed laser

(InSightX3, Spectra Physics, Milpitas, California) tuned to 920nm to visualize both the GRAB5-HT2M (virally expressed) and tdTomato (5-HT neuron marker) signal. Laser intensity was controlled using an acousto-optic modulator coupled to a galvanometer scanning system (Zeiss 810; Germany). Images were acquired using a 40x/0.8NA objective.

Electrical stimulation of 5-HT afferents
Electrical stimuli were delivered through a patch pipette filled with Ringer solution (described above) located within the field of view. The pipette was filled with Alexa 594 for visualization during imaging experiments. Stimuli were 0.1 ms in duration, and consisted of frequencies provided in the text.

Drugs and chemicals
In some experiments, we bath applied either serotonin hydrochloride (Tocris Bioscience (Ellisville, MO; 50mM) or TTX (Tocris Bioscience (Ellisville, MO), 0.5uM).

Data analysis
ROIs were manually annotated and raw fluorescence was extracted using Suite2P (Pachitariu et al., 2016). All analysis was performed with custom Python scripts employing the Numpy and Scipy technical computing libraries. To compute $\Delta F/F_0$ traces, F_0 was calculated as the mean fluorescence from the last three pre-stimulus images. To compute $\int \Delta F/F_0$, in order to capture the full serotonin biosensor response, we integrated the trial-averaged $\Delta F/F_0$ response from stimulus onset to 5 seconds post-stimulus. To compute $max(\Delta F/F_0)$, we computed the value of the trial-averaged $\Delta F/F_0$ response for the same time period as described above.

For full-frame analysis of GRAB sensor responses, to decrease noise, we adopted a spatial pooling approach, first downsampling each 512x512 frame of GRAB sensor responses by a factor of 32 to yield a resolution of 16x16 pixels. Next, for each of these spatially pooled pixels, we computed a $\Delta F/F_0$ value at each time point as described above for ROI analysis. This yielded an image describing the spatial layout of stimulus responses for each stimulus type. To compute the pixel response index, for each frame we first considered the distribution of the $\int \Delta F/F_0$ responses across all pixels for the 5Hz electrical stimulation condition. From this 5Hz distribution we computed a standard deviation, σ_{5Hz}, and mean, μ_{5Hz}. These 5Hz parameters were used to calculate a normalized Z-score metric for the other conditions: $\boldsymbol{z} = (\boldsymbol{r} - \mu_{5Hz})/\sigma_{5Hz}$, where $\boldsymbol{r}$ is a vector across all pixels of $\int \Delta F/F_0$ values for a particular electrical stimulation frequency.

We computed the spatial autocorrelation of these images using the signal.correlate2d function from the Scipy technical computing library, yielding a 2-dimensional autocorrelation function of the same shape as the original stimulus response image, with maximal intensity value in the centre.

To compute spatial masks associated with the 2-dimensional autocorrelation function, we randomly shuffled the spatial location of each response pixel and computed a 2-dimensional autocorrelation function on this shuffled image. We repeated this shuffling 100 times for each image to compute a distribution of null autocorrelation responses across all pixels (excluding the central pixel corresponding to the two images fully aligned). Utilizing this null distribution of

autocorrelation values, we assigned a threshold of p=0.0001 for which experimental autocorrelation values would have to exceed to be considered significant (ie greater than 99.99% of null autocorrelation values). The number and spatial location of these statistically significant autocorrelation values were used to compute the spatial mask and area of autocorrelation.

The pixel correlation across trials was computed by calculating the Pearson correlation coefficient across all pixels for each unique pair of trials (typically 4-6 trials were presented), and then taking the mean of this Pearson r value. The pixel correlation (GRAB vs tdTomato) was computed similarly, but using the GRAB sensor response and the tdTomato signal as input.

Modeling of synaptic dynamics and network simulations

GIRK conductance
The GIRK conductance kinetics were modelled as a sum of exponentials:

$$g_{GIRK}(t|a, \tau_{decay}, \tau_{rise}) = a(e^{-t/\tau_{decay}} - e^{-t/\tau_{rise}}) \tag{1}$$

where $\tau_{decay} > \tau_{rise}$ and $t > 0$.

The parameters $a, \tau_{decay}, \tau_{rise}$ were fit to experimentally-determined 5-HT1AR-mediated GIRK conductances obtained following single-pulse electrical stimulation data (n=6 neurons) using a least-squares approach as follows: We first fit a monoexponential decay function $f(t|\tau_{decay}) = e^{-t/\tau_{decay}}$ to the normalized post-peak decay phase of $g_{GIRK}^{exp}(t)$. This yielded a constrained estimate of the GIRK decay, τ_{decay}. Next, fixing τ_{decay}, we found the values of a, τ_{rise} that minimized the mean-squared error over the whole observation period.

Each estimate was averaged across the population (n=6 neurons):

$$
\begin{aligned}
a &= 0.725 nS \\
\tau_{decay} &= 308.04 ms \\
\tau_{rise} &= 85.97 ms
\end{aligned}
$$

Quantifying 5-HT release dynamics through optogenetic stimulation
In some experiments, we optogenetically stimulated CHETA-expressing 5-HT neurons in slice, while recording 5-HT1AR outward currents from identified 5-HT neurons. By photostimulating at 20Hz and varying the optical pulse number, we could infer basic properties of release dynamics at these 5-HT releasing synapses.

We assume that the current generated by a single pulse is a random variable following a binomial distribution $B(n, p)$, which reflects the activity of n independent 5-HT releasing synapses with release probability p. For a train of successive pulses, we assume n is fixed,

and that each pulse is independent. Then $Q(n_{pulse} = a)$, the distribution of currents given by a pulses, is distributed as a sum of a binomial distributions:

$$Q(n_{pulse} = a) \quad \sim B(a \cdot n, p)$$

Accordingly, the variance of Q as the pulse number a varies is as follows (Sigworth, 1980):

$$var[Q(n_{pulse} = a)] = (p(1-p)n)a + noise \tag{2}$$

Since we observed that the variance of the current did not change as a function of pulse number (Fig. S6), we conclude that the release probability of 5-HT in these conditions can be approximated as being deterministic (i.e., close to $p = 1$). This conclusion nonetheless probably reflects the artefactual increase in release probability obtained when optogenetically activating synaptic terminals (Zhang & Oertner, 2007; see the section below).

Quantifying 5-HT release dynamics through electrical stimulation
We performed experiments where 5-HT afferents were electrically stimulated in acute slice preparations while recording from identified 5-HT neurons. Electrical stimulation intensities were low (in the perithreshold response range), and stimulation frequency was 1Hz. To quantify successes and failures, we computed the total event charge triggered by each stimuli (by integrating current over time), which formed an *event* distribution containing both putative successes and failures. We constructed a surrogate *noise* distribution by performing an identical analysis on a nonstimulated portion of the same recording for each neuron (*i.e.,* integrating equivalent durations of membrane noise where no stimuli were presented). Next, for each neuron we transformed the *event* and *noise* charge distributions into z-scored distributions (where the z-score was computed from that neuron's *noise* distribution). Finally, we pooled z-scored distributions across neurons. This allows us to directly compare electrically evoked responses; responses with a z-score > 3 likely represent successes, while responses with a z-score near 0 probably represent release failures. For electrical stimulation at higher frequencies (1-20Hz), we constructed surrogate noise distributions and event distributions in a manner identical to that described above.

For paired-pulse and pulse-chase stimuli, we quantified responses by first computing total charge elicited over the duration of each of the stimuli (integrating current over time). We next computed the total charge elicited by a single electrical stimulation delivered in isolation, and used this to calculate an expected value for purely linear summation of two pulses (*i.e.,* by multiplying the single stimulation-induced charge by 2). Finally, for each neuron, we normalized the stimulus-evoked charges in the paired-pulse or pulse-chase condition by the expected value for linear summation of two individual pulses.

Short-term plasticity
Short-term plasticity dynamics were modeled using a linear-nonlinear approach called the spike-response plasticity (SRP; Rossbroich et al., 2021). It utilizes a possibly complex efficacy kernel to capture time-varying, spike-triggered changes in release probability (i.e., short-term plasticity). Namely, each presynaptic spike is convolved with the following efficacy kernel:

$$\boldsymbol{k}_\mu(t) = a_s e^{-t/\tau_s} + a_l(e^{-t/\tau_l^+} - e^{-t/\tau_l^-}) \tag{3}$$

To capture the complexity of the dynamics, we chose to implement a kernel consisting of multiple timescales of responses. The short component is controlled by τ_s and a_s, and dictates the rapid short-term facilitation dynamics. The long component is controlled by τ_l^+ and a_l, and dictates the more prolonged short-term depression dynamics. τ_l^- is a shaping term that reduces the effect of short-term depression in the early post-spike phases where short-term facilitation dominates. Since the kernel consists of a sum of exponential functions, it can be conveniently written as a dynamical system. Let k reflects the current efficacy state of the synapse ($k = \boldsymbol{k}_\mu * S$) and is transformed through a nonlinear operation into a spike efficacy, μ. In turn, μ dictates the magnitude of the postsynaptic response given a presynaptic spike, and is implemented as a nonlinear readout of k:

$$
\begin{aligned}
f(x) &= \frac{1}{1+e^{-x}} \\
\mu &= \frac{1}{f(b)} f(k + b)
\end{aligned}
\tag{4}
$$

Here, $f(x)$ is the nonlinear function and b is a baseline term that controls whether the STP dynamics evolve sublinearly or supralinearly.

Fitting short-term plasticity dynamics to experimental data
We start with a set of paired-pulse electrical stimulation experiments where the time interval ranges from 50ms to 1sec. For our experimental dataset, we had computed, for each time interval t_i, a ratio of the total charge transfer for both pulses q_1 and q_2 to the charge transfer for a single pulse delivered separately:

$$R(t_i) = \frac{q_1+q_2}{q_1} = \frac{\mu_1+\mu_2(t_i)}{\mu_1} \tag{5}$$

Where μ_1 and μ_2 relative postsynaptic response amplitudes for the first and second pulse given by the SRP model. We wish to find optimal parameters for the efficacy kernel, $\boldsymbol{k}_\mu(t|a_s, a_l, \tau_s, \tau_l^+, \tau_l^-)$, such that we arrive at a local minimum in a cost function comparing the experimental and theoretical charges for various delays τ_i of the paired-pulse stimulation.

We start by deriving an analytical expression for (5) in terms of the underlying efficacy kernel.
        The first spike's efficacy is taken by evaluating the efficacy kernel at time 0:

$$\mu_1 = \frac{1}{f(b)} f(\boldsymbol{k}_\mu(t = 0) + b)$$

$$\mu_1 = \frac{1}{f(b)} f(a_s + b) \tag{6}$$

The second spike's efficacy is taken by evaluating the sum of an efficacy kernel at time t_i (efficacy from first spike at the time of spike two) and an efficacy kernel at time 0 (second spike's efficacy).

$$\mu_2 = \frac{1}{f(b)} f(\boldsymbol{k}_\mu(t=0) + \boldsymbol{k}_\mu(t=t_i) + b)$$

$$\mu_2 = \frac{1}{f(b)} f(a_s(1 + e^{-t_i/\tau_s}) + a_l(e^{-t_i/\tau_l^+} - e^{-t_i/\tau_l^-}) + b) \tag{7}$$

Substituting (6) and (7) into (5):

$$R(t_i) = \frac{\frac{1}{f(b)} f(a_s + b) + \frac{1}{f(b)} f(a_s(1 + e^{-t_i/\tau_s}) + a_l(e^{-t_i/\tau_l^+} - e^{-t_i/\tau_l^-}) + b)}{\frac{1}{f(b)} f(a_s + b)}$$

$$R(t_i) = \frac{f(a_s+b) + f(a_s(1 + e^{-t_i/\tau_s}) + a_l(e^{-t_i/\tau_l^+} - e^{-t_i/\tau_l^-}) + b)}{f(a_s+b)} \tag{8}$$

Equation (8) and the corresponding experimental results (5) are used to perform least-squares optimization on the parameters $a_s, a_l, \tau_s, \tau_l^+, \tau_l^-, b$. This yields the following parameter estimates:

$$
\begin{aligned}
a_s &= 1.11 \\
a_l &= -1.80 \\
\tau_s &= 0.128s \\
\tau_l^+ &= 1.27s \\
\tau_l^- &= 0.698s \\
b &= -6.22
\end{aligned}
$$

A general expression for spike efficacy given past spiking history
One can also write a solution to the generalized case of calculating the efficacy, μ_n, of a certain presynaptic spike n occurring at time t_n, given a past history of spikes with times $t_1, t_2, t_3, \ldots t_{n-1}$:

$$\mu_n = \frac{1}{f(b)} f\left(\sum_{s=1}^{n} \boldsymbol{k}_\mu(t=t_n - t_s)\right) \tag{9}$$

This equation is used to calculate the model's predictions for the matched pulse-chase experimental data.

Network model of recurrently connected 5-HT neurons receiving LHb input
　　　To construct a simplified model of the DRN network, we took a modular approach where each individual component was fitted to experimental data where possible. Our model consisted of 5000 conductance-based leaky-integrate-and-fire 5-HT neurons, building on recent simplified single-cell modeling of 5-HT neurons (Harkin et al., 2023). 25% of 5-HT

neurons received excitatory input from LHb ($\sum 7.5nS$), while all 5-HT neurons received background Poisson excitation as well as 5-HT1AR-mediated inhibition from other 5-HT neurons ($\sum 2.5nS$). The 5-HT synapse of recurrent inhibitory network were imparted with both the experimentally-fitted event kinetics and the short-term dynamics described above (i.e., from the SRP model).

Individual 5-HT neurons were fit with a simplified leaky-integrate-and-fire model. The fitting was performed experimentally according to previously described methods (Pozzorini et al., 2015). Briefly, identified serotonin neurons from SERT-CRE-tdTomato mice were recorded in whole-cell current-clamp configuration, and a noisy input stimulus was presented through multiple test and training repetitions. These repetitions were used to train a generalized integrate-and-fire model to each individual serotonin neuron by maximizing the likelihood of the model (for full details, see Harkin et al., 2023).

In this work we focused on basic extracted parameters from this model:$g_{leak}, E_{leak}, C_m, t_{refractory}, V_{thresh}$. While Harkin et al., (2023) utilized a current-based generalized-integrate-and-fire model to represent serotonin neuron firing, here we consider a simplified, conductance-based, leaky-integrate-and-fire model of individual serotonin neurons:

$$\frac{dV}{dt} = \frac{-g_{leak}(V-E_{leak})-g_{LHb}(V-E_{AMPA})-g_{pois}(V-E_{AMPA})-g_{GIRK}(V-E_K)-g_{AHP}^{tot}(v-E_K)-I_{const}}{C_m}$$

(10)

With the fitted parameters:

$$
\begin{aligned}
g_{leak} &= 1.4nS \\
C_m &= 0.0876nF \\
t_{refractory} &= 6.5ms \\
V_{thresh} &= -52.33mV
\end{aligned}
$$

And the chosen ionic reversal potentials, which, for the purposes of this LIF model, are considered as fixed parameters:

$$
\begin{aligned}
E_{leak} &= -68.07mV \\
E_K &= -90mV \\
E_{excit} &= 0mV
\end{aligned}
$$

Conductances for $g_{LHb}, g_{pois}, g_{GIRK}, g_{AHP}$ were considered with the following dynamics, where S is a synapse state variable which instantaneously increments by 1 whenever a presynaptic neuron fires, and exponentially decays to 0 with a characteristic time-constant:

$$
\begin{aligned}
g_x &= s_x \cdot g_x^{unitary} \\
\frac{ds}{dt} &= \frac{-s_x}{\tau_x}
\end{aligned}
$$

(11)

where x represents conductances of either *LHb, pois, GIRK* or *AHP*.

The AHP conductance is expressed as a sum of three unitary conductances with distinct timescales.

$$g_{AHP}^{tot} = s_{AHP}^{slow} g_{AHP}^{slow} + s_{AHP}^{med} g_{AHP}^{med} + s_{AHP}^{fast} g_{AHP}^{fast} \qquad (12).$$

The following values were chosen for individual parameters of the model. Note that excitatory decays were chosen to reflect a mixture of both AMPA and NMDA currents. Note that these reflect unitary conductance values, and neurons receive multiple LHb (g_{LHb}) and recurrent GIRK (g_{GIRK}) inputs, as well as a constant background rate of Poisson inputs (g_{pois}).

$$
\begin{aligned}
g_{LHb} &\sim N(\mu = 1.5, \sigma = 0.2)nS \\
g_{pois} &\sim N(\mu = 0.5, \sigma = 0.1)nS \\
g_{GIRK} &= 0.01nS \\
I_{const} &\sim N(\mu = 10, \sigma = 5)pA \\
\lambda_{pois} &= 25Hz \\
\tau_{A+N} &= 20ms \\
\tau_{GIRK} &= 50ms
\end{aligned}
$$

For the AHP dynamics, the following values were extracted from fits on identified 5-HT neurons using a previously described protocol (see Pozzorini et al., 2015; Harkin et al. 2023):

$$
\begin{aligned}
\tau_{AHP}^{fast} &= 3ms \\
\tau_{AHP}^{med} &= 30ms \\
\tau_{AHP}^{slow} &= 300ms \\
g_{AHP}^{fast} &= \frac{0.0281nA}{V_{thresh} - E_k} \\
g_{AHP}^{med} &= \frac{0.0333nA}{V_{thresh} - E_k} \\
g_{AHP}^{slow} &= \frac{0.0210nA}{V_{thresh} - E_k}
\end{aligned}
$$

Note that since the generalized integrate-and-fire model in Pozzorini et al. (2015) extracts a spike-triggered current for the AHP, we convert this to a conductance by dividing by the driving force at spike threshold.

The DRN network model consisted of 5000 5-HT neurons with the LIF dynamics described above. 5-HT neurons were recurrently connected in an unstructured manner with 5% random connection probability and 0.01 nS unitary GIRK synaptic conductance ($\sum 2.5nS$ net GIRK inhibition received by each neuron before facilitation/depression dynamics are considered). Short-term plasticity between 5-HT neuron connections was modelled using the linear-nonlinear description fit to data, described in the preceding section. 25% of DRN 5-HT neurons receive long-range input from a cluster of 500 LHb neurons emitting Poisson spike trains at rates specified in the text, where each LHb neuron and 5-HT neuron is randomly

connected with 1% probability (*i.e.,* each 5-HT neuron receives an average of 5 LHb synapses) with synapse strength specified above. This yields a mean LHb conductance of $\sum 7.5nS$ for each 5-HT neuron.

Network model of recurrently connected 5-HT neurons receiving LHb and A input (Winner-take-all computation)

In the simulations outlined in Fig. 7, we considered a DRN network where 25% of neurons received LHb input, and 25% of neurons received input from a generic area termed Area A. In these simulations, the basic neural architecture and dynamics were identical to that described above, except that 25% of 5-HT neurons now received input from Area A (non-overlapping with the 25% of neurons already receiving LHb input). Area A consisted of 500 neurons emitting Poisson spikes at rates specified in the text, where each A neuron and 5-HT neuron is randomly connected with 1% probability (*ie* each 5-HT neuron receives an average of 5 A synapses). This yields a mean A conductance of $\sum 7.5nS$ for each 5-HT neuron. Synapse strength is identical to that of LHb-5-HT synapses, as are decay kinetics:

$$g_X \quad \sim N(\mu = 1.5, \sigma = 0.2)nS$$
$$\tau_{A+N} \qquad = 20ms$$

Exploration of changes in network connectivity parameters

In some simulations, we varied the fraction of 5-HT neurons innervated by long-range LHb input, from 10-50% (Fig. S14). We sought to compute several network-wide metrics corresponding to recurrent drive, in order to assess whether these metrics were tracking input frequency, or alternately input fraction. The intuition behind these analyses is that the magnitude of the inhibitory synaptic conductance g_{GIRK}, associated with recurrent connections, provides a population-level metric of the strength of recurrence. First, for each simulated 5-HT neuron, we computed the maximum post-stimulus g_{GIRK} conductance, and we then constructed a cumulative distribution of this g_{GIRK} variable across the population (Fig. S14D, S14I). Second, we computed the population mean of g_{GIRK} for each combination of input fraction and input rate (Fig. S14E, S14J).

In other simulations, we varied the spatial structure of recurrent connectivity (local vs global recurrence) to assess its impact on winner-take-all computations (Fig. S16). The probability of recurrent connections was considered as a Gaussian distribution with a peak amplitude of the maximum connectivity (p=0.05), with standard deviations that varied according to the spatial structure of recurrent connectivity. We computed a simple metric relating the standard deviation of the connectivity (in units of neurons) to the total number of neurons of the network, $c = \sigma/n_{5-HT}$, which we report in the text. In these network simulations and simulations in Fig. S17, we sought to additionally compute a metric of the LHb-driven change in the response patterns of A-recipient 5-HT neurons. We compared the rates of these A-recipient neurons when receiving LHb input at a given rate, versus receiving no LHb input. This was computed as $\Delta 5\text{-}HT\ resp. = r_{5-HT}^{A-input}(A\text{-}on\ |\ LHb\text{-}on) - r_{5-HT}^{A-input}(A\text{-}on\ |\ LHb\text{-}off)$. We interpret this metric as quantifying the strength of the winner-take-all interaction and utilize it to explore the parameter space of combinations of rates of A and LHb.

We additionally varied the spatial overlap of inputs from LHb and A (Fig. S17). In these simulations, each input innervated a total of 25% of 5-HT neurons, but the fraction of neurons

receiving overlapping input from both LHb and A, O, is varied as a connectivity parameter across simulations. All other connectivity parameters are identical to in the main simulations. We computed the metric quantifying the strength of winner-take-all interactions as described above.

Finally, in many simulations we removed the short-term plasticity rules at recurrent connections. This was implemented by altering the plasticity kernel to be a flat line over time, such that any presynaptic input would produce a consistent postsynaptic response that was history-independent. All other parameters, including recurrent connectivity, were not altered.

Auditory classical conditioning task

Surgery

All experiments and procedures were performed in accordance with approved procedures and guidelines set forth by the University of Ottawa Animal Care and Veterinary Services. For stereotaxic injections, mice were injected with 0.05 mg/kg buprenorphine 1 hour prior to surgery and anesthetized by inhalation of isoflurane (4% for induction and 1.5-2% for maintenance). The head was shaved, ointment was applied to each eye, and an incision was made above the skull. A dental drill was used to perform skull craniotomies at locations above the coordinates for LHb and DRN described below.

For stereotaxic injections in LHb, we utilized the virus AAV9.CAG.hChR2(H134R)-mCherry.WPRE.SV40 at a concentration of ~10^{-12} GC/mL. Injections were performed using a 10 µl Hamilton Neuros syringe with point style 4 (12° beveled needle point). 0.4 - 0.5 µl was injected per coordinate at a rate of 0.1 – 0.2 µl per minute. Stereotaxic coordinates for injection are as follows for LHb, from Bregma skull landmark, in mm: AP +1.7, ML +/- 0.4, DV -3.1 to -3.3. Following stereotaxic injections, we affixed a custom stainless steel or 3D-printed Acrylonitrile butadiene styrene (ABS) headbar to the anterior surface of the skull with C&B Metabond Quick Adhesive Cement (Parkell).

For optic fiber implantation in DRN, each animal received a fiber (Doric Mono Fibre-Optic Cannula, 4mm length, part number: MFC_100/125-0.37_4mm_ZF1.25_C45) which was implanted at the following coordinates, from Bregma skull landmark in mm: AP-4.6, ML 0, DV -2.9. In some animals, we used an angled injection to avoid the superior saggital sinus (20° angle, from Bregma skull landmark in mm: AP-4.6, ML +1, DV -3.25). Optic fibers were slowly lowered to the desired dorsoventral location, and affixed with C&B Metabond Quick Adhesive Cement (Parkell) followed by Jet Denture repair cement (Lang) to provide structural stability to the implant.

After surgery, animals received topical bupivicaine (2%) and intraperitoneal (IP) carprofen (20 mg/kg). Post-operative care consisted of a further bupivicaine treatment at 4-6h post-surgery, and IP carprofen injection 4-6h post-surgery and for the next two days. Animals were allowed to recover from surgery for one week before water restriction was started. All animals were housed separately.

Behavior, water restriction and optogenetic stimulation

Animals were placed on water restriction for 7 days prior to the start of the task. During this time, they were habituated to the experimenters and the apparatus through handling and brief periods of head fixation while daily water (0.8 - 1 mL) was provided. After habituation, for 2-3 days each mouse was head-fixed for short bouts and 5-10 uL apple juice was delivered to a

lickport every 1 to 8 s without preceding auditory tones. Following this, mice underwent 1-2 weeks of training on the primary auditory classical conditioning task, described below.

The auditory classical conditioning task consisted of 120 randomly interleaved trials presented to each mouse per day, giving a total reward value of ~0.6 - 0.8 mL per mouse per day. Inter-trial intervals were drawn from an exponential distribution with $\lambda = 5s$, where $2s < ITI < 14s$ to prevent excessively long or short ITIs. Exponentially distributed ITIs give the convenient property that trials follow a Poisson distribution in time, and therefore are memoryless (the mice cannot predict trial onset times). To prevent the mice from licking continuously during the baseline periods, new trial starts were contingent on baseline mouse licking of <10% (fractional time). Three trial types were randomly interleaved: a *large-reward trial*, consisting of a 3 kHz tone and a 10uL apple juice reward, a *small-reward trial*, consisting of a 6 kHz tone and 5 uL apple juice reward, and a *no-reward trial*, consisting of a 12 kHz tone and 0 uL apple juice reward. For each trial, the tone lasted 1 s, the trace period between tone and reward delivery was drawn from a Gaussian distribution with $\mu = 2s, \sigma = 4s$ (values of $t_{trace} < 0.5s$ were prohibited), and the apple juice reward was concomitantly delivered with a 500 ms white noise tone to aid learning. Following reward delivery, licking was monitored for 4 s before the trial was terminated. This auditory classical conditioning task was presented daily to mice for 1-2 weeks until criterion performance had been achieved for >2 consecutive days. Criterion performance was defined as a mean lickrate of >0.5 Hz during the trace period of large-reward trials. While many mice displayed mean lickrates which were significantly higher than this (Fig. 8C), this was a convenient lower threshold for criterion performance.

After criterion performance was reached, mice experienced the auditory classical conditioning paradigm while we optogenetically activated the habenulo-raphe pathway in some trials. The trial structure consisted of 120 trials of interleaved large-reward, small-reward and no-reward trials as before. Each trial was composed of the same tone and reward associations and durations as described above, except the period between tone and reward was fixed at 2 s. For the first 40 trials, no optogenetic stimulation was delivered (*baseline block*). For the final 80 trials, optogenetic stimulation was delivered on 40 randomly chosen trials (*optogenetic stimulation block*). Optogenetic stimulation was composed of pulses of 10 ms width, at frequencies noted in the text (5Hz or 20Hz), delivered during the period between tone and reward. Optogenetic stimulation was delivered intracranially via an optic fibre implanted in DRN, to which a patch cable was connected before the session.

The behavioral apparatus consisted of a sound-insulated box containing a Raspberry Pi 3B, custom steel head fixation apparatus and equipment for delivering stimuli and acquiring signals. Lick signals were acquired via a custom 3D-printed combined infrared lickometer and stainless steel lickport. Lickport placement was adjusted daily for each mouse to ensure optimal performance and signal acquisition. A solenoid valve (NResearch 2-way NC isolation valve) gated apple juice delivery, and flow rate was calibrated weekly. Speakers connected to the Raspberry Pi (Logitech Z150) delivered auditory stimuli. We visualized animal behavior with an infrared video camera connected to the Raspberry Pi (Raspberry Pi Camera Module 2 NoIR). In vivo optogenetic stimulation was delivered using a PlexBright Compact LED module and controller (either blue light, 465; or yellow light, 590 nm; as noted in the manuscript; Plexon, Texas) and a PlexBright optical patch cable. A custom script (Python 3.8) on the Raspberry Pi registered lickometer data at 200 Hz while controlling trial structure, auditory stimulus delivery, reward delivery and optogenetic stimulation. Data was stored in hdf5 format.

After behavioral testing was completed for individual mice, they were perfused transcardially with 4% paraformaldehyde, and brains were cut in 50-100 um sections. Immunohistochemistry was performed as described above, and the injection sites and optic fiber tracts were verified.

Data analysis

Data was analyzed with custom Python 3.8 scripts, employing the numpy (van der Walt et al., 2011), scipy (Jones et al., 2001), h5py and matplotlib technical computing libraries. Briefly, raw licking data was binarized and converted into lick onset times. For visualizations (Fig. 8E-F), licking signals were calculated by convolving lick onset times with a gaussian kernel ($\sigma = 50ms$) and averaging across all trials of that type. Optogenetic analyses are described below.

Optogenetically evoked changes in anticipatory (trace) licking were calculated as follows: we considered trials of a given type during the optogenetic stimulation block (last 80 trials), and calculated the mean licking frequency in the period between the tone and reward (anticipatory licking) for each trial. To calculate an optogenetically evoked change in anticipatory licking, the mean anticipatory licking frequency for optogenetically stimulated trials was subtracted from the mean anticipatory licking frequency for non-stimulated interleaved trials during the optogenetic stimulation block. Optogenetically evoked changes in reward licking were calculated in a similar way, but the period considered was the post-reward epoch.

We additionally considered whether optogenetic stimulation modulated behavior beyond the stimulated period (Fig. 8I). For these analyses, we compared nonstimulated trials during the baseline block with nonstimulated interleaved trials during the optogenetic stimulation block. For each trial of a given type, we first computed mean anticipatory licking frequency as described above. Next, to quantify long-term optogenetically evoked changes in anticipatory licking, we subtracted mean anticipatory licking frequency for nonstimulated trials during the optogenetic stimulation block from mean anticipatory licking frequency for trials during the baseline block.

References

Sigworth, F.J. (1980). The variance of sodium current fluctuations at the node of Ranvier. J Physiol 307:97-129.

Pachitariu, M., Stringer, C., Schröder, S., Dipoppa, M., Rossi, L. F., Carandini, M., & Harris, K. D. (2016). Suite2p: beyond 10,000 neurons with standard two-photon microscopy. BioRxiv, 061507.

Figure S1: Validation of SERT-CRE-tdTomato mouse line and LHb injection site. Confocal image of SERTcre **(A)** and TPH2-expressing **(B)** 5-HT neurons in the DRN. **(C)** Merge confocal image of SERTcre and TPH2 channels. **(D)** Bar plot of overlap between SERT-cre and TPH2-expressing 5-HT neurons in the DRN (98.38 +/- 0.8; n = 3, mean +- SE). **(E)** Representative image of LHb injection site (AAV-ChR2-mCherry).

Figure S2: LHb sends axonal projections to the DRN. **(A)** Confocal image of GFP-expressing LHb axons in the DRN with 5-HT neurons identified with SERT-TdT. **(B)** ChR2-infected LHb neuron can be induced to spike with current injection. **(C)** LED photostimulation produces qualitatively similar spiking patterns to those induced by current injection. **(D)** Experimental schematic for long-range optogenetic interrogation. **(E)** Example serotonin neuron which does not receive direct LHb input, but displays a delayed heterosynaptic inhibitory current.

Figure S3: Anatomical location of 5-HT and GABA neurons recorded in the whole-cell configuration from the dorsal raphe nucleus. **(A)** Left, schematic of DRN and the location of cell types within DRN. Right, schematic of the targeted subregion of DRN for electrophysiological experiments. **(B)** Representative subsample of the anatomical locations of 5-HT and GABAergic neurons, projected onto the dorsoventral and mediolateral axes (n=48 neurons). Images were taken using a 5x objective after electrophysiological recording and were registered manually. Circles with shading denote a response as described in the legend, related to electrophysiology experiments described in Figures 1, 2 and 3, as well as Figure S4 and S10. GABA-in, LHb-in and 5-HT-in refer to the observed presence of GABAergic conductances, direct excitatory conductances from long-range LHb activation, and indirect serotonergic inhibitory conductances, respectively. LHb-opto stim and GABA-opto stim refer to optogenetic stimulation experiments described in Fig S10H-K, Fig. 1 and Fig. S4, while elec.-stim refers to electrical stimulation experiments described in Fig. 3 and Fig. S10A-G.

Figure S4: Lateral habenula input fails to recruit strong disynaptic inhibition onto 5-HT neurons located in the ventromedial portion of DRN. **(A)** Experimental setup for panels B-G. Schematic showing viral expression strategy (left), confocal images showing CHETA-YFP and tdTomato expression in DRN (middle), and schematic of whole-cell patch clamp recording setup from SOM neurons, related to panels B and C (right). **(B)** Example trace from CHETA-YFP-expressing 5-HT neuron recorded in voltage clamp, demonstrating photostimulation-evoked inward current. **(C)** Example trace from CHETA-YFP-expressing 5-HT neuron recorded in current clamp, demonstrating photostimulation-evoked spiketrains at 20Hz. **(D)** Schematic of whole-cell patch-clamp recording setup from 5-HT neurons, related to panels E-G. **(E)** Trial-averaged traces from an example 5-HT neuron, recorded in voltage clamp, illustrating photostimulation-driven currents at two holding potentials. Inset depicts zoomed region. **(F)** Latency to peak outward current response at Vm=-40mV across all photostimulation events (n=120 events) for one example neuron. **(G)** I-V curve for photostimulation-driven currents from one example neuron. **(H)** Experimental setup for panels I-O. **(I)** Trial-averaged trace from an example SOM neuron, recorded in voltage clamp, illustrating photostimulation-driven inward currents. Inset depicts zoomed region. **(J)** Latency to peak inward current response across all photostimulation events (n=615 events, 8 neurons). **(K)** Amplitude of peak current response across all photostimulation events (n=615 events, 8 neurons). **(L-M)** Current-clamp traces from two example SOM neurons showing subthreshold responses to photostimulation **(L)** and suprathreshold responses to photostimulation **(M)**. **(N)** Classification of SOM responses to photostimulation across the population of recorded neurons. **(O)** Peak depolarization from baseline for SOM neurons exhibiting subthreshold responses (n=7). Line and shaded area express mean +- standard deviation.

Figure S5: Voltage-clamp experiments provide evidence that the afterhyperpolarization of serotonin neurons is mediated by SK channels, not 5-HT1ARs. **(A)** Representative traces of AHP current (IAHP) driven by a 500 ms depolarization step to 20 mV in voltage clamp. **(B)** Representative scatter plot showing no differences in IAHP amplitude following WAY 100-635 application. **(C)** Representative IAHP traces before and after APAMIN application (300 nM). **(D)** Representative scatter plot showing the IAHP +500 ms amplitudes in response to APAMIN. **(E)** +500 ms IAHP amplitudes before and after APAMIN application (CTL, 91.1 +/- 15.3; APM, 19.6 +/- 5.0; n = 3; p = 0.02, paired two-sided t-test).

Figure S6: Validation of local CHETA optogenetic strategy in DRN. **(A)** Schematic depicting AAV-DIO-ChETA-YFP injection into the DRN of SERTcre-TdT mice. **(B)** Confocal image of ChETA-YFP expression in tdTomato-positive neurons in the DRN. Above, low magnification (scale, 300 μm) ; below, high magnification (scale, 50 μm). **(C)** Schematic and EPI-flourescent image of whole-cell recordings from ChETA-YFP-expressing SERT-TdT neurons. **(D)** Left, LED stimulation of 5-HT neuron spiking with different pulse durations at frequency of 20 Hz. Right, quantification of the probability of photostimulation eliciting a spike as a function of pulse width (n=8 pulses). **(E)** LED stimulation of 5-HT neuron spiking in response to varying number of 10 ms pulses at 20 Hz. **(F)** Representative IV relationship traces and IV relationship (n = 5, mean +- SE). **(G)** Representative traces and scatter plot showing complete blockade of outward current by WAY 100-635. Right, population data is displayed (Pre-WAY, 112.9 +/- 22.9; Post-WAY, 11.5 +/- 1.1; n = 3; p = 0.02, paired two-sided t-test).

A dorsal injection — fluorescent dextran or ventral injection — fluorescent dextran — 18h incubation → **SeeDB clearing + microscopy** → **Cell registration and image processing**

Aq / DRN

B SERT-Cre:tdTom — Dextran (100um) / tdTom (100um) / Overlaid (100um)

C dorsal injection (ex.) — tdTom / Dextran (100um) — dextran-labeled 5-HT — all 5-HT

injection center

ap (um) / ml (um) / dv (um)

D dorsal injection (pop.) — injection center — 200um — inj. sites (~400um, d, v) — ventral injection (pop.) — 200um — dorsal / lateral / lateral / ventral

E pdf — 0.00–0.25 — distance (um) — * p=1.29x10⁻³

F Δpdf from null 5-HT dist. — distance (um) — dextran vs null 5-HT — unstructured — over / under

G 0° / 45° / 90° / 135° / 180° / 225° / 270° / 315° — pdf — dorsal / lateral / lateral / ventral — θ

H 0° / 45° / 90° / 135° / 180° / 225° / 270° / 315° — dextran vs null 5-HT — unstructured

Figure S7: Spatial scale and anatomical features of recurrent connectivity. **(A)** Experimental schematic demonstrating fluorescent dextran labeling approach. **(B)** Representative confocal images of DRN showing dextran injection site and labeled neurons (left), tdTomato neurons (middle) and overlaid image showing colocalization (right, arrows). **(C)** Representative confocal image (left) depicting dextran injection in dorsal DRN (cyan), tdTomato-positive 5-HT neurons (red), and neurons exhibiting colocalization (arrows). Extracted coordinates are shown for colabeled 5-HT neurons (middle) and all 5-HT neurons (right). **(D)** Locations of all colabeled 5-HT neurons from injection center for dorsal injection (left; n=17 neurons, 3 injections) and ventral injection (right; n=21 neurons, 3 injections). Inset depicts anatomical locations of each injection site within DRN. **(E)** Probability density of the distance from colabeled 5-HT neurons to injection center, for dorsal (purple) and ventral (orange) injections (K-S test for statistical comparison). **(F)** Probability density of colabeled neuron distance, expressed as a difference in probability from the distribution of all 5-HT neurons (unstructured, dotted line). **(G)** Polar plot depicting anatomical location of colabeled 5-HT neurons from injection site, for dorsal (purple) and ventral (orange) injections. **(H)** Polar plot of colabeled neuron location as in G, but expressed as a difference in probability from the distribution of all 5-HT neurons (unstructured, dotted line).

Figure S8: Postsynaptic dynamics of recurrent 5-HT1A connections indicate non-desensitizing receptors. **(A)** Representative averaged traces from a single non-CHETA-expressing 5-HT neuron depicting responses to 20Hz optogenetic stimulation of DRN for various pulse numbers. **(B)** Event charge for each neuron as a function of pulse number at 20Hz (n=6). **(C)** Charge mean as a function of pulse number, normalized to the mean at two pulses (n=6, mean +-SD). Dotted red line represents a linear relationship between pulse number and evoked charge. **(D)** Charge variance as a function of pulse number, normalized to the variance at two pulses (n=3, mean +-SD). Dotted red line represents predicted variance for release probability p=1.

Figure S9. Frequency dependence of 5-HT1A connections at suprathreshold stimulation frequencies. **(A)** Experimental setup depicting electrical stimulation of 5-HT1A synapses while recording from an identified 5-HT neuron in the whole-cell configuration. **(B)** Representative averaged traces from a single neuron depicting responses to electrical stimulation at various frequencies, with pulse number held constant. **(C)** IPSC amplitude as a function of frequency, for individual neurons (grey) and the mean of all neurons (blue, n=3). **(D)** Event charge, expressed as a z-score relative to a noise distribution, as a function of frequency. Depicted are individual neurons (grey) and the mean of all neurons (blue, n=3). Green shaded area represents 95% confidence interval for noise distribution.

Figure S10. Frequency-dependent facilitation of serotonin release under low extracellular calcium conditions. Related to Figure 3. **(A)**: Experimental setup for panels B-C depicting whole-cell patch clamp recording from genetically identified 5-HT neuron with electrically evoked 5-HT release, under physiological recording conditions (extracellular calcium and temperature). Inset depicts electrical stimulation parameters for frequency-varying experiments. **(B)** Representative averaged traces from a single neuron depicting responses to electrical stimulation at various frequencies. Red dotted lines denote electrical stimulation, and the shaded area denotes standard deviation across trials. **(C)** Charge z-score as a function of electrical stimulation frequency, with 95% noise C.I shown in green (n=4 neurons). **(D)** Schematic showing electrical stimulation parameters for paired-pulse experiments (E-G). **(E)** Representative averaged traces from a single neuron depicting responses to the paired-pulse electrical stimulation protocol. **(F)** Charge histograms for single-pulse responses and paired-pulse responses at 50ms for one example neuron. **(G)** Post-stimulus charge as a function of paired-pulse time interval (n=4 neurons). Green line depicts expected charge from linear summation. Blue line depicts population average and grey lines are responses of individual neurons. **(H)** Experimental setup for panels I-K depicting channelrhodopsin injection in LHb (left), and whole-cell patch clamp recording from genetically identified 5-HT neuron with electrically evoked 5-HT release, under physiological recording conditions (right). **(I)** Representative averaged traces from one example neuron depicting current responses to optogenetic stimulation of LHb afferents at two frequencies. Note that EPSCs are truncated for clarity to focus on direct comparison of the outward GIRK conductance. **(J)** Representative averaged traces from another example neuron depicting current responses to optogenetic stimulation of LHb afferents at four frequencies. **(K)** Total charge evoked by LHb stimulation as a functino of optogenetic stimulation frequency. Each line depicts one neuron.

Figure S11: GRAB serotonin biosensor imaging reveals that 5-HT release is action potential-dependent. Related to Figure 5. **(A)** Left: Max-projection images of whole-frame tdTomato fluorescence and GRAB fluorescence during 20Hz electrical stimulation (8 pulses). Right: GRAB sensor responses to electrical stimulation at 20Hz before and after TTX washon for the ROIs depicted on the left. Thin traces represent individual trials and thick traces represent means across all trials. **(B)** Example whole-frame tdTomato fluorescence and binned GRAB responses to 20Hz electrical stimulation before and after TTX washon (left), and cumulative distribution of GRAB responses across pixels during 20Hz electrical stimulation before and after TTX washon (right; Kolmogorov-Smirnov test displayed). **(C)** Spatial autocorrelation plots for the GRAB fluorescence frames shown in B. Autocorrelation values are normalized from 0 to 1 for presentation.

Figure S12. Generating a mask for statistically significant autocorrelation regions with a pixel shuffling approach. Related to Figure 5. **(A)** Example region showing whole-frame tdTomato fluorescence (left) and binned GRAB responses to 5, 10 and 20Hz electrical stimulation (8 pulses; right). **(B)** Spatial autocorrelation plots for the frames shown in A. Autocorrelation values are normalized from 0 to 1 for presentation. **(C)** Same as A, but the spatial location of pixels are shuffled. **(D)** Spatial autocorrelation plots for the pixel-shuffled frames shown in C. Autocorrelation values are normalized from 0 to 1 for presentation. **(E)** Probability distribution function of raw autocorrelation values for all pixels, for the spatial autocorrelations performed on pixel-shuffled GRAB responses (grey; computed from D) and for the spatial autocorrelations performed on GRAB responses (colors correspond to labeled electrical stimulation frequencies; computed from B). Pixels to the right of the green threshold have autocorrelation values that statistically exceed those of the shuffled distribution (p=0.0001). **(F)** Spatial mask for statistically significant autocorrelation values in B, where only pixels that exceed the significance threshold compared to the pixel-shuffled noise distribution (p=0.0001, as computed in E) are shown.

Figure S13: Nonlinear transformation of habenulo-raphe afferents generated by recurrent plasticity rules, with tenfold decrease in recurrent inhibitory connection probability (sum 0.25nS conductance received by each serotonin neuron). Related to Figure 6. **(A)** Schematic of the LIF network model. **(B)** Results of a simulation where a 5Hz step input from LHb was delivered to a subset of serotonin neurons. **(C)** Results of a simulation where a 20Hz step input from LHb was delivered to a subset of serotonin neurons. **(D)** Input-output transformation for the transient (0-200ms) response phase, with (black) and without (grey) STP rules at recurrent serotonergic connections. **(E)** Input-output transformation for the stationary (200-400ms) response phase, with (black) and without (grey) STP rules at recurrent serotonergic connections. **(F)** Input-output transformation for the post-response (0-1s after stimulus) phase, with (black) and without (grey) STP rules at recurrent serotonergic connections.

Figure S14. The input-output function of the habenulo-raphe pathway is largely invariant to the fraction of 5-HT neurons innervated by LHb, and this invariance depends on recurrent short-term plasticity. Related to network simulations depicted in Figure 6. Panels A-E depict results of simulations where short-term plasticity at 5-HT-5-HT connections is intact. **(A)** Input-output relationship of 5-HT neurons during the transient (0-200ms) phase of LHb input. Colors denote different fractions of 5-HT neurons receiving LHb input (shown in inset; 10-50%). **(B)** Same as A, but during stationary phase of LHb input (200-400ms). **(C)** Same as A, but during the post-response period of LHb input (0-1s after). **(D)** Cumulative distribution function of GIRK conductance across simulated 5-HT neurons, for 5Hz LHb input (red) and 20Hz LHb input (blue), and for fractions of 5-HT neurons innervated by LHb at 10% (lighter shade) and 50% (darker shade) (all combinations depicted in inset). Note how GIRK conductance depends primarily on LHb input frequency, instead of on LHb input fraction. **(E)** Relationship between LHb input frequency and GIRK conductance (population mean) as LHb input fraction is varied (colors denote distinct input fractions, shown in inset), demonstrating that population GIRK conductance is largely invariant to LHb input fraction. Panels D-H depict results of simulations where short-term plasticity at 5-HT-5-HT connections has been removed (i.e. linearized responses). **(F)** Input-output relationship of 5-HT neurons during the transient (0-200ms) phase of LHb input. Colors denote different fractions of 5-HT neurons receiving LHb input (shown in inset; 10-50%). **(G)** Same as A, but during stationary phase of LHb input (200-400ms). **(H)** Same as A, but during the post-response period of LHb input (0-1s after). **(I)** Cumulative distribution function of GIRK conductance across simulated 5-HT neurons, for 5Hz LHb input (red) and 20Hz LHb input (blue), and for fractions of 5-HT neurons innervated by LHb at 10% (lighter shade) and 50% (darker shade) (all combinations depicted in inset). Note how without short-term plasticity, GIRK conductance depends on both LHb input frequency and LHb input fraction. **(J)** Relationship between LHb input frequency and GIRK conductance (population mean) as LHb input fraction is varied (colors denote distinct input fractions, shown in inset), demonstrating that without short-term plasticity dynamics, the population GIRK conductance is not invariant to LHb input fraction.

Figure S15: Plasticity-dependent winner-take-all dynamics in raphe, with tenfold decrease in recurrent inhibitory connection probability (sum 0.25nS conductance received by each serotonin neuron). **(A)** Simulations with LHb inputs of either 5Hz (left) and 20Hz (middle) were delivered to serotonin neurons during ramp input from area A. Schematic of LIF network model with two inputs (right). Related to Figure 7. **(B)** Mean responses of serotonin neurons to 5Hz (left) and 20Hz (right) input with STP rules intact. **(C)** Raster plot showing serotonin neuron responses to 5Hz (left) and 20Hz (right) input with STP rules intact. **(D)** Input-output function, for input A, with STP rules intact and coincident LHb input at 0, 5 or 20Hz. **(E)** Mean responses of serotonin neurons to 5Hz (left) and 20Hz (right) input with STP rules removed. **(F)** Raster plot showing serotonin neuron responses to 5Hz (left) and 20Hz (right) input with STP rules removed. **(G)** Input-output function, for input A, with STP rules removed and coincident LHb input at 0, 5 or 20Hz.

Figure S16: Winner-take-all dynamics in the DRN persist over a wide range of spatial connectivity profiles for recurrent inhibition. Related to network simulations shown in Figure 7. **(A)** Schematic of network simulation, with LHb and Area A providing input to 5-HT neurons (top). Plot demonstrating how 5-HT-5-HT recurrent connectivity probability varies with distance, as spatial connectivity profiles are parameterized (bottom). **(B)** Simulation with more global recurrent spatial connectivity of c=1/2. Shown are mean firing rates of A and LHb inputs (top); mean firing rates of A-recipient 5-HT neurons (middle); and raster plot depicting firing rates of 2500 5-HT neurons receiving either LHb input (purple) or A input (orange) (depicted is the 50% of 5-HT neurons receiving direct input from either LHb or PFC). **(C)** Simulation with more local recurrent spatial connectivity of c=1/4. Layout of panels is identical to in B. D-E quantify the strength of winner-take-all dynamics as a function of recurrent spatial connectivity profiles, while short-term plasticity at recurrent connections is intact. **(D)** LHb-evoked change in A-recipient 5-HT neuron responses, as a function of the rate of A input, for 5Hz (left) and 20Hz (right) LHb input. Spatial connectivity profile is varied (shown in legend), demonstrating that winner-take-all dynamics are stable across a wide range of spatial recurrent connectivity profiles. **(E)** Strength of winner-take-all effect (quantified as LHb-evoked change in A-recipient 5-HT neuron responses) as A input rate and LHb input rate are varied. Shown are recurrent spatial connectivity values of c=1 (left) and c=1/4 (right). F-G quantify the strength of winner-take-all dynamics as a function of recurrent spatial connectivity profiles, while short-term plasticity at recurrent connections has been removed (i.e. linearized responses). **(F)**: LHb-evoked change in A-recipient 5-HT neuron responses, as a function of the rate of A input, for 5Hz (left) and 20Hz (right) LHb input. Spatial connectivity profile is varied as depicted in the

legend, demonstrating that with short-term plasticity removed, the winner-take-all dynamics vary over the tested range of spatial recurrent connectivity profiles. **(G)** Strength of winner-take-all effect (quantified as LHb-evoked change in A-recipient 5-HT neuron responses) as A input rate and LHb input rate are varied. Shown are recurrent spatial connectivity values of $c=1$ (left) and $c=1/4$ (right). Note that the winner-take-all effect is weaker than with short-term plasticity intact (E).

Figure S17: Winner-take-all dynamics in the DRN persist over a wide range of overlap fractions between competing inputs, a phenomenon which depends on recurrent short-term plasticity. Related to network simulations shown in Figure 7. **(A)** Schematic of network simulation, with LHb and Area A providing input to 5-HT neurons (top). Bottom contains schematic demonstrating how the overlap between LHb and A inputs onto 5-HT neurons is varied from 0-60% . **(B)** Simulation with small input overlap of *o*=0.2. Shown are mean firing rates of A and LHb inputs (top); mean firing rates of A-recipient 5-HT neurons (middle); and raster plot depicting firing rates of 2500 5-HT neurons receiving either LHb input (purple) or A input (orange) (depicted is the 50% of 5-HT neurons receiving direct input from either LHb or PFC). **(C)** Simulation with larger input overlap of *o*=0.6. Layout of panels is identical to in B. D-F quantify the strength of winner-take-all dynamics as a function of input overlap fraction, while short-term plasticity at recurrent connections is intact. **(D)** LHb-evoked change in A-recipient 5-HT neuron responses during the transient phase (0-200ms), as a function of the rate of A input, for 5Hz (left) and 20Hz (right) LHb input. Input overlap is varied (shown in legend). **(E)** LHb-evoked change in A-recipient 5-HT neuron responses during the stationary phase (200-500ms), as a function of the rate of A input, for 5Hz (left) and 20Hz (right) LHb input. Input overlap is varied (shown in legend). **(F)** Strength of winner-take-all effect (quantified as LHb-evoked change in A-recipient 5-HT neuron responses) as A input rate and LHb input rate are varied. Shown are input overlap values of *o*=0 (left) and *o*=0.4 (right), illustrating winner-take-all dynamics (ie negative 5-HT modulation) for these input fractions. G-I quantify the strength of winner-take-all dynamics as a function of input overlap fraction, while short-term plasticity at

recurrent connections has been removed (i.e. linearized responses). **(G)** LHb-evoked change in A-recipient 5-HT neuron responses during the transient phase (0-200ms), as a function of the rate of A input, for 5Hz (left) and 20Hz (right) LHb input. Input overlap is varied (shown in legend). **(H)** LHb-evoked change in A-recipient 5-HT neuron responses during the stationary phase (200-500ms), as a function of the rate of A input, for 5Hz (left) and 20Hz (right) LHb input. Input overlap is varied (shown in legend). **(I)** Strength of winner-take-all effect (quantified as LHb-evoked change in A-recipient 5-HT neuron responses) as A input rate and LHb input rate are varied. Shown are input overlap values of $o=0$ (left) and $o=0.4$ (right), illustrating that winner-take-all dynamics are no longer present with moderate input overlap of $o=0.4$ when short-term plasticity is removed (compare with F).

Figure S18: Recurrent inhibition gates spiking activity of 5-HT neurons. **(A)** Schematic of experimental design. Left, AAV-DIO-ChETA-YFP is injected into the DRN at a 2:1 ratio with Mannitol. Right, whole-cell recordings from ChETA-negative 5-HT neurons. **(B)** Recurrent 5HT1AR-mediated current in ChETA-negative 5-HT neuron. **(C)** Spiking of a 5-HT neuron in response to current injection. Below, raster plot showing the same neuron's activity across trials. **(D)** Spiking of a 5-HT neuron in response to current injection and 20Hz optical activation of CHETA-positive 5-HT neurons. Below, raster plot showing the same neuron's activity across trials.**(E)** Spiking of a 5-HT neuron in response to current injection in the presence of WAY100-635. Below, raster plot showing the same neuron's activity across trials. **(F)** Spiking of a 5-HT neuron in response to current injection and 20Hz optical activation of CHETA-positive 5-HT neurons, in the presence of WAY100-635. Below, raster plot showing the same neuron's activity across trials. **(G)** Smoothed firing rate (normalized) over time for pre- and post-WAY recordings with LED stimulation (mean +- SE, n=3 neurons). **(H)** Cumulative spike fraction in pre- and post-WAY recordings with LED stimulation (n=3 neurons, 2-sample Kolmogorov-Smirnov test: p = 7.4e-15). **(I)** Smoothed firing rate (normalized) over time for pre- and post-WAY recordings without LED stimulation (mean +- SE, n=3 neurons). **(J)** Cumulative spike fraction in pre- and post-WAY recordings with LED stimulation (3 neurons, 2-sample Kolmogorov-Smirnov test: p = 0.85).

Figure S19: LHb-driven, 5HT1AR mediated, recurrent feedforward modulation of 5-HT firing activity in the DRN. **(A)** Schematic of experimental design. Left, AAV-ChR2(H134R)-mCherry is injected into the LHb. Right, whole-cell recordings from TdT-positive 5-HT neurons. **(B)** LHb-driven heterosynaptic 5HT1AR-mediated potential in SERT-TdT 5-HT neuron. **(C)** Spiking of a 5-HT neuron in response to a 2.5 s current step (scale, 20 mV, 500ms). Below, raster plot of the same 5-HT neuron spiking. **(D)** Spiking of a 5-HT neuron in response to a 2.5 s current step with LED activation of ChR2-expressing LHb terminals in the slice for 1 sec in the middle of the current step. Below, raster plot of the same 5-HT neuron spiking. **(E)** Spiking of a 5-HT neuron in response to a 2.5 s current step in the presence of WAY 100-635. Below, raster plot of the same 5-HT neuron spiking. **(F)** In the presence of WAY 100-635, spiking of a 5-HT neuron in response to a 2.5 s current step with LED activation of ChR2-expressing LHb terminals in the slice for 1 sec in the middle of the current step. Below, raster plot of the same 5-HT neuron spiking. **(G)** Instantaneous frequency line (top) and spike count histogram plots (below) from recording in **C**. **(H)** Instantaneous frequency line (top) and spike count histogram plots (below) from recording in **D**. **(I)** Instantaneous frequency line (top) and spike count histogram plots

(below) from recording in **E**. **(J)** Instantaneous frequency line (top) and spike count histogram plots (below) from recording in **F**. **(K)** Instantaneous frequency population plot in pre- and post-WAY recordings with LED stimulation (n=3 neurons, mean +- SE). **(L)** Cumulative frequency population plot in pre- and post-WAY recordings with LED stimulation (3 cells, p = 0.0017, 2-sample Kolmogorov-Smirnov test). **(M)** Instantaneous frequency population plot in pre- and post-WAY recordings in the absence of LED stimulation (n=3 neurons, mean +- SE). **(N)** Cumulative frequency population plot in pre- and post-WAY recordings in the absence of LED stimulation (3 cells, p = 0.07, 2-sample Kolmogorov-Smirnov test).

A

LHb

Small reward trials

B
Lick rate (Hz)
20Hz
S. rew
opto
no-opto
Time from tone (s)

C
Lick rate (Hz)
5Hz
S. rew
opto
no-opto
Time from tone (s)

No reward trials

D
Lick rate (Hz)
20Hz
No rew.
opto
no-opto
Time from tone (s)

E
Lick rate (Hz)
5Hz
No. rew
opto
no-opto
Time from tone (s)

Post-reward stimulation

F
Lick rate (Hz)
L. rew
20Hz
opto
opto
no-opto
Time from tone (s)

G
Δ reward licking (Hz)
N.S.

Response magnitude during large-reward trace optogenetic stimulation

H
opto-evoked Δ trace licking (Hz)
y = 0.14x + -0.13
r = 0.19
y = -0.27x + -0.53
r = -0.54
trace licking (Hz)

I
5Hz opto Δ trace licking (Hz)
y = 0.06x + 0.20
r = 0.05
20Hz opto Δ trace licking (Hz)

Figure S20: The habenulo-raphe pathway disrupts reward-conditioned responses in a value-dependent manner and does not affect unconditioned response. **(A)** Representative example of LHb injection site. **(B)** Mean licking across all small reward trials for a single mouse depicting control trials (black) or 20Hz optogenetic stimulation trials (blue). **(C)** Mean licking across all small reward trials for a single mouse depicting control trials (black) or 5Hz optogenetic stimulation trials (blue). **(D)** Mean licking across all no reward trials for a single mouse depicting control trials (black) or 20Hz optogenetic stimulation trials (blue). **(E)** Mean licking across all no reward trials for a single mouse depicting control trials (black) or 5Hz optogenetic stimulation trials (blue). **(F-G)** depict post-reward optogenetic stimulation experiments. **(F)** Mean licking across all large reward trials for a single mouse depicting control trials (black) or trials where 20Hz optogenetic stimulation was delivered during the post-reward period (blue). **(G)** Change in post-reward licking during optogenetic stimulation trials, compared to pre-stimulated baseline period (n=6 mice, paired two-sided t-test, unstimulated vs stimulated reward licking). **(H)** Correlation between mean trace licking (large reward trials) and optogenetic effect size for 20Hz stimulation (dark blue) and 5Hz stimulation (light blue) for trace stimulation during large reward trials. **(I)** Correlation between effect size for 20Hz stimulation and 5Hz stimulation for trace stimulation during large reward trials.