gUMI-BEAR, a modular, unsupervised population barcoding method to track variants and evolution at high resolution

Short Title: A modular barcoding method to track variants and evolution

Shahar Rezenman[1*], Maor Knafo[1*], Ivgeni Tsigalnitski[1], Shiri Barad[1], Ghil Jona[2], Dikla Levi[2], Orly Dym[3], Ziv Reich[1§] and Ruti Kapon[1§]

[1]Dept. of Biomolecular Sciences, [2]Life Sciences Core Facilities, [3]The Dana and Yossie Hollander Center for Structural Proteomics

Weizmann Institute of Science Rehovot, Israel 7610001

*These authors contributed equally to this work

[§]Corresponding Author

ruti.kapon@weizmann.ac.il

ziv.reich@weizmann.ac.il

Abstract

Cellular lineage tracking provides a means to observe population makeup at the clonal level, allowing exploration of heterogeneity, evolutionary and developmental processes and individual clones' relative fitness. It has thus contributed significantly to understanding microbial evolution, organ differentiation and cancer heterogeneity, among others. Its use, however, is limited because existing methods are highly specific, expensive, labour-intensive, and, critically, do not allow the repetition of experiments. To address these issues, we developed gUMI-BEAR (genomic Unique Molecular Identifier Barcoded Enriched Associated Regions), a modular, cost-effective method for tracking populations at high resolution. We first demonstrate the system's application and resolution by applying it to track tens of thousands of *Saccharomyces cerevisiae* lineages growing together under varying environmental conditions applied across multiple generations, revealing fitness differences and lineage-specific adaptations. Then, we demonstrate how gUMI-BEAR can be used to perform parallel screening of a huge number of randomly generated variants of the *Hsp82* gene. We further show how our method allows isolation of variants, even if their frequency in the population is low, thus enabling unsupervised identification of modifications that lead to a behaviour of interest.

Introduction

With the advent of single-cell techniques, it became clear that significant heterogeneity exists within populations of seemingly identical cells(1–6). However, how individual adaptations translate into success or failure, or how population dynamics are affected by individual variations and their degree, is not yet clear. A number of studies are beginning to shed light on these questions by employing procedures to quantify a population's clonal makeup(6–13). Using these methods, multiple lineages within a population can be distinguished by the introduction of DNA barcodes, short segments of DNA inserted into cells' genomes and passed on to their progenies. By sequencing the barcodes, one can capture a snapshot of the clonal composition of a growing population at any given moment. Successive records can be used to trace the prevalence of individual lineages through time(14), offering insight into adaptation dynamics and fitness changes as they occur *in the context of evolving populations* rather than in each strain on its own.

The study of heterogeneity need not be limited to individuals within a population but could also be considered in the context of a particular gene or pathway. Quantifying the effect of gene-variants allows mapping the fitness landscape of a gene towards the understanding of the biophysical and biochemical requirements of particular regions. Likewise, it can aid in designing disease treatments. Here, the conjugation of DNA barcodes to specific gene variants creates a one-to-one association between barcode and variant and allows to use barcode incidence as a proxy for the abundance of each variant in a population. Thus, the fitness of many variants can be tested together, in one experiment, in the evolutionary setting in which they compete with each other.

Cellular barcoding has thus become invaluable for understanding core evolutionary

processes such as clonal interference(15), differentiation of organs(16) and

heterogeneity development in cancer(1). For variant testing, it has allowed the

assessment of unprecedented numbers of variants(17,18). Its utilisation is, however,

limited because it is generally expensive, labour-intensive, highly specific to each

biological system, and current modes of application do not lend themselves to repetition

- a fundamental requirement in experimental research(19). In addition, for variant

testing, the use of barcoding is restricted to testing single modifications or short

genomic regions in proximity to the barcode due to limitations on sequencing lengths.

Another ability lacking from current barcoding techniques, which prevents them from

being used to study particular adaptations, is the ability to *directly* single out specific

clones for further study. Currently, if particular clones are found of interest, based on the

dynamics of the population, the procedure for distinguishing them for further study

requires first separating single clones using other methods such as streaking or sorting

and then sequencing the barcode of each colony or well to identify the variant. This

procedure can be particularly cumbersome if one is looking for clones that are present

at low frequencies, as the probability of finding a clone goes as its percentage in the

population. Thus, a clone present below 1% requires (on average) more than 100

colonies to be sampled and sequenced, a feat that is both formidable and expensive.

To tackle some of the above limitations, we set out to simplify and streamline the

barcoding process so that it can be applied to a broader range of studies by

incorporating the following capabilities, (1) generation of identical library duplicates, (2)

direct variant identification, (3) precision and modularity in the insertion point of the

barcode and in the number of participating lineages and cells and, (4) potential for

integration into different organisms. In this paper, we describe the method we devised to

achieve these goals, which we named gUMI-BEAR (genomic Unique Molecular

Identifier-Barcode Enriched Associated Regions) and show how it can be utilised to

accomplish high-resolution lineage tracking at a relatively low cost as well as to directly

test millions of variants of a gene in the context of a growing population. While it can be

adapted to other systems, the setup we describe here is particularly applicable to

asexual populations. We note that this method can be applied by any lab versed in

standard molecular biology techniques and thus hope that it will be utilised by labs

specialising in different areas.

Results

Building a barcoded library

gUMI-BEAR is based on the unsupervised barcoding of each cell in an initial population

with a construct, the gUMI-box, that is inserted into the genome using CRISPR-Cas9.

The construct includes a barcode, along with auxiliary elements required for successful

transformation and ones that simplify downstream preparation of the samples for

sequencing (see Fig 1 and Methods).

Fig 1. Schematic illustration gUMI-BEAR

An illustration demonstrating the method in its entirety, from library construction through experimental

design, ending in the preparation of libraries for deep-sequencing. **(a, b)** An initial cellular population is

barcoded by inserting unique 24 bp sequences into its genomes using CRISPR/Cas9 (centre top). (**c**)

Following recovery from transformation, the desired number of live cells are sorted from the population,

(**d**) are allowed to grow, and (**e**) the resulting population is divided into aliquots containing equal numbers

of each barcoded lineage. The aliquots are subjected to the chosen experiment (centre middle), after

which (**f**) the cells undergo DNA extraction. (**g**) A two-step polymerase chain reaction (PCR) targeting the

barcode region (centre bottom) enables amplification of each lineage-associated variant (top) and

subsequent sequencing of the barcode library (bottom).

 <u>Donor and gUMI box construction</u>

The barcode (gUMI) region of the gUMI-box we used contains a 24 bp random DNA

segment preceded by five non-guanine nucleotides to allow efficient cluster generation

by Illumina sequencers. This length of the random segment can be changed according

to the requirements of each experiment, provided that the minimum of 25 base-pairs for

cluster generation is kept. The barcode region is flanked by elements that direct the

construct to its proper genomic locus upon transformation as well as components that

shorten the preparation of samples for sequencing after the experiment (see Fig 1).

Thus, upstream of the gUMI, we added a homology sequence for insertion into the

proper location in the genome (Left Homology Arm, LHA) and the Illumina READ1

sequence that renders the segment ready for attachment of Illumina P5 adapters after

the experiment(20). Downstream of the gUMI we placed a generic linker sequence, to

which READ2 will be attached following the experiment. The gUMI box is completed by

a 20bp sequence that is complementary to a Hygromycin resistance cassette (Fig. 2a).

Fig 2. Construction of the barcoded library

(a) Schematic illustration of the donor DNA construction process **(b)** The number of colonies formed following transformation with donor DNA with (HDR+CRISPR) or without (HDR) a pCAS vector carrying the CRISPR/Cas9 machinery for five biological replicates of each treatment. Utilisation of CRISPR/Cas9 increased the number of colonies 20-fold (t-test between distributions, p = 3.89e-8). **(c)** Percentage of dead cells in a culture (left y-axis) as a function of time following transformation with (circles) and without (squares) donor DNA. Samples were taken at several time points to check membrane integrity using Propidium Iodide staining during the 40-h following transformation for quantification of the percentage of dead cells. The right y-axis presents optical density (OD; orange line) measurements for the culture showing that cells start to grow after 24 h **(d)** Four replicates of a library, A–D, were deep-sequenced to reveal their populations' barcode composition and the log-transformed frequencies of the lineages are

plotted against each other. Pearson correlations were computed between all replicates and the Pearson

coefficient, r, is presented at the bottom of each comparison. Plots on the diagonal present the log-

transformed frequency distribution for each replicate (***p < 0.0001). (**e)** Bar plot of expected and

observed lineage frequencies for two control populations, Mock1 (upper panel) and Mock2 (lower panel).

Eight lineages were isolated from single colonies, and their barcodes were identified using Sanger

sequencing. Their DNA was extracted and mixed in known ratios to create the Mock1 and Mock2 control

populations. Each population was sequenced on two separate occasions, and the results underwent the

same analysis pipeline to reveal its population composition. Each colour represents a single lineage

according to the legend to the right of the graph.

The full donor that is transformed into yeast cells is composed of the gUMI-box,

together with the hygromycin cassette under the TEF1 promoter, and the 3' homology

sequence for genomic insertion (Right Homology Arm, RHA). It is formed by PCR

amplification using the gUMI-box as the forward primer together with a reverse primer

that consists a 20bp region complementary to the Hygromycin resistance cassette box

together with the RHA sequence (Fig. 2a).

As stated, we designed the gUMI-box to also aid the process of preparing the library for

sequencing. We used the linker in a short-cycle PCR step to attach a UMI and a READ2

sequence to each amplicon (bottom centre of Fig. 1). This endows each barcode that

enters the analysis pipeline with a tag (Fig 1g) and prepares it for index attachment.

Then, a second PCR reaction is used to amplify the barcodes to the amount required for

sequencing. The tag introduced in the first PCR provides a means to avoid biases that

may be introduced in the multiple cycles required to produce enough DNA from small

samples and thus increases the accuracy of lineage frequency quantification.

Transformation, sorting and expansion:

To build a modular system that could potentially be applied to different organisms as

well as to any genomic location, we decided to incorporate the gUMI using CRISPR

double-strand break at the integration site(21,22) together with the native yeast

homology-directed repair (HDR) system. Although, in yeast, HDR is usually sufficient for

the integration of desired DNA into the genome by transformation, we found that the

number of successful transformants increased 20-fold (Fig 2b) with the addition of the

CRISPR system, which facilitates the preparation of larger libraries. Equally important,

using CRISPR allowed us to easily change the locus of integration by simply replacing

the gRNA in the pCAS vector and the homologous arms of the donor DNA while leaving

the gUMI intact, thus satisfying our demand for flexibility in application.

To characterise the recovery of the barcoded population following transformation, we

measured the optical density (OD) and the percentage of dead cells in the culture at

several time points up to 40 hours post-transformation (see Methods). We found that

after 20–24 h of recovery, the percentage of dead cells started to decrease while the

OD began to increase (Fig 2c). We thus identified 24 hours as the inflection point,

where each barcode is still unique, and live cells in the population have the ability to

divide such that single cells can produce distinct, identifiable lineages.

One of the important prerequisites we set for the system was to be able to produce

identical copies of libraries to allow comparisons between experiments. This was

achieved using the following procedure. First, we sorted out the desired number of live

cells at the time point we identified previously as the one at which single cells produce

distinct lineages. Then we allowed the required amount of time for the cell population to

expand sufficiently to produce 100 copies of each barcoded yeast (approximately 32

hours) and, finally, divided the culture into 100 aliquots and froze them in glycerol stock

for the experiments. We note that the number of library copies produced can be

changed by modulating the time the culture containing the sorted cells is allowed to

expand and the number of aliquots accordingly. However, care must be taken to keep

the expansion time short enough for the probability of acquiring mutations before the

start of the experiment to be small. A detailed description and protocol of the

transformation process can be found in the methods section.

<u>Validation of the barcoding and analysis procedure</u>:

To assess the effect of various library preparation parameters on lineage makeup, we

created seven mini-libraries that varied in the following parameters: the number of

transformations performed (1 or 10), the number of cells sorted (10×10^3, 55×10^3,

110×10^3) and the expansion times after transformation (31h or 34 h) (Supplementary

Table 1). The compositions of the populations in the resulting mini-libraries were

determined by deep-sequencing the barcodes of four random aliquots from each mini-

library. The distribution of barcodes and the similarity between aliquots were unaffected

by transformation parameters nor by cell sorting (Supplementary table 1),

demonstrating the robustness of the procedure. The number of viable lineages directly

correlated with the number of cells sorted, although the precise number of lineages in

the population was ten times lower than the number of lineages sorted. To maximize the

number of barcodes in the population for further experiments, and given the above

similarities, we pooled all mini-libraries into a larger one, which was then divided into

100 aliquots and frozen until used. To analyse the barcode composition of the resulting

pooled library, we sequenced four of the aliquots. We found that these contain nearly

identical barcode contents distributed log-normally, as shown in Fig 2d. Almost all

26,000 lineages were observed in all four samples (two were missing from one sample),

with lineage frequencies exhibiting high between-aliquot correlations (r = 0.990–0.999; p

< 0.001).

Next, we wanted to verify that our experimental procedures (as described in Methods)

produce an accurate description of the population. We, therefore, constructed two mock

communities of known composition and subjected them to our analysis pipeline. To

generate the mock populations, we first plated our library, picked eight colonies, and

used generic primers spanning the barcode region to amplify and sequence (Sanger)

the barcode of each colony. DNA was then extracted from the identified colonies and

was mixed at two different ratios to construct two mock mixtures simulating different

lineage proportions. Mock1 included equal proportions of all eight lineages, and Mock2

was composed of a non-uniform distribution, 50% of which is made up of lineage F and

the remaining by equal proportions of the residual seven lineages. Fig 2e presents the

expected population structures (bottom illustrations) along with the ones obtained by

sequencing their barcodes on two separate occasions. The average error observed for

the percentage of a lineage within a population is 1.8%, with lineage F showing the

greatest error in Mock2 (7.5%). The fact that errors are almost identical when

comparing the two different sequencing runs indicates that the variations from the

expected structure are due to minor inaccuracies in the mock populations' construction

process rather than sequencing biases and errors.

Application of gUMI-BEAR in two experimental settings:

<u>Tracking evolutionary dynamics</u>

We first tested our system in a microbial evolution experiment setting, which is an area in which cellular barcoding has provided considerable insights (23,24). These types of experiments are usually carried out by serial dilution of a barcoded microbial population or in chemostats However, we wanted to retain as many barcodes as possible by reducing the elimination of low-frequency lineages by repeated bottlenecks, and thus we maintained the populations in turbidostats in a 100ml volume. In this setup, the cell density is maintained within a narrow window, in our case at an OD between 1.8-2.0, by diluting the cultures as soon as they reach the upper limit of the range until they reach the lower one. The cells reach exponential phase at the beginning of the experiment and are maintained in that growth state throughout, receiving rich media for the entirety of the experiment at a quantity that depends on their growth rate.

Cells were grown in the turbidostats in YPD media for 44 days while temperatures were varied three times to induce changes in the structure of the population. These were expected to occur due to differential fitness changes in individual lineages in the population(6,12,14,25,26) . We initially grew the population at 30 °C for two days, raised the temperature to 39 °C for eleven days, then to 41 °C for 27 days, finally reducing it back to 39 °C for the remaining four days of the experiment. Samples were collected throughout the experiment and sequenced to determine the clonal makeup of the population at each time point.

We tested the sequencing depth used in the analyses to rule out an inadequate one,

which would make the results inaccurate, or oversampling, which would increase

sampling costs unnecessarily. To this end, we constructed rarefaction curves for both

the number of unique lineages detected and their distribution using an increasing

number of reads at different time points along the experiment. As seen in Fig 3c and 3d,

for all time points tested, both parameters plateau quickly at a low percentage of reads,

indicating that our sequencing depth is sufficient. Since the distribution of lineage

frequencies is more uniform at the early stages of the experiment (Fig 3d, inset), the

number of reads at which it reaches the plateau increases with time but is still well

below our sequencing depth of ~10 million reads per sample.

Fig 3. Applying the gUMI-BEAR method to track evolutionary dynamics

(**a, b**) Lineage trajectories throughout a 44-day experiment in which temperatures were altered (vertical

dotted lines) to induce fitness fluctuations in the population. (**a**) Each lineage is represented by a line.

Colours were assigned to each lineage based on K-means clustering performed on their trajectories

throughout the experiment (as seen in the key). (**b**) Muller plot where each lineage is represented by a different colour, and the width of the coloured region at each timepoint is proportional to the lineage's relative frequency. (**c, d**) Rarefaction curves for the number of lineages (**c**) and lineage distributions (**d**) obtained by sampling an increasing number of reads at five time points (depicted by the different colours as defined in the key). (**d**) The accuracy of the distribution is quantified by the Wasserstein distance between the distribution obtained with a reduced number of reads and the original distribution (based on 100% of the reads). The inset shows a ridge plot of the frequency distribution of barcodes in the population quantified by sampling an increasing number of reads (y-axis, percentage of population sampled) from Day 1 (blue) and Day 44 (red).

Using our analyses, we were able to follow a population that initially consisted of 26,000 distinct lineages (Fig 3c) distributed log-normally (Fig 3d, inset) and reveal frequency changes of individual lineages. The size of the population dropped to 16,000 after the first day, after which the population underwent several cycles of clonal soft sweeps in which distinct lineages rose in frequency to become dominant (Fig 3a, b), with other lineages going down or remaining at low frequencies. The three temperature shifts induced sharp changes in population structure, the most dramatic manifestation of which was the immediate decline in abundance of the lineages that had been most successful under the previous condition, indicating that they had lost fitness relative to others. Surprisingly, this also occurred when we increased the temperature from 39 °C to 41°C, implying that adaptation to high temperatures does not necessarily occur gradually. Notably, despite the high level of stress, ~2500 of the lineages (~15%) were not diluted out and remained in the population throughout the entire 44 days of the experiment (~128 generations). These results indicate that we were able to follow

changes in the structure of a population consisting of a large number of barcodes at

very low frequencies.

<u>Large scale tracking of gene variants</u>

Next, we wanted to show how gUMI-BEAR can be used for large-scale mutational

screening, where each variant is tagged by a unique barcode, and its competitive

success can be measured in terms of barcode abundance. We chose *Hsp82* to

demonstrate the method's capability to screen genes that are longer than the NGS read

length and because the protein it encodes affects fitness gain and loss via variability

control through chaperone-like activity(27,28). The number of possible *Hsp82* double

mutants is over $2X10^6$ and directed mutagenesis is unrealistic because of the work and

cost involved in producing and testing each variant. However, we were essentially not

limited in the number of variants we tested, which gave us freedom from the need to

design specific mutants and allowed us to, instead, use a random mutagenesis kit

(Genemorph II by Agilent part number 200550, see Materials and Methods). We

performed mutagenesis on the *Hsp82* gene amplified from the *S. cerevisiae* BY4741

strain using the Genemorph kit (forward primer 100 bp upstream of the gene; reverse

primer 100 bp downstream of the gene), with PCR parameters adjusted to produce 0-

4.5 mutations per kb (annealing at 60 °C, 25 cycles, 2 min elongation time, and 90 ng

genomic DNA template).

The resulting variants were conjugated to the gUMI-box, this time with the LHA and

RHA complementary to the *Hsp82* gene. This created a donor molecule ready to be

transformed into the native locus in the yeast genome, replacing the wild-type *Hsp82*

(Fig 4b and Methods). Identical aliquots of the library were prepared in a manner similar

to that described above, which allows exact repeats to be performed as well as to test

the library of variants in other settings. Overall, we produced a library of 1,745,260 yeast

clones, each carrying a randomly mutated and uniquely barcoded *Hsp82* variant.

Fig 4. Applying gUMI-BEAR to track gene variants

(a) Heatmap showing entropy values (4 upper bands, as per the colour bar) for mutations in each

nucleotide in the *Hsp82* variants for various steps in the library construction process. For each position,

mutational proportions were normalised to sequencing depth, and Shannon's entropy was calculated

(dark to bright represents low to high entropy). From top to bottom: The Genemorph panel refers to

Hsp82 variants created by insertion of random mutations using PCR with the Genemorph II kit. The

Donor panel refers to the final donor molecule before transformation (mean entropy of three replicates).

The panels labelled Start and End show the population at the start of the experiment, where cells have

been transformed with the donor molecules and selected for successful transformants (mean of three

replicates), and at the end of the experiment, respectively. The Protein panel shows mutational

occurrence in the translated protein obtained by long-read sequencing of full-length variants (mutations

are identified in white). The Conservation panel shows the conservation score for each position calculated

using the ConSurf[18] server (dark for conserved and bright for variable regions). The insets show two

zoomed-in regions exhibiting different patterns where both conserved and variable regions can have non-

synonymous mutations. (**b**) Donor construction process as described in the Materials and Methods

section. GenemorphF and GenemorphR are primers used to amplify the native *Hsp*82 from the genome;

LHA and RHA stand for Left/Right Homologous Arm and are added to enable correct transformation to

the chosen locus; Gumy2Hyg is a forward primer containing all elements that are added to the donor

molecule including the illumina read1 sequencing binding site (Rd1), a five non-guanine nucleotide

sequence (5xH), the genomic unique molecular identifier (gUMI) and the linker sequence which is used

for later preparation of the barcode molecule for sequencing. Briefly, *Hsp82* variants were produced by

PCR amplification of the gene from the *S. cerevisiae* BY4741 strain using the Genmorph II kit. These

were then conjugated to the gUMI-box, with LHA and RHA complementary to the *Hsp82* gene, which

allowed them to replace the wild-type in the CRISPR/Cas9 reaction (c) The Hsp82 protein dimer (shown

in green and purple). All mutations found by long-read sequencing are shown as red and cyan balls in the

Hsp82 dimer. The image was produced with the Protein Homology/Analogy Recognition Engine (Phyre2).

(**d**) Changes in relative lineage abundances within the yeast population over time. Each line represents

the mean relative abundance of a lineage in the four replicates as a function of time, with error bars

showing the standard deviation between replicates. Red lines mark the seven lineages whose *Hsp*82

variant was amplified using the gUMI sequence as a reverse primer and then Sanger-sequenced. (**e**) A

scheme presenting Sanger sequencing using five forward generic primers (black arrows) distributed

along the *Hsp82* ORF with a specific gUMI reverse complementary primer that was used to obtain a

constructed full-length sequence of the variant (Red). The constructed variants are compared to the full-

length sequence of the variant achieved by long-read sequencing (Loop Genomics) in green (Identified

SNPs are indicated by the white lines).

The ability to test randomly mutated variants has the advantage of not introducing bias

in their design. We wanted to test the effect of the assembly process on the distribution

of mutations and, therefore deep-sequenced amplicons from two stages: after

mutagenesis of the gene and after assembly of the final donor molecule containing the

gUMI-box. A comparison of the sequences of the *Hsp82* variants emerging from the

mutagenesis step and of the donor molecules revealed a decrease in the number of

variants after integration of the ORF. This is to be expected, as the mutagenesis

process is performed in a sequence agnostic manner, whereas genomic integration has

biological constraints and, as such, eliminates some mutations. We then calculated the

distribution of Shannon's information entropy of mutated nucleotides per site (see Fig 4a

and supplementary Fig 4). This distribution was identical to that of the molecules

emerging from the Genemorph process showing that integration into the gUMI-box

during the donor DNA assembly process does not introduce bias (Fig 4a). The mutated

ORF within the donor molecules displayed the same entropy distribution in three

independent assembly replicates, confirming that the process of integration into the

gUMI-box was robust.

To allow variants to compete with each other, we used batch mini-cultures to grow four

replicates at 30°C, sampling and diluting at a rate of 1:100 every 12 hours, for a total of

120 hours. Samples were sequenced to yield the relative abundance of each lineage as

a function of time (Fig 4d). We found the distribution of lineage frequencies to be heavily

skewed throughout the experiment, with 50 lineages comprising almost all (99%) of the

population. Of these, lineages in ranks 25 - 50 were each present at a relative

abundance of 1% or less and all were successfully tracked by the gUMI-BEAR method.

While the frequency of some lineages increased or decreased in the experiment, the

abundance of most lineages remained constant, indicating relatively small fitness

differences. The trajectory of the most successful lineages, namely those ranked among

the top 50 in terms of their abundance, was reproducible between replicates, as

indicated by the high correlations between different replicates (r = 0.89 ± 0.034) and the

small standard deviation (Fig 4d). This signifies that the dynamics were determined by

the variants that entered the experiment and not by adaptations that occurred

throughout it, which could differ between repeats.

To demonstrate the capability of the system to identify variants of interest, we chose

seven lineages for further analyses. As a first step, we used their reverse gUMI

sequences as PCR primers to isolate specific variants. We were able to sequence a

variant ranked as low as the 47th rank in a sample taken after 108 hours, constituting

0.19% of the population, demonstrating how even low-frequency lineages can be

directly isolated from a mixed population (Fig 4 d, e). The complete sequences of the

Hsp82 variants were then determined by Sanger sequencing using five primers

distributed along the gene (Fig 4e). A comparison of the sequences obtained in this

manner with sequences from full-length third-generation sequencing (Loop Genomics)

confirmed the quality of variant sequencing (Fig 4e). Thus, with a single PCR reaction,

performed on the entire population, and five Sanger sequencings, we were able to

distinguish a variant that, using present methods, would require the sequencing of 500

single-cell colonies, *on average*. Having identified variants of interest, one can go on to

test them in isolation by inserting the relevant ORF into wild-type strains.

When we compared short reads obtained from the initial yeast library, after

transformation and expansion, with the long reads from the population after it had

undergone 120 hours of evolution (Fig 4a), there was an overall decline in entropy,

indicating that the experiment began with numerous mutations holding reduced fitness

which were eliminated as growth continued. In the final library, we also found regions

where no mutations had occurred, presumably because mutations in these areas

reduce fitness significantly. The process of growing a population that harbours random

mutations in a gene thus provided a biological filter, at evolutionarily relevant conditions,

that eliminated lethal or function-harming mutations while allowing the propagation of

mutations that were neutral or that provided a fitness advantage.

Discussion

In this work, we describe a single-cell barcoding method that is versatile, easy to

implement, and cost-effective, thus lending itself to a host of experimental scenarios.

These traits are achieved due to a combination of features: Firstly, we used

CRISPR/cas9 coupled with the cellular HDR to integrate the barcode containing gUMI-

box, thereby achieving a high yield of transformants which in turn allows the

construction of libraries of extremely large size. Furthermore, the gUMI-box integration

site can easily be changed by simply designing a new gRNA and a corresponding donor

DNA. Secondly, we perform a sorting step following recovery from the transformation to

obtain control over the number of lineages participating in the experiment. Importantly,

our method allows the assembly of multiple identical replicates of the same initial

population, opening the way for performing precise repeats and for comparisons

between the behaviour of the same populations of cells under different experimental

scenarios. This, in turn, makes it possible to apply barcoding towards investigating the

role of randomness and determinism in highly complex biological systems(22,29,30).

The gUMI-box, together with the simple and efficient protocol we applied to prepare

gUMI-barcoded sequences for deep-sequencing (Fig 1g), eliminates the need for

multiple preparation steps and, at the same time, provides high resolution and accuracy

and supports the use of small samples.

We presented two applications to demonstrate the potential and adaptability of the

gUMI-BEAR method. The first application exhibited its scale and precision in tracking

~16,000 lineages through 30 time points over a period of 44 days and revealed detailed

evolutionary dynamics resulting from the complex environmental regime. We obtained

an intricate picture of the rich population dynamics that arise as multiple lineages adopt

numerous modes of exploration and exhibit varying adaptabilities to changing

environmental conditions. Attaining such a picture through standard population studies,

such as bulk RNA sequencing, whole-genome sequencing, or plating assays, would be

financially and/or analytically impractical.

In the second experiment, we used gUMI-BEAR to explore gene-variant fitness and

dynamics of populations with induced fitness variations. By coupling a gUMI to each

variant, we were able to scan a large number of random mutations produced *in vitro* in

an evolutionarily relevant context in which they compete with each other, eliminating the

need for either fitness testing of monoclonal variants or costly in-depth sequencing.

Due to our ability to produce identical libraries, we were able to observe largely

deterministic dynamics for this population. Importantly, we used reverse-gUMI

sequences as PCR primers to target specific variants and then used Sanger

sequencing to identify the variation without the need for laborious work to single out

variants and expensive long-read sequencing, thereby alleviating the previous limitation

on the complexity of variants that could be tested in parallel methods.

Our method thus allows in-depth exploration of millions of variants of any complexity, all

living and competing as a population, while also providing the ability to explore

interesting variants, even at very low frequency, using simple techniques such as PCR

and Sanger sequencing.

We note that the gUMI-BEAR method can be utilised for variant screening in more

complex systems, consisting of multiple genes, by adding a short sequence to identify

each ORF. Under such a configuration, the relevant genes can be multiplexed to create

an array of randomly mutated genes interacting with each other to enable the study of

complex enzymatic pathways.

Using our ability to produce identical libraries, one can test the dynamics of the same

populations or variants under different treatments. This is relevant as fitness may

depend on environmental factors. Thus, our method allows studying gene-environment

relations on a large scale.

Additional settings where the gUMI-BEAR method can be implemented are the Pharma

and Biotech industries, particularly during the scale-up of fermentation processes and

during the production phase. Specifically, it could be used to identify deleterious

processes before they affect production or to single out the key events that impair large-

scale production and find ways to overcome them (31).

We designed gUMI-BEAR to be an easy-to-use and robust method to track changes in

population makeup and variant fitness. Previously available methods(1,3,10) to track

such changes were not scalable and were difficult to deploy on new lines and modalities

or more costly. A library can be created using the gUMI-BEAR method in a few days

and our results show that it provides an accurate, cost-effective, and versatile means of

tracking subtle changes that propagate into major population shifts in a way that can

easily be adapted to other biological systems.

Materials and Methods

The methods described below were common to both of the experiments conducted,

except where otherwise noted.

Construct assembly

gUMI-box (Genomic Unique Molecular Identifier)

The gUMI-box contains a 24 bp random sequence that serves as a transmissible

barcode for each lineage in the population. The gUMI is flanked by the read 1 Illumina

generic sequence on one side and a 15 bp generic sequence called a linker on the

other side. The linker is used as a target sequence for DNA amplification in the deep-

sequencing library preparation step, such that two-step PCR produces a library

containing all elements necessary for sequencing on Illumina platforms (Fig 1g). An

additional hygromycin-resistance cassette (HGmx6; obtained from the vector

pAG32(32)) is added downstream of the linker sequence to allow selection and to

prevent contamination (Fig 2a). As a final step, two homologous arms are added to the

gUMI-box upstream and downstream of the genomic integration site. All elements are

combined into one linear DNA construct, the gUMI-box, which is then transformed into a

population of yeast cells.

Donor construction for tracking evolutionary dynamics

Donor DNA was built using two primers containing 50 bp sequences inserted upstream

(left homologous arm; LHA) and downstream (right homologous arm; RHA) of the

integration site in the *YBR209W* locus, which encodes a putative protein of unknown

function and a non-essential gene(33). The forward primer consists of the specific LHA

and most components of the gUMI-box. At its 3' end, it contains a complementary

sequence to the HGmx6 cassette from pAG32. The reverse primer consists of the

specific RHA and a sequence complementary to the end of the HGmx6 cassette. PCR

using these two primers produces the full linear DNA used for transformations in later

steps (Fig. 2a). The double-strand break site was determined using CHOPCHOP(22).

The pCAS plasmid was cloned by restriction-free (RF) cloning with two primers

containing the gRNA sequence (5`-TAGAGCGTCAATCAAGAAAG-3`) as described

previously[13.]

<u>Donor construction for large scale tracking of gene variants</u>

Six-step PCR was utilized to obtain a full-length donor DNA comprising 5'-LHA-d*Hsp82*-

gUMI-box-RHA-3'. *HSP82* was amplified from *S. cerevisiae* BY4741 gDNA to obtain a

wild-type copy of the gene. Random mutations were introduced using the Genemorph II

random mutagenesis kit (Agilent, 200550). To achieve a low mutation rate (0–4.5

mutations/kb), 1 µg of the template DNA was amplified using this kit in a 30-cycle PCR

protocol. A series of PCRs containing the gUMI-box and the **Hsp82* were performed to

create the full-length construct (Fig 4b and Supplementary methods).

Two pCAS plasmids were cloned by RF cloning, each with two primers containing the

gRNA sequence for the cleavage site upstream (5'-CAAACAAACACGCAAAGATA-3`)

and downstream of the *Hsp82* gene (5'-AGCTGACACCGAAATGGAAG-3`)

Transformation

<u>For tracking evolutionary dynamics</u>

Competent cells of *S. cerevisiae* BY4741 were prepared and each resulting aliquot was

transformed with donor DNA (5 µg) and cloned pCAS (2 µg) using the protocol

described by Ryan et al(22,34). Following a three-hour recovery in 450 µL of YPD

media at 30 °C, aliquots were mixed and incubated for 20 h in 100 mL YPD broth with

hygromycin B (300 µg/mL) at 30 °C at 220 rpm. To accurately determine the number of

lineages involved in each experiment, following the 20 hr recovery phase, the cells were

stained with propidium iodide[27] to mark perforated cells and the desired number of

viable cells was sorted. The library was grown in 100 mL YPD with hygromycin B and

was allowed to propagate for 32 h (~16 generations). This step was followed by the

division of the library into 100 identical aliquots that were then immediately frozen in

50% Glycerol at -80 °C (Fig 1a). To examine the effects of several parameters on the

barcoded library structure we performed the abovementioned process seven times,

varying the number of competent cells used to build each library, the time allowed for

recovery following transformation, total number of cells sorted following recovery, and

expansion time following recovery. Supplementary table 1 presents the parameters

used in the seven libraries we built.

<u>For large scale tracking of gene variants</u>

Cells were transformed as described above. Following recovery, three aliquots were

incubated for 20 h in 10 mL YPD with hygromycin B at 30 °C at 220 rpm to select

transformants. The library was diluted 100-fold and was allowed to propagate for an

additional 30 h. This step was followed by the division of the library into 100 identical

aliquots that were immediately frozen in 50% Glycerol at -80 °C.

Experimental outline

<u>Tracking evolutionary dynamics</u>

The experiment was performed in a turbidostat (Eppendorf DASbox Mini Bioreactor

System) at a constant volume of 100 mL YPD supplemented with hygromycin B (200

µg/mL). Genetic drift was kept to a minimum by maintaining the concentration of the

culture within a narrow range of 0.9–1.1 (AU), eliminating large dilution steps. Cells

were grown for a total of 44 days The initial temperature of the culture was set to 30 °C.

After 48 hours, the temperature was increased to 39 °C for eleven days, 41 °C for 27

days and reduced back to 39 °C for the remainder of the experiment (Fig 3a, b). A total

of 31 samples were taken throughout the experiment in three replicates, for DNA and

RNA extraction and glycerol stock of viable cells. To prepare the DNA/RNA samples for

later analysis we centrifuged each sample at 8.6g for 2 minutes, removed the

supernatant, and flash frozen them in liquid nitrogen at -80°C. Glycerol samples were

treated similarly, except that following supernatant removal, the cells were resuspended

in YPD with 25% Glycerol.

 <u>Large scale tracking of gene variants</u>

Four library aliquots were mixed to create a homogeneous population and to minimise

minor differences between aliquots. The mix was evenly distributed to four replicates

that were grown in 100 mL Erlenmeyer flasks containing 25 mL YPD with hygromycin B

(200 µg/mL). Environmental conditions were maintained at a constant temperature of 30

°C and constant mixing at 220 rpm on a magnetic stirrer. The experiment spanned five

days such that every 12 hours, as cells approached the end of their exponential growth

phase, a sample was taken, and the culture was diluted 1:100.

579 Sample preparation and sequencing

DNA was extracted using the MasterPure Yeast DNA purification kit (Epicentre,

MPY80200), and concentrations were measured by Qubit. A sub-sample (150 ng) was

taken from each sample to serve as a template for four-cycle PCR in which the barcode

region was linearly amplified, and the internal index and UMI were added (Fig 1g). Five

samples from this reaction were pooled on a Qiagen PCR clean-up column (Cat. 28104)

and eluted in a 50 µL elution buffer. These samples were further cleaned by 1.8×

Ampure XP beads (Cat. A63881). From each pool, 10 µL were used for a 12–17 cycles

PCR using Illumina indexed primers. Samples were purified using 1.8× Ampure XP

beads, and their quality was evaluated using Agilent Tape-station (Cat. 5067-5584) and

Qubit concentration measurements (Cat. Q32853). All samples were diluted to a

concentration of 0.5 ng/µL, and 10 µL of each diluted sample was used to create the

final library. This sample was further diluted to a final concentration of 2 nM in a final

volume of 100 µL. The library was sequenced at the Weizmann Institute of Science's G-

INCPM unit on an Illumina NovaSeq platform with the following cycles: 29|10|10|26.

Preliminary analysis of raw data

Raw data were filtered to remove low-quality reads. Demultiplexing of samples was

performed in two parts using Illumina indices and the internal indices. In order to count

lineage frequency and determine which barcodes originated from a cell and which from

sequencing mistakes, the CD-HIT(35,36) algorithm was used in two steps. In the first,

seed sequences are created from samples taken throughout the experiment. Next,

sequences from each specific time point were compared with the seed sequences list

(using the cd-hit-est2d function). In each step, we allowed up to a two bp mismatch

before assigning the sequence to a new seed. The CD-HIT function clst2txt.pl was used

to create lineage count data-frames for each sample. The final output of this process

contained many clusters of sequences that were each classified as a single lineage for

further analysis. In the final step of the preliminary analysis, custom, in-house R-code

was used to pool together all sequences from the same cluster with the same UMI to

one cell. The final output of the abovementioned pipeline accurately measured the

number of actual, single cells originating from a specific lineage for every time point

after correction of PCR biases. For full script see rezenman/gumi_bear_code github page

Acknowledgements

We thank Shira Holand for the graphic design, Yonatan Nutkewitz for helpful

discussions and help with experiments and Hadas Keren-Shaul for help with next-

generation sequencing.

References

1. Shaffer SM, Dunagin MC, Torborg SR, Torre EA, Emert B, Krepler C, et al. Rare cell variability and drug-induced reprogramming as a mode of cancer drug resistance. Nature. 2017 Jun 7;546(7658):431–5.

2. Locke JCW, Young JW, Fontes M, Hernández Jiménez MJ, Elowitz MB. Stochastic pulse regulation in bacterial stress response. Science. 2011 Oct 21;334(6054):366–9.

3. Ackermann M. A functional perspective on phenotypic heterogeneity in microorganisms. Nat Rev Microbiol 2015 138. 2015 Jul 6;13(8):497–508.

4. Raser JM, O'Shea EK. Noise in Gene Expression: Origins, Consequences, and Control. Science [Internet]. 2005 [cited 2017 Jun 20];309(5743). Available from: http://science.sciencemag.org/content/309/5743/2010.full

5. Pilpel Y. Noise in Biological Systems: Pros, Cons, and Mechanisms of Control. In: Methods in molecular biology (Clifton, NJ) [Internet]. 2011 [cited 2017 Jun 29]. p. 407–25. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21863500

6. Nguyen Ba AN, Cvijović I, Rojas Echenique JI, Lawrence KR, Rego-Costa A, Liu X, et al. High-resolution lineage tracking reveals traveling wave of adaptation in laboratory yeast. Nature. 2019 Nov 21;575(7783):494.

7. Oliver KM, Russell JA, Moran NA, Hunter MS. Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. Proc Natl Acad Sci U S A. 2003 Feb 18;100(4):1803–7.

8. Birgy A, Roussel C, Kemble H, Mullaert J, Panigoni K, Chapron A, et al. Origins and breadth of pairwise epistasis in an α-helix of β-lactamase TEM-1. bioRxiv. 2021 Nov 29;2021.11.29.470435.

9. Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, Sherlock G. Quantitative evolutionary dynamics using high-resolution lineage tracking. Nature. 2015 Feb 25;519(7542):181–6.

10. Rodriguez-Fraticelli A, Morris SA. In preprints: the fast-paced field of single-cell lineage tracing. Development. 2022 Jun 10;149(11):dev200877.

11. Cvijović I, Nguyen Ba AN, Desai MM. Experimental Studies of Evolutionary Dynamics in Microbes. Trends Genet TIG. 2018 Sep;34(9):693–703.

12. Blundell JR, Levy SF. Beyond genome sequencing: Lineage tracking with barcodes to study the dynamics of evolution, infection, and cancer. Genomics. 2014 Dec 1;104(6):417–30.

13. Jasinska W, Manhart M, Lerner J, Gauthier L, Serohijos AWR, Bershtein S.
Chromosomal barcoding of E. coli populations reveals lineage diversity dynamics at
high resolution. Nat Ecol Evol 2020 43. 2020 Feb 24;4(3):437–52.

14. Barrick JE, Yu DS, Yoon SH, Jeong H, Oh TK, Schneider D, et al. Genome
evolution and adaptation in a long-term experiment with Escherichia coli. Nature.
2009;461(7268):1243–7.

15. Park SC, Krug J. Clonal interference in large populations. Proc Natl Acad Sci. 2007
Nov 13;104(46):18135–40.

16. He Z, Maynard A, Jain A, Gerber T, Petri R, Lin HC, et al. Lineage recording in
human cerebral organoids. Nat Methods. 2022 Jan;19(1):90–9.

17. Li C, Zhang J. Multi-environment fitness landscapes of a tRNA gene. Nat Ecol Evol.
2018 Jun;2(6):1025–32.

18. Benjamin R, Giacoletto CJ, FitzHugh ZT, Eames D, Buczek L, Wu X, et al.
GigaAssay – An adaptable high-throughput saturation mutagenesis assay platform.
Genomics. 2022 Jul 1;114(4):110439.

19. Lobkovsky AE, Koonin EV. Replaying the tape of life: Quantification of the
predictability of evolution. Front Genet. 2012;3(NOV):246.

20. Wilkie G. Illumina adapter and primer sequences [Internet]. CVR Bioinformatics.
2015 [cited 2023 Feb 12]. Available from: https://bioinformatics.cvr.ac.uk/illumina-
adapter-and-primer-sequences/

21. Ryan OW, Skerker JM, Maurer MJ, Li X, Tsai JC, Poddar S, et al. Selection of
chromosomal DNA libraries using a multiplex CRISPR system. eLife. 2014 Aug
19;3:e03703.

22. Ryan OW, Poddar S, Cate JHD. CRISPR–Cas9 Genome Engineering in
Saccharomyces cerevisiae Cells. Cold Spring Harb Protoc. 2016 Jun
1;2016(6):pdb.prot086827.

23. Ba ANN, Cvijovic I, Echenique JIR, Lawrence KR, Rego-Costa A, Liu X, et al. High-
resolution lineage tracking reveals travelling wave of adaptation in laboratory yeast.
Nature. 2019 Nov 1;575(7783):494–9.

24. Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, Sherlock G.
Quantitative evolutionary dynamics using high-resolution lineage tracking. Nature.
2015 Mar 12;519(7542):181–6.

25. Messer PW, Petrov DA. Population genomics of rapid adaptation by soft selective
sweeps. Trends Ecol Evol. 2013 Nov 1;28(11):659–69.

26. Nguyen J, Fernandez V, Vicente I. Fernandez, Putri SP, Pontrelli S, Sauer U, et al.
A distinct growth physiology enhances bacterial growth under rapid nutrient
fluctuations. Nat Commun. 2021 Jun 16;12(1):3662–3662.

27. Cowen LE, Lindquist S. Hsp90 Potentiates the Rapid Evolution of New Traits: Drug
Resistance in Diverse Fungi. Science [Internet]. 2005 [cited 2017 Aug 5];309(5744).
Available from: http://science.sciencemag.org/content/309/5744/2185

28. Cote-Hammarlof PA, Fragata I, Flynn J, Mavor D, Zeldovich KB, Bank C, et al. The
Adaptive Potential of the Middle Domain of Yeast Hsp90. Mol Biol Evol. 2021 Feb
1;38(2):368–79.

29. Koonin EV, Wolf YI. Is evolution Darwinian or/and Lamarckian? Biol Direct.
2009;4:42.

30. Raj A, van Oudenaarden A. Nature, nurture, or chance: stochastic gene expression
and its consequences. Cell. 2008 Oct 17;135(2):216–26.

31. Olsson L, Rugbjerg P, Torello Pianale L, Trivellin C. Robustness: linking strain
design to viable bioprocesses. Trends Biotechnol. 2022 Aug 1;40(8):918–31.

32. Addgene: pAG32 [Internet]. [cited 2022 Jan 1]. Available from:
https://www.addgene.org/35122/

33. Fisk DG, Ball CA, Dolinski K, Engel SR, Hong EL, Issel-Tarver L, et al.
Saccharomyces cerevisiae S288C genome annotation: A working hypothesis.
Yeast. 2006 Sep;23(12):857–65.

34. CLONTECH. Yeast Protocols Handbook FOR RESEARCH USE ONLY Yeast
Protocols Handbook [Internet]. 2009 [cited 2022 Jan 1]. Available from:
www.clontech.com

35. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: Accelerated for clustering the next-
generation sequencing data. Bioinformatics. 2012 Dec;28(23):3150–2.

36. Li W, Fu L, Niu B, Wu S, Wooley J. Ultrafast clustering algorithms for metagenomic
sequence analysis. Brief Bioinform. 2012 Nov;13(6):656–68.

Supporting Information Captions

Supplementary Figure 1. Nucleotide composition by position for the gUMI from donor DNA and lineages

Donor DNA prior to transformation (upper panel) and DNA from lineages participating in the experiment (lower panel) were extracted and deep-sequenced to reveal the nucleotide composition for each position in the gUMI (x-axis). The overall height of each base is proportional to its probability (y-axis). Colours represent different nucleotides.

Supplementary Figure 2. Guanine-Cytosine content distributions for the gUMI from three stages in the experiment with no initial fitness variations

GC content was measured based on sequences derived from all lineages at three steps during library construction. (Left, red) Donor DNA prior to transformation. (Middle, green) Lineages at the beginning of the experiment. (Right, blue) Lineages at the end of the experiment. All data are presented in histograms to reveal the GC content distribution in the populations.

Supplementary Figure 3. Mean doubling time and glucose concentration for the experiment in population exhibiting no initial fitness variations

Doubling time for the whole culture was calculated based on dilution rate of the turbidostats and is presented on a log scale (left y-axis, orange line) and ploted as function of time. Glucose concentrations were measured for each sample taken (right y-axis, blue line). The experiment started at an optimal growth temperature of 30°C that was altered to induce stress, as noted by the doted lines.

Supplementary Figure 4. Entropy kernel density estimations for the *HSP82* ORF before and after incorporation into the gUMI-box

The kernel density estimations of the Shannon entropy for nucleotide substitutions in the *HSP82* ORF in the genenmorph amplicon after *in vitro* mutagenesis (black) and three replicates of the assembled donor DNA (green, red and blue) are shown. Three peaks can be observed - first at entropy 0, in which all substations were to the same

nucleotide, second at entropy 1, in which substitutions were equal between one of two

possible substitutions, and finally, the third peak at entropy 1.5 in which an equal

probability for all mutations is observed. The donor DNA distribution mimics the one

produced by the genemorph, indicating that no bias was induced during donor

construction.

Supplementary Table 1. Mini-library parameters

Seven different mini-libraries were prepared while varying the following parameters: No.

transformations - the factor applied to the volumes given in the protocol published by

Ryan et al.[1], namely, 1 µg pCAS, 5 µg donor DNA, and 90 µL competent cells. Hence,

No. of transformations = 10 indicates the use of 10 µg pCAS, 50 µg donor DNA, and

900 µL competent cells; T_{sel}, selection time, being the number of hours the library was

grown for after transformation and before sorting; No. cells sorted, total number (in

thousands) of viable cells sorted following staining with propidium iodide and selection

(see Methods); T_{exp}, expansion time, being the time (hours) the library was allowed to

expand before division into 100 aliquots; No. unique lineages, number of viable lineages

in each mini-library as determined by deep-sequencing.

Supplementary Figure 5. Protein-protein interaction models

a, HSP82 dimer shown in green and purple, Cdc37 cochaperone (pink), and the Cdk4

kinase (yellow) (PDB 5FWK). **b,** HSP82 dimer shown in green and purple and Sba1

Chaperone (in yellow and pink) (PDB 2CG9). All possible mutations are shown in the

HSP82 model as red and cyans balls for the two HSP82 homodimer.

Library preparation

a

b

c

Sorting live cells

d

Expansion

e

Identical aliquotes

Experimental setup

1. Reduced cycle PCR

| Rd1 | gUMI | linker | UMI | index | Rd2 |

2. Standard PCR

| P5 | Rd1 | gUMI | linker | UMI | index | Rd2 | P7 |

| Rd1 | gUMI | linker |

Barcode sequencing

f

Genomic DNA extraction

gRNA + crispr

CAS9

gRNA + crispr

g

Barcode amplification

P5

Index 1

Read 1 Sequencing Primer

Read 2 Sequencing Primer

Index 2

P7

24bp gUMI

a

Genemorph
Donor
Start
End
Protein
Conservaton

-0.4 0.0 0.4 0.8 1.2 1.6

b

Genomic DNA

GenemorphF   GenemorphF
                    hsp82
100bp   2130bp   100bp
                                GenemorphR

GenemorphF
                    hsp82
                                GenemorphR

LHA
                    *hsp82

Gumi2Hyg   HSP82_Read1
pAG32   hphMX
                                RHA

LHA | *hsp82 | Rd1 | Rd1 | 5xH | gUMI | linker | hphMX6 | RHA

LHA
LHA | *hsp82 | Rd1 | Rd1 | 5xH | gUMI | linker | hphMX6 | RHA
                                                              RHA

c

d

Relative frequency

0.08
0.06
0.04
0.02

0.084
0.0811
0.075

0.038

0.02

0.008
0.001

0 24 48 60 72 96 120
Sample time (hr)

e