Title

Multiple climate change factors jointly increased the competitiveness of a C_4 grass against its

C_3 competitor

Author names and affiliations

Gen-Chang Hsu[1,2], Po-Ju Ke[2], Chuan-Kai Ho*[2]

[1]Department of Life Science, National Taiwan University, Taipei, Taiwan

[2]Institute of Ecology and Evolutionary Biology, National Taiwan University, Taipei, Taiwan

***Correspondence:** ckho@ntu.edu.tw

Abstract

C_3 and C_4 plants are the two major terrestrial plant groups that respond differently to temperature, water, and CO_2. Although much research has focused on the individual effect of warming, water availability, and elevated CO_2 on biomass production and competitive interactions between C_3 and C_4 plants, the joint effects of these factors remains underexplored. We grew naturally co-occurring C_3 (*Oplismenus composites*) and C_4 (*Paspalum conjugatum*) grass under three temperature warming scenarios (control, +2°C, and +4°C), two water supply intervals (normal 2 days vs. prolonged 7 days [drought]), and two CO_2 concentrations (ambient 400 ppm vs. elevated 800 ppm) in growth chambers and measured their above- and below-ground dry biomass in monoculture and mixture to quantify their biomass performances and competitive responses. Warming and elevated CO_2 together enhanced the above- and/or below-ground biomass of both C_3 and C_4 grass. Moreover, temperature, water, and CO_2 interacted to increase the below-ground biomass of the C_4 grass. Surprisingly, the C_3 grass performed worse under interspecific competition (mixture biomass) relative to intraspecific competition (monoculture biomass) at elevated CO_2 compared to ambient CO_2, while the reverse is true for the C_4 grass. Furthermore, the C_4 grass was the most competitive against the C_3 grass under simultaneous 4°C warming, drought, and elevated CO_2. Taken together, our results suggest that the competitive balance could potentially shift in favor of C_4 plants under the projected temperature warming, increased drought frequencies, and rising atmospheric CO_2, with an increase in C_4 relative abundance in future plant communities.

Keywords

above- and below-ground biomass, C_3 and C_4 plants, competition, drought, elevated CO_2, global climate change, warming

Introduction

C_3 and C_4 plants are two major terrestrial plant groups that differ in their physiology and response to surrounding environments (Ehleringer and Monson 1993, Yamori et al. 2014). Their relative abundance in the community is closely related to various ecosystem functions and processes such as resource provisioning for herbivores, nutrient cycling, and carbon storage (Still et al. 2003, Warne et al. 2010, Pau et al. 2013). For example, C_3 plants generally have higher nitrogen-to-carbon ratios than C_4 plants (Barbehenn et al. 2004), and therefore their relative biomass production in the habitat may affect the nutritional availability for herbivores and even organisms at higher trophic levels (Warne et al. 2010). Importantly, because C_3 and C_4 plants evolved under different climates and have different adaptations to surrounding environments, understanding how they respond to climate change and the resulting dynamic shifts is crucial for the management of terrestrial ecosystems.

The performances of C_3 and C_4 plants under different environmental conditions can influence their relative competitive abilities (Pearcy et al. 1981, Edwards and Still 2008). Theoretical and empirical studies have suggested that C_3 plants generally perform better and are stronger competitors in cool and well-watered environments, whereas C_4 plants are more adapted to high temperatures and drought conditions and thus more competitive than in hot and arid environments (Teeri and Stowe 1976, Ehleringer and Monson 1993, Sage and Monson 1998, Edwards and Still 2008). In addition to temperature and water availability, C_3 and C_4 plants exhibit differential responses to CO_2 (Pearcy and Ehleringer 1984, Wand et al. 1999). The photosynthesis of C_3 plants are limited by current level of atmospheric CO_2, and increasing CO_2 concentration could stimulate their growth and therefore offer them competitive advantages over C_4 plants (Ward et al. 1999, Poorter and Navas 2003, Lee 2011). These shifts in C_3-C_4 competitive relationships in response to climate change may alter their relative abundance, which could generate profound consequences for community and

ecosystem dynamics (Ehleringer and Monson 1993, Still et al. 2003). Furthermore, different

climate change variables can have counteracting effects on their competitive relationships.

For example, while higher temperatures and drought conditions increase the competitive

ability of C_4 plants against C_3 plants, elevated CO_2 may enhance C_3 plant performance and

thus ameliorate the negative effects of warming and drought on their competitiveness,

highlighting the need of simultaneously considering multiple environmental stressors for

predicting C_3-C_4 plant dynamics under changing climate.

Different environmental factors can interact to exert a more strengthened (synergistic)

or weakened (antagonistic) effect on plants than the simple summation of their individual

effects (Wand et al. 1999). For example, an experiment showed a synergistic effect between

water and CO_2 treatment in C_3-C_4 mixture, where elevated CO_2 had a greater positive effect

on the C_4 biomass under drought compared to well-water conditions (Valerio et al. 2011). On

the other hand, another study found that the growth stimulation of elevated CO_2 on a C_3 crop

was dampened at higher temperatures, indicating an antagonistic effect between warming and

CO_2 (Alberto et al. 1996). While extensive research has focused on the individual effect of

warming, drought, and elevated CO_2 as well as two-factor interactions on C_3 and C_4 plant

performances and competition, studies examining all three factors simultaneously remain

scarce. Because of the concurrent changes in global temperature, precipitation patterns, and

atmospheric CO_2 concentration, investigating how these three factors jointly affect C_3-C_4

plant dynamics could help better predict future plant communities in the face of climate

change.

To examine the joint effects of the aforementioned three climate change factors on the

biomass performances and competitive responses of C_3 and C_4 plants, we grew naturally co-

occurring C_3 (*Oplismenus composites*) and C_4 (*Paspalum conjugatum*) grass in growth

chambers under three temperature warming scenarios, two water supply intervals, and two

CO_2 concentrations. We compared the above- and below-ground dry biomass in monoculture

and mixture to assess their competitive responses under different experimental treatments.

Based on the the ecophysiology of C_3 and C_4 plants, we predict a greater competitiveness of

the C_3 grass under lower temperatures, ample water supply, and elevated CO_2, whereas the C_4

grass will be more competitive under warming, drought, and ambient CO_2.

Materials and Methods

Preparation of plant individuals

Ramets of *O. composites* and *P. conjugatum* were collected from the campus of National

Taiwan University (N25° 00' 57", E121° 32' 19") in October, 2017. These ramets were

planted in seedling nursery pots using fully-mixed peat moss soil (Kekkila natural peat moss

soil) as the substrate. All plant individuals were first cultivated in greenhouse at ambient CO_2

concentration with 25/20°C day/night temperature for two weeks, allowing their recovery and

establishment prior to the start of experiment.

Experimental design

To examine the effects of major climate change factors on the performances and competitive

interactions of the C_3 and C_4 grass, we used a full-factorial design crossing three temperature

warming scenarios, two water supply intervals, and two CO_2 concentrations (see the

following paragraph for more details). Each treatment combination contained four replicate

pots of C_3 monoculture, C_4 monoculture, and C_3-C_4 mixture (each with four individuals; see

the following text), yielding a total of 144 pots (12 treatment combinations × 3 planting

schemes × 4 replicate pots (Fig. 1). To avoid the potential confounding effect of initial size,

similar-sized individuals of each two grasses (C_3 grass: 8.6 ± 4.3 cm; C_4 grass: 6.2 ± 4.5 cm;

mean ± SD) were selected from the greenhouse nursery for the experimental pots. Four

individuals were grown and equally-spaced in each pot (13 cm in height x 12 cm in diameter),

conforming to their densities at the site where the ramets were collected (personal

observations). In the monoculture pots, all four individuals belonged to the same species; in

the mixture pots, two individuals of each species were planted. Each replicate pot was

assigned a pot number prior to the start of the experiment; monoculture and mixture pots

were paired for analysis of competitive responses (Fig. 1). Pots were placed in growth

chambers receiving respective treatments and a constant relative humidity of 70%,

photoperiod of 12:12 light/dark, and photosynthetic photon flux density (PPFD) of 160–200

$\mu mol \cdot M^{-2} \cdot S^{-1}$.

Three warming scenarios were used in the temperature treatment: control (+0°C), 2°C

warming (+2°C), and 4°C warming (+4°C), based on the projection that mean global surface

temperature will be 1.8 ± 0.5°C and 3.7 ± 0.7°C above the 1986–2005 reference period by

2100 under intermediate (Representative Concentration Pathway [RCP] 4.5) and high carbon

emissions scenario (RCP 8.5), respectively (IPCC 2014). To reflect the daily temperature

fluctuations in the field, a control temperature curve was derived using hourly temperature

records in 2016 from the nearest weather station (Taipei weather station; mean: 23.3°C; range:

21.6–25.9°C). Temperature curves for the +2°C and +4°C treatment were obtained by

horizontally shifting the control temperature curve upward by 2°C and 4°C, respectively (Fig.

S1). Two watering regimes were used in the water treatment: 2 days and 7 days, representing

the normal and prolonged precipitation interval (drought). The normal interval of 2 days was

determined based on the mean annual number of precipitation days between 1981 and 2011

(165.5 days per year) using weather records from the Taipei weather station. The water

volume for a single watering event was 160 ml (derived by multiplying the average

precipitation on each rainy day [14.5 mm; mean annual precipitation of 2405 mm divided by 165.5 days] with the pot surface area [110 cm^2]). Two CO_2 concentrations were used in the CO_2 treatment: 400 ppm (ambient) and 800 ppm (elevated), reflecting the current level of atmospheric CO_2 and the projected atmospheric CO_2 between RCP 6.0 (670 ppm) and RCP 8.5 (936 ppm) (IPCC 2014).

Plants were harvested after six weeks of cultivation and separated into above- and below-ground parts. The below-ground parts were carefully washed to remove the soil, with as many fine roots retained as possible. All plant materials were oven-dried at 50°C for seven days. Dry biomass was measured to the third decimal place in the unit of gram (g).

Data analyses

To examine the effects of our experimental treatments on the above- and below-ground dry biomass of the C_3 and C_4 grass, we fitted linear mixed effects models (LMMs) with temperature warming, water supply interval, CO_2 concentration, culture type (monoculture vs. mixture), and their interactions as fixed effects, and pot as a random effect, using the "lmer" function in the *lme4* R package (Bates et al. 2015). Models were fitted separately for the above- and below-ground biomass of the two species (i.e., a total of four models). The response (dry biomass of individual plant) was log-transformed; initial plant height was also included as a covariate. The significance of predictors was assessed via *F*-tests (type III sum of squares) using the "joint_tests" function in the *emmeans* R package (Lenth et al. 2018). Tukey post-hoc comparisons of treatment levels were conducted using also the *emmeans* R package.

To quantify the competitive responses of the C_3 and C_4 grass in each treatment

combination, we first averaged the above-/below-ground dry biomass of the four individuals

in each monoculture pot as well as the biomass of the two individuals of the same species in

each mixture pot. We then calculated the log response ratio (lnRR) of the pot-level mean

biomass in paired monoculture and mixture pots as log(mixture mean biomass / monoculture

mean biomass). This lnRR measures the relative effects of interspecific versus intraspecific

competition received by the focal individual, effectively representing the relative "sensitivity"

of a species to interspecific and intraspecific competition (Bennett et al. 2016). A positive

lnRR indicates a better performance under interspecific competition compared to intraspecific

competition (i.e., less suppression by interspecific competition) and vice versa for a negative

lnRR. The lnRRs were analyzed using linear models with temperature warming, water supply

interval, CO_2 concentration, plant species, and their interactions as predictors. The

significance of predictors was assessed via *F*-tests (type III sum of squares) as above; the

estimated marginal means (EMMs) of lnRRs of the two grass species derived from the linear

models were compared within each treatment combination using the *emmeans* R package. All

analyses were performed in R version 4.1.2 (R Core Team 2021).

Results

175   *Biomass performances of the C_3 and C_4 grass*

(1) C_3 above- and below-ground dry biomass

The above-ground dry biomass of the C_3 grass (*O. composites*) was affected by temperature

warming, water supply interval, and CO_2 concentration (Table 1). The above-ground biomass

was higher under 4°C warming for all but normal water supply interval × ambient CO_2

treatment (Fig. 2), indicating an interaction between temperature and water as well as

between temperature and CO_2 (Table 1). Prolonged water supply interval reduced the

biomass, whereas elevated CO_2 increased the above-ground biomass (Table S1).

The below-ground dry biomass of *O. composites* was affected by temperature

warming and water supply interval but not CO_2 concentration (Table S2). The below-ground

biomass did not differ among temperature treatments at ambient CO_2 but was higher under

4°C warming at elevated CO_2 (Fig. 2), indicating an interaction between temperature and

CO_2 (Table S2). Prolonged water supply interval reduced the below-ground biomass (Table

S1).

(2) C_4 above- and below-ground dry biomass

The above-ground dry biomass of the C_4 grass (*P. conjugatum*) was affected by temperature

warming and CO_2 concentration but not water supply interval (Table 1). The above-ground

biomass was higher under 4°C warming across all water × CO_2 treatments (Fig. 3). Elevated

CO_2 enhanced the above-ground biomass production (Table S1).

Similar to the response of above-ground dry biomass to the experimental treatments,

the below-ground dry biomass of *P. conjugatum* was affected by temperature warming and

CO_2 concentration but not water supply interval (Table S2). The effect of temperature

warming on below-ground biomass varied across water and CO_2 treatments (i.e., Temperature

× Water × CO_2 interaction; Fig. 3; Table S2). Specifically, at normal water supply interval,

the below-ground biomass was lower under 2°C warming compared to control temperature at

ambient CO_2, while there was no difference in biomass among temperature treatments at

elevated CO_2. At prolonged water supply interval, the below-ground biomass was the lowest

under 2°C warming, although the difference between 2°C and 4°C warming was not

statistically significant at ambient CO_2. Elevated CO_2 enhanced the below-ground biomass

production (Table S1).

When examining the effect of culture type, we found a significant Temperature ×

Culture type interaction for the C_4 grass (Table 1; Table S2). In particular, the C_4 above-

ground biomass was considerably higher in mixture than in monoculture (Fig. S2), suggesting

that warming may increase the competitive ability of C_4 grass when grown with C_3 grass.

211   *Competitive responses of the C_3 and C_4 grass*

The lnRRs of above- and below-ground dry biomass were not affected by temperature

warming, water supply interval, or CO_2 concentration; however, the competitive responses of

the two grasses depended on the CO_2 concentration (Species × CO_2 interaction; Table 2; Fig.

4). At ambient CO_2, the C_3 grass had similar above-ground biomass under interspecific

(mixture) and intraspecific (monoculture) competition (lnRR ≈ 0) and higher below-ground

biomass under interspecific competition compared to intraspecific competition (lnRR > 0),

whereas the C_4 grass had lower above- and below-ground biomass under interspecific

competition compared to intraspecific competition (lnRR < 0). In contrast, the competitive

responses reversed at elevated CO_2, with the C_3 grass having lower biomass (lnRR < 0) and

the C_4 grass having higher biomass when grown in mixtures (lnRR > 0) (Fig. 4). These

results suggest that increasing CO_2 concentration could benefit C_4 grass when in competition

with C_3 grass.

Overall, the above- and below-ground competitive responses of the two grass species

in each treatment combination were generally comparable, despite some variations in their

signs and magnitudes (Fig. S3). However, the lnRRs of both above- and below-ground dry

biomass did differ between the two species under a combination of 4°C warming, prolonged

water supply interval, and elevated CO_2, where the C_3 grass performed worse (lnRR < 0) and

the C_4 grass performed better (lnRR > 0) in mixtures compared to their corresponding

monocultures (Fig. 5a; Fig. S3). Indeed, the positive lnRRs of the C_4 grass under multiple

environmental stressors were largely due to an increase in mixture biomass rather than a

reduction in monoculture biomass in comparison to the controlled baseline treatment (Fig.

5b). Together, these results indicate a greater competitive ability of the C_4 grass under higher

temperature, drought condition, and elevated CO_2.

Discussion

We examined the joint effects of three major climate change factors—temperature, water, and

CO_2—on the biomass performances and competitive responses of a C_3 and a C_4 grass in

growth chambers. Warming and elevated CO_2 generally enhanced the above- and/or below-

ground biomass of both grasses, with a synergism between these two factors. In contrast to

theoretical predictions and experimental results (Carter and Peterson 1983, Pearcy and

Ehleringer 1984), the C_3 grass performed worse under interspecific competition (mixture

biomass) relative to intraspecific competition (monoculture biomass) at elevated CO_2, while

the reverse is true for C_4 grass. Furthermore, the C_4 grass outperformed the C_3 grass most

under simultaneous 4°C warming, drought conditions, and elevated CO_2, suggesting that the

competitive balance could potentially shift in favor of C_4 grass under the projected

temperature warming, increased drought frequencies, and rising atmospheric CO_2. Taken

together, our results provide empirical evidence for C_3 and C_4 biomass performances and

competition in response to multiple environment stressors, underlining the importance of

considering multi-variable effects and offering useful insights into C_3-C_4 plant dynamics

under future climate change.

Effects of warming and elevated CO_2 on biomass performances

We found that warming (in particular 4°C warming) increased the above- and below-ground

biomass of the C_3 grass, contradicting the common notion that C_3 grass performance should

not be enhanced, or even potentially be reduced, under higher temperatures (Sage and Kubien

2007). A possible explanation for this observation is that the control temperature used in the

study was below the thermal optimum of photosynthesis for the C_3 grass and therefore

warming indeed led to a further increase in carbon fixation and biomass production.

Additionally, the C_3 grass may shift its thermal optimum as the temperature rises, as research

has suggested that C_3 plants have broader thermal ranges and higher potential for thermal

acclimation (Sage and Kubien 2007, Yamori et al. 2014). Consequently, future climate

warming might not necessarily suppress C_3 plants as generally predicted if they are able to

exhibit plastic thermal acclimation to higher temperatures.

Elevated CO_2 enhanced the above-ground biomass of the C_3 grass as well as the

above- and below-ground biomass of the C_4 grass. The higher C_3 grass biomass at elevated

CO_2 agrees with the widely-documented positive response of C_3 plants to increasing CO_2

(Ainsworth and Long 2005, Wang et al. 2012). In contrast, biomass enhancement of the C_4

grass is surprising as theory predicts that increasing CO_2 concentration may not benefit the

performance of C_4 plants because of their lower CO_2 saturation points for photosynthesis

(Sage 1994, Ghannoum et al. 2000). Nonetheless, empirical evidence regarding the effect of

CO_2 on the performance of C_4 plants is mixed. In fact, some studies have reported biomass

stimulation by CO_2 in C_4 plants (e.g., Owensby et al. 1993, Wand et al. 1999). These findings

indicate that the response of C_4 plants to CO_2 could be species-specific and a future rise in

atmospheric CO_2 may potentially benefit certain C_4 plants.

277   *Interactions among treatment factors*

The interaction between temperature and CO_2 appeared to be the most conspicuous two-way

interaction among the three treatment factors across both grass species (Table 1; Table S2).

The synergistic effect of temperature warming and elevated CO_2 suggests that climate

warming together with increasing atmospheric CO_2 could further enhance the biomass

production of plants regardless of their photosynthetic pathways (Wang et al. 2012). For the

C_3 grass, such a synergism may arise from a greater ameliorating effect of CO_2 on

photorespiration with rising temperature (Long 1991, Dusenge et al. 2019). Furthermore,

elevated CO_2 could increase the thermal optimum of net carbon assimilation rate of the C_3

grass (Sage and Kubien 2007), leading to greater biomass production under simultaneous

warming and elevated CO_2. On the other hand, C_4 plants generally do not respond strongly to

rising CO_2 because of their CO_2 concentration mechanism. The potential mechanisms

underlying the observed positive temperature $\times$ CO_2 interaction in the C_4 grass warrant

further investigations.

Our results revealed a three-way interaction among temperature, water, and CO_2 for

the below-ground biomass of the C_4 grass. There are a few possible explanations for this.

First, warming-induced physiological changes (e.g., reduced stomatal conductance) could

provide cross tolerance to drought (Havaux 1992). Second, elevated CO_2 may increase water

use efficiency and thus help alleviate the impact of water stress (Xu et al. 2013). Third, more

resources may be allocated to below-ground growth to increase water uptake in response to

temperature warming and water limitation (Malik et al. 1979, Sharp and Davies 1989).

Research on how C_3 and C_4 plants respond to multiple stressors has been limited, and our

finding provides experimental evidence that future climate change could affect plant biomass

production in a complex way that might not be predicted from individual factors alone. More

studies on how multiple climate change factors jointly affect C_3 and C_4 plant performances

are needed to verify if such a three-way interaction is a common phenomenon.

304   *Competitive responses under experimental treatments*

Countering previous studies showing greater competitive advantages of C_3 plants at elevated

CO_2 (Marks and Strain 1989, Ziska 2001), our results show that higher CO_2 concentration

could potentially benefit the C_4 grass when in competition with the C_3 grass. Specifically, the

C_3 grass was suppressed more by interspecific competition relative to intraspecific

competition at elevated CO_2 compared to ambient CO_2, whereas the C_4 grass performed

better under interspecific competition relative to intraspecific competition at elevated CO_2

compared to ambient CO_2 (Fig. 4). Such patterns can be partly explained by a greater effect

of elevated CO_2 on the C_4 above-ground biomass in mixture compared to monoculture (CO_2

× Culture type interaction; Table 1). Moreover, elevated CO_2 combined with 4°C warming

and drought resulted in the largest difference in the competitive responses of the two grasses,

with an increased interspecific competitive ability of the C_4 grass against the C_3 grass (Fig. 5).

These results suggest that the competitive balance could potentially shift in favor of C_4 plants

under future climate scenarios, which may have potential impacts on both natural and

agricultural systems. For example, C_4 plants are generally less nutritious than C_3 plants, and

increasing C_4 relative abundance in the habitat could reduce the nutritional availability for

herbivores (Barbehenn et al. 2004). Furthermore, C_3 crops may suffer greater yield loss

because of increasing competition pressure from C_4 weeds (Korres et al. 2016).

Despite some variations in the signs and magnitudes, the competitive responses of the

two grass species were by and large similar in most treatment combinations, indicating

similar relative effects of interspecific versus intraspecific competition received by the two

grasses. Surprisingly, we did not find any main effect of temperature, water, or CO_2 treatment

on the above- and below-ground competitive responses of the two grasses, despite strong

effects of these factors on their biomass production. A lack of treatment effect on the

competitive responses may occur if the two grasses respond to the treatments in a similar

fashion (e.g., positive responses of both C_3 and C_4 grass to elevated CO_2), and consequently

there might be no competitive shift because neither species would gain a net advantage over

the other. This shows that performances of individual species alone may not necessarily

reflect their competitive relationships, which depend largely on the species' relative

performances when grown together.

335   *Conclusions*

Using an experimental approach, our study shows that warming and elevated CO_2 together

enhanced the biomass performances of both C_3 and C_4 grass, and temperature, water, and

CO_2 interacted to affect the C_4 biomass. Moreover, simultaneous warming, drought, and

particularly elevated CO_2, benefited the C_4 grass most when in competition with the C_3 grass,

suggesting potential shifts in their competitive balance along with an increase in C_4 relative

abundance under future climate scenarios. Changes in C_3-C_4 community composition can

have profound consequences for the functioning of natural and agricultural systems,

including primary productivity, nutrient cycling, herbivore nutritional balance, and crop

production. As the effects of climate change are multifaceted and often involve complex

interactions and feedbacks over time and space, long-term and large-scale field experiments

involving more species will represent a promising research avenue that scales up our

laboratory results to better predict C_3-C_4 plant dynamics under changing climate.

Acknowledgements

We thanked William J-A Ou and Jia-Zhen Lin for assisting with experimental work and data

collection. This study was supported by funding from Ministry of Science and Technology,

Taiwan (106-2813-C-002-163-B).

References

Ainsworth, E. A., and S. P. Long. 2005. What have we learned from 15 years of free-air CO2

enrichment (FACE)? A meta-analytic review of the responses of photosynthesis,

canopy properties and plant production to rising CO2. New Phytologist **165**:351-372.

Alberto, A. M., L. H. Ziska, C. R. Cervancia, and P. A. Manalo. 1996. The influence of

increasing carbon dioxide and temperature on competitive interactions between a C3

crop, rice (Oryza sativa) and a C4 weed (Echinochloa glabrescens). Functional Plant

Biology **23**:795-802.

Barbehenn, R. V., Z. Chen, D. N. Karowe, and A. Spickard. 2004. C3 grasses have higher

nutritional quality than C4 grasses under ambient and elevated atmospheric CO2.

Global Change Biology **10**:1565-1575.

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models

using lme4. Journal of Statistical Software 67: 1–48.

Bennett, J. A., K. Riibak, R. Tamme, R. J. Lewis, and M. Pärtel. 2016. The reciprocal

relationship between competition and intraspecific trait variation. Journal of Ecology

104:1410-1420.

Carter, D., and K. Peterson. 1983. Effects of a CO_2-enriched atmosphere on the growth and

competitive interaction of a C_3 and a C_4 grass. Oecologia **58**:188-193.

Dusenge, M. E., A. G. Duarte, and D. A. Way. 2019. Plant carbon metabolism and climate

change: elevated CO_2 and temperature impacts on photosynthesis, photorespiration

and respiration. New Phytologist **221**:32-49.

Edwards, E. J., and C. J. Still. 2008. Climate, phylogeny and the ecological distribution of C4

grasses. Ecology letters **11**:266-276.

Ehleringer, J. R., and R. K. Monson. 1993. Evolutionary and ecological aspects of

photosynthetic pathway variation. Annual Review of Ecology and Systematics

24:411-439.

Ghannoum, O., S. v. Caemmerer, L. H. Ziska, and J. P. Conroy. 2000. The growth response

of C4 plants to rising atmospheric CO2 partial pressure: a reassessment. Plant, Cell &

Environment **23**:931-942.

Havaux, M. 1992. Stress tolerance of photosystem II in vivo: antagonistic effects of water,

heat, and photoinhibition stresses. Plant Physiology **100**:424-432.

IPCC. 2014. Climate change 2014: Synthesis report. In Contribution of Working Groups I, II

and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate

Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)], Geneva,

Switzerland.

Korres, N. E., J. K. Norsworthy, P. Tehranchian, T. K. Gitsopoulos, D. A. Loka, D. M.

Oosterhuis, D. R. Gealy, S. R. Moss, N. R. Burgos, and M. R. Miller. 2016. Cultivars

to face climate change effects on crops and weeds: a review. Agronomy for Sustainable Development **36**:1-22.

Lee, J.-S. 2011. Combined effect of elevated CO2 and temperature on the growth and phenology of two annual C3 and C4 weedy species. Agriculture, ecosystems & environment **140**:484-491.

Lenth, R., H. Singmann, J. Love, P. Buerkner, and M. Herve. 2018. Emmeans: Estimated marginal means, aka least-squares means. R package version **1**:3.

Long, S. 1991. Modification of the response of photosynthetic productivity to rising temperature by atmospheric CO2 concentrations: has its importance been underestimated? Plant, Cell & Environment **14**:729-739.

Malik, R., J. Dhankar, and N. Turner. 1979. Influence of soil water deficits on root growth of cotton seedlings. Plant and soil **53**:109-115.

Marks, S., and B. R. Strain. 1989. Effects of drought and CO2 enrichment on competition between two old-field perennials. New Phytologist **111**:181-186.

Owensby, C. E., P. I. Coyne, J. M. Ham, L. M. Auen, and A. K. Knapp. 1993. Biomass Production in a Tallgrass Prairie Ecosystem Exposed to Ambient and Elevated CO" 2. Ecological applications **3**:644-653.

Öztürk, M., H. Rehder, and H. Ziegler. 1981. Biomass production of C3-and C4-plant species in pure and mixed culture with different water supply. Oecologia **50**:73-81.

Pau, S., E. J. Edwards, and C. J. Still. 2013. Improving our understanding of environmental controls on the distribution of C3 and C4 grasses. Global Change Biology **19**:184-196.

Pearcy, R., and J. Ehleringer. 1984. Comparative ecophysiology of C3 and C4 plants. Plant, Cell & Environment **7**:1-13.

414   Pearcy, R. W., N. Tumosa, and K. Williams. 1981. Relationships between growth,

415   photosynthesis and competitive interactions for a C 3 and C 4 plant. Oecologia

416   **48**:371-376.

417   Poorter, H., and M. L. Navas. 2003. Plant growth and competition at elevated CO2: on

418   winners, losers and functional groups. New Phytologist **157**:175-198.

419   R Core Team. 2021. R: A language and environment for statistical computing. R Foundation

420   for Statistical Computing, Vienna, Austria.

421   Sage, R. F. 1994. Acclimation of photosynthesis to increasing atmospheric CO2: the gas

422   exchange perspective. Photosynthesis research **39**:351-368.

423   Sage, R. F., and D. S. Kubien. 2007. The temperature response of C3 and C4 photosynthesis.

424   Plant, Cell & Environment **30**:1086-1106.

425   Sage, R. F., and R. K. Monson. 1998. C4 plant biology. Elsevier.

426   Sharp, R., and W. Davies. 1989. Regulation of growth and development of plants growing

427   with a restricted supply of water. Page 71   Plants under stress: biochemistry,

428   physiology and ecology and their application to plant improvement.

429   Still, C. J., J. A. Berry, G. J. Collatz, and R. S. DeFries. 2003. Global distribution of C3 and

430   C4 vegetation: carbon cycle implications. Global biogeochemical cycles **17**:6-1-6-14.

431   Teeri, J., and L. Stowe. 1976. Climatic patterns and the distribution of C4 grasses in North

432   America. Oecologia **23**:1-12.

433   Valerio, M., M. Tomecek, S. Lovelli, and L. Ziska. 2011. Quantifying the effect of drought

434   on carbon dioxide-induced changes in competition between a C3 crop (tomato) and a

435   C4 weed (Amaranthus retroflexus). Weed Research **51**:591-600.

436   Wand, S. J., G. Midgley, M. H. Jones, and P. S. Curtis. 1999. Responses of wild C4 and C3

437   grass (Poaceae) species to elevated atmospheric CO2 concentration: a meta-analytic

438   test of current theories and perceptions. Global Change Biology **5**:723-741.

Wang, D., S. A. Heckathorn, X. Wang, and S. M. Philpott. 2012. A meta-analysis of plant
physiological and growth responses to temperature and elevated CO2. Oecologia
169:1-13.

Ward, J., D. T. Tissue, R. B. Thomas, and B. Strain. 1999. Comparative responses of model
C3 and C4 plants to drought in low and elevated CO2. Global Change Biology **5**:857-
867.

Warne, R. W., A. D. Pershall, and B. O. Wolf. 2010. Linking precipitation and C3–C4 plant
production to resource dynamics in higher-trophic-level consumers. Ecology **91**:1628-
1638.

Xu, Z., H. Shimizu, Y. Yagasaki, S. Ito, Y. Zheng, and G. Zhou. 2013. Interactive effects of
elevated CO2, drought, and warming on plants. Journal of Plant Growth Regulation
32:692-707.

Yamori, W., K. Hikosaka, and D. A. Way. 2014. Temperature response of photosynthesis in
C3, C4, and CAM plants: temperature acclimation and temperature adaptation.
Photosynthesis research **119**:101-117.

Ziska, L. H. 2001. Changes in competitive ability between a C4 crop and a C3 weed with
elevated carbon dioxide. Weed Science **49**:622-627.

 Tables and Figures

459 Table 1. Results of the linear mixed models (LMMs) examining the effects of temperature

460 warming, water supply interval, CO_2 concentration, and culture type (monoculture vs.

461 mixture) on the above-ground dry biomass of the C_3 and C_4 grass

Predictor	C_3 grass			C_4 grass		
	F	*df*	*P**	*F*	*df*	*P**
Temperature warming	57.42	2, 103	**< 0.001**	56.66	2, 99	**< 0.001**
Water supply interval	26.13	1, 99	**< 0.001**	1.60	1, 99	0.208
CO_2 concentration	48.7	1, 99	**< 0.001**	18.8	1, 100	**< 0.001**
Culture type	3.54	1, 99	0.063	2.30	1, 99	0.132
Temperature × Water	4.41	2, 99	**0.015**	2.82	2, 99	0.064
Temperature × CO_2	18.07	2, 101	**< 0.001**	3.10	2, 99	**0.049**
Temperature × Culture type	1.07	2, 99	0.346	4.89	2, 100	**0.009**
Water × CO_2	5.74	1, 99	**0.018**	0.73	1, 99	0.394
Water × Culture type	0.39	1, 99	0.534	0.25	1, 99	0.619
CO_2 × Culture type	0.26	1, 102	0.613	4.57	1, 99	**0.035**
Temperature × Water × CO_2	0.13	2, 99	0.881	1.84	2, 99	0.164
Temperature × Water × Culture type	1.97	2, 99	0.145	2.43	2, 100	0.093
Temperature × CO_2 × Culture type	0.90	2, 100	0.409	0.15	2, 99	0.861
Water × CO_2 × Culture type	3.08	1, 99	0.082	0.96	1, 99	0.331
Temperature × Water × CO_2 × Culture type	4.07	2, 99	**0.020**	0.26	2, 99	0.769

462 * Significant effects ($P < 0.05$) are highlighted in bold.

464   Table 2. Results of the linear models examining the effects of temperature warming, water

465   supply interval, CO_2 concentration, and species on the log response ratios (lnRRs) of above-

466   and below-ground dry biomass in mixture relative to monoculture (log[mixture

467   biomass/monoculture biomass])

Predictor	Above-ground			Below-ground		
	F	df	P*	F	df	P*
Temperature warming	0.91	2, 72	0.406	0.59	2, 71	0.556
Water supply interval	2.94	1, 72	0.091	0.40	1, 71	0.528
CO_2 concentration	0.22	1, 72	0.637	0.25	1, 71	0.621
Species	1.59	1, 72	0.211	0.12	1, 71	0.730
Temperature × Water	1.77	2, 72	0.177	1.29	2, 71	0.281
Temperature × CO_2	1.36	2, 72	0.262	1.56	2, 71	0.218
Temperature × Species	1.98	2, 72	0.146	1.61	2, 71	0.207
Water × CO_2	0.74	1, 72	0.392	2.67	1, 71	0.106
Water × Species	0.07	1, 72	0.792	4.97	1, 71	**0.029**
CO_2 × Species	5.66	1, 72	**0.020**	7.87	1, 71	**0.006**
Temperature × Water × CO_2	2.87	2, 72	0.063	2.98	2, 71	0.057
Temperature × Water × Species	0.44	2, 72	0.645	0.46	2, 71	0.631
Temperature × CO_2 × Species	0.31	2, 72	0.736	0.58	2, 71	0.561
Water × CO_2 × Species	1.72	1, 72	0.194	1.43	1, 71	0.235
Temperature × Water × CO_2 × Species	1.60	2, 72	0.210	3.38	2, 71	**0.039**

468 * Significant effects ($P < 0.05$) are highlighted in bold.

Figure 1. An illustration of the experimental design of this study. The experiment was conducted using a factorial design crossing three temperature warming scenarios (control, +2°C, and +4°C), two water supply intervals (normal 2 days vs. prolonged 7 days), and two CO_2 concentrations (ambient 400 ppm vs. elevated 800 ppm). Each treatment combination contained four replicate pots of C_3 (*Oplismenus composites*) and C_4 grass (*Paspalum conjugatum*) in monoculture as well as C_3-C_4 mixture. Four plant individuals (four C_3 or C_4 in monoculture and two C_3 + C_4 in mixture) were grown in each pot. Replicate monoculture and mixture pots in each treatment combination were paired *a priori* by assigning the same pot number for analysis of competitive responses.

483   Figure 2. Above- and below-ground dry biomass of the C_3 grass under different temperature

484   warming, water supply interval, and CO_2 concentration treatments. Tukey post-hoc

485   comparisons were conducted for the three temperature scenarios within each water $\times$ CO_2

486   treatment combination using the estimated marginal means (EMMs) from the LMMs.

487   Different letters indicate statistical significance ($P < 0.05$).

493   Figure 3. Above- and below-ground dry biomass of the C_4 grass under different temperature

494   warming, water supply interval, and CO_2 concentration treatments. Tukey post-hoc

495   comparisons were conducted for the three temperature scenarios within each water $\times$ CO_2

496   treatment combination using the estimated marginal means (EMMs) from the LMMs.

497   Different letters indicate statistical significance ($P < 0.05$).

Figure 4. Interactions between CO_2 concentration and species for the lnRRs of above- and below-ground dry biomass. Points and error bars represent the EMMs and standard errors derived from the linear models. Grey dashed lines (lnRR = 0) indicate equal effects of interspecific and intraspecific competition; a positive lnRR indicates a better performance under interspecific competition compared to intraspecific competition (i.e., weaker suppression by interspecific competition) and vice versa for a negative lnRR.

Figure 5. LnRRs of above- and below-ground dry biomass of the two grass species (a) and above- and below-ground dry biomass of the C_4 grass (b) under two opposite treatment combinations: (1) control temperature × normal water supply interval × ambient CO_2, and (2) 4°C warming × prolonged water supply interval × elevated CO_2. Points and error bars in (a) represent the EMMs and standard errors derived from the linear models; points and error bars in (b) represent original means and standard errors. Asterisks denote a significant difference in lnRRs between the two grass species ($P < 0.05$).