Article

Activity-dependent glassy cell mechanics Ⅰ:

Mechanical properties measured with active microrheology

H. Ebata, 1, K. Umeda, 1 K. Nishizawa, 2 W. Nagao, 1 S. Inokuchi, 1 Y. Sugino, 1 T. Miyamoto, 3,4 and *D. Mizuno 1

1. Department of Physics, Kyushu University, 819-0395 Fukuoka, Japan.
2. Institute of Developmental Biology of Marseille, Campus de Luminy case 907, 13288 Marseille Cedex 09, France.
3. Department of Endocrinology and Metabolism, Faculty of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan
4. Transborder Medical Research Center, University of Tsukuba, Ibaraki, Japan.

* Correspondence: mizuno@phys.kyushu-u.ac.jp; Tel.: +81-92-802-4092

running title：**Critical jamming of living cytoplasm**

Abstract: Active microrheology was conducted in living cells by applying an optical-trapping force to vigorously-fluctuating tracer beads with feedback-tracking technology. The complex shear viscoelastic modulus $G(\omega) = G'(\omega) - iG''(\omega)$ was measured in HeLa cells in an epithelial-like confluent monolayer. We found that $G(\omega) \propto (-i\omega)^{1/2}$ over a wide range of frequencies (1 Hz $< \omega/2\pi <$10 kHz). Actin disruption and cell-cycle progression from G1 to S and G2 phases only had a limited effect on $G(\omega)$ in living cells. On the other hand, $G(\omega)$ was found to be dependent on cell metabolism; ATP-depleted cells showed an increased elastic modulus $G'(\omega)$ at low frequencies, giving rise to a constant plateau such that $G(\omega) = G_0 + A(-i\omega)^{1/2}$. Both the plateau and the additional frequency dependency $\propto (-i\omega)^{1/2}$ of ATP-depleted cells are consistent with a rheological response typical of colloidal jamming. On the other hand, the plateau G_0 disappeared in ordinary metabolically active cells, implying that living cells fluidize their internal states such that they approach the critical jamming point.

Statement of Significance: Intracellular mechanical properties were measured using optical-trap-based microrheology. Despite expectations to the contrary, shear viscoelasticity was hardly affected by reorganization of cytoskeletal structures during cell-cycle progression (G1 to S and G2 phases), nor by artificial disruption of the actin cytoskeleton induced by chemical inhibitors. Rather, the mechanics of cell interiors is governed by the glassy cytoplasm. Cells depleted of ATP solidified, whereas living cells that maintained metabolic activities were more fluid-like. Instead of a completely fluid response, however, we observed a characteristic power-law viscoelasticity $G(\omega) \propto (-i\omega)^{1/2}$ over the whole range of frequencies measured. Based on our current understanding of jamming rheology, we discuss how cells fluidize their internal state in a way that pushes the system towards the critical jamming transition.

Keywords: cell mechanics; microrheology; active fluidization; glassy cytoplasm; cell cycle; cytoskeleton

INTRODUCTION

The mechanics of cells is a fundamental aspect of biology; it governs the dynamics of functional bio-macromolecules that perform various physiological processes (1-3). For instance, motor proteins and other mechanoenzymes change their shapes to catalyze biochemical reactions (4, 5). They also need to be present at the right place at the right time (1); thus their performance depends on the mechanical properties (*i.e.*, fluidity) of the surrounding medium. At the same time, mechanical stability is also necessary for cells in order for them to maintain their complex internal organization. Living cells thus meet these apparently conflicting requirements: dynamic fluidity and mechanical stability. However, the physical machinery underlying the regulation of intracellular mechanics remains elusive (6, 7).

It has been believed that the mechanics of cells is determined by the properties of the cytoskeleton, a network of interlinking protein filaments (8, 9). After decades of intense research, a consensus has been reached that the mechanical properties of the cytoskeleton is described using a theory based on the semiflexible polymer network model (10). However, in living cells, the interstices of the sparse network of cytoskeletons are densely filled with other colloidal constituents such as proteins, nucleic acids, polysaccharides, and their assemblies such as ribosomes (2, 11-13). In contrast to the cytoskeleton, it remains to be elucidated whether and/or how the cytoplasm excluding the cytoskeleton (hereafter referred to as just the cytoplasm) contributes to intracellular mechanics. As it was recently pointed out, cytoplasmic rheology may be similar to that of dense colloidal suspensions close to jamming (3, 7).

As is typical in soft materials, both the cytoskeleton and cytoplasm respond in a highly nonlinear manner to mechanical perturbations. For instance, when subjected to mechanical loads such as shear stress or a locally applied force, semiflexible polymer networks typically stiffen (14-20) while dense colloidal suspensions are fluidized (21-24). Active mechanoenzymes such as molecular motors also apply forces to cytoskeletons (19) which will eventually be transmitted to the surrounding cytoplasm. Conformational changes of enzymes that occur during enzymatic reactions (4, 5) may also mechanically perturb a crowded environment. It is then reasonable to expect that driving the cell interior out of equilibrium will lead to an emergence of a complex intracellular mechanics.

Living cells dynamically change their internal organization during their cell-cycle progression. They replicate chromosomes, double cytoplasmic components and reconstruct cytoskeleton structures (25). Accordingly, it has been reported that the elastic and viscous moduli of cells depend on the cell-cycle phase (26-28). All the factors listed above, *i.e.*, the cytoskeleton, cytoplasm (in the interstices of cytoskeletons), their reorganization during cell-cycle progression, and their metabolic activity, seem to be integrated into intracellular mechanics. Prior studies of intracellular mechanics mostly focused on the cytoskeleton, but attention has rarely been paid to the cytoplasmic contribution. In this study, we aim to investigate how the mechanics of the cytoplasm is regulated by metabolic activities, using interphase cells which make up majority of our body and also cell cultures.

85   Microrheology (MR) is a method to investigate the viscoelastic properties of a medium
86 at microscopic ($\sim$ µm) length scales (29), by tracking the motion of micron-sized tracer beads
87 imbedded in a specimen (30, 31). In active microrheology (AMR) (32, 33), an external force
88 is applied to a tracer bead, and its displacement response is measured. When a sinusoidal
89 force is applied to the bead, we obtain the complex shear viscoelastic modulus
90 $G(\omega) = G'(\omega) - iG''(\omega)$ of the surrounding medium as a function of the angular frequency ω.
91 Here, G' and G'' are the real and imaginary part of the complex modulus $G(\omega)$. AMR can
92 be performed simultaneously by applying an optical-trapping force to the probe particle, and
93 measuring the probe displacements using the back-focal-plane laser interferometry technique
94 (BFPI) (34). Because of the vigorous non-thermal fluctuations, a probe particle in living cells
95 moves out of the laser focus within the experimental time scale. Although a strong laser
96 could trap the probe in the laser focus, it also hinders bead motion, causing large errors in
97 estimating the intrinsic response and fluctuations that should have occurred in the absence
98 of optical trapping. In a prior study, we therefore developed an optical-trap-based MR
99 implemented with a three-dimensional feedback-controlled sample stage (feedback-tracking
100 MR, see supplementary S1) (6, 35). This technique allowed us to stably track a vigorously-
101 fluctuating probe particle in living cells with a laser power as small as 0.4 ~ 1.8 mW,
102 significantly lower than ordinary optical-trapping measurements (33, 36). As seen in Fig. S2,
103 this non-invasive technology is crucial for conducting optical-trap-based MR in living cells.

104   Using this MR technique, we measured the intracellular mechanics of living HeLa cells.
105 To focus on the cytoplasmic mechanics, the contribution of the cytoskeleton was reduced by
106 (1) using probe particles that do not bind to cytoskeletal filaments, and (2) measuring the
107 location where actin cytoskeletons are supposed to be sparse, *i.e.,* the middle of cells cultured
108 in an epithelial-like monolayer sheet. We found that the cell-cycle progression during
109 interphase and cytoskeletal organization merely have a marginal effect on intracellular
110 mechanics measured in this way. The complex shear viscoelastic modulus $G(\omega)$ did not
111 show large variation from cell to cell; they consistently showed a frequency dependence
112 predicted for disordered colloidal jamming $G(\omega) = A(-i\omega)^{1/2}$ (6, 7).

113   On the other hand, the intracellular $G(\omega)$ was highly dependent on metabolic activities.
114 The viscoelasticity was seen to increase in ATP-depleted HeLa cells. A plateau elasticity G_0
115 appeared which is real and constant over the frequency range, *i.e.*, $G(\omega) = G_0 + A(-i\omega)^{1/2}$.
116 We discuss how these observations are consistent with what is expected for a glassy
117 cytoplasm (3, 7, 37). Although it is inferred that the metabolic activity fluidized the
118 cytoplasm (3, 7), a typical fluid-like response $G(\omega) \propto -i\omega\eta$ was not observed in the range of
119 frequencies measured (0.1 Hz ~ 10^4 Hz). Instead, our results imply that the rheological
120 properties typical for critical jamming (38, 39) is achieved during the active fluidization of
121 the glassy cytoplasm in living cells (40).

MATERIALS AND METHODS

Cell preparation

HeLa cells and HeLa/Fucci2 cells (Riken, Cell Bank) were seeded on fibronectin-coated glass-bottom petri dishes in Dulbecco's modified Eagle's medium (Wako, D-Mem, high glucose) with glucose (1 mg/ml), penicillin (100 U/ml), streptomycin (0.1 mg/ml), amphotericin B (250 mg/ml), and 10% fetal bovine serum (FBS) at 37°C. Cells were cultured in a CO_2 incubator until they formed a confluent epithelial-like monolayer sheet. The surface of the probe particles (melamine particles, 1 μm diameter, micro Particles GmbH) were coated with polyethylene glycol (PEG) strands (NANOCS, mPEG-NH2, 1000Da, PG1-AM-1k) (35), PEG coating generally passivates probe surfaces; in aqueous environments, hydrophilic PEG acts as a polymer brush and prevents sticking to other objects or to other molecules. Probe particles were introduced into cells using a gene gun (Bio-Rad, PDS-1000/He). Excess beads that did not enter the cells were removed by washing the dishes with phosphate-buffered saline. After replenishing the dishes with fresh culture medium, they were placed in an CO_2 incubator at least overnight to allow for the recovery from cell damage. Since supplying CO_2 causes additional noise, MR experiments were performed in a CO_2 -independent culture medium (Gibco, L-15 medium) with 10% FBS serum. All measurements were performed at 37°C. For MR experiments under ATP depletion and actin disruption, FBS was not included in the media as stated below.

Actin disruption

To disrupt the actin network, polymerization of actin filaments was inhibited by adding chemical agents (Cytochalasin D 50 μg/ml or Latrunculin B 10 μM). After MR experiments were performed at 37°C in L-15 medium with 10% FBS (w/o chemical agents), the culture medium was replaced with the actin disruption medium (Cytochalasin D or Latrunculin B, without FBS). Cells were then incubated at 37 °C on the MR setup for ~30 min prior to the next MR measurement so that the actin network is disrupted in cells.

ATP depletion

The high energy molecule ATP is mainly produced in cells via two metabolic pathways: glycolytic in the cytoplasm and oxidative phosphorylation in the mitochondria. Therefore, in order to deplete intracellular ATP, cells were cultured in a nutrient-free media with 50 mM 2-deoxy-D-glucose and 10 mM sodium azide (41). 2-deoxy-D-glucose and sodium azide inhibit glycolysis and oxidative ATP production, respectively.

HeLa cells were incubated in the medium for ATP depletion (L-15 medium with 50 mM 2-deoxy-D-glucose, 10 mM sodium azide, without FBS) at 37 °C for ~10 hours prior to the MR measurements so that ATP stored in cells is consumed.

FIGURE 1 (A) Schematic of the feedback-tracking microrheology setup. A fixed probe laser is used to measure u_{QPD} using a quadrant photodiode (QPD). A drive laser and an acousto-optic deflector (AOD) are used to apply an oscillatory external force. The PEG-coated probe particle surrounded by dense colloidal constituents is repelled from the sparse cytoskeletal network. (B)Typical time series of the probe displacement $u(t)$ (black line) and applied force $F(t) = \hat{F}(\omega)e^{-i\omega t}$ (red line). The sinusoidal signal $\hat{u}(\omega)e^{-i\omega t}$ was hidden in the experimental noise (*e.g.* thermal fluctuation) but can be detected by using a lock-in amplifier. (C) Schematic illustration of stage feedback. (D) The displacements of a probe $u(t)$ is obtained by the sum of the displacement of the piezo stage $u_{stage}(t)$ and the displacement from the focus of the probe laser. The focus of the drive laser is moved sinusoidally around the focus of the probe laser, following $L\exp(-i\omega t)$.

Luminescent ATP detection assay

Intracellular ATP was measured using a CellTiter-Glo® Luminescent Cell Viability

Assay kit (Promega, G7570) according to the manufacturer's instructions. Briefly, 5,000 cells

were plated on a 96-well plate and incubated for 24 h at 37°C in 5% CO_2. Following

incubation, the medium was replaced with 50 µl of either: (1) control medium [phenol-red

free DMEM (Thermo Fisher, 31053028) supplemented with 4 mM glutamine (Thermo Fisher,

25030081) and 1% Penicillin-Streptomycin Solution (Sigma-Aldrich, P4333)], or (2) control

medium supplemented with 50 mM 2-deoxyglucose (Nacalai, 10722-11) and 10 mM NaN_3

(WAKO, 199-11095), and then incubated for 7 h at 37°C in 5% CO_2. Subsequently, 50 µl of

reagent was directly added to the well and the contents were mixed for 2 min on an orbital

shaker at 25°C. Luminescence was recorded after 30 min of incubation at 25°C in the dark.

Immunocytochemistry

HeLa cells plated on a poly-lysine-coated glass-bottom dish were fixed with 4%

paraformaldehyde phosphate buffer solution (Wako, 163-20145) for 10 min at 25 °C,

permeabilized with PBS containing 0.5% Triton-100 for 15 min at 25 °C, and stained with

ActinRed™ 555 ReadyProbes™ Reagent (ThermoFisher, R37112) for 30 min at 25 °C.

Subsequently, the cells were incubated with Hoechst 33342 (ThermoFisher, H1399) for 10

min at 25 °C for nuclear staining. The cells were washed three times with PBS after each

step. Imaging was performed using an Eclipse Ti2-E microscope equipped with NIS-

Elements AR imaging software. The representative images were processed by Clarify.ai and

Denoise.ai using NIS-Elements AR 5.30, as previously described (42).

AMR in living cells

The technical details of conventional optical-trapping-based AMR (without feedback

operation) are given elsewhere (33). We will repeat the essence here briefly for the reader's

convenience. As shown in Figs. 1A and B, a small sinusoidal force $F(t) = \hat{F}(\omega)e^{-i\omega t}$ was

applied to a probe particle, by optically trapping it with the drive laser (λ =1064 nm

Nd:YVO$_4$, Coherent Inc., Tokyo, Japan) whose focus position was controlled by an acousto-

optic deflector (AOD; DTSX-400-1064, AA Opto-Electronic, Orsay, France). The probe

displacement synchronous to the applied force $u(t) = \hat{u}(\omega)e^{-i\omega t}$ was measured using back-

focal-plane laser interferometry (BFPI) conducted with a fixed probe laser (λ = 830 nm

IQ1C140, Power Technology, Inc., Alexander, AR, USA) (34). Typical values of $\hat{F}(\omega)$ and

$\hat{u}(\omega)$ are given in Supplementary S6. The frequency response function

$\tilde{\alpha}(\omega) = \alpha'(\omega) + i\alpha''(\omega) \equiv \langle \hat{u}(\omega) \rangle / \hat{F}(\omega)$ is then obtained as a complex quantity. The complex

shear modulus $G(\omega)$ of the surrounding media is also obtained as a complex quantity via the

Stokes relation extended to the sinusoidal response,

$$G(\omega) = 1/6\pi a \tilde{\alpha}(\omega). \tag{1}$$

where a is the radius of the spherical probe particle.

A tracer bead fluctuates vigorously in a living cell because the cytoplasm is driven far
from equilibrium by the energy derived from metabolism. As explained in detail in Refs. (6,
35), such a fluctuating particle was smoothly tracked by analog PID-feedback control of the
piezo-mechanical sample stage on which the specimen was placed. The optical-trapping
force was then stably applied to the probe particle by the oscillating drive laser while
maintaining the probe particle in the focus of the fixed probe laser. As shown in Fig. 1C, the
displacement of the probe particle from the probe laser u_{QPD}, which is measured by BFPI of
the probe laser, is used to control the position of the piezo stage u_{stage} by finding

$$u_{\mathrm{stage}}(t) = \frac{1}{\tau}\int u_{\mathrm{QPD}}dt \,. \tag{2}$$

207   Here, τ (typically set at 10 ms $\sim$ 100 ms) is the delay time for the feedback response of our
208   experimental setup. The displacement u_{FB} of the probe in the coordinate system traveling
209   with the feedback-controlled piezo stage is given as $u_{\mathrm{FB}} = u_{\mathrm{QPD}} + u_{\mathrm{stage}}$ (Fig. 1D). From Eq.
210   (2), the frequency-response relation is obtained as

$$\hat{u}_{\mathrm{FB}}(\omega) = (1 - i\omega\tau)\hat{u}_{\mathrm{stage}}(\omega) = \left(1 - \frac{1}{i\omega\tau}\right)\hat{u}_{\mathrm{QPD}}(\omega) \,, \tag{3}$$

where $^\wedge$ denotes the amplitude of the sinusoidal signal, $e.g.$ $u_{\mathrm{FB}}(t) = \hat{u}_{\mathrm{FB}}(\omega)\exp(-i\omega t)$. From
Eq. (3), it can be seen that $u_{\mathrm{QPD}}(t)$ and $u_{\mathrm{stage}}(t)$ are high- and low-pass filtered from the total
probe displacement $u_{\mathrm{FB}}(t)$. Thus, the slow/large displacements of the probe are tracked by
the piezo stage as u_{stage} whereas the fast/small displacements are measured with the QPD as
u_{QPD}. By measuring either $\hat{u}_{\mathrm{stage}}$ or $\hat{u}_{\mathrm{QPD}}$ as a function of frequencies, we obtain the total
probe response $\hat{u}_{\mathrm{FB}}(\omega)$ under feedback tracking. By estimating the sinusoidal force $\hat{F}_{\mathrm{FB}}(\omega)$
applied to the probe under the feedback (as given in Supplementary S1), we obtain the probe
particle's response function as $\tilde{\alpha}(\omega) = \langle\hat{u}_{\mathrm{FB}}(\omega)\rangle\big/\hat{F}_{\mathrm{FB}}(\omega)$. We then obtain $G(\omega)$ via Eq. (1).
220

Statistical analysis

222   Since the distributions of G' and G'' were non-Gaussian, statistical analysis was performed
223   by using the Mann–Whitney U test, unless otherwise stated in the figure caption. P values <
224   0.05 were considered a significant difference. All the statistical tests were performed using
225   Igor and Matlab software. A list of all P values from the MR experiments is given in
226   Supplementary S7.

RESULTS

Power-law intracellular rheology of untreated HeLa cells

Using feedback-tracking AMR, we measured the shear viscoelastic modulus of HeLa cells
derived from a cervical cancer. Melamine particles ($2a = 1$ μm) were incorporated into cells
forming an epithelial-like confluent monolayer on the surface of a glass-bottom dish.
Particles incorporated at the center of cells between the cell membrane and the nuclear
membrane were used as probes. The average of the complex shear modulus of the untreated
HeLa cells is shown in Fig. 2 together with the standard deviation ($n = 21$). Filled circles in
(A) and open circles in (B) show the real (G') and imaginary (G'') parts of the complex shear

modulus, $G(\omega) = G'(\omega) - iG''(\omega)$, respectively. Curves show raw data before averaging.

Markers are averaged values and bars indicate the log-normal SD. Since the logarithms of

mechanical properties are approximately distributed with Gaussian (41), averages and SDs

were calculated for the logarithm of experimental results. Our results showed a characteristic

frequency dependence of the form $G(\omega) \propto (-i\omega)^{1/2}$. $G'(\omega)$ slightly deviates from the power-

law dependency at low frequencies ($\leq 1\,\mathrm{Hz}$), but this may be an artifact. As shown in

Supplementary S2, low-frequency fluctuations of probe particles in Madin-Darby canine

kidney (MDCK) cells are decreased by prolonged irradiation with a 1.7 mW laser, implying

an increased elasticity at low frequencies. Accordingly, our recent preliminary experiments

FIGURE 2 Intracellular complex shear viscoelastic modulus measured with AMR using $2a = 1$ μm diameter melamine particles. (A) $G'(\omega)$ and (B) $G''(\omega)$ of HeLa cells in a confluent monolayer formed on a glass substrate. Curves are the raw data obtained in 23 different cells. Data shown as dotted lines were excluded from the average (n =21). Since cell cycles were not monitored, cells in different cell-cycle phases (G1, S, and G2) are included. Circles are the statistical average, and bars indicate the log-normal SD. Broken and solid lines show the power-law dependency typical of the glassy cytoplasm ($\propto \omega^{1/2}$) and cytoskeletal networks ($\propto \omega^{3/4}$), respectively. The result follows $\propto \omega^{1/2}$ over a wide range of frequencies, indicating that the intracellular mechanics are predominantly governed by the glasssy cytoplasm. (C) Immunofluorescent images of confluent HeLa cells. The cells were stained with Rhodamine phalloidin and Hoechst33342 to visualize F-actin (red) and the nucleus (blue).

performed more carefully with a reduced laser power (<0.5 mW) in MDCK cells seems to
remove this deviation. This will be reported in more detail elsewhere.
It has been reported that cytoskeletal networks prepared *in vitro* exhibit an elastic plateau
$G(\omega) \propto (-i\omega)^0$ at low frequencies, and a power-law dependency $G(\omega) \propto (-i\omega)^{3/4}$ at higher
frequencies (16, 19, 43, 44). The frequency dependency, which is now accepted as the
hallmark of cytoskeletons (20, 43), is explained by the theory for semi-flexible polymer
networks (45-48). In biological cells, a $G(\omega)$ consistent with this understanding was found
when MR was conducted using probe particles coupled to actin cytoskeletons (6, 41, 49). On
the other hand, the power-law dependence $G(\omega) \propto (-i\omega)^{1/2}$ observed in this study is not
consistent with that expected for the cytoskeleton. It is to be noted that the probe particles
used in this study were coated with polyethylene glycol polymers that generally inhibit
adherence to objects in cells, including the cytoskeleton. Actin cytoskeletons are usually less
expressed in confluent monolayers of cells compared to isolated cells (50). In addition, the
probes were incorporated deep inside of living cells where the actin cytoskeleton is only
sparsely expressed (Fig. 2C), as reported in confluent HeLa cells (51, 52) and also in
confluent epithelial-like monolayers of MCF10A and MDCK (50, 53).
261
Marginal contribution of the actin cytoskeleton to intracellular mechanics
In order to clarify the contribution of cytoskeletons to intracellular $G(\omega)$, we conducted
AMR in HeLa cells in which F-actin was disrupted. After conducting the first AMR
experiments in untreated HeLa cells, actin polymerization was inhibited by adding
Cytochalasin D or Latrunculin B to the culture medium (54, 55). A second round of AMR
measurements was then performed using the same particles as those used in the first AMR
experiments. Since $G(\omega)$ shows a general trend proportional to $(-i\omega)^{1/2}$, results normalized
as $G'/\omega^{1/2}$ are shown in Fig. 3 in order to clarify the effect of the chemical dose at each
frequency. Cytochalasin D at 50 μg/ml did not alter $G(\omega)$ in any frequency range (n = 9, Fig.
3A, and B). Statistical tests at each frequency were conducted for $G'(\omega)$ and $G''(\omega)$ before
and after the addition of Cytochalasin D. P values obtained by the Mann–Whitney U test
were larger than 0.05 for all frequencies measured, which indicates no significant difference.
Similar results were obtained by treating cells with another inhibitor for actin polymerization,
Latrunculin B at 10 μM (Fig. S3). Cells changed their morphology after the inhibitors were
administrated (Fig. 3C). They became more rounded while leaving branched protrusions, lost
cell-cell contact; some cells were even removed from the glass substrate. These observations
indicate that the inhibitors worked effectively and disrupted cytoskeletal organization, since
the actin cytoskeleton is responsible for maintaining the shape of cells and their adhesion to
the substrate (56). We thus conclude that actin cytoskeletons do not contribute to the
intracellular $G(\omega)$ of HeLa cells cultured in a confluent monolayer in any remarkable way.
HeLa cells in confluent monolayers are laterally confined and polarized under normal
conditions. They are roughly as tall as they are wide, and tightly connected to each other. In
such epithelial-like tissue, actin stress fibers are densely expressed at the cell-cell and cell-
substrate interfaces, and they contribute to the organization of the tissue. The disruption of
actin thus remarkably affects cell morphology. In prior studies, mechanical properties of cells

have been measured by using other techniques, *e.g.* AFM (atomic force microscopy) (9, 57-
59), MTC (magnetic twisting cytometry) (37, 41, 60), and others (41, 61, 62). These
techniques have shown that the mechanical properties of cells indeed depend on actin
expression (59). It is therefore commonly believed that the mechanical properties of cells are
directly correlated with their shape and cytoskeletal organization (61, 62), in contrast to our
experimental results. It is to be noted, however, that the techniques used in prior studies
measure the properties of cell surfaces at which actin cytoskeletons are densely expressed.

FIGURE 3 (A) $G'/\omega^{1/2}$ and (B) $G''/\omega^{1/2}$ of HeLa cells, untreated (circles) and treated (triangles) with 50 µg/ml Cytochalasin D. The same probe particles in the same cells were measured before and after adding Cytochalasin D to the culture media (n = 9). Bars indicate the log-normal SD. Similar experiments performed before and after treatment with 10 µM Latrunculin B are given in Fig. S3. (C, D) A microscopy image of HeLa cells. When actin cytoskeletons were disrupted with 50 µg/ml Cytochalasin D, cells strongly contract, leaving branched protrusions. (A, B) The Mann–Whitney U test was used to calculate the P value between untreated cells and cells treated with 50 µg/ml Cytochalasin D. No significant difference was observed for all the frequencies.

Considering that F-actin is less expressed inside of the epithelial cells in a confluent
monolayer, results obtained in this study are reasonable and do not contradict any of the prior
measurements conducted on cell surfaces.
Cell surface elasticities measured using conventional techniques are usually widely
distributed, over orders of magnitude (41, 58, 63). The organization of the actin cytoskeleton
at the cell cortex is heterogeneous and changes with time. Depending on the situation, actin
filaments bundle to form thick fibers, couple to cell membranes, and create tension due to
actomyosin contractile activity, leading to remarkable stiffening as demonstrated **in vivo** (64-
66) and **in vitro** (16, 19, 67). All these and many other factors contribute to variations in the
mechanical properties of cell surfaces. On the other hand, $G(\omega)$ in living cytoplasm is found
to be narrowly distributed, even though they were measured with different probes in different
cells. This observation also supports the claim that the intracellular mechanics of the
epithelium are not strongly associated with the actin cytoskeleton.
307
Cell mechanics during cell-cycle progression (G1 to S and G2 phases)
Next, we performed feedback-tracking AMR in HeLa cells while the cell cycle
progresses during interphase. HeLa cells used here (HeLa/Fucci2) were labelled using the
Fucci system that enables us to identify G1 and S/G2 phases by the fluorescence color in the
cell nucleus (68). In Fig. 4, we show the normalized complex shear modulus (A: $G'/\omega^{1/2}$ and
B: $G''/\omega^{1/2}$) of HeLa cells as a function of frequency. Circles and squares indicate cells at
G1 and S/G2 phases respectively. The intracellular $G(\omega)$ in the S/G2 phase slightly
increased compared to cells at G1 phase; G' and G'' of S/G2 phase were 40% and 30% larger
than those of the G1 phase on average, respectively. Note that these differences were only
marginal but observed because the cell cycle phases could be resolved.
In the literature, the cell-cycle dependency of cell mechanics has been investigated using
atomic force microscopy (AFM) (26) and micropipette aspiration (27), both of which showed
a 50~100% increase of surface elasticity of S-phase cells compared with that of G1-phase
cells. Here, we emphasize that our results and prior studies do not contradict. Our AMR
methodology is indifferent to F-actin, and likely measure the interstices of the sparse network
of the cytoskeleton. On the other hand, prior studies measured the response of the
cytoskeleton at cell surfaces whose expression is likely strongly affected by cell-cycle
progression.
As explained in the introduction, various processes that occur during cell-cycle
progression are inherently mechanical. Especially, cells in the M phase experience drastic
variations in cytoskeletal organization (62, 69), which would affect cell mechanics derived
from the cytoskeleton (28, 70, 71). When the probe particle was coupled to the cytoskeleton,
MR in the M phase could exhibit mechanical properties more evidently different from other
phases; the cell interior underwent softening and the viscoelastic fluid contributions increased
(28). However, AMR experiments were not conducted in M phase cells in this study. Because
mitosis and cytokinesis are highly dynamic processes and complete in a relatively short
period of time, it was not possible to stably track the probe particle in M phase cells with the
feedback technique.

FIGURE 4 (A) $G'/\omega^{1/2}$ of HeLa/Fucci2 cells in G1 (circles, n = 12) and S/G2 phases (squares, n = 9). (B) $G''/\omega^{1/2}$ of HeLa/Fuuci2 cells in G1 (circles) and S/G2 phases (squares). Bars indicate the estimate of the log-normal SD. Remarkable differences were not observed. (C) Schematic illustration of the Fucci system used for identifying cell-cycle phases. The Mann–Whitney U test was used to calculate the P value for differences between the G1 and S/G2 phases. *: P < 0.05. No symbol: P > 0.05.

ATP-dependent intracellular mechanical properties

After finding that the actin cytoskeleton and cell-cycle progression during interphase
have marginal effects on the intracellular mechanics, we now investigate the effect of
metabolism on active mechanics in HeLa cells. In order to regulate the metabolic activity,
cells were incubated with a culture media that causes ATP-depletion (41). Firstly,
intracellular ATP was measured using the luminescent ATP detection assay. Figure 5A
shows that the ATP level significantly decreased by more than an order of magnitude in ATP
depleted cells. On applying AMR, it was found that the ATP-depleted cells showed an elastic

plateau such that $G(\omega) \propto G_0 + A(-i\omega)^{1/2}$ in contrast to $G(\omega) \propto (-i\omega)^{1/2}$ for untreated cells,

as shown in Fig. 5B. To quantitatively evaluate the effect of ATP-depletion on $G(\omega)$, we

fitted $G(\omega)$ of untreated and ATP-depleted cells with $G_0 + A(-i\omega)^{1/2}$ (black dashed lines in

Fig. 5B). The coefficient A hardly changed while the elastic plateau G_0 of ATP-depleted

cells became ~ 10 times larger than that of untreated cells. This result clearly indicated that

only $G'(\omega)$ at low frequency significantly increased due to ATP depletion. In a similar

manner to previous sections, $G'(\omega)/\omega^{1/2}$ and $G''(\omega)/\omega^{1/2}$ of untreated HeLa cells (n = 21,

red dashed line) and ATP-depleted cells (n = 9, black filled circles) are shown in Fig. 5C and

D, respectively. Error bars indicate log-normal SDs except for $G''(\omega)/\omega^{1/2}$ in ATP-depleted

cells. Instead, we show the median, 25% and 75% quantiles for $G''(\omega)/\omega^{1/2}$ in ATP-depleted

cell because $G''(\omega)$ at small ω sometimes showed negative values due to $G'(\omega) \gg G''(\omega)$

and measurement noise. The Mann–Whitney U test for G' in the low frequency range (0.1 to

94 Hz) showed a significant increase of G' upon ATP depletion ($P < 0.05$) compared to

untreated cells.

We note that feedback-tracking MR in cells is most easily conducted when the probe

particle is situated in the homogeneous part of the cell. Experimental errors increase when

the probe particle is in a region close to the nucleus, cell membranes, or other large organelle,

because the inhomogeneity of optical properties in such region disturbs the propagation of

the probe and the drive laser. It frequently occurred that a probe particle that was used in the

first AMR measurement moves to regions where measurement is no longer possible. In Fig.

S4, we compare the limited number of data points (n =3) successfully measured using the

same probe particle in the same cells. All the results showed a remarkable increase in

viscoelasticity when ATP was depleted.

Fig. 5E shows microscope images of HeLa cells taken after ATP depletion. ATP-depleted

cells lose tight adherence to the substrate and to neighboring cells. Cells change their

morphology and take on more rounded shapes due to decreased cytoskeletal tension. To

clarify the effect of ATP-depletion on actin cytoskeletal structures, F-actin was stained using

Rhodamine phalloidin (Fig. 5F). Similar to the untreated cells (Fig. 2C), the actin

cytoskeleton of ATP-depleted cells was also sparse. Thus, we concluded that the actin

cytoskeleton was not responsible for the increase in the elastic plateau during ATP depletion.

Note that *in vitro* cytoskeletons are remarkably stiffened under active tension or pre-stress

(14-16). Consistently, the cell surface elasticity measured with AFM decreases when ATP is

depleted (72, 73) likely because the active stiffening of the cytoskeleton is eased owing to

the relax of actomyosin contraction at the cell cortex (61, 74-76). On the contrary,

intracellular elasticity measured in this study increased significantly when the intracellular

ATP was depleted. This observation again indicates that the cell cortex cytoskeleton is not

relevant to intracellular mechanics. When cells are subjected to the ATP-depletion medium

for a prolonged period, they could enter the path towards cell death (77, 78). The fluorescence

indicator (fluorescence caspase 3) for apoptosis indeed showed positive in HeLa cells

prepared using the similar protocol (Fig. S5). Except for the emergence of the elastic plateau,

however, the mechanical properties typical for intracellular rheology $G(\omega) \propto (i\omega)^{1/2}$ were

still observed for ATP-depleted cells. Therefore, our results indicate that the cytoplasm with
less expression of the cytoskeleton stiffens in a metabolism-dependent manner.

FIGURE 5 (A) Intracellular ATP level in HeLa cells treated with 50 mM 2-deoxyglucose and 10 mM NaN3 (ATP-depleted) for 7 h was measured. Quantification was performed on five independent experiments. All data are presented as mean ± standard deviation. ***: p < 0.001 (Student's t-test). (B) $G'(\omega)$ of untreated and ATP-depleted HeLa cells. $G(\omega)$ of untreated and ATP-depleted HeLa cells are fitted by $G_0 + A(-i\omega)^{1/2}$. Untreated: $G_0 = 136$ Pa. $A = 11.1$ Pa s$^{-0.5}$. ATP-depleted: $G_0 = 13.1$ Pa. $A = 11.2$ Pa s$^{-0.5}$. (C) $G'(\omega)/\omega^{1/2}$ and (D) $G''(\omega)/\omega^{1/2}$ of untreated (red closed circles n = 21) and ATP-depleted HeLa cells (black open squares, n = 9). (C) Bars indicate the log-normal SD. (D) Symbols and bars of $G''(\omega)/\omega^{1/2}$ for untreated cells indicate the log-normal mean and SD, respectively. $G''(\omega)$ of ATP-depleted cells at low frequency sometimes became negative due to $G'(\omega) \gg G''(\omega)$ and measurement noise. Thus, symbols of $G''(\omega)/\omega^{1/2}$ for ATP-depleted cells represent median values, and the bars connect 25% and 75 % quantiles. At low frequencies ($f \leq 94$ Hz), $G'(\omega)$ showed a significant increase after ATP depletion (P < 0.05). (C, D) The Mann–Whitney U test was used to calculate the P value for differences between untreated and ATP depleted cells. ***: P < 0.001. **: P < 0.01. *: P < 0.05. No symbol: P > 0.05.

390
391

(E) Microscope image of ATP-depleted HeLa cells. ATP-depleted HeLa cells became round and lost cell-cell contact. (F) Immunofluorescent images of ATP-depleted HeLa cells. The cells were stained with Rhodamine phalloidin and Hoechst33342 to visualize F-actin (red) and nuclei (blue).

DISCUSSION

It has been believed that mechanical properties of cells are mostly determined by the cytoskeleton (8, 9). This study showed, however, that intracellular mechanics in a confluent epithelium are scarcely affected by the actin cytoskeleton. Note that there are multitudes of other biomacromolecules that are crowded together in living cells. Although their effects on cell mechanics have not been fully appreciated until recently, it was shown that *in vitro* cytoplasm that lacks a cytoskeleton has glass-forming capabilities; the cytoplasm becomes glassy (*i.e.* practically solidified owing to the slowdown of dynamics) when the concentration of polymeric components is increased and/or metabolic activities are inhibited (3, 7, 37, 79). Vigorous fluctuations were observed in cells maintaining ordinary metabolism, but these were remarkably decreased in ATP-depleted cells (3, 36, 41, 80). It was then inferred that the glassy cytoplasm is fluidized by metabolic activities. However, the decrease/increase of fluctuations does not necessarily mean vitrification/fluidization. Even if glass-forming materials are deeply quenched and solidified, they could still flow if they are actively stirred (81). Besides, only observing the spontaneous fluctuations does not tell us how strongly the medium is stirred by metabolic activities. In this study, we directly measured intracellular viscoelasticity with AMR by observing a probe particle's response to a well-determined force. Our direct measurement showed that the cytoplasm in ATP-depleted cells was glassy while cells with ordinary metabolic activities were fluidized.

Cells are constructed of soft materials that respond non-linearly to applied forces. For instance, semiflexible polymer networks (model for cytoskeletons) typically stiffen under tension (14-20) while glassy colloidal suspensions (model for the cytoplasm without the cytoskeleton) are fluidized by mechanical loads (21-24). In living cells, various mechano-

enzymes such as motor proteins perform physiological functions by generating forces
utilizing energy supplied by metabolism. The motor-generated forces then create non-thermal
fluctuations that surpass thermal fluctuations as demonstrated by the violation of the
fluctuation-dissipation theorem, both *in vivo* (6, 36, 82-85) and *in vitro* (16, 19, 33, 86, 87).
Owing to the nonlinear response of constituent materials, mechanical properties of cells are
profoundly altered by the active fluctuations. In the living cytoplasm, vigorous active
fluctuations presumably induce structural relaxations that could not occur with thermal
activation (6, 7), leading to active fluidization. Here, the fluidization of the cytoplasm under
metabolic activities is consistent with the concept of a glassy cytoplasm, but not with
cytoskeletons that stiffen with active metabolism (16, 19, 67).

The complex shear viscoelastic modulus $G(\omega)$ in ordinary HeLa cells showed a
characteristic frequency dependence $G(\omega) = A(-i\omega)^{1/2}$. This dependency may seem to
coincide with the prediction of the Rouse model (88) that describes the high-frequency
dynamics of flexible polymers. In accordance with the theory, cross-linked poly-acrylamide
gels show $G(\omega) \propto (-i\omega)^{1/2}$, but only at high frequencies (6, 89). Networks of cross-linked
semi-flexible polymers usually show $G(\omega) \propto (-i\omega)^{3/4}$, but this changes to $G(\omega) \propto (-i\omega)^{1/2}$
when the network is placed under tension (16, 48). Therefore, polymer networks could also
show $G(\omega) \propto (-i\omega)^{1/2}$ at high frequencies when they are cross-linked. However, the
polymer-network models predict that the prefactor A is proportional to monomer
concentration (47, 88). On the other hand, A for extracted cytoplasm and for cytoplasm in
living cells grow much-more rapidly, super-exponentially and exponentially as a function of
macromolecule concentration, respectively (7). Such rapid growth of viscoelasticity has been
typically observed in glass-forming materials close to the glass transition (90) but not in
polymer networks. Note that the static resilience of cross-linked polymer networks should
not increase at all when the metabolic activity is inhibited: in fact, the opposite should happen
for crosslinked cytoskeletons (16). Also, the mechanical properties of flexible polymer
networks are hardly affected by the kind of mechanical perturbations expected under
physiological conditions (15, 17).

We believe that the universal frequency dependency $G(\omega) \propto (-i\omega)^{1/2}$ of living
cytoplasm also conforms to the concept of a glassy cytoplasm. Recent theoretical studies on
jamming rheology have shown that this frequency dependency may be derived from the
anomalous relaxation modes in the disordered medium close to the jamming transition,
crowded with colloidal objects with slippery interfaces (38, 39). In a critical point scenario,
the system is marginally stable; it displays excess relaxation modes at low frequencies and
becomes highly susceptible to external perturbations (91). When metabolic activity was
decreased in ATP-depleted cells, the elastic plateau arose as $G(\omega) = G_0 + A(-i\omega)^{1/2}$. Similar
viscoelastic behavior with an elastic plateau has been commonly observed in glassy or
jammed suspensions which were quenched beyond jamming (39, 92, 93). However, the
plateau G_0 disappeared when they are driven out of equilibrium under the application of
external flow (40), and only $G(\omega) = A(-i\omega)^{1/2}$ remains, much like in the living cytoplasm.
The glassy suspensions exhibited the rheological properties typical for critical jamming [*i.e.*,

$G(\omega) \propto (-i\omega)^{1/2}$] when they were mechanically driven out of equilibrium. Our results in cells
may indicate that living cytoplasm also reflects such a critical point-like situation.

Cytoskeletons also show glassiness when they are driven out of equilibrium, *e.g.*, when
the network of semiflexible polymers is forced to flow beyond the linear response regime
(94, 95). A glassy worm-like chain model explains such behavior by assuming that the height
of the free-energy barrier for the structural modification of the network is widely spread (96).
Even if thermal activation is not sufficient to overcome the barrier within the experiment
duration, the broad spectrum of the energy landscape becomes apparent in the nonlinear flow
behavior. At equilibrium, the linear response of *in vitro* cytoskeletons is understood as a
network of semi-flexible polymers (33, 46, 97). However, cytoskeletons are driven out of
equilibrium in living cells by active mechano-enzymes (81). With non-thermal perturbation,
structural relaxations are induced in the cytoskeleton of living cells, leading to glassy
cytoskeletal rheology. Such glassy behavior may be described by an effective noise
temperature as determined in the theory of "soft glassy rheology" (22). Indeed, glassy
rheology was observed in cytoskeletons at cell surfaces *in vivo* (60, 63), but in a way distinct
from colloidal crowding. $G(\omega)$ of the cell cortex cytoskeleton showed widely spread power-
law exponents [β for $G(\omega) \propto (-i\omega)^{\beta}$], depending on the situations and the types of cells.

FIGURE 6 Jamming phase diagram. Energy-depleted cells are in a glassy state due to molecular crowding (black circle). The living cytoplasm, however, reaches a marginal glass phase (red circle) due to mechanical force, but does not enter a liquid phase (blue circle).

On the other hand, $G(\omega) = A(-i\omega)^{1/2}$ seems to be universally observed in the
intracellular cytoplasm of MDCK cells, mouse embryonic stem cells, *etc*, as we will report
elsewhere. Not only the power-law exponent ($\beta = 0.5$), but the prefactor A also does not
largely differ for different kind of cells, as long as cells and the probe particles are prepared
such that the intracellular cytoplasm (but not the cytoskeleton) are measured, as done in this
study. Again, we emphasize that these results are consistent with glassy materials close to
jamming. One way to disprove theoretical models listed above is to seek the reason (1) why
this universal behavior appears specifically for cytoplasmic mechanics in living cells and (2)
why normal cells do not fluidize beyond the critical jamming point, but stay there. These
questions may be related to the marginal stability of active cytoplasm; the activity of
mechano-enzymes is regulated in a specific manner to achieve a situation similar to critical
jamming, as we will discuss in a forthcoming paper (Fig. 6) (84).

The interior of a biological cell is an enormously complex system, made up of an
innumerous number of molecule types and is maintained under continuous metabolic
turnover. It is therefore challenging to unequivocally prove a physical model of cells by
solely investigating cells *in vivo*. Even if one inhibits or enhances a specific biochemical
pathway to achieve more detailed control of metabolism, cells try to maintain homeostasis
by bypassing the path in the complex network of metabolism. Thus, it would be necessary to
establish a simplified model system that includes only the essences of living cytoplasm:
activity and crowding. During the last decade, a model system collectively referred as "active
matter" has been intensively investigated. Active matter is composed of objects which
convert the internal or ambient source of energy into mechanical form. During the dissipation
of the input energy, novel collective behavior of the objects was seen to emerge (98, 99). On
the other hand, the physics of densely packed disordered materials (glass and jamming),
especially the mechanism for dynamic arrest and melting, has been a key topic of non-
equilibrium statistical mechanics over 100 years (90, 100). Recently, the active matter and
glass research communities have begun to share a common interest in investigating dense
active matter or active glasses (101) which we believe is a good theoretical model of living
cytoplasm. Whereas fluctuations and dynamics have been investigated so far (102-108),
physical and mechanical properties behind readily observable phenomena have been rarely
measured. In this study, we found that mechanical properties predicted for critical jamming
somehow emerge in the living cytoplasm even though some of the fundamental premises of
jamming theory do not apply. Therefore, to reveal the nature of jamming rheology in the
cytoplasm, simplified dense active matter which allow for control over density, active force
and other conditions is desirable for future work.

CONCLUSION

In this study, we investigated the mechanical properties of HeLa cells in a confluent
epithelial monolayer. The probe particle's displacement in response to a sinusoidal optical-
trapping force was observed using a feedback-tracking microrheology technique (6, 20, 35).
The complex shear modulus $G(\omega)$ of cytoplasm surrounding the probe was then obtained.

In contrast to prior measurements of the cell's cortex, the intracellular shear viscoelastic
modulus showed a frequency dependency $G(\omega) \propto (-i\omega)^{1/2}$, which was seldom affected by
the cell-cycle progression during interphase and the disruption of the actin cytoskeleton.

On the other hand, ATP depletion clearly affected the mechanical properties of the cell
interior; an elastic plateau at low frequencies emerged in ATP-depleted cells. These
observations are consistent with the recently emerging idea that the cell interior is glassy (3,
7). Mechano-enzymes such as molecular motors enhance structural relaxations (yielding) by
actively stirring the cytoplasm, as is typical in glassy media under mechanical loads. Our
results imply that the intracellular cytoplasm is fluidized such that they approach the critical
jamming state, characterized by $G(\omega) \propto (-i\omega)^{1/2}$ (38, 91). Instead of complete fluidization,
the slow relaxation typical of a marginally stabilized system was observed in the range of
frequencies measured. We note that metabolism-dependent glassy mechanics has various
implications for biology. For example, a positive feedback between mechanics and
metabolism may occur in cells, as a cytoplasm fluidized by non-thermal activity may
facilitate metabolic processes (5, 80, 109-112).

ACKNOWLEDGEMENTS
This work was supported by JSPS KAKENHI Grant Number JP22H04848, JP21H01048,
JP20H05536, JP20H00128. We thank Christoph Schmidt in Duke University, and Atsushi
Ikeda in the University of Tokyo for helpful discussions and technical support.

CONFLICTS OF INTEREST The authors declare no conflict of interest.

AUTHOR CONTRIBUTION
H.E., K.U., K.N., W.N., and S.I. collected/analyzed MR data. T.M. performed biochemical
assays and immunocytochemistry. H.E., Y.S., and D.M. analyzed/discussed the results. D.M.
designed/supervised the project, and wrote the manuscript with the support of H. E. and Y.S.

REFERENCES
1. Dix, J. A., and A. S. Verkman. 2008. Crowding effects on diffusion in solutions and
cells. Annu Rev Biophys 37:247-263.
2. Minton, A. P. 2001. The influence of macromolecular crowding and macromolecular
confinement on biochemical reactions in physiological media. J Biol Chem
276(14):10577-10580.
3. Parry, B. R., I. V. Surovtsev, M. T. Cabeen, C. S. O'Hem, E. R. Dufresne, and C.
Jacobs-Wagner. 2014. The Bacterial Cytoplasm Has Glass-like Properties and Is
Fluidized by Metabolic Activity. Cell 156(1-2):183-194.
4. Riedel, C., R. Gabizon, C. A. Wilson, K. Hamadani, K. Tsekouras, S. Marqusee, S.
Presse, and C. Bustamante. 2015. The heat released during catalytic turnover
enhances the diffusion of an enzyme. Nature 517(7533):227-230.

5.   Jee, A. Y., Y. K. Cho, S. Granick, and T. Tlusty. 2018. Catalytic enzymes are active
matter. Proc Natl Acad Sci U S A 115(46):E10812-E10821.
6.   Nishizawa, K., M. Bremerich, H. Ayade, C. F. Schmidt, T. Ariga, and D. Mizuno.
2017. Feedback-tracking microrheology in living cells. Sci Adv 3(9):e1700318.
7.   Nishizawa, K., K. Fujiwara, M. Ikenaga, N. Nakajo, M. Yanagisawa, and D. Mizuno.
2017. Universal glass-forming behavior of in vitro and living cytoplasm. Sci Rep
7(1):15143.
8.   Fletcher, D. A., and R. D. Mullins. 2010. Cell mechanics and the cytoskeleton. Nature
463(7280):485-492.
9.   Rigato, A., A. Miyagi, S. Scheuring, and F. Rico. 2017. High-frequency
microrheology reveals cytoskeleton dynamics in living cells. Nat Phys 13(8):771-775.
10.   Broedersz, C. P., and F. C. MacKintosh. 2014. Modeling semiflexible polymer
networks. Rev Mod Phys 86(3):995-1036.
11.   Zimmerman, S. B., and S. O. Trach. 1991. Estimation of macromolecule
concentrations and excluded volume effects for the cytoplasm of Escherichia coli. J
Mol Biol 222(3):599-620.
12.   Zimmerman, S. B., and A. P. Minton. 1993. Macromolecular Crowding -
Biochemical, Biophysical, and Physiological Consequences. Annu Rev Bioph Biom
22:27-65.
13.   Delarue, M., G. P. Brittingham, S. Pfeffer, I. V. Surovtsev, S. Pinglay, K. J. Kennedy,
M. Schaffer, J. I. Gutierrez, D. Sang, G. Poterewicz, J. K. Chung, J. M. Plitzko, J. T.
Groves, C. Jacobs-Wagner, B. D. Engel, and L. J. Holt. 2018. mTORC1 Controls
Phase Separation and the Biophysical Properties of the Cytoplasm by Tuning
Crowding. Cell 174(2):338-349 e320.
14.   Gardel, M. L., J. H. Shin, F. C. MacKintosh, L. Mahadevan, P. Matsudaira, and D. A.
Weitz. 2004. Elastic behavior of cross-linked and bundled actin networks. Science
304(5675):1301-1305.
15.   Storm, C., J. J. Pastore, F. C. MacKintosh, T. C. Lubensky, and P. A. Janmey. 2005.
Nonlinear elasticity in biological gels. Nature 435(7039):191-194.
16.   Mizuno, D., C. Tardin, C. F. Schmidt, and F. C. Mackintosh. 2007. Nonequilibrium
mechanics of active cytoskeletal networks. Science 315(5810):370-373.
17.   Head, D. A., E. Ikebe, A. Nakamasu, P. Zhang, L. G. Villaruz, S. Kinoshita, S. Ando,
and D. Mizuno. 2014. High-frequency affine mechanics and nonaffine relaxation in
a model cytoskeleton. Phys Rev E Stat Nonlin Soft Matter Phys 89(4):042711.
18.   Head, D. A., and D. Mizuno. 2013. Local mechanical response in semiflexible
polymer networks subjected to an axisymmetric prestress. Phys Rev E Stat Nonlin
Soft Matter Phys 88(2):022717.
19.   Mizuno, D., C. Tardin, and C. F. Schmidt. 2020. Rapid local compression in active
gels is caused by nonlinear network response. Soft Matter 16(40):9369-9382.
20.   Honda, N., K. Shiraki, F. Van Esterik, S. Inokuchi, H. Ebata, and D. Mizuno. 2022.
Nonlinear master relation in microscopic mechanical response of semiflexible
biopolymer networks. New J Phys 24(5):053031.

21.   Liu, A. J., and S. R. Nagel. 1998. Jamming is not just cool any more. Nature
396(6706):21-22.

22.   Sollich, P., F. Lequeux, P. Hebraud, and M. E. Cates. 1997. Rheology of soft glassy
materials. Physical Review Letters 78(10):2020-2023.

23.   Petekidis, G., D. Vlassopoulos, and P. N. Pusey. 2004. Yielding and flow of sheared
colloidal glasses. J Phys-Condens Mat 16(38):S3955-S3963.

24.   Wilson, L. G., A. W. Harrison, W. C. K. Poon, and A. M. Puertas. 2011.
Microrheology and the fluctuation theorem in dense colloids. Epl-Europhys Lett
93(5):58007.

25.   Alberts, B. 2015. Molecular biology of the cell. Garland Science, Taylor and Francis
Group, New York, NY.

26.   Kelly, G. M., J. I. Kilpatrick, M. H. van Es, P. P. Weafer, P. J. Prendergast, and S. P.
Jarvis. 2011. Bone cell elasticity and morphology changes during the cell cycle. J
Biomech 44(8):1484-1490.

27.   Tsai, M. A., R. E. Waugh, and P. C. Keng. 1996. Cell cycle-dependence of HL-60
cell deformability. Biophys J 70(4):2023-2029.

28.   Hurst, S., B. E. Vos, M. Brandt, and T. Betz. 2021. Intracellular softening and
increased viscoelastic fluidity during division. Nature Physics 17(11):1270-1276.

29.   Nijenhuis, N., D. Mizuno, J. A. Spaan, and C. F. Schmidt. 2012. High-resolution
microrheology in the pericellular matrix of prostate cancer cells. J R Soc Interface
9(73):1733-1744.

30.   MacKintosh, F. C., and C. F. Schmidt. 1999. Microrheology. Curr Opin Colloid In
4(4):300-307.

31.   Mason, T. G., K. Ganesan, J. H. vanZanten, D. Wirtz, and S. C. Kuo. 1997. Particle
tracking microrheology of complex fluids. Physical Review Letters 79(17):3282-
3285.

32.   Hough, L. A., and H. D. Ou-Yang. 2002. Correlated motions of two
hydrodynamically coupled particles confined in separate quadratic potential wells.
Phys Rev E Stat Nonlin Soft Matter Phys 65(2 Pt 1):021906.

33.   Mizuno, D., D. A. Head, F. C. MacKintosh, and C. F. Schmidt. 2008. Active and
Passive Microrheology in Equilibrium and Nonequilibrium Systems.
Macromolecules 41(19):7194-7202.

34.   Gittes, F., and C. F. Schmidt. 1998. Interference model for back-focal-plane
displacement detection in optical tweezers. Opt Lett 23(1):7-9.

35.   Sugino, Y., M. Ikenaga, and D. Mizuno. 2020. Optimization of Optical Trapping and
Laser Interferometry in Biological Cells. Appl Sci-Basel 10(14):4970.

36.   Guo, M., A. J. Ehrlicher, M. H. Jensen, M. Renz, J. R. Moore, R. D. Goldman, J.
Lippincott-Schwartz, F. C. Mackintosh, and D. A. Weitz. 2014. Probing the
Stochastic, Motor-Driven Properties of the Cytoplasm Using Force Spectrum
Microscopy. Cell 158(4):822-832.

37.   Zhou, E. H., X. Trepat, C. Y. Park, G. Lenormand, M. N. Oliver, S. M. Mijailovich,
C. Hardin, D. A. Weitz, J. P. Butler, and J. J. Fredberg. 2009. Universal behavior of

the osmotically compressed cell and its analogy to the colloidal glass transition. Proc
Natl Acad Sci U S A 106(26):10632-10637.

38. Tighe, B. P. 2011. Relaxations and rheology near jamming. Phys Rev Lett
107(15):158303.

39. Liu, A. J., S. Ramaswamy, T. G. Mason, H. Gang, and D. A. Weitz. 1996. Anomalous
viscous loss in emulsions. Phys Rev Lett 76(16):3017-3020.

40. Jacob, A. R., A. S. Poulos, A. N. Semenov, J. Vermant, and G. Petekidis. 2019. Flow
dynamics of concentrated starlike micelles: A superposition rheometry investigation
into relaxation mechanisms. J Rheol 63(4):641-653.

41. Hoffman, B. D., G. Massiera, K. M. Van Citters, and J. C. Crocker. 2006. The
consensus mechanics of cultured mammalian cells. Proc Natl Acad Sci U S A
103(27):10259-10264.

42. Miyamoto, T., H. Uosaki, Y. Mizunoe, S. I. Han, S. Goto, D. Yamanaka, M. Masuda,
Y. Yoneyama, H. Nakamura, N. Hattori, Y. Takeuchi, H. Ohno, M. Sekiya, T.
Matsuzaka, F. Hakuno, S. I. Takahashi, N. Yahagi, K. Ito, and H. Shimano. 2021.
Rapid manipulation of mitochondrial morphology in a living cell with iCMM. Cell
Rep Methods 1(4):100052.

43. Koenderink, G. H., M. Atakhorrami, F. C. MacKintosh, and C. F. Schmidt. 2006.
High-frequency stress relaxation in semiflexible polymer solutions and networks.
Phys Rev Lett 96(13):138307.

44. Mason, T. G., T. Gisler, K. Kroy, E. Frey, and D. A. Weitz. 2000. Rheology of F-
actin solutions determined from thermally driven tracer motion. J Rheol 44(4):917-
928.

45. Morse, D. C. 1998. Viscoelasticity of concentrated isotropic solutions of semiflexible
polymers. 1. Model and stress tensor. Macromolecules 31(20):7030-7043.

46. Morse, D. C. 1998. Viscoelasticity of concentrated isotropic solutions of semiflexible
polymers. 2. Linear response. Macromolecules 31(20):7044-7067.

47. Gittes, F., and F. C. MacKintosh. 1998. Dynamic shear modulus of a semiflexible
polymer network. Phys Rev E 58(2):R1241-R1244.

48. Caspi, A., M. Elbaum, R. Granek, A. Lachish, and D. Zbaida. 1998. Semiflexible
polymer network: A view from inside. Physical Review Letters 80(5):1106-1109.

49. Caspi, A., R. Granek, and M. Elbaum. 2002. Diffusion and directed motion in cellular
transport. Phys Rev E Stat Nonlin Soft Matter Phys 66(1 Pt 1):011916.

50. Pavel, M., M. Renna, S. J. Park, F. M. Menzies, T. Ricketts, J. Fullgrabe, A.
Ashkenazi, R. A. Frake, A. C. Lombarte, C. F. Bento, K. Franze, and D. C.
Rubinsztein. 2018. Contact inhibition controls cell survival and proliferation via
YAP/TAZ-autophagy axis. Nat Commun 9(1):2961.

51. Haase, K., and A. E. Pelling. 2013. Resiliency of the plasma membrane and actin
cortex to large-scale deformation. Cytoskeleton (Hoboken) 70(9):494-514.

52. Simone, L. C., S. Caplan, and N. Naslavsky. 2013. Role of phosphatidylinositol 4,5-
bisphosphate in regulating EHD2 plasma membrane localization. Plos One
8(9):e74519.

53. Stevenson, B. R., and D. A. Begg. 1994. Concentration-dependent effects of
cytochalasin D on tight junctions and actin filaments in MDCK epithelial cells. J Cell
Sci 107:367-375.
54. Casella, J. F., M. D. Flanagan, and S. Lin. 1981. Cytochalasin D inhibits actin
polymerization and induces depolymerization of actin filaments formed during
platelet shape change. Nature 293(5830):302-305.
55. Wakatsuki, T., B. Schwab, N. C. Thompson, and E. L. Elson. 2001. Effects of
cytochalasin D and latrunculin B on mechanical properties of cells. J Cell Sci 114(Pt
5):1025-1036.
56. GODMAN, G. C., A. F. MIRANDA, A. D. DEITCH, and S. W. TANENBAUM.
1975. ACTION OF CYTOCHALASIN D ON CELLS OF ESTABLISHED LINES.
J Cell Biol 64:644-667.
57. Haga, H., S. Sasaki, K. Kawabata, E. Ito, T. Ushiki, and T. Sambongi. 2000. Elasticity
mapping of living fibroblasts by AFM and immunofluorescence observation of the
cytoskeleton. Ultramicroscopy 82(1-4):253-258.
58. Wang, A., K. Vijayraghavan, O. Solgaard, and M. J. Butte. 2016. Fast Stiffness
Mapping of Cells Using High-Bandwidth Atomic Force Microscopy. Acs Nano
10(1):257-264.
59. Rotsch, C., and M. Radmacher. 2000. Drug-induced changes of cytoskeletal structure
and mechanics in fibroblasts: an atomic force microscopy study. Biophys J 78(1):520-
535.
60. Fabry, B., G. N. Maksym, J. P. Butler, M. Glogauer, D. Navajas, and J. J. Fredberg.
2001. Scaling the microrheology of living cells. Phys Rev Lett 87(14):148102.
61. Bacabac, R. G., D. Mizuno, C. F. Schmidt, F. C. MacKintosh, J. J. W. A. Van Loon,
J. Klein-Nulend, and T. H. Smit. 2008. Round versus flat: Bone cell morphology,
elasticity, and mechanosensing. J Biomech 41(7):1590-1598.
62. Stewart, M. P., J. Helenius, Y. Toyoda, S. P. Ramanathan, D. J. Muller, and A. A.
Hyman. 2011. Hydrostatic pressure and the actomyosin cortex drive mitotic cell
rounding. Nature 469(7329):226-230.
63. Fabry, B., G. N. Maksym, J. P. Butler, M. Glogauer, D. Navajas, N. A. Taback, E. J.
Millet, and J. J. Fredberg. 2003. Time scale and other invariants of integrative
mechanical behavior in living cells. Phys Rev E Stat Nonlin Soft Matter Phys 68(4 Pt
1):041914.
64. Tee, S. Y., J. Fu, C. S. Chen, and P. A. Janmey. 2011. Cell shape and substrate rigidity
both regulate cell stiffness. Biophys J 100(5):L25-27.
65. Grimm, K. B., H. Oberleithner, and J. Fels. 2014. Fixed endothelial cells exhibit
physiologically relevant nanomechanics of the cortical actin web. Nanotechnology
25(21):215101.
66. Ayala, Y. A., B. Pontes, B. Hissa, A. C. Monteiro, M. Farina, V. Moura-Neto, N. B.
Viana, and H. M. Nussenzveig. 2017. Effects of cytoskeletal drugs on actin cortex
elasticity. Exp Cell Res 351(2):173-181.

67.   Koenderink, G. H., Z. Dogic, F. Nakamura, P. M. Bendix, F. C. MacKintosh, J. H.
721   Hartwig, T. P. Stossel, and D. A. Weitz. 2009. An active biopolymer network
722   controlled by molecular motors. Proc Natl Acad Sci U S A 106(36):15192-15197.

723   68.   Sakaue-Sawano, A., H. Kurokawa, T. Morimura, A. Hanyu, H. Hama, H. Osawa, S.
724   Kashiwagi, K. Fukami, T. Miyata, H. Miyoshi, T. Imamura, M. Ogawa, H. Masai,
725   and A. Miyawaki. 2008. Visualizing spatiotemporal dynamics of multicellular cell-
726   cycle progression. Cell 132(3):487-498.

727   69.   Taubenberger, A. V., B. Baum, and H. K. Matthews. 2020. The Mechanics of Mitotic
728   Cell Rounding. Front Cell Dev Biol 8:687.

729   70.   Fischer-Friedrich, E., Y. Toyoda, C. J. Cattin, D. J. Müller, A. A. Hyman, and F.
730   Jülicher. 2016. Rheology of the Active Cell Cortex in Mitosis. Biophys J 111(3):589-
731   600.

732   71.   Garzon-Coral, C., H. A. Fantana, and J. A. Howard. 2016. A force-generating
733   machinery maintains the spindle at the cell center during mitosis. Science 352:1124–
734   1127.

735   72.   Streppa, L., F. Ratti, E. Goillot, A. Devin, L. Schaeffer, A. Arneodo, and F. Argoul.
736   2018. Prestressed cells are prone to cytoskeleton failures under localized shear strain:
737   an experimental demonstration on muscle precursor cells. Sci Rep 8(1):8602.

738   73.   Embry, A. E., H. Mohammadi, X. Niu, L. Liu, B. Moe, W. A. Miller-Little, C. Y. Lu,
739   L. A. Bruggeman, C. A. McCulloch, P. A. Janmey, and R. T. Miller. 2016.
740   Biochemical and Cellular Determinants of Renal Glomerular Elasticity. Plos One
741   11(12):e0167924.

742   74.   Sutton, T. A., H. E. Mang, and S. J. Atkinson. 2001. Rho-kinase regulates myosin II
743   activation in MDCK cells during recovery after ATP depletion. Am J Physiol-Renal
744   281(5):F810-F818.

745   75.   Yubero, M. L., P. M. Kosaka, A. San Paulo, M. Malumbres, M. Calleja, and J.
746   Tamayo. 2020. Effects of energy metabolism on the mechanical properties of breast
747   cancer cells. Commun Biol 3(1):590.

748   76.   Gupta, S. K., and M. Guo. 2017. Equilibrium and out-of-equilibrium mechanics of
749   living mammalian cytoplasm. J Mech Phys Solids 107:284-293.

750   77.   Eguchi, Y., S. Shimizu, and Y. Tsujimoto. 1997. Intracellular ATP levels determine
751   cell death fate by apoptosis or necrosis. Cancer Res 57(10):1835-1840.

752   78.   Lieberthal, W., S. A. Menza, and J. S. Levine. 1998. Graded ATP depletion can cause
753   necrosis or apoptosis of cultured mouse proximal tubular cells. Am J Physiol
754   274(2):F315-327.

755   79.   Fernandez-Marin, B., I. Kranner, M. San Sebastian, U. Artetxe, J. M. Laza, J. L. Vilas,
756   H. W. Pritchard, J. Nadajaran, F. Miguez, J. M. Becerril, and J. I. Garcia-Plazaola.
757   2013. Evidence for the absence of enzymatic reactions in the glassy state. A case
758   study of xanthophyll cycle pigments in the desiccation-tolerant moss Syntrichia
759   ruralis. J Exp Bot 64(10):3033-3043.

760   80.   Weber, S. C., A. J. Spakowitz, and J. A. Theriot. 2012. Nonthermal ATP-dependent
761   fluctuations contribute to the in vivo motion of chromosomal loci. Proc Natl Acad Sci
762   U S A 109(19):7338-7343.

81.   Fakhri, N., A. D. Wessel, C. Willms, M. Pasquali, D. R. Klopfenstein, F. C.
MacKintosh, and C. F. Schmidt. 2014. High-resolution mapping of intracellular
fluctuations using carbon nanotubes. Science 344(6187):1031-1035.
82.   Ahmed, W. W., E. Fodor, M. Almonacid, M. Bussonnier, M. H. Verlhac, N. Gov, P.
Visco, F. van Wijland, and T. Betz. 2018. Active Mechanics Reveal Molecular-Scale
Force Kinetics in Living Oocytes. Biophys J 114(7):1667-1679.
83.   Turlier, H., D. A. Fedosov, B. Audoly, T. Auth, N. S. Gov, C. Sykes, J. F. Joanny, G.
Gompper, and T. Betz. 2016. Equilibrium physics breakdown reveals the active
nature of red blood cell flickering. Nature Physics 12(5):513-519.
84.   Umeda, K., K. Nishizawa, W. Nagao, Y. Sugino, N. Oyama, H. Ebata, and D. Mizuno.
To be submitted. Activity-dependent glassy mechanics in living cells II; Non-thermal
fluctuations under metabolic activity.
85.   Mizuno, D., R. Bacabac, C. Tardin, D. Head, and C. F. Schmidt. 2009. High-
resolution probing of cellular force transmission. Phys Rev Lett 102(16):168102.
86.   Ariga, T., M. Tomishige, and D. Mizuno. 2018. Nonequilibrium Energetics of
Molecular Motor Kinesin. Phys Rev Lett 121(21):218101.
87.   Bertrand, O. J., D. K. Fygenson, and O. A. Saleh. 2012. Active, motor-driven
mechanics in a DNA gel. Proc Natl Acad Sci U S A 109(43):17342-17347.
88.   Doi, M., and S. F. Edwards. 1986. The theory of polymer dynamics. Clarendon Press;
Oxford University Press, Oxford, New York.
89.   Schnurr, B., F. Gittes, F. C. MacKintosh, and C. F. Schmidt. 1997. Determining
microscopic viscoelasticity in flexible and semiflexible polymer networks from
thermal fluctuations. Macromolecules 30(25):7781-7792.
90.   Angell, C. A., K. L. Ngai, G. B. McKenna, P. F. McMillan, and S. W. Martin. 2000.
Relaxation in glassforming liquids and amorphous solids. J Appl Phys 88(6):3113-
3157.
91.   Berthier, L., and G. Biroli. 2009. Glasses and Aging, A Statistical Mechanics
Perspective on. Encyclopedia of Complexity and Systems Science. R. A. Meyers,
editor. Springer New York, New York, NY, pp. 4209-4240.
92.   Wyss, H. M., K. Miyazaki, J. Mattsson, Z. Hu, D. R. Reichman, and D. A. Weitz.
2007. Strain-rate frequency superposition: a rheological probe of structural relaxation
in soft materials. Phys Rev Lett 98(23):238303.
93.   Krishan, K., A. Helal, R. Hohler, and S. Cohen-Addad. 2010. Fast relaxations in foam.
Phys Rev E Stat Nonlin Soft Matter Phys 82(1 Pt 1):011405.
94.   Semmrich, C., T. Storz, J. Glaser, R. Merkel, A. R. Bausch, and K. Kroy. 2007. Glass
transition and rheological redundancy in F-actin solutions. Proceedings of the
National Academy of Sciences of the United States of America 104(51):20199-20203.
95.   Chapman, C. D., and R. M. Robertson-Anderson. 2014. Nonlinear microrheology
reveals entanglement-driven molecular-level viscoelasticity of concentrated DNA.
Phys Rev Lett 113(9):098303.
96.   Kroy, K., and J. Glaser. 2007. The glassy wormlike chain. New J Phys 9:416.

97. Atakhorrami, M., G. H. Koenderink, J. F. Palierne, F. C. MacKintosh, and C. F.
Schmidt. 2014. Scale-Dependent Nonaffine Elasticity of Semiflexible Polymer
Networks. Physical Review Letters 112(8):088101.

98. Marchetti, M. C., J. F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, M. Rao, and
R. A. Simha. 2013. Hydrodynamics of soft active matter. Rev Mod Phys 85(3):1143-
1189.

99. Bechinger, C., R. Di Leonardo, H. Lowen, C. Reichhardt, G. Volpe, and G. Volpe.
2016. Active Particles in Complex and Crowded Environments. Rev Mod Phys
88(4):045006.

100. Hunter, G. L., and E. R. Weeks. 2012. The physics of the colloidal glass transition.
Rep Prog Phys 75(6):066501.

101. Janssen, L. M. C. 2019. Active glasses. J Phys Condens Matter 31(50):503002.

102. Mongera, A., P. Rowghanian, H. J. Gustafson, E. Shelton, D. A. Kealhofer, E. K.
Carn, F. Serwane, A. A. Lucio, J. Giammona, and O. Campas. 2018. A fluid-to-solid
jamming transition underlies vertebrate body axis elongation. Nature 561(7723):401-
405.

103. Berthier, L., and J. Kurchan. 2013. Non-equilibrium glass transitions in driven and
active matter. Nature Physics 9(5):310-314.

104. Nandi, S. K., R. Mandal, P. J. Bhuyan, C. Dasgupta, M. D. Rao, and N. S. Gov. 2018.
A random first-order transition theory for an active glass. Proceedings of the National
Academy of Sciences of the United States of America 115(30):7688-7693.

105. Oyama, N., T. Kawasaki, H. Mizuno, and A. Ikeda. 2019. Glassy dynamics of a model
of bacterial cytoplasm with metabolic activities. Phys Rev Res 1(3):032038.

106. Mandal, R., P. J. Bhuyan, P. Chaudhuri, C. Dasgupta, and M. Rao. 2020. Extreme
active matter at high densities. Nat Commun 11(1):2581.

107. Mandal, R., P. J. Bhuyan, M. Rao, and C. Dasgupta. 2016. Active fluidization in
dense glassy systems. Soft Matter 12(29):6268-6276.

108. Koyano, Y., H. Kitahata, and A. S. Mikhailov. 2019. Diffusion in crowded colloids
of particles cyclically changing their shapes. Epl-Europhys Lett 128(4):40003.

109. Ross, J. L. 2016. The Dark Matter of Biology. Biophys J 111(5):909-916.

110. Ariga, T., K. Tateishi, M. Tomishige, and D. Mizuno. 2021. Noise-Induced
Acceleration of Single Molecule Kinesin-1. Phys Rev Lett 127(17):178101.

111. Nishizawa, K., N. Honda, M. Ikenaga, S. Inokuchi, Y. Sugino, T. Arigac, and D.
Mizuno. 2021. Feedback Microrheology in Soft Matter.
https://ui.adsabs.harvard.edu/abs/2021arXiv210605119N (posted).

112. Ezber, Y., V. Belyy, S. Can, and A. Yildiz. 2020. Dynein Harnesses Active
Fluctuations of Microtubules for Faster Movement. Nat Phys 16(3):312-316.

841