Targeted insertion and reporter transgene activity at a gene safe harbor of the
human blood fluke, *Schistosoma mansoni*

Wannaporn Ittiprasert[1,*], Max F. Möscheid[2], Cristian Chaparro[3], Victoria H. Mann[1],
Thomas Quack[2], Rutchanee Rodpai[1,4], André Miller[5], Prapakorn Wistiphongpun[1,6],
Watunyoo Buakaew[1,7], Margaret Mentink-Kane[5], Legendré Miller[5], Sarah Schmid[5],
Anastas Popratiloff[8], Christoph G. Grevelding[2,#,*], Christoph Grunau[3, #,*], Paul J.
Brindley[1, #,*]

[1] Department of Microbiology, Immunology & Tropical Medicine, School of Medicine & Health
Sciences, George Washington University, Washington, D.C. 20037

[2] Institute of Parasitology, Biomedical Research Center Seltersberg, Justus Liebig University
Giessen, Giessen, Germany

[3] IHPE, Univ Perpignan Via Domitia, CNRS, IFREMER, Univ Montpellier, Perpignan, France

[4] Department of Parasitology and Excellence in Medical Innovation, and Technology Research
Group, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand

[5] Schistosomiasis Resource Center, Biomedical Research Institute, Rockville, MD, 20850

[6] Faculty of Medical Technology, Rangsit University, Pathum Thani, 12000, Thailand

[7] Cellular and Molecular Immunology Research Unit, Faculty of Allied Health Sciences,
Naresuan University, Phitsanulok, 65000, Thailand

[8] Nanofabrication and Imaging Center, Science & Engineering Hall, George Washington
University, Washington, D.C. 20052

[#] Contributed equally

[*] Correspondence: Wannaporn Ittiprasert, wannaporni@gwu.edu; Paul J. Brindley,
pbrindley@gwu.edu

Abstract

Identification, characterization, and transgene insertion into a genomic safe harbor site (GSH) is
expected to facilitate consistent transgene activity without disruption to the host cell genome
while supporting transgene activity. The location of GSH sites was predicted in the human blood
fluke, *Schistosoma mansoni* using a computational approach, and after which a CRISPR-focused
protocol was developed for insertion of a reporter transgene into a representative GSH.
Ribonuclear protein complexes of Cas9 nuclease, three overlapping guide RNAs, and
phosphorothioate-modified, double-stranded donor DNA encoding green fluorescent protein
driven by a ubiquitously expressed promoter were employed. Gene knockout efficiencies of 24%
and reporter knockin transgene fluorescence of >70% of gene-edited schistosome eggs were
observed. These outcomes advance functional genomics for schistosomes by providing a
tractable path towards transgenic helminths using homology directed repair catalyzed transgene
insertion. This approach should be adaptable to helminths generally.

Main

Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) technology has
revolutionized functional genomics in in biology, medicine, and agriculture[1-3]. Transgenesis
approaches are integral in diverse applications including gene therapy, biotherapeutics,
deciphering host-pathogen interactions, and enhancements in agricultural production. With
progress emanating from model species and cell lines, tools and techniques can frequently be
adapted and transferred to non-model species. Among these are the helminths, which generally
are responsible for neglected tropical diseases that cause substantial mortality and disease[4].
These mostly occur in the global south and are responsible for a disease burden that exceeds that
caused by malaria and tuberculosis. In the current post-genomics era, access to CRISPR-based
transgenesis protocols for functionally characterizing parasitic helminth putative drug and
vaccine targets is a public health research priority. Importantly, progress of this technology in
one helminth, for example the schistosome will facilitate its use in other major invertebrate
clades found within the Protostomia, for which CRISPR-based reverse and forward genetics
have yet to be reported and/or are challenging.

CRISPR enables targeted site-specific mutation(s), obviating an impediment of earlier
transgenesis approaches that rely on lentiviruses[5] and transposons such as *piggyBac*[6]. For
example, these latter approaches could negatively lead to genetic instability, multi-copy
insertion, unstable expression or inactivation of the transgene and interference with the
endogenous gene under investigation. In the process of genome editing, the double stranded
breaks (DSBs) are resolved by several different repair mechanisms, including the predominant
error-prone non-homology end joining (NHEJ) and the templated homologous-directed repair
(HDR) pathways. Sister chromatids provide a natural repair template, whereas the latter also can
be donated by exogenous DNA such a plasmid, oligodeoxynucleotides, and PCR amplicons.
When supplied with double-strand (ds) donor DNA with modifications HDR efficiency can be
markedly improved [7]. The system of CRISPR/Cas-assisted HDR has been applied in
Schistosoma mansoni[8,9] with promoter-free, single strand-deoxynucleotide donors. Multiple
overlapping CRISPR target sites improve precise HDR insertion of large cargoes in embryonic
stem cells[10,11], while modification of 5'-termini of long dsDNA donors enhance HDR, favoring

efficient, single-copy integration through the retention of a monomeric donor confirmation and
thereby facilitating gene replacement and tagging[12].

Here, we undertook a computational investigation of potential genome safe harbor sites in *S.*
mansoni, and adapted CRISPR/Cas9-based approaches to insert a reporter transgene into one of
the predicted GSH situated in euchromatin of chromosome 3, GSH1. The targeted region was
free of repetitive sequences and neighboring long non-coding regions, a situation likely to
minimize off-target effects of CRISPR/Cas activity. Multiple prototypic adjacent motifs within
this region were targeted with two and three programmed single guide RNAs deployed in unison
to enhance mutational efficiency and homology directed repair (HDR) in the presence of a
phosphorothioate-modified donor DNA template. A knock-in (KI) efficiency of 75% was
observed by reporter transgene green protein fluorescence in miracidia within the eggs, when the
RNPs and donor transgene were delivered.

Results

Genome safe harbors predicted in the schistosome genome

We undertook a computational search to predict intergenic genome safe harbor (GSH) sites in *S.*
mansoni based on accepted criteria, newly introduced principles[13], and genome sequence
resources available for this schistosome, that could satisfy benign and stable gene expression.
Based on criteria presented in figure 1a and detailed in the Methods section, in brief regions
proximal to H3K4me3 marks, regions without H3K27me3, ATAC seq-positive regions, regions
free of repetitive sequences, and regions known to be accessible to lentiviral (HIV) insertion, the
draft genome of *S. mansoni*, version 9[14-17] included 6,884 regions with enrichment of H3K4me3
in the absence of H3K27me3, in available developmental stages, i.e. positive for H3K4me3 but
not K3K27me3. In adult schistosomes, we consistently found 10,533 ATAC-Seq positive
regions. There were 4,027 ATAC-Seq regions that overlapped with H3K4me3, and 2,915 regions
overlapped with the ATAC-Seq regions and H3K4me3 but not H3K27me3. Forty-two
unambiguous HIV integration sites were identified in these regions, and eight genes were < 4 kb
upstream or downstream from these integration sites[18]. To identify intergenic GSH, rather than
intragenic GSH given the transgene insertion in the former should be safer[13], we located 10,149
intergenic regions[19]. There were 9,985 regions beyond 2 kb upstream of the transcription start
site (TSS) and within 11 kb of an HIV integration site, and 8,837 regions outside long non-
coding-RNA (lncRNA), which were intersected to 95,587 unique intergenic regions beyond 2 kb
of lncRNAs of ≥ 100 bp in length. Next, 200 regions were identified intersecting with the
merged ATAC/ H3K4me3 signals and, of these, four were situated ≤ 11 kb from HIV integration
sites. These four regions satisfied our criteria. These were termed GSH1 (1,416 bp in length;
location, chromosome 3:13380432-13381848), GSH2 (970 bp; chromosome 2: 15434976-
15435954), GSH3 (752 bp; chromosome 2: 9689988-9690739), and GSH4 (138 bp;
chromosome 3: 13381901-13382038), respectively (Fig. 1), with the naming based in rank order
on their size from longest (GSH1) to shortest (GSH4). We note that several protein-coding gene
loci were situated in the vicinity of these gene-free GSHs, although these genes were > 2kb
distant from any GSH: Smp_052890 [uncharacterized protein], Smp_150460 [copper transport
protein], Smp_071830 [uncharacterized protein], Smp_245610 [uncharacterized protein] and
Smp_131070 [condensing complex] (Figs. 1b-d). Most of there were as yet uncharacterized

proteins, and may be non-essential genes based on ortholog to essential genes known from model
eukaryotes[20].

Efficiency of programmed mutation at GSH1 enhanced by multiple guide RNAs

Having predicted the location of these GSHs, we proceeded to investigate the efficiency of
programmed mutation and reporter transgene activity at GSH1, chromosome 3: 13380432-
13381848. Multiple and/or overlapping guide RNAs were employed, an approach that enhances
KI efficiency in some mammalian cell lines and embryos[10,11]. Among guide RNAs (gRNA)
exhibiting on-target specificity for GSH1, three overlapping synthetic guide RNAs, sgRNA1,
sgRNA2 and sgRNA3, which lacked self-complementarity and off-target matches to the
reference *S. mansoni* genome (Figs. 1b,e, Fig. 2a), were selected from among CRISPR/Cas9
target sites predicted by CHOPCHOP [21,22]. The ribonucleoprotein complexes (RNPs) of Cas9
nuclease and sgRNA were assembled, after which four discrete mixtures of the three RNPs were
used. Three of the mixtures included two RNPs, *i.e.* dual RNPs mixtures (RNP1+RNP2,
RNP2+RNP3, and RNP1+RNP3) and the fourth included the three RNPs, i.e. triple RNPs
(RNP1+RNP2+RNP3).

The mixture of the dual and triple RNPs, in the presence of the donor DNA template encoding
EGFP, was delivered to schistosome eggs by electroporation (EP). The transfected eggs were
maintained in culture for 15 days, after which EGFP expression was quantified. Thereafter,
genomic DNAs and total RNAs were extracted from the eggs. Efficiency of genome editing,
both in mock-treated controls and experimental groups, was assessed using DECODR[23] analysis
of Sanger sequence chromatograms of amplicons that spanned the CRISPR/Cas9 programmed
DSBs. Analysis of PCR products amplified from schistosome genomic DNA using primers
(indel-F, indel-R) flanking the DSBs (Fig. 2a) revealed programmed knockout (KO) efficiency,
as assessed by indel (insertions-deletions)-bearing alleles resulting from the dual gRNAs, as
follows: KO frequencies at GSH1 of 5.4% (range of 0.8-10.4%), 3.6% (range of 1.2-19.3%) and
12.6% (range of 4.9-19.3%) for RNP1+ RNP2, RNP2 + RNP3 and RNP1 + RNP3, respectively
(Fig. 2b). The dual gRNA RNPs mixtures induced short deletions of one to several nucleotides in
length at the predicted DSB sites for sgRNA1, 2 and/or 3 (Figs. 2c-e). Mutations were not
evident in amplicons from the control groups, *i.e.* wild type (no treatment) and mock treatment
(not shown). The triple RNPs mixture, where the three sgRNAs shared at least six overlapping
nucleotides, induced 23.9% KO (range of 2.4 - 71.9%), higher than any mixture of the dual
RNPs (Fig. 2b).

The CRISPR efficiencies at each target sites varied among the RNP mixtures, as did the indel
profiles (Fig. S1). KO efficiency at each target resulting from dual RNPs was lower generally
than for the triple guide RNPs. At target site 1, estimated KO rates were 6.9% (4.1-10.4%),
13.8% (6-19.3%) and 26.5% (14-33.2%) using RNP1+RNP2, RNP1+RNP3 and
RNP1+RNP2+RNP3, respectively. There were 3.9% (0.8%-5.8%), 1.8% (1.2%-2.4%) and
14.2% (2.4%-16.8%) mutations on target 2 using RNP1+RNP2, RNP2+RNP3 and
RNAP1+RNP2+RNP3, respectively. Mutation efficiencies at target 3 using RNP2+RNP3 and
RNP1+RNP3 were 13.9% (6%-19.3%) and 11.3% (6%-19.3%). The mutational profile of the
indels, as assessed from Sanger sequencing chromatograms of the target amplicons, was mostly
deletions rather than insertions (Figs. S1b-e). Notably, deletions up to 115 nt in length were
identified among the biological replicates with the triple overlapping RNPs (Fig. 2e, Fig. S1e).

KO efficiency was assessed using at least five independent biological replicates: Combining
three overlapping sRNAs induced an aggregate mutation efficiency (23.9%) higher than that
obtained with any of the dual RNPs, 5.4%, 3.6%, 12.6% ($P \leq 0.001$, one-way ANOVA). The KO
efficiency of the RNP1+RNP3 group was higher than that of either of the other dual RNPs
($P \leq 0.01$) (Fig. 2b).

Three overlapping guide RNAs enhanced efficiency of CRISPR knockin

As multiple sgRNAs with overlapping sequences can enhance CRISPR/Cas9-mediated HDR
efficiency[10] and given that triple overlapping sgRNAs performed better than dual gRNAs in
initiating programmed mutation at GSH1 in eggs (Fig. 2b), we investigated KI of a reporter
transgene at GSH1 with the triple overlapping sgRNA/RNPs mixture (Figs. 3a,b). We employed
the gene encoding enhanced green fluorescent protein (EGFP) driven by the promoter of the
endogenous *S. mansoni* ubiquitin gene and its cognate terminator region as the repair template
for programmed HDR (Fig. 3a). The donor template included symmetrical homology arms
specific for GSH1, located on the 5´-flank of target site 1 and the 3´-flanking region of target site
3 (Fig. 3b). The donor template was delivered as linearized long double-strand DNA (lsDNA) of
4,451 bp in length. Aiming to optimize precise and efficient single-copy integration of the donor
transgene into GSH1 by HDR, the 5´ termini of the DNA donor amplicons were chemically
modified[12] to shield the donor template from multimerization and from integration at the DSB
via the non-homologous end-joining (NHEJ) repair pathway (Fig. 3a).

At the outset, we investigated the impact of length of the homology arms (HA) by comparing the
performance of the donor repair template bearing homology arms of increasing length of 200,
400 and 600 bp. Dual (RNP1+RNP3) and triple (RNP1+RNP2+RNP3) sgRNA mixtures were
used in this investigation. EGFP expression was not evident in eggs electroporated with lsDNA
donors with 200 and 400 bp HAs at about day 5 after transfection (not shown). By contrast, we
observed a few EGFP-positive eggs, although in < 1% of them, with the lsDNA donor with 600
bp HA (not shown). Subsequently, given this outcome, we changed the focus of the investigation
for EGFP expression to transfection with the donor transgene flanked by 600 bp HA using the
triple guides RNP mixture (RNP1+RNP2+RNP3) and to monitoring expression of the EGFP
transgene for up to 15 days. Thereafter, on examination using spectrally resolved, confocal laser
scanning microscopy, the EGFP signal was detected in the eggs of the experimental group,
which received the CRISPR materials including the lsDNA donor with 600 bp HA. EGFP signal
remained until 15 days, when the experiment ended. EGFP signals were not observed in the
negative control groups, although the autofluorescence characteristic of schistosome eggs was
apparent[24]. The EGFP signal was detected in the lsDNA donor control (without RNPs) for
several days, indicating that extrachromosomal lsDNA expressed EGFP transiently after the
transfection.

Next, we investigated the nature of the programmed KI at GSH1 by PCR-based analysis for the
presence of the expected amplicons spanning the 5´ and 3´ flanks of the donor transgene i.e.,
flanking the ubiquitin promoter, EGFP, and the ubiquitin terminator sequences. At the F 5´
flank, we used a forward primer specific for the genome upstream of the 5´ HA paired with a
reverse primer specific for the ubiquitin promoter (Fig. 3b). For the 3´ integration junction, were
used a reverse primer specific for a site downstream of the 3´ terminus of the HA paired with a
forward primer specific for the ubiquitin terminator. Fragments representing the 3´ KI and 5´ KI

integration regions of 983 bp and 728 bp, respectively, were observed in the experiment
treatment groups but not in the other (control) groups (Fig. 3c). Expression of the EGFP
transgene was monitored using DNase-treated RNA, to eliminate the possibility of lsDNA donor
contamination and genomic DNA. EGFP transcripts were observed in the KI experimental
group, although some variability in transcript abundance among the biological replicates was
seen based on the signals obtained for *Sm*GAPDH, which served as the reference gene for
integrity of the RNAs (Fig. 3d).

Reporter transgene expressed in schistosome eggs

EGFP positivity and intensity were quantified using spectral laser scanning confocal
fluorescence microscopy[24]. Active transgene expression was confirmed within miracidia
developing inside transfected eggs (Figs. 4a, b). EGFP appeared to be expressed by numerous
diverse cells throughout the developing larva whereas morphological malformation was not
observed in transgenic eggs and their enclosed larvae. More intense EGFP fluorescence was
consistently recorded and quantified at 509 nm in eggs from the experimental treatment group
(Figs. 4b1, 2) than the mock control eggs and in eggs transfected solely with donor template
(Figs. 4a1, 2). Subsequently, on day 15 following transfection with the repair template in the
presence or absence of the RNPs mixture, we quantified EGFP intensity in eggs that contained a
miracidium by normalization with the EGFP signal from lsDNA only transfected eggs.
Fluorescence intensity differed markedly between these two groups; by 15 days after
transfection, 75% of miracidia in eggs from the experimental group (RNPs/lsDNA) emitted
EGFP fluorescence whereas 25% of eggs containing a miracidium transfected with the donor
lsDNA repair template only emitted EGFP (Figs. 4c, d) ($P \leq 0.001$; $n = 402$ eggs in experimental
group, $n = 397$ eggs in the lsDNA only group, aggregated from four independent biological
replicates). In addition, by scoring the intensity of fluorescence at 509 nm[24], the emission
wavelength for EGFP, as shown figures 4c1,c3, c4 (green curve), established by subtracting the
signal at 509 nm from the autofluorescence spectrum, especially at the egg shell, the EGFP-
specific signal in the control lsDNA donor repair template treatment group, mean = 1,290
arbitrary units (AU) [25] (range, 856-1,1713) was lower than the experimental group transfected
with the triple, overlapping guide RNPs mixture in the presence of the donor repair template,
6,905 AU (range, 4,973 to 8,963)($P \leq 0.001$) (Fig. S2). Moreover, emission of EGFP was not
detected in the control groups, *i.e.* mock-transfected and WT eggs (not shown). Lastly, diverse
cells and tissues of the developing miracidium expressed the fluorescence reporter gene, i.e.
EGFP was not obviously localized to specific cells (Figs. 4b,c).

Discussion

To advance functional genomics for helminths, here we predicted genome safe harbor sites in *S.*
mansoni, optimized conditions for delivery and structure of transgene cargo, inserted the reporter
transgene into a predicted intergenic genome safe harbor (GSH) by programmed CRISPR/Cas9
homology-directed repair, and quantified transgene activity using confocal imaging of emission
spectra specific for EGFP. Programmed editing with RNPs and three overlapping guide RNAs
targeting GSH1 delivered 24% mutation efficiency that was accompanied by transgene
fluorescence activity in 75% of miracidia in the gene-edited schistosome eggs.

Our criteria to predict GSH in the schistosome included location in euchromatin to avoid
silencing of the transgene, unique genome-target sequence to minimize off-target events,
avoidance of lncRNA encoding genes, presence of epigenetic marks for open chromatin
structure, and the absence of epigenetic marks indicating heterochromatin. We termed the
intergenic sites GSH1, -2, -3 and -4, which were located on chromosomes 2 and 3. (*S. mansoni*
has seven pairs of autosomes and one pair of sex chromosomes, Z and W). In addition, we
assessed the GSH1 locus for CRISPR/Cas9 integration, gene editing and over expression of
EGFP. We edited GSH1 using two or three ribonucleoprotein complexes of Cas9 endonuclease
with the overlapping guide RNAs. Overlapping CRISPR target sites were selected by the
CHOPCHOP web-based tool for CRISPR/Cas9 directed mutagenesis. RNPs of triple
overlapping guides delivered significantly higher CRISPR/Cas9 efficiency, and larger sized
deletion mutations than RNPs using dual guides. this outcome would also enhance the efficiency
of HDR based on reports with rodents, where it has been speculated that shorter rather than
longer indels are predominate after Cas9 catalyzed cleavage and the shorter indels enhance
HDR[10,26]. Thereafter, we deployed this approach to targeted knockin of a long (4,551 bp)
doubled stranded DNA donor template encoding the green fluorescence reporter driven by the *S.*
mansoni ubiquitin gene promoter and terminator elements.

These findings indicated that GSH1 represented a promising safe harbor site for forward
genetics-focused genomics within this schistosome. Efficient HDR resulted from the
combination of multiple target site overlapping RNPs programmed to cleave GSH1 in the
presence of a repair template protected by chemical modifications. Notably, 75% of mature eggs
exhibited reporter transgene fluorescence in the miracidium developing within the eggshell, and
significantly more fluorescence than seen in the control eggs transfected with donor template but
not with the RNPs, ~5%. EGFP signals were not present in the control, untreated wild type egg,
which by 9-10 days following transfection exhibits minimal background fluorescence[39]. Our
approach to circumvent the autofluorescence emitted by schistosome eggs, which can confound
detection of GFP transgene activity, used confocal laser spectral imaging and linear
unmixing[27,28], an approach to our knowledge previously not used to facilitate quantification of
GFP specific emission to resolve overlap between the GFP and endogenous fluorophores of the
egg of the helminth.

This study confirmed that GSH1 is a prospective safe harbor site for germline transgenesis,
although further validation will surely be informative. Limitations with the other prospective
GSHs, GSH2, -3 and 4 included absence of efficient CRISPR/Cas9 sites and predicted off-
targets which precluded further analysis,, although comparisons of the utility of all four GSH
including with other CRISPR tools such as Cas12a[9] and base editors[29] can be considered.
Although distance from known genes was one of our criteria to predict a schistosome GSH,
intragenic sites rather than intergenic GSH may in some cases exhibit expedient attributes for
functional genomics in *S. mansoni* where detrimental impacts to the fitness the schistosome may
be far less consequential, for example for *in vitro*-focused studies. Despite drawbacks for clinical
use in human gene therapy[30], several intragenic safe harbors are used in experimental human
gene therapy, including *CCR5*, *AAVSI*, and *Rosa26*, and all reside within gene-rich loci and
whereas they have been targeted with therapeutic gene cargo, including for example the insertion
of *FANCA* at *AAVS1* in CD34+ hematopoietic progenitors from Fanconi anemia patients[31].
Intergenic sites nonetheless are inherently safer given coding regions or other elements are not
disrupted by transgene insertion[30]. Recently, additional sites, named *Rogi 1* and *Rogi 2*, located

in intergenic regions, conform with additional criteria focusing on enhancement of safety for use
in human gene therapy, including location distant from endogenous genes, lncRNAs, enhancers,
and telomeres[13].

These findings are consequential in that they advance functional genomics and forward genetics
for a hitherto unmet challenge to manipulate a pathogen of global public health significance.
They confirm that transgenes can be targeted to safe integration sites to endow individual stages
or populations of these pathogens with novel functions, with broad potential for basic and
translational studies[32-34]. Whereas these findings deal with somatic transgenesis of the
schistosome larva, use of the same approach for transfection of the newly laid egg of *S. mansoni*,
a stage that at its origin includes a single zygote (surrounded by vitelline yolk cells) and which,
therefore, is a window to the germline[37,38] and therefore facilitate germline transgenesis to derive
lines of transgenic parasites carrying gain- or loss-of-function mutations. In addition, the editing
methods developed can be adapted for knock-out approaches of genes of interest, in
schistosomes, and likely other platyhelminths, for which genome sequences are available to be
analyzed for prospective genome safe harbors. The information presented here enabled novel
insight into efficient transgenesis and forward genetics for *S. mansoni* and will promote forward
genetics approaches in functional genomics for parasitic helminths.

Online content

Methods

Computational search for gene safe harbors in *Schistosoma mansoni*

We undertook a genome analysis focusing of intergenic (gene-free) regions to identify
prospective GSHs, using similar approaches as those used on the human genome[13]. We aimed to
locate a GSH, a site that would facilitate stable expression of the integrated transgene free of
interference from the host genome and which, in parallel, integrates and transcribes the transgene
without negative consequences or loss of fitness for the host cell[33]. The search for GSHs
deployed included several criteria, First, its location should be adjacent to peaks of H3K4me3, a
histone modification associated with euchromatin and transcription start sites[35]. Second, it should
not be near or not containing H3K27me3 in any developmental stage, a histone mark associated
with heterochromatin[35]. Third, as the schistosome genome contains highly repetitive elements[15],
the GSH site should be located in a unique tract of the genome sequence. Fourth, it should reside
in open, euchromatic chromatin accessible to Tn5 transposase as assessed from ATAC-
sequencing, which provides a positive display of transposase integration events[36]; consequently,
safe harbor candidate regions should deliver an ATAC-sequence signal. Fifth, in the vicinity of
known HIV integration sites, given that HIV integrates preferentially into euchromatin in human
cells[37], we anticipated that HIV integration into the schistosome genome may likewise indicate a
region of euchromatin (Fig. 1a).

To predict loci conforming to the criteria, pooled ChIP-seq data for H3K4me3 and K3K27me2
from previous studies were aligned against on *S. mansoni* genome data (version 9 on the date of
analysis). ATAC-seq was performed as described with modification[19]. Peakcalls of ChIP-seq and
ATAC-Seq were done with ChromstaR[13,35,36,38] and stored as Bed files. Bed files were used to
identify the presence of H3K4me3 and absence of H3K27me3 in adults, miracidia, *in vitro*

sporocysts, cercariae and *in vitro* schistosomula with bedtools intersect. Thereafter, ATAC-seq
data from adult male and adult female worms (two replicates each) were intersected to find
common ATAC-positive regions. H3K4me3-only (H3K27me3-absent) common to all stages and
ATAC signals were intersected to find common regions. Next, the HIV integration sites were
identified by using data from ERR33833.8[51]. Reads were mapped to the lentivirus genome (HIV-
1 vector pNL-3, accession AF324493.2) using Bowtie2 with default parameters. Paired reads
were extracted where one end mapped to HIV and the other end mapped to schistosome genome
at a unique location. Genes from the BED files above that located ≤ 11 kb HIV-1 integration
sites were identified with bedtools closestbed. Gene expression data for these genes were
obtained using the metanalysis tool, https://meta.schisto.xyz/analysis/, of Lu and Berriman[39].

Computational searches that addressed these criteria predicted, *a priori*, gene free (intergenic)-
GSH (Fig. 1), given that transgene integration into an existing gene could diminish fitness of the
genetically modified cell[40,41]. We defined genes as protein coding sequences and sequences
coding for long non-coding RNA (lncRNA). In view of our goal to use CRISPR/Cas mediated-
HDR to insert the transgene, we searched preferentially for unique sequences, to obviate off-
target gene modification, and excluded gene free-regions composed of repetitive sequences.
Those unique sequences were also annotated outside lncRNA, regions beyond putative
promotors that we deemed as 2 kb upstream of the transcription termination site (TTS), and the
regions close to peaks of H3K4me2 in all parasite stages which never contained H3K27me3. The
regions overlapping with ATAC-seq positive sites with ≤11 kb distance from HIV integration
sites also were included. (The HIV-1 genome is ~10 kb in length.) BEDtools were used to
delimit 2 kb upstream regions (FlankBed). Annotations of 16,583 lncRNA were pooled from
http://verjolab.usp.br/public/schMan/schMan3/macielEtAl2019/files/macielEtAt2019.bed12[42].
Repeats were masked with RepeatMasker V4.1.0 using a specific repeat library produced with
RepeatModeler2 V2.0.1 and stored as a GFF file. BED files with coordinates outside these
annotations were generated by BedTools complementBed. BedTools Multiple Intersect was used
to identify regions that are common to unique regions (complement of repeatmasker), intergenic
regions, ≥ 2 kb upstream and outside of lncRNA. Only regions which a length ≥ 100 bp were
retained. These regions were intersected with merged H3K4me3-only common to all
developmental stages and ATAC signals (euchromatic signal). BedTools ClosestBed was used to
determine distance to the nearest HIV provirus integration.

Schistosome eggs

Mice (female, Swiss Webster) infected with *S. mansoni* were obtained from the Schistosomiasis
Resource Center (Biomedical Research Institute, Rockville, MD) within seven days of infection
by cercariae (180 cercariae/mouse/percutaneous infection). The mice were housed at the Animal
Research Facility of George Washington University, which is accredited by the American
Association for Accreditation of Laboratory Animal Care (AAALAC no. 000347) and has the
Animal Welfare Assurance on file with the National Institutes of Health, Office of Laboratory
Animal Welfare, OLAW assurance number A3205. All procedures employed were consistent
with the Guide for the Care and Use of Laboratory Animals. The Institutional Animal Care and
Use Committee of the George Washington University approved the protocol used for
maintenance of mice and recovery of schistosomes.

Mice were euthanized six to seven wk after infection, after which schistosomes were recovered
by portal vein perfusion with 150 mM NaCl, 15mM sodium citrate, pH 7.0. The liver was
resected, homogenized with a. tissue blender, and the homogenate incubated with collagenase at
37°C for 18 h. Thereafter, schistosome eggs from the digested livers were recovered by Percoll
gradient centrifugation, as described[43]. Eggs isolated from livers, termed "liver eggs", LE[44], were
cultured in 20% inactivated bovine serum, 2% antibiotic/antimycotic at 5% CO_2, 37°C overnight
before being subjected to transfection with the CRISPR materials.

Guide RNAs, ribonucleoprotein complexes

Here, we focused on GSH1, located on *S. mansoni* chromosome 3; 13231955-13233370 (Fig.
1d), an intergenic safe harbor site of 1,416 nt, the longest in length of the four GSH (Figs. 1 b–d
). Guide RNAs (gRNA) for GSH1 were designed with the assistance of the CHOPCHOP[21,22,45]
tools, using the annotated *S. mansoni* genome[14], to predict target sites, off-targets, and efficiency
of CRISPR/Cas9 programmed cleavage. CHOPCHOP predicted three overlapping guide RNAs
targeting three cleavage sites withinGSH1 with the three DSBs at six to 12 nt apart from each
other. All three overlapping gRNAs lacked off-target sites and lacked self-complementarity. The
three gRNAs were located on the forward strand of GSH1 at nucleotide positions 605-624, 617-
636, and 623-642, respectively (Fig. 2a). We termed the predicted DSBs as target 1, target 2 and
target 3. Synthetic guide RNAs (sgRNA), specifically Alt-R CRISPR-Cas9 sgRNA chemically
modified to enhance functional stability, and recombinant *Streptococcus pyogenes* Cas9
nuclease, specifically Alt-R HiFi Cas9, which includes nuclear localization sequences (NLS),
were sourced from Integrated DNA Technologies, Inc. (IDT, Coralville, IA). Each
ribonucleoprotein complex (RNP) was complexed in the separate tube, with Cas9 and a single
sgRNA at 1:1 ratio in 25 µl Opti-MEM; 10 µl of 1 µg/µl sgRNA (Opti-ME as diluent) was
mixed with 10 µl of 1 µg/µl Cas9 (in Opti-MEM) by gentle pipetting and incubated for 10 min at
23°C to allow the RNP to assemble.

Doubled stranded DNA donor

A plasmid vector, pUC-Ubi-EGFP-ubi, was constructed by chemical synthesis of the donor
transgene and its ligation into pUC (Azenta Life Sciences, Chelmsford, MA). The inserted
sequence included homology arms of 600 bp length corresponding to GSH1 at 22-621 nt (5´-
homology arm) and 640-1239 nt (3´-homology arm), respectively, flanking the in frame
expression cassette of the *S. mansoni* ubiquitin promoter (2,056 bp), EGFP (717 bp), and the
ubiquitin terminator (578 bp). Plasmid DNA was amplified using Phusion High-Fidelity DNA
Polymerase (New England Bio-Labs, Ipswich, MA, cat no. M0530) with primers specific for the
5´ and 3´ termini of the homology arms of 200, 400 or 600 bp in length (Table S2). Thus, three
forms of the donor transgene DNA were prepared, with homology arms of increasing length –
200, 400 and 600 bp. The primers employed to amplify the transgene DNA from the pUC-Ubi-
EGFP-ubi plasmid were 5×phosphorothioate-modified to enhance stability of the amplified
DNA. (5´-modified long dsDNA donor (lsDNA) enhances HDR and favors efficient single-copy
integration by its retention of a monomeric conformation[12].) (Fig. 3a). PCRs were carried out in
reaction volumes of 50 µl in 200 µM dNTPs, 0.5µM of each primer, 100 ng pUC-Ubi-EGFP-ubi,
3% DMSO and 1 unit of Phusion DNA polymerase, with thermocycling of 98°C, 30 sec, 30
cycles of 98°C, 10 sec, 55°C, 30 sec, 72°C, 3 min, and final extension at 72°C, 10 min.

Amplificons were isolated using the NucleoSpin Gel and PCR Cleanup and gel extraction kit
(Takara, San Jose, CA, cat no. 740609), eluted in 30 µl nuclease-free water, and the long
stranded (ls) DNA donor transgene stored at -20°C until used.

Transfection of schistosome eggs

Ten thousand eggs (LE) of *S. mansoni* were washed three times with chilled (4°C) 1×PBS before
transfer into a chilled electroporation cuvette (4 mm electrode gap, BTX, Holliston, MA) with
Opti-MEM as the electroporation buffer. Each 25 µl of RNP along with the lsDNA donor were
immediately added into the cuvette with the schistosome eggs, to a total cuvette volume of 300
µl with Opti-mem: specifically, for the dual guide RNA/RNPs, group 1) 25 µl RNP1+25 µl
RNP2, group 2) 25 µl RNP2+25 µl RNP3, and group 3) 25 µl RNP1+25 µl RNP3.. In groups
with the lsDNA, 10 µg of this donor DNA was added into the cuvette before bringing the final
volume to 300 µl/cuvette. Transfection of schistosome eggs with CRISPR materials was
undertaken using square wave electroporation (Electro SquarePorator ECM 830, BTX), with a
single 125 volt pulse for 20 ms[8,9,46]. The transfected eggs were transferred to culture medium, as
above.

Nucleic acids

To recover genomic DNA and total RNA, eggs from each replicate were triturated in ~100 µl
DNA/RNA Shield solution (Zymo Research, cat no. R1100, Irvine, CA) using a motor-driven
homogenizer fitted with a disposable pestle and collection tube (Biometer II, Bunkyo-ku, Tokyo,
Japan). DNA was isolated from 50% of the homogenate, and RNA from the remainder. 250 µl
DNAzol® ES (Molecular Research Center, Inc., Cincinnati, OH) was dispensed into the
homogenate, and DNA recovered according to the manufacturer's protocol. Total RNA was
extracted from the homogenate by adding 250 µl RNAzol RT (Molecular Research Center, Inc.).
Yields and purity were assessed quantified by spectrophotometry (NanoDrop One
Spectrophotometer, Thermo Fisher Scientific), based on the ratios of absorbance at 260/280 and
at 260/230 nm[47].

Analysis of CRISPR on-target efficiency

Amplicons of GSH1 spanning the programed DSBs were obtained using population genomic
DNA (above) and primers termed 'indel-F and indel-R primers' that cover the region flanking
expected double strand break of all the CRISPR target sites (Fig. 2a, Table S1). Amplification
products were purified (NucleoSpin Gel and PCR Cleanup and gel extraction kit, Takara; cat no.
740609) and the nucleotide sequences determined by Sanger cycle sequencing (Azenta Life
Sciences, South Plainfield, NJ). Chromatograms of the sequence traces of experimental and
control group(s) was compared using DECODR[23] at default parameters with single and multiple
CRISPR target analysis (Figs. 2b-2f, S1a). The pattern of indels in was confirmed by TIDE
analysis of Sanger sequence reads [48] (Figs. S1b-e).

Detection of integration of the transgene into the schistosome genome

Integration of the donor transgene at GSH1 was analyzed by PCR, using GoTaq G2 DNA
polymerase (Promega, Madison, WI) and two pairs of primers; one primer located on the GSH1

using specific primers upstream (5' KI-F) or downstream (3' KI-R) of the homology arms paired
with primers specific for the transgene (5' KI-R or 3' KI-F) (Fig. 2b, Table S2), as described[9].
The PCR cycling conditions were 95°C, 2 min, 40 cycles 94°C, 15 sec, 58°C 30 sec, 72°C, 60
sec, and the amplification products were separated by electrophoresis and stained with ethidium
bromide. The expected product sizes for the 5' and 3' integration site-specific amplicons were
728 and 983 bp, respectively, and the amplification control (amplified by indel-F and indel-R
primers), expected product size, 764 bp (Fig. 2c).

Quantification of transgene expression in schistosome eggs

To examine the mRNA expression of EGFP, total RNAs extracted from LE were exposed to
DNase to eliminate residual genomic DNAs and donor lsDNA donor were transcribed into
cDNA using the Maxima First Strand cDNA synthesis kit with DNase (Thermo Fisher
Scientific). The qPCR was performed using the GoTaq® G2 DNA polymerase (cat no. M7841,
Promega, Madison, WI) with the specific primers; EGFP-F 5´-atggtgagcaagggcgagg-3´and
EGFP-R 5´-cttgtacagctcgtccatgcc-3´ (Fig. 3b) with expected amplicon at 717 bp. *S. mansoni*
GAPDH (Smp_056970) served as the reference gene. The specific primer for GAPDH-specific
oligos: GAPDH-F; 5´-atgggacatttccaggcgag-3´, GAPDH-R; 5´-ccaacaacgaacatgggtgc-3´,
expected amplicon of 285 bp in length. PCR cycling conditions: 95°C, 2 min, 25 cycles 94°C,
15 sec, 58°C, 30 sec, 72°C, 30 sec, after which amplification products were examined, as above.

Quantification of fluorescence by spectral imaging and linear unmixing

Spectral and spatial distribution of EGFP fluorescence were assessed using confocal laser
scanning microscopy, using a Zeiss LSM710 Meta detector fitted Axiovert 200 (Carl Zeiss, Jena,
Germany). Images were collected with the C-Apochromat 20×, 1.2 NA water immersion
objective. Spectroscopic measurements were performed in response to excitation by 458 nm
(16.5 μW) Arion laser line and 633 nm He/Ne laser line (Lasos Lasertechnik, Jena, Germany),
which were used for focus and transmission mode imaging. Emission was detected with a
spectral META detector at 16 channels, scanning at 477 nm through 638 nm, simultaneously. A
hurdle for the detection of EGFP using fluorescence microscopy is presented by the
autofluorescence known to originate from the schistosome eggshell[49,50,51], with vitelline cells the
likely origin of this emission[52]. To surmount this hurdle, the EGFP spectrum emitted from within
each eggshell was investigated by selecting the area to be examined, specifically the entire
miracidium, and collecting multispectral images of the miracidium within the eggshell using the
LSM Image Examiner. The images collected were assessed for EGFP by subtracting regions
emitting autofluorescence from the fluorescence signal collected for entire surface area of the
egg: Specifically, the total EGFP intensity at 509 nm[52] was calculated using the Zeiss Zen (black
edition) software module from ~400 eggs (~100 eggs from each of four biological replicates) in
each of the control and experimental groups.

Databases. The sequence reads are available at GenBank BioProject XXX, SRA reads YYY to
ZZZ.

Acknowledgements

Schistosome-infected mice were provided by the NIAID Schistosomiasis Resource Center of Biomedical Research Institute, Rockville, MD through NIH-NIAID contract HHSN272201700014I for distribution through BEI Resources. This work was supported by Wellcome Trust award 107475/Z/15/Z (PI, Karl F Hoffmann). This study is set within the framework of the « Laboratoire d'Excellence (LabEx) » TULIP (ANR-10-LABX-41) with the support of LabEx CeMEB, an ANR « Investments d'avenir » program (ANR- 10-LABX-04-01) and the Environmental Epigenomics Core Service at IHPE.

Author contributions

PB, WI, CG and CGG conceived the study. CC and CG identified gene safe harbors, MFM and TQ contributed promoter/ terminator cloning and analysis, RR optimized conditions for knock-in of donor transgenes, and PW and WB performed mutation analyses. WI planned and performed the programmed gene editing, knockout and transgene knock-in analysis. VM, AM, LM, SS and MM-K maintained the schistosome life cycle. MFM, AP and WI analyzed confocal micrographs and fluorescence. WI, CG and PB drafted the manuscript. All authors contributed to the writting and all approved the final version of the paper.

Competitive interests

The authors declare no competing interests.

 Figure legends

Figure 1. Bioinformatic identification of sites of genomic safe harbors and their CRISPR
programmed targets in the genome of *Schistosoma mansoni*

Panel **a.** Table shows criteria and rationale used to computationally predict genome safe harbors
(GSHs). **b-d.** Chromosomal locations and lengths of four candidate GSH sites (blue boxes),
located on chromosome 2 (GSH2 and GSH3) and chromosome 3 (GSH1, GSH4) of *S. mansoni*.
In addition, coding sequences of genes identified in the vicinity of the four (gene-free) GSHs are
shown, which were mostly uncharacterized proteins (red boxes with connected red line). Black
and white scale bars at the top of each panel each represent one megabase pairs in length. **e.**
Features of the GSHs including chromosomal location, length, programmed CRISPR/Cas9
targets, guide RNAs, and their predicted on- and off-target specificities from the CHOPCHOP
design toolbox.

Figure 2. Efficiency of programmed knockout enhanced using three overlapping guide
RNAs. Panel **a**. Schematic map of sites of the overlapping guide RNAs (blue, red and green
arrows for target 1, -2 and -3, respectively) within GSH1 (yellow box), along with primer
locations for indel analysis (purple arrows). The black arrows indicate the site programmed
double stranded break (DSB) programmed by sgRNA1, 2 and 3. **b**. Efficiency of CRISPR
knockout at GSH1 in eggs of *S. manson*i, as assessed with the Deconvolution of Complex DNA
Repair (DECODR) algorithm using distance, following transfection with dual and triple
overlapping guide RNPs; RNP1+RNP2 (blue dots), RNP2+RNP3 (red), RNP1+RNP3 (green),
and RNP1+RNP2+RNP3 (purple). Significantly higher CRISPR efficiency obtained with the
three overlapping guide RNPs, mean = 23.6%, than the other groups (P<0.001). Among the
groups transfected with dual RNPs, efficiency obtained with the RNP1+RNP3 treatment group,
mean = 12.6%, was significantly higher than either of the other groups, RNP1+RNP2 at 5.4%
and RNP2+RNP3 at 3.6% (P<0.01; one-way ANOVA with 95% confidence intervals, five or six
independent biological replicates; GraphPad Prism). **c-e**. Representative alleles, in the
schistosome genome, bearing indels at the target site in GSH1 (panel a) following transfection
with dual guide RNPs, as a gauge of efficiency CRISPR-catalyzed gene editing. The reference
wild type (WT) allele is shown above the mutant (KO) allele. KO alleles were identified during
examination of nucleotide sequences for small deletions, 1-6 nt in length (- dash) or
insertion/substitution (black boxes) The vertical black line boxes show targeted PAM
(protospacer adjacent motif) trinucleotides. **f**. A representative example of an KO allele bearing a
large sized deletion resulting from transfection with the triple guide RNPs.

Figure 3. Targeted transgene insertion and expression in the schistosome egg. Programmed
CRISPR/Cas9 insertion (knock-in, KI) of a donor repair template of 4.451 kb in length, encoding
an EGFP transgene driven by the *S. mansoni* ubiquitin promoter and terminator. Panel **a.**
Topology of double-stranded DNA donor prepared from a primer pair with 5´ 5x-
phosphorothioate modification. The donor template encoded the *S. mansoni* ubiquitin promoter
(pink bar) driving expression of the EGFP reporter gene (green) and ubiquitin terminator (pink)
and was flanked at its termini with symmetrical 600 bp homology arms (black bars). The
homology arm on the left (HAL) was situated 600 bp upstream of the position of sgRNA1 and
the homology arm on the right (HAR) was sited 600 bp of downstream of the prototypic adjacent
motif of sgRNA 3. **b.** Schematic illustration of the wild type (WT) and mutant (KI) alleles after

double stranded breaks programmed by the overlapping sgRNAs 1, 2 and 3 (scissors). PCR
primer sites are indicated by the purple arrows. **c.** Targeted KI of the transgene detected by
genomic PCR analysis using 5′KI (728 bp) or 3′KI (983 bp) specific pairs of primers. Negative
controls for KI included WT, mock transfected, donor only treatment or KO (only RNPs
transfection) groups, none of which had been transfected with RNPs. Lanes KI-1, KI-2 and KI-3
show amplicons from three independent biological replicates of programmed KI of the donor
transgene. Other lanes show the outcome of RT-qPCRs from schistosome RNA with donor DNA
electroporation (without CRISPR materials - nuclease or guide RNAs). The integrity of the
genomic DNAs was confirmed by the presence of the expected amplicon of 784 bp in all lanes of
the gel labeled "control PCR" using control primer (Fig. 3b).

Primer-dimer and/or non-specific PCR band(s) from DNA donor transfected-eggs were seen ≤
100 bp in size. **d.** Expression of the EGFP transcript (expected amplicon size, 717 bp) as
assessed by RT-qPCR following programmed KI of the transgene into GSH1. The integrity of
the RNAs was assessed by analysis of transcripts of the reference gene, *Sm*GAPDH, with an
expected amplicon size of 285 bp. Double-stranded DNA donor was used as the positive PCR
template for EGFP. Transcription of GAPDH was seen in all treatment and control groups (lanes
1-3 and KI-1 to KI-3 in bottom panel) but not in the donor group.

Figure 4. Markedly higher numbers of EGFP-positive eggs following programmed knock-
in at GSH1 as assessed by spectral image analysis. Confocal laser scanning micrographs:
Panel **a**, eggs exhibiting background signal (autofluorescence) from the control group, *i.e.* eggs
transfected only with the lsDNA donor repair template; **a1, a2,** representative images from
biological replicates. **b.** Eggs expressing EGFP in the experimental group transfected with RNPs
and the donor repair template; **b1, b2**, representative images from two biological replicates.
Many eggs expressed EGFP with the broad range in intensity of fluorescence ranging from
higher intensity (green arrow) and lower levels (yellow arrow) following programmed KI. Eggs
expressed EGFP observed from day 5 until day 15 when the experiment was terminated and
taken the micrographs. **c.** Linear mixing analysis revealed EGFP fluorescence in miracidia within
the eggshell in these representative micrographs; **c1**, plot showing the emission spectrum of
EGFP (green curve) with a peak signal at 509 nm, and, also showing the autofluorescence
background (purple curve). **d.** To assess expression of EGFP, eggs were counted (scored) as
EGFP-positive when >20% of the surface area of the miracidium (within the eggshell) was
EGFP positive; representative EGFP-positive eggs are shown in panels c3 and c4 (green eggs).
On day 15 following transfection, eggs were scored and counted for EGFP positivity in each of
four independent biological replicates (~100 eggs from each replicate) of KI (RNPs + lsDNA)
and lsDNA only eggs. About 25% of the eggs in the lsDNA only control group were scored as
EGFP-positive (range, 19 to 32%), whereas 75% of eggs the KI group expressed EGFP (range,
68 to 79%); P <0.001; two-tailed t = 69.87, df = 142; difference between means (EGFP-KI –
only donor) ± SEM, 49.7 ± 0.7, 95% CI, 48.3 to 51.1. There was marked decrease of
background autofluorescence in eggs from WT and mock-transfection groups after the linear
unmixing normalization of the fluorescence spectrum.

Supporting information

Figure S1. a. CRISPR efficiency (percentage) on individual target sites as estimated using the
Deconvolution of Complex DNA Repair (DECODR) algorithm using distance from dual

gRNAs; RNP1+RNP2 (blue dots), RNP2+RNP3 (green), RNP1+RNP3 (yellow) and triple
gRNAs; RNP1+RNP2+RNP3 (pink dots), plotted from 5-6 independent biological replicates.
The % indels on each target varied among RNP combinations:| transfection with triple RNPs
lead to highest CRISPR efficiencies on each target site 1, 2 and 3, 26.5%, 14.2% and 31.2%,
respectively, compared with the dual RNP groups. The % indels resulting by dual RNPs on
target 1 were 6.7% and 13.8%; target 2, 3.9% and 1.7%, and target 3, 5.4% and 11.3%. **b-e,**
examples of indel patterns resulting from the dual RNPs and triple RNPs. The graphs plotted by
DECODR software confirmed the indel profiles similar to those obtained with TIDE
(supplement figures 1b-1e).

Figure S2. Emission spectral intensity for eggs, scanned from 477 to 638 nm, normalized
fluorescence spectral intensity from control eggs (transfected with donor repair template)
exhibiting higher intensity than autofluorescence; these eggs were also scored as EGFP-positive,
and with a normalized EGFP intensity mean, 1290 au (range, 856 - 1712.8); experimental group,
normalized-EGFP intensity, mean 6905 au (range 4971.5 – 8963.1); $P < 0.001$, unpaired t-test, n
= 402; difference between means of experimental and control group eggs ± SEM, 5651 ± 57.40,
95% CI, 5502 to 5728).

602
603   **Table S1.** Primer list used in PCRs to amplify DNA donor, investigate EGFP transcript,
604   programmed knockout and knockin.

References

1   Zhu, H., Li, C. & Gao, C. Applications of CRISPR-Cas in agriculture and plant
biotechnology. *Nat Rev Mol Cell Biol* **21**, 661-677 (2020).
https://doi.org:10.1038/s41580-020-00288-9
2   Zaib, S., Saleem, M. A. & Khan, I. CRISPR-Cas9 Genome Engineering: Trends in
Medicine and Health. *Mini Rev Med Chem* **22**, 410-421 (2022).
https://doi.org:10.2174/1389557521666210913112030
3   Wang, F. & Qi, L. S. Applications of CRISPR Genome Engineering in Cell Biology.
Trends Cell Biol **26**, 875-888 (2016). https://doi.org:10.1016/j.tcb.2016.08.004
4   Hotez, P. J. *et al.* Helminth infections: the great neglected tropical diseases. *J Clin Invest*
118, 1311-1321 (2008). https://doi.org:10.1172/JCI34261
5   Naldini, L. *et al.* In vivo gene delivery and stable transduction of nondividing cells by a
lentiviral vector. *Science* **272**, 263-267 (1996).
https://doi.org:10.1126/science.272.5259.263
6   Li, X. *et al.* piggyBac transposase tools for genome engineering. *Proc Natl Acad Sci U S*
A **110**, E2279-2287 (2013). https://doi.org:10.1073/pnas.1305987110
7   Guo, J. C. *et al.* Highly Efficient CRISPR/Cas9-Mediated Homologous Recombination
Promotes the Rapid Generation of Bacterial Artificial Chromosomes of Pseudorabies
Virus. *Front Microbiol* **7**, 2110 (2016). https://doi.org:10.3389/fmicb.2016.02110
8   Ittiprasert, W. *et al.* Programmed genome editing of the omega-1 ribonuclease of the
blood fluke, Schistosoma mansoni. *Elife* **8:e41337** (2019).
https://doi.org:10.7554/eLife.41337
9   Ittiprasert, W. *et al.* RNA-Guided AsCas12a- and SpCas9-Catalyzed Knockout and
Homology Directed Repair of the Omega-1 Locus of the Human Blood Fluke,
Schistosoma mansoni. *Int J Mol Sci* **23** (2022). https://doi.org:10.3390/ijms23020631
10   Jang, D. E. *et al.* Multiple sgRNAs with overlapping sequences enhance CRISPR/Cas9-
mediated knock-in efficiency. *Exp Mol Med* **50**, 1-9 (2018).
https://doi.org:10.1038/s12276-018-0037-x
11   Acosta, S., Fiore, L., Carota, I. A. & Oliver, G. Use of two gRNAs for CRISPR/Cas9
improves bi-allelic homologous recombination efficiency in mouse embryonic stem cells.
Genesis **56**, e23212 (2018). https://doi.org:10.1002/dvg.23212
12   Gutierrez-Triana, J. A. *et al.* Efficient single-copy HDR by 5' modified long dsDNA
donors. *Elife* **7** (2018). https://doi.org:10.7554/eLife.39468
13   Aznauryan, E. *et al.* Discovery and validation of human genomic safe harbor sites for
gene and cell therapies. *Cell Rep Methods* **2**, 100154 (2022).
https://doi.org:10.1016/j.crmeth.2021.100154
14   Berriman, M. *et al.* The genome of the blood fluke Schistosoma mansoni. *Nature* **460**,
352-358 (2009). https://doi.org:10.1038/nature08160
15   Protasio, A. V. *et al.* A systematically improved high quality genome and transcriptome
of the human blood fluke Schistosoma mansoni. *PLoS Negl Trop Dis* **6**, e1455 (2012).
https://doi.org:10.1371/journal.pntd.0001455
16   Howe, K. L. *et al.* WormBase 2016: expanding to enable helminth genomic research.
Nucleic Acids Res **44**, D774-780 (2016). https://doi.org:10.1093/nar/gkv1217
17   Howe, K. L., Bolt, B. J., Shafie, M., Kersey, P. & Berriman, M. WormBase ParaSite - a
comprehensive resource for helminth genomics. *Mol Biochem Parasitol* **215**, 2-10
(2017). https://doi.org:10.1016/j.molbiopara.2016.11.005

18 Suttiprapa, S., Rinaldi, G. & Brindley, P. Genetic manipulation of schistosomes -
Progress with integration competent vectors. *Parasitology* **139**, 641-650 (2011).
https://doi.org:10.1017/S003118201100134X
19 Augusto, R. C. *et al.* A simple ATAC-seq protocol for population epigenetics. *Wellcome*
Open Res **5**, 121 (2020). https://doi.org:10.12688/wellcomeopenres.15552.2
20 Luo, H. *et al.* DEG 15, an update of the Database of Essential Genes that includes built-in
analysis tools. *Nucleic Acids Res* **49**, D677-D686 (2021).
https://doi.org:10.1093/nar/gkaa917
21 Labun, K. *et al.* CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome
editing. *Nucleic Acids Res* **47**, W171-W174 (2019). https://doi.org:10.1093/nar/gkz365
22 Montague, T. G., Cruz, J. M., Gagnon, J. A., Church, G. M. & Valen, E. CHOPCHOP: a
CRISPR/Cas9 and TALEN web tool for genome editing. *Nucleic Acids Res* **42**, W401-
407 (2014). https://doi.org:10.1093/nar/gku410
23 Bloh, K. *et al.* Deconvolution of Complex DNA Repair (DECODR): Establishing a
Novel Deconvolution Algorithm for Comprehensive Analysis of CRISPR-Edited Sanger
Sequencing Data. *CRISPR J* **4**, 120-131 (2021). https://doi.org:10.1089/crispr.2020.0022
24 Marcek Chorvatova, A., Kirchnerova, J., Cagalinec, M., Mateasik, A. & Chorvat, D., Jr.
Spectrally and spatially resolved laser-induced photobleaching of endogenous flavin
fluorescence in cardiac myocytes. *Cytometry A* **95**, 13-23 (2019).
https://doi.org:10.1002/cyto.a.23591
25 Abudayyak, M., Jannuzzi, A. T., Ozhan, G. & Alpertunga, B. Investigation on the toxic
potential of Tribulus terrestris in vitro. *Pharm Biol* **53**, 469-476 (2015).
https://doi.org:10.3109/13880209.2014.924019
26 Aoshima, T. *et al.* Modification of improved-genome editing via oviductal nucleic acids
delivery (i-GONAD)-mediated knock-in in rats. *BMC Biotechnol* **21**, 63 (2021).
https://doi.org:10.1186/s12896-021-00723-5
27 Dickinson, M. E., Bearman, G., Tille, S., Lansford, R. & Fraser, S. E. Multi-spectral
imaging and linear unmixing add a whole new dimension to laser scanning fluorescence
microscopy. *Biotechniques* **31**, 1272, 1274-1276, 1278 (2001).
https://doi.org:10.2144/01316bt01
28 Lee, H., Kim, J., Kim, H. H., Kim, C. S. & Kim, J. Review on Optical Imaging
Techniques for Multispectral Analysis of Nanomaterials. *Nanotheranostics* **6**, 50-61
(2022). https://doi.org:10.7150/ntno.63222
29 Newby, G. A. *et al.* Base editing of haematopoietic stem cells rescues sickle cell disease
in mice. *Nature* **595**, 295-302 (2021). https://doi.org:10.1038/s41586-021-03609-w
30 Pavani, G. & Amendola, M. Targeted Gene Delivery: Where to Land. *Front Genome Ed*
2, 609650 (2021). https://doi.org:10.3389/fgeed.2020.609650
31 Diez, B. *et al.* Therapeutic gene editing in CD34(+) hematopoietic progenitors from
Fanconi anemia patients. *EMBO Mol Med* **9**, 1574-1588 (2017).
https://doi.org:10.15252/emmm.201707540
32 Hoffmann, K. F., Brindley, P. J. & Berriman, M. Medicine. Halting harmful helminths.
Science **346**, 168-169 (2014). https://doi.org:10.1126/science.1261139
33 Cox, D. B., Platt, R. J. & Zhang, F. Therapeutic genome editing: prospects and
challenges. *Nat Med* **21**, 121-131 (2015). https://doi.org:10.1038/nm.3793
34 Douglas, B. *et al.* Immune System Investigation Using Parasitic Helminths. *Annu Rev*
Immunol **39**, 639-665 (2021). https://doi.org:10.1146/annurev-immunol-093019-122827

35   Zhang, T., Cooper, S. & Brockdorff, N. The interplay of histone modifications - writers
that read. *EMBO Rep* **16**, 1467-1481 (2015). https://doi.org:10.15252/embr.201540945
36   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J.
Transposition of native chromatin for fast and sensitive epigenomic profiling of open
chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* **10**, 1213-1218
(2013). https://doi.org:10.1038/nmeth.2688
37   Schroder, A. R. *et al.* HIV-1 integration in the human genome favors active genes and
local hotspots. *Cell* **110**, 521-529 (2002). https://doi.org:10.1016/s0092-8674(02)00864-4
38   Sadelain, M., Papapetrou, E. P. & Bushman, F. D. Safe harbours for the integration of
new DNA in the human genome. *Nat Rev Cancer* **12**, 51-58 (2011).
https://doi.org:10.1038/nrc3179
39   Lu, Z. & Berriman, M. Meta-analysis of RNA-seq studies reveals genes responsible for
life stage-dominant functions in Schistosoma mansoni. *bioRxiv*, 308189
(2018). https://doi.org:10.1101/308189
40   Pavani, G. & Amendola, M. Targeted Gene Delivery: Where to Land. *Front Genome Ed*
2, 609650 (2020). https://doi.org:10.3389/fgeed.2020.609650
41   Pavani, G. & Amendola, M. Corrigendum: Targeted Gene Delivery: Where to Land.
Front Genome Ed **3**, 682171 (2021). https://doi.org:10.3389/fgeed.2021.682171
42   Cortes, M. F. *et al.* Community-acquired methicillin-resistant Staphylococcus aureus
from ST1 lineage harboring a new SCCmec IV subtype (SCCmec IVm) containing the
tetK gene. *Infect Drug Resist* **11**, 2583-2592 (2018).
https://doi.org:10.2147/IDR.S175079
43   Dalton, J. P., Day, S. R., Drew, A. C. & Brindley, P. J. A method for the isolation of
schistosome eggs and miracidia free of contaminating host tissues. *Parasitology* **115 (Pt**
1), 29-32 (1997). https://doi.org:10.1017/s0031182097001091
44   Rinaldi, G. *et al.* Germline transgenesis and insertional mutagenesis in Schistosoma
mansoni mediated by murine leukemia virus. *PLoS Pathog* **8**, e1002820 (2012).
https://doi.org:10.1371/journal.ppat.1002820
45   Labun, K., Montague, T. G., Gagnon, J. A., Thyme, S. B. & Valen, E. CHOPCHOP v2: a
web tool for the next generation of CRISPR genome engineering. *Nucleic Acids Res* **44**,
W272-276 (2016). https://doi.org:10.1093/nar/gkw398
46   Hulme, B. J. *et al.* Schistosoma mansoni alpha-N-acetylgalactosaminidase (SmNAGAL)
regulates coordinated parasite movement and egg production. *PLoS Pathog* **18**, e1009828
(2022). https://doi.org:10.1371/journal.ppat.1009828
47   Wilfinger, W. W., Mackey, K. & Chomczynski, P. Effect of pH and ionic strength on the
spectrophotometric assessment of nucleic acid purity. *Biotechniques* **22**, 474-476, 478-
481 (1997). https://doi.org:10.2144/97223st01
48   Brinkman, E. K. *et al.* Easy quantification of template-directed CRISPR/Cas9 editing.
Nucleic Acids Res **46**, e58 (2018). https://doi.org:10.1093/nar/gky164
49   Domingo, M., Mais, R. F., Weiskopf, R. & Fink, S. Detection of schistosome ova by dark
field fluorescence microscopy. *Gastroenterology* **54**, 884-886 (1968).
50   Lu, Z. *et al.* Isolation, enrichment and primary characterisation of vitelline cells from
Schistosoma mansoni obtained by the organ isolation method. *Int J Parasitol* **45**, 663-672
(2015). https://doi.org:10.1016/j.ijpara.2015.04.002
51   Collins, J. J., 3rd, King, R. S., Cogswell, A., Williams, D. L. & Newmark, P. A. An atlas
for Schistosoma mansoni organs and life-cycle stages using cell type-specific markers

743    and confocal microscopy. *PLoS Negl Trop Dis* **5**, e1009 (2011).
744     https://doi.org:10.1371/journal.pntd.0001009
745  52  Wu, Q., Feng, Z. & Hu, W. Reduction of autofluorescence in whole adult worms of
746    Schistosoma japonicum for immunofluorescence assay. *Parasit Vectors* **14**, 532 (2021).
747     https://doi.org:10.1186/s13071-021-05027-3
748

Figure 1

a

GSH criteria	Rationale
• Close to peaks of H3K4me3 in all developmental stages of the parasite	• a histone modification site that is associated with euchromatin and transcription start sites for highly expression of transgene
• Do not contain H3K27me3 in any developmental stage	• a histone modification that is associated with heterochromatin in any of the life stages to avoid silencing genes to be integrated
• Unique sequences	• Avoid CRISPR/Cas off target
• Euchromatin accessible to Tn5 integration	• Positive display of integration events
• Located outside annotated genes and long non-coding RNA, outside putative promotor regions where *more than 2 kb upstream* of the TTS and within 11 kb of HIV integration site	• Avoid gene disruption during cell differentiation
	• Avoid disrupting essential function of endogenous gene(s)
	• HIV integration sites known from *S. mansoni* genome to additionally support predictions of euchromatic regions

e

S. mansoni GSH genome, version 9	chromosome	location	length (nt)	CRISRP/Cas9 target sequence(s) predicted by CHOPCHOP
GSH1	3	13231955-13233370	1,416	Efficient CRISPR/Cas9, overlapping gRNAs without predicted off-targets: GGAACTCGGTAGATGAAGGC[TGG] ATGAAGGCTGGTTCAATAGC[TGG] GCTGGTTCAATAGCTGGAAT[TGG]
GSH2	2	13399775-13400744	967	One region only contains CRISPR/Caa9 target with PAM, CAGCATTACCTGTGACTTAC[TGG]G[TGG]
GSH3	2	7708870-7709621	752	Modest CRISPR/Cas9 efficiency (<35%) overlapping gRNAs without predicted off-targets : TGTAAAATGGCAACAGTTCT [TGG] ATGGCAACAGTTCTTGGTGT [TGG] ACAGTTCTTGGTGTTGGACC [TGG]
GSH4	3	13233423-13233560	138	Only one CRISPR/Cas9 target was predicted with very low CRISPR efficiency. Several CRISPR/Cas9 targets were predicted but these contain off targets on chromosome 1, 2, 3, 5 and ZW

Figure 2

a 5'-PPPPP
HAL | ubiquitin promoter | EGFP | ubiquitin terminator | HAR
5'-PPPPP
5' phosphorothioate modified donor containing the 4.451 kb transgene

b
sgRNA2 sgRNA3
control-F sgRNA1
wild type
HAL HAR
control-R

knock-in
5' KI-F
5' KI-R
EGFP-F
ubiquitin promoter | EGFP | ubiquitin terminator
Transgene 4.451 kb
EGFP-R
3' KI-F
3' KI-R

c

3' KI-PCR | 5' KI-PCR
M WT mock donor KO K1-1 KI-2 KI-3 | WT mock donor KO KI-1 KI-2 KI-3
← 983 bp
← 728 bp

control PCR
M WT mock donor KO K1-1 KI-2 KI-3
← 784 bp

d
+ve | donor transfection only | | | M | KI-1 KI-2 KI3
1 2 3
← 717 bp eGFP
} primer-dimer/non-specific products
← 285 bp *Sm*GAPDH

Figure 4

Supplement Table 1. Primer list used in PCRs to amplify DNA donor, investigate EGFP transcript, programmed knockout and knockin

Primer	5' to 3' sequence
indel-F	TGTTATCGTCCGTCGCTTCA
indel-R	GCGTTCAAACATTGCCCACT
600 bp HA-F	AGATTGCTAGAAATTTATGAAAG
600 bp HA-R	ACTGCCGAATTTATAATATTTGG
400 bp HA-F	GATGAGTGCATCGATCGATTAC
400 bp HA-R	GAATGATGTTGGAAGTCAGA
200 bp HA-F	CGCTTCAATATTGTTTTTTGC
200 bp HA-R	CACACCTTCAAAACAATGTTTC
EGFP-F	ATGGTGAGCAAGGGCGAGG
EGFP-R	CTTGTACAGCTCGTCCATGCC
SmGAPDH-F	ATGGGACATTTCCAGGCGAG
SmGAPDH-R	CCAACAACGAACATGGGTGC
5' KI-F	AGGGTTTTGGTTCGCAGGAT
5' KI-R	CGGAGACAATCTGGAAAGGTCA
3' KI-F	GTCCGCGTAATCGTCGTTACTA
3' KI-R	GTGGTTCCATACTATGCAGTTTCC