Tyramine and its *Amtyr1* receptor modulate attention in honey bees (*Apis mellifera*)

Joseph S Latshaw, Reece E Mazade, Mary Petersen, Julie A Mustard[1], Irina Sinakevitch[2], Lothar Wissler, Xiaojiao Guo[3,4], Chelsea N. Cook[5], Hong Lei, Jürgen Gadau[6], Brian H Smith[*]

School of Life Sciences, PO Box 874501, Arizona State University, Tempe, AZ 85287

The authors wish to dedicate this work to the memory of Professor Hans-Joachim Pflüger, who was a wonderful friend and a great colleague. He made significant contributions to understanding the roles of tyramine and octopamine in insect behavior.

Abstract: Animals must learn to ignore stimuli that are irrelevant to survival and attend to ones that enhance survival. When a stimulus regularly fails to be associated with an important consequence, subsequent excitatory learning about that stimulus can be delayed, which is a form of non-associative conditioning called 'latent inhibition'. Honey bees show latent inhibition toward an odor they have experienced without association with food reinforcement. Moreover, individual honey bees from the same colony differ in the degree to which they show latent inhibition, and these individual differences have a genetic basis. To investigate the mechanisms that underly individual differences in latent inhibition, we selected two honey bee lines for high and low latent inhibition, respectively. We crossed those lines and mapped a Quantitative Trait Locus (QTL) for latent inhibition to a region of the genome that contains the tyramine receptor gene *Amtyr1*[1]. We then show that disruption of *Amtyr1* signaling either pharmacologically or through RNAi qualitatively changes the expression of latent inhibition but has little effect on appetitive conditioning, and these results suggest that AmTYR1 modulates inhibitory processing in the CNS. Electrophysiological recordings from the brain during pharmacological blockade are consistent with a model that AmTYR1 indirectly regulates Hebbian plasticity at inhibitory synapses in the CNS. Finally, sequencing *Amtyr1* and its up and downstream genomic region for high and low line workers suggested that individual differences might arise from variation in transcriptional control rather than structural changes in the coding region. Our results therefore identify a distinct modulatory pathway for this type of non-associative learning, and we propose a model for how it modulates Hebbian plasticity at defined synapses in the CNS. We have shown elsewhere how this modulation also underlies potentially adaptive intracolonial learning differences among individuals that benefit colony survival.

Introduction

The ability to learn predictive associations between stimuli and important events, such as food or threats, is ubiquitous among animals (*1*), and it may underlie more complex

---

[1] We use *Amtyr1* to denote the gene and AmTYR1 the receptor throughout the text.

cognitive capabilities (*1, 2*). This ability arises from various forms of associative and oper-
ant conditioning(*3*). However, the absence of reward also provides important information
for learning about stimuli, because all animals must use this information to redirect a lim-
ited attention capacity to more important stimuli (*4*). One important mechanism for learning
to ignore irrelevant stimuli is called latent inhibition (*5*). After an animal is presented with
a stimulus several times without reinforcement, learning is delayed or slower when that
same stimulus is reinforced in a way that would normally produce robust excitatory con-
ditioning. For example, when honey bees are repeatedly exposed to a floral odor without
association to food rewards, their ability to subsequently learn an excitatory association
of this odor with a reward is delayed or reduced (*6*). While many studies in the honey bee
have focused around how the presence of reward shapes learning and memory (*7*), eval-
uating this important form of nonassociative learning has not received as much attention
(*6, 8*). Yet, like in all animals, it plays an important ecological role in the learning repertoire
of honey bees. The presence of unrewarding flowers in an otherwise productive patch of
flowers (*9*), or the unreinforced presence of an odor in the colony (*10*), can influence for-
agers' choices of flowers during foraging trips.

Moreover, individual honey bees from the same colony differ in the degree to which
they exhibit several learning traits (*11-15*), including latent inhibition(*12*). Several studies
of different forms of learning have demonstrated that individual differences are heritable
(*12*). Individuals showing different learning phenotypes occur within the same colony be-
cause a queen mates with up to 20 drones (males)(*16*), and thus honey bee colonies
typically contain a mixture of many different paternal genotypes. This within-colony ge-
netic diversity of learning capacities may reflect a colony level trait that allows the colony
to react and adapt to rapidly changing resource distributions (*17, 18*).

Our objective here was to evaluate the genetic and neural mechanisms that underlie
individual differences for latent inhibition in honey bees. We show that a major locus sup-
porting individual differences maps to a location in the honey bee genome previously iden-
tified in independent mapping studies as being important for latent inhibition(*19*) as well
as for sugar and pollen preferences in foragers (*20, 21*). Disruption of a tyramine receptor
encoded by *Amtyr1* in this region changes expression of latent inhibition in a way that
suggests that intact signaling via the *Amtyr1* pathway is important for modulating plasticity

at inhibitory synapses. Furthermore, electrophysiological analyses combined with block-
ade of the AmTYR1 receptor in the antennal lobe – the first synaptic center along the
olfactory pathway – decreased antennal lobe responsiveness to odor and blocked a neu-
ral correlate of latent inhibition. Finally, sequencing the gene failed to reveal mutations in
the coding regions that would affect protein function, leading to the conclusion that varia-
tion across workers could arise from differential gene expression through transcriptional
regulation. We discuss how these data strongly imply a functional role for *Amtyr1* signaling
in modulating expression of attention via latent inhibition. These findings are also im-
portant for understanding the strategies colonies use to explore for and exploit pollen and
nectar resources(*22*).

Results

We used two genetic lines of honey bees that had been bred for high (inhibitor) or low
(noninhibitor) expression of latent inhibition. These lines were independently selected us-
ing identical methods to a previous study that had successfully bred high and low
lines(*19*). We evaluated 523 recombinant drones generated from a single hybrid queen
produced from a cross between a drone from a noninhibitor line and a queen from an
inhibitor line (Fig. 1). Honey bee drones are ideal for behavior genetic studies because
they are haploid progeny that develop from an unfertilized egg laid by the queen. We then
selected 94 high and 94 low performing drones for the Quantitative Trait Locus (QTL)
analysis, which identified one significant locus (Fig. 2A). The QTL mapped to the same
genomic region identified in a previous study of latent inhibition (called '*lrn1*') using an
independent inhibitor and noninhibitor cross and different (RAPD-based) genetic markers
(*19*). This is the same genomic region that has been identified in studies of foraging pref-
erences of honey bees (*21, 23*), where it has been called *pln2* for its effect on pollen
versus nectar preferences and in modulating sensitivity to sucrose (*24*). Clearly this ge-
nomic region has major effects on several foraging-related behaviors.

Fig. 1. Ev[...] om an F1 queen. **(A)** Drone honey bees were first evaluated [...] als to an odor (A; gray bars) followed by sucrose reinforcem[...] ces robust associative conditioning expressed as odor-indu[...] ER) (*25*). All drones that showed no PER response on the firs[...] he following two trials were selected for the subsequent fam[...] nsured that only drones motivated to respond to sucrose a[...] re selected. Approximately 10% of honey bees fail to show evidence of learning in PER conditioning using the collection methods described in Materials and Methods. The familiarization phase involved 40 4-sec exposures to a different odor (X; black bars) using a 5 min interstimulus interval. These conditions are sufficient for generating latent inhibition that lasts for at least 24 h (*6*). Finally, the test phase involved 6 exposures to X followed by sucrose reinforcement. **(B)** Frequency distribution of 523 drones evaluated in the test phase. The x-axis shows the summed number of responses over 6 conditioning trials. Fewer responses correspond to stronger latent inhibition. A total of 94 drones were selected in each tail of the distribution. 'Inhibitor' drones showed zero through 3 responses, and 'Noninhibitor' drones showed 5 or 6 responses. **(C)** Acquisition curves for the 94 inhibitor and noninhibitor drones. Approximately half of the noninhibitor drones showed spontaneous responses on the first trial, which is typical for noninhibitors in latent inhibition studies of honey bees (*6*). All of the drones in that category showed responses on trials 2 through 6. In contrast, inhibitor drones showed delayed acquisition to the now familiar odor.

When we analyzed the gene list within the confidence intervals of this QTL, one gene - *Amtyr1* - in particular stood out (Fig. 2A,B). That gene encodes a biogenic amine

receptor for tyramine (AmTYR1) (*26*) that is expressed in several regions of the honey

bee brain (*27-29*). AmTYR1 is most closely related to the insect alpha-2-adrenergic-like

octopamine receptors and the vertebrate alpha-2-adrenergic receptors (*30*). Activation

of AmTYR1 reduces cAMP levels in neurons that express it. We specifically considered

AmTYR1 for more detailed evaluation for several reasons. Tyramine affects sucrose

sensitivity in honey bees(*31*), and nurses and foragers differ in AmTYR1 expression(*32*).

Mutations in the orthologous tyramine receptor in fruit flies disrupt odor-guided innate

behaviors to repellants (*33*). Tyramine is also the direct biosynthetic precursor to octopa-

mine(*34*), which has been widely implicated in sucrose-driven appetitive reinforcement

learning in the honey bee (*35, 36*). Therefore, Ventral Unpaired Medial neurons, which

lie on the median of the subesophageal ganglion in the honey bee brain (*28, 37-39*), and

which form the basis for the appetitive reinforcement pathway must produce tyramine in

the process of making octopamine. Recent analyses indicate these neurons in locusts

and fruit flies also release both neuromodulators when activated (*40, 41*). Finally, octo-

pamine and tyramine affect locomotor activity in the honey bee(*42*).

Fig. 2. Single-Nucleotide Polymorphism (SNP) mapping of high and low recombinant drones. (A) Markers from linkage group 1.55 surrounding one significant QTL (est941 with a LOD score of 2.6). **(B)** Partial list of genes within 10 cM of the marker showing the location of *Amtyr1*.

We then evaluated whether non-synonymous mutations in the coding sequence might change the functionality of the receptor. We performed a detailed genomic analysis of the 40kb region including the *Amtyr1* gene, a 2kb upstream, and a 0.5kb downstream non-coding region. SNP frequency in the coding sequence (CDS) was relatively low compared to the genome wide SNP frequency, and all 46 SNPs in the coding regions in any of the sequenced eight individual worker genomes represented synonymous substitutions, i.e., these SNPs do not change the sequence of the encoded protein. Thus, phenotypic differences are not caused by structural changes in the tyramine receptor protein itself. We did, however, find an increased SNP frequency in introns, the up- and downstream non-coding regions and the 3' untranslated region (UTR). If *Amtyr1* is involved in latent inhibition, these variations might be linked to the changes in the regulation of *Amtyr1* gene expression, e.g., by changes in transcription factor binding sites or the stability of the mRNA, which might eventually be responsible for the observed phenotypic differences.

Disruption of *Amtyr1* affects expression of latent inhibition

To further examine the role of *Amtyr1* signaling in latent inhibition, we performed a series of behavioral experiments that involved treatment of honey bees either with the tyramine receptor antagonist yohimbine (*43*) or with a Dicer-substrate small interfering (Dsi) RNA of the receptor (NCBI Reference Sequence: NM_001011594.1) to disrupt translation of mRNA into AmTYR1 (*28, 44*). For these experiments we used unselected worker honey bees from the same background population used for selection studies, which ensured that workers used for behavioral assays would represent a mixture of inhibitor and non-inhibitor phenotypes. Therefore, treatment could increase or decrease the mean level of latent inhibition in this population. Training involved two phases (Fig. 3A). First, during the 'familiarization' phase honey bees were identically exposed over 40 trials to odor X without reinforcement. Our previous studies have shown that this procedure produces robust latent inhibition. The second 'test' phase involved measurement of latent inhibition. During this phase odor X and a 'novel' odor N were presented on separate trials. Both odors were associated with sucrose reinforcement in a way that produces robust

173 appetitive conditioning (*25*). Latent inhibition would be evident if responses to odor X were
174 lower than the responses to the novel odor N. Injections of yohimbine directly into brains
175 occurred either prior to the familiarization phase (Fig. 3A,B) or prior to the test phase (Fig.
176 3C).

177
Fig. 3. Blockade of the tyramine receptor with yohimbine modulated expression of latent
inhibition. (A) Acquisition during the test phase in two injection groups of honey bees familiarized
to air as a control procedure to evaluate the effects of yohimbine on excitatory conditioning. The
conditioning protocol is shown at the top. In this experiment (and in Fig 3**B** and **C**) we omitted the
first phase (Fig. 1**A**), which does not affect expression of latent inhibition(*6*) and is only necessary
when subjects are being selected for development of genetic lines. One group was injected (arrow)
with saline (orange circles; n = 37 animals) and the other with yohimbine (blue triangles; n=35)
prior to familiarization. Because there was no odor presented during familiarization (open box),
odors during the test phase were both 'novel' when conditioned. The test phase in this experiment
(also in Fig 3**B** and **C**) differed from the test phase in Figure 1. For this design, each subject was
equivalently conditioned to both odors on separate, pseudorandomly interspersed trials. Acquisi-
tion to both odors in both injection groups was evident as a significant effect of trial (X^2=47.5, df=3,

p<0.001). None of the remaining effects (odor, injection or any of the interaction terms) were sig-
nificant (p>0.05). **(B)** As in **A**, except both groups (orange saline: n=36; blue yohimbine: n=36) in
this experiment were familiarized to odor; each odor (gray and black boxes; see Methods) was
familiarized in approximately half of the animals in each injection group. In this design, each indi-
vidual was equivalently conditioned to both odors during the test phase; latent inhibition is evident
when the response to the novel odor is greater than to the familiar odor. Injection was prior to odor
familiarization. **(C)** As in **B**, except injection of saline (n=32) or yohimbine (n=30) occurred prior to
the test phase. Statistical analysis of data sets in **B** and **C** yielded a significant interaction (X^2=7.4,
df=1, p <0.01) between injection (saline vs yohimbine) and odor (novel vs familiar) that was the
same in both experiments, as judged by the lack of a significant odor*injection*experiment inter-
action term (p>0.05). There was a higher response to the novel odor than to the familiar odor, but
only in the saline injected groups. The lower rate of acquisition in **C** (X^2=64.0,1,p<0.01) could be
due to performance of this experiment at a different time of year, or to injections immediately prior
to testing, which affects levels PER conditioning in honey bees but leaves intact relative differ-
ences between groups (*45*).

The first experiment provided an important control procedure to evaluate whether yo-
himbine affects excitatory conditioning. This procedure involved familiarization to air,
which does not induce latent inhibition to odor (*6*). Honey bees familiarized to air learned
the association of both odors with sucrose reinforcement equally well (Fig. 3A). The re-
sponse to each odor significantly increased, as expected, across trials (X^2=47.5, df=3,
p<0.001). Moreover, there was no effect of injection with saline versus yohimbine; the re-
sponse levels to all four odors across the saline and yohimbine injection groups were
equivalent. Therefore, blockade of tyramine signaling does not affect excitatory condition-
ing, which is an important control for the effects about to be described. This control pro-
cedure also shows that yohimbine at 10^{-4} M probably does not affect receptors for other
biogenic amines, such as octopamine, dopamine and serotonin, all of which have been
shown to have specific effects on appetitive olfactory learning in honey bees (*35, 46-49*).

Yohimbine treatment affected the expression of latent inhibition in both treatments that
involved familiarization to odor (the interaction between novel vs familiar odor and saline vs
yohimbine injection: X^2=7.4, df=1, p <0.01). First, in the saline controls, honey bees re-
sponded more often to odor N than to X after injection of saline prior to familiarization or

prior to testing (Fig. 3B,C circles). The response to the familiar odor was lower than the
response to the novel odor on most trials, including spontaneous responses on the first
trial. Injection of yohimbine eliminated the difference in response to the novel and familiar
odors. Moreover, the responses to both odors after yohimbine treatment were significantly
lower than, or at least equal to, the response to the familiar odor in the respective saline
controls. This pattern could not arise from blockade of excitatory learning about N, be-
cause excitatory learning was unaffected in the air preexposure controls (Fig 3A). Instead,
the yohimbine-induced pattern was specific to the treatments in which one odor was fa-
miliar.

This result implies that blockade of AmTYR1 modulates latent inhibition to a familiar
odor and that the effect now generalizes to the novel odor. Finally, the relative effect of
yohimbine treatment, i.e. reduction of PER response rate, is similar when it is injected
either prior to familiarization (Fig. 3B) or prior to testing (Fig. 3C). This pattern, that is, the
same effect prior to acquisition or testing, is similar to the action of octopamine blockade
on excitatory conditioning (*35*).

Although the results with yohimbine were promising, we were concerned that yohim-
bine can have effects on other receptors, specifically on an α2-adrenergic-like octopamine
receptor (*30*) and on an excitatory tyramine receptor AmTYR2*(43).* Therefore, we decided
to disrupt *Amtyr1* expression via injection of *Amtyr1* DsiRNA in order to provide an inde-
pendent method to test the role of AmTYR1 in producing latent inhibition (Fig. 4). Yohim-
bine blocks the receptor, whereas dsiAmtyr1 disrupts production of the receptor protein.
Similar outcomes with the two different methods would increase confidence in the result.
For the behavioral experiments we used the same procedure as above for yohimbine
except that the mixture of three *Amtyr1* DsiRNA constructs was injected 20 hours prior to
conditioning because of the time frame needed for the DsiRNA to target mRNA. Because
of that time frame, and because injection of yohimbine prior to either phase produced
equivalent results, we performed injections of *Amtyr1* DsiRNA only prior to familiarization.
As a control we used a scrambled sequence of *Amtyr1* (DsiScr). Use of DsiScr controls
for possible nonspecific effects arising from any aspect of the injection.

Fig. 4 Disruption of translation of the tyramine receptor by DsiRNA also modulated expression of latent inhibition. This experiment was identical to that shown in Fig 3A,B, except injections were performed with a mixture of Dsi *Amtyr1*RNA (dsRNA; arrow) 24 hour prior to behavioral training and testing. The control for this experiment was a scrambled sequence of the *Amtyr1* RNA, Dsiscr (scr). **(A)** After treatment with Dsiscr (N=17) or DsiRNA (N-19) and familiarization to air, acquisition to both odors was significant across trials (X^2=62.7,7,p<<0.01). There was also a significant effect of injection (X^2=8.8, 1, p<0.01). However, the odor*injection interaction was not significant. **(B)** Same as in A, except familiarization was to odor (Dsiscr (N=17) DsiRNA (N=13)). The injection*odor interaction was significant (X^2=7.8; 1; p<0.01. Quantitative PCR analysis of *Amtyr1* mRNA levels in brains revealed lower levels of mRNA in Dsi*Amtyr1* injected animals (0.046+/-0.006) than in Dsiscr injected animals (0.142+/0.028).

Injection of *Amtyr1* DsiRNA produced the same effects as yohimbine. After familiarization to air as a control, both groups of foragers learned the association of both odors (X^2=62.7, 7, p<<0.01; Fig. 4A), although there was a slight decrement in response rate in DsiRNA injected animals (X^2=62.7, 7, p<<0.01; see discussion below). In contrast, after familiarization to one of the odors, learning of both the novel and familiar odors was poor in the *Amtyr1* DsiRNA injected group (Fig. 4B). But expression of latent inhibition was normal – i.e. responses to the novel odor exceeded the responses to the familiar odor in the DsiScr group. As before the interaction between odor and injection was significant (X^2=7.8, 1, p<0.01).

In conclusion, both behavioral experiments support the hypothesis that *Amtyr1* affects expression of latent inhibition without affecting excitatory conditioning. However, the results at first glance seemed counterintuitive. Blockade and disruption of *Amtyr1* did not

attenuate latent inhibition by, for example, increasing the responsiveness to the familiar
odor. Instead, treatment with yohimbine or *Amtyr1* DsiRNA reduced responsiveness to
the novel odor. This result is consistent with *Amtyr1* modulating inhibition involved in, for
example, identified inhibitory processes in the antennal lobes and or the CNS(*50*). Spe-
cifically, and as we propose below in a model, it would prevent the inhibition from becom-
ing too strong, and possibly keep it at a set point between very strong and very weak.

Disruption of *Amtyr1* signaling affects neural codes for odors in the antennal lobe

Because of this intriguing result, we performed additional experiments to investigate
the mechanism in more detail. Our prior studies of odor coding identified neural manifes-
tations of latent inhibition in early synaptic processing of the antennal lobes of the honey
bee brain(*51, 52*). Familiarization to an odor X caused a mixture of a novel odor N and X
to become much more like N(*52*). That is, neural information about familiar odors like X is
filtered out of mixtures. Furthermore, responses to any novel odor are enhanced after
familiarization to X(*51*), which is a form of novelty detection. These effects in the antennal
lobes could arise because of expression of AmTYR1 in presynaptic terminals of sensory
axons in the honey bee antennal lobes(*28*), where activation of AmTYR1 would decrease
cAMP levels(*26*) and likely decrease release of acetylcholine at synapses.

We therefore chose to analyze the effect of yohimbine treatment on odor processing in
the antennal lobes by recording electrophysiological responses to odors prior to and after
familiarization in combination with yohimbine treatment. We used yohimbine in these ex-
periments because of the more rapid onset (minutes versus hours) compared to DsiRNA
treatment. This first experiment did not employ familiarization to odor. Prior to yohimbine
treatment, recordings from 71 units across 4 animals revealed responses to odors that
ranged from no detectable change in spike activity with odor presentation to a robust in-
crease in spiking activity (Fig.5A). After yohimbine treatment, responses decreased pre-
cipitously, although spiking activity was still detectable (Fig. 5B). This decrease in re-
sponse is consistent with AmTYR1 being involved in regulation of inhibition in networks
of the antennal lobe, which are the first order processing centers for odors. Hypothetically
at least, when AmTYR1 is blocked by yohimbine, excitation of inhibitory networks in the
antennal lobe increases and drives down PN responses.

Fig. 5 Yohimbine disrupts processing in the honeybee antennal lobe. **A-B**. Perfusion of yohimbine solution (50μM in physiological saline) into honey bee head capsule caused antennal lobe units to decrease response magnitude to odor stimuli (2-octanone and 1-hexanol) in general. Odor was delivered through a solenoid valve that was open at time zero and lasted for 4 secs. **C**. In control experiments where yohimbine was not applied, most of the units were responsive to both hexanol and octanone, but 39% were biased towards octanone (purple dots above the diagonal line), i.e. showing stronger response to octanone than to hexanol. During the familiarization protocol, these units were familiarized to hexanol 40 times with 1 min interval (arrow down) and were tested again with hexanol and octanone 10 min after the last odor stimulation in the familiarization phase. The test results show 54% of units responded more strongly to octanone (orange dots), which is a novel odor in this protocol. The 15% increase is statistically significant (*McNemar* test with Yates's correction, d.f.=1, Chi-square=5.939, p<0.02) (asterisks on purple and orange bars, N=99). **D**. When the familiarization protocol was used with saline versus yohimbine perfusion, the response bias towards novel odor was disrupted, showing a significant decrease in comparison with the familiar odor (*McNemar* test with Yates's correction, d.f.=1, Chi-square=11.13, p<0.001) (asterisks on grey bars, N=56).

We then evaluated whether continuous perfusion of the brain with saline or yohimbine *during odor familiarization* would interrupt how latent inhibition is manifested in the antennal lobe by potentiation of responses to novel odors, as we have reported(*51*). Indeed, yohimbine treatment modified how neurons respond to novelty. Using the same familiarization protocol as in Figs 3 and 4, but under conditions of saline perfusion, we found that 39% of units (N=99) responded more strongly to the Novel odor before the familiarization to an odor (Fig.5C, purple dots in upper panel; purple bar in Fig.5D). After familiarization, this percentage increased significantly to 54% (Fig.5C, orange dots in lower panel; orange bar in Fig.5D) (*McNemar* test with Yates's correction, d.f.=1, Chi-square=5.939, p<0.02). Hence familiarization increased bias towards the novel odor in neurons that were more responsive to that odor to begin with, which is consistent with our earlier results(*51*). In different experiments where yohimbine was perfused, the familiarization protocol not only did not increase bias towards the Novel odor, it significantly decreased the original bias from 49% (N=56) to 14% (Fig.5D, grey bars) (*McNemar* test with Yates's correction, d.f.=1, Chi-square=11.13, p<0.001), suggesting that yohimbine interrupted this neural manifestation of latent inhibition in the antennal lobes.

The tyramine/octopamine ratio in the brain is also associated with latent inhibition

A recent report implicated the release of dopamine in driving reward seeking behavior (*53*). In order to evaluate whether dopamine might be involved in latent inhibition, and whether change in release of octopamine and/or tyramine might contribute to our behavioral results, we reanalyzed previously published data (*54*) on levels of dopamine, serotonin, octopamine and tyramine in individual brains of 81 foragers collected from an unselected genetic background used for selection of lines for expression of latent inhibition. The foragers were collected as 'scouts' or 'recruits'. Scouts were defined as the first bees to explore a new landscape into which their colony had been moved. Recruits were defined as foragers that were exploiting resources once they were found. All scouts and recruits were trained for latent inhibition in the laboratory, and then classified as to whether they showed strong or weak latent inhibition based on learning a novel and familiar odor(*54*).

Of the biogenic amines (Figs. 6A,B), only tyramine showed differences between scouts
and recruits (see (*54*) for methods and a more complete analysis of these data). Differ-
ences in dopamine or serotonin levels were not significant. For the current purpose, we
re-analyzed the data to focus on the ratios of tyramine to octopamine and dopamine to
serotonin (serotonin was used a reference for dopamine levels in (*53*)). Scouts that
showed strong latent inhibition also had significantly lower ratio of tyramine to octopamine
than recruits, and that ratio was also lower than scouts and recruits that showed weak
latent inhibition (Figs. 6C,D). There were no significant differences in the dopamine to
serotonin ratios. Thus, there is an interaction of tyramine and octopamine *production* with
behavioral division of foraging labor and expression of latent inhibition. However, dopa-
mine, serotonin and their ratios do not appear to be involved in latent inhibition.

Fig. 6. Biogenic amine levels in individual brains of scout and recruit foragers that ex-
pressed strong or weak latent inhibition. A-B. Absolute levels of octopamine (blue) and tyra-
mine (green) and of dopamine (red) and serotonin (yellow) in individual forager brains. **C.** Ratios
of tyramine/octopamine. In foragers that exhibited strong latent inhibition, the ratio was signifi-
cantly lower in scouts (N=25) than recruits (N=13) (Wilcoxon W=56.0, $p<0.05$). Ratios did not differ
in scouts (N=24) and recruits (N=19) that exhibited weak latent inhibition ($p>0.05$) **D**. Ratios of

dopamine/serotonin did not differ in either the strong or weak groups (p>0.05). Sample sizes the
same as in C.

Discussion

Our results have identified the genetic and neural underpinnings that modulate an im-
portant form of learning and memory in the brain. All animals need to learn about stimuli
in their environment. Latent inhibition is important for redirecting limited attention capacity
away from unimportant, inconsequential stimuli and refocusing it toward novel stimuli
about which the animal knows little or nothing. Two independent QTL mapping studies
have now identified the genetic locus that contains *Amtyr1* as important for regulating
individual variation in attention(*19*). There are other loci in the genome that show associ-
ations with the behavior, and there are also other unidentified genes in the same locus.
Nevertheless, our manipulation of *Amtyr1* function using both pharmacology and DsiRNA
treatments confirm its association with behavioral expression of latent inhibition.

There are a few points that need to be kept in focus at this point in our understanding
of *Amtyr1*. First, it is not a *latent inhibition gene*. Instead, it is a gene that has broad effects
on foraging-related behaviors that include its effect on expression of latent inhibition. In
that sense it has major effects on a broader behavioral syndrome that includes effects on
sucrose sensitivity (*24, 29, 31*), preferences for nectar and/or pollen (*21*), behavioral caste
differences (*32*), reproductive physiology (*55*) and learning (*12*). It remains to be deter-
mined how *Amtyr1* can have such broad effects. Part of the answer undoubtedly lies in its
expression in several areas of the brain (*28*). Additionally, as we discuss below, it is likely
modulating afferent processing that, in the case we describe here, drives Hebbian plas-
ticity.

Second, the effects of *Amtyr1* specifically on expression of latent inhibition likely arise
by combining its expression in sensory as well as more central areas of the brain. That is,
it is unlikely that there is a single locus in the brain that underlies latent inhibition. We have
shown for honey bees(*28*), for example, that AmTYR1 is on presynaptic terminals of Ol-
factory Sensory Neuron axons in the antennal lobes, where they provide excitation to
dendrites of Local GABAergic Inhibitory Interneurons (LN) and Projection Neurons (PN)
(Fig. 7A). AmTYR1 is also on presynaptic terminals of Projection Neuron axons that

terminate in the mushroom body calyces. In the antennal lobe, where our electrophysio­logical and imaging studies have focused (*51, 52, 56, 57*), familiarization to an odor causes the neural representation of a mixture that contains the familiar odor to become more like novel odors in the mixture (*52*). And it potentiates the response to the novel odor(*51*). We assume, but have yet to show experimentally, that this bias combines with how AmTYR1 affects processing in the mushroom bodies, where olfactory information converges with information from other sensory modalities. These higher-order effects of AmTYR1 could underlie why individuals from genetic lines selected in the laboratory for odor-based latent inhibition show differential attention to different sensory stimuli associ­ated with feeders when tested in free flying conditions in the field (*22*).

Figure 7. Hypothetical model of how *Amtyr1* could modulate hebbian plasticity in feed-for­ward and lateral inhibition to modulate latent inhibition. A. Circuitry of two glomeruli (Figure 7 A has been adapted from Figure P1 from (*58*)) that underlies latent inhibition (habituation) in the fruit fly antennal lobe, and which was also proposed via a computational model to underlie changes in antennal lobe responses to familiar and novel odors in the honey bee (*52*). In the fruit fly(*58*), ORNs release Ach (purple) to coactivate PNs and LNs, which both express ACh receptors. LNs release both GABA (red) and glutamate (green) onto synapses with PNs. NMDA receptors

on PNs potentiate the LN/PN synapses via a retrograde signal (green arrows) to LNs. This hebbian

plasticity increases inhibition of PNs by LNs and thus decreases excitation of the PN. LN/PN co-

activation in other glomeruli is either non-existent (e.g.(*58*)), too low to produce potentiation of the

LN/PN synapse, or strong enough to produce inhibition to odors other than the familiar odor.

AmTYR1 receptors on presynaptic terminals of ORNs (*28*) are shown as black triangles. Here we

do not show the source for tyramine. We assume it could be from the VUM neuron, which has

terminals in the cortex of every glomerulus, where ORN terminals overlap with PN and LN den-

drites(*59*). B-D. Increasing activation of AmTYR1 (pink triangles) progressively decreases release

of ACh and lowers coactivation of LNs and PNs. Decreased release of glutamate reaches a

threshold (D) below which it fails to modify the LN/PN synapse via a decreased retrograde signal,

although ACh release might still activate the glomeruli. Graphs inset in each figure show activation

of AmTYR1 (x-axis) verses LTP-based hebbian plasticity (y-axis). Low activation of AmTYR1 pro-

duces high coactivation-based LTP (B) and vice-versa (D). Arrows show the hypothetical relation-

ship in each figure. Gray lines represent the predicted acquisition levels to novel (solid) and famil-

iar (dashed) odors given the strength of Hebbian plasticity in each figure. For these lines the x-

axis would be 'Trial' and y-axis 'Percent proboscis extension', as in Figs. 3 and 4.

The precise relationship of *Amtyr1* to latent inhibition is different from what is normally

expected from disruption of a gene that underlies a behavior. We expected that disruption

of *Amtyr1* function would reduce or eliminate latent inhibition; that is, learning about a

familiar odor (X) would rise to equal learning about the novel odor. Instead, the response

to the novel odor was reduced to equal that to the familiar odor. This reduction was specific

to familiarization treatment, so it is dependent on plasticity. It cannot be explained by non-

specific – e.g. toxic – effects of treatment, because the same treatments did not reduce

excitatory conditioning in the absence of familiarization to an odor. Moreover, the same

effect was evident using two very different means for disruption of *Amtyr1* signaling.

We propose that *Amtyr1* modulates neural plasticity that reduces attention to a familiar

odor. *Amtyr1* maintains coactivation of LNs and PNs in the antennal lobe at a set point

between the extremes where it becomes too strong (e.g. when *Amtyr1* is disrupted) or too

weak (*Amtyr1* strongly activated). Given that activation of *Amtyr1* reduces cAMP levels, it

would be expected that this would reduce excitability of axon terminals. Hypothetically

then, activation of *Amtyr1* could reduce excitatory drive of post-synaptic processes on

PNs and LNs in, for example, the antennal lobes, and possibly between PN axons and

intrinsic neurons of the mushroom bodies(*28*).

A model of how *Amtyr1* could modulate inhibition in antennal lobe networks

We can now propose a model for how *Amtyr1* could act in the antennal lobes, and

possibly also as an important component for encoding latent inhibition in distributed net-

works in the mushroom body. The model would have to explain why air (mechanosensory)

stimulation alone does not reduce subsequent acquisition to odors (Figs. 3A, 4A) as much

as familiarization to an odor (Figs. 3B,C, 4B). Mechanosensory stimulation produces fast,

transient responses in antennal lobe glomeruli, whereas odor stimulation produces more

robust, longer lasting responses(*60, 61*). These differences in response could underlie the

difference in response to air versus odor in our analyses, particularly in driving activity

dependent plasticity at LN-to-PN synapses that we describe below.

We propose that effects *specific to odor familiarization* occur by amplifying hebbian

plasticity between PNs and LNs in the antennal lobe networks (Fig. 7A). A computational

model of the honey bee antennal lobe identified hebbian plasticity at Local GABAergic

Inhibitory Interneuron (LN) synapses onto Projection Neurons (PN) as the most likely lo-

cus of plasticity to give rise to the observed biasing of an odor mixture to be less like a

familiar odor and more like a novel odor (*52*). If ORy in Fig 7A is not activated by an odor

that activates ORx, then after familiarization to X a novel odor that activates ORy would

be better able to suppress X in a mixture because of strengthened LN-to-PNx synapses

(see Fig 7 in (*52*)). A potentiated novel odor would strengthen this effect. Work with fruit

flies also showed that glutamate/NMDA receptor dependent hebbian plasticity at the same

synapses underlies odor habituation ((*58*); the same process as latent inhibition just dif-

ferently named).

In our model (Fig. 7A-D), coactivation of PNs and LNs via excitation from OR axon

terminals would produce plasticity at the LN-to-PN synapses. Low activation of AmTYR1

(Fig. 7B) would lead to strong input from sensory axon terminals that would maximally

activate LNs and PNs, thus leading to strong Hebbian plasticity at LN-to-PN synapses.

That is, it would lead to the strong inhibition of PNs in our manipulations that blocked or

disrupted AmTYR1 signaling. At moderate levels (Fig. 7C) this plasticity would give rise

to normal latent inhibition in our high lines just as it does in fruit flies. Strong activation of

AmTYR1 would weaken the activity and reduce or prevent plasticity (Fig. 7D), and hence

lead to learning about both the novel and familiar odors (such as in our genetic lines that

do not show strong latent inhibition; (*12*)). How well this model works might depend on a

nonlinear thresholding function for coactivation that drives hebbian plasticity (AmTYR1-

LTP relationships and expected learning curves shown in Figs. 7B-D).

Furthermore, strong activation of lateral projections of the inhibitory interneuron (Fig.

7A) might cause generalization of latent inhibition to novel odors, as we have observed

with disruption of AmTYR1 signaling. Many of the monomolecular odorants that have

been used to study olfaction and olfactory learning in honey bees have neural activity

patterns that partially overlap (*62*), such as with the representations for hexanol and 2-

octanone. For each odor, PNs in a few of the 160+ glomeruli of the antennal lobe are

highly activated, and a subset of other PNs are activated to a lesser degree. Hypotheti-

cally at least, if the familiar odor activates some of the ORs from glomeruli that also code

for the novel odor, then coactivation of those PNs with lateral inhibition could cause

Hebbian plasticity at those synapses too. Under normal circumstances the excitation of

those ORs might be too low to potentiate inhibition (Fig. 7C,D). But when AmTYR1 sig-

naling is too low or disrupted (Fig 7B), coactivation would increase enough to drive the

plasticity and reduce responses to novel odors.

We have linked expression of latent inhibition to hebbian plasticity in synapses from

inhibitory LNs to PNs in the antennal lobe. Although any behavioral phenomenon likely

arises from distributed neural networks in the brain, we focused on the effect of familiari-

zation in the antennal lobes because of our prior work analyzing odor processing and

latent inhibition there in the honey bee (*52*), and because of reports of latent inhibition in

the same networks in the fruit fly (*58*). In particular, we report neural effects (suppression

of potentiation to the novel odor) that are produced by the same familiarization treatment

that produces behavioral latent inhibition, and like the behavior is affected by blockade or

disruption of AmTYR1. We do not specifically identify the type of LN represented in Fig.

7, but it could be likely that it would belong to the group of heteroLNs(*63*), which receive

excitatory inputs in one glomerulus and transmit inhibition to other glomeruli. Computa-

tional modeling suggests that the interglomerular connectivity of heteroLNs should be

based on 'functional networks' defined by overlapping glomerular activity patterns to sim-

ilar odors(*50*). However, Fig 7 only represents a minimal part of the network that we feel

is needed to convey how it hypothetically modulates Hebbian plasticity. It will be useful to

consider the broader network as represented in Sinakevitch et al (*28*) to more clearly un-

derstand how AmTYR1 functions in the antennal lobe, and how it may function in the

calyces of the mushroom bodies as well.

We show in Figure 5 that treatment with yohimbine to block AmTYR1 reduced re-

sponses to odor in general, which is consistent with increased excitation from ORN axon

terminals driving inhibition, as represented in Fig 7. In fact, this reduction (in the antennal

lobe and possibly the mushroom bodies) could be the reason for slightly lower acquisition

in Fig 4A under dsRNA treatment. If true, the reduction in unit responsiveness did not

completely block excitatory conditioning to odors.

We show the effect of odor familiarization in the antennal lobe as potentiation of the

response to a novel odor, which is consistent with our earlier report(*51*). AmTYR1 block

coupled to familiarization decreased potentiation of responses to the novel odor relative

to the increase normally observed in controls. Although we show that potentiation after

familiarization depends on activation state of AmTYR1, we do not, and at this point for

lack of data we cannot, represent in Figure 7 what gives rise to it. Potentiation could occur

via an as yet unknown mechanism intrinsic to the antennal lobe neural networks(*28*). Al-

ternatively, it could occur via identified feedback pathways to the antennal lobe (*64, 65*)

from neural mechanisms in the mushroom bodies that are known to produce potentiation

to novel stimuli in fruit flies(*66*). Under normal circumstances AmTYR1 is functional and

moderating the Hebbian plasticity at LN-to-PN synapses at levels consistent with Figs 7C

and 7D, where novel odors are learned well (likely aided by potentiation). Blocking or

disrupting AmTYR1 puts the network in a state consistent with Fig 7B, where responses

to both types of odors are affected by Hebbian plasticity at LN-to-PN synapses – including

potentiated novel odors given the overlap in sensory representations.

Finally, it remains to be determined what the source for tyramine could be, and which

of the local interneuron types might be the ones mediating latent inhibition in the antennal

lobes. VUM neurons are an obvious possibility for the source of tyramine, since tyramine

is a direct precursor to octopamine released by VUM neurons(*67*). However, this hypoth-

esis would depend on tyramine being released at almost constant, low levels without stim-

ulation of VUM by taste receptors sensitive to sugars.

Ideas and Speculation

The modulatory role for inhibition that we propose for *Amtyr1* could help to explain its

broad involvement in many different behaviors. Differing degrees of *Amtyr1* activation in

different neural circuits in the central or peripheral nervous system might regulate activity

in those circuits to drive behaviors in one direction or the other; for example, toward high

or low sensitivities to sucrose(*31*), preferences for pollen versus nectar(*23*), and states of

worker reproductive physiology(*55*).

We were initially drawn to *Amtyr1* because of its relationship potentially to the release

of tyramine by identified VUM neurons, which have been implicated in excitatory condi-

tioning through release of octopamine (*35, 36*). VUM neurons must make tyramine in the

process of making octopamine, and they likely release both biogenic amines. In particular,

the dynamic balance between octopamine and tyramine is important for regulating insect

behaviors (*40, 41*). It is intriguing to now propose and eventually test whether a balance

between octopamine and tyramine release from VUM neurons is critical for driving atten-

tion in one direction or another depending on association with reinforcing contexts. In this

model, activation of VUM neurons would release octopamine to drive excitatory associa-

tion between odor and reinforcement. At the same time, release of tyramine would sup-

press excitatory drive onto inhibition. Both processes could synergistically drive the asso-

ciation. Furthermore, if there is a low level of background tyramine release from VUM

when unstimulated, it would explain why in the *Amtyr1* DsiRNA injected group in Fig 4A

responded slightly lower than the Dsiscr control group.

Interestingly, we have identified a potential interaction in the ratio of tyramine to octo-

pamine between foraging role (scouts versus recruits) and expression of latent inhibition.

The lower tyramine-to-octopamine ratios in scouts would potentially activate this receptor

even less that it would normally be, yielding stronger inhibition according to the model

described above. Further analyses need to test this prediction in more detail and evaluate

its role in the foraging ecology of honey bees.

Finally, why do individuals in colonies under quasi-natural conditions differ in expres-

sion of latent inhibition, and presumably in the functioning of *Amtyr1*? We have used this

naturally occurring and selectable genetic variation to establish colonies composed of dif-

ferent mixtures of genotypes(*22, 68*). The mixture of genotypes in the colony affects

whether and how quickly colonies discover new resources via an attention-like process

operant in individual for-

agers(*68*). We have there-

fore proposed that genetic

variation leading to colony

level variation in *Amtyr1*

expression represents a

balance between explora-

tion for and exploitation of

resources. The precise

balance of genotypes

would give colonies flexi-

bility to respond to chang-

ing resource distributions

over the life of the colony.

Box 1. Important questions that need to be ad-
dressed in honey bees and other animal mod-
els, such as the fruit fly, and in computational
models:

- To what extent is tyramine constantly released
  at a background level, such that it modulates
  activity of AmTYR1?
- What is the balance of octopamine and tyra-
  mine during odor stimulation and when the
  odor is associated with reinforcement?
- How does the action of AmTYR1 in distributed
  networks, such as the antennal lobes and
  mushroom bodies, coordinate to produce be-
  havioral expression of latent inhibition (Fig. 7)?
- How is *amtyr1* activity regulated by other genes
  in a network, and by epigenetic factors in the
  environment?

Materials and Methods

Selection of honey bee lines for differences in latent inhibition. We established high

and low latent-inhibition lines by conditioning drone and virgin queen honey bees to odors

in three different conditioning phases(*12*). The first phase involved selection of drones or

queens that could successfully learn to associate an odor with sucrose reinforcement,

which established that the honey bees were motivated to learn. This initial excitatory con-

ditioning does not affect generation or expression of latent inhibition. The second 'famil-

iarization' phase involved 40 unreinforced odor exposures for 4 sec each; this new odor

(black box; X) was different and discriminable from the first odor. The third and final phase

involved conditioning honey bees to the now familiar odor X associated with sucrose

reinforcement in a way that normally produces robust associative conditioning (*25*). Strong

latent inhibition should slow the rate of learning to X. Drones and queens that exhibited

this 'inhibitor' phenotype (defined as zero or one response to X over six conditioning trials)

(*12*) were mated using standard instrumental insemination techniques (*69*) for honey bees

to create a high (inhibitor) latent inhibition line. Drones and queens that learned X quickly

(five positive responses to X over six trials) were also mated to produce a low (noninhibi-

tor) latent inhibition line. Our previous studies have shown that worker progeny from in-

hibitor and non-inhibitor matings showed significant correlation in expression of latent in-

hibition to their parents (*12, 70*).

Recombinant drones Male honey bees (drones) were produced from a cross between

genetic lines selected for high and low expression of latent inhibition(*12, 17*). Hybrid

queens were reared from a cross of a queen from the inhibitor line instrumentally insemi-

nated(*71*) with sperm from a single drone from the noninhibitor line. These queens were

then allowed to mate naturally to increase longevity in a colony. Natural mating involves

mating with several different drones. However, since drones arise from unfertilized eggs,

the haploid (drone) genotype involves only recombination of the genotypes of the high

and low lines in the hybrid queen. A single hybrid queen was then selected to produce

drones. Sealed drone brood from the hybrid queen was placed in a small nucleus colony.

Queen excluder material (wire mesh that does not permit the passage of queens or

drones) was used to confine the emerging drones to the upper story. Upon emergence,

drones were individually marked on the thorax with enamel paint for later identification,

and then marked drones were co-fostered in a single outdoor colony until collected for

behavioral conditioning.

Mature drones were collected from the colony upon returning from mating flights during

the late afternoon the day before testing. Returning drones gathered on a piece of queen

excluder material blocking the colony entrance and were put into small wooden boxes

with queen excluder material on each side. They were then fed a small amount of honey

and placed in a queenless colony overnight. The following morning drones were secured

in a plastic harness using a small piece of duct tape (2 mm x 20 mm) placed between the

head and the thorax (*25*). All drones were then kept at room temperature for two hours.

They were then screened for their motivation to feed by lightly touching a small drop of

2.0 M sucrose solution to the antennae. Drones that extended their proboscis were se-
lected for training.

Foragers Female pollen foragers (workers) were captured at the colony entrance as de-
scribed above. Each bee was chilled to 4˚ C, restrained in a harness and fed to satiation
with 1.0 M sucrose. The next day bees were tested for motivation by stimulation of their
antennae with 2.0 M sucrose; bees that extended their proboscis were used in experi-
ments shown in Figures 3 and 4.

Conditioning protocols

Familiarization: Familiarization to the odor was done as described in (*6*). Restrained bees
were placed in individual stalls where a series of valves regulated odor delivery via a
programmable logic controller (PLC) (Automation Direct). Hexanol and 2-octanone were
used either as pure odorants or diluted to 2.0 M in hexanes with odor treatments counter-
balanced across animals. Odor cartridges were made by applying 3.0 μl of odorant onto
a piece of filter paper (2.5 x 35 mm) and inserting the filter paper into a 1 ml glass syringe.
The odor cartridge was then connected to a valve regulated by the PLC that shunted air
through the cartridge for 4 s once the automated sequence was initiated. Odor preexpo-
sure in all experiments involved 40 unreinforced presentations of odor for 4 s using a 5
min (Fig 1) or 30 s (Fig 3) intertrial interval (ITI). All odor cartridges were changed for fresh
ones after every 10 uses to avoid odor depletion(*72*). The use of pharmacological treat-
ment necessitated the use of a shorter ITI to avoid having the drug wear off before the
end of preexposure. Our previous studies have revealed that latent inhibition is robust
over this range of ITIs and odor concentrations (*6*).

Proboscis Extension Response conditioning: All PER learning paradigms used for testing
used a 5 min ITI. An acquisition trial consisted of a 4 s presentation of an odor, the condi-
tioned stimulus (CS, black or gray bars), followed by presentation of a 0.4 μl drop of 1.0
M sucrose solution, the unconditioned stimulus (US, triangles in Figs 1, 3 and 4). Three
seconds after onset of the CS the US was delivered using a Gilmont ® micrometer syringe.
The US was initially delivered by gently touching the antennae to elicit proboscis extension
and subsequent feeding. Once a bee began to extend its proboscis at the onset of CS
delivery, it was no longer necessary to touch the antennae prior to feeding.

We used two different procedures for testing latent inhibition after familiarization. For

evaluation of recombinant drones (Fig 1), subjects were conditioned to the familiarized

odor (X) as the CS over 6 forward pairing trials. The second procedure (Figs 3 and 4)

involved use of a within animal control protocol. After familiarization all subjects received

equivalent PER conditioning to 2 odors, one was the familiarized odor (X) and the other

was a novel odor (N) that honey bees can easily discriminate from the familiarized

odor(*73*). Odors were presented in a pseudorandomized order (NXXNXNNX or

XNNXNXXN) across trials such that equal numbers of animals received N or X on the first

trial. Pharmacological treatment required the use of a control procedure involving famil-

iarization to air to evaluate the degree to which expression of excitatory conditioning was

affected by drug treatment (Figs. 3A and 4A).

Linkage analysis Upon completion of the training paradigm, 523 drones were placed in

individual 1 ml micro-centrifuge tubes and stored at -70° C. Genomic DNA extraction fol-

lowed a standard protocol developed for honey bees (*74*). For SNP analysis, DNA was

selected from 94 drones that exhibited the highest level of latent inhibition (0, 1, 2, or 3

responses over the six test trials) and from another 94 drones that exhibited the lowest

level (5 or 6 responses). Analysis of the 188 samples was conducted by Sequenom, Inc.

San Diego, California.

The linkage map was built with a set of 311 SNP markers. The list of selected markers

was provided by Olav Rueppell from previous studies examining the genetic architecture

of foraging behavior and sucrose response thresholds (*75, 76*). The 74 SNPs segregating

in our mapping population were used for a QTL analysis. Map positions for markers in

linkage group one were determined using the *Apis mellifera* 4.0 genome. The software

MultiPoint 1.2 (http://www.mulitqtl.com) was used to determine the actual recombination

frequencies for markers in linkage group 1. Recombination frequencies were then con-

verted to centiMorgans using the Kosambi mapping function. The actual mapping dis-

tances in our mapping population were used in the QTL analysis. QTL analysis was per-

formed with MapQTL 4.0. Interval mapping and MQM mapping revealed one significant

QTL. Genomewide significant thresholds for $p<0.05$ (LOD = 2.6) and $p<0.01$ (LOD = 3.2)

were determined using an implemented permutation test (1000 runs).

Pharmacological and DsiAmTyr1 treatments. Yohimbine hydrochloride (Sigma) was diluted to 10^{-4} M in saline (5 mM KCl, 10 mM NaH$_2$PO$_4$, pH 7.8). We chose a concentration of yohimbine that has been shown to be effective in our previous study of its effect on honey bee behavior (*42*). One µl of drug or saline alone was injected into the brain through the median ocellus using a Hamilton syringe (Hamilton; Reno, NV). Training began 15 min after injection, as this time has been shown to be effective in other drug studies using the same methodology (*6, 48, 77*).

For DsiRNA studies, we used sequences and protocols developed previously for a study of *Amtyr1* receptor distribution in the brain, which in that study were used to show that the anti-*Amtyr1* antibodies specifically recognized the receptor(*28*). We used a Dicer-substrate small interfering (Dsi) RNA of the *AmTyr1* receptor (NCBI Reference Sequence: NM_001011594.1) to knockdown *AmTyr1* mRNA receptor in the brain. We used the mix-ture of three DsiAmTyr1 constructs designed by the tool in IDT technology(*28*) (Table 1). As a control we used a scrambled (dsiScr) version of the *Amtyr1* sequence. A 138 nano-liter injection of a 100microM mixture of dsiAmTyr1 or dsiScr (Nanoinject 2000) was made into the middle ocellus 18-20 hours before behavioral tests. All injections were done blind so that the investigator doing behavioral tests was not aware of the content of the injection. After the tests brains without optic lobes were dissected out and homogenized each in TRIzol (Invitrogen) (N=27 for bees injected with dsiScr and N=32 for bees injected with DsiAmTyr1). Then, the total mRNA from each injected brain was extracted separately using the manufacturer's protocol for TRIzol method (Invitrogen). Contaminating genomic DNA was removed using DNA-free^{TM} kit (Ambion, AM1906). RNA quantity and purity was evaluated using a NanoDrop (NanoDrop 2000). Expression of AmTyr1 was quantified us-ing QuantiFAST SYBR Green RT-PCR kit (QIAGEN) on Applied Biosystem 7900 cycler (ASU DNA Facilities) with the protocol provided by the kit for a 384-well plate. The primers for quantitative real-time PCR assays were: AmTyr1_F 5'-GTTCGTCGTATGCTGGTTGC-3', AmTyr1_R 5'- GTAGATGAGCGGGTTGAGGG-3' and for reference gene AmActin_F 5'- TGCCAACACTGTCCTTTCTG-3' AmActin_R 5'-AGAATTGACCCACCAATCCA-3'(*78*).

Table 1

Nucleotide sequences of sense and antisense strands of control DsiSCR and AmTyr1 DsiRNA.

____

DsiRNA	Sequences
DsiScr	5'- GAGUCCUAAGUUAACCAAGUCACAGCA-3
	3'- CUCAGGAUUCAAUUGGUUCAGUGUCGU-5'
DsiTyr1_N	5'-AGCGUGACGUUGGAUUGACGAGAGC-3'
	3'-CCUCGCACUGCAACCUAACUGCUCUCG-5'
DsiTyr1_T1	5'-CCUGUGCAAAUUGUGGCUAACCUGC-3'
	3'-GUGGACACGUUUAACACCGAUUGGACG-5'
DsiTyr_C	5'-CAACGCUUGUUUAUUGCAUCUAUCG-3'
	3'-CCGUUGCGAACAAAUAACGUAGAUAGC-5'

All injections were done blind so that the investigator doing behavioral tests was not
aware of the content of the injection.

Electrophysiological recordings from the antennal lobe. Extracellular recordings were
performed in the antennal lobes with a 16-channel probe (NeuroNexus, Ann Arbor, MI).
Spike waveforms were digitized with a RZ2 system at a sampling rate of 20KHz (Tucker-
Davis Technologies, Alachua, FL). After a stable recording was achieved, the honeybee
preparation was first stimulated with two presentations of each of the following odors: 1-
hexanol (Hex) and 2-octanone (Oct). The duration of each pulse was 4 sec, and two
minutes of recovering time were allowed between two pulses. During the pre-exposure
phase, 40 pulses of Oct were delivered with inter-pulse interval of 60 sec, after which 10
min recovery was given before testing. Upon completion of each experiment, extracellular
spike waveforms are exported to Offline Sorter program (Plexon Inc, Dallas, TX) which
classifies the similar waveforms into individual clusters (units) based on the relatedness
of waveforms' projection onto a 3D space derived from the first three principle components
that capture the most variation of the original waveforms. To increase the discriminating
power, the original waveforms are grouped in a tetrode configuration, matching the phys-
ical design of the recording probe, i.e. 16 recording sites are distributed in two shanks in
a block design of 2x4. Each block is called a tetrode. Statistical separation of waveform

clusters, representing individual neurons or units, is aided with visual inspection, all implemented in the Offline Sorter program. Once satisfied with the clustering results, the time stamps of waveforms are then exported to Neuroexplorer program (Plexon Inc, Dallas, TX) and Matlab (Mathworks, Natick, MA) for further analysis.

Yohimbine (Millipore-Sigma, St. Louis, MO.), was diluted in saline (50 µM), which was perfused into the head capsule through a T-tube switch. Repeated stimulation with Oct started fifteen minutes after perfusion; by then the slowing-down of spiking activities were often noticeable. Care was taken not to introduce any air bubble into the tubing when switching from the syringe containing saline to the syringe containing the yohimbine solution. The water level in the two syringes was intentionally kept the same in order to maintain a similar perfusing rate upon switching. The drug solution was kept flowing through the honeybee preparation until the end of protocol, which usually lasted for about 2 hours. No saline wash was attempted in this protocol due to the long time required for the recording sessions.

Sequencing the *Amtyr1* region of the genome. We studied single nucleotide polymorphisms (SNPs) in full-genome sequences of eight *A. mellifera* workers (four high pollen hording and four low pollen hording). For each individual, Illumina short reads were mapped against the *Apis mellifera* genome assembly version 4.5 (*79*) using bwa version 0.5.9-r16 (*80*). An average 25x genome coverage per individual allowed the identification of high-quality SNPs in each individual against the reference genome. SNPs were identified with SAMtools version 0.1.17-r973:277 (*81*) enforcing a minimum quality score of 20 (base call accuracy ≥ 99%).

Statistical analysis. To analyze the effects in behavioral experiments, we used a generalized linear model with binomial error distribution and logit transformation to perform a logistic regression. The response variable is binomal (0,1). Trial is an ordered variable. We were most interested in testing the hypothesis that injection of yohimbine and dsiRNA before familiarization treatment would impact latent inhibition, so we focused on the interactions between trial, injection, injection time (before pre-exposure or before acquisition), and odor (novel or pre-exposed odor). To explore significant interactions

further, we performed a tukey post hoc test using the package emmeans. All analyses were performed in R version 4.2.0 using RStudio version 2022.07.1.

References

1. C. Heyes, Simple minds: a qualified defence of associative learning. *Philos Trans R Soc Lond B Biol Sci* **367**, 2695-2703 (2012).
2. A. Dickinson, Associative learning and animal cognition. *Philos Trans R Soc Lond B Biol Sci* **367**, 2733-2742 (2012).
3. N. Mackintosh, *Conditioning and Associative Learning*. (Clarendon Press, 1983).
4. R. Lubow, *Latent Inhibition and Conditioned Attention Theory*. (Cambridge University Press, Cambridge University Press, 1989), pp. 324.
5. R. E. Lubow, Latent inhibition. *Psychological Bulletin* **79**, 398-407 (1973).
6. S. B. Chandra, G. A. Wright, B. H. Smith, Latent inhibition in the honey bee, Apis mellifera: Is it a unitary phenomenon? *Anim Cogn* **13**, 805-815 (2010).
7. B. H. Smith, G. A. Wright, K. S. Daly, in *The Biology of Floral Scents,* N. Dudareva, E. Pichersky, Eds. (CRC Press, Boca Raton, FL, 2006), pp. 263-295.
8. C. I. Abramson, M. E. Bitterman, Latent inhibition in honeybees. *Animal Learning & Behavior* **14**, 184-189 (1986).
9. S. Seefeldt, R. J. De Marco, The response of the honeybee dance to uncertain rewards. *J Exp Biol* **211**, 3392-3400 (2008).
10. V. M. Fernández, A. Arenas, W. M. Farina, Volatile exposure within the honeybee hive and its effect on olfactory discrimination. *Journal of Comparative Physiology A* **195**, 759-768 (2009).
11. C. Brandes. (Behavior Genetics, 1991), vol. 21(3), pp. 271-294.
12. S. B. Chandra, J. S. Hosler, B. H. Smith, Heritable variation for latent inhibition and its correlation with reversal learning in honeybees (Apis mellifera). *Journal of Comparative Psychology* **114**, 86-97 (2000).
13. V. Finke, D. Baracchi, M. Giurfa, R. Scheiner, A. Avarguès-Weber, Evidence of cognitive specialization in an insect: proficiency is maintained across elemental and higher-order visual learning but not between sensory modalities in honey bees. *J Exp Biol* **224**, (2021).
14. B. H. Smith, C. I. Abramson, T. R. Tobin, Conditional withholding of proboscis extension in honeybees (Apis mellifera) during discriminative punishment. *Journal of Comparative Psychology* **105**, 345-356 (1991).
15. E. Pamir, P. Szyszka, R. Scheiner, M. P. Nawrot, Rapid learning dynamics in individual honeybees during classical conditioning. *Front Behav Neurosci* **8**, 313 (2014).
16. P. R., *The Mechanisms of Social Evolution*. (Harvard Univ Press, Boston, MA, 2013).
17. J. S. Latshaw, B. H. Smith, Heritable variation in learning performance affects foraging preferences in the honey bee (Apis mellifera). *Behavioral Ecology & Sociobiology* **58**, 200-207 (2005).
18. T. Mosqueiro *et al.*, Task allocation and site fidelity jointly influence foraging regulation in honeybee colonies. *R Soc Open Sci* **4**, 170344 (2017).

19. S. B. Chandra, G. J. Hunt, S. Cobey, B. H. Smith, Quantitative trait loci associated with
reversal learning and latent inhibition in honeybees (Apis mellifera). *Behavior Genetics* **31**,
275-285 (2001).
20. G. J. Hunt *et al.*, Behavioral genomics of honeybee foraging and nest defense.
Naturwissenschaften **94**, 247-267 (2007).
21. R. E. Page, Jr. *et al.*, Genetic dissection of honeybee (Apis mellifera L.) foraging behavior.
Journal of Heredity **91**, 474-479 (2000).
22. C. N. Cook *et al.*, Individual learning phenotypes drive collective behavior. *Proc Natl Acad*
Sci U S A **117**, 17949-17956 (2020).
23. G. J. Hunt, R. E. Page, Jr., M. K. Fondrk, C. J. Dullum, Major quantitative trait loci affecting
honey bee foraging behavior. *Genetics* **141**, 1537-1545 (1995).
24. T. Pankiw, K. D. Waddington, R. E. Page, Jr. , Modulation of sucrose response thresholds
in honey bees (Apis mellifera L.): influence of genotype, feeding, and foraging experience.
Journal of Comparative Physiology A-Sensory Neural & Behavioral Physiology **187**, 293-
301 (2001).
25. M. E. Bitterman, R. Menzel, A. Fietz, S. Schafer, Classical conditioning of proboscis
extension in honeybees (*Apis mellifera*). *Journal of Comparative Psychology* **97**, 107-119
(1983).
26. W. Blenau, S. Balfanz, A. Baumann, Amtyr1: characterization of a gene from honeybee
(*Apis mellifera*) brain encoding a functional tyramine receptor. *J Neurochem* **74**, 900-908
(2000).
27. J. A. Mustard, P. T. Kurshan, I. S. Hamilton, W. Blenau, A. R. Mercer, Developmental
expression of a tyramine receptor gene in the brain of the honey bee, *Apis mellifera*. *J Comp*
Neurol **483**, 66-75 (2005).
28. I. T. Sinakevitch, S. M. Daskalova, B. H. Smith, The Biogenic Amine Tyramine and its
Receptor (AmTyr1) in Olfactory Neuropils in the Honey Bee (*Front Syst Neurosci* **11**, 77
(2017).
29. M. Thamm *et al.*, Neuronal distribution of tyramine and the tyramine receptor AmTAR1 in
the honeybee brain. *J Comp Neurol* **525**, 2615-2631 (2017).
30. W. Blenau, J. A. Wilms, S. Balfanz, A. Baumann, AmOctα2R: Functional Characterization
of a Honeybee Octopamine Receptor Inhibiting Adenylyl Cyclase Activity. *Int J Mol Sci*
21, (2020).
31. R. Scheiner *et al.*, Learning, gustatory responsiveness and tyramine differences across nurse
and forager honeybees. *J Exp Biol* **220**, 1443-1450 (2017).
32. R. Scheiner, L. Kulikovskaja, M. Thamm, The honey bee tyramine receptor AmTYR1 and
division of foraging labour. *J Exp Biol* **217**, 1215-1217 (2014).
33. M. Kutsukake, A. Komatsu, D. Yamamoto, S. Ishiwa-Chigusa, A tyramine receptor gene
mutation causes a defective olfactory behavior in *Drosophila melanogaster*. *Gene* **245**, 31-
42 (2000).
34. T. Roeder, in *Annual Review of Entomology*. (2005), vol. 50, pp. 447-477.
35. T. Farooqui, K. Robinson, H. Vaessin, B. H. Smith, Modulation of early olfactory
processing by an octopaminergic reinforcement pathway in the honeybee. *Journal of*
Neuroscience **23**, 5370-5380 (2003).
36. M. Hammer, An identified neuron mediates the unconditioned stimulus in associative
olfactory learning in honeybees. *Nature* **366**, 59-63 (1993).

37.   I. Sinakevitch, M. Niwa, N. J. Strausfeld, Octopamine-like immunoreactivity in the honey
bee and cockroach: comparable organization in the brain and subesophageal ganglion. *J*
Comp Neurol **488**, 233-254 (2005).

38.   I. Sinakevitch, G. R. Bjorklund, J. M. Newbern, R. C. Gerkin, B. H. Smith, Comparative
study of chemical neuroanatomy of the olfactory neuropil in mouse, honey bee, and human.
Biol Cybern **112**, 127-140 (2018).

39.   S. Kreissl, S. Eichmuller, G. Bicker, J. Rapus, M. Eckert, Octopamine-Like
Immunoreactivity In the Brain and Subesophageal Ganglion Of the Honeybee. *Journal of*
Comparative Neurology **348**, 583-595 (1994).

40.   N. L. Kononenko, H. Wolfenberg, H. J. Pfluger, Tyramine as an independent transmitter
and a precursor of octopamine in the locust central nervous system: an
immunocytochemical study. *J Comp Neurol* **512**, 433-452 (2009).

41.   N. Schützler *et al.*, Tyramine action on motoneuron excitability and adaptable
tyramine/octopamine ratios adjust. *Proc Natl Acad Sci U S A* **116**, 3805-3810 (2019).

42.   B. L. Fussnecker, B. H. Smith, J. A. Mustard, Octopamine and tyramine influence the
behavioral profile of locomotor activity in the honey bee (*Apis mellifera*). *J Insect Physiol*
52, 1083-1092 (2006).

43.   T. Reim *et al.*, AmTAR2: Functional characterization of a honeybee tyramine receptor
stimulating adenylyl cyclase activity. *Insect Biochem Mol Biol* **80**, 91-100 (2017).

44.   X. Guo, Y. Wang, I. Sinakevitch, H. Lei, B. H. Smith, Comparison of RNAi knockdown
effect of tyramine receptor 1 induced by dsRNA and siRNA in brains of the honey bee,
Apis mellifera. *J Insect Physiol* **111**, 47-52 (2018).

45.   B. Gerber *et al.*, Honey bees transfer olfactory memories established during flower visits
to a proboscis extension paradigm in the laboratory. *Animal Behaviour* **52**, 1079-1085
(1996).

46.   G. A. Wright *et al.*, Parallel reinforcement pathways for conditioned food aversions in the
honeybee. *Curr Biol* **20**, 2234-2240 (2010).

47.   M. Hammer, R. Menzel, Multiple sites of associative odor learning as revealed by local
brain microinjections of octopamine in honeybees. *Learning & Memory* **5**, 146-156 (1998).

48.   A. R. Mercer, R. Menzel, The effects of biogenic amines on conditioned and unconditioned
responses to olfactory stimuli in the honeybee *Apis mellifera*. *Journal of Comparative*
Physiology **145A**, 363-368. (1982).

49.   G. Bicker, R. Menzel, Chemical codes for the control of behaviour in arthropods. *Nature*
337, 33-39 (1989).

50.   C. Linster, S. Sachse, C. G. Galizia, Computational modeling suggests that response
properties rather than spatial position determine connectivity between olfactory glomeruli.
J Neurophysiol **93**, 3410-3417 (2005).

51.   H. Lei *et al.*, Novelty detection in early olfactory processing of the honey bee, Apis
mellifera. *PLoS One* **17**, e0265009 (2022).

52.   F. F. Locatelli *et al.*, Nonassociative plasticity alters competitive interactions among
mixture components in early olfactory processing. *Eur J Neurosci* **37**, 63-79 (2013).

53.   J. Huang *et al.*, Food wanting is mediated by transient activation of dopaminergic signaling
in the honey bee brain. *Science* **376**, 508-512 (2022).

54.   C. N. Cook *et al.*, Individual differences in learning and biogenic amine levels influence
the behavioural division between foraging honeybee scouts and recruits. *J Anim Ecol* **88**,
236-246 (2019).

55.   Y. Wang, G. V. Amdam, B. C. Daniels, R. E. Page, Tyramine and its receptor TYR1 linked
behavior QTL to reproductive physiology in honey bee workers (Apis mellifera). *J Insect*
Physiol **126**, 104093 (2020).
56.   P. C. Fernandez, F. F. Locatelli, N. Person-Rennell, G. Deleo, B. H. Smith, Associative
conditioning tunes transient dynamics of early olfactory processing. *J Neurosci* **29**, 10191-
10202 (2009).
57.   F. F. Locatelli, P. C. Fernandez, B. H. Smith, Learning about natural variation of odor
mixtures enhances categorization in early olfactory processing. *J Exp Biol* **219**, 2752-2762
(2016).
58.   S. Das *et al.*, Plasticity of local GABAergic interneurons drives olfactory habituation. *Proc*
Natl Acad Sci U S A **108**, E646-654 (2011).
59.   I. T. Sinakevitch *et al.*, Apis mellifera octopamine receptor 1 (AmOA1) expression in
antennal lobe networks of the honey bee (Apis mellifera) and fruit fly (Drosophila
melanogaster). *Front Syst Neurosci* **7**, 70 (2013).
60.   H. Tuckman, J. Kim, A. Rangan, H. Lei, M. Patel, Dynamics of sensory integration of
olfactory and mechanical stimuli within the response patterns of moth antennal lobe
neurons. *J Theor Biol* **509**, 110510 (2021).
61.   H. Tuckman, M. Patel, H. Lei, Effects of Mechanosensory Input on the Tracking of
Pulsatile Odor Stimuli by Moth Antennal Lobe Neurons. *Front Neurosci* **15**, 739730
(2021).
62.   M. Paoli, G. C. Galizia, Olfactory coding in honeybees. *Cell Tissue Res* **383**, 35-58 (2021).
63.   C. Fonta, X. J. Sun, C. Masson, Morphology and Spatial-Distribution of Bee Antennal Lobe
Interneurons Responsive to Odors. *Chem Senses* **18**, 101-119 (1993).
64.   S. Kirschner *et al.*, Dual olfactory pathway in the honeybee, Apis mellifera. *J Comp Neurol*
499, 933-952 (2006).
65.   A. Hu, W. Zhang, Z. Wang, Functional feedback from mushroom bodies to antennal lobes
in the Drosophila olfactory pathway. *Proc Natl Acad Sci U S A* **107**, 10262-10267 (2010).
66.   D. Hattori *et al.*, Representations of Novelty and Familiarity in a Mushroom Body
Compartment. *Cell* **169**, 956-969.e917 (2017).
67.   T. Roeder, Tyramine and octopamine: ruling behavior and metabolism. *Annu Rev Entomol*
50, 447-477 (2005).
68.   B. H. Smith, C. N. Cook, Experimental psychology meets behavioral ecology: what
laboratory studies of learning polymorphisms mean for learning under natural conditions,
and vice versa. *J Neurogenet* **34**, 178-183 (2020).
69.   S. W. Cobey, D. R. Tarpy, J. Woyke, Standard methods for instrumental insemination of
Apis mellifera queens. *Journal of Apicultural Research* **52(4)**, 1-18 (2013).
70.   H. J. Ferguson, Cobey, S., Smith, B.H., Sensitivity to a change in reward is heritable in the
honey bee, Apis mellifera. *Animal Behaviour* **61**, 527-534 (2001).
71.   S. Cobey, Comparison studies of instrumentally inseminated and naturally mated honey
bee queens and factors affecting their performance. *Apidologie* **38**, 390-410 (2007).
72.   B. H. Smith, C. M. Burden, A proboscis extension response protocol for investigating
behavioral plasticity in insects: application to basic, biomedical, and agricultural research.
J Vis Exp, e51057 (2014).
73.   B. H. Smith, R. Menzel, The use of electromyogram recordings to quantify odorant
discrimination in the honey bee, Apis mellifera. *Journal of Insect Physiology* **35**, 369-375
(1989).

74.   G. J. Hunt, R. E. Page, Jr. , Linkage map of the honey bee, Apis mellifera, based on RAPD
markers. *Genetics* **139**, 1371-1382 (1995).
75.   O. Rueppell *et al.*, The genetic architecture of sucrose responsiveness in the honeybee (Apis
mellifera L.). *Genetics* **172**, 243-251 (2006).
76.   O. Rueppell *et al.*, The genetic architecture of the behavioral ontogeny of foraging in
honeybee workers. *Genetics* **167**, 1767-1779 (2004).
77.   R. Menzel, A. Heyne, C. Kinzel, B. Gerber, A. Fiala, Pharmacological dissociation between
the reinforcing, sensitizing, and response-releasing functions of reward in honeybee
classical conditioning. *Behavioral Neuroscience* **113**, 744-754 (1999).
78.   S. Tuzmen, J. Kiefer, S. Mousses, Validation of short interfering RNA knockdowns by
quantitative real-time PCR. *Methods Mol Biol* **353**, 177-203 (2007).
79.   M. C. Munoz-Torres *et al.*, Hymenoptera Genome Database: integrated community
resources for insect species of the order Hymenoptera. *Nucleic Acids Res* **39**, D658-662
(2011).
80.   H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**, 1754-1760 (2009).
81.   H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
2079 (2009).

Funding

National Institutes of Health NIGMS (R01GM113967)

- Brian H Smith

National Science Foundation CRCNS (2113179)

- Brian H Smith

National Science Foundation NeuroNex (1559632)

- Brian H Smith

Department of Energy (SC0021922)

- Brian H Smith

The funders had no role in study design, data collection and interpretation, or the decision to

submit the work for publication.