Predicting predator-prey interactions in terrestrial endotherms using random forest

John Llewelyn[1,2], Giovanni Strona[3], Christopher R. Dickman[4], Aaron C. Greenville[5,6,7], Glenda M. Wardle[5], Michael S. Y. Lee[8,9], Seamus Doherty[1,2], Farzin Shabani[10], Frédérik Saltré[1,2], Corey J. A. Bradshaw[1,2]

[1]Global Ecology | *Partuyarta Ngadluku Wardli Kuu*, College of Science and Engineering, Flinders University, GPO Box 2100, Adelaide, South Australia 5001, Australia

[2]ARC Centre of Excellence for Australian Biodiversity and Heritage. EpicAustralia.org.au

[3]Faculty of Biological and Environmental Sciences, University of Helsinki, Viikinkaari 1, Biocentre 3, 00790, Helsinki, Finland

[4]Desert Ecology Research Group, School of Life and Environmental Sciences, University of Sydney, New South Wales 2006, Australia

[5]School of Life and Environmental Sciences, University of Sydney, Sydney, New South Wales 2006, Australia

[6]School of Life Sciences, University of Technology Sydney, Sydney, New South Wales 2007, Australia

[7]National Environmental Science Program, Threatened Species Recovery Hub, University of Sydney, Sydney, New South Wales 2006, Australia

[8]College of Science and Engineering, Flinders University, GPO Box 2100, Adelaide, SA 5001, Australia

[9]Earth Sciences Section, South Australian Museum, North Terrace, Adelaide, SA 5000, Australia

[10]Department of Biological and Environmental Sciences, College of Arts and Sciences, Qatar University, PO Box 2713, Doha, Qatar

Corresponding Author: John Llewelyn (johnsllewelyn@gmail.com)

Predicting predator-prey interactions in terrestrial endotherms using random forest

Abstract

Species interactions play a fundamental role in ecosystems. However, few ecological communities have complete data describing such interactions, which is an obstacle to understanding how ecosystems function and respond to perturbations. Because it is often impractical to collect empirical data for all interactions in a community, various methods have been developed to infer interactions. Machine learning is increasingly being used for making interaction predictions, with random forest being one of the most frequently used of these methods. However, performance of random forest in inferring predator-prey interactions in terrestrial vertebrates and its sensitivity to training data quality remain untested. We examined predator-prey interactions in two diverse, primarily terrestrial vertebrate classes: birds and mammals. Combining data from a global interaction dataset and a specific community (Simpson Desert, Australia), we tested how well random forest predicted predator-prey interactions for mammals and birds using species' ecomorphological and phylogenetic traits. We also tested how variation in training data quality—manipulated by removing records and switching interaction records to non-interactions—affected model performance. We found that random forest could predict predator-prey interactions for birds and mammals using ecomorphological or phylogenetic traits, correctly predicting up to 88% and 67% of interactions and non-interactions in the global and community-specific datasets, respectively. These predictions were accurate even when there were no records in the training data for focal species. In contrast, false non-interactions for focal predators in training data strongly degraded model performance. Our results demonstrate that random forest can

25   identify predator-prey interactions for birds and mammals that have few or no interaction

26   records. Furthermore, our study provides guidance on how to prepare training data to

27   optimise machine-learning classifiers for predicting species interactions, which could help

28   ecologists (*i*) address knowledge gaps and explore network-related questions in data-poor

29   situations, and (*ii*) predict interactions for range-expanding species.

Introduction

Species interactions are essential for maintaining biodiversity (Thompson 1999) because they

determine how energy and nutrients flow among organisms (Traill et al. 2010, Allan et al.

2021) and affect the distribution, survival, and abundance of species (Van der Putten et al.

2010). Species interactions also determine how ecological disturbances—such as those

involving invasive species, habitat modification, and climate change—reverberate through

ecological communities (Tylianakis et al. 2008, Blois et al. 2013). This is because

environmental changes that directly alter one or a subset of species in a community can have

flow-on effects for other species via interaction networks (Kaneryd et al. 2012, Morton et al.

2022). Given the need to predict and mitigate human-driven biodiversity declines, interest in

modelling interaction networks for biological communities is increasing (Bohan et al. 2017,

Strydom et al. 2021; Strona and Bradshaw 2022). However, most species interactions in most

ecological communities have not been documented (Hortal et al. 2015, Jordano 2016, Caron

et al. 2022). Indeed, it is usually logistically impossible to measure how all species interact in

a community—even when only one type of interaction (e.g., predator-prey) is considered

(Brousseau et al. 2018, Strona 2022). In addition to the challenge of identifying long-

established interactions, the rapid and widespread translocation of species across the planet

due to migratory responses to climate change and movement by humans means we also need

to identify interactions that could occur between species that previously did not co-occur, and

therefore had not previously had the opportunity to interact (Smith and Phillips 2006,

Valdovinos et al. 2018).

To overcome gaps in species interaction data, various methods of inferring interactions

using species traits/proxies can be applied, such as quantile-regression trophic niche-space

modelling (Gravel et al. 2013, Llewelyn et al. 2021), Bayesian linear modelling (Caron et al.

2022), and several machine-learning approaches (e.g., *K* nearest-neighbour—(Desjardins-

Proulx et al. 2017, Pichler et al. 2020), neural networks/deep learning methods—(Fricke et al.

2022), random forests—(Sydenham et al. 2021), and ensemble methods—(Poisot 2022)). Of

these, random forest is a popular method due to its ability to learn nonlinear interactions

between predictor variables and predict interactions accurately (Desjardins-Proulx et al.

2017, Laigle et al. 2018, Sydenham et al. 2021). Random forests have been used to predict

various interaction types, including plant-pollinator (Ornai and Keasar 2020, Sydenham et al.

2021), parasite-host (Kotula et al. 2020), and predator-prey interactions (Desjardins-Proulx et

al. 2017, Parravicini et al. 2020). They have also been used to identify which species' traits

are most important for predicting whether species interact (Dellinger et al. 2019, Klomberg et

al. 2022). Using training data consisting of interacting and non-interacting species, random

forests can learn how ecomorphological traits, phylogenetic position, and/or taxonomy (as a

proxy for phylogenetic position; Laigle et al. 2018) relate to which species interact. However,

questions remain regarding the ability of random forests to predict different types of

interactions in different taxonomic groups, which species' data/traits are best for inferring

interactions, and how to fine-tune training datasets (Poisot 2022). Furthermore, investigating

ways to optimise training data for predicting species interactions using random forest will

likely help guide procedures for optimising other machine learning methods.

Identifying predator-prey interactions is central to community ecology because these

interactions influence both top-down and bottom-up processes such as energy transfer and

extinction cascades (Sinclair et al. 2003). Random forests have been applied to predict such

interactions, but these applications have largely been limited to fish communities (Parravicini

et al. 2020, Strona et al. 2021) and soil invertebrates (Desjardins-Proulx et al. 2017, Laigle et

al. 2018). Indeed, the performance of random forests for inferring predator-prey interactions

among terrestrial vertebrates has not been assessed despite this group including many species

of high conservation concern (Tilman et al. 2017, Cox et al. 2022) and of important

ecological function (e.g., ecological engineers; Samson and Knopf 1996).

If we are to model terrestrial ecosystems realistically to predict and minimise the risk of

extinction cascades, reliable inference of trophic interactions is essential. However, trophic-

related traits and network structure differ among terrestrial vertebrates, fish, and invertebrates

(Liem 1984, Liem 1990, Shurin and Gruner 2006, Miller-ter Kuile et al. 2022), suggesting

traits and methods (such as random forest and other machine learning techniques) that

effectively predict interactions in fish and invertebrates might not necessarily work well for

terrestrial vertebrates. These vertebrates are diverse structurally, phylogenetically, and

functionally, potentially making the 'rules' determining interactions complex and difficult for

machine-learning classifiers to identify (whereas just one trait—body size—largely

determines trophic interactions in fish due to their gape limitation; Gravel et al. 2013,

Morales-Castilla et al. 2015). Furthermore, interaction data for many terrestrial vertebrate

species have been recorded opportunistically or not at all (although detailed diet data do exist

for some species; Pringle and Hutchinson 2020), potentially leading to sparse (i.e., missing

most prey for most predators) and taxonomically biased datasets for training and testing

random forests. Having sparse interaction records for predators in training data is problematic

because machine learning classifiers such as random forests need to be trained with data that

include both interacting (e.g., predator and prey) and non-interacting species pairs, with non-

interacting species pairs usually assigned by assuming that undocumented interactions imply

no interaction (i.e., pseudo-non-interactions generated from presence-only data; Pichler et al.

2020). When applied to incomplete (sparse) datasets, this process can identify undocumented

but truly interacting species pairs as non-interacting pairs (but see Strona et al. 2021).

Importantly, such 'false non-interactions' in training data likely reduce the reliability of

interactions and non-interactions inferred from random forests and other classifiers.

Similarly, random forests and other classifiers trained on datasets that are taxonomically

biased might perform poorly when making predictions for species not represented in the

training data (Strydom et al. 2022).

We designed analyses to test whether random forest could accurately predict predator-prey

interactions within and between two diverse terrestrial vertebrate classes—birds and

mammals—using both a global (*GloBI*) and a single-community (Simpson Desert, Australia)

predator-prey interaction dataset. We also assessed (*i*) the importance of ecomorphological

traits and phylogenetically based latent traits (i.e., as proxies for phylogenetically conserved

traits associated with trophic interactions; Benandi et al. 2022) for inference accuracy, (*ii*)

whether prediction accuracy improved by removing predators from the training dataset that

had few interaction records (i.e., those with scarce diet records and were, therefore, prone to

generating false non-interactions), and (*iii*) the sensitivity of model performance to the

quality of the training data. To test sensitivity to the quality of training data, we compared

two scenarios that could occur as a result of sparse training data: (a) a reduction in the

number of interaction and non-interaction records used to train the models, and (b) a

reduction in the number of interaction records and an increase in the number of false non-

interaction records used to train the models. We also investigated how these changes affected

model performance when they were restricted to different components of the training dataset

(i.e., testing the effect of taxonomic bias on performance).

Our results confirm that: (*i*) random forests can predict interactions in birds and mammals,

(*ii*) models trained on interaction datasets including either ecomorphological traits or

phylogenetic position (or both) can perform well, (*iii*) removing predators from the training

data that have few recorded interactions therein can improve prediction accuracy, (*iv*) models

can identify the prey of predators that have no interaction records in the training data, and (*v*)

model performance is especially sensitive to false non-interactions in the training data that

involve records of predatory species for which predictions are being made (i.e., focal

predators). Our results highlight the unrealised potential for the application of random forests

for predicting interactions among terrestrial vertebrates. Furthermore, they show to which

components of the training data random forests are most sensitive, thereby identifying the

most efficient ways to increase accuracy via fine-tuning of training data—highlighting steps

that would likely also improve species-interaction inference by machine learning classifiers

more generally.

Methods

We used two interaction datasets combined with ecomorphological and phylogenetic

information to test the performance of random forests for predicting predator-prey

interactions in birds and mammals (Supporting information Figure S1 shows flowchart of

dataset preparation). We also tested how the performance of these models was influenced by

the type of species data used, the filtering of training data by removing predators with few

interaction records in that data, and the quality of the training data. The two interaction

datasets were: (1) a dataset from the ***Glob**al **B**iotic **I**nteractions database*

(globalbioticinteractions.org; Poelen et al. 2014) supplemented with interaction data from

diet studies done in Australia (Llewelyn 2022), and (2) an ecosystem-specific dataset from

the Simpson Desert in Australia that focuses on seven predators for which detailed dietary

information is available (Llewelyn 2022). We focused on birds and mammals rather than on

all tetrapods due to the availability of detailed trait (Wilman et al. 2014) and phylogenetic

(vertlife.org) information for birds and mammals (detailed trait databases for reptiles and

amphibians are far from complete), thereby providing a wide range of traits for predicting

interactions.

Datasets

Global data

We extracted all predator-prey interactions involving birds and mammals from *GloBI*

(accessed 29/05/2020 and 26/06/2020 for birds and mammals as consumers, respectively).

However, *GloBI* contains few records of species found in Oceania (including Australia)

relative to other regions of the world: of the 3329 unique species combinations in the records

we extracted from *GloBI*, only 109 involved a predator and prey species both found in

Oceania. Supervised learners, including random forests, need to be trained on data that

capture the variation present in the species for which predictions are being made, and so

geographic and taxonomic biases in training data could reduce prediction accuracy. We

therefore added interactions involving Simpson Desert species and their relatives to the

GloBI data (Supporting information). We added data for these species by merging two

additional datasets with *GloBI*: (*i*) predator-prey records from the Simpson Desert that did not

involve the seven focal predators (focal predator = species for which we made predictions),

and (*ii*) interactions of the focal Simpson Desert predators with non-Simpson Desert prey

elsewhere in Australia (i.e., from diet studies of the focal predators outside the Simpson

Desert; no interactions between the focal predators and Simpson Desert species were

included in the global/training data). In both cases, we built these datasets by aggregating

predator-prey records from diet studies (Supporting information). We then combined these

additional interaction datasets with the *GloBI* data (adding 363 records) to create an enhanced

global interaction dataset (Llewelyn 2022).

To predict species interactions using random forests, we required a training dataset that

included interacting and non-interacting pairs of species. Like most interaction datasets, our

global dataset did not include non-interactions i.e., since relationships are based on presence-

only data. We therefore generated pseudo-non-interactions ('non-interactions' henceforth) by

taking all possible two-species combinations from the species in our global interaction

dataset and removing those combinations documented as 'interacting' (supporting

information). We also removed combinations where either the 'predator' does not consume

vertebrates (e.g., strictly herbivorous species could not be potential predators) or the

combination involved species not found on the same continent (using the `rgbif` package in

R; Chamberlain et al. 2022). We removed these non-interactions because they would not help

models predict which sympatric species known predators preyed upon (i.e., the type of

interactions we aimed to predict in the Simpson Desert). Although the non-interactions we

generated would have included many sympatric, truly non-interacting species combinations,

they would have also included false non-interactions (species combination assigned as not

interacting, but that do indeed interact) as well as allopatric species pairs that would interact

if they were sympatric.

Random forests use explanatory variables such as species' traits to learn and predict

associations (Desjardins-Proulx et al. 2017). We therefore added trait data to all the predator

and prey species in the interaction and non-interaction datasets (Supporting information).

These traits included: body mass, time of activity (whether the species displays nocturnal,

diurnal, and/or crepuscular activity), ability to fly, resource groups consumed (plants,

vertebrates, and/or invertebrates), and percentage use of different habitat strata (8 habitat

categories for birds and 5 for mammals; Wilman et al. 2014; Supporting information). We

added phylogenetic information by calculating eigenvectors describing a species' position in

the phylogeny (eigenvector mapping; Guénard et al. 2013), a potential proxy for explanatory

variables which are phylogenetically conserved. We extracted phylogenies from

VertLife.org, and used the `MPSEM` (Guénard and Legendre 2022) and `ape` (Paradis and

Schliep 2019 p. 0) R packages to handle phylogenetic data and extract eigenvectors.

We also identified which species pairs in the non-interaction data included a potential prey

species from within the predator's preferred prey-size range (Supporting information). We

did this so the ratio of non-interaction records from inside *versus* outside the preferred size

range in training datasets could be adjusted to optimise performance of random forests. This

is analogous to optimising species distribution models by adjusting the ratio of pseudo-

absences sampled from among *versus* outside observed occurrences (Barbet-Massin et al.

2012). To identify a preferred size range, we log-transformed ($\log_e$) predator and prey body

masses to linearise body-size relationships (Gravel et al. 2013). We then resampled the

interaction dataset to create 100 training and testing dataset pairs using all 3692 interaction

records in each of the 100 pairs (75:25 interaction records in training:testing datasets).

To each training set we fitted a series of upper and lower quantile regressions capturing

the middle 0.99 to 0.8 quantiles at 0.01 quantile increments. We included a prey mass ×

predator taxonomic class interaction in the regression models to allow different body-size

relationships for predatory birds and mammals (Llewelyn et al. 2021). We then combined

each of the 100 testing sets with an equal number of species pairs randomly sampled from the

non-interaction set (also with body mass $\log_e$-transformed), and applied the quantile

regression models (i.e., upper and lower quantile regressions combinations) built in the

previous step to the testing sets. Species combinations in the testing sets that included a

potential prey species from within the upper and lower quantile ranges derived from the

regressions were predicted as prey, and those outside these limits were excluded as prey.

We assessed model performance by comparing predictions to documented interactions (i.e., asking whether the quantile regression models correctly identified which combinations came from the interaction and non-interaction datasets), and used these results to calculate the true skill statistic to identify the optimal quantile regression model for assigning predator-prey interactions. The true skill statistic is a metric that can effectively evaluate the performance of classifiers such as those predicting species interactions (Poisot 2022). A true skill statistic = 1 indicates a model that perfectly predicts interactions and non-interactions, whereas when it is 0, the model performs no better than assigning links randomly. The model capturing the middle 0.84 quantiles performed best (true skill statistic = 0.26), so we used that model to classify species pairs in the non-interaction dataset as those that included a potential prey from within the predator's preferred size range and those that included a potential prey from outside the predator's preferred size range.

Simpson Desert testing data

It is unclear how the accuracy of random forests trained and tested on data including species from many different communities (such as was the case for our global datasets) translates to model performance when applied to single ecological communities (Parravicini et al. 2020). We therefore tested model performance when applied to seven common predators from the Simpson Desert, a community we chose because it has been intensively studied for over 30 years and has detailed information on species composition and the diets of some predators. We used the species documented at established field sites in the Simpson Desert as the species assemblage (i.e., potential prey), rather than all species documented in the Simpson Desert bioregion, because the former includes sympatric species that have the potential to interact whereas the latter would likely include species that seldom have the opportunity to interact.

The Simpson Desert bird and mammal assemblage from these sites includes 64 vertebrate-

consuming bird and mammal species (49 birds and 15 mammals). However, the diets of

many of these species have not been studied in detail. When testing the performance of

random forests for predicting predator-prey interactions, we therefore restricted the Simpson

Desert testing dataset to seven predatory species with detailed dietary information from

studies both in the Simpson Desert and other arid regions in Australia. These seven focal

predators included four birds—*Aquila audax*, *Falco berigora*, *Hamirostra melanosternon*,

Tyto alba, and three mammals—*Canis dingo*, *Felis catus*, *Vulpes vulpes*. *Felis catus*

(domestic cats) and *Vulpes vulpes* (red fox) are recent introductions to Australia (since

European colonisation in 1788), *Canis dingo* (dingo) was brought to Australia by people >

3000 years ago (Fillios et al. 2012), and the remaining focal predators naturally occur in

Australia.

To construct the dataset, we aggregated data from published and unpublished diet studies

(excluding species consumed as carrion) of the focal predators in the Simpson Desert

(Supporting information; Llewelyn 2022). We also included records from diet studies done

outside the Simpson Desert involving species pairs found in the Simpson Desert (i.e., we

assumed that if trophic interactions occurred between the species in other communities, they

would also occur in the Simpson Desert; Llewelyn 2022), and records involving the focal

predators and Simpson Desert species in the global interaction dataset (we removed these

records from the global dataset). None of the studies we used to build the Simpson Desert

interaction dataset reported non-interactions. We therefore generated non-interactions by

assuming any species pairs involving the focal predator and the species recorded from the

Simpson Desert field sites (157 bird and mammal species) that have not been documented as

predator and prey are not predator and prey. We added the same trait and phylogenetic

information to the Simpson Desert data as described for the global dataset. However, we did

not include a column indicating if non-interacting species pairs in the Simpson Desert data

involved potential prey from within the predator's preferred size range because we only used

the Simpson Desert dataset to test model performance (i.e., we did not need to optimise the

sampling of these non-interactions because they were not used for training random forests).

Modelling

We used the `ranger` package (Wright and Ziegler 2017) in R and the Flinders University

high performance computer (Flinders 2021) to apply random forests using the model-based

recursive partitioning algorithm (Breiman 2001). In the first phase, we optimised the random

forest models and tested how performance was affected by (a) the number and choice of

variables included in the random forests, and (b) removing predators that had few

documented interactions in the training data (i.e., those whose diet is likely not adequately

captured in the data). In the second phase, we tested how changes to the quality of the

training dataset—including removing records and increasing the number of false non-

interactions—influenced model performance (Supporting information shows flowchart of

modelling steps and what different models were used for). When optimising models, we

assessed model performance by comparing predictions to documented interactions and used

these results to calculate the true skill statistic (we also calculated other performance metrics

to confirm our results; see Supporting information). A true skill statistic of 1 indicates a

model that perfectly predicts interactions and non-interactions, a score > 0.75 is 'excellent',

between 0.4 to 0.75 is 'fair to good', and < 0.4 indicates 'poor' performance (Fleiss et al.

2003).

1. Model optimisation and variables

We used the global dataset and cross-validation to optimise training datasets and model

parameters, and we compared performance of models trained using: (*i*) only

ecomorphological traits, (*ii*) only phylogenetic eigenvectors, or (*iii*) both types of data

(Supporting information). We also compared performance of models that incorporated many

versus few explanatory variables/traits to test whether more variables resulted in higher

accuracy or overfitting. There were 42 variables in the many-variable ecomorphological

model (21 each for predator and prey; Llewelyn 2022) because there were 21

ecomorphological variables in the databases we built (Supporting information). To be

consistent, we used 42 phylogenetic eigenvectors in the many-variable phylogenetic model

(first 21 phylogenetic eigenvectors for predator and prey). In the few-variable models, we

included 10 variables (a number we chose arbitrarily that is substantially smaller than the 42

variables in the many-variable models). These 10 variables were the most important

ecomorphological or phylogenetic variables (5 each for predator and prey) for predicting

interactions, as determined by permutation, which measures the impact of changing a

variable's value on model accuracy. To identify the most important variables, we sequentially

removed the least-important variable and repeated the permutation and removal steps until 10

variables remained (retained variables for each model listed in Supporting information;

Llewelyn 2022). Although a variable's importance in predicting interactions does not imply

causation (Arif and MacNeil 2022), knowing which traits are most important for predicting

interactions can help researchers avoid spending time and resources collecting data on traits

that are uninformative. The many- and few-variable models that incorporated both types of

data simply combined the variables from the respective ecomorphological and phylogenetic

models (i.e., either 84 or 20 variables in total).

We also tested whether performance was enhanced or eroded by removing records from

the global dataset involving predators with few (< 5) interaction records. We posited that

predators that had only a small proportion of their diet documented in the dataset could bias

results and lead to the generation of many false non-interactions because, for example,

models might be unduly influenced by predators that had not been studied in detail – rather

than matching of predator and prey traits. Removing records involving these predators

reduced the number of interactions from 3,692 in the original global dataset to 3,424 in the

modified global dataset, and reduced the number of non-interactions from 454,924 to 89,184.

To optimise training datasets and model parameters, we divided the global dataset's

interactions (original and modified data) into training and testing datasets (ratio: 75:25

training:testing), and combined them with non-interaction data. The number of non-

interaction records added to the testing datasets equalled the number of interaction records in

that dataset, whereas we varied the number of non-interactions in the training datasets to

identify the optimal (highest true skill statistic) ratio. We replicated the process of generating

training and testing datasets each time we applied a random forest in the parameter-

optimization and cross-validation steps described below.

We weighted non-interactions such that their combined weight was equal to that of the

observed interactions because models built with weighted non-interactions performed slightly

better than those built using unweighted non-interactions (Supporting information; similar to

results from species distribution modelling; Barbet-Massin et al. 2012). These weightings

determine the probability an interaction/non-interaction is selected in the bootstrap samples

for the trees built from the training datasets. We optimised the training datasets and random

forests in terms of (*i*) number of non-interactions in the training data (see above), (*ii*) the ratio

of non-interactions from inside *versus* outside preferred size range in the training data, (*iii*)

the probability threshold for assigning interactions—because random forests can estimate

probabilities for binary responses such as interact *versus* not interact (Malley et al. 2012), (*iv*)

the number of variables randomly sampled as candidates at each split in the trees, (*v*) the

number of decision trees in the random forest, and (*vi*) the maximum depth of the trees.

We optimised these parameters by iterating through values for each parameter

individually, running 5 random forests at each value (on 5 resampled training and testing

datasets) and calculating the mean true skill statistic for models at each value. We identified

the parameter values that gave models with the highest true skill statistic. We separately

optimised the 12 models that differed in terms of included variables (i.e., the many- *versus*

few-variable ecomorphological, phylogenetic, and combined data models) and whether the

original or modified global dataset was used (Supporting information shows optimised

parameter sets). We then compared performance of these optimised models by applying them

to 100 training and testing datasets generated using the global dataset cross-validation design

described above, and calculated the true skill statistic for each of these 100 cases.

After comparing performance of the twelve models with the global dataset, we tested their

performance when trained on the global dataset (with hyper-parameters set to optimise

performance, as identified in the previous step) and applied to the Simpson Desert data

(Supporting information). Again, we trained and applied 100 random forests for each of the

twelve models, and assessed their performance with the true skill statistic. We resampled

non-interactions for the training dataset in these 100 analyses because there were more non-

interactions than needed, whereas we used all available interaction records in the global

datasets (3,692 and 3,424 for the original and modified global datasets, respectively) in each

of the 100 training datasets.

2. Quality of training dataset and model performance

To test the influence of the size of the training dataset, taxonomic coverage, and prevalence

of false non-interactions (true interactions classified as non-interactions) on model

performance, we reran the analyses with the optimised model that performed best on the

Simpson Desert data (i.e., the many ecomorphological and phylogenetic variable model

trained on the modified global dataset; Figure 1f; Supporting information) but varied the

number of records, taxonomic coverage, and prevalence of false non-interactions in the

training datasets (Supporting information).

We deliberately reduced the quality of training data in eight ways to test specific

sensitivity hypotheses. We modified the entire training dataset by (*i*) randomly removing

records and (*ii*) randomly switching interaction records to non-interactions (false non-

interactions). We hypothesised that model performance would be sensitive to removal of

training data because having more records helps the models learn associations between

species traits and trophic links, and we hypothesised that models would be sensitive to false

non-interactions because random forests would learn the wrong association when trained with

erroneous data. We also tested what impact data removal and false non-interactions had when

they were restricted to subsets of the training data that included either: (*iii*, *iv*) focal prey, (*v*,

vi) focal predators, or (*vii*, *viii*) interactions where neither prey nor predators were

focal/Simpson Desert species. We hypothesised that model performance would be especially

sensitive to changes involving focal predators and prey because these records involve the

species for which predictions are made. Here, the associations between these species' traits

and trophic interactions should be especially important for the model to learn. Furthermore,

we hypothesised that model performance would be sensitive, albeit less so, to removing

records and increasing the proportion of false non-interactions in non-Simpson Desert (non-

focal) interactions because these interactions teach the random forests trait-trophic

associations that also apply to Simpson Desert species. Assessing the sensitivity to changes in

the quality of training data involving the focal species is important because it assesses the

suitability of random forests for inferring interactions of species that have few/no interaction

records, a common phenomenon.

For each of the eight ways training dataset quality was reduced, we made these changes at

1% increments relative to the initial number of records in the modified group (e.g., 1%

increments of the total number of interactions involving the focal predator), generating 10

training datasets at each increment. We trained the random forests on these reduced-quality

datasets, applied them to the Simpson Desert data, and calculated true skill statistics 10 times

at each increment. We plotted results to make visual comparison of the sensitivity of model

performance to each of the eight types of artificial variation in the quality of training data,

and we compared model performance when all relevant data were modified to quantify the

impacts of each type of modification.

We also examined the mechanisms underlying changes in model performance by testing

how removing records or increasing false non-interactions for the focal predator influenced

the: (*i*) relative probability (or suitability) of interacting assigned by the random forest to each

potential prey for each predator (i.e., asking whether changes to the training dataset affected

which species were identified as the most likely prey), and (*ii*) mean probability assigned to

potential prey for each predator (i.e., asking whether changes to the training dataset cause an

overall increase or decrease in the probabilities assigned to potential prey). If lower model

performance reflects a systematic shift in all the probabilities (upward or downward), rather

than a change in which potential prey are identified as the most likely, the models could still

be used to predict a predator's prey because the rank order of probability of different prey

remains similar irrespective of absolute model performance and absolute probability.

To test for changes in relative prey suitability, we measured the correlation (Pearson's *r*)

between prey probabilities assigned by models trained on reduced-quality training datasets

with that assigned by models trained on the full training dataset (Supporting information). To

test for changes in mean probability assigned to potential prey overall, we subtracted the

mean probability assigned to potential prey for each predator as calculated by the full

training-data models from that calculated by the models trained on the reduced-quality

training datasets (Supporting information). For both analyses, we reduced the quality of the

training dataset at 1% increments (10 replicate training datasets/models at each increment)

and plotted the results to compare visually. We hypothesised that removing records and

including more false non-interactions in the training dataset would disrupt the assignment of

relative suitability and reduce assigned probabilities overall.

Results

Variables and model performance

Random forests trained and tested on the global data had mean true skill statistics ranging

from 0.53 to 0.77 (Figure 1a,c,e,g), indicating 'good' to 'excellent' performance (Fleiss et al.

2003). These models tended to perform better when we used the original global interaction

dataset rather than the modified dataset that had predators with few prey records removed

(mean difference in true skill statistic between models = 0.14; Figure 1a and Figure 1c *versus*

1e and 1g). Using few or many variables in the phylogenetic or

phylogenetic/ecomorphological variable models had little effect on performance when

applied to the global datasets (difference in mean true skill statistic < 0.02 in all cases; Figure

1), but there was a slight decrease in performance associated with using few variables in the

ecomorphological-only models (difference in mean true skill statistic = 0.08 and 0.06

between the many- and few-variable ecomorphological models when the original or modified

global dataset was used, respectively; Figure 1a *versus* 1c, Figure 1e *versus* 1g).

The models trained on the global dataset and applied to the Simpson Desert data yielded

mean true skill statistics between 0.25 and 0.45, indicating 'poor' to 'fair' performance

(Fleiss et al. 2003; Figure 1). Those that used both phylogenetic and ecomorphological

variables performed well when trained on the modified global dataset, yielding mean true

skill statistics of 0.43 ± 0.02 (mean $\pm$ standard deviation) and 0.45 ± 0.02 when we used few

or many variables, respectively (Figure 1f,h). Using few *versus* many variables did not

strongly affect the performance of the phylogenetic-only or phylogenetic- and

ecomorphological-variable models (difference in mean true skill statistic < 0.07 in all four

cases; Figure 1). In contrast, the few-variable ecomorphological models did not perform as

well as their many-variable counterparts (difference in mean true skill statistic > 0.13

between the many- and few-variable ecomorphological models when using either the original

or modified global datasets Figure 1b *versus* 1d, Figure 1f *versus* 1h).

Using the model that performed best on the Simpson Desert data (i.e., the many-variable

model with phylogenetic and ecomorphological traits trained on the modified global dataset;

Figure 1f), performance predicting interaction for recently introduced predators—cats and

foxes—was better than that for native predators (true skill statistic $= 0.56 \pm 0.03$ and $0.41 \pm$

0.02 for recently introduced and native predators, respectively; Figure 2).

Effects of training dataset size, false non-interactions, and taxonomic coverage

We altered the global (modified) training dataset by randomly removing records and

changing interaction records to false non-interactions, applying these changes both to the

entire training dataset as well as to three subsets: (*i*) records involving the focal prey (i.e.,

species found in the Simpson Desert), (*ii*) records involving the focal predators (i.e., the 7

Simpson Desert predators for which we predicted interactions), and (*iii*) records involving

species not found in the Simpson Desert. Removing records reduced performance when

modifying either the entire training dataset (Figure 3a; true skill statistic reduced to 0 when

all records removed because random forests cannot run without training data) or the focal-

predator component only (Figure 3e; true skill statistic reduced to 0.24 ± 0.01 [mean ±

standard deviation] when all focal-predator records were removed). However, these effects

became pronounced only when > 80% of the interaction and non-interaction records were

removed (Figure 3a,e), which equates to removing > 15,750 or 1,240 records from the entire

or focal-predator components of the training datasets, respectively. Removing records

involving the focal prey or non-Simpson Desert species had only weak or no effect on model

performance (Figure 3c,g; true skill static = 0.36 ± 0.02 and 0.46 ± 0.01 when removing all

focal prey or all non-Simpson Desert species, respectively).

The effect of false non-interactions on model performance was stronger than that of

removing records. False non-interactions assigned at random to interaction records in either

the entire training dataset or the component of the training dataset involving focal predators

degraded model performance more rapidly (Figure 3b,f; true skill static reduced to 0 and 0.04

± 0.01 for models when changing all records or all records involving focal predators to non-

interactions, respectively). These declines were triggered when changing > 40% of the

interaction records in the relevant group to false non-interactions (Figure 3b,f), which equates

to 1,370 records in the entire training-dataset analysis and 235 in the focal-predator analysis.

False non-interactions in the training data involving the focal prey (Figure 3d) or non-

Simpson Desert species (Figure 3h) caused slight declines in model performance (true skill

static reduced to 0.36 ± 0.03 and 0.34 ± 0.02 when changing all interactions records to false

non-interactions for focal prey species and non-Simpson Desert species, respectively).

Focal-predator training data and the mechanisms underlying changes in model

performance

The correlation between probabilities assigned to potential prey by models trained on the full

versus record-removed global datasets showed a gradual decrease as more interactions

involving the focal predator were removed from the training data, and this decline became

more rapid when removing > 80% of records (Figure 4a). The correlation between these

probabilities reduced to 0.59 ± 0.02 when all records involving focal predators were removed

from the training dataset, indicating that models identified many of the same species as prey

even when there were no records involving the focal predator in the training data. The mean

probability assigned to potential prey for each predator was reduced by removing records of

the focal predator from the training data and, like the correlation results, this change was

small until 80% of records were removed (Figure 4c). After removing all records, the

probabilities assigned to potential prey were on average 0.17 ± 0.01 lower compared to those

assigned by models trained on the full dataset.

Both the correlation and change in mean probability assigned to potential prey were more

sensitive to false non-interactions involving the focal predators than to removing their records

from the training dataset, showing more rapid declines (Figure 4a *versus* 4b, Figure 4c *versus*

4d). When changing all interactions involving the focal predators in the training data to non-

interactions, the correlation between probabilities assigned to prey decreased to 0.27 ± 0.03

(Figure 4b) and the mean probabilities assigned to potential prey dropped by 0.31 ± 0.01

(Figure 4d).

Discussion

Our results demonstrate that random forests can predict the prey of predatory birds and

mammals even when few or no interaction records involving the focal predator are included

in training data (Figure 4a). Applying random forests to predict predator-prey interactions for

these predators (and terrestrial vertebrates more generally) to facilitate network modelling of

terrestrial vertebrate systems is therefore supported (Brousseau et al. 2018). However, our

analyses highlight that caution is needed in applying the random forest and other machine-

learning approaches depending on the quality, quantity, and type of training data available, and in the face of these limitations we suggest that filtering training data could improve model performance (Figure 5).

We found that random forests could predict bird and mammal interactions in both our global and ecosystem-specific datasets (including for native and recently introduced predators). Models based on ecomorphological, phylogenetic, or both types of traits, and those using few or many traits, could perform similarly well (Figure 1)—consistent with previous studies showing that either data type can be used for predicting interactions in other taxa (Faisal et al. 2010, Morales-Castilla et al. 2015, Brousseau et al. 2018, Pomeranz et al. 2019, Elmasri et al. 2020) and that few traits/variables are required to make accurate predictions (Desjardins-Proulx et al. 2017). Because phylogenetic information is available for most terrestrial vertebrates (e.g., VertLife.org), ecological/morphological trait databases are increasingly common and comprehensive (Jones et al. 2009, Oliveira et al. 2017, e.g., Herberstein et al. 2022), and interaction data for many species are available from open-source databases (e.g., globalbioticinteractions.org; web-of-life.es) and publications, random forests and other machine learning methods can now easily be applied to such data to predict predator-prey interactions for most vertebrate communities, and aid in the ecological network and invasive species modelling required to guide conservation decisions.

We found that model performance was robust to the random removal of records from the training dataset. In fact, we had to remove > 80% of records (>15,750 individual records) before an obvious change in performance occurred (Figure 3a). Thus, models that use large training datasets are unlikely to be sensitive to random additions or removal of interaction and non-interaction records.

The removal of records involving focal predators only caused a prominent decline in model performance when we removed > 80% (Figure 3e) of records in that group, but this

translates to fewer (1,240) records in total needing to be removed (as compared to the random

removal of records). Conversely, the complete removal of records involving focal-prey

species or non-focal species had little impact on model performance (Figure 3c,g). Thus,

declines in model performance in the random-removal analyses were largely driven by the

loss of focal predator records. This reduction in model performance following removal of

focal-predator records was due to every prey species being assigned consistently lower

probabilities and a slight shuffling of prey preference (Figure 4a,c). Because *relative* prey

preferences were largely maintained even when all training records involving the focal

predator were removed from training data (mean Pearson $r = 0.59$; Figure 4a), our results

suggest prey species could be predicted for predators that do not have any interaction data,

which is likely the case for many understudied, rare, or extinct species. The ability to predict

trophic interactions for extinct species accurately has important applications for

understanding past extinction events (Ripple and Van Valkenburgh 2010, Pires et al. 2015,

Llewelyn et al. 2021), while the capacity to predict interactions for rare species could help

identify cryptic factors that threaten their persistence (Smith and Phillips 2006, Doherty et al.

2016). However, additional work to assess the requirements of training data for predicting

interactions for species without interaction records is still required. For example, it is unclear

how phylogenetically and/or ecologically similar species in the training data need to be to the

focal species for models to be able to predict trophic interactions accurately.

In contrast to removal, false non-interactions in the training data reduced model

performance more rapidly, triggering lower performance when > 40% of the interaction

records were changed to non-interactions. This sensitivity highlights an important issue: most

studies generate non-interactions by assuming that undocumented interactions imply true

non-interaction; however, this assumption could generate many false non-interactions,

especially if interaction records are sparse (i.e., if many of the predators in the training data

have only a small proportion of their interactions documented). To avoid generating false

non-interactions, one can restrict non-interactions in training datasets to species combinations

where the predator is too small to eat the potential prey and/or the potential predator comes

from a low trophic level (Strona et al. 2021). However, applying this approach could reduce

the capacity of random forests to identify unsuitable prey from within a predator's preferred

prey-size range or for high trophic-level species. Indeed, the optimal ratio of non-interactions

sampled from within *versus* outside preferred size range in the model that performed best on

our Simpson Desert data (i.e., the many-variable, both data-types model) was 5:2 (Supporting

information), suggesting that non-interactions of suitably-sized potential prey should be

included in training data.

Alternatively, filtering training data by removing predators that have few interaction

records could help avoid false non-interactions because many of the interactions/prey of these

predators are likely missing (i.e., they have sparse interaction records, although these species

would also include some predators with narrow diet breadths). When applied to the Simpson

Desert data, our models trained on the global dataset modified by removing predators with <

5 interaction records performed better than those trained on the original/full global dataset

(Figure 3). This result might reflect a reduction of false non-interactions in the modified

training dataset. Thus, removing predators/consumers that are missing most of their

interactions from training datasets could be an effective way to filter training data and

improve model performance.

Model performance was especially sensitive to false non-interactions in the training data

involving focal predators. This type of change in data quality disrupted relative prey

preferences and lowered probabilities assigned to interactions more extensively than did

reducing the number of records involving focal predators (Figure 4a *versus* 4b, Figure 4c

versus 4d), reinforcing the importance of avoiding false non-interactions—especially for

focal predators—when generating pseudo-non-interactions for training data; otherwise, such

data can change which species are identified as most-likely prey. Although removing

predators/consumers with few interaction records from training data and/or restricting the

generation of non-interactions based on relative size and trophic level are two simple

methods for avoiding false non-interactions as we discussed, there are other strategies that

could be applied to the entire- or focal-predator-component of training datasets to minimise

the number of false non-interactions. For example, false non-interactions could be minimised

by generating non-interactions on a case-by-case basis (similar to the 'target-group absences'

approach used in species distribution modelling; Barbet-Massin et al. 2012), an approach

shown to improve the accuracy of predator-prey interactions predicted by deep-learning

models (Fricke et al. 2022). We generated non-interactions for the global dataset using

interaction data from different regions, studies, and sources, but this approach could generate

false non-interactions, such as when pairs of species are allopatric in studies that contributed

to the training data and so are designated as non-interacting, but the component species

interact when/where they are sympatric. This approach could also generate misleading non-

interactions, such as pairs of currently allopatric species that might interact should future

conditions force them to become sympatric. Instead, training data non-interactions could be

generated based on individual studies or locations of recorded interactions describing what a

predator does and does not eat in a particular area. Such a nuanced approach would identify

more reliable and informative non-interactions than would be achieved by assuming that any

undocumented pairs from an incomplete global database are truly not predator and prey.

Our models performed better with the global dataset than with the Simpson Desert data.

This difference might indicate that the random forests learnt patterns in the global dataset that

are not as effective when re-applied to specific communities. For example, the random forests

might have learnt from the global dataset that predators with particular traits do not eat prey

with particular traits, but predators or prey with those traits might not be present in the

Simpson Desert. In other words, learning the relationship did not help the random forest

predict interactions in the Simpson Desert. Alternatively, random forests might learn which

predators had few or many prey records within the global dataset—another pattern that would

be less effective at identifying interactions and non-interactions when applied to the Simpson

Desert predators due to the similarity among focal predators in terms of number of

interactions in the training data (60% of predators in the original/full global dataset had < 5

interaction records whereas all Simpson Desert focal predators had > 10 interaction records in

the global dataset). This might explain why our models did not perform as well on the global

data when we used the modified dataset rather than the original/full global dataset for training

and testing, because we had removed predators with few interaction records from the

modified dataset. Geographic biases in the global training dataset could also lead to weaker

performance on our ecosystem-specific dataset (Tuia et al. 2022). Irrespective of what caused

the difference in model performance, our results indicate that performance based on global

datasets might not be a strong indicator of performance when applied to single communities.

Previous studies applying random forests in aquatic and invertebrate communities have

tended to report higher performance than our models achieved (mean true skill statistic

consistently > 0.6; Desjardins-Proulx et al. 2017, Laigle et al. 2018, Brousseau et al. 2018,

Parravicini et al. 2020, Strona et al. 2021). There are several possible explanations for this

discrepancy, including previous studies (*i*) pooling prey species into broad resource

categories (Parravicini et al. 2020), (*ii*) using higher-quality training data (Laigle et al. 2018,

Parravicini et al. 2020), (*iii*) including resource predictions for a wider range of trophic

groups, including easily-excluded non-interactions such as herbivores and/or detritivores

eating other animals (Desjardins-Proulx et al. 2017), and (*iv*) focusing on taxa whose

interactions might be easier to predict than for terrestrial vertebrates (although further testing

is required to evaluate this possibility). Another potential means of improving the

performance of random forests that neither we nor others have yet addressed is to include

detail on intraspecific variation in traits and interactions, including ontogenetic shifts

(González-Varo and Traveset 2016).

Random forest and other machine learning approaches are useful and flexible tools that

have great potential in terms of predicting species interactions. While global interaction

databases are a valuable resource for training and testing such models, they currently include

only a small fraction of real-world species interactions. Furthermore, for the consumers that

are included in such databases, in most cases only a small subset of their trophic interactions

are documented, thereby increasing the likelihood of generating false-non-interactions.

Databases therefore need to be expanded, and a framework for generating non-interactions

that avoids or minimises false non-interactions and retains informative non-interactions is

required (Figure 5 shows a proposed framework). By making such changes, the application of

random forests and other machine-learning methods (including ensemble approaches) can be

further honed to infer species interactions more accurately.

References

Allan, J. D. et al. 2021. Energy Flow and Nutrient Cycling in Aquatic Communities. - In:
Allan, J. D. et al. (eds), Stream Ecology : Structure and Function of Running Waters.
Springer International Publishing, pp. 357–381.

Arif, S. and MacNeil A. 2022. Predictive models aren't for causal inference. Ecol. Lett. 25:
1741-1745.

Barbet-Massin, M. et al. 2012. Selecting pseudo-absences for species distribution models:
how, where and how many? - Methods Ecol. Evol. 3: 327–338.

Benandi, G. et al. 2012. Quantitative predictions of interactions in bipartite networks based
on traits, abundance, and phylogeny? – Am. Nat. 3: 327–338.

Blois, J. L. et al. 2013. Climate Change and the Past, Present, and Future of Biotic
Interactions. - Science 341: 499–504.

Bohan, D. A. et al. 2017. Next-Generation Global Biomonitoring: Large-scale, Automated
Reconstruction of Ecological Networks. - Trends Ecol. Evol. 32: 477–487.

Breiman, L. 2001. Random forests. – Mach. Learn 45:5-32.

Brousseau, P.-M. et al. 2018. Trait matching and phylogeny as predictors of predator-prey
interactions involving ground beetles (M Ferrari, Ed.). - Funct. Ecol. 32: 192–202.

Caron, D. et al. 2022. Addressing the Eltonian shortfall with trait-based interaction models. -
Ecol. Lett. 25: 889–899.

Chamberlain, S. et al. 2022. rgbif: interface to the global biodiversity information facility.

Cox, N. et al. 2022. A global reptile assessment highlights shared conservation needs of
tetrapods. - Nature: 1–6.

Dellinger, A. S. et al. 2019. Beyond buzz-pollination – departures from an adaptive plateau
lead to new pollination syndromes. - New Phytol. 221: 1136–1149.

Desjardins-Proulx, P. et al. 2017. Ecological interactions and the Netflix problem. - PeerJ 5:
e3644.

Doherty, T. S. et al. 2016. Invasive predators and global biodiversity loss. - Proceedings of
the National Academy of Sciences 113: 11261–11265.

Elmasri, M. et al. 2020. A hierarchical Bayesian model for predicting ecological interactions
using scaled evolutionary relationships. - Ann. Appl. Stat. 14: 221–240.

Faisal, A. et al. 2010. Inferring species interaction networks from species abundance data: A
comparative evaluation of various statistical and machine learning methods. - Ecol.
Inform. 5: 451–464.

Fillios, M. et al. 2012. The impact of the dingo on the thylacine in Holocene Australia. -
World Archaeology 44: 118–134.

Fleiss, J.L. et al. 2003. Statistical methods for rates and proportions. – John Wiley & Sons,
Ltd.

Flinders University 2021. Deep Thought (HPC).
https://doi.org/10.25957/flinders.hpc.deepthought

Fricke, E. C. et al. 2022. Collapse of terrestrial mammal food webs since the Late
Pleistocene. - Science 377: 1008–1011.

González-Varo, J. P. and Traveset, A. 2016. The labile limits of forbidden interactions. -
Trends Ecol. Evol. 31: 700–710.

Gravel, D. et al. 2013. Inferring food web structure from predator-prey body size
relationships (R Freckleton, Ed.). - Methods Ecol. Evol. 4: 1083–1090.

Guénard, G. and Legendre, P. 2022. MPSEM: Modeling phylogenetic signals using
eigenvector maps.

Guénard, G. et al. 2013. Phylogenetic eigenvector maps: a framework to model and predict
species traits. - Methods in Ecology and Evolution 4: 1120–1131.

Herberstein, M. E. et al. 2022. AnimalTraits - a curated animal trait database for body mass,
metabolic rate and brain size. - Sci Data 9: 265.

Hortal, J. et al. 2015. Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity. -
Annu. Rev. Ecol. Evol. Syst. 46: 523–549.

Iturbide, M. et al. 2015. A framework for species distribution modelling with improved
pseudo-absence generation. - Ecological Modelling 312: 166–174.

Jones, K. E. et al. 2009. PanTHERIA: a species-level database of life history, ecology, and
geography of extant and recently extinct mammals: *Ecological Archives* E090-184
(WK Michener, Ed.). - Ecology 90: 2648–2648.

Jordano, P. 2016. Chasing ecological interactions. - PLoS Biol. 14:e1002559.

Kaneryd, L. et al. 2012. Species-rich ecosystems are vulnerable to cascading extinctions in an
increasingly variable world: Extinction Cascades in a Variable World. - Ecol. Evol. 2:
858–874.

Klomberg, Y. et al. 2022. Spatiotemporal variation in the role of floral traits in shaping
tropical plant-pollinator interactions. - Ecol. Lett. 25: 839–850.

Kotula, H. J. et al. 2020. Predicting direct and indirect non-target impacts of biocontrol
agents using machine-learning approaches.

Laigle, I. et al. 2018. Species traits as drivers of food web structure. - Oikos 127: 316–326.

Liem, K.F. 1984. Functional versatility, speciation, and niche overlap: are fishes different? –
Trophic interactions within aquatic ecosystems: 269-305. Routledge.

Liem, K.F. 1990. Aquatic versus terrestrial feeding modes: possible impacts on the trophic
ecology of vertebrates. – Am. Zool. 30: 209-221.

Llewelyn, J. 2022. Random forests for predicting predator-prey interactions in terrestrial
vertebrates. - GitHub and Zenodo: DOI: 10.5281/zenodo.7037189.

Llewelyn, J. et al. 2022. Sahul's megafauna were vulnerable to plant-community changes due
to their position in the trophic network. - Ecography 2022: DOI: 10.1111/ecog.06089.

Lobo, J. M. et al. 2010. The uncertain nature of absences and their importance in species
distribution modelling. - Ecography 33: 103–114.

Malley, J.D. et al. 2012. Probability machines. – Methods Inf. Med. 51: 74-81.

Mateo, R. G. et al. 2010. Profile or group discriminative techniques? Generating reliable
species distribution models using pseudo-absences and target-group absences from
natural history collections: Reliability of models using pseudo-absences and target-
group absences. - Divers. Distrib. 16: 84–94.

Miller-ter Kuile et al. 2022. Predator-prey interactions of terrestrial invertebrates are
determined by predator body size and species identity. – Ecology: e3634

Morales-Castilla, I. et al. 2015. Inferring biotic interactions from proxies. - Trends Ecol.
Evol. 30: 347–356.

Morton, D. N. et al. 2022. Merging theory and experiments to predict and understand
coextinctions. - Trends Ecol. Evol. in press.

Oliveira, B. F. et al. 2017. AmphiBIO, a global database for amphibian ecological traits. -
Sci. Data 4: 170123.

Ornai, A. and Keasar, T. 2020. Floral complexity traits as predictors of plant-bee interactions
in a Mediterranean pollination web. - Plants 9: 1432.

Paradis, E. and Schliep, K. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. - Bioinformatics 35: 526–528.

Parravicini, V. et al. 2020. Delineating reef fish trophic guilds with global gut content data
synthesis and phylogeny (P Jordano, Ed.). - PLoS Biol. 18: e3000702.

Pichler, M. et al. 2020. Machine learning algorithms to infer trait-matching and predict
species interactions in ecological networks. - Methods Ecol. Evol. 11: 281–293.

Pires, M. M. et al. 2015. Pleistocene megafaunal interaction networks became more
vulnerable after human arrival. - Proc. R. Soc. B 282: 20151367.

Poisot, T. 2022. Guidelines for the prediction of species interactions through binary
classification. - EcoEvoRxiv: 16.

Poelen, J.H. et al (2014) Global biotic interactions: an open infrastructure to share and
analyze species-interaction datasets. – Ecol. Inform. 24: 14-8-159.

Pomeranz, J. P. F. et al. 2019. Inferring predator–prey interactions in food webs. - Methods
Ecol. Evol. 10: 356–367.

Pringle, R. M. and Hutchinson, M. C. 2020. Resolving Food-Web Structure. - Annu. Rev.
Ecol. Evol. Syst. 51: annurev-ecolsys-110218-024908.

Ripple, W. J. and Van Valkenburgh, B. 2010. Linking top-down forces to the Pleistocene
megafaunal extinctions. - BioScience 60: 516–526.

Samson, F. B. and Knopf, F. L. 1996. Ecosystem Management: Selected Readings. - Springer
New York.

Shurin, J. B. and Gruner, D. S. 2006. All wet or dried up? Real differences between aquatic
and terrestrial food webs. - Proc. R. Soc. B 273: 1-9.

Sinclair, A. R. E. et al. 2003. Patterns of predation in a diverse predator–prey system. -
Nature 425: 288–290.

Smith, J. G. and Phillips, B. L. 2006. Toxic tucker: the potential impact of Cane Toads on
Australian reptiles. - Pac. Conserv. Biol. 12: 40–49.

Strona, G. 2022. Hidden Pathways to Extinction. - Springer International Publishing.

Strona, G. and Bradshaw, C. J. A. 2022. Coextinctions dominate future vertebrate losses from
climate and land use change. – Sci Adv 8: eabn4345.

Strona, G. and Bradshaw, C. J. A. 2018. Co-extinctions annihilate planetary life during
extreme environmental change. - Sci Rep 8: 16724.

Strona, G. et al. 2021. Ecological dependencies make remote reef fish communities most
vulnerable to coral loss. - Nat. Commun. 12: 7282.

Strydom, T. et al. 2021. A roadmap towards predicting species interaction networks (across
space and time). - Proc. R. Soc. B 376: 20210063.

Strydom, T. et al. 2022. Food web reconstruction through phylogenetic transfer of low-rank
network representation. – Methods Ecol. Evol. 2022: DOI: 10.1111/2041-
210X.13835.

Sydenham, M. A. K. et al. 2021. MetaComNet: A random forest-based framework for
making spatial predictions of plant-pollinator interactions. - Methods Ecol. Evol. in
press.

Thompson, J. N. 1999. The evolution of species interactions. - Science 284: 2116–2118.

Tilman, D. et al. 2017. Future threats to biodiversity and pathways to their prevention. -
Nature 546: 73–81.

Traill, L. W. et al. 2010. Mechanisms driving change: altered species interactions and
ecosystem function through global warming. - J. Anim. Ecol. 79: 937–947.

Tuia, D. et al. 2022. Perspectives in machine learning for wildlife conservation. – Nat.
Commun. 13: 792.

Tylianakis, J. M. et al. 2008. Global change and species interactions in terrestrial ecosystems.
- Ecol. Lett. 11: 1351–1363.

Valdovinos, F. S. et al. 2018. Species traits and network structure predict the success and
impacts of pollinator invasions. - Nat. Commun. 9: 2153.

Van der Putten, W. H. et al. 2010. Predicting species distribution and abundance responses to
climate change: why it is essential to include biotic interactions across trophic levels.
- Phil. Trans. R. Soc. B 365: 2025–2034.

Wilman, H. et al. 2014. EltonTraits 1.0: Species-level foraging attributes of the world's birds
and mammals: *Ecological Archives* E095-178. - Ecology 95: 2027–2027.

Wright, M. N. and Ziegler, A. 2017. ranger: A fast implementation of Random Forests for
high dimensional data in C++ and R. - J. Stat. Soft. in press.

Figure captions

Figure 1. Density plots comparing the performance (true skill statistic) of random forest models predicting predator-prey interactions in birds and mammals. The random forests used either ecomorphological traits, phylogenetic eigenvectors, or both types of data, and were applied to two interaction datasets: a global interaction dataset (panels in the left column) and a specific ecosystem dataset (7 predators and their potential prey/sympatric species in the Simpson Desert; panels in the right column). We also assessed the effects of using many variables (42, 42, and 84 variables, for ecomorphological, phylogenetic eigenvector, and both data-type models) or few variables (10, 10, and 20 variables, for ecomorphological, phylogenetic eigenvector, and both data-type models), and restricting the training dataset to predators with ≥ 5 prey records (modified training dataset). Panels (a) and (b) show results from models that use the complete training dataset and many variables; (c) and (d) show results from models that use the complete training dataset and few variables; (e) and (f) show results from models that use the modified training dataset and many variables; and (g) and (h) show results from models that use the modified training dataset and few variables.

Figure 2. The five most- and least-likely mammalian and bird prey for the red fox (*Vulpes vulpes*) in the Simpson Desert predicted by random forest. Solid blue arrows indicate prey that were predicted and observed, dashed blue arrows show prey that were predicted but not observed (potentially indicating interactions that occur but have not yet been recorded), and red lines indicate species that were not predicted or observed as prey. The five most-likely prey according to the many-variable phylogenetic and ecomorphological model (run 100 times and trained on the modified training data) were the common wallaroo (*Osphranter robustus*), long-haired rat (*Rattus villosissimus*), red kangaroo (*Osphranter rufus*), Eurasian coot (*Fulica atra*), and sandy inland mouse (*Pseudomys hermannsburgensis*). The five least-likely prey were the little eagle (*Hieraaetus morphnoides*), brown goshawk (*Accipiter fasciatus*), whistling kite (*Haliastur sphenurus*), wedge-tailed eagle (*Aquila audax*), and spotted harrier (*Circus assimilis*).

Figure 3. Effect of training-dataset quality on the performance of random forest for predicting predator-prey interactions in birds and mammals. Each panel shows performance when modifying a percentage of either the whole or specific taxonomic groups in the training dataset. Panels (a) and (b) show performance when changing the whole dataset, panels (c) and (d) show performance when restricting changes to focal prey, panels (e) and (f) show performance when restricting changes to focal predators, and panels (g) and (h) show performance when restricting changes to non-focal (non-Simpson Desert) species. Panels in the left column show the effects of removing records from the training data, whereas panels in the right column show the effects of switching interaction records to non-interactions (false non-interactions). We trained random forests on the modified global predator-prey interaction dataset (i.e., restricted to predators with ≥ 5 prey records) and applied them to seven predators and their potential prey from the Simpson Desert. Plot points indicate performance (according to the true skill statistic) when applied to the Simpson Desert predators.

Figure 4. Effects of the quality of the training dataset on (*i*) the predicted relative suitability of prey and (*ii*) the mean probabilities assigned to potential prey by random forests. Panels in the left column show the effects of reducing the number of interactions records involving the focal predator in the training dataset, whereas panels in the right column show the effect of switching interaction records involving the focal predator to non-interactions (false non-interactions). Panels (a) and (b) show the correlation between prey probabilities assigned by

the models trained on the changed datasets with that assigned by models trained on the
complete (modified) training dataset. Panels (c) and (d) show the mean probability
(suitability) assigned to potential prey by models trained on the changed datasets minus those
values assigned by the models trained on the complete (modified) training datasets. We
trained these random forests on the modified global predator-prey interaction dataset
(restricted to predators with ≥ 5 prey records) and applied them to seven predators and their
potential prey from the Simpson Desert.
871

872 **Figure 5**. Framework for preparing and optimising random forest models for predicting
873 species interactions. Most of these steps and methods are also relevant to the application of
874 other machine-learning approaches for inferring species interactions.
875

random forest for predicting community interactions

steps	methods

1 — obtain interaction & non-interaction data for training

extract documented interaction and non-interaction records
from online databases, publications, unpublished data, and/or targeted collection of new data via experiments or field observations (see below for non-interactions)

prioritise collecting interaction data for focal predators & ecologically similar species

generate pseudo-non-interactions
Few studies report non-interactions; pseudo-non-interactions usually assigned instead by assuming undocumented interactions imply no interaction. Strategies to assign reliable (truly non-interacting) & informative pseudo-non-interactions required. For example:
1. generate non-interactions from individual studies rather than from pooled data;
2. include only co-occurring species if predictions needed for co-occurring species;
3. remove predators with few interaction records;
4. restrict non-interactions based on trait-matching rules (e.g., body size, trophic level)

important to avoid incorrect non-interactions involving focal predators

2 — add trait data to training dataset

collate ecomorphological/phylogenetic data
sources: online databases, publications, unpublished data, targeted collection of new trait data

3 — optimise training dataset

optimise interactions:non-interactions ratio (dataset balance)
out-of-bag design; vary interactions:non-interactions ratio; compare performance of random forests trained & tested on these datasets

optimise ratio of suitable to unsuitable predator/prey species combinations in non-interaction data
based on trait matching & out-of-bag approach (see above)

4 — optimise model hyper-parameters

identify & tune important hyper-parameters
e.g., # variables at each tree split, # trees, probability threshold for assigning interactions, max tree depth, max/min leaf node size

5 — build dataset for making predictions

collate data for focal species/ecological community (focal species = those for which predictions needed)
add same trait and/or phylogenetic variables as in training data & arrange into species pairs for making predictions

6 — train model on optimised training dataset & apply to focal species pairs to make predictions