Title: Spring growth variation in pioneer and fibrous roots in *Abies sachalinensis* seedlings from

provenances with contrasting snow cover environments

Author names and affiliations: Tetsuto Sugai[1*], Wataru Ishizuka[2], Taiga Masumoto[3], Izuki Endo[4,5],

Jun'ichiro Ide[5], Saki Fujita[6], Makoto Kobayashi[7], Naoki Makita[3]

1: Forestry and Forest Products Research Institute, Hokkaido Research Center, Sapporo, Japan

2: Hokkaido Research Organization, Forestry Research Institute, Bibai, Japan

3: Faculty of Science, Shinshu University, Matsumoto, Japan

4: School of Human Science and Environment, University of Hyogo, Himeji, Japan

5: Department of Applied Chemistry and Bioscience, Chitose Institute of Science and Technology,

Chitose, Japan

6: Forestry and Forest Products Research Institute, Tsukuba, Japan

7: Field Science Center for Northern Biosphere, Hokkaido University, Horonobe, Japan

*Corresponding authors: Forestry and Forest Products Research Institute, Hokkaido Research Center,

062-0045, Sapporo, Japan, Telephone: +81-11-590-5527, Fax: +81-11-851-4167, Email:

sugai922@affrc.go.jp

Shortened title: SUGAI ET AL.— SPRING ROOT GROWTH AND PROVENANCE VARIATION

Keywords: phenology, local adaptation, soil space exploration, snow, boreal forest

Abstract:

Premise of research. In boreal forests with various snow cover conditions, local adaptation can be

expected in spring root exploration due to soil freezing or competition for the fundamental niche.

However, few studies have investigated the adaptive strategies for acquiring soil space and resources after

snow thawing. To get insights for the eco-evolutionary significance of root growth dynamics in boreal

regions, this study evaluated the intraspecific variation of morphology and structures in pioneer roots and

fibrous roots.

Methodology. Validation model specie was *Abies sachalinensis*, a representative boreal conifer

species with local adaptation. Five-year-old seedlings derived from an eastern provenance with low snow

cover and a northern provenance with high snow cover were grown for two years in a common garden of

high snow cover region. To predict spring growth patterns of roots, seedlings were dug up in early spring

just after snow thawing and late spring after one month has passed. Pioneer roots exploring soil space and

fibrous roots acquiring soil resource were respectively collected, and their functions were evaluated by

morphology and structures indicating its construction costs and maturity.

Pivotal results. In pioneer roots, morphological and structural difference between sampling months

were significant only in northern provenance, and high construction cost and well-developed structure

were observed in early spring. This represents the intraspecific variation in reaction norms of pioneer

roots to spring soil environment, indicating the soil space exploration in northern provenance may start in

early spring or earlier. In fibrous roots, differences of specific root length and area between sampling

months were significant only in eastern provenance, and low construction cost were observed in early

spring. This may reflect the plastic responses of fibrous roots to the condition of high snow cover region.

Conclusions. This study indicated that local adaptation to contrasting snow cover condition may lead

to the differentiation of spring root growth dynamics for the competitive or conservative strategies. Our

results address better understandings of the mechanisms of niche acquisition in evergreen conifer.

Introduction

In the cool-temperate zone, temporal variability of environments is recognized as a season. To the

seasonal dynamic, a developmental pattern of traits is often adaptive (Ghalambor et al. 2007, Forrest and

Miller-Rushing 2010). The phenological optima are expected to be specific to the regional environments,

causing the intraspecific variation of phenology because of local adaptation (De Lisle et al. 2022). Since

the intraspecific variation in phenology is the consequence of reaction norms to seasonal stimuli

(West-Eberhard 2003), its patterns provide cues for predicting how organisms seasonally react to climate

changes (Franks et al. 2014). In trees, the intraspecific variation in the development of above-ground

traits has been elucidated. The significant intraspecific variation has been often observed in specific

seasons (West-Eberhard 2003), such as the timing of bud burst (Mediavilla & Escudero 2009), the leaf

longevity (De Kort et al. 2016), freezing tolerances (Noordermeer et al. 2021), but also in the seasonal

pattern itself, such as photosynthetic activities (Fréchette et al. 2020, Sugai et al. 2023) and stem growth

(Sanmiguel-Vallelado et al. 2021). However, the body of studies on the intraspecific variation in the

below-ground dynamics is still growing yet, for example, the non-structural carbon (Wang et al. 2018),

the growth pattern of fine roots (Ma et al. 2022). Moreover, little information is available regarding the

root phenology (McCormack et al. 2015, Tamura et al. 2022, Liu et al. 2022). The insights derived from

root phenology and its intraspecific variation would shed light on the ecological significance for

understanding the mechanisms of niche acquisition and their evolutionary aspects (Begon et al. 2006).

Soil temperature has been considered as the driver of root development given sufficient

precipitation (Ding et al. 2020). The sensitivity of fine root elongation to soil temperature was variable

between tree species (Pregitzer et al. 2000). In evergreen conifer species, the relatively high quantity of

fine roots has been recognized in spring (Persson 1978), when the fine root elongation rate of conifers

was correlated only below 8 °C of soil temperature (Wang et al. 2018). This fine root elongation at

relatively low soil temperatures, where other species including deciduous trees would not develop (Steele

et al. 1997), may reflect the ecological characteristics as late successional species who can acquire niche

early and stay in for the long term (Makoto et al. 2020). These studies indicated that the root phenology

would have the potential to be evolved, and particularly boreal evergreen conifers would have developed

the root dynamics with relatively low soil temperature dependence at the beginning of the growing season

when low soil temperatures begin to increase in spring (Ding et al. 2020).

In soils with heterogeneous nutrient and physical properties, roots often prioritize exploration

rather than acquisition (de Kroon & Visser 2003, Begon et al. 2006). Among the various root taxonomies,

pioneer roots are visually and functionally distinguishable from fibrous roots (Figure 1). Compared with

fibrous roots, pioneer roots have been known to have a thick, well-developed structure and requires high

construction costs (Zadworny & Eissenstat 2011, Montagnoli et al. 2021) which lead to the tolerances to

anaerobic conditions with superior transportability and the exploration function of soil space. On the other

hand, fibrous roots often show a relatively thin diameter and cortex layers without lignified woody cells,

which enable them to efficiently absorb soil water and nutrients, and symbioses with mycorrhizal fungi

(de Kroon & Visser 2003). The clear sequential stages of development are recognized in these roots

(Wilcox 1954), where pioneer roots develop first, and then fibrous roots often emerge laterally from the

maturity zone of pioneer roots. However, the intraspecific variations in the growth patterns of these root

types have not been verified simultaneously so far.

This study investigated the morphology and structure in different root types in a boreal evergreen

conifer species, Sakhalin fir (*Abies sachalinensis*). In Sakhalin fir, the local adaptation has been reported

mainly in above-ground traits (e.g., Ishizuka et al. 2021, Sugai et al. 2023), which would represent

responsible association with winter climate environments. In boreal forests, the regional variation in

spring soil temperature depends on snow cover conditions (Fukuzawa et al. 2021, Makoto et al. 2022). In

high snow cover regions, soil temperature and resource availability would increase rapidly (Bardgett et al.

2005) more than low snow cover regions because soil suddenly faced relatively high air temperatures

when the thermal interception by snow cover ends. In low snow cover regions, however, the rapid root

elongation can lead to a risk of encountering soil freezing. We expected the fast development of pioneer

roots would be adaptive for high snow cover provenances since the competition for underground

resources in spring will be intense. If so, a relatively thick and well-developed pioneer roots may be

observed in high snow cover provenance. On the other hand, fibrous roots branching later than pioneer

roots may be dominated by seasonal stimuli without genetic variation to optimize the balance between

construction costs and its benefit (de Kroon & Visser 2003). Here, we elucidated the intraspecific

variation in the functional morphological traits; specific root length (ratio of root length to mass), specific

root area (ratio of root surface area to mass) and root tissue density (ratio of root mass to volume), and the

anatomical traits; central cylinder, cortex, and epidermis, which could exhibit its functions and the

resource availability (McCormack et al. 2015, Yahara et al. 2019). We conducted a common garden trial,

where seven-year-old seedlings originating from high and low snow cover provenances were dug up in

early spring just after snow thawing and late spring after one month has passed to predict spring growth

patterns of roots.

Materials and methods

Study site and material

A common garden trial was conducted at the nursery of the Forestry Research Institute, Hokkaido

Research Organization, located at central Hokkaido (43°29′N, 141°85′E; 45 m a.s.l.). As an experimental

material, Sakhalin fir (*Abies sachalinensis*) was adopted in this study. Sakhalin fir distribution is on a

northern island in Japan, Hokkaido in which various environmental conditions are recognized along with

a geographic gradient, e.g., maximum snow depth from 20 cm to more than 400 cm mainly from eastern

side to western side (Supplemental material Appendix 1). In the most regions of Hokkaido, snow begins

to thaw in mid-March, and it has widely disappeared by early May. Previous studies have reported the

genetic variation of Sakhalin fir in above-ground traits such as home advantage in growth performance,

frost tolerance of winter buds and disease resistance (Sakai 1983, Eiga 1984, Ishizuka et al. 2021). In

addition, boreal evergreen conifers, including Sakhalin fir, have a relatively early onset and late cessation

of roots development from mid-April when covered snow just started melting to mid-December when

snow have started covering soil (Satoh 1995). Thus, Sakhalin fir is a valuable model for testing the

adaptive developmental patterns of different root types. In this common garden trial, open-pollinated

progeny of selected mothers (family lines) were planted. Plant materials were sown in 2015 using seeds

produced in a seed orchard of Sakhalin fir in 2014 and transplanted twice at spring of 2017 and 2019.

Tested progeny were selected from 8 family lines derived from two provenances with contrasting

environmental condition, i.e., (i) the eastern-edge provenance ($43°07' \pm 0.01$N, $145°03' \pm 0.01$E) and (ii)

the northern provenance ($44°35' \pm 0.08$N, $142°32' \pm 0.01$E). At the time of second transplanting in 2019,

the experimental plots of approximately 2.2 m $\times$ 2.7 m were set up with three replications in the same

field with the growing field before transplanting, with seedlings planted approximately 50 cm apart in

each. Basic meteorological information on the selected family lines and the cultivated experimental

environmental conditions were summarized (Supplemental material Appendix 1, Supplemental material

Appendix 2). After transplantation, seedlings were cultivated with commercial fertilizer and automatic

irrigation during two growing seasons except from November to May during 2019−2020.

In a previous study, pioneer roots of Sakhalin fir seedlings were observed in late March under snow

cover at the same site of this experiment (Satoh 1995), which was validated again in this study

(Supplemental material Appendix 3). Based on them, seedlings were dug up both in April immediately

after snow thawing, and in May when the soil temperature has increased sufficiently after one month has

passed since the snow was thawing. A total of 48 seven-year old seedlings of seven-year-old (24

seedlings per month) were selected as samples for the following analysis. Then, 12 seedlings in northern

provenance and 12 seedlings of eastern provenance were collected from 3 plots (4 seedlings of northern

and eastern provenances per plot) in each month. Of these, three individuals were excluded from samples

in May due to abnormal growth conditions, so the final number of seedlings collected was 45 in this study.

Additionally, the seedlings of both northern and eastern provenance continued to grow without being

sampled in spring were collected in December 2021 and February 2022 (Supplemental material Appendix

3).

Root morphology

Immediately after cutting the above-ground parts of each individual seedling, a below-ground part

was carefully collected by shovels for each individual. The attached soil with a below-ground part was

removed by tap water within the bucket which was kept for the following drymass measurement. After

cleaning the whole root system, the length and number of pioneer roots in the below-ground part were

measured. In this study, pioneer root was distinguished from other fibrous roots based on the diameter

and white color from the elongation zone to maturity zone of a root (Figure 1, Supplemental material

Appendix 4). From each individual, five pioneer roots were randomly selected from all the pioneer roots

to measure the length of its root length, which would indicate the function to explore soil space and/or its

maturity (Zadworny & Eissenstat 2011, Montagnoli et al. 2021). The length of the pioneer root was

measured as the length from the root end to the elongation zone with bright white by a ruler to the nearest

1 mm. The number of all pioneer roots per individual was counted, including the number of this root in

stored soil water in the bucket. After pioneer root measurements were completed, intact fibrous roots and

pioneer roots were randomly collected from whole roots of individuals to evaluate morphological and

anatomical characteristics, respectively. Finally, the roots were dried and used for drymass measurements,

where all roots contained the stored soil were also used. The fibrous and pioneer roots collected for

morphometrics were scanned with a flatbed scanner (GT-X980, EPSON, Tokyo, Japan) immediately after

collection. The scanned images were analyzed using root image analysis software (WinRHIZO, REGENT

Inc, Quebec, Canada) to obtain root length, surface area, volume, and average diameter. The dry weight

of roots completed scans was measured by 0.1 mg increments. Then, specific root length (SRL, m g^{-1}),

and specific root area (SRA, cm^2 g^{-1}) were obtained as the ratio of dry weight to length and area of roots,

and root tissue density (RTD, g cm^{-3}) was obtained as the ratio of volume to dry weight of roots.

Root anatomy

The fibrous and pioneer roots for evaluating tissue structure were impregnated with 2%

glutaraldehyde solution immediately after collection. After one day of permeation of the fixative solution

into the tissue, the tissue was stored in 80% ethanol solution. To prepare sections from the samples, the

samples were initially traversed mainly from the maturation zone with a cutter, and a few centimeters of

the sample were fixed horizontally on a freezing stage (MC-802A, YAMATO, Saitama, Japan). Sections

with a thickness of 50 µm were cut on a microtome (REM-700, YAMATO, Saitama, Japan) and stained

with iodine solution for approximately 30 seconds. The sections were then observed under an optical

microscope and the tissue structure was captured (VH-5500, KEYENCE, Osaka, Japan). This process was

repeated approximately three times per sample, and the image with the clearest structural characteristics

among the images obtained was used for subsequent image analysis.

The image analysis of root sections was performed by a free image processing software (ImageJ,

NIH, USA) in this study. As root tissue structures, diameter, cortex, and central cylinder were measured

in the obtained section image process (Supplemental material Appendix 4). These characteristics were

measured at three locations from a single image and their average value was calculated. Since the linear

relationships between a diameter of root section and average size of each cell as well as the other

anatomical values were observed, these anatomical values in each sample were converted to relative ones

to the diameter of root section, respectively.

Mass and structural fraction

For the pioneer root quantity at a whole plant level, total weight was estimated from SRL and total

root length estimated from the total number and average length per seedling. All the sampled roots were

dried in oven at 75 °C for approximately seven days and weighed. The roots used for the weight

measurement were divided into woody roots with a diameter of 1.5 mm or more since the average

diameter of the identified woody roots ranged up to approximately 1.5 mm in this study (Supplemental

material Appendix 5). The other part except for woody root, where root branching was difficult to

identify, was classified as a rootstock. The dry mass of fibrous roots was obtained by subtracting the

estimated dry weight of the pioneer roots from the weights of the other roots except for woody roots and

rootstock.

Statistical analysis

All statistical analyses were performed by an open-source package for statistical computing (R

ver.4.1.2, R Core Team 2019). In this study, missing values accounted for maximum of approximately

30% in the data of pioneer roots and a minimum of approximately 8% in fibrous roots. Since the complete

case analyses with the process of excluding missing values should led to biased (Ellington et al. 2014), we

alternatively adopted the multiple imputation method using the *mice* package (Nissen et al. 2019). The

effects of sampling months and the provenances (i.e., genetic variation) on the following traits; total root

dry mass, mass fraction of each root types, average length and total number of pioneer roots, SRL, SRA,

RTD, average diameter, cortex, and central cylinder were tested by Two-way ANOVA in the following

generalized linear mixed model. Because of the unbalanced data structure in this study, a type III variance

analysis was performed using the *car* package. For this analysis, all values were used after logarithm

transformation. The modeling was conducted by *lme4* package. The model was constructed as follows;

$$Y_{ijkl} = M_i + G_j + M_i \times G_j + \varepsilon_{ij} + \varepsilon_k + \varepsilon_l$$

where Y_{ijkl} denotes the measured values of traits of progeny of k-th family line derived from j-th region at

i-th sampling months and located in l-th plots. M_i denotes the effect of sampling months, G_j denotes the

effect of provenances, and $M_i \times G_j$ denotes the interaction effect of sampling months and provenances.

For the random effects, ε_{ij}, ε_k, and ε_l denote the family lines of progeny, the plot replications, and

error term, respectively. To evaluate differences between any of the groups, the multiple comparison

adjusted by Tukey method was conducted as a post-hoc test using the *emmeans* package. A significant

level of 5% was adopted for all statistical tests in this study.

Results

While any significant changes were not observed in total root dry mass, the mass fraction of

pioneer roots were significantly increased from April to May regardless of provenances ($p <0.05$, Figure

2, Supplemental material Appendix 7). The mass fraction of fibrous roots was slightly but significantly

increased from April to May only in norther provenance ($p <0.05$, Figure 2). While there were not

significant difference in average length of pioneer roots (Figure 3), it tended to increase from April to

May in eastern provenance while decrease in northern provenance. On the other hand, the total number of

pioneer roots was significantly increased only in northern provenance ($p < 0.05$, Figure 3).

In pioneer roots, significant increases of SRL and SRA from April to May were observed only in

northern provenance ($p < 0.05$, Figure 4, Supplemental material Appendix 8). Any significant changes of

RTD were not observed in pioneer roots. In contrast, SRL and SRA of fibrous roots were significantly

decreased from April to May only in eastern provenance ($p < 0.05$, Figure 4). Contrary to pioneer roots,

RTD of fibrous roots was significantly increased from April to May regardless of provenances ($p < 0.05$,

Figure 4, Supplemental material Appendix 8).

The interaction effect between sampling months and provenances was significant in average

diameter of pioneer roots ($p < 0.05$, Supplemental material Appendix 8). While there were not significant

difference in average diameter of pioneer roots (Figure 5), its pattern of diameter change from April to

May was similar to the average length (Figure 3), namely it tended to increase from April to May in

eastern provenance while decrease in northern provenance. The interaction effect between sampling

months and provenances was also significant in relative size of central cylinder of pioneer roots ($p < 0.05$,

Supplemental material Appendix 8). Only in northern provenance, the relative size of central cylinder was

significantly decreased from April to May ($p < 0.05$, Figure 5). In contrast, any significant change of

average diameter, the relative size of central cylinder, and cortex and epidermis was not detected in

fibrous roots.

Discussion

Spring dynamics of pioneer roots

Previously, the quantitative dynamics of pioneer roots have been elucidated (Nakahata 2020,

Montagnoli et al. 2021) with respect to the seasonal changes of the soil temperature (Ding et al. 2020),

whereas relatively few studies have investigated the seasonal variation of construction costs and

structures of pioneer roots (Zadworny & Eissenstat 2011). Only in the seedlings originating from heavy

snow cover provenance (i.e., northern provenance), morphology and structure of pioneer roots

significantly varied between April and May (Figure 4, Figure 5), while the quantity of pioneer roots was

increased during the experimental periods regardless of provenances (Figure 2). To the best of our

knowledge, this is the first empirical data regarding the intraspecific differences in spring growth

dynamics of pioneer roots. These results would consist with previous studies (Satoh 1995) as well as our

preliminary survey (Supplemental material Appendix 3). In fact, the increase in the number of root

development in response to conditions of increasing soil temperature (Camm & Harper 1991). While root

growth is fundamentally activated as soil temperature increases from winter to spring in areas with

adequate precipitation (Pregitzer et al. 2000), it has been reported that the threshold of soil temperature at

which roots can grow in conifers is approximately 0 to 4 °C (Wang et al. 2018, Ding et al. 2020). Beyond

our expectation, however, the fact that the pioneer roots were observed even in April should raise the

question as to when the pioneer roots of April emerged. At our experimental site, where the ground

temperature remained near 0°C due to the thick snow cover and its insulation, the well-developed pioneer

roots were observed (Supplemental material Appendix 3). It is thus possible that changes in the physical

environment other than soil temperature, such as soil moisture conditions associated with snow thawing

(Sanmiguel-Vallelado et al. 2021), and finer-scale nutrient conditions (Pregitzer et al. 2000) are relevant

to spring root dynamics.

Spring dynamics of fibrous roots

As soil nutrients are generally heterogeneous, choosing the location for nutrient absorption is

important for plants to optimize the balance between construction costs of roots and benefits. Thus, the

plasticity of fibrous roots is expected to have the ability to regulate construction costs (de Kroon & Visser

2003, Begon et al. 2006). In accordance with our hypothesis, RTD of fibrous roots in both provenances

were increased from April to May (Figure 4). We also observed the non-significant changes in average

diameter and other structural traits in May (Figure 5). These results can demonstrate the matured fibrous

roots would develop from April to May. This would partly support the sensitivity to drought as well as

soil temperature of fibrous roots (Ding et al. 2020). Although we did not directly evaluate soil water

condition (Supplemental material Appendix 2), the soil was relatively dry in May, a month after the snow

thawed out in April. It has been reported that high RTD would denote the stress resistance and long life of

roots (Weemstra et al. 2016)

Implications for the intraspecific variation in spring root dynamics

Understanding phenotypic development is synonymous with understanding its evolutionary

processes (West-Eberhard 2003). The intraspecific variation due to local adaptation would represent a

trade-off between growth and survival (Csilléry et al. 2020). In high snow cover regions, soil temperature

and resource availability would increase rapidly (Bardgett et al. 2005) more than low snow cover regions

because soil suddenly faced relatively high air temperatures when the thermal interception by snow cover

ends (Fukuzawa et al. 2021, Makoto et al. 2022). As a result, after thawing of snow cover, the niche

suddenly become available and competition for soil space and resources intensifies (Begon et al. 2006). In

May, we observed the pioneer roots with lower construction costs and immature central cylinder,

indicating that it would be the relatively young pioneer roots with inferior function of soil space

exploration. This interception was further supported by the quantitative dynamics of pioneer roots (Figure

2, Figure 3), suggesting that the newly pioneer roots were developed from April with response to

increasing relatively lower soil temperature, i.e., from below 8 °C, approximately (Wang et al. 2018).

These results would support our hypothesis. Namely, the intraspecific variation in the ability to explore

soil space may occurs between high snow cover regions with high competition and low snow cover

regions with severe soil freezing.

On the other hand, in fibrous roots, SRL and SRA were significantly decreased from April to May

only in eastern provenance (Figure 3), indicating the construction costs was low and the function of soil

nutrients absorption was high in April. We suspect these results may be plastic responses in eastern

provenance to the environment of common garden with high snow cover condition. This would be partly

supported by the difference of root morphology between growing sites, i.e., the common garden and

eastern provenance site (Supplemental material Appendix 6), where the average values ($\pm$ standard error)

of SRL and SRA in eastern provenance site were 18.9 ± 1.67 m g^{-1} and 320.79 ± 9.97cm^2 g^{-1} in April.

Although the statistical comparison was not able to be performed, the relatively higher values of these

parameters were observed in common garden rather than a provenance site in April. This may be

associated with the tolerance to soil freezing or low temperature of fibrous roots in eastern provenance

(Sakai & Larcher 1987, Ambroise et al. 2020). Given that roots are the least freeze-tolerant of all plant

organs and soil freezing can be a selection pressure (Ambroise et al. 2020), the contrasting snow cover

environments may consequently be realized as the intraspecific variation in freezing tolerance in fibrous

roots. If so, since the soil temperature in April at common garden may be relatively high for seedlings of

eastern provenance, they may start the development of nutrient absorption even just after snow thawed

out. In other aspect, recent studies have reported dramatic changes in soil microbial communities due to

spring snow thawing (Broadbent et al. 2021), which was roughly common in the study site (Supplemental

material Appendix 1). Although this may be the chance of nutrient absorption for eastern provenance,

further experimental approach should be needed to verify whether there was the intraspecific variation in

the tolerance to soil freezing or low temperature of fibrous roots.

Conclusion

This study elucidated the spring growth dynamics of pioneer roots and fibrous roots in Sakhalin fir

seedlings originating from provenances with contrasting snow cover environments. The obtained results

provided the eco-evolutional insights for mechanistical processes in niche acquisition under variable

snow cover environments in boreal regions. The difference in competitive versus conservative soil

environments due to variable snow cover condition may be responsible for the intraspecific variation in

root function to explore soil space. Contrary to our expectation, in low snow cover provenance, the

intraspecific variation was observed in morphological traits in fibrous roots, perhaps reflecting the plastic

responses to growing sites with high snow cover environments. Since we did not reveal the critical factors

explaining why root developmental patterns varied between original provenances, future studies are

expected to find the genetic evidence associated with the intraspecific variation of spring root dynamics.

The insights should enhance the understanding of the adaptive phenology in trees to changing

environments.

Acknowledgements: We thank the staffs of Hokkaido Research Organization for maintaining the

common garden trial and its plant materials and helping the measurement. The authors declare no

conflicts of interest associated with this manuscript. The datasets analyzed during the current study are

available from the corresponding author on reasonable request. TS conceived and managed the study

project. WI provided and managed the experimental materials and filed. TS, WI, TM, IE, JI, MK, NM

conducted the field samplings with the methodology of sampling and survey developed by MK and NM.

All authors analyzed the samples. TS statistically analyzed the data and wrote the draft of manuscript, and

all other authors provided the editorial advice.

Literature Cited

Ambroise, V, Legay, S, Guerriero, G, Hausman, JF, Cuypers, A, Sergeant, K (2020) The roots of plant

frost hardiness and tolerance. Plant and Cell Physiology, 61:3–20 doi: 10.1093/pcp/pcz196

Bardgett, RD, Bowman, WD, Kaufmann, R, Schmidt, SK (2005) A temporal approach to linking

aboveground and belowground ecology. Trends in ecology & evolution, 20:634–641 doi:

10.1016/j.tree.2005.08.005

Begon, M, Townsend, CR, Harper, JL (2006) Resources. In *Ecology: from individuals to ecosystems*.

Wiley, New York, Chapter 3, pp.65–101.

Broadbent, AA, Snell, HS, Michas, A, Pritchard, WJ, Newbold, L, Cordero, I, Goodall, T, Schallhart, N,

Kaufmann, R, Griggiths, RI, Schloter, M, Bahn, M, Bardgett, RD (2021) Climate change alters

temporal dynamics of alpine soil microbial functioning and biogeochemical cycling via earlier

snowmelt. The ISME journal, 15:2264–2275 doi: 10.1038/s41396-021-00922-0

Camm, EL, Harper, GJ (1991) Temporal variations in cold sensitivity of root growth in cold-stored white

spruce seedlings. Tree Physiology, 9:425–431 doi: 10.1093/treephys/9.3.425

Csilléry, K, Ovaskainen, O, Sperisen, C, Buchmann, N, Widmer, A, Gugerli, F (2020) Adaptation to local

climate in multi-trait space: evidence from silver fir (*Abies alba* Mill.) populations across a

heterogeneous environment. Heredity, 124:77–92 doi: 10.1038/s41437-019-0240-0

De Kort, H, Vander Mijnsbrugge, K, Vandepitte, K, Mergeay, J, Ovaskainen, O, Honnay, O (2016)

Evolution, plasticity and evolving plasticity of phenology in the tree species *Alnus glutinosa*. Journal

of evolutionary biology, 29:253–264 doi: 10.1111/jeb.12777

de Kroon, H, Visser, EJ (2003) Distribution of Roots in Soil, and Root Foraging Activity. In *Root ecology*.

Springer, Berlin, Chapter 2, pp.33–60. doi: 10.1007/978-3-662-09784-7_2

De Lisle, SP, Mäenpää, MI, Svensson, EI (2022) Phenotypic plasticity is aligned with phenological

adaptation on both micro□and macroevolutionary timescales. Ecology letters, 25:790–801 doi:

10.1111/ele.13953

Ding, Y, Schiestl-Aalto, P, Helmisaari, HS, Makita, N, Ryhti, K, Kulmala, L (2020) Temperature and

moisture dependence of daily growth of Scots pine (*Pinus sylvestris* L.) roots in Southern Finland.

Tree Physiology, 40:272–283 doi: 10.1093/treephys/tpz131

Eiga, S (1984) Ecological study on the freezing resistance of Saghalin fir (*Abies sachalinensis* Mast.) in

Hokkaido. Bull. For. Tree Breed. Inst. 2:61–107 (in Japanese with English abstract)

359 Ellington, EH, Bastille☐Rousseau, G, Austin, C, Landolt, KN, Pond, BA, Rees, EE, Robar, N, Murray,

360 DL (2015) Using multiple imputation to estimate missing data in meta☐regression. Methods in

361 Ecology and Evolution, 6:153-163. doi: 10.1111/2041-210X.12322

362 Forrest, J, Miller-Rushing, AJ (2010) Toward a synthetic understanding of the role of phenology in

363 ecology and evolution. Philosophical Transactions of the Royal Society B: Biological

364 Sciences, 365:3101–3112 doi: 10.1098/rstb.2010.0145

365 Franks, SJ, Weber, JJ, Aitken, SN (2014) Evolutionary and plastic responses to climate change in

366 terrestrial plant populations. Evolutionary applications, 7:123–139 doi: 10.1111/eva.12112

367 Fréchette E, Chang CYY, Ensminger I (2020) Variation in the phenology of photosynthesis among

368 eastern white pine provenances in response to warming. Global Change Biology, 26:5217–5234 doi:

369 10.1111/gcb.15150

370 Fukuzawa, K, Tateno, R, Ugawa, S, Watanabe, T, Hosokawa, N, Imada, S, Shibata, H (2021) Timing of

371 forest fine root production advances with reduced snow cover in northern Japan: implications for

372 climate-induced change in understory and overstory competition. Oecologia, 196:263–273 doi:

373 10.1007/s00442-021-04914-x

374 Ghalambor, CK, McKay, JK, Carroll, SP, Reznick, DN (2007) Adaptive versus non☐adaptive phenotypic

375 plasticity and the potential for contemporary adaptation in new environments. Functional ecology,

376 21:394–407 doi: 10.1111/j.1365-2435.2007.01283.x

Ishizuka, W, Kon, H, Kita, K, Kuromaru, M, Goto, S (2021) Local adaptation to contrasting climatic

conditions in Sakhalin fir (*Abies sachalinensis*) revealed by long-term provenance trials. Ecological

Research, 36:720–732 doi: 10.1111/1440-1703.12232

Liu, H, Wang, H, Li, N, Shao, J, Zhou, X, van Groenigen, KJ, Thakur, MP (2022) Phenological

mismatches between above-and belowground plant responses to climate warming. Nature climate

change, 12:97–102 doi: 10.1038/s41558-021-01244-x

Ma, T, Parker, T, Unger, S, Gewirtzman, J, Fetcher, N, Moody, ML, Tang, J (2022) Responses of root

phenology in ecotypes of *Eriophorum vaginatum* to transplantation and warming in the Arctic.

Science of The Total Environment, 805, 149926 doi: 10.1016/j.scitotenv.2021.149926

Makoto, K, Wilson, SD, Sato, T, Blume☐Werry, G, Cornelissen, JH (2020) Synchronous and

asynchronous root and shoot phenology in temperate woody seedlings. Oikos, 129:643–650 doi:

10.1111/oik.06996

Makoto, K, Templer, PH, Katayama, A, Seki, O, Takagi, K (2022) Early snowmelt by an extreme

warming event affects understory more than overstory trees in Japanese temperate forests. Ecosphere,

13:e4182 doi: 10.1002/ecs2.4182

McCormack, ML, Gaines, KP, Pastore, M, Eissenstat, DM (2015) Early season root production in

relation to leaf production among six diverse temperate tree species. Plant and Soil, 389:121–129 doi:

10.1007/s11104-014-2347-7

Mediavilla, S, Escudero, A (2009) Ontogenetic changes in leaf phenology of two co-occurring

Mediterranean oaks differing in leaf life span. Ecological Research, 24:1083–1090 doi:

10.1007/s11284-009-0587-4

Nakahata, R (2020) Pioneer root invasion and fibrous root development into disturbed soil space observed

with a flatbed scanner method. Trees, 34:731–743 doi: 10.1007/s00468-020-01953-4

Nissen, J, Donatello, R, Van Dusen, B (2019) Missing data and bias in physics education research: A case

for using multiple imputation. Physical Review Physics Education Research, 15:020106. doi:

10.1103/PhysRevPhysEducRes.15.020106

Noordermeer, D, Velasco, V, Ensminger, I (2021) Autumn warming delays downregulation of

photosynthesis and does not increase risk of freezing damage in interior and coastal Douglas-fir

seedlings. Frontiers in Forest Global Change, 4:688534 doi: 10.3389/ffgc.2021.688534

Persson, H (1978) Root dynamics in a young Scots pine stand in Central Sweden. Oikos, 30:508–519 doi:

10.2307/3543346

Pregitzer, KS, King, JS, Burton, AJ, Brown, SE (2000) Responses of tree fine roots to temperature. New

Phytologist, 147:105–115 doi: 10.1046/j.1469-8137.2000.00689.x

R Core Team (2019) R development core team. R: a language and environment for statistical computing,

55:275–286.

Sakai, A (1983) Comparative study on freezing resistance of conifers with special reference to cold

adaptation and its evolutive aspects. Canadian Journal of Botany, 61:2323–2332 doi: 10.1139/b83-255

Sakai, A, Larcher, W (1987) Frost Resistance in Plants. In: *Frost Survival of Plants.* Springer, Berlin,

Chapter 6, pp.138–173.

Sanmiguel-Vallelado, A, Camarero, JJ, Morán-Tejeda, E, Gazol, A, Colangelo, M, Alonso-González, E,

López-Moreno, JI (2021) Snow dynamics influence tree growth by controlling soil temperature in

mountain pine forests. Agricultural and Forest Meteorology, 296:108205 doi:

10.1016/j.agrformet.2020.108205

Satoh, T (1995) Fundamental studies on the root growth of trees in Hokkaido. Bulletin of the Hokkaido

Forestry Research Institute, 31:1–54. (in Japanese with English abstract)

Steele, SJ, Gower, ST, Vogel, JG, Norman, JM (1997) Root mass, net primary production and turnover in

aspen, jack pine and black spruce forests in Saskatchewan and Manitoba, Canada. Tree physiology,

17:577–587 doi: 10.1093/treephys/17.8-9.577

Sugai, T, Ishizuka, W, Watanabe, T (2023) Landscape gradient of autumn photosynthetic decline in *Abies*

sachalinensis seedlings. Journal of Forestry Research, 34:187–195 doi: 10.1007/s11676-022-01592-0

Tamura, A, Oguma, H, Fujimoto, R, Kuribayashi, M, Makita, N (2022) Phenology of fine root and shoot

using high frequency temporal resolution images in a temperate larch forest. Rhizosphere, 22, 100541

doi: 10.1016/j.rhisph.2022.100541

Wang, Y, Mao, Z, Bakker, MR, Kim, JH, Brancheriau, L, Buatois, B, Leclerc, R, Selli, L, Rey, H,

Jourdan, C, Stokes, A (2018) Linking conifer root growth and production to soil temperature and

carbon supply in temperate forests. Plant and Soil, 426:33–50 doi: 10.1007/s11104-018-3596-7

Weemstra, M., Mommer, L., Visser, EJ, van Ruijven, J., Kuyper, TW, Mohren, GM, Sterck, FJ (2016)

Towards a multidimensional root trait framework: a tree root review. New Phytologist, 211:1159-1169.

doi: 10.1111/nph.14003

West-Eberhard, MJ (2003) Development. In: *Developmental plasticity and evolution*. Oxford University

Press, Oxford, Chapter 5, pp.89–90.

Wilcox, H (1954) Primary organization of active and dormant roots of noble fir, *Abies procera*. American

Journal of Botany, 41:812–821 doi: 10.2307/2438547

Yahara, H, Tanikawa, N, Okamoto, M, Makita, N (2019) Characterizing fine-root traits by species

phylogeny and microbial symbiosis in 11 co-existing woody species. Oecologia, 191, 983-993 doi:

10.1007/s00442-019-04546-2

Zadworny, M, Eissenstat, DM (2011) Contrasting the morphology, anatomy and fungal colonization of

new pioneer and fibrous roots. New Phytologist, 190:213–221 doi: 10.1111/j.1469-8137.2010.03598.x

Figure legends

Figure 1: Diagram of representative pioneer roots and fibrous roots in seedlings of *Abies sachalinensis*.

(a) Underground part of a seedling with pioneer roots denoted by pink arrowheads. (b) Individual pioneer

roots with thick and white root tips and fibrous roots with thin and dark-brown root tips. (c-d) Section

images showing internal structure of a pioneer root (c), and a fibrous root (d), which were prepared as a

thickness of 50 µm and stained with iodine solution.

Figure 2: Mass variation at a whole root level between sampling months (APR: April, MAY: May) and

provenances (grey: Eastern provenance, white: Northern provenance). Different letters denote significant

difference in each panel (p <0.05).

Figure 3: Variation in the average length (a) and total root number (b) of pioneer roots between sampling

months and provenances.

Figure 4: Morphological variation between sampling months and provenances in pioneer roots (a-c) and

fibrous roots (d-f), respectively. SRL: specific root length, SRA: specific root area, RTD: root tissue

density.

Figure 5: Average diameter and structural variation between sampling months and provenances in

pioneer roots (a-c) and fibrous roots (d-f), respectively.

(a) A whole root level

(b) Individual root level

(c) Structure of pioneer root

(d) Structure of fibrous root

2 cm

5 mm

300 μm

100 μm

(a) Total root drymass (g)

(b) Mass fraction of pioneer root (%)

(c) Mass fraction of fibrous root (%)

Eastern provenance
Northern provenance

Pioneer root

(a) SRL (m g^{-1})

Fibrous root

(d) SRL (m g^{-1})

(b) SRA (cm^2 g^{-1})

(e) SRA (cm^2 g^{-1})

(c) RTD (g cm^{-3})

(f) RTD (g cm^{-3})

Pioneer root

(a) Average diameter (mm)

(b) Central cylinder (%)

(c) Cortex + Epidermis (%)

Fibrous root

(d) Average diameter (mm)

(e) Central cylinder (%)

(f) Cortex + Epidermis (%)

(a) Average length (cm)

(b) Total root number