Title: Odor and reward-evoked GABAergic neuronal activity in the basal forebrain influences olfactory-

guided behavior in mice

Abbreviated title: Basal forebrain GABA impacts olfactory behavior

Authors: Elizabeth H. Moss[1,3*], Evelyne K. Tantry[1], Elaine Le[1], Katie L. Brandel-Ankrapp[2], Benjamin R.

Arenkiel[1]

Affiliations: [1]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston TX,

97030, USA; [2]Neuroscience Graduate Program, Baylor College of Medicine, Houston, TX, 97030, USA,

[3]Present Address: Department of Anesthesiology and Perioperative Medicine, Oregon Health &

Science University, Portland, OR, 97239, USA

*Correspondence to mosse@ohsu.edu

Key words: Olfaction, Basal Forebrain, Parvalbumin, Somatostatin, go/no-go, decision-making

Pages: 49

Figures: 5

Tables: 3

Word Count: Abstract: 187, Introduction: 650 words, Discussion: 1379

Conflict of Interest Statement: The authors declare no competing financial interests.

Acknowledgements: This work was supported by the NIH through NINDS and the BRAIN Initiative

(U01NS111692 and R01NS078294 to BRA) and NIDCD and the BRAIN Initiative (K99DC019505 to

EHM).

ABSTRACT

Sensory perception relies on the flexible detection and interpretation of stimuli across variable contexts,

conditions, and behavioral states. The basal forebrain is a hub for behavioral state regulation, supplying

dense cholinergic and GABAergic projections to various brain regions involved in sensory processing.

Of GABAergic neurons in the basal forebrain, parvalbumin (PV) and somatostatin (SST) subtypes

serve opposing roles towards regulating behavioral states. To elucidate the role of basal forebrain

circuits in sensory-guided behavior, we investigated GABAergic signaling dynamics during odor-guided

decision-making. We used fiber photometry to record cell type-specific basal forebrain activity during an

odor discrimination task and correlated temporal patterns of PV and SST neuronal activity with olfactory

task performance. We found that while both PV-expressing and SST-expressing GABAergic neurons

were activated by odors, PV neurons were selectively suppressed by reward whereas SST neurons

were activated. Notably, chemogenetic inhibition of BF SST neurons modestly altered decision bias to

favor reward-seeking while optogenetic inhibition of BF PV neurons during odor presentations improved

discrimination accuracy. Together, these results suggest that the bidirectional activity of GABAergic

basal forebrain neuron subtypes distinctly influence perception and decision-making during olfactory

guided behavior.

SIGNIFICANCE STATEMENT

This study reveals distinct roles for basal forebrain GABAergic neurons in odor perception and odor-

guided decision-making. Fiber photometry shows that basal forebrain parvalbumin-expressing neurons

are selectively suppressed by rewards, while somatostatin-expressing neurons are activated,

establishing the unique recruitment of these GABAergic neurons during behavioral reinforcement.

Chemogenetic and optogenetic interventions demonstrate divergent roles for these neuronal subtypes

in reward-seeking behavior and odor perception. This research provides new insights into how

GABAergic neurons in the basal forebrain shape sensory perception and decision-making.

INTRODUCTION

Behavioral states potently influence neural circuit function and control information processing from the level of sensory perception to cognition, learning, and memory. The basal forebrain (BF) is involved in behavioral state regulation, with BF circuits controlling directed attention (Sarter and Bruno, 1999; Sarter et al., 2001; Hasselmo and McGaughy, 2004; Herrero et al., 2008), sensory processing (Parikh et al., 2007; Chaudhury et al., 2009; Pinto et al., 2013; Bendahmane et al., 2016; Gritton et al., 2016; Ogg et al., 2018), feeding behavior (Herman et al., 2016; Patel et al., 2019), reward prediction and salience (Lau and Salzman, 2008; Lin and Nicolelis, 2008; Hangya et al., 2015; Ledbetter et al., 2016; Zhang et al., 2019), learning and memory (Muir et al., 1993; Devore et al., 2015; Harrison et al., 2016; Zheng et al., 2018), and arousal (Semba, 1991; Szymusiak, 1995; Sarter and Bruno, 1999; Szymusiak et al., 2000; Lau and Salzman, 2008; Anaclet et al., 2015; Xu et al., 2015). BF projections robustly target both sensory and cognitive circuits and their degeneration is thought to underlie cognitive deficits in disease (Bigl et al., 1982; Grothe et al., 2010; Ballinger et al., 2016; Chaves-Coira et al., 2016; Gielow and Zaborszky, 2017). During learning, BF responses to sensory stimuli are modulated by reward expectation (Lin and Nicolelis, 2008; Hangya et al., 2015; Zhang et al., 2019), supporting that BF circuits integrate sensory input with internal information about reward expectation. Additionally, inhibition of the BF impairs attention in response to motivationally salient stimuli (Tashakori-Sabzevar and Ward, 2018). Intriguingly, both cholinergic and non-cholinergic BF neurons show distinct activity in response to sensory stimuli, and play different roles in reward-based learning (Hangya et al., 2015; Patel et al., 2019; Böhm et al., 2020; Nunez-Parra et al., 2020; Hanson et al., 2021). Thus, to better understand how BF circuits mediate state-dependent effects on sensory perception and related behaviors, it is necessary to determine the dynamic activity patterns of specific cell types in the BF and how they are modulated across different behavioral contexts.

The mouse olfactory system is an ideal model for examining the influence of BF modulation on sensory perception and decision-making behavior because both the olfactory bulb and olfactory cortex receive significant cholinergic and GABAergic input from the BF (Woolf et al., 1986; Zaborszky et al.,

1986; Nunez-Parra et al., 2013; Sanz Diez et al., 2019; Böhm et al., 2020). Olfactory behaviors,

including odor discrimination and odor-guided decision-making can be measured via convenient

behavioral assays. Studies that have examined the impact of BF neurons on odor processing have

found that inhibiting BF GABAergic neurons impairs fine odor discrimination in a cross-habituation test

(Nunez-Parra et al., 2013), and ablating BF cholinergic neurons impairs discrimination in a go/no-go

task (Zheng et al., 2022). Moreover, both cholinergic and GABAergic BF neurons respond to odors and

reinforcement in go/no-go discrimination tasks (Hangya et al., 2015; Nunez-Parra et al., 2020). We

have previously shown that BF GABAergic neuronal activity peaks during odor presentations and is

suppressed by reward delivery in a go/no-go task – dynamics which mirror changes in BF cholinergic

tone (Hanson et al., 2021). This demonstrates that BF GABAergic neurons are acutely and

bidirectionally regulated during odor discrimination tasks with positive reinforcement. These

experiments, however, did not differentiate subtypes of GABAergic neurons. Towards understanding

how specific BF cell types respond to sensory stimulation, decision-making, and discrimination learning,

we used fiber photometry to record BF PV and SST neuronal activity while mice performed an

olfactory-cued go/no-go discrimination task. Our data reveal distinct patterns of neuronal activity in PV

and SST BF neurons that are uniquely correlated with different aspects of task performance. Further,

chemogenetic inhibition of BF SST neurons modestly increased decision bias in favor of reward-

seeking, while optogenetic inhibition of BF PV neurons during odor presentations improved odor

discrimination. Together these findings highlight the unique contributions of BF PV and SST neurons to

odor-guided task performance and decision making.

RESULTS

BF PV neurons are excited by odor and suppressed by reward delivery

To assess PV neuronal responses during odor discrimination learning, we selectively expressed the

genetically encoded calcium indicator GCaMP8s in BF PV neurons by injecting an AAV expressing

Cre-dependent GCaMP8s (AAV hsyn-FLEX-GCaMP8s) into the BF of PV-Cre mice. At the same time,

we placed a fiberoptic implant over the horizontal limb of the diagonal band of Broca (HDB) in the BF

(**Figure 1A**). Implant and viral targeting were verified via post hoc immunofluorescence imaging

(**Figure 1B, C**). Implanted mice were then trained and tested on a go/no-go paradigm as previously

described (Hanson et al., 2021). During training, mice learned to distinguish an S+ odorant associated

with a water reward from an S- odorant not associated with reward. In this odor-cued go/no-go

discrimination task, there were four trial outcomes depending on the odor presented and the decision to

go or not go for a reward. Retrieving the water reward upon S+ cue presentation was considered a

"Hit". Not retrieving water after S+ presentation was considered a "Miss". Attempting to retrieve water

upon S- cue presentation was a "False Alarm". Finally, initiating a new trial after S- presentation was

considered a "Correct Reject" (**Figure 1D**). Punishment for False Alarm trials consisted of a 4 second

time-out period in which they were unable to initiate another trial. Mice were trained on the go/no-go

task for 10-14 days prior to testing and fiber photometry recording. After training, mice were tested on

novel odor pairings (**Table 1** in methods) while neuronal activity was recorded with fiber photometry. To

quantify behavioral performance and decision making we used the signal detection theory measures of

d' and criterion (C). d' measures the sensitivity of odor discrimination with higher d' values indicating

more accurate discrimination between the S+ and S- odors and chance equaling zero. Criterion

measures the bias in decision-making toward reward seeking (go) or non-reward-seeking (no-go)

behavior with negative C values indicating a bias towards reward-seeking behavior. Notably, in our

go/no-go task, S+ and S- odors are presented with equal frequency so that optimal decision bias in the

task is 0. However, mice performed this task with a significantly negative decision bias (i.e., a bias

towards reward-seeking, C: -0.11 ± 0.05, $p < 0.05$ one sample t test) likely because the positive

possibility of a water reward outweighed the negative possibility of a timeout punishment. Notably, all

mice effectively learned to discriminate new pairs of S+ and S- odors with performance significantly

above chance within 400 trials (d': 2.30 ± 0.17, $p < 0.0001$, $t = 14.56$, $df = 26$, one-sample t test).

To determine HDB PV neuronal activity across phases of the go/no-go task and different trial

outcomes, photometry traces were first aligned to odor port entry times and sorted by trial type. We

observed a distinct pattern of PV neuronal activity over the course of a go/no-go trial, with excitation

during odor presentations across all trial types, and suppression following reward delivery in Hit trials

(**Figure 1E**). The peak in z-scored PV neuronal activity during odor presentation was equivalent across

all four trial types (**Figure 1F**, Hit: 1.73 ± 0.21, Miss: 1.71 ± 0.20, Correct Reject: 1.64 ± 0.20, False

Alarm: 1.68 ± 0.21 z score, p = 0.40, F = 0.93, df = 23, RM one-way ANOVA) suggesting that the

response was not odor specific (S+ vs. S-), predictive of behavioral choice (go vs. no-go), or predictive

of accuracy (correct vs incorrect decisions) on individual trials. To further examine the response of PV

neurons to reward seeking and reinforcement, we realigned Hit and False Alarm trials to the timing of

reward port entry (**Figure 1G**). Quantifying the area under the curve (AUC) for the first 5 seconds after

reward port entry showed that PV neuronal activity was strongly suppressed in Hit, but not False Alarm

trials (**Figure 1H**, Hit: -2.52 ± 0.50, False Alarm: 2.60 ± 0.40 z score*s, p < 0.001, t = 6.88, df = 5,

paired t test), implying that the suppression was a consequence of obtaining the water reward. Notably,

this pattern of odor-evoked excitation and reward-evoked suppression mirrored the changes in both the

cholinergic tone and general GABAergic neuronal activity we have previously described in the HDB

(Hanson et al., 2021). These data indicate that PV neurons in the HDB respond bidirectionally to the

odor cue and positive reinforcement during go/no-go task performance, thus raising the question of

whether PV neuronal activity controls, or is modulated by, task performance across sessions and/or

learning within sessions.

To determine the extent to which PV neuronal activity was related to task performance, we

examined correlations between the odor response magnitude, reward response magnitude, behavioral

performance, and decision making. Using linear regression and Pearson's correlation to examine the

relationship between PV neuronal activity and sensitivity across sessions, we found that d' was

negatively correlated with odor response magnitude in all trial types (**Figure 1I**, Hit slope = -0.41 ± 0.17,

r = -0.44, p < 0.05; Miss slope = -0.42 ± 0.14, r = -0.52, p < 0.01; Correct Reject slope = -0.44 ± 0.17, r

= -0.45, p < 0.05; False Alarm slope = -0.43 ± 0.17, r = -0.45, p < 0.05). The slopes and intercepts

between trial types were not significantly different (slopes: p = .99, F = 0.005, df = 100, pooled slope = -

0.43; intercepts: p = 0.92, F = 0.16, df = 103, pooled intercept = 2.83). These data show that in

sessions where the average odor response among HDB PV neurons was higher, overall performance

on the discrimination task was worse. Using the same analysis to compare decision bias to PV

neuronal activity, we found that C was not correlated with the magnitude of the odor-evoked peak for

any trial type (Hit slope = -0.76 ± 0.5, r = -0.29, p = 0.15; Miss slope = -0.81 ± 0.4, r = -0.36, p = 0.07;

Correct Reject slope = -0.76 ± 0.5, r = -0.28, p = 0.15; False Alarm slope = -0.62 ± 0.5, r = -0.23, p =

0.25). The slopes and intercepts were also not different between trial types (slopes: p = .99, F = 0.03, df

= 100, pooled slope = -0.74; intercepts: p = 0.94, F = 0.12, df = 103, pooled intercept = 1.87). Together

these data help distinguish between neuronal activity patterns related to decision making behavior and

task difficulty (i.e. the discriminability of individual odor pairs). While these data do not indicate whether

the magnitude of the PV odor response influences odor discrimination, they do suggest that the

observed odor-evoked peak in HDB PV neuronal activity is related to odor discriminability, but not

decision-making behavior in this go/no-go task.

Next, we examined whether PV neuronal activity in response to reward delivery was related to

odor discriminability or decision-making behavior. Using the same linear regression and Pearson's

correlation analyses, we found no correlation between d' and the magnitude of the reward-evoked

suppression in Hit or False Alarm trials (**Figure 1J,** Hit slope = -0.37 ± 0.48, r = -0.15, p = 0.45; False

Alarm slope = -0.131 ± 0.52, r = -0.05, p = 0.80). However, we noted a significant difference between

the intercepts of the regressions for Hit and False Alarm trials (Hit intercept = -1.68 ± 1.06, False Alarm

intercept = 2.90 ± 1.13; p < 0.0001, F = 100.4, df = 51), confirming that the difference in AUC between

Hit and False Alarm trials is consistent across sessions with a range of d'. At the same time, we found a

significant negative correlation between C and the magnitude of reward evoked suppression in Hit trials

(**Figure 1K,** slope = -4.08 ± 1.10, r = -0.59, p < 0.01). This correlation was absent in unrewarded False

Alarm trials (slope = -0.84 ± 1.44, r = -0.12 p = 0.57), suggesting that the suppression of PV neuronal

activity after reward delivery is related to decision making-behavior. However, these data do not reveal

whether the greater suppression of BF PV neurons drives more optimal decision-making behavior or

vice versa. Intercepts were also significantly different between Hit and False Alarm trials (Hit intercept =

-2.88 ± 1.10, False Alarm intercept = 2.54 ± 1.44; p < 0.0001, F = 113, df = 51), confirming that the

reward-evoked suppression of PV neuronal activity was present across sessions with a range of

decision biases.

Correlations between session averages of d' and criterion do not help determine whether odor

and/or reward-evoked suppression of PV neurons is modulated by changes in performance over the

course of a session (as performance changes with learning) or whether it is fixed within a session.

While these data establish relationships between behavior and PV neuronal activity across sessions,

we also sought to determine whether and how PV activity patterns were modulated over the course of

odor learning. To examine this, we first subdivided sessions into blocks, including the first 20 trials of an

odor pair, and best blocks, including the block of 20 trials with the highest accuracy in a session. As

expected, accuracy in the best blocks was consistently higher than accuracy in the first blocks, showing

that mice learned to discriminate novel odor pairs over the course of the sessions (**Figure 1L,** best

block – first block = 0.31 ± 0.02, p < 0.0001, t = 18.99, df = 26). Interestingly, we found that odor

evoked response peaks were not different between first and best blocks (**Figure 1M**). However, the

reward-evoked suppression of PV neuronal activity was significantly greater in the best blocks

compared to the first blocks (**Figure 1N, O**, best block – first block = -3.16 ± 0.84, p < 0.001, t = 3.78, df

= 26, paired t test). Together these findings indicate that the reward-related suppression of PV neurons,

but not odor response magnitudes, changed with learning.

SST neurons are excited by both odor presentation and reward delivery

Within the BF, PV neurons receive inhibitory input from neighboring SST neurons. This disinhibitory

circuit contributes to the opposing effects that BF SST and PV neurons have on arousal state (Xu et al.,

2015). Given the nature of this connectivity, we next asked whether inhibition from SST neurons may

contribute to the reward-evoked suppression that we observed in PV neurons. To evaluate how SST

neurons selectively respond during odor discrimination learning, we again used fiber photometry during

the go/no-go task. For this, we selectively expressed GCaMP in BF SST neurons and implanted a fiberoptic over the HDB (**Figure 2A, B**). As with PV-Cre, mice reliably learned S+/S- odor-reward associations (d' = 1.75 $\pm$ 0.26, p < 0.0001, t = 6.65, df = 9, one-sample t test). Aligning the photometry traces to odor port entry times revealed that SST neurons were excited by odor presentation (**Figure 2C**), and that this excitation was equivalent across trial types (**Figure 2D**, Hit = 1.39 $\pm$ 0.24, Miss = 1.46 $\pm$ 0.29, Correct Reject = 1.30 $\pm$ 0.20, False Alarm = 1.25 $\pm$ 0.23, p = 0.06, F = 3.85, df = 31, RM one-way ANOVA). Responses were similar to what we observed in PV neurons upon odor presentation. Notably, however, when traces were aligned to reward port entry (**Figure 2E**) we found that SST neurons were also activated by reward delivery (**Figure 2F**, Hit AUC = 0.21 $\pm$ 0.13, False Alarm AUC = -0.132 $\pm$ 0.14, p < 0.01, t = 5.16, df = 7, paired t test), in contrast to the suppression we observed in PV neurons. These data show that SST and PV neurons in the HDB exhibit opposing responses to reward delivery. Together, they support the hypothesis that SST and PV neurons serve opposing purposes in go/no-go task reinforcement. This raises the question of how SST neuronal activity is correlated with behavioral performance and decision-making, and to what extent the correlations oppose those observed in PV neurons.

With the goal of directly comparing the two primary GABAergic neuronal subtypes in the HDB, we examined correlations between the SST odor response magnitude, reward response magnitude, behavioral performance, and decision making. In contrast to PV neurons, we found no correlation between d' and SST odor response magnitude (Hit slope = 0.12 $\pm$ 0.10, r = 0.17, p = 0.23; Miss slope = 0.12 $\pm$ 0.14, r = 0.13, p = 0.38; Correct Reject slope = 0.09 $\pm$ 0.10, r = 0.12, p = 0.39; False Alarm slope = 0.14 $\pm$ 0.11, r = 0.19, p = 0.19), nor did we observe differences between trial types in the relationship between odor response magnitude and d' (slopes: p = .99, F = 0.04, df = 189, pooled slope = 0.12; intercepts: p = 0.07, F = 2.43, df = 192, pooled intercept = 1.47). However, we found significant positive correlations between odor response magnitude and response bias for Hit, Miss, and False Alarm Trials (**Figure 2G,** Hit slope = 0.42 $\pm$ 0.16, r = 0.30, p < 0,05; Miss slope = 0.47 $\pm$ 0.18, r = 0.31, p < 0.01; Correct Reject slope = 0.29 $\pm$ 0.16, r = 0.22, p = 0.07; False Alarm slope = 0.36 $\pm$ 0.17, r = 0.25, p <

0.05). The slopes and intercepts were not different between trial types (slopes: p = .88, F = 0.22, df =

273, pooled slope = 0.38; intercepts: p = 0.26, F = 1.34, df = 276, pooled intercept = 1.83). These data

imply that a larger odor-evoked response may be correlated with a more optimal decision bias (i.e.

closer to zero). However, without a substantial number of sessions with decision bias above zero, we

cannot determine whether the relationship would continue increasing linearly as criterion values

become suboptimal in the positive direction. Therefore, it remains a question whether larger S+ odor

responses in SST neurons are related to optimal decision-making behavior, or whether larger odor

responses reflect decision biases away from reward seeking behavior.

When traces were aligned to reward port entry, we found a slight negative correlation between

the magnitude of the reward-related response (AUC) and d' across sessions in Hit but not False Alarm

trials (Hit slope = -0.2 $\pm$ 0.09, r = -0.37, p < 0.05, False Alarm slope = 0.005 $\pm$ 0.09, r = 0.01, p = 0.95).

Taken together with our finding that reward-related excitation of SST neurons is larger in Hit trials than

in False Alarm trials, this analysis suggested that the difference between Hit and False Alarm trials was

greatest in sessions with lower overall d' values (indicating worse task performance). Comparing

reward response magnitudes to decision bias, we found no correlation between criterion and reward

response for Hits or False Alarms (Hit slope = -0.17 $\pm$ 0.17, r = -0.14, p < 0.32, False Alarm slope = -

0.14 $\pm$ 0.14, r = -0.13, p = 0.34). However, we did observe a significant difference between the

intercepts of the regressions for Hit vs. False Alarm trials (**Figure 2H,** Hit intercept = -0.02 $\pm$ 0.15, False

Alarm intercept = -0.24 $\pm$ 0.12, p < 0.05, F = 5.60, df = 109), confirming that increased reward

responses in Hit trials compared to False Alarms remained consistent across a range of decision

biases.

Next, to determine how patterns of SST neuronal activity were modulated during the learning of

a new S+/S- odor pair, we examined how odor and reward responses changed within go/no-go

sessions, as mice improved at the discrimination task. To examine performance-related changes, we

subdivided sessions into first blocks and best blocks (as described for PV neuronal recordings), with

first blocks showing significantly lower accuracy (**Figure 2I,** Best block – first block = 0.31 $\pm$ 0.02, p <

0.0001, t = 14.8, df = 30, paired t test). Aligning first and best block Hit trials to trial initiation revealed no

differences in odor-evoked peaks in SST neuronal activity (**Figure 2J**), suggesting that SST odor

responses are determined by the identity of the odor pair, and are not modulated by learning. However,

aligning Hit trials to reward port entry revealed a significant reduction in reward-related excitation of

SST neurons in the best blocks (Best Block AUC: -1.50 ± 0.48) compared to the first blocks (First Block

AUC: 1.305 ± 0.63, p < 0.001, t = 4.43, df = 30) (**Figure 2K, L**). Taken together with the negative

correlation between session average d' values and reward-related excitation (**Supplemental Figure**

2C), these data suggest that reward responses of SST neurons are modulated by performance, with

improvements in discrimination leading to a reduction in reward-related excitation both across odor

pairs with a range of difficulties, and within sessions, as mice learn to discriminate new S+ and S- odor

pairs.

DREADD-mediated inhibition of SST neurons does not impact firing of neighboring BF neurons

We next sought to determine how directly manipulating BF GABAergic signaling impacted odor

discrimination and decision-making behavior. Towards this, we selectively expressed the inhibitory

DREADD hM4Di in BF SST neurons, first confirming that treatment with the DREADD ligand CNO

showed targeted and ligand-dependent inhibition (**Figure 3A-C**). We chose SST neurons for

manipulation because they have been shown previously to make local inhibitory synapses onto each of

the other major cell types within the HDB, including PV GABAergic, glutamatergic, and cholinergic

neurons (Zaborszky and Duque, 2000a; Xu et al., 2015). To determine the direct effect of CNO

treatment on hM4Di expressing SST neurons, we made acute brain slices harboring the HDB for whole

cell electrophysiological recordings. After current clamping mCherry labeled, hM4Di-expressing

neurons (**Figure 3D**) we made stepped current injections from -100 to 180 pA before and after CNO

application (10 μM). Prior to CNO, SST neurons showed fast-spiking patterns with narrow action

potentials as previously described (Anaclet et al., 2018) (**Figure 3E**). As expected, we found that CNO

significantly inhibited action potential firing in hM4Di expressing HDB SST neurons at current steps of

60 pA or greater (**Figure 3F**), though we did not observe systematic hyperpolarization with CNO (ΔVm -

2.18 ± 1.95, p = 0.33) (**Figure 3F, inset**). To examine whether inhibition of SST neurons impacted firing

properties of neighboring HDB neurons, we also patched and current clamped unlabeled, non-hM4Di-

expressing neurons (**Figure 3G**). Unlabeled neurons were slow spiking and susceptible to

depolarization block, suggesting that they represented cholinergic or glutamatergic subtypes (**Figure**

3H). We found that CNO treatment neither reduced, nor enhanced the firing of the non-hM4Di

expressing neurons in the HDB (**Figure 3I**). CNO application did, however, cause depolarization in non-

hM4Di-expressing neurons (ΔVm 3.22 ± 0.73, p < 0.01) (**Figure 3I, inset**). Notably, the observed

depolarization did not alter the overall firing properties of non-hM4Di-expressing neurons. While these

data indicate that reducing the ability of SST neurons to fire action potentials does not significantly

impact the ability of the other HDB neurons to fire action potentials when driven, they do not show how

DREADD-mediated inhibition of SST neurons impacts HDB circuit activity *in vivo* or during behavior.

DREADD inhibition of SST neurons modestly increases decision bias in go/no-go task

To directly examine the role of BF SST neurons in odor discrimination and decision-making behavior,

we next selectively targeted hM4Di expression to BF SST neurons in the HDB, the magnocellular

preoptic nucleus (MCPO), and substantia innominata (SI) – regions forming a complex within the BF

that is densely interconnected with olfactory brain regions (Zheng et al., 2018) (**Figure 4A**). We then

trained mice on the go/no-go odor discrimination task and tested how they performed in sessions with

or without CNO treatment. In a subset of mice, mCherry instead of hM4Di-mCherry was selectively

expressed in SST neurons to serve as controls for CNO treatment. After training, mice were tested on

new S+/S- odor pairs for consecutive days until they either (1) successfully learned the new odor

associations (achieved two consecutive blocks of 85% accuracy), or (2) failed to meet the criteria for

learning three days in a row (**Figure 4B**). Odor pairs were then chosen to span a range of difficulties

and were approximately matched in difficulty between CNO and Vehicle treated sets of odors (Table 2

in methods). Mice were treated with either CNO (3.5 mg/kg) or Vehicle (1% DMSO in saline) for each

day of alternating odor pairs (**Figure 4B**). Sessions across multiple days of the same odor pair received

the same CNO or Vehicle treatment and were concatenated into single sessions from which sensitivity

and decision bias measures (d' and C) were calculated. Despite our earlier finding that discrimination

sensitivity was inversely correlated with reward related excitation in SST neurons, we found that

inhibiting SST neurons did not impact odor discrimination at the level of whole sessions (**Figure 4C,**

Hm4Di CNO-Vehicle = -0.06 $\pm$ 0.0.18, p = 0.74, t = 0.34, df = 7; Control CNO-Vehicle = 0.14 $\pm$ 0.0.19, p

= 0.49, t = 0.75, df = 5, paired t tests). Additionally, inhibiting SST neurons did not impact the rate at

which mice learned new odor associations (**Figure 4D**). These data suggest that odor and reward

evoked excitation of HDB SST neurons does not directly impact odor discrimination or odor

discrimination learning. Notably, however, we found that decision bias was subtly, but significantly

altered by CNO treatment in hM4Di expressing mice (**Figure 4E,** Hm4Di CNO-Vehicle = -0.11 $\pm$ 0.03, p

< 0.05, t = 3.38, df = 7; Control CNO-Vehicle = 0.04 $\pm$ 0.05, p = 0.47, t = 0.78, df = 5, paired t tests).

Thus, when HDB SST neurons were inhibited, mice showed a slightly stronger bias toward reward

seeking behavior (i.e. a more negative C value). This bias towards reward seeking moved decision-

making further away from the optimal decision bias of zero for our go/no-go task. Together, these data

suggest that HDB SST may affect reward-seeking without directly impacting odor discrimination or

learning in the context of a freely moving, self-initiated, go/no-go discrimination task.

Optogenetic manipulations of PV neurons influence odor discrimination but not decision bias

To investigate the role of BF PV neurons in odor discrimination and decision-making, we targeted

conditional ChR2-GFP (AAV ef1a-Flex-hChR2-eYFP) to the BF neurons in the HDB of PV-cre mice and

placed fiber optic implants above the region to selectively activate the PV neurons (**Figure 5A-C**). Mice

were then trained on the go/no-go odor discrimination task while tethered to an optogenetic cable. After

training, mice received one of three optogenetic stimulation conditions during different go/no-go testing

sessions with novel S+/S- odor pairs (**Figure 5D, Table 3**). In the first condition, mice received 5

seconds of 20 Hz stimulation each time they received a reward on a Hit trial (**Figure 5D**, reward-locked

stimulation). In the second condition, mice received 5 seconds of 20 Hz stimulation every 20 seconds

regardless of where they were in a behavioral trial (**Figure 5D**, regular stimulation). In the last control

condition, mice were tethered to a fiberoptic cable but received no stimulation (**Figure 5D**, no

stimulation). Mice were randomized to different combinations of stimulus conditions and odor pairs.

Optogenetic stimulation time locked to reward delivery, during a time when PV neuronal activity is

typically suppressed (**Figure 1 G-J**), did not impact the overall accuracy of odor discrimination

compared to (**Figure 5E**, $p = 0.19$, $F = 1.97$, $df = 26$, RM one-way ANOVA). Optogenetic stimulation

also did not affect decision-bias, as C values across the three stimulation types showed no significant

difference (**Figure 5F,** $p = 0.14$, $F = 2.23$, $df = 26$, RM one-way ANOVA).

To determine the acute impact of BF PV neuronal activity on odor discrimination, we next targeted the

inhibitory light activated chloride channel iChloC (AAV-ef1a-Flex-iChloC-2A-dsRed) to BF PV neurons

(**Figure 5G-I**). We then placed bilateral fiberoptics over the HDB and trained mice to perform the go/no-

go task as described above. After training, mice received one of three optogenetic stimulation

conditions (**Figure 5J**) during different go/no-go testing sessions with the same set of S+/S- odor pairs

used in stimulation experiments (**Table 3**). In the first condition, mice received 500 ms of continuous

stimulation upon entering the odor port and triggering odor delivery on every trial (**Figure 5J**, All). In the

second condition, mice received 500 ms of continuous stimulation on 50% of the trials selected at

random (**Figure 5J,** Random). In the third condition, mice received 500 ms of continuous stimulation

every 20 seconds regardless, analogous to the "regular stimulation" condition for optogenetic

stimulation (**Figure 5J**, Control). Mice were randomized to different combinations of stimulus conditions

and odor pairs. Surprisingly, we found that optogenetically inhibiting BF PV neurons during all odor

presentations within a session improved odor discrimination (**Figure 5K**). D' values were significantly

higher in sessions where BF PV neurons were inhibited during all odor presentations compared to

sessions where only 50% of trials triggered inhibition and sessions where inhibition was not time-locked

to odor presentations ($p < 0.01$, $F = 11.8$, $df = 14$, RM one-way ANOVA). Inhibition of BF PV neurons

did not, however, affect decision-bias (**Figure 5L,** $p = 0.35$, $F = 1.18$, $df = 14$, RM one-way ANOVA)

indicating that the reduction in errors was balanced across miss and false alarm-type errors. Together,

these results highlight distinct roles for different BF GABAergic neuronal subtypes in different aspects

of odor discrimination behavior including odor discrimination and decision making.

DISCUSSION

Particular attention has been paid to the role of BF cholinergic neurons in regulating behavioral

state. BF cholinergic neurons mediate effects of attention and arousal at downstream circuits, and are

disproportionately affected by neurodegenerative diseases like Alzheimer's and Parkinson's (Sarter and

Bruno, 1997; Levin and Simon, 1998; Sarter et al., 2001; Grothe et al., 2010; Parrao et al., 2012). BF

cholinergic neurons respond directly to sensory cues, positive and negative reinforcement during

learning, and changes in behavioral states across multiple timescales (Phillis, 1968; Szymusiak, 1995;

Sarter and Bruno, 1999; Hasselmo and McGaughy, 2004; Parikh et al., 2007; Chaudhury et al., 2009;

Hangya et al., 2015; Xu et al., 2015; Ruivo et al., 2017; Patel et al., 2019). Activity of non-cholinergic

neurons in the BF is also strongly correlated with behavioral state, sensory perception, and

performance on behavioral tasks (Hassani et al., 2009; Devore et al., 2015; Hangya et al., 2015; Xu et

al., 2015; Harrison et al., 2016). Previous studies using auditory, visual, and olfactory cues have found

increased activity in BF neurons in response to conditioned stimuli, positive and negative

reinforcement, and in anticipation of salient stimuli (Lin and Nicolelis, 2008; Hangya et al., 2015; Zheng

et al., 2018; Nunez-Parra et al., 2020). BF neuronal activity is also modulated by expectation, with more

surprising reinforcement eliciting larger increases in firing for both cholinergic (Hangya et al., 2015) and

non-cholinergic BF neurons (Lin and Nicolelis, 2008; Zheng et al., 2018).

In the context of the olfactory system, BF cholinergic and GABAergic neurons send extensive

projections to the olfactory bulb (OB) (Zaborszky et al., 1986). GABAergic projections preferentially

target specific GABAergic interneurons in the OB, where they influence fine odor discrimination, signal

to noise ratios, and the survival of adult born neurons (Böhm et al., 2020; Hanson et al., 2020; de Saint

Jan, 2022). A direct comparison of the impact of BF cholinergic and GABAergic projections on OB

activity reveals that BF GABAergic neurons preferentially increase odor-evoked mitral cell firing and

suppress spontaneous activity, while cholinergic neurons increase spontaneous, sniff-locked, and odor

evoked mitral cell firing (Böhm et al., 2020). In olfactory tasks, both cholinergic and non-cholinergic BF

neurons respond to conditioned stimuli, unconditioned stimuli, and both positive and negative

reinforcement (Hangya et al., 2015; Nunez-Parra et al., 2020; Hanson et al., 2021), with a significant

proportion of BF neurons showing suppression in response to odor stimuli and positive reinforcement

(Nunez-Parra et al., 2020). Additionally, we have recently shown that BF GABAergic activity is

powerfully suppressed after positive reinforcement (Hanson et al., 2021). Previous studies, however,

did not differentiate between subtypes of non-cholinergic neurons and were, therefore, unable to

discern distinct roles of BF GABAergic neuronal subtypes on odor-guided behavior.

Here, we separately targeted PV and SST GABAergic neurons in the BF, with the goal of

determining how they uniquely contribute to BF olfactory discrimination, learning, and decision-making

behavior. We showed that, while both neuronal subtypes are excited by odor stimuli, each subtype

responds in opposite ways to positive reinforcement. Specifically, SST neurons were briefly excited by

reward delivery, while PV neurons showed a sustained suppression after obtaining a reward. The

opposite responses of PV and SST neurons to reinforcement mirrors the opposing roles of PV and SST

neurons in behavioral state regulation. While opposing responses in PV and SST populations may be

driven, in part, by direct inhibition of PV neurons by SST neurons in the BF, the congruency of their

responses to odor stimuli suggest that this relationship is more complicated. It is possible, for example,

that the inhibitory input from SST to PV neurons is gated differently in response to external sensory

inputs than it is for internal state and/or reinforcement information. While our findings do not suggest

specific gating mechanisms, they do reveal a clear divergence in how BF circuits encode sensory

inputs and reinforcement feedback.

The opposing responses of PV and SST neurons to reward delivery are intriguing because they

mirror the opposite responses of BF GABAergic neurons to changes in behavioral state (Xu et al.,

2015; Harrison et al., 2016). This suggests that the two cell types play distinct roles in the response to

positive reinforcement during discrimination learning, though whether each cell type's activity impacts

discrimination or alters decision making is not clear. Correlating the magnitude of each cell type's

reward-related responses to discrimination sensitivity and decision bias provides insight into their

potential roles, if not information about causal mechanisms. Importantly, correlations observed between

neuronal activity and behavioral performance may be due to neuronal activity causing behavioral

changes, behavioral changes causing changes in BF neuronal activity, or a larger contextual or state

change impacting both neuronal activity and behavior. SST neurons are intriguing components of the

BF circuit because they directly inhibit each of the other major cell types in the BF, including

glutamatergic, cholinergic, and PV GABAergic neurons. Notably, these previously described patterns of

synaptic connectivity in the BF were revealed with optogenetic circuit mapping. In all its utility,

optogenetic circuit mapping leverages artificial, channel rhodopsin-mediated stimulation of SST

neurons and, thus, does not indicate the extent to which SST synapses affect the activity of

downstream BF neurons during normal circuit activity. To help better determine the impact of SST

neuronal excitability on other cell types of the BF, we expressed an inhibitory DREADD in SST neurons

and measured action potential firing and resting membrane potentials in both DREADD and non-

DREADD expressing neurons. While CNO treatment effectively inhibited action potential firing of SST

neurons driven by current injections, it did not systematically alter action potential firing in neighboring

non-DREADD expressing neurons. However, CNO treatment did lead to a slight increase in the resting

membrane potential of non-DREADD expressing neurons. This suggests that SST neurons may be

tonically inhibiting neighboring BF neurons, and that DREADD-mediated inhibition of SST neurons

relieves some of this inhibition, raising the resting potential of neighboring neurons.

To investigate whether SST neuronal activity was driving behavioral changes, as suggested by

the correlation analysis, we chemogenetically inhibited SST neurons during a subset of go/no-go

testing sessions. Based on our observed correlations, we predicted that (1) if SST excitation during

odor exposure was driving more optimal decision-making behavior, then inhibiting SST neurons would

lead to larger (more negative) decision biases, (2) if higher reward-related excitation of SST neurons

was causing poor discrimination, then inhibiting SST neurons would improve sensitivity, and (3) if a

reduction in SST reward-related excitation facilitated learning, then inhibiting SST neurons would

increase the rate at which discrimination sensitivity (d') improved within sessions. Comparing go/no-go

discrimination testing sessions in which BF SST neurons were chemogenetically inhibited to sessions

in which BF SST neurons were not inhibited, we found that DREADD-mediated inhibition of SST

neurons led to a modest reduction in criterion scores with no change in discrimination within or across

sessions. Together, these data suggest that SST neuronal activity influences reward-seeking behavior

without altering discrimination ability or discrimination learning directly.

Finally, to investigate whether PV neuronal was driving decision-making behavior or influencing

odor discrimination, we incorporated optogenetic manipulations of BF PV neurons into the go/no-go

assay. Although we had previously found that suppression of PV neuronal activity was correlated with

reward delivery, optogenetic stimulation of the BF PV neurons in the post-reward period did not affect

either decision-bias or odor discrimination. Instead, optogenetic inhibition of PV neurons during odor

presentations (blunting the odor-evoked peak in BF PV neuronal activity) effectively increased session

accuracy without affecting decision bias. These data indicate that inhibiting BF PV neurons during odor

presentations improves odor discrimination, and they agree with the negative correlation we observed

between odor response peak magnitudes and session d' averages (**Figure 1I**). Together, these results

suggest that BF PV activity likely has a direct, causal impact on odor discrimination.

In conclusion, our findings highlight unique activity patterns exhibited by BF GABAergic PV and

SST neurons in response to olfactory stimuli and reward reinforcement, and provide valuable insights

into the processing of odor and reward information within the BF. Our results underscore the nuanced

contributions of distinct BF GABAergic neuronal subtypes to the complex landscape of odor-guided

decision-making behavior. Together, this work advances our understanding of the neural mechanisms

underlying sensory perception and decision-making processes and sheds light on the multifaceted role

played by specific neuronal subpopulations in the BF.

MATERIALS AND METHODS

Mice

Mice were maintained on a 12-h light-dark cycle and were treated in compliance with the ethical guidelines of the US Department of Health and Human Services and approved by IACUC at Baylor College of Medicine. Both male and female mice over 2 months of age were included for all experiments. SST-Cre (JAX#: 13044) and PV-Cre (JAX#: 17320) mice were originally purchased from Jackson Laboratories.

Surgery and viral constructs

Surgical procedures were performed under general anesthesia using isoflurane (4% v/v) induction and maintained under isoflurane with O2 (1-2% v/v) inhalation. Following anesthetic induction, mice were secured to a stereotaxic frame and administered 5 mg/kg of meloxicam prior to surgery. All procedures were carried out using to Angle 2 software (Leica Biosystems) to set and locate coordinates. Craniotomies were performed over the horizontal diagonal band of Broca (HDB) at the coordinates from Bregma: ML +/- 1 mm; AP 0.7 mm; DV -5.4 mm. For fiber photometry, to target the HDB for viral expression of calcium-indicator GCaMP6f, cohorts of PV-Cre (N = 6) and SST-Cre (N = 8) mice were unilaterally injected with 250 nL of AAV2/9 hSyn-Flex-GCaMP8s-WPRE. Following viral injection, a custom fiber optic implant (0.48 na, 200 um core diameter, RWD systems) was placed so that the tip of the fiberoptic was 0.1 mm dorsal to the HDB coordinates used for viral injection. Implants were then fixed with Metabond dental cement (Parkell).

For optogenetic stimulation, PV-Cre mice (N = 6) were bilaterally injected with 250 nL of AAV-Efla-DIO-ChR2(h134R)::EYFP-WPRE-pA (serotype DJ8) at the same coordinates. For optogenetic inhibition, PV-Cre mice (N = 5) were injected at the same coordinates with 250 nL of AAV-ef1a-Flex-iChloC-2A-dsRed (Addgene plasmid: 70762, serotype DJ8, packaged by the BCM Neuroconnectivity Core). Following viral injection, the same fiber optic implants used for photometry were inserted, bilaterally at 5-degree angles

and placed so that the tips of the fiberoptics were 0.3 mm dorsal to the HDB coordinates used for viral

injection. Implants were then fixed with Metabond dental cement (Parkell).

For inhibitory DREADD expression, SST-Cre mice were bilaterally injected with 250 nL of AAV-DIO-

hM4Di-mCherry (N = 8) or AAV-DIO-mCherry (N = 6) at the same HDB coordinates. All mice were given

a 2-week recovery period prior to the start of behavioral testing. HDB/MCPO/SI targeting was verified in

all cases with post hoc immunofluorescence imaging of viral expression.

Go/no-go training

Freely moving mice were trained on an olfactory-cued go/no-go discrimination task using olfactory

conditioning boxes (Med Associates) controlled by MedPC software. Conditioning boxes included one

odor delivery port and one water reward port, both equipped with IR beams to record port entries. The

odor delivery port received a constant stream of clean room air, cleared by a vacuum. Odors (Sigma,

Tables 1-3) were injected into the steady air stream by opening valves to one of two odor reservoirs,

allowing the air stream to pass through the head space of the reservoir carrying the volatized odorant.

Reservoirs were 20 mL in volume and included 2.5 mL of odorant diluted to 1% by volume in mineral oil.

Water rewards were dispensed at 5uL in volume in the separate reward port upon reward port entry only

on correctly decided Hit trials. Conditioning boxes were housed within larger behavior isolation boxes

(Med Associates). Box fans in the isolation boxes continuously circulated ambient air away from the

conditioning boxes and out of the isolation boxes to reduce the accumulation of ambient odor.

Over the course of 10-14 days, mice underwent 5 stages of behavior-shaping as previously

described (Hanson et al., 2021). Mice were water restricted to no less than 85% of their body weight

during the 48 hours prior to the start of training. Mice were first trained to seek water rewards, then to

initiate trials by poking their nose into the odor port. Mice were then trained to respond to an S+ odor cue

(1% Eugenol in mineral oil) to receive a water reward. Upon forming this association, the S- odor cue

(1% Methylsalicylate in mineral oil) was introduced, in which mice were trained to refrain from seeking a

reward and instead initiate another trial. Trials required mice to sample the odor cues for 100ms or more

before responding to the cue and were categorized based on the S+/- odor cue and the mouse's

response. When mice were presented with the S+ odor cue, trials were considered "Hits" when mice

sought a water reward within 5s and "Misses" if mice did not enter the reward port. Trials in which the S-

odor cue was delivered were considered "False Alarms" when mice incorrectly attempted to retrieve water

and "Correct Rejects" if mice did not seek water. False alarms resulted in a 4-second time out

punishment. Accuracy was calculated as percent correct trials (hit + correct reject trials) for blocks of 20

trials. For each training session, mice completed 10-20 blocks. Training was considered complete after

mice completed a session with the training odors (Eugenol = S+, Methysalicylate = S-) in which they

achieved higher than 85% accuracy in at least 2 consecutive blocks. Testing sessions followed the same

pattern as the last stage of training, but new odor pairs were introduced, requiring mice to learn new S+

and S- odor associations (Tables 1-3).

Go/no-go testing with fiber photometry

For go/no-go testing with concurrent photometry mice were attached to a fiberoptic cable, connected to

a two-channel fiber photometry recording system (Doric) via a rotary joint in the top of the behavior box.

Odor pairs for photometry experiments were chosen randomly for each mouse and session from the list

in Table 1 so that each odor was never presented on more than one day. The timing of trial initiations

and reward port entries were measured by IR beams in the odor and reward ports. Port entry timings,

odor presentations, and trial outcomes were recorded with (Med Associates). The timing of each odor

presentation was output to the photometry recording system as a TTL pulse lasting the duration of an

odor presentation allowing synchronization of the photometry and behavior. Sessions were excluded if

mice failed to complete at least 100 trials and if the average d' value across the whole session was less

than 0.25.

Table 1: Odors used in go/no-go experiments with fiber photometry

Odor	Molecular Weight (g/mol)	Vapor Pressure (mmHg)	Functional Group	Cat. Number
Isovaleric Acid	102.13	0.44	Carboxylic acid 5C	31024
(+) Carvone	150.22	0.16	Cyclic ketone 10C 1 double bond	124931
(-) Carvone	150.22	0.16	Cyclic ketone 10C 1 double bond	22070
1-Methylpyrrole	81.12	15.00	Cyclic nitrogen	509973
Acetophenone	120.15	0.40	Aromatic ketone	200905
Guaiacol	124.14	0.10	Aromatic ether, alcohol	253200
P-Toyl Acetate	150.17	0.16	Aromatic ester, 7C	307300
Amyl Acetate	130.00	4.00	Ester 7C	504009
1-Octen-3-ol	128.22	0.53	Alcohol, 8C	280518
(+) Limonene	136.23	2.00	Cyclic 10C 2 double bond	183164
(-) Limonene	136.23	2.00	Cyclic 10C 2 double bond	218367
Benzaldehyde	106.12	1.00	Aromatic aldehyde	B1334
Isoamyl Acetate	130.19	4.00	Ester 7C	205508
1,4-Cineole	154.25	1.93	Cyclic oxygen, ether	27393
2-Methypyrazine	94.12	9.69	Pyrazine	75608
2-Ethylphenol	122.16	0.15	Aromatic alcohol	44000
Citral	152.24	0.22	Aldehyde	83007
2,6 Dimethyl-5-heptenal	140.22	0.62	Aldehyde, 8C	238902
alpha-Pinene	136.24	3.00	Cyclic 7C 1 double bond	147524
Cyclohexanol	100.16	0.66	Cyclic alcohol	105899
Allyl Sulfide	114.00	9.22	Sulfide	A35801
1-Butenol	74.12	7.00	Alcohol, 4C	281549
1-Pentanol	88.15	44.6	Alcohol, 5C	138975
2-Heptanol	116.20	1.00	Alcohol, 7C	328804
1-Hexanol	102.18	1.00	Alcohol 6C	H13303
1-Heptanol	116.20	0.22	Alcohol 7C	H2805

1-Octanol	130.23	0.14	Alcohol 8C	297887
Butylbenzene	134.22	1.03	Aromatic 10C	686859
Cyclobutanecarboxylic Acid	100.12	0.18	Cyclic carboxylic acid 5C	C95609
P-Cymene	134.21	1.50	Aromatic 10C	30039
Decane	142.29	1.43	Hydrocarbon	30540
Diethylbenzene	134.22	1.03	Aromatic 10C	D91055
2,3-Hexanendione	114.14	10.00	Ketone, 6C	255801
alpha-Phellandrene	136.24	1.86	Cyclic 10C 2 double bond	285609
Isoamyl Butyrate	158.00	0.95	Ester 9C	206008
Menthone	154.00	0.50	Cyclic ketone 10C	218235
2-Heptanone	114.18	3.00	Ketone 7C	537683
Anisole	108.00	3.54	Aromatic ether	123226

Fiber optic cables (0.48 na, 400 um core diameter) connected light emitting diodes (465 nm

experimental and 405 nm isosbestic excitation wavelengths) to a filter cube that separating excitation

and emission wavelengths (Doric). Excitation wavelengths were directed along a fiber optic tether (0.48

na, 200 um core diameter) toward the mouse through a rotary connector attached to the behavior box.

The same fiber carried emission wavelengths from the mouse to the filter cube, which was then directed

to a femtowatt photodetector (Newport) through a fiber optic cable (0.48 na, 600 um core diameter). 405

and 465 nm excitations were controlled by Doric Studio software in "lock-in" mode and the emission

evoked by each channel was separated into distinct Ca^{2+}-dependent and isosbestic GCaMP traces in

Doric Studio. Traces were then exported to MATLAB where Ca^{2+}-dependent GCaMP traces were scaled

to the isosbestic traces, isosbestic signals were subtracted to remove artifacts, and the subtracted traces

were z-scored. Signals were then aligned to odor presentations and subdivided into individual go/no-go

trials using the TTL output from the behavior apparatus.

Z-scored photometry traces from individual trials within a go/no-go session were sorted based on

trial type and outcome. Additional realignment of photometry traces based on reward port entry time was

carried out in MATLAB using output from MedPC software. Mean traces were calculated for each session

by trial type and averaged across animals and sessions. Error was calculated as the 95% confidence

interval across sessions. Peak odor response values were calculated as the maximum value of the z-

scored photometry trace during the odor presentations on individual trials and then averaged across the

session. Session averages were then averaged for each animal and mouse-averaged values were

compared across trial types using one way ANOVA. For photometric recordings of GCaMP from SST

neurons, the area under the curve (AUC) of the reward-related response was calculated for Hit and False

Alarm traces by summing the z score from reward port entry to 1 second after reward port entry. For

recordings from PV neurons, reward-related suppression was calculated as the sum of the z-scored trace

from reward port entry to 5 seconds after reward port entry. Values were not calculated for Miss and

Correct Reject trials because those trials did not involve reward port entry. For analyses subdividing

sessions to examine within-session learning (**Figure 1L-O, Figure 2I-L**), traces were extracted from

single blocks of 20 trials. The "first block" included the first 20 trials from a session. The "best block" was

considered the block of 20 trials with the highest accuracy. If more than one block achieved the same

level of high accuracy, traces from the first instance of high accuracy were used. AUC values were then

calculated from Hit trials within the "first block" and the "best block" of a session and compared with a

paired t-test.

Go/no-go testing with chemogenetic and optogenetic manipulations

For go/no-go experiments involving DREADD mediated inhibition of SST neurons, odor pairs were

presented in the order shown in Table 2. The same odor pairs were presented on consecutive days until

mice achieved at least two consecutive blocks of 85% accuracy, when the odor pair was considered

learned, or until the mice failed to learn for three consecutive days. Sessions of the same odor pair were

then concatenated into a single session spanning up to three days. Concatenated sessions were

excluded from analysis if mice failed to complete at least 100 trials. Mice were treated each day, 15

minutes before starting the go/no-go task, with either CNO (3.5 mg/kg in 1% DMSO, Tocris) or vehicle

(1% DMSO in saline) by I. P. injection. Mice were habituated to I. P. injections of 1% DMSO during the

last stage of go/no-go training. CNO or vehicle treatment was consistent within each concatenated

session such that each session of the same odor pair (up to three consecutive sessions) was associated

with the same drug treatment (**Figure 4B**). As an additional control, a subset of mice was injected with a

non-DREADD expressing virus (AAV-ef1a-FLEX-mCherry) and subjected to the same experimental

paradigm.

For go/no-go testing with optogenetics stimulation of PV neurons, PV-Cre mice injected with

conditional ChR2 and implanted with fiberoptics were attached to a bifurcated fiber optic cable attached

to commutator at the top of the experimental box that connected to a Doric 473 nm laser. The laser was,

in turn, connected to an AMPI Master-8 pulse stimulator that cued the laser to pulse at 20 Hz for 10 ms

based on either internal programming or specific behavioral inputs from the mice, depending on the

stimulation condition. Laser power was modulated so that the output of the fiber connecting to the

fiberoptic implant was 2 mW. Behavioral sessions were run once per day on weekdays. Each session,

mice performed the go/no-go behavioral task using one of 12 novel S+/S- odor pairs (**Table 3**). Each

session corresponded to one of three different stimulation conditions randomized across sessions and

mice so that different mice received different combinations of odor pairs and stimulation conditions. In

the "reward-locked stimulation" experimental condition, reward delivery triggered the onset of a 5 second

burst of 20 Hz laser light stimulation. In the "regular stimulation" control condition, 5 second pulses of 20

Hz laser light were triggered every 20 seconds regardless of the state of mouse in the behavioral task.

In the "no stimulation" control condition, mice were tethered to the bilateral fiberoptic, but no laser light

was delivered. Each mouse underwent multiple sessions of all three conditions but was only exposed to

each odor pair once.

For go/no-go testing with optogenetics inhibition of PV neurons, PV-Cre mice were injected with

conditional iChloC and implanted with fiberoptics then attached to a bifurcated fiber optic cable and

commutator that connected to a Doric 473 nm laser as for optogenetic stimulation. For iChloc

experiments, an AMPI Master-8 pulse stimulator cued the laser to output a single 500 ms duration pulse

based on either internal programming or specific behavioral inputs from the mice, depending on the

stimulation condition. Behavioral sessions were run once per day on weekdays. Each session, mice

performed the go/no-go behavior task using one of 12 novel S+/S- odor pairs (**Table 3**), similar to

optogenetic stimulation experiments. Each session corresponded to one of three different stimulation

conditions randomized across sessions and mice so that different mice received different combinations

of odor pairs and stimulation conditions. In the "all trials" experimental condition, odor presentations

triggered the onset of the 500 ms pulse of laser light. In the "random trials" odor presentations triggered

laser pulses 50% of the time at random. In the control condition, 500 ms pulses of laser light were

delivered every 20 seconds regardless of the state of the mouse in the behavioral task. Each mouse

underwent multiple sessions of the three conditions but was only exposed to each odor pair once.

Table 2: Odors used in go/no-go experiments with DREADD-mediated inhibition of SST neurons

Pair #	Odor	S+/S-	Treatment	Cat. Number
1	Anisole	S+	Vehicle	123226
	Acetophenone	S-		200905
2	Isoamyl Acetate	S+	CNO	205508
	2-Heptanone	S-		537683
3	2-Methypyrazine	S+	Vehicle	75608
	alpha-Pinene	S-		147524
4	(+) Carvone	S+	CNO	124931
	(-) Carvone	S-		22070
5	(+) Limonene	S+	Vehicle	183164
	(-) Limonene	S-		218367
6	2-Ethylphenol	S+	CNO	44000
	1,4-Cineole	S-		27393
7	1-Butenol	S+	Vehicle	281549
	1-Pentanol	S-		138975
8	1-Hexanol	S+	CNO	H13303

	1-Heptanol	S-		H2805
9	2,3 Hexanendione	S+	Vehicle	255801
	Citral	S-		83007
10	Isoamyl Butyrate	S+	CNO	206008
	Menthone	S-		218235

625 **Table 3: Odors used in go/no-go experiments with optogenetic stimulation and inhibition of PV**

626 **neurons**

Pair #	Odor	S+/S-	Cat. Number
1	Acetophenone	S+	200905
	Anisole	S-	123226
2	Isoamyl Acetate	S+	205508
	Isoamyl Butyrate	S-	206008
3	(+) Carvone	S+	124931
	(-) Carvone	S-	22070
4	(+) Limonene	S+	183164
	(-) Limonene	S-	218367
5	2-Ethylphenol	S+	44000
	1,4-Cineole	S-	27393
6	2-Heptanone	S+	537683
	Menthone	S-	218235
7	2,3 Hexanendione	S+	255801
	Citral	S-	83007
8	2-Methypyrazine	S+	75608
	alpha-Pinene	S-	147524
9	1-Butanol	S+	281549
	1-Pentanol	S-	138975
10	1-Hexanol	S+	H13303

	1-Heptanol	S-	H2805
11	Guaiacol	S+	253200
	Eucalyptol	S-	246506
12	Trans-cinnamaldehyde	S+	239968
	Methyl Acetate	S-	296996

Quantification of go/no-go task performance

Go/no-go task performance was quantified using the signal detection theory measures d prime ($d' =$

$Z(Hit\ Rate) - Z(False\ Alarm\ Rate)$) and criterion ($C = -1(Z(Hit\ Rate) + Z(False\ Alarm\ Rate))$)

(Stanislaw and Todorov, 1999). d' measures the sensitivity of discrimination and reflects overall task

performance. C measures discrimination bias. More negative C values reflect a bias towards reward

seeking, values close to 0 reflect optimal decision bias, and positive values reflect bias away from reward-

seeking, in favor of trial reinitiation. To show change in sensitivity over time within a session, d' was

calculated over a rolling window of 40 trials. Session summary d' values were calculated as the maximum

d' over a 40-trial rolling window. This was done to avoid early trials where mice were learning new S+

and S- associations and later trials where mice may have disengaged with the task. For experimental

and control mice in SST DREADD inhibition experiments, d' and C were calculated for each odor pair

and values were averaged for each mouse separated by treatment condition. Mean values for each

mouse from CNO treatment sessions were compared to mean values from vehicle treatment sessions

by paired t-tests. For optogenetic stimulation and inhibition experiments, d' and C were calculated for

each odor pair and values were averaged for each mouse separated by stimulation conditions.

Stimulation conditions were then compared using repeated measures one-way ANOVA and Tukey

corrections for multiple comparisons.

Electrophysiology

For slice electrophysiological recording experiments, mice (N = 3) were deeply anesthetized with

isoflurane then transcardially perfused with ice-cold artificial cerebrospinal fluid (aCSF) solution

containing (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH2P04, 1 MgCl2, 2 CaCl2, 25 glucose, and 25

bicarbonate (pH 7.3, 295 mOsM). Brains were removed and transferred into ice-cold cutting solution

containing (in mM): 2.5 KCl, 1.25 NaH2PO4, 10 MgSO4, 0.5 CaCl2, 234 sucrose, 11 glucose, and 26

bicarbonate. Cutting solution was continuously bubbled with 95% CO2 / 5% O2. Brains were mounted

with cyanoacrylate glue and immediately submerged in oxygenated cutting solution on a Leica VT1200

vibratome. 300 um coronal sections were made at a cutting speed of 0.1 mm/s. Slices including the

HDB were removed to a slice recovery chamber of oxygenated aCSF at 37C for at least 30 minutes.

Following recovery, slices were slowly returned to room temperature for 30 minutes before recording.

For whole-cell current clamp recordings, slices were submerged in a recording chamber and

continuously perfused with room temperature oxygenated aCSF at ~2 mL/minute. Cells were visualized

with DIC optics (Olympus BX50WI) and fluorescence imaging (Q Capture). Once visualized, cells were

whole-cell patched in current-clamp configuration. Recording electrodes (3–7 megaohms) were pulled

from thin-walled borosilicate glass capillaries (inner diameter: 1.1 mm, outer diameter: 1.5 mm) with a

horizontal micropipette puller (Sutter Instruments). Current-clamp internal solution contained (in mM):

120 K gluconate, 20 HEPES, 10 EGTA, 2 MgATP, and 0.2 NaGTP (with 0.4% biocytin by weight, pH to

7.3 with KOH, 285 mOsM). Recordings were made using PClamp software (Axon) with an Axon

MultiClamp 700B amplifier digitized at 10 kHz (Axon Digidata 1440A). Once a whole cell patch was

achieved, cells were allowed to equilibrate for five minutes before resting membrane potential was

recorded and current injections were performed. Cells were then injected with 1 s current steps from -

100 pA to +180 pA in intervals of 20 pA. Resting membrane potential was recorded and current

injections were repeated three times. After recording current injections at baseline, CNO (10 uM) was

washed added to the bath and allowed to circulate for 10 minutes. Resting membrane potential was

recorded and current injections were repeated three times in the presence of CNO. During all

recordings, access resistance was continuously monitored, and cells where the access resistance

changed by more than 20% were excluded from analysis. N = 5 fluorescently labeled cells and N = 7

unlabeled cells. Traces were then exported to MATLAB where resting membrane potential and spike

counts were calculated with custom scripts.

All cells were dialyzed with 0.4% biocytin for the duration of the recording and patched neurons

were saved for post hoc imaging. After recordings, electrodes were withdrawn slowly allowing the cells

to reseal and form an outside-out patch. Slices were then allowed to equilibrate in the recording

chamber for 5 minutes before being transferred to 4% PFA. After patching and cell-filling, slices were

fixed overnight in 4% PFA at 4º C. Slices were then washed 3x in 0.1% PBS-T for 30 min each. After

washing, slices were incubated in 10% normal goat serum blocking buffer for 2 hours at room

temperature. Slices were then incubated in streptavidin conjugated to Alexa:647 (1:500, Invitrogen)

overnight at 4C. The next day, slices were washed 3x in 0.1% PBS-T for 30 min, then mounted on

glass slides using 500 um spacers (Electron Microscopy Sciences) filled with mounting media without

DAPI (Southern Biotech). Slices were imaged on a Leica SP8 confocal with a 20x objective to verify

colocalization with DREADD expression. Z stacks of filled cells were reconstructed in FIJI.

Immunofluorescence and imaging

For antibody staining of GCaMP in mice used for photometry experiments, mice were deeply

anesthetized then transcardially perfused with PBS followed by 4% PFA. Brains were removed and

immersion fixed in 4% PFA overnight at 4C. Brains were transferred to 30% sucrose and allowed to

equilibrate, then they were frozen and sectioned at 40 um on a cryostat (Leica). Sections were washed

in 0.3% PBS-T, then incubated in a blocking solution composed of 10% normal goat serum, 0.3% PBS-

T, and 3M glycine for 1 hour at room temperature. Following blocking, slices were incubated in primary

antibody (Chicken a GFP, 1:1000, Abcam, ab13970) diluted in blocking buffer overnight at 4C. The next

day slices were washed 3x in 0.1% PBS-T then incubated in secondary antibody (Goat a Chicken:488,

1:1000, Invitrogen, A32931) for 2 hours at room temperature. Slices were then washed 3x in 0.3%

PBS-T, transferred to 0.5x PBS, and mounted on glass slides with DAPI-containing mounting media

(Southern Biotech). For fluorescence imaging of DREADD-mCherry and iChloC-dsRed expression, the

endogenous mCherry and dsRed fluorescence was sufficient for imaging and no antibody staining was

performed. Sections were imaged on a Leica DM6000B epifluorescence microscope. For

characterization of DREADD expression in mice used for behavioral experiments (**Figure 4**), three

equivalent sections including the HDB/MCPO/SI complex were selected from each mouse and the

region of viral expression was outlined manually in FIJI and superimposed onto the matching coronal

section schematic (**Figure 4A**).

Experimental Design and Statistical Analyses

All statical analyses were caried out in Prism (Graphpad). Tests performed along with relevant F, t, and

degrees of freedom are reported in the text. For insignificant comparisons, exact p values are reported.

Significant p values are reported as $p < 0.5$, 0.01, 0.001, or 0.0001 as appropriate. Errors reported in

the text are SEM unless otherwise noted. For comparisons across four trial types where values were

matched by animal and session, repeated measures one-way ANOVAs were used with Tukey

corrections for multiple comparisons. For comparisons across two trial types or groups where values

were matched by animal and session, paired t tests were used. For comparisons to a null hypothesis

value, one sample t tests were used. When mouse average values are reported, overall averages and

errors are calculated from mouse average values. For correlation analyses, correlations are reported as

Pearson's R values. Lines of best fit, confidence intervals, slopes, intercepts, and significance values in

were calculated with simple linear regressions. For analyses comparing CNO and vehicle treatment in

control and experimental animals paired t tests were performed separately for control and experimental

mice to compare CNO to vehicle treated sessions. Data from odor pairs for which a mouse performed

fewer than 100 total trials were excluded from all analyses. Four sessions of go/no-go behavior were

excluded from analyses because of behavioral apparatus malfunctioning and three sessions were

excluded for improper laser parameters.

FIGURES and FIGURE LEGENDS

Figure 1: BF PV neurons are excited by odor and suppressed by reward delivery.

A. Schematic showing injection of Cre-dependent viral construct and fiberoptic implant in PV-Cre mice

(top). Highlighted regions of the BF include the horizontal limb of the diagonal band of broca (HDB), the

magnocellular preoptic nucleus (MCPO) and substantia innominata (SI). An experimental timeline

shows timing of surgery, behavioral training, and behavioral testing with fiber photometry imaging

(bottom). **B.** Example slice image showing the implant tract and expression of GCaMP. **C.** Viral

expression (green) and implant tip localization (dashed line) in the HDB. **D.** Schematic of olfactory-

cued go/no-go behavioral paradigm showing possible stimuli, decisions, and outcomes of individual

trials. **E.** Average z-scored GCaMP traces of the four trial types listed in D overlaid with 95%

confidence intervals (shaded regions). **F.** Peak signal amplitudes of the four trial types. Pastel circles

represent the peak odor response of each testing session. White circles represent the average peak

odor response of each animal. Line shows mean across animals. Error bars show SEM across animals.

G. Average z-scored GCaMP traces for Hit and False Alarm trials aligned to the time of reward port

entry with 95% confidence intervals overlaid (shaded regions). Yellow shaded area represents the time

over which the area under the curve (AUC) was quantified. **H.** Area under the curve (AUC) of Hit and

False Alarm traces across testing sessions. Transparent circles represent the AUC of each testing

session. White circles represent the average AUC of each animal. Lines show mean across animals

and error bars show SEM across animals. *** $p < 0.001$ paired t test across N = 6 animals. **I.**

Correlation of the peak odor responses with discrimination sensitivity (d') by trial type, across sessions.

Circles show values of odor response peaks for all trials of a given type within a session plotted against

the d' value for that session. Lines of best fit (solid lines) are shown for each trial type with 95%

confidence intervals (dashed lines). Hash tags indicate slopes of best fit lines that are significantly

different from zero. # $p < 0.05$, ## $p < 0.01$. Bracketed n.s. indicates that no best fit line slopes or

intercepts are significantly different across trial types. **J.** Correlation of the reward response area under

the curve (AUC) with discrimination sensitivity (d') by reward seeking trial type, across sessions. Circles

show values of odor response peaks for hit and false alarm trials within a session plotted against the d'

value for that session. Lines of best fit (solid lines) are shown for each trial type with 95% confidence

intervals (dashed lines). Slopes of each best fit line are not significantly different from zero (n.s.).

Bracketed asterisks indicate a significant difference between the y intercepts of the best fit lines (*** $p <$

0.001). **K.** Correlation of the reward response AUC with decision bias (C) by reward seeking trial type.

Circles show values of odor response peaks for hit and false alarm trials within a session plotted

against the C value for that session. Lines of best fit (solid lines) are shown for each trial type with 95%

confidence intervals (dashed lines). Hash tags indicate a best fit line slope significantly different from

zero (## $p < 0.01$). Bracketed asterisks indicate a significant difference between the y intercepts of the

best fit lines (*** $p < 0.001$). **L.** Accuracy (% correct trials) from first block of 20 trials in a session

(purple) compared to the block of 20 trials with highest accuracy in a session (yellow). Circles show

values for individual sessions with lines and error bars showing mean and SEM. **** $p < 0.0001$, paired

t test. **M.** Average traces (solid lines), aligned to the time of odor port entry (trial initiation), from hit

trials in first blocks (purple) and highest accuracy blocks (yellow) with 95% confidence intervals (shaded

regions). Horizontal line indicates baseline subtracted z score = 0. **N.** Average traces (solid lines),

aligned to the time of reward port entry, from hit trials in first blocks (purple) and highest accuracy

blocks (yellow) with 95% confidence intervals (shaded regions). Horizontal line indicates baseline

subtracted z score = 0. **O.** Quantification of the reward-related response (AUC) in hit trials from first

(purple) and best blocks (yellow). Circles show values for individual sessions with lines and error bars

showing mean and SEM. *** $p < 0.001$.

Figure 2: BF SST neurons are excited by odor and reward delivery.

A. Example slice image showing the implant tract and expression of GCaMP in the BF. **B.** Viral

expression (green) and fiberoptic implant tip localization (dashed line) in the HDB. **C.** Average z-scored

GCaMP traces of the four trial types across all sessions overlaid with 95% confidence intervals (shaded

regions). **D.** Peak signal amplitudes of the four trial types. Light circles represent the peak odor

response of each testing session. White circles represent the average peak odor response of each

animal with mean and SEM of mouse averages. **E.** Average z-scored GCaMP traces for Hit and False

Alarm trials, aligned to the time of reward port entry, with 95% confidence intervals (shaded regions).

Yellow shaded area represents the time over which the area under the curve (AUC) of the reward

response was calculated. **F.** Area under the curve (AUC) of Hit and False Alarm traces across testing

sessions. Transparent circles represent the AUC of each testing session. White circles represent the

average AUC of each animal with mean and SEM of mouse averages. ** $p < 0.01$ paired t test across N

= 8 animals. **G.** Correlation of peak odor response with decision bias (C) by trial types across sessions.

Circles show values of odor response peaks for all trials of a given type within a session plotted against

the C value for that session. Lines of best fit (solid lines) are shown for each trial type with 95%

confidence intervals (dashed lines). Hash tags indicate a best fit line slope significantly different from

zero (# $p < 0.05$). Bracketed n.s. indicates that no best fit line slopes are significantly different from

each other. **H.** Correlation of the reward response AUC with decision bias (C) by reward seeking trial

type. Circles show values of odor response peaks for hit and false alarm trials within a session plotted

against the C value for that session. Lines of best fit (solid lines) are shown for each trial type with 95%

confidence intervals (dashed lines). Slopes of each best fit line are not significantly different from zero

(n.s.). Bracketed asterisk indicates a significant difference between the y intercepts of the best fit lines

(* $p < 0.05$). **I.** Accuracy (% correct trials) from first block of 20 trials in a session (purple) compared to

the block of 20 trials with highest accuracy in a session (yellow). Circles show values for individual

sessions with lines and error bars showing mean and SEM. **** $p < 0.0001$, paired t test. **J.** Average

traces (solid lines), aligned to the time of odor port entry (trial initiation), from hit trials in first blocks

(purple) and highest accuracy blocks (yellow) with 95% confidence intervals (shaded regions).

Horizontal line indicates baseline subtracted z score = 0. **K.** Average traces (solid lines), aligned to the

time of reward port entry, from hit trials in first blocks (purple) and highest accuracy blocks (yellow) with

95% confidence intervals (shaded regions). Horizontal line indicates baseline subtracted z score = 0.

L. Quantification of the reward-related response (AUC) in hit trials from first (purple) and best blocks

(yellow). Circles show values for individual sessions with lines and error bars showing mean and SEM.

*** $p < 0.001$, paired t test.

Figure 3: DREADD inhibition of SST neurons in slice does not impact firing of neighboring BF

neurons. **A.** Schematic showing bilateral injection of Cre-dependent hM4Di viral construct into basal

forebrain regions including the horizontal limb of the diagonal band of broca (HDB), magnocellular

preoptic nucleus (MCPO), and substantia innominata (SI) of SST-Cre mice. **B.** Example of hM4Di-

mCherry expression in BF with the HDB outlined. **C.** Inset from B showing hM4Di-mCherry expression

in HDB. **D.** Example cell fill and hM4Di-mCherry expression from patched hM4Di-expressing

(mCherry+) SST neuron. **E.** Example responses to current injections in aCSF (top, purple) and after

CNO (10 μM) wash on (yellow middle). Bottom panel shows current injection schematic for example

traces. Traces are from cell shown in D. **F.** For hM4Di-expressing neurons: Quantification of action

potentials fired at each current injection in aCSF (purple) and after CNO wash on (yellow). n = 5 cells.

Circles show mean $\pm$ SEM. ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, paired t test. Inset shows resting

membrane potential of each cell before (purple) and after (yellow) wash on of CNO. **G.** Example cell fill

and hM4Di-mCherry expression from patched neuron not expressing hM4Di. **H.** Example responses to

current injections in aCSF (top, purple) and after CNO (10 μM) wash on (yellow middle). Bottom panel

shows current injection schematic for example traces. Traces are from cell shown in G. **I.** For patched

neurons not expressing hM4Di: Quantification of action potentials fired at each current injection in

aCSF (purple) and after CNO wash on (yellow). Circles show mean $\pm$ SEM. Inset shows resting

membrane potential of each cell before (purple) and after (yellow) wash on of CNO. n = 7 cells. ** $p <$

0.01, paired t test.

Figure 4: DREADD inhibition of SST neurons increases decision bias in the go/no-go task.

A. Quantification of DREADD expression in each animal used for behavioral experiments. Shaded regions represent areas containing cells expressing hM4Di-mCherry across three coronal sections spanning AP from bregma -0.22 – 0.74 mm. Regions are color coded by animal (N = 8). **B.** Timeline of surgery, behavioral training, and behavioral testing with DREADD-mediated inhibition (top) and schematic of the CNO or vehicle injections during testing on different odor pairs (bottom). **C.** Average d' for individual sessions (transparent circles, n = 30-40) and mouse averages (white circles, N = 6-8) with mean and SEM of mouse averages. **D.** Performance of mice in the testing phase quantified as a rolling window of sensitivity (d') across trials. Averages of all CNO-treatment sessions are shown in yellow with 95% confidence intervals (shaded yellow). Averages of all vehicle-treatment sessions are shown in purple with 95% confidence intervals (shaded purple). **E.** Average criterion values for individual sessions (transparent circles) and mouse averages (white circles) with mean and SEM of mouse averages. * $p < 0.05$ paired t test between CNO and vehicle-treated sessions (n = 5 sessions per treatment per mouse) with hM4Di-expressing animals (N = 8 mice). n.s. $p > 0.05$ paired t test

842   between CNO and vehicle-treated sessions (n = 5 sessions per treatment per mouse) with mCherry-

843   expressing control animals (N = 6 mice).

Figure 5: Optogenetic manipulations of PV neurons during odor cue and reward.

A. Schematic showing bilateral injection of Cre-dependent ChR2 viral construct and fiberoptic implant in PV-Cre mice. Green-highlighted regions of the BF include the horizontal limb of the diagonal band of broca (HDB), the magnocellular preoptic nucleus (MCPO) and substantia innominata (SI). **B.** Example slice image showing the implant tract, expression of ChR2-eYFP (green), and DAPI (blue). **C.** Viral expression (green) and implant tip localization (dashed line) in the HDB. **D.** Timeline of surgery, behavioral training, and behavioral testing with optogenetic stimulation (top) and schematic outlining the different optogenetic stimulation experimental conditions (bottom). **E.** Summary d' for individual sessions, calculated as the maximum d' from a 40-trial rolling window over the course of a session (see methods) (transparent circles, n = 19 no stimulation sessions, 18 regular stimulation sessions, and 30 reward-locked stimulation sessions). Mouse averages of summary d' values are shown as white circles with mean and SEM (N = 9 mice, p values from RM one-way ANOVA with multiple comparisons using Tukey correction). **F.** Average criterion values for individual sessions (transparent circles) and mouse averages (white circles) with mean and SEM of mouse averages. **G.** Schematic showing bilateral injection of Cre-dependent iChloC viral construct and fiberoptic implant in PV-Cre mice. Red-highlighted regions of the BF include the horizontal limb of the diagonal band of broca (HDB), the magnocellular preoptic nucleus (MCPO) and substantia innominata (SI). **H.** Example slice image showing the implant tract and expression of iChloC-dsRed. **I.** Viral expression and implant tip localization in the HDB. **J.** Timeline of surgery, behavioral training, and behavioral testing with optogenetic inhibition (top) and schematic outlining the different optogenetic inhibition experimental conditions (bottom). **K.** Summary d' for individual sessions, calculated as the maximum d' from a 40-trial rolling window over the course of a session (transparent circles, n = 8 control sessions, 14 random trial stimulation sessions, and 10 all trial stimulation sessions). Mouse averages of summary d' values are shown as white circles with mean and SEM (N = 5 mice, Control vs. All p = 0.029, Random vs All p = 0.041, *p < 0.05, RM one-way ANOVA with multiple comparisons using Tukey correction). **L.** Average criterion values for individual sessions (transparent circles) and mouse averages (white circles) with mean and SEM of mouse averages.

REFERENCES

Agostinelli LJ, Geerling JC, Scammell TE (2019) Basal forebrain subcortical projections. Brain Struct

Funct 224:1097–1117.

Anaclet C, de Luca R, Venner A, Malyshevskaya O, Lazarus M, Arrigoni E, Fuller PM (2018) Genetic

activation, inactivation, and deletion reveal a limited and nuanced role for somatostatin-containing

basal forebrain neurons in behavioral state control. Journal of Neuroscience 38:5168–5181.

Anaclet C, Pedersen NP, Ferrari LL, Venner A, Bass CE, Arrigoni E, Fuller PM (2015) Basal forebrain

control of wakefulness and cortical rhythms. Nat Commun 6:8744.

Ballinger EC, Ananth M, Talmage DA, Role LW (2016) Basal forebrain cholinergic circuits and signaling

in cognition and cognitive decline. Neuron 91:1199–1218.

Bendahmane M, Ogg MC, Ennis M, Fletcher ML (2016) Increased olfactory bulb acetylcholine bi-

directionally modulates glomerular odor sensitivity. Sci Rep 6:1–13.

Bigl V, Woolf NJ, Butcher LL (1982) Cholinergic projections from the basal forebrain to frontal, parietal,

temporal, occipital, and cingulate cortices: a combined fluorescent tracer and acetylcholinesterase

analysis. Brain Res Bull 8:727–749.

Böhm E, Brunert D, Rothermel M (2020) Input dependent modulation of olfactory bulb activity by HDB

GABAergic projections. Sci Rep 10:1–15.

Castillo PE, Carleton A, Vincent J-D, Lledo P-M (1999) Multiple and opposing roles of cholinergic

transmission in the main olfactory bulb. Journal of Neuroscience 19:9180–9191.

Chaudhury D, Escanilla O, Linster C (2009) Bulbar acetylcholine enhances neural and perceptual odor

discrimination. Journal of Neuroscience 29:52–60.

Chaves-Coira I, Barros-Zulaica N, Rodrigo-Angulo M, Núñez Á (2016) Modulation of specific sensory

cortical areas by segregated basal forebrain cholinergic neurons demonstrated by neuronal tracing

and optogenetic stimulation in mice. Front Neural Circuits 10:28.

de Saint Jan D (2022) Target-specific control of olfactory bulb periglomerular cells by GABAergic and

cholinergic basal forebrain inputs. Elife 11:e71965.

Devore S, Pender-Morris N, Dean O, Smith D, Linster C (2015) Basal forebrain dynamics during

nonassociative and associative olfactory learning. J Neurophysiol 115:423–433.

Do JP, Xu M, Lee S-H, Chang W-C, Zhang S, Chung S, Yung TJ, Fan JL, Miyamichi K, Luo L (2016)

Cell type-specific long-range connections of basal forebrain circuit. Elife 5:e13214.

Elaagouby A, Ravel N, Gervais R (1991) Cholinergic modulation of excitability in the rat olfactory bulb:

effect of local application of cholinergic agents on evoked field potentials. Neuroscience 45:653–

662.

Fisher RS, Buchwald NA, Hull CD, Levine MS (1988) GABAergic basal forebrain neurons project to the

neocortex: the localization of glutamic acid decarboxylase and choline acetyltransferase in feline

corticopetal neurons. Journal of Comparative Neurology 272:489–502.

Gielow MR, Zaborszky L (2017) The input-output relationship of the cholinergic basal forebrain. Cell

Rep 18:1817–1830.

Goard M, Dan Y (2009) Basal forebrain activation enhances cortical coding of natural scenes. Nat

Neurosci 12:1444–1449.

Gracia-Llanes FJ, Crespo C, Blasco-Ibáñez JM, Nacher J, Varea E, Rovira-Esteban L, Martínez-

Guijarro FJ (2010) GABAergic basal forebrain afferents innervate selectively GABAergic targets in

the main olfactory bulb. Neuroscience 170:913–922.

Gritton HJ, Howe WM, Mallory CS, Hetrick VL, Berke JD, Sarter M (2016) Cortical cholinergic signaling

controls the detection of cues. Proceedings of the National Academy of Sciences 113:E1089–

E1097.

Grothe M, Zaborszky L, Atienza M, Gil-Neciga E, Rodriguez-Romero R, Teipel SJ, Amunts K, Suarez-

Gonzalez A, Cantero JL (2010) Reduction of basal forebrain cholinergic system parallels cognitive

impairment in patients at high risk of developing Alzheimer's disease. Cerebral Cortex 20:1685–

1695.

Hangya B, Ranade SP, Lorenc M, Kepecs A (2015) Central cholinergic neurons are rapidly recruited by

reinforcement feedback. Cell 162:1155–1168.

Hanson E, Brandel-Ankrapp KL, Arenkiel BR (2021) Dynamic cholinergic tone in the basal forebrain

reflects reward-seeking and reinforcement during olfactory behavior. Front Cell Neurosci

15:635837.

Hanson E, Swanson J, Arenkiel BR (2020) GABAergic Input From the Basal Forebrain Promotes the

Survival of Adult-Born Neurons in the Mouse Olfactory Bulb. Front Neural Circuits 14.

Harrison TC, Pinto L, Brock JR, Dan Y (2016) Calcium imaging of basal forebrain activity during innate

and learned behaviors. Front Neural Circuits 10:36.

Hassani OK, Lee MG, Henny P, Jones BE (2009) Discharge profiles of identified GABAergic in

comparison to cholinergic and putative glutamatergic basal forebrain neurons across the sleep–

wake cycle. Journal of Neuroscience 29:11828–11840.

Hasselmo ME (1999) Neuromodulation: acetylcholine and memory consolidation. Trends Cogn Sci

3:351–359.

Hasselmo ME, McGaughy J (2004) High acetylcholine levels set circuit dynamics for attention and

encoding and low acetylcholine levels set dynamics for consolidation. Prog Brain Res 145:207–

231.

Herman AM, Ortiz-Guzman J, Kochukov M, Herman I, Quast KB, Patel JM, Tepe B, Carlson JC, Ung K,

Selever J (2016) A cholinergic basal forebrain feeding circuit modulates appetite suppression.

Nature 538:253–256.

Herrero JL, Roberts MJ, Delicato LS, Gieselmann MA, Dayan P, Thiele A (2008) Acetylcholine

contributes through muscarinic receptors to attentional modulation in V1. Nature 454:1110–1114.

Kim T, Thankachan S, McKenna JT, McNally JM, Yang C, Choi JH, Chen L, Kocsis B, Deisseroth K,

Strecker RE, Basheer R, Brown RE, McCarley RW (2015) Cortically projecting basal forebrain

parvalbumin neurons regulate cortical gamma band oscillations. Proceedings of the National

Academy of Sciences 112:201413625.

Lau B, Salzman CD (2008) Noncholinergic neurons in the basal forebrain: often neglected but

motivationally salient. Neuron 59:6–8.

Ledbetter NM, Chen CD, Monosov IE (2016) Multiple mechanisms for processing reward uncertainty in

951   the primate basal forebrain. Journal of Neuroscience 36:7852–7864.

952   Levin ED, Simon BB (1998) Nicotinic acetylcholine involvement in cognitive function in animals.

953   Psychopharmacology (Berl) 138:217–230.

954   Lin S-C, Nicolelis MAL (2008) Neuronal ensemble bursting in the basal forebrain encodes salience

955   irrespective of valence. Neuron 59:138–149.

956   Ma M, Luo M (2012) Optogenetic Activation of Basal Forebrain Cholinergic Neurons Modulates

957   Neuronal Excitability and Sensory Responses in the Main Olfactory Bulb. The Journal of

958   Neuroscience 32:10105–10116.

959   Mandairon N, Ferretti CJ, Stack CM, Rubin DB, Cleland TA, Linster C (2006) Cholinergic modulation in

960   the olfactory bulb influences spontaneous olfactory discrimination in adult rats. European Journal

961   of Neuroscience 24:3234–3244.

962   Markram H, Segal M (1990) Long-lasting facilitation of excitatory postsynaptic potentials in the rat

963   hippocampus by acetylcholine. J Physiol 427:381–393.

964   McDonald AJ, Muller JF, Mascagni F (2011) Postsynaptic targets of GABAergic basal forebrain

965   projections to the basolateral amygdala. Neuroscience 183:144–159.

966   Muir JL, Page KJ, Sirinathsinghji DJS, Robbins TW, Everitt BJ (1993) Excitotoxic lesions of basal

967   forebrain cholinergic neurons: effects on learning, memory and attention. Behavioural brain

968   research 57:123–131.

969   Nunez-Parra A, Maurer RK, Krahe K, Smith RS, Araneda RC (2013) Disruption of centrifugal inhibition

970   to olfactory bulb granule cells impairs olfactory discrimination. Proceedings of the National

971   Academy of Sciences 110:14777–14782.

972   Nunez-Parra A, Rio C-D, Christian A, Huntsman MM, Restrepo D (2020) The basal forebrain modulates

973   neuronal response in an active olfactory discrimination task. Front Cell Neurosci 14:141.

974   Ogg MC, Ross JM, Bendahmane M, Fletcher ML (2018) Olfactory bulb acetylcholine release

975   dishabituates odor responses and reinstates odor investigation. Nat Commun 9:1–11.

Parikh V, Kozak R, Martinez V, Sarter M (2007) Prefrontal acetylcholine release controls cue detection

on multiple timescales. Neuron 56:141–154.

Parrao T, Chana P, Venegas P, Behrens MI, Aylwin ML (2012) Olfactory deficits and cognitive

dysfunction in Parkinson's disease. Neurodegener Dis 10:179–182.

Patel JM, Swanson J, Ung K, Herman A, Hanson E, Ortiz-Guzman J, Selever J, Tong Q, Arenkiel BR

(2019) Sensory perception drives food avoidance through excitatory basal forebrain circuits. Elife

8.

Phillis JW (1968) Acetylcholine release from the cerebral cortex: its role in cortical arousal. Brain Res

7:378–389.

Pinto L, Goard MJ, Estandian D, Xu M, Kwan AC, Lee S-H, Harrison TC, Feng G, Dan Y (2013) Fast

modulation of visual perception by basal forebrain cholinergic neurons. Nat Neurosci 16:1857–

1863.

Rothermel M, Carey RM, Puche A, Shipley MT, Wachowiak M (2014) Cholinergic inputs from Basal

forebrain add an excitatory bias to odor coding in the olfactory bulb. Journal of Neuroscience

34:4654–4664.

Rothermel M, Wachowiak M (2014) Functional imaging of cortical feedback projections to the olfactory

bulb. Front Neural Circuits 8:73.

Ruivo LMT-G, Baker KL, Conway MW, Kinsley PJ, Gilmour G, Phillips KG, Isaac JTR, Lowry JP, Mellor

JR (2017) Coordinated acetylcholine release in prefrontal cortex and hippocampus is associated

with arousal and reward on distinct timescales. Cell Rep 18:905–917.

Sanz Diez A, Najac M, De Saint Jan D (2019) Basal forebrain GABAergic innervation of olfactory bulb

periglomerular interneurons. J Physiol 597:2547–2563.

Sarter M, Bruno JP (1997) Cognitive functions of cortical acetylcholine: toward a unifying hypothesis.

Brain Res Rev 23:28–46.

Sarter M, Bruno JP (1999) Cortical cholinergic inputs mediating arousal, attentional processing and

dreaming: differential afferent regulation of the basal forebrain by telencephalic and brainstem

afferents. Neuroscience 95:933–952.

Sarter M, Bruno JP, Turchi J (1999) Basal forebrain afferent projections modulating cortical

acetylcholine, attention, and implications for neuropsychiatric disorders. Ann N Y Acad Sci

877:368–382.

Sarter M, Givens B, Bruno JP (2001) The cognitive neuroscience of sustained attention: where top-

down meets bottom-up. Brain Res Rev 35:146–160.

Semba K (1991) The cholinergic basal forebrain: a critical role in cortical arousal. The basal

forebrain:197–218.

Stanislaw H, Todorov N (1999) Calculation of signal detection theory measures. Behavior research

methods, instruments, & computers 31:137–149.

Szymusiak R (1995) Magnocellular nuclei of the basal forebrain: substrates of sleep and arousal

regulation. Sleep 18:478–500.

Szymusiak R, Alam N, McGinty D (2000) Discharge patterns of neurons in cholinergic regions of the

basal forebrain during waking and sleep. Behavioural brain research 115:171–182.

Tashakori-Sabzevar F, Ward RD (2018) Basal forebrain mediates motivational recruitment of attention

by reward-associated cues. Front Neurosci 12:786.

Tingley D, Alexander AS, Quinn LK, Chiba AA, Nitz DA (2015) Cell assemblies of the basal forebrain.

Journal of Neuroscience 35:2992–3000.

Tsuno Y, Kashiwadani H, Mori K (2008) Behavioral state regulation of dendrodendritic synaptic

inhibition in the olfactory bulb. Journal of Neuroscience 28:9227–9238.

Villar PS, Hu R, Araneda RC (2020) Long-range GABAergic inhibition modulates spatiotemporal

dynamics of the output neurons in the olfactory bulb. bioRxiv.

Woolf NJ, Hernit MC, Butcher LL (1986) Cholinergic and non-cholinergic projections from the rat basal

forebrain revealed by combined choline acetyltransferase and Phaseolus vulgaris leucoagglutinin

immunohistochemistry. Neurosci Lett 66:281–286.

Xu M, Chung S, Zhang S, Zhong P, Ma C, Chang W-C, Weissbourd B, Sakai N, Luo L, Nishino S

(2015) Basal forebrain circuit for sleep-wake control. Nat Neurosci 18:1641–1647.

Zaborszky L, Carlsen J, Brashear HR, Heimer L (1986) Cholinergic and GABAergic afferents to the

olfactory bulb in the rat with special emphasis on the projection neurons in the nucleus of the

horizontal limb of the diagonal band. Journal of Comparative Neurology 243:488–509.

Zaborszky L, Duque A (2000a) Local synaptic connections of basal forebrain neurons. Behavioural

brain research 115:143–158.

Zaborszky L, Duque A (2000b) Local synaptic connections of basal forebrain neurons. Behavioural

brain research 115:143–158.

Zant JC, Kim T, Prokai L, Szarka S, McNally J, McKenna JT, Shukla C, Yang C, Kalinchuk A v,

McCarley RW (2016) Cholinergic neurons in the basal forebrain promote wakefulness by actions

on neighboring non-cholinergic neurons: an opto-dialysis study. Journal of Neuroscience 36:2057–

2067.

Zhang K, Chen CD, Monosov IE (2019) Novelty, salience, and surprise timing are signaled by neurons

in the basal forebrain. Current Biology 29:134–142.

Zheng Y, Feng S, Zhu X, Jiang W, Wen P, Ye F, Rao X, Jin S, He X, Xu F (2018) Different subgroups

of cholinergic neurons in the basal forebrain are distinctly innervated by the olfactory regions and

activated differentially in olfactory memory retrieval. Front Neural Circuits 12:99.

Zheng Y, Tao S, Liu Y, Liu J, Sun L, Zheng Y, Tian Y, Su P, Zhu X, Xu F (2022) Basal Forebrain-Dorsal

Hippocampus Cholinergic Circuit Regulates Olfactory Associative Learning. Int J Mol Sci 23:8472.

1047