Time-Resolved Interactome Profiling Deconvolutes Secretory Protein Quality

Control Dynamics

Madison T. Wright[1], Bibek Timalsina[1,2], Valeria Garcia Lopez[2], & Lars Plate[1,2,3,*]

[1]Department of Chemistry, Vanderbilt University, Nashville, TN, USA

[2]Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[3]Lead Contact

*Correspondence: lars.plate@vanderbilt.edu

SUMMARY

Many cellular processes are governed by protein-protein interactions that require tight spatial and temporal regulation. Accordingly, it is necessary to understand the dynamics of protein interactions to fully comprehend and elucidate cellular processes and pathological disease states. To map de novo protein-protein interactions with time-resolution at an organelle-wide scale, we developed a quantitative mass-spectrometry method, time-resolved interactome profiling (TRIP). We apply TRIP to elucidate aberrant protein interaction dynamics that lead to the protein misfolding disease congenital hypothyroidism. We deconvolute altered temporal interactions of the thyroid hormone precursor thyroglobulin with pathways associated with hypothyroidism pathophysiology such as Hsp70/90 assisted folding, disulfide/redox processing, and N-glycosylation. Functional siRNA screening identified VCP and TEX264 as key protein degradation components whose inhibition selectively rescues mutant prohormone secretion. Ultimately, our results provide insight into the temporal coordination of protein homeostasis, and our TRIP method can have broad application in investigating protein folding diseases and cellular processes.

HIGHLIGHTS

- Method to map de novo protein-protein interactions with temporal resolution
- Characterization of the time-resolved interactome of a protein folding disease state

• siRNA screening identifies VCP and TEX264 as major regulators of thyroglobulin

processing

• VCP pharmacological inhibition rescues mutant thyroglobulin secretion

INTRODUCTION

Protein-protein interactions drive functional diversity within cells and are often closely

connected to the observed phenotypes for cellular processes and disease states

(Bludau and Aebersold, 2020). Protein homeostasis (proteostasis) is a critical cellular

process that relies on tightly regulated protein interactions. The proteostasis network

(PN), consisting of protein folding chaperones, trafficking, and degradation components,

maintains the integrity of the proteome by ensuring the appropriate trafficking and locali-

zation of properly folded proteins while recognizing misfolded, potentially detrimental

states and routing them for degradation (Behnke et al., 2016; Karagöz et al., 2019;

Needham et al., 2019; Pohl and Dikic, 2019). The concerted action of hundreds of pro-

teostasis factors is referred to as protein quality control (PQC). Perturbations to the pro-

teostasis network through genetic, age-related, or environmental factors manifest in

several disease states including amyloidosis, neurodegeneration, cancer, and others

(Kuo et al., 2021; Marinko et al., 2021; McDonald et al., 2022; Taldone et al., 2020;

Wright et al., 2021).

Identifying and quantifying protein-protein interactions has been critical for com-

prehending the pathogenesis of these disease states. Approaches include yeast two-

hybrid systems, co-immunoprecipitation coupled with western blot analysis,

Luminescence-based Mammalian IntERactome (LUMIER) assay, as well as affinity pu-

rification – mass spectrometry (AP-MS) (Piette et al., 2021; Rizzolo and Houry, 2019;

Rizzolo et al., 2017; Taipale et al., 2012, 2014; Wright and Plate, 2021). These methods

have been powerful for mapping steady-state proteostasis interactions to disease

states, yet most lack the ability to measure interaction dynamics over time. Proximity la-

beling mass spectrometry (BioID & APEX-MS) has had limited use to spatiotemporally

resolve protein-protein interactions only following protein maturation, as synchronization

of newly synthesized protein populations is challenging to achieve (Lobingier et al.,

2017; Perez Verdaguer et al., 2022). Unnatural amino acid incorporation methods, such

as biorthogonal non-canonical amino acid tagging (BONCAT), pulsed azidohomo-

alanine (PALM), or heavy isotope labeled azidohomoalanine (HILAQ) can identify newly

synthesized proteins but have not focused on a single endogenously expressed protein

or group of proteins in the context of disease (Bagert et al., 2014; van Bergen et al.,

2021; Dieterich et al., 2006; Howden et al., 2013; Ma et al., 2017, 2018; McClatchy et

al., 2015).

Similar DNA and RNA labeling approaches are used to temporally resolve nas-

cent DNA-protein and RNA-protein interactions in cell culture and whole organisms. Iso-

lation of proteins on nascent DNA (iPOND) has revealed the timing of DNA-protein in-

teractions during replication and chromatin assembly (Cortez, 2017; Munden et al.,

2022; Sirbu et al., 2011). Thiouracil crosslinking mass spectrometry (TUX-MS) and viral

cross-linking and solid-phase purification (VIR-CLASP) are used to study the timing of

RNA-protein interactions during viral infection (Kim et al., 2020; Phillips et al., 2016). In

contrast, no methods were previously available to identify de novo protein-protein inter-

actions of newly synthesized proteins in a client specific manner with temporal resolu-

tion, which has motivated our efforts here.

In earlier work, we mapped the interactome of the secreted thyroid prohormone

thyroglobulin (Tg), comparing the WT protein to secretion-defective mutations impli-

cated in congenital hypothyroidism (CH) (Wright et al., 2021). Tg is a heavily post-trans-

lationally modified, 330 kDa prohormone that is necessary to produce triiodothyronine

(T3) and thyroxine (T4) thyroid specific hormones. Tg biogenesis relies extensively on

distinct interactions with the PN to facilitate folding and eventual secretion (Citterio et

al., 2019; Coscia et al., 2020). Our previous results identified topological changes in Tg-

PN interactions among CH-associated Tg mutants compared to WT. Nonetheless, the

lack of temporal resolution precludes more mechanistic discernment of these changes

in PN interactions. While some changes may simply correlate with disease pathogene-

sis, others may be directly responsible for the aberrant PQC and secretion defect of the

mutant Tg variants.

To address this shortcoming, we developed time-resolved interactome profiling

(TRIP) to capture and quantify interactions between Tg and interacting partners

throughout the life cycle of the protein. We found that Tg mutants are characterized by

both discrete changes with select PN components and broad temporal alterations

across Hsp70/90 and disulfide/redox processing pathways. Moreover, we find that these

perturbations are correlated with alterations in interactions with degradation compo-

nents. We coupled our TRIP method with functional siRNA screening and uncovered

that VCP and TEX264 are two key regulators of Tg processing. VCP and TEX264 inhi-

bition or silencing in thyroid cells resulted in rescued secretion of mutant Tg, represent-

ing – to our knowledge – the first restorative approach based upon proteostasis modula-

tion to increase mutant Tg secretion.

RESULTS

TRIP temporally resolves Tg interactions with PQC components

To develop the time-resolved interactome profiling method, we envisioned a two-stage

enrichment strategy utilizing pulsed biorthogonal unnatural amino acid labeling and

functionalization. After pulse labeling with homopropargylglycine (Hpg), samples can be

collected across time points throughout the chase period (Beatty et al., 2006; Kiick et

al., 2001). After conjugating the Hpg alkyne using copper-catalyzed alkyne-azide cy-

cloaddition (CuAAC) the client protein of interest is globally captured and enriched using

immunoprecipitation. The second enrichment step can then utilize a biotin-streptavidin

pulldown to capture the pulsed, Hpg-labeled, and CuAAC conjugated population, en-

riching samples into time-resolved fractions for quantitative liquid chromatography – tan-

dem mass spectrometry (LC-MS/MS) analysis with tandem mass tag (TMTpro) multi-

plexing (Figure 1A) (Li et al., 2020; Thompson et al., 2019).

Thyroglobulin was chosen as model secretory client protein, and we generated

isogenic Fisher rat thyroid cells (FRT) cells that stably expressed WT Tg or mutant vari-

ants (A2234D and C1264R) (Figures S1A and S1B; Table S1). WT Tg is readily

secreted from the FRT cells while the C1264R Tg shows only minimal residual secretion

(~2% of WT when detected after immunoprecipitation) and A2234D is fully secretion de-

ficient, consistent with prior characterization of these variants (Figures S1B-S1D) (Hishi-

numa et al., 1999; Pardo et al., 2009). We first set out to determine if Tg could tolerate

immunoprecipitation after pulse Hpg labeling, dithiobis(succinimidyl propionate) (DSP)

crosslinking, and CuAAC conjugation with a trifunctional tetramethylrhodamine

(TAMRA)-Azide-Polyethylene Glycol (PEG)-Desthiobiotin probe. We showed previously

that a C-terminal FLAG-tag is tolerated by Tg and allows efficient immunoprecipitation,

while DSP crosslinker aids in capturing transient protein-protein interactions that take

place during Tg processing (Wright et al., 2021). We pulse-labeled FRT cells expressing

WT Tg for 4 hours with Hpg, performed DSP crosslinking, CuAAC, and tested our two-

stage enrichment strategy via western blot analysis (Figure 1A). The additional steps

did not significantly affect immunoprecipitation efficiency and allowed for robust purifica-

tion of WT Tg and well validated interacting proteins (Figure 1A, Box 1-Box 3; Figure 1B

and 1C). We benchmarked our TRIP approach by monitoring Tg interactors HSPA5,

HSP90B1, and PDIA4 (Baryshev et al.; Menon et al., 2007; Wright et al., 2021). Moreo-

ver, DSP crosslinking was necessary to capture these interactions after the dual purifi-

cation and stringent wash steps (Figure 1A, Box 4; Figure 1D).

Next, we investigated whether TRIP could temporally resolve interactions with

these PN components. We pulse labeled WT Tg FRT cells with Hpg for 1 hr, followed

by a 3 hr chase in regular media capturing time points in 30-minute intervals and ana-

lyzing via western blot (Figure 2A). For WT Tg, interactions with HSPA5 peaked within

the first 30 minutes of the chase period and rapidly declined, in line with previous obser-

vations, but PDIA4 interactions were not detectable by western blot analysis (Figure 2B)

(Kim and Arvan, 1995; Menon et al., 2007). To test the ability of TRIP to distinguish

temporal differences in PQC interactions across mutant Tg variants, we performed TRIP

on FRT cells expressing C1264R Tg, a known patient mutation implicated in CH (Hishi-

numa et al., 1999; Kanou et al., 2007). For C1264R, interactions with HSPA5 were

highly abundant at the 0 hr time point, peaked later in the chase period at 2 hours, and

then rapidly declined. A similar temporal profile was also observed for HSP90B1. Addi-

tionally, interactions with PDIA4 were detectable for C1264R and were found to gradu-

ally increase throughout the first 1.5 hours of the chase period, before rapidly declining

(Figure 2C). We noticed similar temporal profiles for HSPA5 and PDIA4 to our western

blot analysis, when measured via TMTpro LC-MS/MS as further outlined below (Figure

2D and 2E).

These data highlight the utility of TRIP to not only identify changes in protein in-

teractions over time but monitor how these interactions differ for a given protein of inter-

est across disease states. Moreover, these data corroborate our previous findings that

steady-state interactions with HSPA5, HSP90B1 and PDIA4 are elevated for C1264R

Tg. (Wright et al., 2021).

TRIP identifies altered temporal dynamics associated with Tg processing

We benchmarked the utility of our TRIP approach to temporally resolve previously iden-

tified and novel interactors, as the Tg interactome has not been fully characterized in

native tissue. TRIP was carried out in FRT cells stably expressing WT, A2234D, and

C1264R Tg. We focused on A2234D and C1264R Tg as they present with distinct de-

fects in Tg processing and mutations are localized in structurally distinct domains (Hishi-

numa et al., 1999; Kanou et al., 2007). Following the Hpg pulse-chase labeling scheme

and dual affinity purification, the time-resolved Tg fractions were trypsin/Lys-C digested

and labeled individually with isobaric TMTpro tags. Subsequently, two biological repli-

cate TRIP time courses (0, 0.5, 1, 1.5, 2, and 3 hr) could be pooled into a 16plex

TMTpro sample that was analyzed by LC-MS/MS (Figures 2A and 3A).

Aside from the experimental samples, we utilized a (–) biotin pulldown booster

(or carrier) channel with cells that were pulse-labeled with Hpg, underwent CuAAC func-

tionalization, and immunoprecipitation, but did not undergo biotin-streptavidin enrich-

ment. This booster sample acted to 1) aid in increased peptide/protein identification -

compared to the much lower abundant chase samples; and 2) benchmark Tg interac-

tors in FRT cells compared to our previously published dataset (Figure 3A) (Petelski et

al., 2021; Tsai et al., 2020). When comparing the booster channel to our (–)Hpg nega-

tive control samples, most previously identified Tg interactors were strongly and signifi-

cantly enriched (Figure S2; Table S2). Our dataset in this study showed appreciable

overlap between our previous results in HEK293T cells identifying 75 of the previous

171 Tg interactors, and identifying 198 new interactors (Figure 3B). We noticed several

ribosomal and proteasomal subunits, trafficking factors, and lysosomal components that

were not identified in our previous dataset (Figure S2; Table S2). We then took our list

of previously identified interactors and PQC components found to be enriched in the (–)

biotin pulldown samples compared to (–) Hpg and carried these proteins forward to

time-resolved analysis utilizing the Hpg-chase samples.

To map the time-resolved Tg PN interactome changes, we considered the Hpg

chase samples (0 – 3 hr) and compared the enrichment of Tg interactors to the (–) Hpg

control. The enrichments were normalized to Tg protein levels to account for gradual

changes due to Tg secretion or degradation (Figure 2B). We organized the interactors

according to distinct PN pathways known to influence Tg processing (Figures S3 and

S4; Tables S3-S5). We found that both A2234D and C1264R exhibited prolonged inter-

actions with components across Hsp70/90 and disulfide/redox processing pathways

(Figures 3C and 3D; Figure S4). Particularly, A2234D and C1264R showed increased

interactions with HSPA5, HSP90B1, HYOU1, DNAJC3, and SDF2L1 throughout the

chase period (Figure 3D). Conversely, WT interactions peaked at the 0 hr time point

and consistently tapered off for many of these components, with DNAJC3 and SDF2L1

mainly being enriched at the 0 hr time point (Figure 3D; Figures S3 and S4). We even

noticed altered temporal interactions for previously unidentified components including

EMC1 and EMC4 of the ER membrane complex (EMC) and CCDC47 of the PAT com-

plex, all of which are involved membrane protein processing within the ER (Figures 3E;

Figure S4) (Chitwood and Hegde, 2020; McGilvray et al., 2020a; Shurtleff et al., 2018).

WT Tg interactions with both CCDC47 and EMC1 peak at the end of the chase period,

yet for C1264R this temporal profile is inversed peaking at the 0 hr time point and slowly

tapering off. Nonetheless, it remains unclear why EMC and PAT complex components

might engage with luminal Tg during biosynthesis and folding.

Similar divergent temporal interactions were observed in the case of disulfide/re-

dox processing components. PDIA3, PDIA4, PDIA6, and ERP29 have all been heavily

implicated in Tg processing (Baryshev et al., 2006; di Jeso et al., 2005, 2014; Menon et

al., 2007). While WT interactions with these components showed similar trends as

Hsp70/90 chaperoning components, peaking at the 0 hr time point and consistently de-

creasing, prolonged mutant interactions peaked only at 1-1.5 hr time points for both

A2234D and C1264R (Figure 3F and 3G). Tg engages with several prolyl isomerization

and hydroxylation enzymes, although previous differences between mutant and WT in-

teractions were quite small and disparate across constructs (Wright et al., 2021). With

the utilization of TRIP, we discovered that previously identified interactors such as NID1,

P4HA1, and SERPINH1 show distinct patterns of altered temporal profiles from mutant

to WT Tg (Figure 3H).

TRIP defines link between glycan processing and ER-phagy pathways.

One area of particular interest was the crosstalk and correlation between interac-

tions with glycan processing components and degradation pathways. The link between

the glycosylation state of ER clients and ER-associated degradation (ERAD) is well es-

tablished, whereas more recently defined autophagy at the ER (ER-phagy) or ER to ly-

sosome-associated degradation (ERLAD) represent alternative degradation mecha-

nisms for ER clients (Chiritoiu et al., 2020; Christianson et al., 2008, 2012; Fregno et al.,

2018, 2021). Previously, we showed that A2234D and C1264R differ in interactions with

N glycosylation components, particularly the oligosaccharyltransferase (OST) complex.

Efficient A2234D degradation required both STT3A and STT3B isoforms of the OST,

which mediate co-translational or post-translational N-glycosylation, respectively. TRIP

revealed differential interactions with glycosylation components that may lead to altered

degradation dynamics. Many glycan processing enzymes, lectin chaperones, and sev-

eral subunits of the OST complex, including STT3A and STT3B were identified (Figure

3I, Figure S4). While STT3A interactions across all constructs showed similar temporal

profiles, we observed prolonged interactions for lectin chaperones CALR, CANX with

mutant Tg. The most striking difference observed was with the "gatekeeping" glycosyl-

transferase UGGT1. This protein regulates glycoprotein engagement and folding

through the CANX/CALR cycle by re-glycosylating ER clients, and thus triggering reen-

gagement with the lectin chaperone cycle (Lamriben et al., 2016). UGGT1 interactions

with WT remain moderate from 0.5-1.5 hrs and peak at 3 hrs, while interactions for

A2234D and C1264R peaked earlier and were more pronounced throughout the chase

period. These differential interactions with UGGT1 may suggest changes in monitoring

of the Tg folded state. Moreover, CANX has been directly linked to emerging mecha-

nisms of ERLAD for glycosylated ER clients (Forrester et al., 2019; Fregno et al., 2018).

Proteasomal and autophagic degradation pathways exhibited widely different in-

teractions dynamics for WT and mutant Tg (Figure 3J-3L). The ERAD associated lectin

OS9, and ATPase VCP/p97 both peaked at the 3 hr chase time point for WT Tg (Figure

3K). Conversely, A2234D exhibited early VCP interactions that decreased throughout

the chase, with OS9 interactions peaking at the 2 hr time point. For C1264R, we ob-

served much stronger and prolonged VCP interactions, as well as additional interactions

with the ERAD associated mannosidase EDEM3 and E3 ubiquitin ligase adaptor SEL1L

(Christianson et al., 2008, 2012) (Figure 3K).

Our original dataset identified several lysosomal components, yet it was unclear

how Tg might be delivered to the lysosome. Recent work has identified several selective

ER-phagy receptors, and highlighted ER-phagy mechanisms for the clearance of mu-

tant prohormones and other destabilized clients from the ER (Chen et al., 2021; Cun-

ningham et al., 2019). It was intriguing then to identify several lysosomal and autophagy

related components and observe differential temporal profiles across Tg constructs (Fig-

ure 3L, Figure S4). Both WT and C1264R Tg showed appreciable interactions with lyso-

somal components, while A2234D interactions with these components were more

sparse. Interactions with several lysosomal cathepsin proteases and glycan trimming

enzymes exhibited altered temporal profiles from WT to C1264R. We found that interac-

tions with these components peaked for WT at 1 hr, while C1264R interactions peaked

earlier and remained more stable throughout the chase period (Figure 3L and S4). Yet,

the most intriguing observation was the enrichment of three different ER-phagy recep-

tors, ATL3, CCPG1, and RTN3 between WT and C1264R, along with the RTN3 adaptor

protein PGRMC1 (Chen et al., 2019, 2021; Grumati et al., 2017; Liang et al., 2018;

Smith et al., 2018). ATL3 showed similar interaction patterns between WT and C1264R

with interactions rising and peaking at 1-1.5 hr time points before beginning to taper off.

On the other hand, CCPG1 and RTN3 were found to specifically interact with C1264R,

with RTN3 interactions peaking at 0 hr and then decreasing, while CCPG1 interactions

remained more stable (Figure 3L). Another striking observation was that the temporal

profile of EMC interactions for C1264R correlated with RTN3, PGRMC1, CTSB, and

CTSD interactions (Figure 3E and 3L). This is particularly interesting as there are few

known adaptor or accessory proteins that aid in client specific ER-phagy or ERLAD.

Overall, these data display the ability of our TRIP workflow to assess the tem-

poral dynamics associated with PQC, providing further insight into the dysregulation of

interactions within CH dyshormonogenesis. It is worth noting that many interactors, in-

cluding CANX, ERP29, and VCP amongst others, previously did not show any apprecia-

ble changes in interactions from WT to mutant Tg when using traditional AP-MS meth-

ods, yet TRIP was able to identify and highlight these unseen changes over time

(Wright et al., 2021).

siRNA screening discovers key regulators of construct specific Tg processing

To investigate and better comprehend the functional implications of Tg interactors and

other PQC components we developed an RNA interference screening platform. Moreo-

ver, we hoped to identify factors whose modulation may act to rescue mutant Tg secre-

tion. HEK293 cells were engineered to stably express nanoluciferase-tagged Tg con-

structs (Tg-NLuc) and screened against 167 Tg interactors and related PN components.

The NLuc tag allowed us to monitor changes to both intracellular Tg abundance in the

cell lysates, and Tg levels in the conditioned media to assess secretion rates in a 96-

well format. Importantly, the NLuc-tag did not alter the secretion of WT Tg, and CH-as-

sociated mutants maintained the same secretion deficiency (Figure 4A, Figures S1E-

S1K) (England et al., 2016). We identified 4 genes whose silencing modulated Tg-NLuc

abundance in lysate (Figures 4B, Figure S5A; Table S6). Silencing of NAPA (α-SNAP)

and LMAN1 were found to increase WT Tg-NLuc lysate abundance but had no effects

on the two mutants. α-SNAP is a member of the Soluble N-ethylmaleimide-sensitive

factor Attachment Protein (SNAP) family and plays a critical role in vesicle fusion and

docking, while LMAN1 is a mannose-specific lectin that functions as a glycoprotein

cargo receptor for ER-to-Golgi trafficking (Marinko et al., 2021; Song et al., 2017; Zhao

et al., 2007). For mutant Tg-NLuc constructs we found cathepsin Z (CTSZ) silencing de-

creased A2234D Tg-NLuc lysate abundance, while β-glucoronidase (GUSB) silencing

increased C1264R lysate abundance (Figure 4B, Figure S5A; Table S6).

When monitoring secreted Tg-NLuc levels in the conditioned media, we identified

8 genes whose silencing led to differential changes in Tg-NLuc secretion (Figures 4C,

Figure S5B; Table S6). α-SNAP, neuraminidase 1 (NEU1), and nodal modulator 2

(NOMO2) silencing led to decreases in Tg-NLuc secretion, with α-SNAP & NEU1 affect-

ing WT, and NOMO2 affecting A2234D in a construct specific manner. The decrease in

WT Tg secretion upon α-SNAP silencing complemented the results of increased WT ly-

sate levels, indicating a shift from secretion towards retention. NEU1 is a lysosomal en-

zyme that cleaves terminal sialic acid residues on proteins and lipid species

(Pshezhetsky and Hinek, 2011; Ryuzono et al., 2016). While Tg contains terminal sialic

residues after glycan maturation through the Golgi complex, it is unclear how silencing

of NEU1 might decrease secretion of WT Tg (Citterio et al., 2019). NOMO2 is one of

three close homologs of the NOMO gene believed to share similar functions, with

NOMO recently identified as a component of the TMCO1-ribosome complex/translocon

involved in membrane protein biogenesis and found to be a close interactor with ER-

phagy receptor TEX264 (An et al., 2019; McGilvray et al., 2020b).

Remarkably, we identified 6 genes whose silencing rescued mutant Tg-NLuc se-

cretion in a construct specific manner. α-SNAP silencing increased secretion of A2234D

Tg-NLuc. This contrast to the reduction in WT secretion may suggest an alternative role

for α-SNAP in regulating mutant Tg processing as other proteins involved in vesicular

fusion and trafficking have been implicated in ER-phagy (Cui et al., 2019; Liang et al.,

2020). Silencing of LEPRE1 (P3H1), an ER-resident prolyl hydroxylase, increased

C1264R Tg-NLuc secretion but not WT nor A2234D (Vranka et al., 2004). Silencing of

several protein degradation genes robustly increased mutant Tg secretion. VCP silenc-

ing increased both A2234D and C1264R Tg-NLuc secretion (Figure 4C, Figure S5B;

Table S6). As mentioned previously, VCP is associated with ERAD but also aids in sev-

eral diverse cellular functions including the interplay between proteasomal and lysoso-

mal degradation (Christianson et al., 2008, 2012; Hill et al., 2021). HERPUD1, TEX264

and RTN3 silencing selectively increased C1264R secretion, but did not alter WT nor

A2234D secretion. HERPUD1 is a ubiquitin-like protein and associates with VCP during

ERAD (Christianson et al., 2012; Needham et al., 2019; Okuda-Shimizu and Hender-

shot, 2007). The ER-phagy receptors RTN3 and TEX264 localize to sub domains of the

ER to facilitate degradation of specific ER clients and organellular regions (An et al.,

2019; Chen et al., 2021; Chino et al., 2019; Cunningham et al., 2019; Fielden et al.,

2022; Grumati et al., 2017). Of note, while the ER-phagy receptor CCPG1 was identified

in our mass spectrometry dataset siRNA silencing of CCPG1 did not significantly alter

338 Tg-NLuc abundance in lysate or media, nor did silencing of SEC62 or RETREG1

339 (FAM134B), two additional ER-phagy receptors found to regulate ER dynamics (Figure

340 S5; Table S6).

341  This is the first study to broadly investigate the functional implications of Tg in-

342 teractors and other PQC network components on Tg processing through siRNA screen-

343 ing. Particularly, with many of these components being involved in protein degradation

344 these results may suggest that the blockage of degradation pathways can broadly res-

345 cue the secretion of A2234D and C1264R mutant Tg, a phenomenon similarly found in

346 other protein folding diseases (Vij et al., 2006).

348 **Trafficking and degradation factors selectively regulate Tg processing in thyroid**

349 **cells**

350 We examined the hits from the siRNA screen in FRT cells stably expressing Tg con-

351 structs to test whether their silencing exhibited similar phenotypes in thyroid specific tis-

352 sue. First, WT Tg FRT cells were transfected with siRNAs targeting LMAN1 and NAPA

353 (α-SNAP), and changes in Tg lysate and media abundance were monitored. Silencing

354 of α-SNAP led to a ~50% increase in WT Tg lysate abundance, while α-SNAP and

355 LMAN1 silencing both led to marginal decreases in WT Tg secretion after 4 hrs, con-

356 sistent with the results in HEK293 cells (Figure 5A and 5B). ^{35}S pulse-chase analysis

357 also confirmed that α-SNAP silencing significantly increased lysate retention by 15%

358 and decreased secretion by 18% (Figure S7A and S7B).

Next, C1264R Tg FRT cells were transfected with siRNAs targeting VCP, RTN3,

TEX264, HERPUD1, and LEPRE1 (P3H1) and lysate and media Tg abundance were

similarly monitored. RTN3, TEX264, and HERPUD1 silencing marginally, yet signifi-

cantly decreased C1264R lysate abundance (Figure 5C and 5D). Moreover, VCP and

TEX264 silencing significantly increased C1264R secretion by 3-fold and 2-fold respec-

tively after 8 hrs, in line with our results in HEK293 cells (Figures 5C and 5D). In con-

trast, silencing of RTN3 significantly decreased C1264R Tg secretion by 5-fold, opposite

to the increased secretion observed in HEK293 cells. Efficient silencing of all proteosta-

sis factors in FRT cells was also confirmed (Figure S6).

Mutant Tg is selectively enriched with the ER-phagy receptor TEX264

Intrigued by the finding that TEX264 silencing increased C1264R Tg secretion without

affecting WT Tg, we asked whether TEX264 exhibited differential interactions with Tg

constructs. HEK293 Tg-NLuc cells were transfected with either a fluorescent GFP con-

trol, or C-terminal FLAG-tagged ER-phagy receptors SEC62 and TEX264 and FLAG co-

immunoprecipitations (co-IP) were performed. We then used the NLuc assay and West-

ern blotting to monitor Tg enrichment in the SEC62 or TEX264 co-IP (Figure 6A).

SEC62 silencing did not affect the processing of any Tg constructs in our siRNA screen

and therefore should act as a negative control (Figure S5; Table S6). We found that co-

IP of SEC62 did not yield any appreciable enrichment of Tg compared to GFP control.

Conversely, co-IP of TEX264 resulted in the enrichment of all Tg constructs compared

to GFP control when monitored by NLuc assay, with C1264R and A2234D being

significantly enriched. C1264R Tg exhibited a 3-fold increase in interactions compared

to WT Tg and almost a 2-fold increase compared to A2234D Tg (Figure 6B). This in-

crease in C1264R enrichment was also observable by western blot analysis (Figure

6C).

Together, these data suggest that not only does TEX264 play a role in Tg pro-

cessing, but CH-associated Tg mutants are selectively targeted to TEX264. Further-

more, ER-phagy should be considered a major degradative pathway in Tg processing,

as other studies have mainly focused on Tg degradation through ERAD (Menon et al.,

2007; Tokunaga et al., 2000).

Pharmacological VCP Inhibition Selectively Rescues C1264R Tg Secretion

Considering the exciting finding that silencing of degradation factors by siRNA rescued

C1264R secretion, we sought to investigate whether pharmacological modulation of se-

lect Tg processing components could similarly improve Tg secretion. There are no se-

lective inhibitors currently available for TEX264, but several for VCP. We monitored

C1264R Tg secretion from FRT cells in the presence of VCP inhibitors ML-240, CB-

5083, and NMS-873 and found treatment with ML-240 was able to significantly increase

C1264R Tg secretion, corroborating our siRNA silencing data (Figure 7A and 7B) (Chou

et al., 2013, 2014; Zhou et al., 2015). Further, VCP inhibition in FRT cells with ML-240

did not alter C1264R Tg lysate abundance. To investigate whether this rescue of

C1264R Tg secretion with ML-240 treatment was specific to C1264R Tg, we also moni-

tored WT Tg abundance. In contrast, we observed that ML-240 significantly reduced

WT Tg abundance in both lysate and media 4 and 8 hrs after treatments (Figures 7C

and 7D).

We hypothesized that ML-240 treatment may differentially regulate Tg degrada-

tion and processing in a construct specific manner. We turned to a ^{35}S pulse-chase as-

say to fully characterize Tg degradation and processing dynamics with ML-240 treat-

ment and found that lysate abundance of C1264R Tg was not significantly changed at

the 4 hr time point with ML-240 treatment compared to DMSO (Figure 7E; Figure S7C

and S7D). This paralleled the previous results with VCP silencing, and ML-240 treat-

ment under steady-state conditions in FRT cells. Excitingly, we found a 10-fold increase

in C1264R Tg secretion at the 4 hr time point with ML-240 treatment compared to

DMSO with no significant change in C1264R Tg degradation in the presence of ML-240

(Figure 7E, Figure S7D). Conversely, for WT Tg ML-240 treatment led to a significant

increase in lysate accumulation at 4 hrs compared to DMSO (Figure 7F; Figure S7E

and S7F). This increase in lysate abundance correlated with a significant decrease in

WT Tg secretion from 67% to 4% in the presence of ML-240 compared to DMSO, with-

out altering WT degradation (Figure S7F).

Pharmacological VCP Inhibition is Associated with Temporal Remodeling of the

Tg Interactome

We were curious if our TRIP method was capable of capturing changes in interactions

across PQC pathways that mediate the rescued secretion of C1264R Tg. Hence, TRIP

of C1264R Tg was carried out in the presence of ML-240 to monitor temporal changes

across key PQC pathways (Figures S8A-S8C; Tables S4 and S5). Both glycan pro-

cessing and Hsp70/90 chaperoning pathways exhibited broad decreases in C1264R Tg

interactions in ML-240 treated samples compared to the untreated controls (Figure 7G

and 7H). Particularly, CANX and UGGT1 interactions tapered off more rapidly within 0.5

hr compared to untreated C1264R. In contrast, interactions with key Hsp70/90 chaper-

ones components remained relatively steady throughout the chase period before peak-

ing at the 3 hr time point (Figure 7I and 7J). Conversely, C1264R interactions with co-

chaperones DNAJB11 and DNAJC10 decreased and mimicked the WT Tg temporal

profile (Figure 3D and 7J).

Interactions with disulfide/redox processing and prolyl isomerization and hydrox-

ylation components exhibited milder but marked declines at intermediate time points

with ML-240 treatment compared to untreated samples (Figure 7K and 7L; Figure S8C).

PDIA6 and TXNDC12 interactions peaked at the 0 hr time point and tapered off

throughout the chase period, while PDIA4 interactions remained relatively low before

peaking at the 3 hr time point (Figure 7L). Conversely, ERP29 and PDIA3 interactions

remained largely unchanged in the presence of ML-240. These more drastic changes in

glycan processing and Hsp70/90 chaperone interactions may highlight that extensive

early interactome remodeling during N-glycosylation and Hsp70/90 dependent folding

could be a key rescue mechanism. Conversely, remodeling with disulfide/redox pro-

cessing components during secretion rescue of C1264R involves more discrete

changes in interactions, and prolyl isomerization and hydroxylation likely play a minor

role during this secretion rescue.

We monitored the activation of UPR markers in the presence of ML-240 and

found that 3 hr treatment with ML-240 left HSPA5 and ASNS expression unchanged in

C1264R Tg cells but led to a significant decrease in WT Tg cells. Only DNAJB9 tran-

scripts showed a small, but significant increase in both WT Tg and C2164R Tg FRT

cells (Figure S8D). Moreover, ML-240 did not significantly alter cell viability after 3

hours, as measured by propidium iodide staining (Figure S8E-S8G). Overall, these re-

sults suggest that ML-240 can selectively rescue C1264R Tg secretion, and that this

rescue is correlated with unique temporal changes in C1264R Tg PN interactions, but

not global remodeling of the ER proteostasis network due to UPR activation.

DISCUSSION

The timing of protein-protein interactions implicated in cellular processes and patho-

genic states remains pivotal to our understanding of disease mechanisms. Nonetheless,

the temporal measurement of protein-protein interactions has remained difficult and

proven to be a bottleneck in elucidating the coordination of complex biological pathways

such as the proteostasis network. Here, we have presented an approach utilizing a two-

stage enrichment strategy to temporally resolve protein-protein interactions implicated in

PQC. Furthermore, we utilized a functional genomic screen to further characterize and

investigate the implications of these protein-protein interactions on Tg processing. This

combination of our novel TRIP approach coupled with functional screening deconvo-

luted previously established PQC mechanisms for Tg processing while also providing

new paradigms within PQC pathways that are critical for secretion of the prohormone.

TRIP has allowed identification and resolution of unique temporal changes in Tg

interactions with glycan processing components including STT3B, CANX, CALR, and

UGGT1, while contrasting WT to mutant Tg variants. These changes subsequently cor-

relate with altered interactions with ERAD components EDEM3, OS9, SEL1L, and VCP.

Moreover, we identified a broader scope of Tg interactors in thyroid tissue, including

ER-phagy receptors ATL3, CCPG1, RTN3, and the RTN3 adaptor protein PGRMC1.

Identification of these receptors establishes a direct link from Tg processing to ER-

phagy or ERLAD degradation mechanisms. Additionally, glycan dependent and inde-

pendent mechanisms have been established for the degradation of ERAD resistant ER

clients (Fregno et al., 2021; He et al., 2021). This overlap in degradation pathways may

be similarly at play in the case of Tg biogenesis and processing, especially for mutants

forming difficult to degrade aggregates (di Jeso et al., 2005, 2014; Kim et al., 1993;

Menon et al., 2007).

An important finding was that TRIP was capable of resolving subtle temporal in-

teraction changes, for example with CANX, ERP29, and VCP, which were otherwise

masked in steady-state interactomics data. Nonetheless, there are some limitations

within our TRIP methodology. TRIP relies on a two-stage purification strategy which in-

creases sample handling and limits the amount of protein that can be subsequently en-

riched. Both add inherent variability to the workflow. Furthermore, pulsed labeling with

unnatural amino acids has been shown to slow portions of protein translation (Bagert et

al., 2014; Dieterich et al., 2006; Kiick et al., 2001; Lang and Chin, 2014). Additionally,

within our dataset we noticed that the temporal profiles for ribosomal and proteasomal

subunits, trafficking and lysosomal components are inherently difficult to measure

across experiments. To efficiently measure these components temporally the TRIP

methodology will require further optimization.

The functional implications of protein-protein interactions can be difficult to de-

duce, especially in the case of PQC mechanisms containing several layers of redun-

dancy across stress response pathways, paralogs, and multiple unique proteins sharing

similar functions (Bludau and Aebersold, 2020; Braakman and Hebert, 2013; Karagöz et

al., 2019; Wright and Plate, 2021). This led us to establish a siRNA screening platform

to broadly investigate the functional implications of PQC components to complement

our TRIP data. With this assay we found the trafficking factor NAPA (α-SNAP) was

heavily implicated in WT Tg secretion. Most strikingly, we found that VCP and TEX264

were implicated in C1264R processing, and siRNA silencing of either led to rescue of

C1264R secretion. Rescue of mutant Tg trafficking from ER to Golgi, but not secretion,

with low-temperature correction had been documented previously (Kim et al., 1996).Yet,

this is the first study, to our knowledge, that identified a restorative approach to mutant

Tg secretion. To expound upon this, our findings that pharmacological VCP inhibition

selectively rescues C1264R secretion compared to WT, and mutant Tg was selectively

enriched with TEX264 further corroborated the TRIP data establishing that Tg mutants

undergo differential interactions with degradation pathways compared to WT Tg. Similar

phenomena have been observed for the partitioning and differential interactions of other

prohormone proteins, such as proinsulin and proopiomelanocortin, and proteasome re-

sistant polymers of alpha1-antitrypsin Z-variant. RTN3 is implicated in these differential

interactions and is shown to be selective for these prohormones compared to other ER-

phagy receptors (Chen et al., 2021; Cunningham et al., 2019; Fregno et al., 2018).

While A2234D and C1264R Tg were preferentially enriched with TEX264 compared to

WT, it remains unclear what other accessory proteins may be necessary for the recogni-

tion of TEX264 clients (An et al., 2019; Chino et al., 2019). Furthermore, TEX264 func-

tion in both protein degradation and DNA damage repair further complicates RNAi

based investigations (Fielden et al., 2022). Finally, as we discovered that pharmacolog-

ical VCP inhibition can rescue C1264R secretion yet is detrimental for WT processing, it

is unclear whether VCP may exhibit distinct functions for WT and mutant Tg PQC.

We used our TRIP method to monitor the changes in interactions associated with

the rescue of C1264R with pharmacological VCP inhibition. We found that this rescue is

correlated with broad temporal changes in interactions across both glycan processing

and Hsp70/90 chaperone pathways, while exhibiting more discrete changes with select

disulfide/redox processing components and leaving interactions with prolyl isomerization

and hydroxylation components largely unchanged. Mapping these temporal changes in

response to pharmacological VCP inhibition and C1264R rescue highlights the capabili-

ties of TRIP to not only resolve protein-protein interactions across disease states but

also identify compensatory mechanisms that may take place with drug treatment or

other modulating techniques like gene inhibition or activation that can correct or modu-

late disease states. Consequently, TRIP may find broad applicability for delineating the

proteostasis deficiencies that give rise to diverse protein misfolding diseases and eluci-

dating other cellular interactome dynamics.

STAR*METHODS

Detailed methods are provided in the online version of this paper and included the fol-

lowing

- KEY RESOURCES TABLE

- LEAD CONTACT AND METERIALS AVAILABILITY

- METHOD DETAILS

  o Plasmid Production and Antibodies

  o Cell Line Engineering

  o Time-Resolved Interactome Profiling

  o Liquid Chromatography – Tandem Mass Spectrometry

  o Immunoblotting and SDS-PAGE

  o qRT-PCR

  o siRNA Screening Assay

  o FRT siRNA Validation Studies

  o 35 S Pulse Chase Assay

  o VCP Pharmacological Inhibition Studies

  o TEX264 & Tg-NLuc Co-Immunoprecipitation Studies

- QUANTIFICATION AND STATISTICAL ANALYSIS

  o Mass Spectrometry Interactomics and TMT Quantification

- DATA AND CODE AVAILABILITY

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at #####

ACKNOWLEDGEMENTS

M.T.W. was supported by the National Science Foundation – Graduate Research Fel-

lowship Program. This work was also funded by an NIGMS R35 award

(1R35GM133552). This content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

AUTHOR CONTRIBUTIONS

M.T.W. and L.P. designed experiments. All authors performed experiments. M.T.W. and

L.P. wrote the manuscript. All authors contributed feedback for the manuscript.

DECLARATION OF INTEREST

The authors declare no competing interests.

REFERENCES

An, H., Ordureau, A., Paulo, J.A., Shoemaker, C.J., Denic, V., and Harper, J.W. (2019). TEX264 Is an Endoplasmic Reticulum-Resident ATG8-Interacting Protein Critical for ER Remodeling during Nutrient Stress. Mol Cell *74*, 891-908.e10. https://doi.org/10.1016/j.molcel.2019.03.034 PMID - 31006537.

Bagert, J.D., Xie, Y.J., Sweredoski, M.J., Qi, Y., Hess, S., Schuman, E.M., and Tirrell, D.A. (2014). Quantitative, Time-Resolved Proteomic Analysis by Combining Bioorthogonal Noncanonical Amino Acid Tagging and Pulsed Stable Isotope Labeling by Amino Acids in Cell Culture. Molecular & Cellular Proteomics *13*, 1352–1358. https://doi.org/10.1074/mcp.m113.031914 PMID - 24563536.

Baryshev, M., Sargsyan, E., and Mkrtchian, S. (2006). ERp29 is an essential endoplasmic reticulum factor regulating secretion of thyroglobulin. Biochem Biophys Res Commun *340*, 617–624. https://doi.org/10.1016/j.bbrc.2005.12.052.

Baryshev, M., Sargsyan, E., Wallin, G., Lejnieks, A., Furudate, S., Hishinuma, A., and Mkrtchian, S. Unfolded protein response is involved in the pathology of human congenital hypothyroid goiter and rat non-goitrous congenital hypothyroidism. J Mol Endocrinol *32*, 903 920. https://doi.org/10.1677/jme.0.0320903 PMID - 15171721.

Beatty, K.E., Liu, J.C., Xie, F., Dieterich, D.C., Schuman, E.M., Wang, Q., and Tirrell, D.A. (2006). Fluorescence Visualization of Newly Synthesized Proteins in Mammalian Cells. Angewandte Chemie International Edition *45*, 7364–7367. https://doi.org/10.1002/ANIE.200602114.

Behnke, J., Mann, M.J., Scruggs, F.-L., Feige, M.J., and Hendershot, L.M. (2016). Members of the Hsp70 Family Recognize Distinct Types of Sequences to Execute ER Quality Control. Mol Cell *63*, 739 752. https://doi.org/10.1016/j.molcel.2016.07.012 PMID - 27546788.

van Bergen, W., Heck, A.J.R., and Baggelaar, M.P. (2021). Recent advancements in mass spectrometry–based tools to investigate newly synthesized proteins. Curr Opin Chem Biol *66*. https://doi.org/10.1016/J.CBPA.2021.07.001.

Bludau, I., and Aebersold, R. (2020). Proteomic and interactomic insights into the molecular basis of cell functional diversity. Nat Rev Mol Cell Biol *21*, 327–340. https://doi.org/10.1038/s41580-020-0231-2.

Braakman, I., and Hebert, D.N. (2013). Protein Folding in the Endoplasmic Reticulum. Cold Spring Harb Perspect Biol *5*, a013201–a013201. https://doi.org/10.1101/cshperspect.a013201.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G. v., Clark, N.R., Ma'ayan, A., and Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics *14*, 128. https://doi.org/10.1186/1471-2105-14-128.

Chen, Q., Xiao, Y., Chai, P., Zheng, P., Teng, J., and Chen, J. (2019). ATL3 Is a Tubular ER-Phagy Receptor for GABARAP-Mediated Selective Autophagy. Curr Biol *29*, 846-855.e6. https://doi.org/10.1016/j.cub.2019.01.041.

Chen, Y.J., Knupp, J., Arunagiri, A., Haataja, L., Arvan, P., and Tsai, B. (2021).
PGRMC1 acts as a size-selective cargo receptor to drive ER-phagic clearance of mu-
tant prohormones. Nat Commun *12*. https://doi.org/10.1038/S41467-021-26225-8.

Chino, H., Hatta, T., Natsume, T., and Mizushima, N. (2019). Intrinsically Disordered
Protein TEX264 Mediates ER-phagy. Mol Cell *74*, 909-921.e6.
https://doi.org/10.1016/j.molcel.2019.03.033 PMID - 31006538.

Chiritoiu, M., Chiritoiu, G.N., Munteanu, C.V.A., Pastrama, F., Erwin Ivessa, N., and
Petrescu, S.M. (2020). EDEM1 Drives Misfolded Protein Degradation via ERAD and Ex-
ploits ER-Phagy as Back-Up Mechanism When ERAD Is Impaired. Int J Mol Sci *21*.
https://doi.org/10.3390/IJMS21103468.

Chitwood, P.J., and Hegde, R.S. (2020). An intramembrane chaperone complex facili-
tates membrane protein biogenesis. Nature *584*, 630–634.
https://doi.org/10.1038/s41586-020-2624-y.

Chou, T.F., Li, K., Frankowski, K.J., Schoenen, F.J., and Deshaies, R.J. (2013). Struc-
ture–Activity Relationship Study Reveals ML240 and ML241 as Potent and Selective In-
hibitors of p97 ATPase. ChemMedChem *8*, 297–312.
https://doi.org/10.1002/CMDC.201200520.

Chou, T.F., Bulfer, S.L., Weihl, C.C., Li, K., Lis, L.G., Walters, M.A., Schoenen, F.J., Lin,
H.J., Deshaies, R.J., and Arkin, M.R. (2014). Specific Inhibition of p97/VCP ATPase and
Kinetic Analysis Demonstrate Interaction between D1 and D2 ATPase Domains. J Mol
Biol *426*, 2886–2899. https://doi.org/10.1016/J.JMB.2014.05.022.

Christianson, J.C., Shaler, T.A., Tyler, R.E., and Kopito, R.R. (2008). OS-9 and GRP94
deliver mutant α1-antitrypsin to the Hrd1–SEL1L ubiquitin ligase complex for ERAD. Nat
Cell Biol *10*, 272–282. https://doi.org/10.1038/ncb1689 PMID - 18264092.

Christianson, J.C., Olzmann, J.A., Shaler, T.A., Sowa, M.E., Bennett, E.J., Richter,
C.M., Tyler, R.E., Greenblatt, E.J., Harper, J.W., and Kopito, R.R. (2012). Defining hu-
man ERAD networks through an integrative mapping strategy. Nat Cell Biol *14*, 93–105.
https://doi.org/10.1038/ncb2383 PMID - 22119785.

Chung, N., Zhang, X.D., Kreamer, A., Locco, L., Kuan, P.-F., Bartz, S., Linsley, P.S.,
Ferrer, M., and Strulovici, B. (2008). Median Absolute Deviation to Improve Hit Selection
for Genome-Scale RNAi Screens. J Biomol Screen *13*, 149–158.
https://doi.org/10.1177/1087057107312035 PMID - 18216396.

Citterio, C.E., Targovnik, H.M., and Arvan, P. (2019). The role of thyroglobulin in thyroid
hormonogenesis. Nat Rev Endocrinol *15*, 323–338. https://doi.org/10.1038/s41574-019-
0184-8 PMID - 30886364.

Cortez, D. (2017). Proteomic Analyses of the Eukaryotic Replication Machinery. In
Methods in Enzymology, pp. 33–53.

Coscia, F., Taler-Verčič, A., Chang, V.T., Sinn, L., O'Reilly, F.J., Izoré, T., Renko, M.,
Berger, I., Rappsilber, J., Turk, D., et al. (2020). The structure of human thyroglobulin.
Nature *578*, 1–4. https://doi.org/10.1038/s41586-020-1995-4 PMID - 32025030.

Cui, Y., Parashar, S., Zahoor, M., Needham, P.G., Mari, M., Zhu, M., Chen, S., Ho, H.-
C., Reggiori, F., Farhan, H., et al. (2019). A COPII subunit acts with an autophagy re-
ceptor to target endoplasmic reticulum for degradation. Science (1979) *365*, 53–60.
https://doi.org/10.1126/science.aau9263.

Cunningham, C.N., Williams, J.M., Knupp, J., Arunagiri, A., Arvan, P., and Tsai, B.
(2019). Cells Deploy a Two-Pronged Strategy to Rectify Misfolded Proinsulin Aggre-
gates. Mol Cell *75*, 442-456.e4. https://doi.org/10.1016/J.MOLCEL.2019.05.011.

Dieterich, D.C., Link, A.J., Graumann, J., Tirrell, D.A., and Schuman, E.M. (2006). Se-
lective identification of newly synthesized proteins in mammalian cells using bioorthogo-
nal noncanonical amino acid tagging (BONCAT). Proceedings of the National Academy
of Sciences *103*, 9482–9487. https://doi.org/10.1073/pnas.0601637103.

England, C.G., Ehlerding, E.B., and Cai, W. (2016). NanoLuc: A Small Luciferase Is
Brightening Up the Field of Bioluminescence. Bioconjug Chem *27*, 1175–1187.
https://doi.org/10.1021/acs.bioconjchem.6b00112 PMID - 27045664.

Fielden, J., Popović, M., and Ramadan, K. (2022). TEX264 at the intersection of au-
tophagy and DNA repair. Autophagy *18*, 40.
https://doi.org/10.1080/15548627.2021.1894059.

Fonslow, B.R., Niessen, S.M., Singh, M., Wong, C.C.L., Xu, T., Carvalho, P.C., Choi, J.,
Park, S.K., and Yates, J.R. (2012). Single-step inline hydroxyapatite enrichment facili-
tates identification and quantitation of phosphopeptides from mass-limited proteomes
with MudPIT. J Proteome Res *11*, 2697–2709.
https://doi.org/10.1021/PR300200X/SUPPL_FILE/PR300200X_SI_002.XLS.

Forrester, A., de Leonibus, C., Grumati, P., Fasana, E., Piemontese, M., Staiano, L.,
Fregno, I., Raimondi, A., Marazza, A., Bruno, G., et al. (2019). A selective ER-phagy ex-
erts procollagen quality control via a Calnexin-FAM134B complex. EMBO J *38*.
https://doi.org/10.15252/embj.201899847.

Fregno, I., Fasana, E., Bergmann, T.J., Raimondi, A., Loi, M., Soldà, T., Galli, C., D'An-
tuono, R., Morone, D., Danieli, A., et al. (2018). ER-to-lysosome-associated degradation
of proteasome-resistant ATZ polymers occurs via receptor-mediated vesicular transport.
EMBO J *37*. https://doi.org/10.15252/EMBJ.201899259.

Fregno, I., Fasana, E., Soldà, T., Galli, C., and Molinari, M. (2021). N-glycan processing
selects ERAD-resistant misfolded proteins for ER-to-lysosome-associated degradation.
EMBO J e107240. https://doi.org/10.15252/embj.2020107240 PMID - 34152647.

Grumati, P., Morozzi, G., Hölper, S., Mari, M., Harwardt, M.-L.I.E., Yan, R., Müller, S.,
Reggiori, F., Heilemann, M., and Dikic, I. (2017). Full length RTN3 regulates turnover of
tubular endoplasmic reticulum via selective autophagy. Elife *6*, e25555.
https://doi.org/10.7554/elife.25555 PMID - 28617241.

He, L., Kennedy, A.S., Houck, S., Aleksandrov, A., Quinney, N.L., Cyr-Scully, A., Cho-
lon, D.M., Gentzsch, M., Randell, S.H., Ren, H.Y., et al. (2021). DNAJB12 and Hsp70

triage arrested intermediates of N1303K-CFTR for endoplasmic reticulum-associated
autophagy. Mol Biol Cell *32*, 538–553. https://doi.org/10.1091/mbc.E20-11-0688.

Hill, S.M., Wrobel, L., Ashkenazi, A., Fernandez-Estevez, M., Tan, K., Bürli, R.W., and
Rubinsztein, D.C. (2021). VCP/p97 regulates Beclin-1-dependent autophagy initiation.
Nat Chem Biol *17*, 448–455. https://doi.org/10.1038/S41589-020-00726-X.

Hishinuma, A., Takamatsu, J., Ohyama, Y., Yokozawa, T., Kanno, Y., Kuma, K., Yo-
shida, S., Matsuura, N., and Ieiri, T. (1999). Two Novel Cysteine Substitutions (C1263R
and C1995S) of Thyroglobulin Cause a Defect in Intracellular Transport of Thyroglobulin
in Patients with Congenital Goiter and the Variant Type of Adenomatous Goiter1. J Clin
Endocrinol Metab *84*, 1438–1444. https://doi.org/10.1210/jcem.84.4.5633.

Howden, A.J.M., Geoghegan, V., Katsch, K., Efstathiou, G., Bhushan, B., Boutureira,
O., Thomas, B., Trudgian, D.C., Kessler, B.M., Dieterich, D.C., et al. (2013). QuaNCAT:
quantitating proteome dynamics in primary cells. Nat Methods *10*, 343–346.
https://doi.org/10.1038/nmeth.2401 PMID - 23474466.

di Jeso, B., Park, Y., Ulianich, L., Treglia, A.S., Urbanas, M.L., High, S., and Arvan, P.
(2005). Mixed-Disulfide Folding Intermediates between Thyroglobulin and Endoplasmic
Reticulum Resident Oxidoreductases ERp57 and Protein Disulfide Isomerase. Mol Cell
Biol *25*, 9793–9805. https://doi.org/10.1128/MCB.25.22.9793-9805.2005.

di Jeso, B., Morishita, Y., Treglia, A.S., Lofrumento, D.D., Nicolardi, G., Beguinot, F.,
Kellogg, A.P., and Arvan, P. (2014). Transient Covalent Interactions of Newly Synthe-
sized Thyroglobulin with Oxidoreductases of the Endoplasmic Reticulum*. Journal of Bi-
ological Chemistry *289*, 11488–11496. https://doi.org/10.1074/JBC.M113.520767.

Kanou, Y., Hishinuma, A., Tsunekawa, K., Seki, K., Mizuno, Y., Fujisawa, H., Imai, T.,
Miura, Y., Nagasaka, T., Yamada, C., et al. (2007). Thyroglobulin Gene Mutations Pro-
ducing Defective Intracellular Transport of Thyroglobulin Are Associated with Increased
Thyroidal Type 2 Iodothyronine Deiodinase Activity. J Clin Endocrinol Metab *92*, 1451–
1457. https://doi.org/10.1210/JC.2006-1242.

Karagöz, G.E., Acosta-Alvear, D., and Walter, P. (2019). The Unfolded Protein Re-
sponse: Detecting and Responding to Fluctuations in the Protein-Folding Capacity of
the Endoplasmic Reticulum. Cold Spring Harb Perspect Biol a033886.
https://doi.org/10.1101/cshperspect.a033886 PMID - 30670466.

Kiick, K.L., Weberskirch, R., and Tirrell, D.A. (2001). Identification of an expanded set of
translationally active methionine analogues in Escherichia coli. FEBS Lett *502*, 25–30.
https://doi.org/10.1016/S0014-5793(01)02657-6.

Kim, P.S., and Arvan, P. (1995). Calnexin and BiP act as sequential molecular chaper-
ones during thyroglobulin folding in the endoplasmic reticulum. J Cell Biol *128*, 29–38.
https://doi.org/10.1083/JCB.128.1.29.

Kim, B., Arcos, S., Rothamel, K., Jian, J., Rose, K.L., McDonald, W.H., Bian, Y., Rea-
soner, S., Barrows, N.J., Bradrick, S., et al. (2020). Discovery of Widespread Host

Protein Interactions with the Pre-replicated Genome of CHIKV Using VIR-CLASP. Mol
Cell *78*, 624-640.e7. https://doi.org/10.1016/j.molcel.2020.04.013.

Kim, P.S., Kim, K.R., and Arvan, P. (1993). Disulfide-linked aggregation of thyroglobulin
normally occurs during nascent protein folding. American Journal of Physiology-Cell
Physiology *265*, C704–C711. https://doi.org/10.1152/ajpcell.1993.265.3.c704 PMID -
8214027.

Kim, P.S., Kwon, O.Y., and Arvan, P. (1996). An endoplasmic reticulum storage disease
causing congenital goiter with hypothyroidism. J Cell Biol *133*, 517–527.
https://doi.org/10.1083/jcb.133.3.517.

Kuo, C.C., Chiang, A.W.T., Baghdassarian, H.M., and Lewis, N.E. (2021). Dysregulation
of the secretory pathway connects Alzheimer's disease genetics to aggregate formation.
Cell Syst *12*, 873-884.e4. https://doi.org/10.1016/J.CELS.2021.06.001.

Lamriben, L., Graham, J.B., Adams, B.M., and Hebert, D.N. (2016). N-Glycan-based ER
Molecular Chaperone and Protein Quality Control System: The Calnexin Binding Cycle.
Traffic *17*, 308–326. https://doi.org/10.1111/tra.12358 PMID - 26676362.

Lang, K., and Chin, J.W. (2014). Cellular Incorporation of Unnatural Amino Acids and
Bioorthogonal Labeling of Proteins. Chem Rev *114*, 4764–4806.
https://doi.org/10.1021/cr400355w.

Li, J., Vranken, J.G. van, Vaites, L.P., Schweppe, D.K., Huttlin, E.L., Etienne, C.,
Nandhikonda, P., Viner, R., Robitaille, A.M., Thompson, A.H., et al. (2020). TMTpro rea-
gents: a set of isobaric labeling mass tags enables simultaneous proteome-wide meas-
urements across 16 samples. Nat Methods *17*, 399–404.
https://doi.org/10.1038/s41592-020-0781-4 PMID - 32203386.

Liang, J.R., Lingeman, E., Ahmed, S., and Corn, J.E. (2018). Atlastins remodel the en-
doplasmic reticulum for selective autophagy. Journal of Cell Biology *217*, 3354–3367.
https://doi.org/10.1083/JCB.201804185.

Liang, J.R., Lingeman, E., Luong, T., Ahmed, S., Muhar, M., Nguyen, T., Olzmann, J.A.,
and Corn, J.E. (2020). A Genome-wide ER-phagy Screen Highlights Key Roles of Mito-
chondrial Metabolism and ER-Resident UFMylation. Cell *180*, 1160-1177.e20.
https://doi.org/10.1016/j.cell.2020.02.017.

Lobingier, B.T., Hüttenhain, R., Eichel, K., Miller, K.B., Ting, A.Y., von Zastrow, M., and
Krogan, N.J. (2017). An Approach to Spatiotemporally Resolve Protein Interaction Net-
works in Living Cells. Cell *169*, 350-360.e12. https://doi.org/10.1016/j.cell.2017.03.022.

Ma, Y., McClatchy, D.B., Barkallah, S., Wood, W.W., and Yates, J.R. (2017). HILAQ: A
Novel Strategy for Newly Synthesized Protein Quantification. J Proteome Res *16*,
2213–2220. https://doi.org/10.1021/acs.jproteome.7b00005 PMID - 28437088.

Ma, Y., McClatchy, D.B., Barkallah, S., Wood, W.W., and Yates, J.R. (2018). Quantita-
tive analysis of newly synthesized proteins. Nat Protoc *13*, 1744–1762.
https://doi.org/10.1038/S41596-018-0012-Y.

Marinko, J.T., Wright, M.T., Schlebach, J.P., Clowes, K.R., Heintzman, D.R., Plate, L.,
and Sanders, C.R. (2021). Glycosylation limits forward trafficking of the tetraspan mem-
brane protein PMP22. Journal of Biological Chemistry *296*.
https://doi.org/10.1016/J.JBC.2021.100719.

McClatchy, D.B., Ma, Y., Liu, C., Stein, B.D., Martínez-Bartolomé, S., Vasquez, D., Hell-
berg, K., Shaw, R.J., and Yates, J.R. (2015). Pulsed Azidohomoalanine Labeling in
Mammals (PALM) Detects Changes in Liver-Specific LKB1 Knockout Mice. J Proteome
Res *14*, 4815–4822. https://doi.org/10.1021/acs.jproteome.5b00653 PMID - 26445171.

McDonald, E.F., Sabusap, C.M.P., Kim, M., and Plate, L. (2022). Distinct proteostasis
states drive pharmacologic chaperone susceptibility for Cystic Fibrosis Transmembrane
Conductance Regulator misfolding mutants. Mol Biol Cell
https://doi.org/10.1091/mbc.E21-11-0578.

McGilvray, P.T., Anghel, S.A., Sundaram, A., Zhong, F., Trnka, M.J., Fuller, J.R., Hu,
H., Burlingame, A.L., and Keenan, R.J. (2020a). An ER translocon for multi-pass mem-
brane protein biogenesis. Elife *9*, 1–43. https://doi.org/10.7554/ELIFE.56889.

McGilvray, P.T., Anghel, S.A., Sundaram, A., Zhong, F., Trnka, M.J., Fuller, J.R., Hu,
H., Burlingame, A.L., and Keenan, R.J. (2020b). An ER translocon for multi-pass mem-
brane protein biogenesis. Elife *9*, 1–43. https://doi.org/10.7554/ELIFE.56889.

Menon, S., Lee, J., Abplanalp, W.A., Yoo, S.-E., Agui, T., Furudate, S., Kim, P.S., and
Arvan, P. (2007). Oxidoreductase Interactions Include a Role for ERp72 Engagement
with Mutant Thyroglobulin from the rdw/rdw Rat Dwarf. Journal of Biological Chemistry
282, 6183–6191. https://doi.org/10.1074/jbc.M608863200.

Munden, A., Wright, M.T., Han, D., Tirgar, R., Plate, L., and Nordman, J.T. (2022). Iden-
tification of replication fork-associated proteins in Drosophila embryos and cultured cells
using iPOND coupled to quantitative mass spectrometry. Sci Rep *12*, 1–11.
https://doi.org/10.1038/s41598-022-10821-9.

Needham, P.G., Guerriero, C.J., and Brodsky, J.L. (2019). Chaperoning Endoplasmic
Reticulum-Associated Degradation (ERAD) and Protein Conformational Diseases. Cold
Spring Harb Perspect Biol a033928. https://doi.org/10.1101/cshperspect.a033928
PMID - 30670468.

Okuda-Shimizu, Y., and Hendershot, L.M. (2007). Characterization of an ERAD path-
way for nonglycosylated BiP substrates, which require Herp. Mol Cell *28*, 544–554.
https://doi.org/10.1016/J.MOLCEL.2007.09.012.

Pardo, V., Vono-Toniolo, J., Rubio, I.G.S., Knobel, M., Possato, R.F., Targovnik, H.M.,
Kopp, P., and Medeiros-Neto, G. (2009). The p.A2215D Thyroglobulin Gene Mutation
Leads to Deficient Synthesis and Secretion of the Mutated Protein and Congenital Hy-
pothyroidism with Wide Phenotype Variation. J Clin Endocrinol Metab *94*, 2938–2944.
https://doi.org/10.1210/jc.2009-0150.

Perez Verdaguer, M., Zhang, T., Surve, S., Paulo, J.A., Wallace, C., Watkins, S.C.,
Gygi, S.P., and Sorkin, A. (2022). Time-resolved proximity labeling of protein networks

associated with ligand-activated EGFR. Cell Rep *39*, 110950.
https://doi.org/10.1016/J.CELREP.2022.110950/ATTACHMENT/0568B930-8965-47D5-
BC4E-950606E2E486/MMC8.XLSX.

Petelski, A.A., Emmott, E., Leduc, A., Huffman, R.G., Specht, H., Perlman, D.H., and
Slavov, N. (2021). Multiplexed single-cell proteomics using SCoPE2. Nat Protoc *16*,
5398–5425. https://doi.org/10.1038/s41596-021-00616-z.

Phillips, S.L., Soderblom, E.J., Bradrick, S.S., and Garcia-Blanco, M.A. (2016). Identifi-
cation of Proteins Bound to Dengue Viral RNA In Vivo Reveals New Host Proteins Im-
portant for Virus Replication. MBio *7*. https://doi.org/10.1128/mBio.01865-15.

Piette, B.L., Alerasool, N., Lin, Z.-Y., Lacoste, J., Lam, M.H.Y., Qian, W.W., Tran, S.,
Larsen, B., Campos, E., Peng, J., et al. (2021). Comprehensive interactome profiling of
the human Hsp70 network highlights functional differentiation of J domains. Mol Cell
https://doi.org/10.1016/j.molcel.2021.04.012 PMID - 33957083.

Pohl, C., and Dikic, I. (2019). Cellular quality control by the ubiquitin-proteasome system
and autophagy. Science (1979) *366*, 818–822. https://doi.org/10.1126/science.aax3769
PMID - 31727826.

Pshezhetsky, A. v., and Hinek, A. (2011). Where catabolism meets signalling: neuram-
inidase 1 as a modulator of cell receptors. Glycoconjugate Journal 2011 28:7 *28*, 441–
452. https://doi.org/10.1007/S10719-011-9350-5.

Rizzolo, K., and Houry, W.A. (2019). Multiple functionalities of molecular chaperones re-
vealed through systematic mapping of their interaction networks. Journal of Biological
Chemistry *294*, 2142–2150. https://doi.org/10.1074/JBC.TM118.002805.

Rizzolo, K., Huen, J., Kumar, A., Phanse, S., Vlasblom, J., Kakihara, Y., Zeineddine,
H.A., Minic, Z., Snider, J., Wang, W., et al. (2017). Features of the Chaperone Cellular
Network Revealed through Systematic Interaction Mapping. Cell Rep *20*, 2735–2748.
https://doi.org/10.1016/j.celrep.2017.08.074 PMID - 28903051.

Ryuzono, S., Takase, R., Oishi, K., Ikeda, A., Chigwechokha, P.K., Funahashi, A., Ko-
matsu, M., Miyagi, T., and Shiozaki, K. (2016). Lysosomal localization of Japanese
medaka (Oryzias latipes) Neu1 sialidase and its highly conserved enzymatic profiles
with human. Gene *575*, 513–523. https://doi.org/10.1016/J.GENE.2015.09.028.

Sabusap, C.M., Wang, W., McNicholas, C.M., Chung, W.J., Fu, L., Wen, H., Mazur, M.,
Kirk, K.L., Collawn, J.F., Hong, J.S., et al. (2016). Analysis of cystic fibrosis–associated
P67L CFTR illustrates barriers to personalized therapeutics for orphan diseases. JCI In-
sight *1*. https://doi.org/10.1172/jci.insight.86581.

Sirbu, B.M., Couch, F.B., Feigerle, J.T., Bhaskara, S., Hiebert, S.W., and Cortez, D.
(2011). Analysis of protein dynamics at active, stalled, and collapsed replication forks.
Genes Dev *25*, 1320–1327. https://doi.org/10.1101/gad.2053211.

Smith, M.D., Harley, M.E., Kemp, A.J., Wills, J., Lee, M., Arends, M., Kriegsheim, A.
von, Behrends, C., and Wilkinson, S. (2018). CCPG1 Is a Non-canonical Autophagy

Cargo Receptor Essential for ER-Phagy and Pancreatic ER Proteostasis. Dev Cell *44*,
217-232.e11. https://doi.org/10.1016/j.devcel.2017.11.024 PMID - 29290589.

Song, H., Orr, A., Duan, M., Merz, A.J., and Wickner, W. (2017). Sec17/Sec18 act
twice, enhancing membrane fusion and then disassembling cis-SNARE complexes.
Elife *6*. https://doi.org/10.7554/ELIFE.26646.

Taipale, M., Krykbaeva, I., Koeva, M., Kayatekin, C., Westover, K.D., Karras, G.I., and
Lindquist, S. (2012). Quantitative Analysis of Hsp90-Client Interactions Reveals Princi-
ples of Substrate Recognition. Cell *150*, 987–1001.
https://doi.org/10.1016/j.cell.2012.06.047 PMID - 22939624.

Taipale, M., Tucker, G., Peng, J., Krykbaeva, I., Lin, Z.-Y., Larsen, B., Choi, H., Berger,
B., Gingras, A.-C., and Lindquist, S. (2014). A Quantitative Chaperone Interaction Net-
work Reveals the Architecture of Cellular Protein Homeostasis Pathways. Cell *158*,
434–448. https://doi.org/10.1016/j.cell.2014.05.039.

Taldone, T., Wang, T., Rodina, A., Pillarsetty, N.V.K., Digwal, C.S., Sharma, S., Yan, P.,
Joshi, S., Pagare, P.P., Bolaender, A., et al. (2020). A Chemical Biology Approach to
the Chaperome in Cancer—HSP90 and Beyond. Cold Spring Harb Perspect Biol *12*,
a034116. https://doi.org/10.1101/cshperspect.a034116.

Thompson, A., Wölmer, N., Koncarevic, S., Selzer, S., Böhm, G., Legner, H., Schmid,
P., Kienle, S., Penning, P., Höhle, C., et al. (2019). TMTpro: Design, Synthesis, and Ini-
tial Evaluation of a Proline-Based Isobaric 16-Plex Tandem Mass Tag Reagent Set.
Anal Chem *91*, 15941–15950. https://doi.org/10.1021/acs.analchem.9b04474 PMID -
31738517.

Tokunaga, F., Brostrom, C., Koide, T., and Arvan, P. (2000). Endoplasmic reticulum
(ER)-associated degradation of misfolded N-linked glycoproteins is suppressed upon in-
hibition of ER mannosidase I. Journal of Biological Chemistry *275*, 40757–40764.
https://doi.org/10.1074/jbc.m001073200 PMID - 10984471.

Tsai, C.-F., Zhao, R., Williams, S.M., Moore, R.J., Schultz, K., Chrisler, W.B., Pasa-
Tolic, L., Rodland, K.D., Smith, R.D., Shi, T., et al. (2020). An Improved Boosting to Am-
plify Signal with Isobaric Labeling (iBASIL) Strategy for Precise Quantitative Single-cell
Proteomics. Molecular & Cellular Proteomics *19*, 828–838.
https://doi.org/10.1074/mcp.RA119.001857.

Vij, N., Fang, S., and Zeitlin, P.L. (2006). Selective inhibition of endoplasmic reticulum-
associated degradation rescues DeltaF508-cystic fibrosis transmembrane regulator and
suppresses interleukin-8 levels: therapeutic implications. J Biol Chem *281*, 17369–
17378. https://doi.org/10.1074/JBC.M600509200.

Vranka, J.A., Sakai, L.Y., and Bächinger, H.P. (2004). Prolyl 3-Hydroxylase 1, Enzyme
Characterization and Identification of a Novel Family of Enzymes. Journal of Biological
Chemistry *279*, 23615–23621. https://doi.org/10.1074/JBC.M312807200.

Wong, M.Y., and Shoulders, M.D. (2019). Targeting defective proteostasis in the colla-
genopathies. Curr Opin Chem Biol *50*, 80–88.
https://doi.org/10.1016/J.CBPA.2019.02.021.

Wright, M.T., and Plate, L. (2021). Revealing functional insights into ER proteostasis
through proteomics and interactomics. Exp Cell Res *399*.
https://doi.org/10.1016/J.YEXCR.2020.112417.

Wright, M.T., Kouba, L., and Plate, L. (2021). Thyroglobulin Interactome Profiling De-
fines Altered Proteostasis Topology Associated With Thyroid Dyshormonogenesis. Mo-
lecular & Cellular Proteomics *20*, 100008. https://doi.org/10.1074/mcp.ra120.002168
PMID - 33208410.

Zhang, X., Smits, A.H., van Tilburg, G.B.A., Ovaa, H., Huber, W., and Vermeulen, M.
(2018). Proteome-wide identification of ubiquitin interactions using UbIA-MS. Nature
Protocols 2018 13:3 *13*, 530–550. https://doi.org/10.1038/nprot.2017.147.

Zhao, C., Slevin, J.T., and Whiteheart, S.W. (2007). Cellular functions of NSF: Not just
SNAPs and SNAREs. FEBS Lett *581*, 2140–2149. https://doi.org/10.1016/J.FEB-
SLET.2007.03.032.

Zhou, H.J., Wang, J., Yao, B., Wong, S., Djakovic, S., Kumar, B., Rice, J., Valle, E., So-
riano, F., Menon, M.K., et al. (2015). Discovery of a First-in-Class, Potent, Selective,
and Orally Bioavailable Inhibitor of the p97 AAA ATPase (CB-5083). J Med Chem *58*,
9480–9497. https://doi.org/10.1021/ACS.JMEDCHEM.5B01346.

STAR*METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
KDEL	Enzo Life Sciences	ADI-SPA-827
M2-FLAG	Sigma Aldrich	F1804
PDIA4	Protein Tech	14712-1-AP
TG	Proteintech	21714-1-AP
GAPDH	GeneTex	GTX627408
Tubulin-Rhodamine	BioRad	12004165
Goat anti-mouse Star-bright700	Bio-Rad	12004158
Goat anti-rabbit IRDye800	LI-COR	926-32211
Chemicals, Peptides, & Recombinant Proteins		
NMS-873	Cayman Chemical	17674
CB-5083	Cayman Chemical	19311

ML-240	Cayman Chemical	17373
dithiobis(succinimidyl propionate) (DSP)	Thermo Scientific	22585
2-(4-((Bis((1-(*tert*-butyl)-1*H*-1,2,3-triazol-4-yl)methyl)amino)methyl)-1*H*-1,2,3-triazol-1-yl)acetic acid (BTTAA)	Click Chemistry Tools	1236
Carboxytetramethylrhodamine (TAMRA)-Azide-Polyethylene Glycol (PEG)-Desthiobiotin	BroadPharm	BP-22475
DharmaFECT1 Transfection Reagent	Dharmacon	T-2001
Lipofectamine 3000	Invitrogen	L3000
G1 Anti-DYKDDDDK affinity resin	GenScript	L00432
High-Capacity Streptavidin agarose resin	Pierce	20357
Commercial Assays		
Nano-Glo Luciferase Assay System	Promega	N1110
Deposited Data		
Proteomics RAW and analyzed data	This study	
Experimental Models: Cell Lines		
Human HEK293 Flp-In	Thermo Fisher	R75007
Fischer Rat Thyroid (FRT) Flp-In	(Sabusap et al., 2016)	N/A
Recombinant DNA		
Tg-FLAG - pcDNA3.1+/C-(K)-DYK	Genscript	OHu20241
pcDNA5/FRT	(Sabusap et al., 2016)	N/A
pOG44	(Sabusap et al., 2016)	N/A
pcDNA5/FRT-Tg-FLAG	This study	N/A
pcDNA5/FRT-Tg-NLuc	This study	N/A
pMRX-INU-TEX264-FLAG	Addgene	128258
pMRX-INU-SEC62-FLAG	Addgene	128263
Sequence-Based Reagents		
Site-Directed Mutagenesis Primers	This study	See Table S1
qPCR Primers	This study	See Table S1

siGenome Cherry Picked Library	Dharmacon	See Table S6
Software		
Bio-Rad Image Lab	Bio-Rad	https://www.bio-rad.com/en-us/product/image-lab-software
Prism 9	GraphPad	https://www.graphpad.com/scientific-software/prism/
Proteome Discoverer 2.4	Thermo Fisher	https://www.thermofisher.com
CFX Maestro	Bio-Rad	https://www.bio-rad.com/en-us/product/cfx-maestro-software-for-cfx-real-time-pcr-instruments
FlowJo	BD Biosciences	https://www.flowjo.com/solutions/flowjo/downloads

LEAD CONTACT AND MATERIALS AVAILABILITY

All unique/stable reagents generated, including plasmids and cell lines are available

from the Lead Contact, Lars Plate (lars.plate@vanderbilt.edu) with a complete Materials

Transfer Agreement.

METHOD DETAILS

Plasmid Production and Antibodies

Tg-FLAG in pcDNA3.1+/C-(K)-DYK plasmid was purchased from Genscript (Clone ID

OHu20241). The Tg-FLAG gene was then amplified and assembled with an empty

pcDNA5/FRT expression vector using a HiFi DNA assembly kit (New England BioLabs,

E2621). This plasmid then underwent site-directed mutagenesis to produce pcDNA5-

C1264R-Tg-FLAG, and pcDNA5-A2234D-Tg-FLAG plasmids (Table S1).

An oligonucleotide fragment encoding Nanoluciferase (NLuc) was ordered from

Genewiz. To generate pcDNA5/FRT-Tg-NLuc plasmids the Tg gene was amplified from

the pcDNA5/FRT-Tg-FLAG plasmid and assembled with the NLuc fragment using a HiFi

DNA assembly kit (New England BioLabs). To generate the respective mutant construct

plasmids for A2234D and C1264R Tg, site directed mutagenesis was performed using a

Q5 polymerase (New England BioLabs). All oligonucleotide sequences can be found in

Table S1.

Cell Line Engineering

FRT cells were cultured in Ham's F-12, Coon's Modification (F12) media (Sigma, cat.

No. F6636) supplemented with 5% fetal bovine serum (FBS), and 1% penicillin

(10,000U) / streptomycin (10,000 µg/mL). To generate FRT flp-Tg-FT cells, 5×10^5 cells

cultured for 1 day were cotransfected with 2.25 µg flp recombinase pOG44 plasmid and

0.25 µg of FT-Tg pcDNA5 plasmid using Lipofectamine 3000. Cells were then cultured

in media containing Hygromycin B (100 µg/mL) to select site-specific recombinants. Re-

sistant clonal lines were sorted into single cell colonies using flow cytometry and

screened for FT-Tg expression (Table S1 and Figures S1A-S1D).

To generate HEK293 flp-Tg-NLuc cells, 5×10^5 cells cultured for 1 day were co-

transfected with 2.25 µg flp recombinase pOG44 plasmid and 0.25 µg of NLuc-Tg

pcDNA5 plasmid using Lipofectamine 3000. Cells were then cultured in media contain-

ing Hygromycin B (100 µg/mL) to select site specific recombinants. Polyclonal lines

953 were screened for Tg-NLuc expression and furimazine substrate turnover (Figures S1E-

954 S1K).

956 **Time-Resolved Interactome Profiling**

957 Fully confluent 15cm tissue culture plates of FRT cells were used. Cells were washed

958 with phosphate buffered saline (PBS) and depleted of methionine by incubating with

959 methionine free Dulbecco's Modified Eagle Medium (DMEM) supplemented with 5% di-

960 alyzed fetal bovine serum (FBS), 1% L-glutamine (2mM final concentration), 1% L-cys-

961 teine (200 µM final concentration), and 1% penicillin (10,000U) / streptomycin (10,000

962 µg/mL) for 30 minutes. Cells were then pulse labeled with Hpg-enriched DMEM supple-

963 mented with with 1% Hpg (200µM final concentration), 5% dialyzed FBS, 1% L-gluta-

964 mine (200mM final concentration), 1% L-cysteine (200µM), and 1% penicillin (10,000U)

965 / streptomycin (10,000 µg/mL) for 1 hour. After pulse labeling cells were washed with

966 F12 media containing 10-fold excess methionine (2mM final concentration). Cells were

967 then cultured in normal F12 media supplemented with 5% FBS and chased for the

968 specified time-points. Cells were harvested by washing with PBS and then crosslinked

969 with 0.5mM DSP in PBS at 37°C for 10 minutes. Crosslinking was quenched by the ad-

970 dition of 100mM Tris pH 7.5 at 37°C for 5 minutes. Lysates were prepared by lysing in

971 Radioimmunoprecipitation assay (RIPA) buffer (50mM Tris pH 7.5, 150mM NaCl, 0.1%

972 SDS, 1% Triton X-100, 0.5% deoxycholate) with protease inhibitor cocktail (Roche,

973 4693159001). Protein concentration was normalized to 1mg/mL using a BCA assay

974 (Thermo Scientific, 23225) and cell lysates underwent click reactions with the addition of

0.8mM copper sulfate (diluted from a 20mM stock in water), 1.6mM BTTAA (diluted

from a 40mM stock in water), 5mM sodium ascorbate (diluted from a 100mM stock in

water), and 100uM TAMRA-Azide-PEG-Desthiobiotin ligand (diluted from a 5mM stock

in DMSO). Samples were allowed to react at 37° for 1 hour while shaking at 750rpm.

Cell lysates were then precleared on 4B sepharose beads (Sigma, 4B200) at 4°C for 1

hour while rocking. Precleared lysates were immunoprecipitated with G1 Anti-

DYKDDDDK affinity resin overnight at 4°C while rocking. Resin was washed four times

with RIPA buffer, and proteins were eluted twice in 100μL immunoprecipitation elution

buffer (2% SDS in PBS) by heating at 95°C for 5 minutes. Eluted samples from FLAG

immunoprecipitations were then diluted with PBS to reduce the final SDS concentration

to 0.2%. The solutions then underwent streptavidin enrichment with high-capacity strep-

tavidin agarose resin (Pierce, 20359) for 2 hours at room temperature while rotating.

The resin was then washed with 1mL each of 1% SDS, 4M Urea, 1M sodium chloride,

followed by a final wash with 1% SDS (all wash buffers dissolved in PBS). Resin was

frozen overnight at -80°C and samples were then eluted twice with 100μL biotin elution

buffer (50mM Biotin in 1% SDS in PBS) by heating at 37°C and shaking at 750rpm for 1

hour. Eluted streptavidin enrichment samples were precipitated in methanol/chloroform,

washed three times with methanol, and air dried. Protein pellets were then resuspended

in 3μL 1% Rapigest SF Surfactant (Waters, 186002122) followed by the addition of

10μL of 50mM HEPES pH 8.0, and 34.5μL of water. Samples were reduced with 5mM

tris(2-carboxyethyl)phosphine (TCEP) (Sigma, 75259) at room temperature for 30

minutes and alkylated with 10mM iodoacetimide (Sigma, I6125) in the dark at room

temperature for 30 minutes. Proteins were digested with 0.5 µg of trypsin/Lys-C prote-

ase mix (Pierce, A40007) by incubating for 16-18 hours at 37°C and shaking at 750rpm.

Peptides were reacted with TMTpro 16plex reagents (Thermo Fisher, 44520) in 40% v/v

acetonitrile and incubated for one hour at room temperature. Reactions were quenched

by the addition of ammonium bicarbonate (0.4% w/v final concentration) and incubated

for one hour at room temperature. TMT labeled samples were then pooled and acidified

with 5% formic acid (Fisher, A117, v/v). Samples were concentrated using a speedvac

and resuspended in buffer A (97% water, 2.9% acetonitrile, and 0.1% formic acid, v/v/v).

Cleaved Rapigest SF surfactant was removed by centrifugation for 30 minutes at

21,100 x g.

For TRIP analysis coupled with ML-240 treatment, C1264R Tg FRT cells were

processed as described above with ML-240 (10µM) supplemented in Hpg-pulse media

and throughout the chase period.

1010

Liquid Chromatography – Tandem Mass Spectrometry

MudPIT microcolumns were prepared as previously described (Fonslow et al., 2012).

Peptide samples were directly loaded onto the columns using a high-pressure chamber.

Samples were then desalted for 30 minutes with buffer A (97% water, 2.9% acetonitrile,

0.1% formic acid v/v/v). LC-MS/MS analysis was performed using an Exploris480

(Thermo Fisher) mass spectrometer equipped with an Ultimate3000 RSLCnano system

(Thermo Fisher). MudPIT experiments were performed with 10µL sequential injections

of 0, 10, 30, 60, and 100% buffer C (500mM ammonium acetate in buffer A), followed

by a final injection of 90% buffer C with 10% buffer B (99.9% acetonitrile, 0.1% formic

acid v/v) and each step followed by a 140 minute gradient from 4% to 80% B with a flow

rate of 500nL/minute on a 20cm fused silica microcapillary column (ID 100 μm) ending

with a laser-pulled tip filled with Aqua C18, 3μm, 125 Å resin (Phenomenex). Elec-

trospray ionization (ESI) was performed directly from the analytical column by applying

a voltage of 2.2kV with an inlet capillary temperature of 275°C. Data-dependent acquisi-

tion of mass spectra was carried out by performing a full scan from 400-1600m/z at a

resolution of 120,000. Top-speed data acquisition was used for acquiring MS/MS spec-

tra using a cycle time of 3 seconds, with a normalized collision energy of 32, 0.4m/z iso-

lation window, automatic maximum injection time and 100% normalized AGC target, at

a resolution of 45,000 and a defined first mass (m/z) starting at 110. Peptide identifica-

tion and TMT-based protein quantification was carried out using Proteome Discoverer

2.4. MS/MS spectra were extracted from Thermo Xcalibur .raw file format and searched

using SEQUEST against a Uniprot rat proteome database supplemented with the hu-

man thyroglobulin gene (accessed 03/2014 and containing 28860 entries). The data-

base was curated to remove redundant protein and splice-isoforms. Searches were car-

ried out using a decoy database of reversed peptide sequences and the following pa-

rameters: 20ppm peptide precursor tolerance, 0.02 Da fragment mass tolerance, mini-

mum peptide length of 6 amino acids, trypsin cleavage with a maximum of two missed

cleavages, dynamic methionine modification of +15.995 Da (oxidation), dynamic protein

N-terminus +42.011 Da (acetylation), -131.040 (methionine loss), -89.030 (methionine

loss + acetylation), static cysteine modification of +57.0215 Da (carbamidomethylation),

and static peptide N-terminal and lysine modifications of +304.2071 Da (TMTpro

16plex).

1043

Immunoblotting and SDS-PAGE

Cell lysates were prepared by lysing in RIPA buffer with protease inhibitor cocktail

(Roche) and protein concentrations were normalized using a BCA assay (Thermo Sci-

entific). Lysates were then denatured with 1X Laemmli buffer + 100mM DTT and heated

at 95°C for 5 minutes before being separated by SDS-PAGE. Samples were transferred

onto poly-vinylidene difluoride (PVDF) membranes (Millipore, IPFL00010) for immunob-

lotting and blocked using 5% non-fat dry milk dissolved in tris buffered saline with 0.1%

Tween-20 (Fisher, BP337-100) (TBS-T). Primary antibodies were incubated either at

room temperature for 2 hours, or overnight at 4°C. Membranes were then washed three

times with TBS-T and incubated with secondary antibody in 5% non-fat dry milk dis-

solved in TBS-T either at room temperature for 1 hour or overnight at 4°C. Membranes

were washed three times with TBS-T and then imaged using a ChemiDoc MP Imaging

System (BioRad). Primary antibodies were acquired from commercial sources and used

at the indicated dilutions in immunoblotting buffer (5% bovine serum albumin (BSA) in

Tris-buffered saline pH 7.5, 0.1% Tween-20, and 0.1% sodium azide): KDEL (1:1000),

M2 anti-FLAG (1:1000), PDIA4 (1:1000), thyroglobulin (1:1000). Tubulin-Rhodamine pri-

mary antibody was obtained from commercial sources and used at 1:10000 dilution in

5% milk in Tris-buffered saline pH 7.5, 0.1% Tween-20 (TBS-T). Secondary antibodies

were obtained from commercial sources and used at the indicated dilutions in 5% milk

in TBS-T: Goat anti-mouse Star-bright700 (1:10000), Goat anti-rabbit IRDye800

(1:10000).

1065

qRT-PCR

RNA was prepared from cell pellets using the Quick-RNA miniprep kit (Zymo Research).

cDNA was synthesized from 500ng total cellular RNA using random hexamer primer

(IDT), oligo-dT primer (IDT), and M-MLV reverse transcriptase (Promega). qPCR analy-

sis was performed using iTaq Universal SYBR Green Supermix (Bio-Rad) combined

with primers for genes of interest and reactions were run in 96-well plates on a Bio-Rad

CFX qPCR instrument. Data analysis was then carried out in CFX Maestro (Bio-Rad).

Primer sequences for genes of interest are found in Table S1.

1074

siRNA Screening Assay

Proteins previously identified as Tg interactors, and other key proteostasis network

components were selected and targeted using an siGENOME SMARTpool siRNA li-

brary (Dharmacon) (Wright et al., 2021). HEK293 cells stably expressing Tg-NLuc con-

structs were seeded into 96-well plates at 2.5×10^4 cells/well and transfected using Dhar-

maFECT 1 following the DharmaFECT1 protocol (Dharmacon) with a 25nM siRNA con-

centration. Approximately 36 hours after transfection, cells were washed with PBS and

replenished with fresh DMEM media. After 4 hours, Tg-NLuc abundance in the lysate

and media were measured using the nano-glo luciferase assay system according to the

manufactures protocol (Promega). Four controls were included for the experiments

including a non-targeting siRNA control, a siGLO fluorescent control to monitor transfec-

tion efficiency, a vehicle control containing transfection reagents but lacking any siR-

NAs, and a lethal TOX control. Data was median normalized across individual 96-well

plates (Chung et al., 2008). Data represents two independent experiments for WT-NLuc

and A2234D-NLuc, and three independent experiments for C1264R NLuc. Cutoff crite-

ria for hits were set to those genes that increased or decreased Tg-NLuc abundance in

lysate or media by 3σ.

1092

FRT siRNA Validation Studies

For siRNA silencing follow-up experiments, Tg FRT cells were seeded into 6-well dishes

at 6.0×10^5 cells/well transfected using DharmaFECT 1 following the DharmaFECT pro-

tocol (Dharmacon) with a 25nM siRNA concentration. Approximately 36 hours after

transfection, cells were washed with PBS and harvested for qRT-PCR. For qRT-PCR

siRNA target transcript levels were normalized to a GAPDH loading control to monitor

siRNA silencing efficiency. For immunoblot analysis approximately 36 hours after trans-

fection cells were washed with PBS and replenished with fresh DMEM media. After 4

hours in the case of WT Tg and 8 hours in the case of C1264R or A2234D Tg, cells and

media samples were harvested. Cells were lysed with 1mL of RIPA with protease inhibi-

tor cocktail (Roche), and lysate and media samples were subjected to immunoprecipita-

tion with G1 Anti-DYKDDDDK affinity resin overnight at 4°C while rocking. After three

washes with RIPA buffer, protein samples were eluted with 3× Laemmli buffer with 100

mM DTT heating at 95ºC for 5 min. Immunoblot quantification was performed using Im-

age Lab Software (BioRad).

1108

^{35}S Pulse Chase Assay

Confluent 6-well dishes of FRT cells (approximately 1×10^6/well) were metabolically la-

beled in DMEM depleted of methionine and cysteine and supplemented with EasyTag

^{35}S Protein Labeling Mix (Perkin Elmer, NEG772007MC), glutamine, penicillin/strepto-

mycin, and 10% FBS at 37ºC for 30 minutes. Afterward, cells were washed twice with

F12 media containing 10X methionine and cysteine, followed by a burn off period of 10

minutes in normal F12 media. Cells were then chased for the respective time periods

with normal F12 media, lysed with 500uL of RIPA buffer with protease inhibitor cocktail

(Roche) and 10 mM DTT. Cell lysates were diluted with 500uL of RIPA buffer with pro-

tease inhibitor cocktail (Roche) and subjected to immunoprecipitation with G1 anti-

DYKDDDDK affinity resin overnight at 4ºC. After three washes with RIPA buffer, protein

samples were eluted with 3× Laemmli buffer with 100 mM DTT heating at 95ºC for 5

minutes. Eluted samples were then separated by SDS-PAGE, and gels were dried and

exposed on a storage phosphor screen. Radioactive band intensity was then measured

using a Typhoon Trio Imager (GE Healthcare) and quantified by densitometry in Image

Lab (BioRad).

For pulse chase analysis coupled with ML-240 treatment, cells were pre-treated

with vehicle (0.1% DMSO) or ML-240 (10µM) for 15 minutes prior to ^{35}S pulse labeling.

Vehicle (0.1% DMSO) or ML-240 (10µM) treatment was then maintained throughout the

pulse labeling and chase period.

1129

VCP Pharmacological Inhibition Studies

For follow-up experiment with VCP inhibitors, confluent 6-well dishes of C1264R Tg-FT

FRT cells were washed with PBS and fresh F12 media supplemented with ML-240

(10µM), CB-5083 (5µM), NMS-873 (10µM), or vehicle (0.1% DMSO) was added and in-

cubated for 8 hours. For WT Tg-FT cells, confluent 6-well dishes were similarly used,

washed with PBS and fresh F12 media supplemented with ML-240 (10µM) or vehicle

(0.1% DMSO) was added and incubated for 4 or 8 hours. Cells were lysed with 1mL of

RIPA with protease inhibitor cocktail (Roche), and lysate and media samples were sub-

jected to immunoprecipitation with G1 Anti-DYKDDDDK affinity resin overnight at 4°C

while rocking. After three washes with RIPA buffer, protein samples were eluted with 3×

Laemmli buffer with 100 mM DTT heating at 95°C for 5 minutes. Immunoblot

quantification was performed using Image Lab Software (BioRad).

Viability with ML-240 was monitored using propidium iodide staining. Briefly, cells

were treated with either ML-240 (10µM) or vehicle (0.1% DMSO) for 4 hours before be-

ing harvested and stained with propidium iodide (1µg/mL) at room temperature for 15

minutes in the dark. Cells permeabilized with 0.2% Triton were used as a positive stain-

ing control. Data analysis was then carried out in FlowJo (BD Biosciences).

1147

TEX264 & Tg-NLuc Co-Immunoprecipitation Studies

HEK293 flp-Tg-NLuc cells were cultured at 1.0×10^5 cells/well in 12-well tissue culture

dishes for 1 day and transfected with either a fluorescent control, c-terminal FLAG-tag

TEX264, or c-terminal FLAG-tag SEC62 using a calcium phosphate method. Confluent

plates were harvested by lysing with 300uL of TNI buffer (50mM Tris pH 7.5, 150 mM

NaCl, 0.5% IGEPAL CA-630 (Sigma-Aldrich)) and protease inhibitor (Roche). Lysates

were sonicated for 10 minutes at room temperature and normalized using a BCA Assay

(Thermo Scientific). Cell lysates were then precleared on 4B sepharose beads (Sigma,

4B200) at 4°C for 1 hour while rocking, then immunoprecipitated with G1 Anti-

DYKDDDDK affinity resin overnight at 4°C while rocking. Resin was washed four times

with TNI buffer and resuspended in 250 µL of TNI buffer. 50 µL aliquots were taken and

measured using the nano-glo luciferase assay system according to the manufactures

protocol (Promega). For differential enrichment, statistical analysis was performed in

Prism 9 (GraphPad) using a one-way ANOVA with post hoc Tukey's multiple testing

corrections.

For western blot analysis HEK293 flp-Tg-NLuc 1.0×10^6 cells/dish in 10 cm tissue

culture dishes for 1 day and transfected with either a fluorescent control, TEX264, or

SEC62 using a calcium phosphate method. Cells were collected and lysed using 300

µL of TNI buffer with protease inhibitor (Roche) sonicated at room temperature for 10

minutes. Lysates were normalized, precleared, and immunoprecipitated as described

above. Proteins were eluted using 6X Laemmli buffer + 100mM DTT, separated by

SDS-PAGE and transferred to PVDF membrane (Millipore) for immunoblotting.

1170

QUANTIFICATION AND STATISTICAL ANALYSIS

Mass Spectrometry Interactomics and TMT Quantification

To identify Tg interactors (-) biotin pulldown vs (-) Hpg samples were processed using the DEP pipeline (Zhang et al., 2018). Enriched proteins were determined based on those with a log2 fold change of 2σ and Benjamini-Hochberg adjusted p-value (false discovery rate) of 0.05. For pathway enrichment analysis of identified proteins, EnrichR was used and GO Cellular Component and Molecular Function 2018 terms were used to differentiate secretory pathway associated proteins from background (Chen et al., 2013). For time resolved analysis, data were processed in R with custom scripts. Briefly, TMT abundances across chase samples were normalized to Tg TMT abundance as described previously and compared to (-) Hpg samples for enrichment analysis (Wright et al., 2021). Inconsistencies in the quantification of Tg bait protein were observed for replicate 5 of WT Tg Hpg-chase samples, likely due to sample loss during the enrichment. Hence, this replicate was excluded from further time-resolved analysis. All analysis scripts are available as described in the Data and Code Availability section.

DATA AND CODE AVAILABILITY

Mass spectrometry spectrum and result files are available via ProteomeXchange under identifier PXD035681. Raw TMT quantification data is available in Table S7. Code used for data analysis and generation of figures is available at github.com/wrighmt1/2022_TRIP.

1193

Figure Legends

Figure 1. Time-Resolved Interactome Profiling to Identify Interactions with New

Synthesized Proteins.

(A) Key steps necessary for the TRIP workflow utilizing pulsed unnatural amino acid la-

beling with Hpg (Box 1), labeled protein functionalization with CuAAC click chemistry

(Box 2), global immunoprecipitation (Box 3), followed by streptavidin-biotin enrichment

into time-resolved fractions (Box 4).

(B-D) Western blot analysis of the two-stage purification strategy with continuous Hpg

labeling. FRT cells stably expressing WT Tg were pulse-labeled with Hpg (200µM) for 4

hrs and crosslinked with DSP (0.5mM) for 10 minutes to capture transient proteostasis

network interactions (+ Xlink). Lysates were functionalized with a TAMRA-Azide-PEG-

Desthiobiotin probe using copper catalyzed azide-alkyne cycloaddition (CuAAC) and

subjected to the dual affinity purification scheme as described.

(B) FLAG IP inputs as described in Box 2 in (A). Top shows fluorescence image of rho-

damine labeled proteins, and bottom shows immunoblots of Tg (IB: FLAG) and interac-

tors HSP90B1, HSPA5, PDIA4 and loading control GAPDH.

(C) FLAG IP elutions as described in Box 3 in (A). Top shows fluorescence image of

rhodamine labeled Tg and total Tg (IB: FLAG). Bottom shows immunoblots of interac-

tors (HSP90B1, HSPA5, and PDIA4). Samples were then subjected to biotin pulldown.

(D) Biotin pulldown elutions as described in Box 4 in (A). Top shows fluorescence image

of rhodamine labeled Tg and total Tg (IB: FLAG). Bottom shows immunoblots of interac-

tors (HSP90B1, HSPA5, and PDIA4).

1216

Figure 2. TRIP Temporally Resolves Tg Interactions with Protein Quality Control

Components

(A) Workflow for TRIP protocol utilizing western blot or mass spectrometric analysis of

time-resolved interactomes. Samples were pulse-labeled with Hpg (200μM final concen-

tration) for 1 hour, then chased in regular media for specified time points, and cross

linked with DSP (0.5mM) for 10 minutes to capture transient proteoastasis network in-

teractions. Lysates were functionalized with a TAMRA-Azide-PEG-Desthiobiotin probe

using copper catalyzed azide-alkyne cycloaddition (CuAAC). Chase samples were pro-

cessed through the dual affinity purification TRIP workflow and processed for mass

spectrometry or western blot analysis. (-) Hpg samples were unlabeled and served as

negative controls.

(B and C) TRIP western blot analysis of WT Tg-FT (B) and C1264R Tg-FT (C). Samples

were processed as described above in (A). FLAG IP Input panel shows immunoblot of

Tg (IB:FLAG), and interactors HSP90B1, HSPA5, PDIA4 and loading control GAPDH.

Biotin pulldown elution panel shows fluorescence image of rhodamine labeled Tg and

immunoblot of Tg interactors HSP90B1 and HSPA5. Validated Tg interactors show

higher and delayed enrichment with the misfolded C1264R Tg mutant in (C) compared

to WT in (B).

(D and E) Plots showing the log2 fold change enrichment of select Tg interactors

HSPA5 and PDIA4 compared across constructs. Samples were processed as described

above in (A) and analyzed by mass spectrometry. Temporal trends from western blot

analysis are recapitulated with mass spectrometry analysis.

1239

Figure 3. TRIP Identifies Altered Temporal Dynamics Associated with Tg Pro-

cessing

(A) TMTpro labeling scheme for TRIP experiments including a (-) Hpg control channel to

identify enriched interactors and a (-) Biotin pulldown channel to act as a booster (or

carrier). Chase samples are pulse labeled with Hpg (200µM) for 1 hour. Cells are then

harvested at specified time points and cross linked with DSP (0.5mM) for 10 minutes to

capture transient proteoastasis network interactions. Lysates are functionalized with a

TAMRA-Azide-PEG-Desthiobiotin probe using copper catalyzed azide-alkyne cycloaddi-

tion (CuAAC). Chase samples are processed through the dual affinity purification TRIP

workflow and processed for mass spectrometry. (-) biotin pulldown samples were col-

lected after the first FLAG co-IP and serve as positive control and booster channel. (-)

Hpg samples were processed through the dual affinity purification TRIP workflow ,in-

cluding 3 hr chase period, serving as negative controls to assess background.

(B) Venn diagram showing the overlap in proteostasis components identified as Tg in-

teractors here compared to our previous data set (Wright et al., 2021; Zhang et al.,

2018). (-) biotin pulldown vs (-) Hpg samples were used to identify Tg interactors in FRT

cells.

(C) Plot showing the relative pathway enrichment for Hsp70/90 assisted folding & chap-

eroning interactors with WT, A2234D, or C1264R Tg constructs. For this and all

subsequent panels, the average log2 fold change enrichment value across time points

for a given interactor were used to scale data. Positive enrichment was scaled from 0 to

1. Lines represent the median scaled enrichment for the group of interactors and

shades correspond to the first and third quartile range. All source data for this and sub-

sequent panels can be found in Table S6.

(D) Heatmap showing the relative enrichment for individual Hsp70/90 assisted folding &

chaperoning interactors with WT, A2234D, or C1264R Tg constructs. Relative enrich-

ment scaled as described above in (C). Source data found in Table S6.

(E) Heatmap showing the relative enrichment for select membrane biogenesis interac-

tors with WT and C1264R Tg constructs.

(F) Plot showing the relative pathway enrichment for disulfide/redox processing interac-

tors with WT, A2234D, or C1264R Tg constructs. Relative enrichment scaled and plot-

ted as described above in (C). Lines and shades are defined as in in (C).

(G) Heatmap showing the relative enrichment for individual disulfide/redox processing

interactors with WT, A2234D, or C1264R Tg constructs.

(H) Heatmap showing the relative enrichment for select prolyl isomerization & hydrox-

ylation interactors with WT, A2234D, or C1264R Tg constructs.

(I) Heatmap showing the relative enrichment for select glycan processing Tg interactors

with WT, A2234D, or C1264R Tg constructs.

(J) Plot showing the relative pathway enrichment for proteasomal degradation Tg in-

teractors with WT, A2234D, or C1264R Tg constructs. Lines and shades are defined as

in in (C).

(K) Heatmap showing the relative enrichment for select proteasomal degradation Tg in-

teractors with WT, A2234D, or C1264R Tg constructs.

(L) Heatmap showing the relative enrichment for select autophagy Tg interactors with

WT, A2234D, or C1264R Tg constructs.

1285

Figure 4. siRNA Screening Deconvolutes Interactions and Key Regulators of Con-

struct Specific Tg Processing

(A) siRNA screening workflow utilizing NLuc-tagged Tg to monitor lysate and media

abundance. Approximately 36 hours after transfection with 25nM siRNAs cells were re-

plenished with fresh media and Tg-NLuc abundance in lysate and media was measured

after 4 hours using the nano-glo luciferase assay system.

(B) Violin plots showing the relative Tg-NLuc abundance changes in lysate with siRNA

knockdown of select genes. Tg-NLuc abundance in lysate was measured 4 hours after

replenishing with fresh media using the nano-glo luciferase assay system. Cutoff criteria

for hits were set to those genes that increased or decreased Tg-NLuc abundance in ly-

sate or media by 3σ. Hits are labeled within the plot. Source data found in Table S7.

(C) Violin plots showing the relative Tg-NLuc abundance changes in media with siRNA

knockdown of select genes. Sample processing and cutoff criteria to identify hits are de-

scribed above in (B). Source data found in Table S7.

1300

1301　**Figure 5. Trafficking and Degradation Factors Selectively Regulate Tg Processing**

1302　**in FRT cells**

1303 (A and B) Western blot analysis of WT Tg-FT secretion from FRTR cells after transfec-

1304 tion with select siRNAs hits from initial screening data set. Cells were transfected with

1305 25nM siRNAs for 36 hours, media exchanged and conditions for 4 hrs, Tg-FT was im-

1306 munoprecipitated from lysate and media samples, and Tg-FT amounts were analyzed

1307 via immunoblotting.

1308 (C and D) Western blot analysis of C1264R Tg-FT secretion from FRT cells transfected

1309 with select siRNA hits from the initial screening data set. Samples were processed as in

1310 (A). All statistical testing performed using an unpaired student's t-test with Welch's cor-

1311 rection. *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$.

1312

1313 **Figure 6. Mutant Tg is Selectively Enriched with the ER-phagy Receptor TEX264**

1314 (A) Schematic for western blot and NLuc analysis for identifying TEX264-Tg interac-

1315 tions. HEK293T cells stably expression WT or mutant Tg-NLuc were transfected with

1316 FLAG-tagged ER-phagy receptors TEX264, SEC62, or GFP control. After FLAG Co-IP,

1317 samples were analyzed using the using the nano-glo luciferase assay system or im-

1318 munoblot analysis to monitor Tg enrichment.

1319 (B) Luminescence data from FLAG-CoIP and nano-glo luciferase analysis. Tg is selec-

1320 tively enriched with TEX264 compared to GFP fluorescent control, or SEC62. Mutant Tg

1321 exhibits higher enrichment compared to WT. Data represented as mean ± SEM. Statisti-

1322 cal testing was performed using a one-way ANOVA with post hoc Tukey's multiple test-

1323 ing corrections, *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$.

1324 (C) Western blot analysis of FLAG Co-IP samples. Top panel shows IP inputs and bot-

1325 tom panel IP elutions with Tg (IB:Tg), ER-phagy receptors (IB:FLAG and IB:TEX264)

1326 and loading control Tubulin. Tg is selectively enriched with TEX264 compared to GFP

1327 fluorescent control, or SEC62, with C1264R Tg exhibiting higher enrichment compared

1328 to WT.

1329

Figure 7. Pharmacological VCP Inhibition Selectively Rescues C1264R Tg Secre-

tion through Interactome Remodeling

(A and B) Western blots analysis of C1264R Tg-FT FTR cells treated with VCP inhibi-

tors ML-240 (10µM), CB-5083 (5µM), NMS-873 (10µM), or vehicle (0.1% DMSO) for 8

hours. Lysate and media samples were subjected to immunoprecipitation and analyzed

via immunoblotting. Data is normalized to the median C1264R Tg-FT abundance of

DMSO treated samples. Data represented as mean ± SEM. Statistical testing performed

using an unpaired student's t-test with Welch's correction, $*p<0.05$, $**p<0.005$,

$***p<0.0005$.

(C and D) Western blots analysis of WT Tg-FT FTR cells treated with ML-240 for 4 and

8 hours. Lysate and media samples were processed and analyzed as described above

(A and B). Data is normalized to the median WT Tg-FT abundance of DMSO treated

samples. Data representation and statistical testing as in (B)

(E) ^{35}S Pulse-chase analysis of C1264R Tg-FT FTR cells with ML-240 treatment. Cells

were pre-treated with ML-240 or DMSO for 15 minutes prior to pulse labeling with ^{35}S

for 30 minutes and chased for 4 hours with DMSO or ML-240 treatment. Data is repre-

sented as mean ± SEM. Full data panel available in Figure S7C and S7D.

(F) ^{35}S Pulse chase analysis of WT Tg-FT FRT cells with ML-240 (10µM) treatment.

Samples processed as described above in (E). Data is represented as mean ± SEM.

Full data panel available in Figure S7E and S7F.

(G-L) TRIP analysis showing the relative enrichment for select Tg interactors across

proteostasis pathways between untreated and ML-240 (10µM) treated C1264R Tg.

Cells were treated with ML-240 (10µM) and analyzed utilizing the TRIP workflow. The

average log2 fold change enrichment values across time points for a given interactor

are used to scale data. Positive enrichment values were scaled from 0 to 1, while nega-

tive enrichment values were scaled from 0 to -1. The enrichment is shown for Glycan

Processing (G and H), Hsp70/90 Assisted Folding (I-J), and Disulfide/Redox Processing

(K-L). Aggregate pathway enrichment is displayed in (G), (I), and (J), where lines repre-

sent the median scaled enrichment for the group of interactors and shades correspond

to the first and third quartile range. Heatmaps with individual enrichments of interactors

is shown in (H), (J), and (L). Full data panel available in Figure S8A-S8C.

1361

Figures

Figure 1

A

Box 1

Cytosol

Hpg unnatural amino acid labeling & in situ protein crosslinking

Endoplasmic Reticulum Lumen

Box 2

Cell lysis & CuAAC functionalization (FLAG co-immunoprecipitation input)

CuAAC

Detection & Purification

Box 3

Hsp70/90
Thyroglobulin
CANX

FLAG Antibody Bead

Hsp70/90
Thyroglobulin
CANX

FLAG co-immunoprecipitation & elution (biotin pulldown input)

Box 4

biotin pulldown & elution

Thyroglobulin
Hsp70/90
CANX

Streptavidin Bead

Hsp70/90
Thyroglobulin
CANX

B

Hpg + − +
Xlink − + +

Total Protein

Rhodamine Probe

FLAG IP Input

Thyroglobulin
250 kDa   IB: FLAG

Interactors
100 kDa   HSP90B1   IB: KDEL
75 kDa   HSPA5
75 kDa   IB: PDIA4
37 kDa   IB: GAPDH

C

Hpg + − +
Xlink − + +

Thyroglobulin
250 kDa   Rhodamine Probe

IB: FLAG

Biotin PD Input
250 kDa

Interactors
100 kDa   HSP90B1   IB: KDEL
75 kDa   HSPA5
75 kDa   IB: PDIA4

D

Hpg + − +
Xlink − + +

Thyroglobulin
250 kDa   Rhodamine Probe

IB: FLAG
250 kDa

Interactors
100 kDa   HSP90B1   IB: KDEL
75 kDa   HSPA5
75 kDa   IB: PDIA4

Figure 2.

Figure 3.

Figure 4.

Figure 5.

A

WT Tg-FT   IB: FLAG

C

C1264R Tg-FT   IB: FLAG

B

Lysate

Media

D

Lysate

Media

Figure 6.

A

B

Figure 7.