**A unique case in which *Kimoto*-style fermentation was completed with**

***Leuconostoc* as the dominant genus without transitioning to *Lactobacillus***

**Authors**

Kohei Ito,[1*] Ryo Niwa,[1,2] Yuta Yamagishi,[1,3] Ken Kobayashi,[1] Yuji Tsuchida,[4] Genki Hoshino,[4]

Tomoyuki Nakagawa,[5] Takashi Watanabe[6]

**Affiliation**

1. BIOTA inc., Tokyo, 101-0022, Japan

2. Graduate School of Medicine, Kyoto University, Kyoto, 606-8501, Japan

3. Department of Life Science, College of Science, Rikkyo University, Tokyo, 171-8501, Japan

4. Tsuchida Sake Brewery, Gunma, 378-0102, Japan

5. Faculty of Applied Biological Sciences, Gifu University, Gifu, 501-1193, Japan

6. Gunma Industrial Technology Center, Gunma 379-2147, Japan

* Corresponding author (e-mail: kohei@biota.ne.jp)

**Abstract**

The *Kimoto*-style fermentation starter is a traditional preparation method of *sake*

brewing. In this process, specific microbial transition patterns have been observed within

nitrate-reducing bacteria and lactic acid bacteria during the production process of the

fermentation starter. We have characterized phylogenetic compositions and diversity of the

bacterial community in a *sake* brewery performing the *Kimoto*-style fermentation. Comparing

the time-series changes with other *sake* breweries previously reported, we found a novel type

of *Kimoto*-style fermentation which the microbial transition differed significantly from other

breweries during the fermentation step. Specifically, the lactic acid bacteria, *Leuconostoc* spp.

was a predominant species in the late stage in the preparation process of fermentation starter,

on the other hand, *Lactobacillus* spp., which plays a pivotal role in other breweries, was not

detected in this analysis. The discovery of this new variation of microbiome transition in

*Kimoto*-style fermentation has further deepened our understanding of the diversity of *sake*

brewing.

[**Keywords:** Microbiome, Japanese-*sake*, *Kimoto*-style, Lactic acid bacteria, Fermentation]

**INTRODUCTION**

*Sake* is a traditional alcoholic beverage made from rice in Japan. In a preliminary stage

of a main brewing process, a fermentation starter with purely cultured yeast is produced to

prevent microbial contamination and poor fermentation and to promote smooth alcoholic

fermentation. The production process is as follows; First, *Aspergillus oryzae*, which secretes

amylases, is propagated on steamed rice to make *koji*. Then, *koji*, steamed rice, and water are

mixed in an open-top tank. Fermentation of this mixture produces a fermentation starter. The

fermentation starter is further mixed with *koji*, steamed rice, and water, and after a 3-5 week

fermentation process, the fermentation mash or starter is produced. The fermentation starter is

separated into *sake* and spent rice by a filter press to complete the *sake*.

The fermentation starter is divided into three styles, *Sokujo*, *Kimoto* and *Yamahai*.

*Sokujo*-style fermentation starter is a modern method to make the starter culture with the

addition of food-grade lactic acid. *Kimoto*-style fermentation starter is the traditional

preparation method of the starter culture and is manufactured under highly acidic conditions by

inducing the growth of nitrate-reducing bacteria and lactic acid bacteria properly. *Yamahai*-

style is similar to *Kimoto*-style but made without grinding rice. Lactic acid inhibits

contaminations of unintended yeasts and bacteria from external environments into the

fermentation starter.

The microbiome compositions during *Kimoto*-style fermentation starter production

show standard transitions as follows. Nitrate-reducing bacteria, which were reported to come

from water (1), initially grows and produces nitrite, thereby inhibiting the growth of

microorganisms that are less tolerant to nitrite. At the same time, lactic acid bacteria, especially

*Leuconostoc* spp., which grow at low temperatures and have fewer nutrient requirements,

increase, and then lactic acid bacteria such as *Latilactobacillus sakei*, which require strict

nutrients, occupy the microbiome compositions as a predominant species. These steps are

known as a common microbial transition in *sake* brewing (2-4).

However, it was reported that some fermentation starters brewed by *Kimoto*-style in

several *sake* breweries show distinctive microbial transitions and chemical changes and that

this is one of the reasons for producing unique *sake* flavors among breweries, even if they use

the same production process (5-8). A possible reason to explain this fact is differences in

*Kuratsuki* microorganisms (microorganisms living in *sake* breweries) and the introduction of

diverse microorganisms from outside of tanks during the *sake* brewing process (9,10).

The *Tsuchida Sake* Brewery (Gunma, Japan) is one of a few breweries that produce

the *Kimoto*-style fermentation starter without adding yeasts and fully relies on *Kuratsuki*

microorganisms to produce the fermentation starter. In addition, compared to the *Kimoto*-style

fermentation in other *sake* breweries, this *sake* brewery is characterized by not using any food

additives such as brewers alcohol, enzyme reagents, or activated charcoal. Furthermore, this

brewery does not use sake-brewing rice but table rice for *sake* brewing. For the above reasons,

the microbial community and its transition in the fermentation starter of the brewery were

considered to be divergent from previous studies.

In this study, we focused on the fermentation mechanism of the *Kimoto*-style

fermentation starter from the *Tsuchida Sake* Brewery and analyzed its microbial community

during the fermentation process in detail using its 16S rRNA amplicon sequencing. The *Kimoto*-

style fermentation starter from the *Tsuchida Sake* Brewery possesses a distinctive microbial

community compared with other *Kimoto*-style fermentation starters; specifically, the dominant

genus of lactic acid bacteria was *Leuconostoc*, and the switch of the predominant lactic acid

bacteria, from *Leuconostoc* to *Lactobacillus*, was not observed in the fermentation process of

the *Kimoto*-style fermentation starter. Here we report a new profile of microbial transition in

the *Kimoto*-style fermentation starter.

**MATERIALS AND METHODS**

**Sample collection**     Samples were collected for each day 1, 3, 5, 7, 9, 13, 17, 22, 28,

and 33 after starting the fermentation of the fermentation starter. All samples were collected in

duplicate and between October and November 2021. Fermentation starter samples used in this

study were provided by *Tsuchida Sake* Brewing Company (Gunma, Japan). All samples were

immediately frozen and stored until DNA extraction.

**Measurement of basic chemical components**     Temperatures and chemical

compositions of the fermentation starter were measured. Alcohol (ALC) and Baumé degree

(Be) were measured using a DA-155 vibrating density meter for alcoholic beverages (Kyoto

Electronics Manufacturing Co., Ltd., Kyoto, Japan), while Acidity (TA) and Amino Acid

Content (AA) were obtained using a CHA-700 multi-sample changer (Kyoto Electronics

Manufacturing Co., Ltd.). The nitrite concentration was measured using a Merck Millipore

MQuant nitrite test (Merck KGaA, Darmstadt, Germany).

**Total DNA extraction and high throughput sequencing**     Samples were subjected

to 750ul of Lysis buffer, vortexed for 10 min and heat-treated at 100°C for 10 min, centrifuged,

and the supernatant was transferred to MORA beads for automated purification on a Beckman

Coulter GenFindv2 after mechanical fragmentation at a maximum speed of 3 min on an MM-

400.     Finally,     DNA     was     eluted     with     80ul     of     sterile     water.     341F     (5'-

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGGNGGCWGCAG-

3')                              and                              806R                              (5'-

GTCTCGTGGGCTCGGGAGATGTGTATAAGAGACAGGACTACHVGGGGTATCTAA

TCC-3') primers were utilized to amplify the V3-V4 region of the 16S rRNA gene by PCR

(11,12). Thermal cycling conditions were 95 °C for 3 min; 32 cycles of 95 °C for 30 s, 55 °C

for 30 s, and 72 °C for 30 s; and a final extension at 72 °C for 5 min. A second PCR was

performed to add sequencing adapters and dual-index barcodes to distinguish amplicons from

each sample using the same reaction conditions with eight cycles. Preparation of libraries and

sequencing were performed by paired-end sequencing of 300 bp on the Illumina MiSeq

platform (Illumina, Inc., San Diego USA) at GenomeRead Inc. in Kagawa, Japan.

**Microbiome analysis**    The QIIME2 (version 2021.02) (13) platform was used for

microbiome analysis. FASTQ files were imported into the QIIME2 platform. Sequences were

processed using qiime dada2 denoise-paired command for quality control and classification into

amplicon sequencing variants (ASVs).  Taxonomic analysis was run on SILVA database SSU

138 by qiime feature-classifier classify-sklearn (14-16). Sequence reads that were not classified

as any species after phylogenetic classification (Unassigned) and reads classified as chloroplast

and mitochondria were excluded from further analyses. Since the genus *Latilactobacillus* is a

taxon which relatively recently derived from the genus *Lactobacillus* (17), the SILVA database

SSU 138 used in this study does not reflect the reclassification of the genus *Latilactobacillus*,

so it was assigned as *Lactobacillus* in this study.

**Processing for comparative microbiome analysis**    Sequence data from BioProject

PRJDB12939 was used for comparative analysis (8). Since 341F (5'-

CCTACGGGGNGGCWGCAG-3') and 805R (5'-GACTACHVGGGGTATCTAATCC-3')

were used in the study, removal of one base from the 5' end of the reverse read was performed

to match the read lengths of the samples obtained in our study. fastp v.0.20.1 (18) was deployed

for this processing. The method described above was repeated to perform taxonomic analysis.

The depth of sequence reads (Features) differed among samples, and to normalize them, we

subsampled from each sample to 5,000 reads each. Samples with fewer than 5,000 reads (S22,

23, 24, 25, 26, 28) were excluded from the diversity analysis (Table S1). Sub-sampling is an

approach for inferring microbiome differences between samples and has been reported to be a

suitable analytical method when analyzing new data sets (19). To evaluate the effect of

sequence read counts on microbiome diversity assessment, we examined changes in the value

of the Shannon index over a range of reading counts from 0-5,000 by rarefaction curves. The

Rarefaction curve of the Shannon index leveled off when the number of leads was just under

500 (Table S2). Therefore, this investigation suggests that no significant changes in microbial

diversity are due to the subsampling of 5,000 reads from each sample.

**Statistical Analysis** Kruskal-Wallis tests were used to compare the alpha diversity

(Shannon diversity index) among samples. To compare differences in Beta diversity (Weighted

UniFrac distance) between samples, for all PERMANOVA analyses, 5,000 trials were

performed to assess the statistical significance. Q-values < 0.05 after multiple testing

corrections were considered statistically significant. All multiple testing corrections were

performed by computing FDRs using the Benjamini–Hochberg method, and Q-values (adjusted

P-values) < 0.05 were considered statistically significant.

<center>**RESULTS**</center>

**Chemicals and Temperature Change in fermentation starters** The chemical data

and the average temperature of the *sake* starter are shown in Fig. 1. The amount of nitrite started

to rise on day 7 and reached its peak on day 11. On day 14 it was undetectable; Titratable acidity

(TA), a value representing total acidity, increased on day 15, followed by an increase in ethanol

concentration on day 27. In addition, a rapid decrease in glucose was observed along with an

increase in ethanol concentration on day 27. These changes in chemical data and the average

temperature are generally observed in *Kimoto*-style fermentation, so we suggest that the

fermentation proceeded well in the *Kimoto*-style fermentation starter of the *Tsuchida Sake*

Brewery.

**Phylogenetic composition of the microbiome found in the fermentation starter** To

track the microbial transition in the fermentation starter during the fermentation, we performed

16S rRNA amplicon sequencing. Results demonstrated large changes in relative abundances of

the bacterial genera (Genus) during the early to late stages (Fig. 2).

The genera with average relative abundances higher than 5% in the early stage (day 1

to day 7) of the fermentation starters were *Methylobacterium-Methylorubrum* (23.0%),

*Anaerobacillus* (16.7%), *Bacillus* (9.4%), and *Pseudomonas* (9.3%) (Table S3). The relative

abundances of these bacterial genera decreased significantly within days 7-9.

*Methylobacterium* spp. was reported to utilize methanol as a source of carbon and energy and

is widely found on the leaf surface of plants, including rice (20,21). *Anaerobacterium* spp. is

salt-tolerant and halophilic and has been found in lakes and soils in arid regions (22,23). It is

reported that *Bacillus* spp. may live in *sake* breweries as *Kuratsuki* bacteria (24). *Bacillus* spp.

have also been detected in *koji* and are known to be associated with the production of 4-vinyl

guaiacol according to previous studies (9,25,26).

The bacterial genera with average relative abundances higher than 5% in the late stage

(days 17-33) were *Rahnella1* (18.1%), *Serratia* (13.9%), *Hafnia-Obesumbacterium* (8.6%), and

*Carnobacterium* (6.2%) (Table S3). *Leuconostoc, Rahnella, Serratia, Hafnia-*

*Obesumbacterium*, and *Carnobacterium* were the bacterial genera that accounted for a large

proportion of the bacteria found in the late stage (days 17-33). *Leuconostoc* spp. has been

detected in several fermented foods (27,28). *Serratia* spp. have been identified by previous

studies from *Yamahai-moto* (6), *Rahnella* spp. from *Yamahai* (29), and Chinese sauerkraut (30).

*Hafnia-Obesumbacterium* spp. has been detected in brewer's yeast along with *Rahnella* spp.,

suggesting that it may produce high pH beer by inhibiting fermentation reaction (31,32).

*Carnobacterium* spp. are frequently detected in the natural environment and food products,

previous studies reported producing of bacteriocins in some species (33).

**Comparison with other breweries**     To characterize the microbial transition in this

brewery, we compared our data with a previous study that investigated the microbial transition

in four *sake* breweries (8). Time-series changes in the microbial diversity (Alpha diversity;

Shannon index) for each sample are shown (Table 1). *Tsuchida Sake* brewery was confirmed

to have a higher microbial diversity (Shannon index) of the fermentation starters (Q-value <

0.05) compared to the other breweries (Table S4).

A Principal Coordinate Analysis (PCoA) by Weighted UniFrac distance to visualize the

differences (Beta diversity) among the microbiomes of each sample showed that the *Tsuchida*

*Sake* Brewery formed a different cluster (q-value < 0.007) than the other breweries (Fig. 3 and

Table S5).

The relative abundances of the genus *Lactobacillus* and *Leuconostoc* in each brewery

during the preparation are shown in the line graphs (Fig. 4). The genus *Lactobacillus* became

the dominant species in all other breweries, but the relative abundance of the genus

*Lactobacillus* did not increase during the entire preparation in the *Tsuchida Sake* Brewery. On

the other hand, the genus *Leuconostoc* increased in only three breweries, and in all but the

*Tsuchida Sake* Brewery, their abundance eventually decreased and shifted to the genus

*Lactobacillus*. In contrast, in the *Tsuchida Sake* Brewery, the genus *Leuconostoc* maintained a

constant high relative abundance during the fermentation.

## DISCUSSION

**209**

**210**      **Comparative analysis revealed unique dynamics of lactic acid bacteria in the**

**211** ***Tsuchida Sake* Brewery**     A previous study reported three major microbial transition profiles

**212** in sake brewing from the point of switches occurring in lactic acid bacteria: (1) lactic acid

**213** bacteria increase overall, but *Lactobacillus* spp. remain more abundant than *Leuconostoc* spp.,

**214** (2) *Lactobacillus* spp. are more abundant in the beginning, but *Leuconostoc* spp. become

**215** dominant later, and (3) *Leuconostoc* spp. remain more abundant than *Lactobacillus* spp. (8).

**216** Surprisingly, in the brewery we sampled, the genus *Leuconostoc* was detected as the dominant

**217** species even on day 33, the last day of brewing, and only low abundances of the genus

**218** *Lactobacillus* was detected throughout the entire preparation, indicating a microbial transition

**219** profile differs from previous reports.

**220**      In addition, another study reported that *Kimoto*-style fermentation is characterized by

**221** the detection of multiple lactic acid bacteria compared to *Sokujo-moto* (29), so the process of

**222** Kimoto-style fermentation itself may contribute to the diversity of taste in different sake

**223** breweries. *Sokujo-moto* is a modern type of fermentation starter which includes lactic acid to

**224** maintain a low pH, thus preventing microbial contamination from the brewery (5). As indicated

**225** in this study, specific microbiome transitions were observed that were different from the

**226** microbial transitions reported by Takahashi *et al.*, suggesting that the microbiome is unique in

**227** individual breweries and that these microbial transition factors that characterize them need to

**228** be investigated (8).

**229**      Some studies reported that only one lactic acid bacteria appears and the transition does

**230** not occur (34), and no *Lactobacillus* spp. are found (35), but to the best of my knowledge this

**231** is the first study proving this trend of microbiome shift in a manner of uncultured analysis

**232** method.

**233**      A previous study revealed that *L. sakei* has a more stringent amino acid requirement

than *L. mesenteroides*, *Lactobacillus* sp., and di-tripeptides including asparagine, which is

produced by *koji* mold degradation, are growth factors, and that pH and temperature affect the

growth of *L. sakei* (8,36,37). In this study, the peak of nitrite concentration was relatively late

at day 11, and the increase in acidity (TA; Titratable acidity) was delayed accordingly,

suggesting a longer survival period of adventitious bacteria that were initially introduced (Fig.

2). Therefore, there is a possibility that the growth of *L. sakei* was inhibited by specific changes

in nutrients and temperature.

In addition, D-amino acids are attracting attention as new taste components of *sake*, D-

alanine is reported to be produced by amino acid racemases from *L. mesenteroides* with low

temperature (38). *Leuconostoc* spp. capable of high D-amino acid production have also been

isolated (28), therefore, the presence of a high abundance of *Leuconostoc* spp. may contribute

significantly to the taste of *sake*, and need to isolate and culture *Leuconostoc* spp. found in this

brewery for bacterial genome sequencing and detection of unique metabolism pathways.

**Nitrate-reducing bacteria are a major factor in microbial transitions**    In this study,

*Pseudomonas* spp. was detected in the middle stage, and the peak of nitrite reaction was

observed on the 11th day (Fig. 1). On the other hand, previous studies on *Yamahai-moto*

fermentation starters, which is a sake brewing recipe similar to *Kimoto*-style and use lactic acid

bacteria for maintaining the high acidity in the starter, reported cases where *Pseudomonas* spp.

were not detected and no nitrite production was observed (5,39). It is considered that the genus

*Pseudomonas* is easily lysed by exposure to alcohol due to its cell surface structure (40).

Therefore, we believe that *Pseudomonas* spp. was detected until the middle stage, where the

alcohol level rose relatively late, after day 20.

The structure of the microbiome changed significantly before and after the production

of nitrite, suggesting that nitrate-reducing bacteria may also affect microbial transitions and

other factors in the early stage (Fig. 1 and 2). This suggests that the presence or absence of

nitrate-reducing bacteria such as *Pseudomonas* spp. may be one of the factors that cause

differences in the microbial transitions in different breweries.

**Environmental microbiome mapping in the *sake* brewery**   This study allowed us

to detect the bacterial communities with phylogenetic composition during culturing of the

fermentation starter comprehensively. In the early stage, a complex and varied microbiota is

constructed as adventitious bacteria are contaminated by the built environments, tools, and raw

materials used in the brewery. Since the sake brewing process is conducted in an open system,

a variety of adventitious bacteria may contaminate the brewing *sake* from building

environments, tools, and raw materials used in the brewery. Investigating the microbiome from

architectural surfaces and tools in *sake* breweries may reveal the origin of adventitious bacteria

in early sake. It has been suggested that these adventitious bacteria may affect the quality and

taste of sake, so they need to be clarified (29,41). There are still many unexplained aspects of

*Kuratuski* microorganisms, and we believe that this scientific elucidation of the traditional

Japanese liquor will provide significant insights into food microbiology.

**Acknowledgment**

Samples were collected by Mr. Kota Watanabe, Mr. Keitaro Nozaki, Ms. Chinami Fujita,

Ms. Aimi Kurihara, Mr. Tsutomu Watanabe, Mr. Hiroaki Igarashi, Mr. Yuki Taguchi, and Ms.

Mariko Kanazawa of the *Tsuchida Sake* Brewery. Amplicon sequencing was performed by

GenomeRead Inc. We thank Morgenrot Inc. for providing the computational environment for

the analysis. We thank Mr. Masaomi Yanagisawa of Gunma Prefectural Technical Center for

their advice. R.N. is a graduate student of Medical Innovation Program at Kyoto University and

Microbial transition in the *Kimoto*-style starter

284    supported by JST SPRING, Grant Number JPMJSP2110.

**Data availability**

287       The BioSample, DRA/SRA, and BioProject accession numbers for the sequence

288    reported here are SAMD00513369-SAMD00513388, DRR393497-DRR393516, and

289    PRJDB13924 respectively.

## References

1. **Ashizawa, T., and Saito, Y.:** Yamahai shubo ni okeru biseibutsugaku-teki kenkyu (part 12) [Microbiological research on yamahai-shubo (12)], J. Brew. Soc. Jpn, **61,** 638–642. (1966) (in Japanese).

2. **Katagiri, H., and Kitahara, K.:** Studies of lactic acid bacteria isolated from seed mash (part 1), J. Agric. Chem. Soc. Jpn, **10,** 942–951. (1934) (in Japanese).

3. **Obayashi, A., and Kitahara, K.:** On the factor determining the flora of lactic acid bacteria in the starter of sake, J. Agric. Chem. Soc. Jpn, **33,** 839–843. (1959) (in Japanese).

4. **Masuda, Y., Noguchi, T., Takahashi, T., Iguchi, A., Osawa, R., and Mizoguchi, H.:** DGGE and PFGE analysis of lactic acid bacterial succession during *Kimoto* making, Seibutsu-Kogaku Kaishi, **90,** 684–690. (2012) (in Japanese).

5. **Koyanagi, T., Nakagawa, A., Kiyohara, M., Matsui, H., Tsuji, A., Barla, F., Take, H., Katsuyama, Y., Tokuda, K., Nakamura, S., and other 4 authors:** Tracing microbiota changes in yamahai-moto, the traditional Japanese *sake* starter. Biosci. Biotechnol. Biochem., **80,** 399–406. (2016).

6. **Tsuji, A., Kozawa, M., Tokuda, K., Enomoto, T., Koyanagi, T.:** Robust Domination of *Lactobacillus sakei* in microbiota during traditional Japanese *sake* starter yamahai-moto fermentation and the accompanying changes in metabolites, Curr. Microbiol., **75,** 1498–1505. (2018).

7. **Nguyen, T. H. N., Wang, W.-Y., Huang, W-L., Huang, C.-L., Chiang, T.-Y.:** Metagenomics analyses of microbial dynamics associated with putative flavor development in mash fermentation of *sake*, LWT, **163,** 113570. (2022).

8. **Takahashi, M., Morikawa, K., Kita, Y., Shimoda, T., Akao, T., Goto-Yamamoto, N.:** Changes in bacterial and chemical components and growth prediction for *Lactobacillus sakei* during Kimoto-style fermentation starter preparation in *sake* brewing: a

comprehensive analysis. Appl. Environ. Microbiol., **87,** e02546-20. (2021).

9.   **Bokulich, N.A., Ohta, M., Lee, M., and Mills, D.A.:** Indigenous bacteria and fungi drive

traditional kimoto *sake* fermentations. Appl. Environ. Microbiol., **80,** 5522–5529. (2014).

10.   **Akaike, M., Miyagawa, H., Kimura, Y., Terasaki, M., Kusaba, Y., Kitagaki, H., and**

**Nishida, H.:** Chemical and bacterial components in sake and sake production process,

Curr. Microbiol., **77,** 632–637. (2020).

11.   **Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., and**

**Glöckner, F.O.:** Evaluation of general 16S ribosomal RNA gene PCR primers for

classical and next-generation sequencing-based diversity studies. Nucleic Acids Res., **41,**

e1. (2013).

12.   **Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A.,**

**Turnbaugh, P.J., Fierer, N., and Knight, R.:** Global patterns of 16S rRNA diversity at

a depth of millions of sequences per sample. Proc. Natl. Acad. Sci. U S A, **108,** 4516–

4522. (2011).

13.   **Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA,**

**Alexander H, Alm EJ, Arumugam M, Asnicar F, and other 102 authors:**

Reproducible, interactive, scalable and extensible microbiome data science using QIIME

2, Nat. Biotechnol., **37,** 852–857. (2019).

14.   **Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and**

**Glöckner, F.O.:** The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. Nucleic Acids Res., 41(Database issue), D590-596.

(2013).

15.   **Bokulich, N.A., Kaehler, B.D., Rideout, J.R., Dillon M., Bolyen, E., Knight, R.,**

**Huttley, G.A., and Caporaso, J.G.:** Optimizing taxonomic classification of marker-gene

amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome, **6,** 90.

(2018).

16.  **Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., and Holmes,**

**S.P.:** DADA2: High-resolution sample inference from Illumina amplicon data, Nat.

Methods, **13,** 581–583. (2016).

17.  **Zheng, J., Wittouck, S., Salvetti, E., Franz, C.M.A.P., Harris, H.M.B., Mattarelli, P.,**

**O'Toole, P.W., Pot, B., Vandamme, P., Walter, J., and other 5 authors:** A taxonomic

note on the genus Lactobacillus: Description of 23 novel genera, emended description of

the genus *Lactobacillus Beijerinck* 1901, and union of *Lactobacillaceae* and

*Leuconostocaceae*, Int. J. Syst. Evol. Microbiol., **70,** 2782–2858. (2020).

18.  **Chen, S., Zhou, Y., Chen, Y., and Gu, J.:** Fastp: an ultra-fast all-in-one FASTQ

preprocessor, Bioinformatics, **34,** i884–i890. (2018).

19.  **Hughes, J.B, and Hellmann, J.J.:** The application of rarefaction techniques to molecular

inventories of microbial diversity, Methods Enzymol., **397,** 292–308. (2005).

20.  **Yurimoto, H., Iguchi, H., Di Thien, D.T., Tani, A., Okumoto, Y., Ota, A., Yamauchi,**

**T., Akashi, T., and Sakai, Y.:** Methanol bioeconomy: promotion of rice crop yield in

paddy fields with microbial cells prepared from natural gas-derived $C_1$ compound, Microb.

Biotechnol., **14,** 1385–1396. (2021).

21.  **Lai, K., Thai Nguyen N., Miwa, H., Yasuda, M., Huu Nguyen H., and Okazaki, S.:**

Diversity of *Methylobacterium* spp. in the rice of the Vietnamese Mekong delta. Microbes

Environ., **35,** ME19111. (2020).

22.  **Zavarzina, D.G., Tourova, T.P., Kolganova, T.V., Boulygina, E.S., and Zhilina, T.N.:**

Description of *Anaerobacillus alkalilacustre* gen. nov., sp. nov.—Strictly anaerobic

diazotrophic *Bacillus* isolated from soda lake and transfer of *Bacillus arseniciselenatis*,

*Bacillus macyae*, and *Bacillus alkalidiazotrophicus* to *Anaerobacillus* as the new

combinations *A. arseniciselenatis* comb. nov., *A. macyae* comb. nov., and *A.*

*alkalidiazotrophicus* comb. nov. Microbiology (Moscow), **78,** 723–731. (2009).

23. **Blum, J.S., Bindi, A.B., Buzzelli, J., Stolz, J.F., and Oremland, R.S.:** *Bacillus*

*arsenicoselenatis*, sp. nov., and *Bacillus selenitireducens*, sp. nov.: two haloalkaliphiles

from Mono Lake, California that respire oxyanions of selenium and arsenic. Arch.

Microbiol., **171,** 19–30. (1998).

24. **Kanamoto, E., Terashima, K., Shiraki, Y., and Nishida, H.:** Diversity of *Bacillus*

isolates from the *sake* brewing process at a *sake* brewery, Microorganisms, **9,** 1760.

(2021).

25. **Kaneoke, M.:** 4-Vinylguaiacol formation in *sake*, J. Brew. Soc. Jpn, **109,** 320–326. (2014)

(in Japanese).

26. **Sun, L.H., Lv. S.W., Yu, F., Li, S.N., and He, L.Y.:** Biosynthesis of 4-vinylguaiacol

from crude ferulic acid by *Bacillus licheniformis* DLF-17056. J. Biotechnol., **281,** 144–

149. (2018).

27. **Endo, A.:** Diversity of lactic acid bacteria in fermented products, Jpn. J. Lactic Acid

Bacteria, **22,** 87–92. (2011) (in Japanese).

28. **Kato, S., and Oikawa, T.:** Genome sequence of *Leuconostoc mesenteroides* LK-151

isolated from a Japanese *sake* cellar as a high producer of D-amino acids, Genome

Announc., **5,** e00661-17. (2017).

29. **Terasaki, M., Fukuyama, A., Takahashi, Y., Yamada, M., and Nishida, H.:** Bacterial

DNA detected in Japanese rice wines and the fermentation starters, Curr. Microbiol., **74,**

1432-1437. (2017).

30. **Zhou, Q., Zang, S., Zhao, Z., and Li, X.:** Dynamic changes of bacterial communities and

nitrite character during northeastern Chinese sauerkraut fermentation. Food Sci.

Biotechnol., **27,** 79–85. (2017).

31. **Vaughan, A., O'Sullivan, T., and van Sinderen, D.:** Enhancing the microbiological

stability of malt and beer–A review, J. Inst. Brew., **111,** 355–371, (2005).

32.   **Ashtavinayak, P., and Elizabeth, H.A.:** Review: Gram negative bacteria in brewing.

Adv. Microbiol., **6,** 195–209, (2016).

33.   **Leisner, J.J., Laursen, B.G., Prévost, H., Drider, D., and Dalgaard, P.:**

*Carnobacterium*: positive and negative effects in the environment and in foods, FEMS

Microbiol. Rev., **31,** 592–613. (2007).

34.   **Ashizawa, T.:** Yamahai shubo ni okeru biseibutsugaku-teki kenkyu (part 10)

[Microbiological research on yamahai-shubo (10)], J. Brew. Soc. Jpn, **60,** 900–903. (1965)

(in Japanese).

35.   **Momose, H., and Kamao, A.:** Lactic acid cocci isolated from Moto (*sake* starter) prepared

by traditional method, J. Brew. Soc. Jpn, **88,** 76–80. (1993) (in Japanese).

36.   **Mizoguchi, H.:** Quality of *sake* characterized by lactic acid bacterial flora in traditional

yeast starter (*Kimoto*), J. Brew. Soc. Jpn, **108,** 382–388. (2013) (in Japanese).

37.   **Yamaji, E., Furukawa, K., Mizoguchi, H., and Hara, S.:** Growth factors required for

the predominance of *Lactobacillus sakei* over *Leuconostoc mesenteroides* in *kimoto*, J.

Brew. Soc. Jpn, **100,** 281–288. (2005) (in Japanese).

38.   **Oikawa, T.:** "D-Amino acid", a new ingredient effecting on *sake* taste, J. Brew. Soc. Jpn,

**110,** 189–197. (2015) (in Japanese).

39.   **Terasaki, M., Miyagawa, S., Yamada, M., and Nishida, H.:** Detection of bacterial DNA

during the process of *sake* production using sokujo-moto. *Curr. Microbiol.*, **75,** 874–879.

(2018).

40.   **Terasaki, M., Nishida, H.:** Bacterial DNA diversity among clear and cloudy sakes, and

*sake-kasu*. Open Bioinformatics J., 13, 74–82. (2020).

41.   **Nishida, H.:** *Sake* brewing and bacteria inhabiting *sake* breweries, Front. Microbiol., **12,**

602380. (2021).

**Figure legends**

FIG. 1.  Time-series data of chemical concentrations, degree, and average temperature. Average

temperature (A), Baume degree (B), Alcohol (C), Glucose (D), Amino acid content (E),

Acidity (F), Nitrite (G).

FIG. 2. Taxonomic composition of microbiomes during *Kimoto*-style fermentation starter in

426      *sake* brewing. The top 15 genera are listed, and the rest are noted as "Remainder".

429 FIG. 3. Principal coordinate analysis (PCoA) of 5 *sake* breweries. Samples are compared using

430 the weighted UniFrac distance metrics.

FIG. 4. Line plots of relative abundances of *Lactobacillus* and *Leuconostoc*.

--------

**Tables**

TABLE 1. Time-series changes in Shannon index for each sample.

# Microbial transition in the *Kimoto*-style starter

| days | sample_type | brewery | Shannon index |
|---|---|---|---|
| 5 | shubo | Tsuchida | 4.083626084 |
| 7 | shubo | Tsuchida | 4.483739465 |
| 9 | shubo | Tsuchida | 4.00914674 |
| 9 | shubo | Tsuchida | 3.260447879 |
| 13 | shubo | Tsuchida | 3.955204079 |
| 13 | shubo | Tsuchida | 3.953384193 |
| 17 | shubo | Tsuchida | 2.683731987 |
| 17 | shubo | Tsuchida | 2.54526448 |
| 22 | shubo | Tsuchida | 3.246816955 |
| 22 | shubo | Tsuchida | 2.970864866 |
| 28 | shubo | Tsuchida | 2.966174624 |
| 28 | shubo | Tsuchida | 3.072856664 |
| 33 | shubo | Tsuchida | 2.925714709 |
| 33 | shubo | Tsuchida | 3.056336844 |
| 3 | shubo | A | 0.075588383 |
| 6 | shubo | A | 0.47124225 |
| 9 | shubo | A | 1.748384977 |
| 10 | shubo | A | 1.517717158 |
| 11 | shubo | A | 1.252412288 |
| 12 | shubo | A | 1.496213641 |
| 13 | shubo | A | 1.682703746 |
| 15 | shubo | A | 1.732575907 |
| 16 | shubo | A | 1.486121819 |
| 3 | shubo | B | 2.38193983 |
| 6 | shubo | B | 2.933967722 |
| 7 | shubo | B | 3.090082101 |
| 8 | shubo | B | 3.294985987 |
| 9 | shubo | B | 3.142318413 |
| 10 | shubo | B | 2.2285432 |
| 12 | shubo | B | 1.835421019 |
| 13 | shubo | B | 2.113163833 |
| 14 | shubo | B | 2.012528982 |
| 3 | shubo | C | 1.837678435 |
| 4 | shubo | C | 2.042470055 |
| 5 | shubo | C | 1.28555966 |
| 7 | shubo | C | 1.261663511 |
| 8 | shubo | C | 1.221165987 |
| 9 | shubo | C | 1.037990141 |
| 11 | shubo | C | 1.10982519 |
| 13 | shubo | C | 1.085215814 |
| 16 | shubo | C | 1.016089562 |
| 3 | shubo | D | 1.812960613 |
| 6 | shubo | D | 2.714883385 |
| 8 | shubo | D | 3.393236556 |
| 9 | shubo | D | 2.492923589 |
| 10 | shubo | D | 2.097860823 |
| 11 | shubo | D | 2.214411678 |
| 13 | shubo | D | 2.233278552 |
| 3 | shubo | D2 | 0.541406715 |
| 5 | shubo | D2 | 0.55204851 |
| 6 | shubo | D2 | 0.460579254 |
| 7 | shubo | D2 | 0.725185898 |
| 9 | shubo | D2 | 2.65814023 |
| 10 | shubo | D2 | 1.727365975 |
| 11 | shubo | D2 | 1.722444728 |
| 12 | shubo | D2 | 1.675929835 |
| 14 | shubo | D2 | 1.64604447 |
| 3 | shubo | E | 0.806481485 |
| 5 | shubo | E | 1.332881408 |
| 6 | shubo | E | 2.553896711 |
| 7 | shubo | E | 1.644847183 |
| 9 | shubo | E | 1.56711282 |
| 11 | shubo | E | 1.427864233 |
| 12 | shubo | E | 1.328343618 |
| 13 | shubo | E | 1.247446865 |
| 16 | shubo | E | 1.325637366 |