Plants stand still but hide: imperfect and heterogeneous

detection is the rule when counting plants

Authors: Jan Perret[1], Aurélien Besnard[1], Anne Charpentier[1], Guillaume Papuga[2]

1. CEFE, Univ Montpellier, CNRS, EPHE-PSL University, IRD, Montpellier, France

2. AMAP, IRD, CIRAD, CNRS, INRAE, Univ Montpellier, Montpellier, France

Corresponding author: Jan Perret (jan.perret@gmail.com)

Orcid:

- Besnard A.: https://orcid.org/0000-0002-2289-9761

- Charpentier A.: https://orcid.org/0000-0002-8588-6001

- Papuga G.: https://orcid.org/0000-0002-7803-2219

- Perret J.: https://orcid.org/0000-0002-1912-6995

Abstract

1. The estimation of population size and its variation largely relies on counts of individuals, generally carried out within spatial units such as quadrats or sites. Missing individuals during counting (i.e. imperfect detection) results in biased estimates of population size and trends. Imperfect detection has been shown to be the rule in animal studies, and most studies now correct for this bias by estimating detection probability. Yet this correction remains exceptional in plant studies, suggesting that most plant ecologists implicitly assume that all individuals are always detected.

2. To assess if this assumption is valid, we conducted a field experiment to estimate individual detection probability in plant counts conducted in 1x1 m quadrats. We selected 30 herbaceous plant species along a gradient of conspicuousness at 24 sites along a gradient of habitat closure, and asked groups of observers to count individuals in 10 quadrats using three counting methods requiring progressively increasing times to complete (quick count, unlimited count and cell count). In total, 158 participants took part in the experiment, allowing an analysis of the results of 5,024 counts.

3. Over all field sessions, no observer succeeded in detecting all the individuals in the 10 quadrats. The mean detection rate was 0.44 (ranging from 0.11 to 0.82) for the quick count, 0.59 for the unlimited count (range 0.18–0.87) and 0.74 for the cell count (range 0.46–0.94).

4.   Detection probability increased with the conspicuousness of the target species and

decreased with the density of individuals and habitat closure. The observer's experience

in botany had little effect on detection probability, whereas detection was strongly

affected by the time observers spent counting. Yet although the more time-consuming

methods increased detection probability, none achieved perfect detection, nor did they

reduce the effect on detection probability of the variables we measured.

5.   *Synthesis and applications.* Given that plant detection was always imperfect and highly

heterogeneous, rather than rely on raw counts, plant ecologists should use methods that

estimate the detection probability of individuals to assess the size, temporal or spatial

trends of plant populations.

Keywords: conservation; demography; detectability; field methods; N-mixture; observer

effect; plant survey; population monitoring.

Introduction

The size of populations and their variation over time and space are of central importance in ecology and conservation. These parameters are required to understand ecosystem dynamics (Ripple & Beschta, 2012; Watson & Estes, 2011), assess population viability (Dennis et al., 1991) and trigger conservation policies and actions (e.g. IUCN, 2019). Their estimation largely relies on counts of individuals, generally made in spatial units such as quadrats or sites (Elphick, 2008; Yoccoz et al., 2001). Therefore, the accuracy of these counts is crucial, as counting errors would result in biased estimates of population sizes and trends.

When carrying out counts, observers can miss some individuals, in effect counting fewer individuals than were present in the sampling units. This phenomenon of imperfect detection, well known in animal studies, leads to underestimating population sizes (Williams et al., 2002). If detection is imperfect but constant over time and space, raw counts can provide unbiased estimates of population trends or spatial variation (Yoccoz et al., 2001). However, if detection varies over time (e.g. across the sampling season or between years) or space (e.g. between habitats), the raw differences between counts performed at different sampling sessions or in different sampling units are a mixture of differences in abundance and detection (Kéry et al., 2009). For example, a decrease in population size over the years may be hidden as observers improve their ability to detect individuals over time (Kendall et al., 1996). Or one might conclude that there are fewer individuals in closed than in open habitats because they are harder to detect in closed habitats (Buckland et al., 2008). Even differences as small as 4–8% in detection dramatically increase the risk of erroneously concluding that population size varies over time or space (Archaux et al., 2012).

Since the 1970s, many studies on vertebrates have shown that imperfect detection is the rule

and that detection probability tends to vary in time and space. Detection probability depends

on a number of factors: the observation process (both the observation method and the survey

effort), the visibility of the target species, and the characteristics of its habitat (Kéry et al.,

2005; Kéry & Schmidt, 2008; Petitot et al., 2014). Consequently, studies including an estimate

of detection probability, which corrects for the bias resulting from imperfect detection, have

become increasingly common for vertebrates and, to a lesser extent, invertebrates. Yet they

remain exceptional for plants (Kellner & Swihart, 2014), suggesting that most plant ecologists

implicitly assume that all individuals are always detected, or that differences in detection

probability are negligible. It is true that, unlike most animals, plants are immobile, and

observers may thus consider it unlikely to miss individuals if careful sampling is conducted.

This assumption led John Harper to make the famous statement "Plants stand still and wait to

be counted," comparing the ease of studying plant and animal populations (Harper, 1977).

Although detection has almost never been investigated in plants at the individual level, some

studies have investigated it at the species level, assessing species occurrence in a given area

using two main approaches: occupancy or time-to-detection designs. In the first approach, the

survey duration is fixed, and the detection probability of the species when surveying quadrats

or sites is estimated from repeated surveys. In the second approach, detection probability is

estimated based on the survey time required to detect the species. These studies found that

the detection of plant species was almost always imperfect, and that detection probability

varied depending on ecological and observational variables. For instance, species detection

varied according to the morphology of the target species (Chen et al., 2009, 2013; Garrard et

al., 2013), with conspicuous growth forms and phenological stages raising detection

probability. A high density of individuals increased the probability of detecting the species

(Bornand et al., 2014), while the closure of the surrounding habitat decreased it (Alexander et

al., 2012; Garrard et al., 2008). Species detection probability also increased with survey effort

(Chen et al., 2009) and often varied between observers (Garrard et al., 2008; Moore et al.,

2011). While these studies confirm that detection issues exist when assessing plant species

occurrence, do these issues affect individual detection to a similar extent?

Concerning detection at the individual level, the only studies that have been conducted on

plants used marked individuals in so-called 'capture–recapture' designs (Lebreton et al., 1992)

or the time-to-detection of individuals (Hauser et al., 2022). These studies found that

individual detection was imperfect, although individuals were generally searched for in

relatively small, intensively surveyed plots (Alexander et al., 1997; Andrieu et al., 2017; Kéry

& Gregg, 2003). In most studies using marked individuals, individual detection probability

varied with the phenological stage of individuals and sometimes with habitat closure (Andrieu

et al., 2017). Hauser et al. (2022) showed that flowering individuals were detected faster than

non-flowering ones, and that the abundance of individuals of species resembling the target

species delayed detection. Observers with recent experience in searching for the target

species detected it faster, confirming at the individual level what has been demonstrated at

the species level. Therefore, it is particularly concerning that individual detection probability

has never been estimated for counts of unmarked plants, despite this being one of the most

commonly used methods for estimating population sizes and trends in plant ecology (Reisch

et al., 2018).

To assess if imperfect detection of individuals is indeed the rule in plant counts, we conducted

a field experiment in which observers counted plants in 1x1 m quadrats. In total, the

experiment covered 30 species, 300 quadrats and involved 158 observers who made 5,024

counts. The questions we wanted to answer were: (1) Are all individuals detected in plant

counts? (2) Is individual detection probability homogeneous in time and space? (3) Does

increasing the survey effort result in higher detection probability, and could this allow perfect

detection to be reached? To explore these questions, we tested whether detection varies

according to ecological variables (species conspicuousness, true population density and

surrounding habitat closure) and observational variables (counting method and experience of

observers), which are likely to vary over time and space, as well as testing varying degrees of

survey effort.

Materials & methods

1 – Selection of sites, species and observers

We selected 30 herbaceous plant species from 24 sites throughout France in Mediterranean,

Alpine and temperate climatic regions (Appendix 1). We chose species with various growth

forms and phenological stages to cover a gradient from cryptic to conspicuous species. As the

experiment focused on counting individuals, we selected only species for which 'individuals'

(ramets or genets depending on the species) could be unambiguously differentiated. We

avoided species difficult to identify so that all could be identified at first glance without

needing to check identification criteria on each detected individual. We selected the study

sites to cover a gradient of habitat closure, from bare ground to dense meadows, to include

all possible combinations of species conspicuousness and habitat closure (Fig. 1).

Participants were colleagues, researchers, undergraduate students and professionals involved

in flora conservation. We selected 4 to 8 observers for each field session, including at least

one with several years of field experience in botany, one with no experience in botany, and

intermediate profiles. Observers were allowed to participate in up to three sessions, but these

sessions had to be several weeks apart. This resulted in 171 participations by 158 unique

observers. We conducted the fieldwork from 7 October 2020 to 6 August 2021.

Figure 1: Diagram of the study's experimental design. We conducted 30 field sessions, each focusing on a different species in a habitat with a different closure level, to cover the widest possible range of combinations of species conspicuousness and habitat closure. For each field session, we placed ten quadrats on the population to form a density gradient and chose 4–8 observers to have a gradient of experience in botany. The observers carried out three counting methods successively (quick, unlimited and cell counts). We then counted the true number of individuals by systematically digging out the individuals detected by the observers and then conducting an additional survey for any individuals that remained unnoticed. Picture A shows the experiment in progress at a Mediterranean site, and picture B shows a quadrat with the cells.

<u>Field set up and observer training</u>

The fieldwork consisted of one-day sessions, during which a group of observers counted the

number of individuals of a target species in a habitat at one of the 24 sites. Before the

observers' arrival, we placed ten 1x1 m quadrats on the target species population. We chose

quadrat locations to ensure that (i) two were free of individuals of the target species (so that

observers would not always expect to find individuals in the quadrats), (ii) they covered a

relatively linear gradient of densities up to a maximum of 150 individuals (when this density

could be found on the site). We labelled the quadrats from 1 to 10 and fixed them to the

ground. When the observers arrived, we explained how the counts would be conducted and

how to identify the target species at all phenological stages and differentiate it from the other

species present at the study site. We then trained them to count on additional quadrats to

check that everyone had understood the instructions. This training phase lasted until all

observers felt comfortable with species identification and the counting methods. The whole

training process usually took about an hour.

Counting methods

Each observer had to count the number of individuals of the target plant in each quadrat using

three different methods, requiring a progressively increasing amount of survey time. In the

first round, participants had 30 seconds to estimate the number of individuals in each quadrat

(the quick count). In the second round, they had to count the number of individuals without

any time limit (the unlimited count). We then divided the quadrats into 100 cells of 10x10 cm

(Fig. 1B), and observers performed the third round, where they had to count the number of

individuals in each cell without any time limit (the cell count). This last method forced

observers to survey the quadrat homogeneously and has been regularly used in the plant

survey literature (e.g. Gauthier et al., 2017, 2019). For the unlimited and cell counts, observers

recorded the time they spent counting in each quadrat. To avoid memory effects, we collected

the sheets after each round, and observers surveyed the quadrats in a different order,

approaching it from a different side at each round. The observers conducted the three counts

on the ten quadrats in 1.5 to 5 hours (on average 3.25 hours).

Exhaustive count and measurement of covariates

After the observers had left, we carried out a final count to determine the true number of

individuals present in each quadrat. This was done by searching for all the individuals detected

per 10x10 cm cell by the observers during the third round and collecting them. Once all

individuals detected by at least one observer were removed, we searched again for any

potentially undetected individuals. This final count took between 1 and 7 hours (3 hours on

average), depending on habitat closure and the number of individuals of the target species in

the quadrats. We calculated the detection probability of individuals by dividing the number of

individuals detected by the observers by the true number of individuals present in each

quadrat.

We measured three variables to investigate their effect on detection probability: an index of

species conspicuousness (measured for each field session), an index of habitat closure

(measured for each quadrat) and the experience in botany of the observers. All the variables

and interactions we investigated in the following analysis are presented in Table 1, as well as

the underlying hypotheses. The way we measured the variables is detailed in Appendix 2.

192   3 – Statistical analysis

Overall, 300 quadrats were surveyed during the experiment by, on average, 5.7 observers,

resulting in 5,024 counts. We removed from the analysis 34 quadrats that contained no

individuals of the target species, as these do not provide information on detection probability.

We withdrew four additional quadrats that unexpectedly contained more than 300 individuals

to avoid the leverage effect due to these density outliers (Appendix 3). In some cases (N = 140,

representing 2.8% of the counts), the observers recorded more individuals than were truly

present. We thus carried out the analyses on two datasets, one in which we removed these

counts and one in which we set these counts to the true number of individuals. As there was

no discernible difference in the results, we present the results of the former analysis in the

main text and the latter in Appendix 7. After all the filtering steps, 4,459 counts remained in

the dataset from which we had removed the observations with excess detections. We also

performed the analyses keeping only the experienced botanists to check that our results were

not driven by an excess of novice observers in our sample. Despite some minor differences,

our overall results remained unchanged (Appendix 8).

We used generalised linear mixed models (GLMMs) with a binomial distribution and a logit

link function to analyse the correlation between detection probability and the variables we

measured. To take into account the repeated measurements at each quadrat, we included

random intercepts for the observers, the field sessions and the quadrats (nested within the

field session). We standardised all the variables prior to model fitting. We started by fitting a

model including all the variables and some of their interactions (see Table 1) and performed a

backward variable selection procedure, dropping every variable and interaction one by one,

keeping only the variables that decreased the Akaike information criterion (AIC) value of the

model (Burnham & Anderson, 2002). For the selected model, we computed the proportion of

variance explained by the model and the proportion of variance explained by the model's fixed

effects, using the theoretical variance method from Nakagawa et al. (2017). Analyses were

performed in R version 3.6.1 (R Core Team, 2019), using packages *lme4* (Bates et al., 2022) to

fit the models and *ggeffects* (Lüdecke et al., 2022) to plot the predictions.

Table 1: Ecological variables (species conspicuousness, habitat closure, true density),

observational variables (counting method, experience in botany), and interaction between

variables tested for their effect on the detection probability of individuals, with the hypothesis

of the underlying mechanism. The measurement method of all variables is detailed in

Appendix 2.

Variable	Description	Measurement method	Hypothesis	Type	Observed values
Species conspicuousness	Conspicuousness of the species at the time of the field session	Expert judgement based on the mean height of individuals and the presence of colourful flowers	Tall and colourful species are easier to detect, which increases their detection probability	Ordinal	1, 2, 3, 4
Habitat closure	Closure of the vegetation in the quadrat	Composite variable created from the vegetation height and the proportion of the area covered by vegetation in the quadrats (Appendix 2)	Dense vegetation masks individuals, which lowers detection probability	Continuous	-1.86–3.54
True density	Number of individuals present in the quadrat	Number of individuals counted in the quadrat by the experimenter at the end of the experiment	High density requires greater concentration and makes it difficult to distinguish between individuals, reducing detection probability	Count	0–295
Counting method	Counting method used	Fixed by the experimental design, each observer successively used the three counting methods on the ten quadrats	Counting methods requiring more survey time yield higher detection probability	Categorical	Quick count, unlimited count, cell count
Experience in botany	General experience in botany of the observer (not specific to the target species)	Self-evaluated by the observers before the experiment using a standardised form	Observers with more experience in botany have higher detection probability	Ordinal	1, 2, 3, 4, 5
Quadrat order + (Quadrat order)²	Position of the quadrat in the round	Defined by the experimental design, the order in which the observers investigated the quadrats was fixed in advance and different for each observer and round. We added the squared variable to specify a quadratic relationship.	Learning increases detection probability over time, but fatigue decreases it	Continuous	1, 2, 3, …, 10

Counting time	Time spent counting by the observer	Each observer timed the duration of counting	Spending more time increases detection probability	Continuous	Quick count: 0.5 min Unlimited count: 0–11 min Cell count: 0–41.58 min
Species conspicuousness x Habitat closure			Detection probability is less affected by habitat closure for conspicuous species	Interaction term	
Species conspicuousness x Counting method			Detection probability obtained with counting methods requiring more survey effort is less affected by species conspicuousness	Interaction term	
Habitat closure x Counting method			Detection probability obtained with counting methods requiring more survey time is less affected by habitat closure	Interaction term	
Experience in botany x Counting method			The difference in detection probability between experts and beginners is larger for the quick count than for the other two methods	Interaction term	

True density x Counting method			Counting methods requiring more survey time make it easier to distinguish between individuals, which decreases the negative effect of true density	Interaction term	
(Quadrat order + (Quadrat order)^2) x Counting method			Counting methods requiring more survey time increase fatigue, which decreases detection probability	Interaction term	
Species conspicuousness x Counting time x Counting method		No coefficient is estimated for the quick count as the time was fixed	Spending an additional extra minute further increases detection probability for species that are easy to detect	Interaction term	
Habitat closure x Counting time x Counting method		No coefficient is estimated for the quick count as the time was fixed	Spending an additional extra minute further increases detection probability in sparse vegetation	Interaction term	

Results

Over the 30 field sessions, no observer succeeded in detecting all the individuals in all the quadrats. Averaged over all quadrats and observers, the mean detection rate per field session was 0.44 (ranging from 0.11 to 0.82) for the quick count, 0.59 for the unlimited count (from 0.18 to 0.87) and 0.74 for the cell count (from 0.46 to 0.94).

Figure 2: Coefficients of standardised variables in the best model. Dots represent estimated coefficients, and lines represent their 95% confidence intervals. Coefficients with a non-significant effect are shown in grey. The reference counting method is the quick count, in which the observers had 30 seconds to estimate the number of individuals in the quadrat. The coefficients presented for the two other methods indicate the difference (on the logit scale) in mean detection probability obtained with those methods compared to the quick count. The variable 'counting time' appears twice because we manually coded the interaction between the counting time and the counting method, so no slope parameter is estimated for the quick count.

After the backward selection procedure, the best model included all the tested variables and
interactions except the squared quadrat order (Appendix 5). All ecological and observational
variables were significantly correlated with detection probability in at least some conditions,
i.e. for some counting methods or some levels of the other variables (Fig. 2; Appendix 6).
Regarding the ecological variables, as expected, the more closed the habitat, the lower the
detection probability (Fig. 3). The more conspicuous the species, the higher the detection
probability, except in entirely open habitats, where the detection probability was constant
and high regardless of the conspicuousness of the species (Fig. 3). Detection probability
decreased steeply with true density, in a similar way for the three counting methods (Fig. 4).
Regarding the observational variables, spending more time sharply increased detection
probability for the unlimited and cell counts. Habitat closure moderated this trend, i.e.
spending an extra minute counting increased detection probability less in closed habitats than
in open habitats. Experienced observers achieved slightly higher detection probability than

inexperienced ones, and this difference was stronger for the cell count than for the unlimited

count. In contrast, there was no significant correlation between experience and detection

probability for the quick count. For the unlimited and cell counts, detection probability slightly

increased with the position of the quadrat in the round, while there was no correlation for the

quick count (Fig. 4).

Figure 3: Predicted detection probability depending on species conspicuousness (1 = least conspicuous; 4 = most conspicuous), habitat closure (-1.5 = most open habitats; 2.5 = most closed habitats), the counting method and the time spent counting. Predictions were made for a quadrat containing 100 individuals of the species of interest, located in the fifth position in the round for an observer who rated his/her level of experience in botany at 2.5 out of 5. Lines represent the predicted mean detection probability, and the shading represents 95% confidence intervals.

Observers achieved higher detection probability with the cell count than the unlimited count,
which yielded higher detection probability than the quick count. However, this came at the
cost of a major increase in the time spent counting. The counting time per quadrat was fixed
at 30 seconds for the quick count, and the mean counting time averaged over all sessions and
observers was 1.8 minutes (ranging from 0.3 to 5.7) for the unlimited count and 8.2 minutes
(from 0.9 to 21.6) for the cell count. While the time spent increased, the slope between
detection probability and the other variables (e.g. observer experience, vegetation closure)
was not reduced. Some variables had roughly the same correlation with detection probability
regardless of the counting method (e.g. true density), while others had an even steeper slope
(e.g. experience in botany).

The model predicts that reaching a detection probability over 0.95 (i.e. 'almost perfect
detection') is only possible in a very open habitat (typically with <7% vegetation cover and
median vegetation height <1 cm) no matter the species' conspicuousness. In a less open
habitat (e.g. vegetation cover <90%, median vegetation height <6 cm), the target species must
be very conspicuous, i.e. a plant above 20 cm in height or 10 cm but with colourful flowers.

Regardless of the species and habitat conditions, reaching this detection probability requires

using the cell count, a true density below 100 individuals per quadrat and a counting time per

quadrat of at least 20 minutes.

Figure 4: Predicted detection probability of individuals depending on the experience in botany of the observer (A), the true density (B) and the position of the quadrat in the round (C). The other variables included in the model are held constant at the following levels: species conspicuousness = 2, habitat closure = 0, quick count time = 0.5 min, unlimited count time = 3 min, cell count time = 3 min. Lines represent the mean detection probability of individuals, and the shading represents 95% confidence intervals.

Substantial variation in detection probability between field sessions remained unexplained by

the model, even after controlling for species and habitat characteristics, as the variance of this

random effect was $\sigma_{session}^2 = 0.58$. There was also unexplained variation between quadrats

($\sigma_{quadrat}^2 = 0.42$). The unexplained between-observer variation was the smallest, with a

variance of $\sigma_{observer}^2 = 0.16$. The proportion of variance explained by the model was

$R^2_{GLMM(c)} = 0.35$, and the proportion of variance explained by the model's fixed effects was

$R^2_{GLMM(m)} = 0.12$. In other words, detection probability remained heterogeneous even after

considering the effect of the seven variables included in the model and accounting for

unexplained inter-session, inter-plot and inter-observer variance.

283   **Discussion**

This study provides the first estimate of the detection probability of individuals in plant counts,

using a detection experiment with a cross-taxa and cross-habitat design. Our results

demonstrated that detection is always imperfect when counting plants, even in small (1 m²)

quadrats with unlimited counting time. Detection probability was also highly heterogeneous

and varied with the species, habitat, observer and counting method. Methods involving more

survey time increased the mean detection probability, but did not achieve perfect detection,

nor did they reduce the effect of the ecological and observational variables on detection

probability. Even after controlling for the effects of all the variables, detection remained highly

heterogeneous. We, therefore, argue that plant ecologists should not rely on the use of raw

counts and should always take into account detection issues when studying plant populations.

1.   Evidence for imperfect detection of plants

In our experiment, while using common plant counting methods (Reisch et al., 2019), not a

single observer detected all the individuals present in the quadrats of a field session. Even

with counts unlimited in time, the predicted detection probability in average conditions (i.e.

for the average values of all variables included in the model with the unlimited count) did not

exceed 0.59 (95% confidence interval: 0.50–0.67). Thus, our results not only confirm that

imperfect detection is the rule when counting plant individuals, but show that detection

probability can even be relatively low. While these values may be surprising, detection issues

have been previously documented for plants when searching for a species (e.g. Chen et al.,

2013), and a recent study conducted on large quadrats (20x20 m) suggested that imperfect

detection may also occur for plant individuals (Hauser et al., 2022). Furthermore, the

detection probabilities we found for plants are higher than those commonly found for animals.

For example, detection probabilities of 0.04–0.32 have been found in bird point counts (Royle,

2004), 0.07–0.41 for lizards counted along transects (Kéry et al., 2009), 0.06–0.41 for

salamanders counted along stream banks (Dodd & Dorazio, 2004), and 0.06–0.34 for

freshwater mussels searched on riverbeds (Carey et al., 2019). Nonetheless, this does not

discount that detection remains far from exhaustive for plants and thus should not be

overlooked.

2. Effects of the ecological variables

Our results showed that species conspicuousness positively affects detection probability,

while habitat closure negatively affects it. This general result was expected, as it has also been

observed for plants in occupancy and time-to-detection studies (Alexander et al., 2012;

Garrard et al., 2008; Hauser et al., 2022). In open habitats, detection probability was the same

regardless of the species' conspicuousness. In contrast, differences between species appeared

with increasing habitat closure, reaching up to a 0.42 difference in detection probability

321 between the extreme levels of species conspicuousness in the most closed habitats (estimated

322 difference for the unlimited count with 3 min counting time). This shows that habitat closure

323 drives the detection probability of plant individuals more than species conspicuousness.

324 Furthermore, the negative effect of the density of individuals on detection probability was

325 unexpected in its magnitude (e.g. a predicted detection probability of 0.67 in a quadrat with

326 20 individuals to 0.37 in a quadrat with 300 individuals, using the unlimited count in average

327 conditions) and occurred whatever the counting method. This phenomenon has never been

328 reported to our knowledge, even for animal taxa, but may be due to the Weber–Fechner law

329 regarding how humans perceive numbers (see Nieder, 2020, for a summary). Indeed, studies

330 on human cognition have shown that it becomes increasingly difficult to estimate the number

331 of items in an area as the number grows, and that there is a general tendency to

332 underestimate it. It is plausible that this phenomenon seldom applies to animal counts, as

333 these do not usually involve counting immobile items in high density as is the case for plants.

3. Observer effect

336 The experience in botany of observers had little effect on detection probability compared to

337 the ecological variables and the counting time. In average conditions, the difference in

338 detection probability between novices and the most experienced observers was 0.04 for the

339 quick count, 0.09 for the unlimited count and 0.17 for the cell count, and confidence intervals

340 largely overlapped except for the cell count (Appendix 6). The advantages of having

341 experience in botany when searching for individuals of a target species may be the ability to

342 more quickly identify an individual once it has been detected, to make fewer identification

errors, and possibly to survey the study area more efficiently by knowing where to search for

the species (Archaux et al., 2006). These advantages were probably minor in our experiment,

given that the observers were trained to identify the target species before starting the counts,

all species were easy to identify, and an efficient surveying strategy was of little use in 1x1 m

quadrats. In this context, the observers' task can be seen as searching for small objects in a

noisy environment, and it seems plausible that their performance was mainly driven by their

ability to stay focused during a repetitive task, their motivation, or their fatigue during the

experiment. However, larger differences between observers might have been observed if we

had involved participants with recent experience in searching for the target species (Hauser

et al., 2022) due to the well-known 'search image' effect, whereby foraging predators improve

their probability of detecting the type of prey they are trained to detect (Ishii & Shimada,

2009). If this effect indeed applies to plant counts, population trends estimated from raw

counts will be biased if the same observers always make the counts, as their detection

probability will increase with time.

4. Comparison of the counting methods

As expected, increasing the survey effort, measured by the time spent counting, massively

increased detection probability (Archaux et al., 2006; Hauser et al., 2022). In average

conditions, the estimated detection probability was 0.36 for the quick count (30 sec counting

time), 0.54 for the unlimited count (1.8 min average counting time) and 0.73 for the cell count

(8.2 min). In addition, for the unlimited and cell counts, observers spending more time

counting achieved higher detection probability (e.g. for the unlimited count in average

conditions, detection probability increased from 0.50 at 1 min to 0.83 at 10 min).

The gain in detection associated with counting individuals per cell instead of the whole

quadrat could be due to two reasons: the cells force the observer to survey the whole surface

area of the quadrat uniformly, and also might reduce uncertainty whether or not an individual

has already been counted, which frequently occurs when individuals are close to each other.

However, contrary to our expectations, increasing survey effort did not reduce the effect of

the ecological variables on detection probability, the most striking being the density of

individuals, which had an almost identical effect on detection probability for all three counting

methods.

5. Unexplained variance and implications of detection probability heterogeneity

Of the random effects included in the model, the most unexplained variation in detection

probability was at the inter-session and inter-quadrat levels. Since we only measured three

synthetic variables to characterise the differences between sessions and quadrats (species

conspicuousness, habitat closure and true density), we expected that unexplained variance

would remain at these levels. For example, we did not measure colour and height

heterogeneity, for the habitat or for the target species, although these variables certainly

impact detection probability (Garrard et al., 2013; Hauser et al., 2022).

The fact that the unexplained inter-observer variation was the lowest of the random effects is

an encouraging result for future studies, as it means that there are few differences in

detection probability between observers compared to the differences between species or

over a habitat closure gradient. Yet overall, more than half (65%) of the variance in detection

probability remained unexplained by our model. Thus, observers had heterogeneous

detection probability between quadrats even after controlling for several variables, which

suggests that factors varying over time for each observer, such as concentration or motivation,

probably explain a large part of the variation in detection probability.

6.   Recommendations for counting plants

Our study shows for plants what has been documented many times for animals: that detection

probability of individuals is less than one and is usually heterogeneous in space and time,

whatever the counting method and survey effort. Using counts affected by imperfect

detection as a proxy for the density of individuals to make comparisons over time and space

is only valid under the hypothesis of constant detection. Our findings show that multiple

ecological and observational variables cause heterogeneity in detection probability. While

observational variables can be standardised, it seems unlikely that ecological variables (i.e.

species conspicuousness, habitat closure and true density of individuals) will be constant in

any study, even those conducted at limited temporal and spatial scales. Furthermore, trying

to achieve almost perfect detection (i.e. >0.95) by increasing survey efforts to fix the issue of

detection heterogeneity seems unrealistic given the resource constraints of most ecology and

conservation studies. We therefore argue that temporal or spatial trends in plant populations

should not be estimated using raw counts, as they could then reflect a trend in ecological

variables rather than the population's fate.

Several methods can be used to estimate individual detection probability and thus estimate

unbiased population sizes or trends (e.g. Lebreton et al., 1992; Buckland et al., 2001). In

particular, the N-mixture method uses repeated counts of unmarked individuals on sampling

units (e.g. quadrats or sites) to estimate detection probability from the differences in the

number of individuals detected between counts (Royle, 2004). With this method, estimates of

population sizes or trends are more precise if detection probability is high and homogeneous

in time and space, if the mean density of individuals is high, and if the sample size, i.e. the

number of sampling units and count repetitions, is high (Royle, 2004; Ficetola et al., 2018).

Thus, for studies using the N-mixture method, no matter the studied population and the

counting method used, we recommend setting a fixed counting time, as leaving it to the

observer's choice will add heterogeneity to detection probability, resulting in less precise

estimates.

In terms of allocating survey effort, there are three options: use a more time-consuming

counting method to increase detection probability, make more count repetitions, or survey

more sampling units. The only way to be sure of the optimal design is to conduct a pilot study

to estimate the mean density of individuals and their detection probability in the population

of interest. However, Ficetola et al. (2018) have shown that when detection probability is

above 0.10, the most advantageous strategy is usually to make three count repetitions and to

increase the number of sampling units as much as possible. Conversely, when detection

probability is below 0.10, it is better to increase it by using a more time-consuming counting

method or to increase the number of count repetitions. Thus, we recommend counting

individuals in quadrats without cells with a fixed short counting time (e.g. 0.5–1 min), as this

method will yield a detection probability above 0.10 for most species and allow the most

sampling units to be surveyed. More time-consuming counting methods should be used only

if the study population grows in a very closed habitat and individuals are inconspicuous, or if

travel costs resulting from increasing the number of sampling units is a strong constraint.

Acknowledgements

435 We would like to warmly thank the 167 volunteers (Appendix 9) who participated in the

436 experiment, including during the preliminary tests. We also thank the volunteers who helped

437 organise the field sessions by finding study sites or participants, and Virginie Pons for her help

438 in transcribing the data.

Authors' contributions

441 J.P., A.B., A.C. and G.P. conceived the study and designed the methodology; J.P. and G.P.

442 organised and conducted the field sessions; J.P. analysed the data and led the writing of the

443 manuscript. All authors contributed critically to the drafts and gave final approval for

444 publication.

Data availability statement

447 All data and code used for this work are available from the GitHub repository:

448 https://github.com/JanPerret/Individual_detection_in_plant_counts

Conflict of Interest statement

All authors declare no conflict of interest.

References

Alexander, H. M., Reed, A. W., Kettle, W. D., Slade, N. A., Roels, S. A. B., Collins, C. D., & Salisbury, V. (2012). Detection and Plant Monitoring Programs: Lessons from an Intensive Survey of Asclepias meadii with Five Observers. *Plos One*, *7*(12), e52762. https://doi.org/10.1371/journal.pone.0052762

Alexander, H. M., Slade, N. A., & Kettle, W. D. (1997). Application of mark-recapture models to estimation of the population size of plants. *Ecology*, *78*(4), 1230-1237. https://doi.org/10.1890/0012-9658(1997)078[1230:AOMRMT]2.0.CO;2

Andrieu, E., Besnard, A., Fréville, H., Vaudey, V., Gauthier, P., Thompson, J. D., & Debussche, M. (2017). Population dynamics of Paeonia officinalis in relation to forest closure: From model predictions to practical conservation management. *Biological Conservation*, *215*, 51-60. https://doi.org/10.1016/j.biocon.2017.08.010

Archaux, F., Gosselin, F., Berges, L., & Chevalier, R. (2006). Effects of sampling time, species richness and observer on the exhaustiveness of plant censuses. *Journal of Vegetation Science*, *17*(3), 299-306. https://doi.org/10.1658/1100-9233(2006)017[0299:EOSTSR]2.0.CO;2

Archaux, F., Henry, P.-Y., & Gimenez, O. (2012). When can we ignore the problem of imperfect detection in comparative studies? *Methods in Ecology and Evolution*, *3*(1), 188-194.

https://doi.org/10.1111/j.2041-210X.2011.00142.x

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., Dai, B.,

Scheipl, F., Grothendieck, G., Green, P., Fox, J., Bauer, A., & Krivitsky, P. N. (2022). *lme4:*

Linear Mixed-Effects Models using "Eigen" and S4 (1.1-29).

Bornand, C. N., Kery, M., Bueche, L., & Fischer, M. (2014). Hide-and-seek in vegetation: Time-

to-detection is an efficient design for estimating detectability and occurrence. *Methods*

in Ecology and Evolution, *5*(5), 433-442. https://doi.org/10.1111/2041-210X.12171

Buckland, S. T., Anderson, D. R., Burnham, K., Laake, J. L., Borchers, D. L., & Thomas, L. (2001).

Introduction to Distance Sampling: Estimating Abundance of Biological Populations.

Oxford University Press.

Buckland, S. T., Marsden, S. J., & Green, R. E. (2008). Estimating bird abundance: Making

methods   work.   *Bird   Conservation   International*,   *18*(S1),   S91-S108.

https://doi.org/10.1017/S0959270908000294

Burnham, K. P., & Anderson, D. R. (2002). *Model Selection and Multimodel Inference: A*

Practical Information Theoretic Approach (2nd ed.). Springer-Verlag.

Carey, C. S., Jones, J. W., Butler, R. S., Kelly, M. J., & Hallerman, E. M. (2019). A Comparison of

Systematic Quadrat and Capture-Mark-Recapture Sampling Designs for Assessing

Freshwater   Mussel   Populations.   *Diversity*,   *11*(8),   127.

https://doi.org/10.3390/d11080127

Chen, G., Kery, M., Plattner, M., Ma, K., & Gardner, B. (2013). Imperfect detection is the rule

rather than the exception in plant distribution studies. *Journal of Ecology*, *101*(1), 183-

191. https://doi.org/10.1111/1365-2745.12021

Chen, G., Kery, M., Zhang, J., & Ma, K. (2009). Factors affecting detection probability in plant

distribution   studies.   *Journal   of   Ecology*,   *97*(6),   1383-1389.

https://doi.org/10.1111/j.1365-2745.2009.01560.x

Dennis, B., Munholland, P. L., & Scott, J. M. (1991). Estimation of Growth and Extinction

Parameters for Endangered Species. *Ecological Monographs*, *61*(2), 115-143.

https://doi.org/10.2307/1943004

Dodd, C. K., & Dorazio, R. M. (2004). Using counts to simultaneously estimate abundance and

detection probabilities in a salamander community. *Herpetologica*, *60*(4), 468-478.

https://doi.org/10.1655/03-60

Elphick, C. S. (2008). How you count counts: The importance of methods research in applied

ecology. *Journal of Applied Ecology*, *45*(5), 1313-1320. https://doi.org/10.1111/j.1365-

2664.2008.01545.x

Ficetola, G. F., Romano, A., Salvidio, S., & Sindaco, R. (2018). Optimizing monitoring schemes

to detect trends in abundance over broad scales. *Animal Conservation*, *21*(3), 221-231.

https://doi.org/10.1111/acv.12356

Garrard, G. E., Bekessy, S. A., McCarthy, M. A., & Wintle, B. A. (2008). When have we looked

hard enough? A novel method for setting minimum survey effort protocols for flora

surveys. *Austral Ecology*, *33*(8), 986-998. https://doi.org/10.1111/j.1442-

9993.2008.01869.x

Garrard, G. E., McCarthy, M. A., Williams, N. S. G., Bekessy, S. A., & Wintle, B. A. (2013). A

general model of detectability using species traits. *Methods in Ecology and Evolution*, *4*(1),

45-52. https://doi.org/10.1111/j.2041-210x.2012.00257.x

Gauthier, P., Pons, V., Fisogni, A., Murru, V., Berjano, R., Dessena, S., Maccioni, A., Chelo, C.,

de Manincor, N., Doncieux, A., Papuga, G., & Thompson, J. D. (2019). Assessing

vulnerability of listed Mediterranean plants based on population monitoring. *Journal for*

Nature Conservation, *52*(125758), 1-11. https://doi.org/10.1016/j.jnc.2019.125758

Gauthier, P., Pons, V., Letourneau, A., Klesczewski, M., Papuga, G., & Thompson, J. D. (2017).

Combining population monitoring with habitat vulnerability to assess conservation status

in populations of rare and endangered plants. *Journal for Nature Conservation*, *37*, 83-95.

https://doi.org/10.1016/j.jnc.2017.03.005

Harper, J. L. (1977). *Population Biology of Plants*. Academic Press.

Hauser, C. E., Giljohann, K. M., McCarthy, M. A., Garrard, G. E., Robinson, A. P., Williams, N. S.

G., & Moore, J. L. (2022). A field experiment characterizing variable detection rates during

plant surveys. *Conservation Biology*, 36(3), e13888. https://doi.org/10.1111/cobi.13888

Ishii, Y., & Shimada, M. (2009). The effect of learning and search images on predator–prey

interactions. *Population Ecology*, *52*(1), 27. https://doi.org/10.1007/s10144-009-0185-x

IUCN Standards and Petitions Committee. (2019). *Guidelines for Using the IUCN Red List*

Categories and Criteria-Version 14.

Kellner, K. F., & Swihart, R. K. (2014). Accounting for Imperfect Detection in Ecology: A

Quantitative Review. *Plos One*, *9*(10), e111436.

https://doi.org/10.1371/journal.pone.0111436

Kendall, W. L., Peterjohn, B. G., & Sauer, J. R. (1996). First-Time Observer Effects in the North

American Breeding Bird Survey. *The Auk*, *113*(4), 823-829.

https://doi.org/10.2307/4088860

Kery, M., Dorazio, R. M., Soldaat, L., van Strien, A., Zuiderwijk, A., & Royle, J. A. (2009). Trend

estimation in populations with imperfect detection. *Journal of Applied Ecology*, *46*(6),

1163-1172. https://doi.org/10.1111/j.1365-2664.2009.01724.x

Kery, M., & Gregg, K. B. (2003). Effects of life-state on detectability in a demographic study of

the terrestrial orchid Cleistes bifaria. *Journal of Ecology*, *91*(2), 265-273.

https://doi.org/10.1046/j.1365-2745.2003.00759.x

Kery, M., Royle, J. A., & Schmid, H. (2005). Modeling avian abundance from replicated counts

using binomial mixture models. *Ecological Applications*, *15*(4), 1450-1461.

https://doi.org/10.1890/04-1120

Kery, M., & Schmidt, B. R. (2008). Imperfect detection and its consequences for monitoring for

conservation. *Community Ecology*, *9*(2), 207-216.

https://doi.org/10.1556/ComEc.9.2008.2.10

Lebreton, J.-D., Burnham, K. P., Clobert, J., & Anderson, D. R. (1992). Modeling Survival and

Testing Biological Hypotheses Using Marked Animals: A Unified Approach with Case

Studies. *Ecological Monographs*, *62*(1), 67-118. https://doi.org/10.2307/2937171

Lüdecke, D., Aust, F., Crawley, S., & Ben-Shachar, M. S. (2022). *ggeffects: Create Tidy Data*

Frames of Marginal Effects for "ggplot" from Model Outputs (1.1.2).

Moore, J. L., Hauser, C. E., Bear, J. L., Williams, N. S. G., & McCarthy, M. A. (2011). Estimating

detection-effort curves for plants using search experiments. *Ecological Applications*,

21(2), 601-607. https://doi.org/10.1890/10-0590.1

Nakagawa, S., Johnson, P. C. D., & Schielzeth, H. (2017). The coefficient of determination R2

and intra-class correlation coefficient from generalized linear mixed-effects models

revisited and expanded. *Journal of The Royal Society Interface*, *14*(134), 20170213.

https://doi.org/10.1098/rsif.2017.0213

Nieder, A. (2020). The Adaptive Value of Numerical Competence. *Trends in Ecology &*

Evolution, *35*(7), 605–617. https://doi.org/10.1016/j.tree.2020.02.009

Petitot, M., Manceau, N., Geniez, P., & Besnard, A. (2014). Optimizing occupancy surveys by

maximizing detection probability: Application to amphibian monitoring in the

Mediterranean region. *Ecology and Evolution*, *4*(18), 3538-3549.

https://doi.org/10.1002/ece3.1207

R Core Team. (2019). *R: A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Reisch, C., Schmid, C., & Hartig, F. (2018). A comparison of methods for estimating plant

population size. *Biodiversity and Conservation*, *27*(8), 2021-2028.

https://doi.org/10.1007/s10531-018-1522-1

Ripple, W. J., & Beschta, R. L. (2012). Trophic cascades in Yellowstone: The first 15 years after

wolf reintroduction. *Biological Conservation*, *145*(1), 205-213.

https://doi.org/10.1016/j.biocon.2011.11.005

Royle, J. A. (2004). N-Mixture Models for Estimating Population Size from Spatially Replicated

Counts. *Biometrics*, *60*(1), 108–115. https://doi.org/10.1111/j.0006-341X.2004.00142.x

Watson, J., & Estes, J. A. (2011). Stability, resilience, and phase shifts in rocky subtidal

communities along the west coast of Vancouver Island, Canada. *Ecological Monographs*,

81(2), 215-239. https://doi.org/10.1890/10-0262.1

Williams, B. K., Nichols, J. D., & Conroy, M. J. (2002). *Analysis and Management of Animal*

Populations: Modeling, Estimation, and Decision Making. Academic Press.

Yoccoz, N. G., Nichols, J. D., & Boulinier, T. (2001). Monitoring of biological diversity in space

and time. *Trends in Ecology & Evolution*, *16*(8), 446-453. https://doi.org/10.1016/S0169-

5347(01)02205-4