Plants stand still but hide: imperfect and heterogeneous

detection is the rule when counting plants

Authors: Jan Perret[1], Aurélien Besnard[1], Anne Charpentier[1], Guillaume Papuga[2]

1. CEFE, Univ Montpellier, CNRS, EPHE-PSL University, IRD, Montpellier, France

2. AMAP, IRD, CIRAD, CNRS, INRAE, Univ Montpellier, Montpellier, France

Number of words: 8,427

Corresponding author: Jan Perret (jan.perret@gmail.com), Centre d'Ecologie Fonctionnelle

et Evolutive, 1919 route de Mende, 34293 Montpellier, France.

Abstract

1. The estimation of population size and its variation across space and time largely relies on counts of individuals, generally carried out within spatial units such as quadrats or sites. Missing individuals during counting (i.e. imperfect detection) results in biased estimates of population size and trends. Imperfect detection has been shown to be the rule in animal studies, and most studies now correct for this bias by estimating detection probability. Yet this correction remains exceptional in plant studies, suggesting that most plant ecologists implicitly assume that all individuals are always detected.

2. To assess if this assumption is valid, we conducted a field experiment to estimate individual detection probability in plant counts conducted in 1x1 m quadrats. We selected 30 herbaceous plant species along a gradient of conspicuousness at 24 sites along a gradient of habitat closure, and asked groups of observers to count individuals in 10 quadrats using three counting methods requiring progressively increasing times to complete (quick count, unlimited count and cell count). In total, 158 participants took part in the experiment, allowing an analysis of the results of 5,024 counts.

3. Over all field sessions, no observer succeeded in detecting all the individuals in the 10 quadrats. The mean detection rate was 0.44 (ranging from 0.11 to 0.82) for the quick count, 0.59 for the unlimited count (range 0.18–0.87) and 0.74 for the cell count (range 0.46–0.94).

4.　Detection probability increased with the conspicuousness of the target species and

decreased with the density of individuals and habitat closure. The observer's experience

in botany had little effect on detection probability, whereas detection was strongly

affected by the time observers spent counting. Yet although the more time-consuming

methods increased detection probability, none achieved perfect detection, nor did they

reduce the effect on detection probability of the variables we measured.

5.　*Synthesis.* Our results show that detection is imperfect and highly heterogeneous when

counting plants. To avoid biased estimates when assessing the size, temporal or spatial

trends of plant populations, plant ecologists should use methods that estimate the

detection probability of individuals rather than relying on raw counts.

Keywords: conservation; demography; detectability; field methods; N-mixture; observer

effect; plant survey; population monitoring.

Introduction

Population size and the variability of population size over time and space are of central importance in ecology and conservation. These parameters are required to understand ecosystem dynamics (Ripple & Beschta, 2012; Watson & Estes, 2011), assess population viability (Dennis et al., 1991) and trigger conservation policies and actions (e.g. IUCN, 2019). Population size estimates largely rely on counts of individuals, generally made in spatial units such as quadrats or sites (Elphick, 2008; Yoccoz et al., 2001). Therefore, the accuracy of these counts is crucial, as counting errors would result in biased estimates of population sizes and trends.

When carrying out counts, observers can miss some individuals, in effect counting fewer individuals than were present in the sampling units. This phenomenon of imperfect detection, well known in animal studies, leads to underestimating population sizes (Williams et al., 2002). If detection is imperfect but constant over time and space, raw counts can provide unbiased estimates of population trends or spatial variation (Yoccoz et al., 2001). However, if detection varies over time (e.g. across the sampling season or between years) or space (e.g. between habitats), the raw differences between counts performed at different sampling sessions or in different sampling units are a mixture of differences in abundance and detection (Kéry et al., 2009). For example, a decrease in population size over the years may be hidden as observers improve their ability to detect individuals over time (Kendall et al., 1996). Or one might conclude that there are fewer individuals in closed than in open habitats because they are harder to detect in closed habitats (Buckland et al., 2008). Even differences as small as 4–8% in detection dramatically increase the risk of erroneously

concluding that population size varies over time or space (Archaux et al., 2012).

Since the 1970s, many studies on vertebrates have shown that imperfect detection is the

rule and that detection probability tends to vary in time and space. Detection probability

depends on a number of factors: the observation process (both the observation method and

the survey effort), the visibility of the target species, and the characteristics of its habitat

(Kéry et al., 2005; Kéry & Schmidt, 2008; Petitot et al., 2014). Consequently, studies including

an estimate of detection probability, which corrects for the bias resulting from imperfect

detection, have become increasingly common for vertebrates and, to a lesser extent,

invertebrates. Yet, they remain exceptional for plants (Kellner & Swihart, 2014), suggesting

that most plant ecologists implicitly assume that all individuals are always detected or that

differences in detection probability are negligible. It is true that, unlike most animals, plants

are immobile, and observers may thus consider it unlikely to miss individuals if careful

sampling is conducted. This assumption led John Harper to make the famous statement

'Plants stand still and wait to be counted', comparing the ease of studying plant and animal

populations (Harper, 1977).

Although detection has almost never been investigated in plants at the individual level, some

studies have investigated it at the species level, assessing species occurrence in a given area

using two main approaches: occupancy or time-to-detection designs. In the first approach,

the survey duration is fixed, and the detection probability of the species when surveying

quadrats or sites is estimated from repeated surveys. In the second approach, the detection

probability is estimated based on the survey time required to detect the species. These

studies found that the detection of plant species was almost always imperfect and that

detection probability varied depending on ecological and observational variables. For

example, species detection varied according to the morphology of the target species (Chen

et al., 2009, 2013; Garrard et al., 2013), with conspicuous growth forms and phenological

stages raising detection probability. A high density of individuals increased the probability of

detecting the species (Bornand et al., 2014), while the closure of the surrounding habitat

decreased it (Alexander et al., 2012; Garrard et al., 2008). Species detection probability also

increased with survey effort (Chen et al., 2009) and often varied between observers

(Archaux et al., 2006; Garrard et al., 2008. Although these studies confirm that detection

issues exist when assessing the occurrence of plant species, do these issues affect individual

detection to a similar extent?

Concerning detection at the individual level, the only studies that have been conducted on

plants used marked individuals in so-called 'capture–recapture' designs (Lebreton et al.,

1992) or the time-to-detection of individuals (Hauser et al., 2022). These studies found that

individual detection was imperfect, although individuals were generally searched for in

relatively small, intensively surveyed plots (Alexander et al., 1997; Andrieu et al., 2017; Kéry

& Gregg, 2003). In most studies using marked individuals, individual detection probability

varied with the phenological stage of individuals and sometimes with habitat closure (Moore

et al., 2011; Andrieu et al., 2017). Hauser et al. (2022) showed that flowering individuals

were detected faster than nonflowering ones, and that the abundance of individuals of

species resembling the target species delayed detection. Observers with recent experience

in searching for the target species detected it faster, confirming at the individual level what

has been demonstrated at the species level. Therefore, it is of particular concern that

individual detection probability has never been estimated for counts of unmarked plants,

despite this being one of the most commonly used methods for estimating population sizes

and trends in plant ecology (Elzinga et al., 1998; Reisch et al., 2018). In addition, many

studies in plant ecology estimate vital rates by counting unmarked individuals per life stage

(e.g. Molano-Flores & Bell, 2012; Giljohann et al., 2017). In theory, data from capture-

recapture designs could be used to estimate detection probability and correct estimates

from counts of unmarked individuals. To our knowledge, this has never been done, probably

because it would be very impractical since both observation processes would have to be

strictly equivalent regarding detection and because estimating detection probability directly

from counts of unmarked plants is usually less labour–intensive (Royle, 2004).

To assess whether imperfect detection of individuals is indeed the rule in plant counts, we

conducted a field experiment in which observers counted plants in 1x1 m quadrats, a

commonly used quadrat size (Elzinga et al., 1998; Gauthier et al., 2017, 2019). The

experiment covered 30 species and 300 quadrats, and involved 158 observers who made

5,024 counts. Our study had three aims: (i) to estimate the detection probability of

individuals in plant counts; (ii) to estimate how detection probability varies in time and

space; (iii) to determine if increasing the survey effort results in a higher detection

probability, and could allow reaching perfect detection. To meet these objectives, we tested

whether detection varies according to ecological variables (species conspicuousness, true

population density and surrounding habitat closure) and observational variables (counting

method and experience of observers), which are likely to vary over time and space, as well

as testing varying degrees of survey effort.

Materials & methods

1 – Selection of sites, species and observers

We selected 30 herbaceous plant species from 24 sites throughout France in Mediterranean, Alpine and temperate climatic regions (Appendix 1). We chose species with various growth forms and phenological stages to cover a gradient from cryptic to conspicuous species. As the experiment focused on counting individuals, we selected only species for which 'individuals' (ramets or genets depending on the species) could be unambiguously differentiated. We avoided species difficult to identify so that all could be identified at first glance without having to check identification criteria on each detected individual. We selected study sites to cover a gradient of habitat closure, from bare ground to dense meadows, to include all possible combinations of species conspicuousness and habitat closure (Fig. 1).

Participants were volunteers selected among our colleagues, researchers, undergraduate students and professionals involved in flora conservation. We selected 4 to 8 observers for each field session, including at least one with several years of field experience in botany, one with no experience in botany, and intermediate profiles. Observers were allowed to participate in up to three sessions, but these sessions had to be several weeks apart. This resulted in 171 participations by 158 unique observers. We conducted the fieldwork from 7 October 2020 to 6 August 2021.

<u>Figure 1:</u> Diagram of the study's experimental design. We conducted 30 field sessions, each focusing on a different species in a habitat with a different closure level, to cover the widest possible range of combinations of species conspicuousness and habitat closure. For each field session, we placed ten quadrats on the population to form a density gradient and chose 4–8 observers to have a gradient of experience in botany. The observers carried out three counting methods successively (quick, unlimited and cell counts). We then counted the true number of individuals by systematically digging out the individuals detected by the observers and then conducting an additional survey for any individuals that remained unnoticed. Picture A shows the experiment in progress at a Mediterranean site and picture B shows a quadrat with the cells.

<u>Field setup and observer training</u>

The fieldwork consisted of one-day sessions, during which a group of observers counted the

number of individuals of a single target species in a habitat at one of the 24 sites. The

availability of participants restricted us to organise the field sessions. Whenever we had

enough observers available somewhere, we prospected the landscape nearby to find an

appropriate site (i.e. easy to access and with a habitat corresponding to what we needed to

complete the habitat closure gradient). Then we searched for a suitable species, meeting our

general criteria and completing the species conspicuousness gradient. Before the observers'

arrival, we placed ten 1x1 m quadrats on the target species population. We chose quadrat

locations to ensure that (i) two were free of individuals of the target species (so that

observers would not always expect to find individuals in the quadrats), and (ii) they covered

a relatively linear gradient of densities up to a maximum of 150 individuals (when possible).

It happened regularly that quadrats meant to be empty contained a few individuals and

those with the highest densities contained more than 150 individuals. We labelled the

quadrats from 1 to 10 and fixed them to the ground. When the observers arrived, we

explained how the counts would be conducted and how to identify the target species at all

phenological stages and differentiate it from the other species present at the study site. We

then trained them to count on additional quadrats to check that everyone had understood

the instructions. This training phase lasted until all observers felt comfortable with species

identification and the counting methods. The entire training process usually took about an

hour.

Counting methods

Each observer had to count the number of individuals of the target plant in each quadrat

using three different methods, requiring a progressively increasing amount of survey time. In

the first round, participants had 30 seconds to estimate the number of individuals in each

quadrat (the quick count). In the second round, they had to count the number of individuals

without any time limit (the unlimited count). We then divided the quadrats into 100 cells of

10x10 cm (Fig. 1B), and observers performed the third round, where they had to count the

number of individuals in each cell without any time limit (the cell count). This last method

forced observers to survey the quadrat homogeneously and has been regularly used in the

plant survey literature (e.g. Gauthier et al., 2017, 2019). For the unlimited and cell counts,

observers recorded the time they spent counting in each quadrat. To avoid memory effects,

after each round we collected the sheets on which the observers recorded their

measurements, and observers surveyed the quadrats in a different order, approaching it

from a different side at each round. The observers conducted the three counts on the ten

quadrats in 1.5 to 5 hours (on average 3.25 hours).

<u>Exhaustive count and measurement of covariates</u>

After the observers had left, we carried out a final count to determine the true number of

individuals present in each quadrat. This was done by searching each 10x10 cm cell for the

maximum number of individuals detected by the observers during the third round and

collecting them. Once the maximum number of individuals detected by the observers in each

cell had been removed, we searched again for any potentially undetected individuals. This

final count took between 1 and 7 hours (3 hours on average), depending on habitat closure

and the number of individuals of the target species in the quadrats. Despite our verification

method, we might have missed a few individuals during the final count, but we will refer to

this as the 'true number of individuals' for the sake of simplicity. We calculated the detection

probability of individuals by dividing the number of individuals detected by the observers by

the true number of individuals present in each quadrat.

We measured three variables to investigate their effect on detection probability: an index of

species conspicuousness (measured for each field session), an index of habitat closure

(measured for each quadrat), and the experience in botany of the observers. For simplicity's

sake, we used synthetic indices to characterise species conspicuousness and habitat closure.

This allowed us to limit the number of variables and interactions in the model (e.g. species

conspicuousness combines the average size of individuals, the presence of vividly coloured

flowers, and the interaction of these variables). All the variables and interactions that we

investigated in the following analysis are presented in Table 1, as well as the underlying

hypotheses. The way we measured the variables is detailed in Appendix 2.

3 – Statistical analysis

In total, 300 quadrats were surveyed during the experiment by on average 5.7 observers,

resulting in 5,024 counts. We removed 34 quadrats from the analysis that contained no

individuals of the target species, as these do not provide information on detection

probability. We withdrew four additional quadrats that unexpectedly contained more than

300 individuals to avoid the leverage effect due to these density outliers (Appendix 3).

Keeping these quadrats in the dataset would have led to a reduction in the slope of the

density effect, which could result from a nonlinear relationship at high densities, but with

only four quadrats, we could not reliably test this hypothesis. In some cases (N = 140,

representing 2.8% of the counts), the observers recorded more individuals than were truly

present (Appendix 4). We thus carried out the analyses on two datasets, one in which we

removed these counts and one in which we set these counts to the true number of

individuals. As there was no discernible difference in the results, we present the results of

the former analysis in the main text and the latter in Appendix 7. After all filtering steps,

4,319 counts remained in the dataset from which we had removed the observations with

excess detections. We also performed the analyses keeping only the experienced botanists

to verify that our results were not driven by an excess of novice observers in our sample.

Despite some minor differences, our overall results remained unchanged (Appendix 8).

We used a generalised linear mixed model (GLMMs) with a binomial distribution and a logit

link function to analyse the correlation between detection probability and the variables we

measured and some of their interactions. There could be dependence between detections

due to individuals forming clusters. However, observers often detected some but not all

individuals within clusters. Furthermore, the aggregation of individuals was not very strong

at the quadrat scale. Hence, the dependence between detections is probably relatively

small, and we consider the binomial model appropriate. We designed the experiment to

disentangle the effects on detection probability of species conspicuousness, habitat closure

and survey effort. We had precise hypotheses about the mechanisms explaining these

effects (detailed in Table 1) that included interactions between variables, and we collected a

large dataset to test all our hypotheses. Therefore, we fitted a single model including all the

variables and interactions corresponding to our hypotheses and used their statistical

significance for inference. To take into account the repeated measurements at each quadrat,

we included random intercepts for the observers, the field sessions and the quadrats (nested

within the field session). As the majority of observers participated in only one field session,

we did not add a random effect interaction between observers and field sessions.

Furthermore, we did not add random effects on the slopes because we had no specific

hypothesis that would justify it. We standardised all the variables prior to model fitting and

assessed the model's goodness-of-fit using the package DHARMa (Hartig, 2022). We

computed the proportion of variance explained by the model and the proportion of variance

explained by the model's fixed effects, using the theoretical variance method of Nakagawa

et al. (2017). Analyses were performed in R version 3.6.1 (R Core Team, 2019), using

packages *lme4* (Bates et al., 2022) to fit the model and *ggeffects* (Lüdecke et al., 2022) to

plot the predictions.

<u>Table 1:</u> Ecological variables (species conspicuousness, habitat closure, true density),

observational variables (counting method, experience in botany, quadrat order, counting

time), and interactions between variables tested for their effect on the detection probability

of individuals, with our hypothesis about the underlying mechanisms. The measurement

method of all variables is detailed in Appendix 2.

Variable	Description	Measurement method	Hypothesis	Type	Observed values
Species conspicuousness	Conspicuousness of the species at the time of the field session	Expert judgement based on the mean height of individuals and the presence of colourful flowers	Tall and colourful species are easier to detect, which increases their detection probability	Ordinal	1, 2, 3, 4
Habitat closure	Closure of the vegetation in the quadrat	Composite variable created from the vegetation height and the proportion of the area covered by vegetation in the quadrats (Appendix 2)	Dense vegetation masks individuals, which lowers detection probability	Continuous	-1.86–3.54
True density	Number of individuals present in the quadrat	Number of individuals counted in the quadrat by the experimenter at the end of the experiment	High density requires greater concentration and makes it difficult to distinguish between individuals, reducing detection probability	Count	0–295
Counting method	Counting method used	Fixed by the experimental design, each observer successively used the three counting methods on the ten quadrats	Counting methods requiring more survey time yield higher detection probability	Categorical	Quick count, unlimited count, cell count
Experience in botany	General experience in botany of the observer (not specific to the target species)	Self-evaluated by the observers before the experiment using a standardised form	Observers with more experience in botany have higher detection probability	Ordinal	1, 2, 3, 4, 5

Quadrat order	Position of the quadrat in the round	Defined by the experimental design, the order in which the observers investigated the quadrats was fixed in advance and different for each observer and round.	Learning increases detection probability over time, or fatigue decreases it	Continuous	1, 2, 3, …, 10
Counting time	Time spent counting by the observer	Each observer timed the duration of counting	Spending more time increases detection probability	Continuous	Quick count: 0.5 min Unlimited count: 0–11 min Cell count: 0–41.58 min
Species conspicuousness x Habitat closure			Detection probability is less affected by habitat closure for conspicuous species	Interaction term	
Species conspicuousness x Counting method			Detection probability obtained with counting methods requiring more survey effort is less affected by species conspicuousness	Interaction term	
Habitat closure x Counting method			Detection probability obtained with counting methods requiring more survey time is less affected by habitat closure	Interaction term	
Experience in botany x Counting method			The difference in detection probability between experts and beginners is larger for the quick count than for the other two methods	Interaction term	

True density x Counting method			Counting methods requiring more survey time make it easier to distinguish between individuals, which decreases the negative effect of true density	Interaction term	
Quadrat order x Counting method			The effect of fatigue or learning on detection probability is stronger for counting methods requiring more survey time	Interaction term	
Species conspicuousness x Counting time x Counting method		No coefficient is estimated for the quick count as the time was fixed	Spending an additional extra minute further increases detection probability for species that are easy to detect	Interaction term	
Habitat closure x Counting time x Counting method		No coefficient is estimated for the quick count as the time was fixed	Spending an additional extra minute further increases detection probability in sparse vegetation	Interaction term	

Results

Over the 30 field sessions, no observer succeeded in detecting all the individuals in all the quadrats. Averaged over all quadrats and observers, the mean detection rate per field session was 0.44 (ranging from 0.11 to 0.82) for the quick count, 0.59 for the unlimited count (from 0.18 to 0.87), and 0.74 for the cell count (from 0.46 to 0.94).

 Coefficients of standardised variables in the model. Dots represent estimated coefficients, and lines represent their 95% confidence intervals. Coefficients with a nonsignificant effect are shown in grey. The reference counting method is the quick count, in which observers had 30 seconds to estimate the number of individuals in the quadrat. The coefficients presented for the other two methods indicate the difference (on the logit scale) in mean detection probability obtained with those methods compared to the quick count. The variable 'counting time' appears twice because we manually coded the interaction between the counting time and the counting method, so no slope parameter is estimated for the quick count.

The goodness-of-fit of the fitted model was very good (Appendix 5). All ecological and
observational variables were significantly correlated with detection probability in at least
some conditions, i.e. for some counting methods or some levels of the other variables (Fig.
2; Appendix 6). Regarding the ecological variables, as expected, the more closed the habitat,
the lower the detection probability (Fig. 3). The more conspicuous the species, the higher
the detection probability, except in entirely open habitats, where the detection probability
was constant and high regardless of the conspicuousness of the species (Fig. 3). Detection
probability decreased steeply with the true density, in a similar way for the three counting
methods (Fig. 4). Regarding the observational variables, spending more time sharply
increased detection probability for the unlimited and cell counts. Habitat closure moderated
this trend, i.e. spending an extra minute counting increased detection probability less in
closed habitats than in open habitats. Experienced observers achieved slightly higher
detection probability than inexperienced observers, and this difference was stronger for the

cell count than for the unlimited count. In contrast, there was no significant correlation

between experience and detection probability for the quick count. For the unlimited and cell

counts, detection probability slightly increased with the position of the quadrat in the round,

while there was no correlation for the quick count (Fig. 4).

Figure 3: Predicted detection probability depending on species conspicuousness (1 = least conspicuous; 4 = most conspicuous), habitat closure (-1.5 = most open habitats; 2.5 = most closed habitats), the counting method, and the time spent counting. Predictions were made for a quadrat containing 100 individuals of the species of interest, located in the fifth position in the round for an observer who rated his level of experience in botany at 2.5 out of 5. The maximum counting time shown for the unlimited count (11 min) and the cell count (39 min) is the maximum time an observer spent counting on a single quadrat during the experiment. Lines represent the predicted mean detection probability and the shading represents 95% confidence intervals.

Observers achieved higher detection probabilities with the cell count than with the

unlimited count, which yielded higher detection probabilities than with the quick count.

However, this came at the cost of a major increase in the time spent counting. The counting

time per quadrat was fixed at 30 seconds for the quick count, and the mean counting time

averaged over all sessions and observers was 1.8 minutes (ranging from 0.3 to 5.7) for the

unlimited count and 8.2 minutes (from 0.9 to 21.6) for the cell count. While the time spent

increased, the slope between detection probability and the other variables (e.g. observer

experience, vegetation closure) was not reduced. Some variables had roughly the same

correlation with detection probability regardless of the counting method (e.g. true density),

while others had an even steeper slope (e.g. experience in botany).

Within the limits of the counting times we observed during the experiment, the model

predicts that reaching a detection probability over 0.95 (i.e. 'almost perfect detection') is

only possible by using the cell count for quadrats with a true density below 100 individuals

and a counting time per quadrat of at least 20 minutes. In a very open habitat (typically with

<7% vegetation cover and median vegetation height <1 cm), this can be achieved no matter

the species' conspicuousness. In a less open habitat (e.g. vegetation cover <90%, median

vegetation height <6 cm), the target species must be very conspicuous, i.e. a plant above 20

cm in height or 10 cm but with colourful flowers.

<u>Figure 4:</u> Predicted detection probability of individuals, depending on the experience in botany of the observer (A), the true density (B) and the position of the quadrat in the round (C). The other variables included in the model are held constant at the following levels: species conspicuousness = 2, habitat closure = 0, quick count time = 0.5 min, unlimited count time = 3 min, cell count time = 3 min. Lines represent the mean detection probability of individuals and the shading represents 95% confidence intervals.

Substantial variation in detection probability between field sessions remained unexplained

by the model, even after controlling for species and habitat characteristics, as the variance

of this random effect was $\sigma_{session}{}^2 = 0.58$. There was also unexplained variation between

quadrats ($\sigma_{quadrat}{}^2 = 0.42$). The unexplained between-observer variation was the smallest,

with a variance of $\sigma_{observer}{}^2 = 0.16$. The proportion of variance explained by the model was

$R^2{}_{GLMM(c)} = 0.35$, and the proportion of variance explained by the model's fixed effects was

$R^2{}_{GLMM(m)} = 0.12$. In other words, detection probability remained heterogeneous even

after considering the effect of the seven variables included in the model and accounting for

unexplained inter-session, inter-plot and inter-observer variance.

320 Discussion

This study provides the first estimate of the detection probability of individuals in plant

counts, using a detection experiment with a cross-taxa and cross-habitat design. Our results

demonstrated that imperfect detection is the rule when counting plants, even in small (1 m²)

quadrats with unlimited counting time. Detection probability was also highly heterogeneous

and varied with the species, habitat, observer and counting method. Methods involving

longer survey time increased the mean detection probability, but did not achieve perfect

detection, nor did they reduce the effect of the ecological and observational variables on

detection probability. Even after controlling for the effects of all the variables, detection

remained highly heterogeneous. Therefore, we argue that plant ecologists should not rely on

the use of raw counts and should always take into account detection issues when studying

plant populations.

1. Evidence for imperfect detection of plants

In our experiment, while using common plant counting methods (Reisch et al., 2019), not a

single observer detected all the individuals present in all the quadrats of a field session. Even

with counts unlimited in time, the predicted detection probability under average conditions

(i.e. for the average values of all variables included in the model with the unlimited count)

did not exceed 0.59 (95% confidence interval: 0.50–0.67). Thus, our results not only confirm

that imperfect detection is the rule when counting plant individuals, but also show that

detection probability can even be relatively low. Although these values may be surprising,

detection issues have been previously documented for plants when searching for a species

(e.g. Chen et al., 2013), and a recent study conducted on large quadrats (20x20 m) suggested

that imperfect detection may also occur for plant individuals (Hauser et al., 2022).

Furthermore, the detection probabilities we found for plants are higher than those

commonly found for animals. For example, detection probabilities of 0.04–0.32 have been

found in bird point counts (Royle, 2004), 0.07–0.41 for lizards counted along transects (Kéry

et al., 2009), 0.06–0.41 for salamanders counted along stream banks (Dodd & Dorazio,

2004), and 0.06–0.34 for freshwater mussels searched on riverbeds (Carey et al., 2019).

Nevertheless, this does not discount the fact that detection remains far from exhaustive for

plants and thus should not be overlooked.

2. Effects of the ecological variables

Our results showed that species conspicuousness positively affects detection probability,

while habitat closure negatively affects it. This general result was expected, as it has also

been observed for plants in occupancy and time-to-detection studies (Alexander et al., 2012;

Garrard et al., 2008; Hauser et al., 2022). In open habitats, detection probability was the

same regardless of the species' conspicuousness. In contrast, differences between species

appeared with increasing habitat closure, reaching up to a 0.42 difference in detection

probability between the extreme levels of species conspicuousness in the most closed

habitats (estimated difference for the unlimited count with 3 min counting time). It shows

that habitat closure drives the detection probability of plant individuals more than species

conspicuousness. Furthermore, since both species conspicuousness and habitat closure

usually vary over time and space, our results show that heterogeneous detection probability

is common and should be considered as the rule rather than the exception. Estimating

temporal or spatial trends using raw counts when detection is heterogeneous is likely to

yield biased estimates and should therefore be avoided by all means.

The negative effect of the density of individuals on detection probability was unexpected in

its magnitude (e.g. a predicted detection probability of 0.67 in a quadrat with 20 individuals

to 0.37 in a quadrat with 300 individuals, using the unlimited count in average conditions)

and occurred regardless of the counting method. Although it would have supported our

results, we did not find this phenomenon reported in the literature, even for animal taxa,

where density-dependent detection probabilities usually result from changes in individual

behaviour in response to density variations (Veech et al., 2016), and not to the observer

detecting fewer individuals. In our case, it may be due to the Weber–Fechner law regarding

how humans perceive numbers (see Nieder, 2020, for a summary). Indeed, studies on

human cognition have shown that it becomes increasingly difficult to estimate the number

of items in an area as the number grows, and that there is a general tendency to

underestimate it. It is plausible that this phenomenon seldom applies to animal counts, as

these do not usually involve counting immobile items in high density, as is the case for

plants.

3. Observer effect

The experience in botany of observers had little effect on detection probability compared to

the ecological variables and the counting time. In average conditions, the difference in

detection probability between novices and the most experienced observers was 0.04 for the

quick count, 0.09 for the unlimited count, and 0.17 for the cell count, and confidence

intervals largely overlapped except for the cell count (Appendix 6). The advantages of having

experience in botany when searching for individuals of a target species may be the ability to

more quickly identify an individual once it has been detected, to make fewer identification

errors, and possibly to survey the study area more efficiently by knowing where to search for

the species (Archaux et al., 2006). These advantages were probably minor in our experiment,

since observers were trained to identify the target species before starting counts, all species

were easy to identify, and an efficient surveying strategy was of little use in 1x1 m quadrats.

In this context, the observers' task can be seen as searching for small objects in a noisy

environment, and it seems plausible that their performance was mainly driven by their

ability to remain focused during a repetitive task, their motivation, or their fatigue during

the experiment. Larger differences between observers might have been observed in larger

quadrats, as the surveying strategy might then be important. Overall, it is likely that the

variables we studied have a similar effect on the detection probability regardless of quadrat

size. In larger quadrats, the observation process has an additional layer of complexity as

observers have to walk in the quadrat and might follow different surveying strategies.

Furthermore, the differences between observers might have been greater if we had involved

participants with recent experience in searching for the target species (Hauser et al., 2022)

due to the well-known 'search image' effect, whereby foraging predators improve their

probability of detecting the type of prey they are trained to detect (Ishii & Shimada, 2009). If

this effect indeed applies to plant counts, population trends estimated from raw counts will

be biased if the same observers always make the counts, as their detection probability will

increase with time.

410   4. Comparison of the counting methods

As expected, increasing the survey effort, measured by the time spent counting, massively

increased detection probability (Archaux et al., 2006; Hauser et al., 2022). In average

conditions, the estimated detection probability was 0.36 for the quick count (30 sec counting

time), 0.54 for the unlimited count (1.8 min average counting time), and 0.73 for the cell

count (8.2 min). In addition, for the unlimited and cell counts, observers spending more time

counting achieved a higher detection probability (e.g. for the unlimited count in average

conditions, detection probability increased from 0.50 at 1 min to 0.83 at 10 min).

The gain in detection associated with counting individuals per cell instead of the whole

quadrat could be due to two reasons: the cells force the observer to survey the whole

surface area of the quadrat uniformly, and might also reduce uncertainty whether or not an

individual has already been counted, which frequently occurs when individuals are close to

each other. However, contrary to our expectations, increasing survey effort did not reduce

the effect of the ecological variables on detection probability, the most striking being the

density of individuals, which had an almost identical effect on detection probability for all

three counting methods.

5. Unexplained variance and implications of detection probability heterogeneity

Of the random effects included in the model, the most unexplained variation in detection

probability was at the inter-session and inter-quadrat levels. Since we only measured three

synthetic variables to characterise the differences between sessions and quadrats (species

conspicuousness, habitat closure and true density), we expected that unexplained variance

would remain at these levels. For example, we did not measure colour and height

heterogeneity, for the habitat or for the target species, although these variables certainly

impact detection probability (Garrard et al., 2013; Hauser et al., 2022).

The fact that the unexplained inter-observer variation was the lowest of the random effects

is an encouraging result for future studies, as it means that there are few differences in

detection probability between observers compared to the differences between species or

over a habitat closure gradient. Yet, overall, more than half (65%) of the variance in

detection probability remained unexplained by our model. Thus, observers had a

heterogeneous detection probability between quadrats even after controlling for several

variables, suggesting that factors that vary over time for each observer, such as

concentration or motivation, probably explain a large part of the variation in detection

probability.

6. Recommendations for counting plants

Our study shows for plants what has been documented many times for animals: that

detection probability of individuals is less than one and is usually heterogeneous in space

and time, regardless of the counting method and survey effort. Using counts affected by

imperfect detection as a proxy for the density of individuals to make comparisons over time

and space is only valid under the hypothesis of constant detection. Our findings show that

multiple ecological and observational variables cause heterogeneity in detection probability.

While observational variables can be standardised, it seems unlikely that ecological variables

(i.e. species conspicuousness, habitat closure and true density of individuals) will be constant

in any study, even those conducted at limited temporal and spatial scales. Furthermore,

trying to achieve almost perfect detection (i.e. >0.95) by increasing survey efforts to fix the

issue of detection heterogeneity seems unrealistic given the resource constraints of most

ecology and conservation studies. As we studied a wide variety of species and habitats, we

trust that our results are transferable to most biomes and species types, provided the

scientific context is comparable. The spatial scale must be equivalent in terms of quadrat size

(i.e. small enough that it is not necessary to walk inside during the survey) and the studied

species must be of a similar life form (e.g. herbaceous species or tree seedlings). We

therefore argue that temporal or spatial trends in plant populations should not be estimated

using raw counts, as they could then reflect a trend in ecological variables rather than the

population's fate.

The detection probability depends on the conditions of the observation process and the

studied system; thus, it has to be estimated within each study and cannot be extrapolated

from our results. Several methods can be used to estimate individual detection probability,

and thus estimate unbiased population sizes or trends (e.g. Lebreton et al., 1992; Buckland

et al., 2001). In particular, the N-mixture method uses repeated counts of unmarked

individuals on sampling units (e.g. quadrats or sites) to estimate detection probability from

the differences in the number of individuals detected between counts (Royle, 2004). With

this method, estimates of population sizes or trends are more precise if detection probability

is high and homogeneous in time and space, if the mean density of individuals is high, and if

the sample size, i.e. the number of sampling units and count repetitions, is high (Royle, 2004;

Ficetola et al., 2018). Thus, for studies using the N-mixture method, no matter the studied

population and the counting method used, we recommend setting a fixed counting time as

leaving it to the observer's choice will add heterogeneity to detection probability, resulting

in less precise estimates.

In terms of allocating survey effort, there are three options: use a more time-consuming

counting method to increase detection probability, perform more count repetitions, or

survey more sampling units. The only way to be sure of the optimal design is to conduct a

pilot study to estimate the mean density of individuals and their detection probability in the

population of interest. However, Ficetola et al. (2018) have shown that when the detection

probability is above 0.10, the most advantageous strategy is usually to make three count

repetitions and to increase the number of sampling units as much as possible. Conversely,

when the detection probability is below 0.10, it is better to increase it by using a more time-

consuming counting method or to increase the number of count repetitions. Thus, we

recommend counting individuals in quadrats without cells with a fixed short counting time

(e.g. 0.5–1 min), as this method will yield a detection probability above 0.10 for most species

and allow the most sampling units to be surveyed. More time-consuming counting methods

should be used only if the study population grows in a very closed habitat and the individuals

are inconspicuous, or if travel costs resulting from increasing the number of sampling units is

a strong constraint.

Acknowledgements

496 We would like to warmly thank the 167 volunteers (Appendix 9) who participated in the

497 experiment, including during the preliminary tests. We are grateful to the volunteers who

498 helped organise the field sessions by finding study sites or participants, Virginie Pons for her

499 help in transcribing the data, and Solal Boutoux for his help in designing Fig. 1.

Authors' contributions

502 J.P., A.B., A.C. and G.P. conceived the study and designed the methodology; J.P. and G.P.

503 organised and conducted the field sessions; J.P. analysed the data and led the writing of the

504 manuscript. All authors critically contributed to the drafts and gave their final approval for

505 publication.

Data availability statement

508 All data and code used for this work are available from the GitHub repository:

509 https://github.com/JanPerret/Individual_detection_in_plant_counts

Conflict of Interest statement

All authors declare no conflict of interest.

References

Alexander, H. M., Reed, A. W., Kettle, W. D., Slade, N. A., Roels, S. A. B., Collins, C. D., & Salisbury, V. (2012). Detection and Plant Monitoring Programs: Lessons from an Intensive Survey of Asclepias meadii with Five Observers. *Plos One*, *7*(12), e52762. https://doi.org/10.1371/journal.pone.0052762

Alexander, H. M., Slade, N. A., & Kettle, W. D. (1997). Application of mark-recapture models to estimation of the population size of plants. *Ecology*, *78*(4), 1230-1237. https://doi.org/10.1890/0012-9658(1997)078[1230:AOMRMT]2.0.CO;2

Andrieu, E., Besnard, A., Fréville, H., Vaudey, V., Gauthier, P., Thompson, J. D., & Debussche, M. (2017). Population dynamics of Paeonia officinalis in relation to forest closure: From model predictions to practical conservation management. *Biological Conservation*, *215*, 51-60. https://doi.org/10.1016/j.biocon.2017.08.010

Archaux, F., Gosselin, F., Berges, L., & Chevalier, R. (2006). Effects of sampling time, species richness and observer on the exhaustiveness of plant censuses. *Journal of Vegetation Science*, *17*(3), 299-306. https://doi.org/10.1658/1100-9233(2006)017[0299:EOSTSR]2.0.CO;2

Archaux, F., Henry, P.-Y., & Gimenez, O. (2012). When can we ignore the problem of imperfect detection in comparative studies? *Methods in Ecology and Evolution*, *3*(1),

188-194. https://doi.org/10.1111/j.2041-210X.2011.00142.x

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., Dai, B.,

Scheipl, F., Grothendieck, G., Green, P., Fox, J., Bauer, A., & Krivitsky, P. N. (2022). *lme4:*

Linear Mixed-Effects Models using "Eigen" and S4 (1.1-29).

Bornand, C. N., Kery, M., Bueche, L., & Fischer, M. (2014). Hide-and-seek in vegetation: Time-

to-detection is an efficient design for estimating detectability and occurrence. *Methods*

in Ecology and Evolution, *5*(5), 433-442. https://doi.org/10.1111/2041-210X.12171

Buckland, S. T., Anderson, D. R., Burnham, K., Laake, J. L., Borchers, D. L., & Thomas, L.

(2001). *Introduction to Distance Sampling: Estimating Abundance of Biological*

Populations. Oxford University Press.

Buckland, S. T., Marsden, S. J., & Green, R. E. (2008). Estimating bird abundance: Making

methods work. *Bird Conservation International*, *18*(S1), S91-S108.

https://doi.org/10.1017/S0959270908000294

Carey, C. S., Jones, J. W., Butler, R. S., Kelly, M. J., & Hallerman, E. M. (2019). A Comparison

of Systematic Quadrat and Capture-Mark-Recapture Sampling Designs for Assessing

Freshwater Mussel Populations. *Diversity*, *11*(8), 127.

https://doi.org/10.3390/d11080127

Chen, G., Kery, M., Plattner, M., Ma, K., & Gardner, B. (2013). Imperfect detection is the rule

rather than the exception in plant distribution studies. *Journal of Ecology*, *101*(1), 183-

191. https://doi.org/10.1111/1365-2745.12021

Chen, G., Kery, M., Zhang, J., & Ma, K. (2009). Factors affecting detection probability in plant

distribution studies. *Journal of Ecology*, *97*(6), 1383-1389.

https://doi.org/10.1111/j.1365-2745.2009.01560.x

Dennis, B., Munholland, P. L., & Scott, J. M. (1991). Estimation of Growth and Extinction

556  Parameters for Endangered Species. *Ecological Monographs*, *61*(2), 115-143.

557  https://doi.org/10.2307/1943004

558 Dodd, C. K., & Dorazio, R. M. (2004). Using counts to simultaneously estimate abundance

559  and detection probabilities in a salamander community. *Herpetologica*, *60*(4), 468-478.

560  https://doi.org/10.1655/03-60

561 Elphick, C. S. (2008). How you count counts: The importance of methods research in applied

562  ecology. *Journal of Applied Ecology*, *45*(5), 1313-1320. https://doi.org/10.1111/j.1365-

563  2664.2008.01545.x

564 Elzinga, C. L., & Salzer, D. W. (1998). Measuring & Monitoring Plant Populations. U.S.

565  Department of the Interior, Bureau of Land Management.

566 Ficetola, G. F., Romano, A., Salvidio, S., & Sindaco, R. (2018). Optimizing monitoring schemes

567  to detect trends in abundance over broad scales. *Animal Conservation*, *21*(3), 221-231.

568  https://doi.org/10.1111/acv.12356

569 Garrard, G. E., Bekessy, S. A., McCarthy, M. A., & Wintle, B. A. (2008). When have we looked

570  hard enough? A novel method for setting minimum survey effort protocols for flora

571  surveys. *Austral Ecology*, *33*(8), 986-998. https://doi.org/10.1111/j.1442-

572  9993.2008.01869.x

573 Garrard, G. E., McCarthy, M. A., Williams, N. S. G., Bekessy, S. A., & Wintle, B. A. (2013). A

574  general model of detectability using species traits. *Methods in Ecology and Evolution*,

575  *4*(1), 45-52. https://doi.org/10.1111/j.2041-210x.2012.00257.x

576 Gauthier, P., Pons, V., Fisogni, A., Murru, V., Berjano, R., Dessena, S., Maccioni, A., Chelo, C.,

577  de Manincor, N., Doncieux, A., Papuga, G., & Thompson, J. D. (2019). Assessing

578  vulnerability of listed Mediterranean plants based on population monitoring. *Journal for*

579  *Nature Conservation*, *52*(125758), 1-11. https://doi.org/10.1016/j.jnc.2019.125758

Gauthier, P., Pons, V., Letourneau, A., Klesczewski, M., Papuga, G., & Thompson, J. D. (2017).

Combining population monitoring with habitat vulnerability to assess conservation

status in populations of rare and endangered plants. *Journal for Nature Conservation*,

37, 83-95. https://doi.org/10.1016/j.jnc.2017.03.005

Giljohann, K. M., McCarthy, M. A., Keith, D. A., Kelly, L. T., Tozer, M. G., & Regan, T. J. (2017).

Interactions between rainfall, fire and herbivory drive resprouter vital rates in a semi-

arid ecosystem. *Journal of Ecology*, *105*(6), 1562–1570. https://doi.org/10.1111/1365-

2745.12768

Harper, J. L. (1977). *Population Biology of Plants*. Academic Press.

Hauser, C. E., Giljohann, K. M., McCarthy, M. A., Garrard, G. E., Robinson, A. P., Williams, N.

S. G., & Moore, J. L. (2022). A field experiment characterizing variable detection rates

during plant surveys. *Conservation Biology*, 36(3), e13888.

https://doi.org/10.1111/cobi.13888

Ishii, Y., & Shimada, M. (2009). The effect of learning and search images on predator–prey

interactions. *Population Ecology*, *52*(1), 27. https://doi.org/10.1007/s10144-009-0185-x

IUCN Standards and Petitions Committee. (2019). *Guidelines for Using the IUCN Red List*

Categories and Criteria-Version 14.

Kellner, K. F., & Swihart, R. K. (2014). Accounting for Imperfect Detection in Ecology: A

Quantitative Review. *Plos One*, *9*(10), e111436.

https://doi.org/10.1371/journal.pone.0111436

Kendall, W. L., Peterjohn, B. G., & Sauer, J. R. (1996). First-Time Observer Effects in the North

American Breeding Bird Survey. *The Auk*, *113*(4), 823-829.

https://doi.org/10.2307/4088860

Kery, M., Dorazio, R. M., Soldaat, L., van Strien, A., Zuiderwijk, A., & Royle, J. A. (2009). Trend

estimation in populations with imperfect detection. *Journal of Applied Ecology*, *46*(6),

1163-1172. https://doi.org/10.1111/j.1365-2664.2009.01724.x

Kery, M., & Gregg, K. B. (2003). Effects of life-state on detectability in a demographic study

of the terrestrial orchid Cleistes bifaria. *Journal of Ecology*, *91*(2), 265-273.

https://doi.org/10.1046/j.1365-2745.2003.00759.x

Kery, M., Royle, J. A., & Schmid, H. (2005). Modeling avian abundance from replicated counts

using binomial mixture models. *Ecological Applications*, *15*(4), 1450-1461.

https://doi.org/10.1890/04-1120

Kery, M., & Schmidt, B. R. (2008). Imperfect detection and its consequences for monitoring

for conservation. *Community Ecology*, *9*(2), 207-216.

https://doi.org/10.1556/ComEc.9.2008.2.10

Lebreton, J.-D., Burnham, K. P., Clobert, J., & Anderson, D. R. (1992). Modeling Survival and

Testing Biological Hypotheses Using Marked Animals: A Unified Approach with Case

Studies. *Ecological Monographs*, *62*(1), 67-118. https://doi.org/10.2307/2937171

Lüdecke, D., Aust, F., Crawley, S., & Ben-Shachar, M. S. (2022). *ggeffects: Create Tidy Data*

Frames of Marginal Effects for "ggplot" from Model Outputs (1.1.2).

Molano-Flores, B., & Bell, T. J. (2012). Projected population dynamics for a federally

endangered plant under different climate change emission scenarios. *Biological*

Conservation, *145*(1), 130–138. https://doi.org/10.1016/j.biocon.2011.10.022

Moore, J. L., Hauser, C. E., Bear, J. L., Williams, N. S. G., & McCarthy, M. A. (2011). Estimating

detection-effort curves for plants using search experiments. *Ecological Applications*,

21(2), 601-607. https://doi.org/10.1890/10-0590.1

Nakagawa, S., Johnson, P. C. D., & Schielzeth, H. (2017). The coefficient of determination R2

and intra-class correlation coefficient from generalized linear mixed-effects models

revisited and expanded. *Journal of The Royal Society Interface*, *14*(134), 20170213.

https://doi.org/10.1098/rsif.2017.0213

Nieder, A. (2020). The Adaptive Value of Numerical Competence. *Trends in Ecology &*

Evolution, *35*(7), 605–617. https://doi.org/10.1016/j.tree.2020.02.009

Petitot, M., Manceau, N., Geniez, P., & Besnard, A. (2014). Optimizing occupancy surveys by

maximizing detection probability: Application to amphibian monitoring in the

Mediterranean region. *Ecology and Evolution*, *4*(18), 3538-3549.

https://doi.org/10.1002/ece3.1207

R Core Team. (2019). *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing. https://www.R-project.org/

Reisch, C., Schmid, C., & Hartig, F. (2018). A comparison of methods for estimating plant

population size. *Biodiversity and Conservation*, *27*(8), 2021-2028.

https://doi.org/10.1007/s10531-018-1522-1

Ripple, W. J., & Beschta, R. L. (2012). Trophic cascades in Yellowstone: The first 15 years after

wolf reintroduction. *Biological Conservation*, *145*(1), 205-213.

https://doi.org/10.1016/j.biocon.2011.11.005

Royle, J. A. (2004). N-Mixture Models for Estimating Population Size from Spatially

Replicated Counts. *Biometrics*, *60*(1), 108–115. https://doi.org/10.1111/j.0006-

341X.2004.00142.x

Veech, J. A., Ott, J. R., & Troy, J. R. (2016). Intrinsic heterogeneity in detection probability

and its effect on N-mixture models. *Methods in Ecology and Evolution*, 7(9), 1019–1028.

https://doi.org/10.1111/2041-210X.12566

Watson, J., & Estes, J. A. (2011). Stability, resilience, and phase shifts in rocky subtidal

communities along the west coast of Vancouver Island, Canada. *Ecological Monographs*,

81(2), 215-239. https://doi.org/10.1890/10-0262.1

Williams, B. K., Nichols, J. D., & Conroy, M. J. (2002). *Analysis and Management of Animal*

Populations: Modeling, Estimation, and Decision Making. Academic Press.

Yoccoz, N. G., Nichols, J. D., & Boulinier, T. (2001). Monitoring of biological diversity in space

and time. *Trends in Ecology & Evolution*, *16*(8), 446-453. https://doi.org/10.1016/S0169-

5347(01)02205-4

658