NUDT6, the Antisense Protein of FGF2 Gene, Plays a Depressogenic Role by Promoting Inflammation and Suppressing Neurogenesis without Altering FGF2 Signaling

Burak Uzay[1,2], Fatma Özlem Hökelekli[1,3,a], Murat Yılmaz[1,4,a], Emre Cem Esen[1], Koray Başar[5], Aslıhan Bahadır-Varol[1], Yavuz Ayhan[5], Turgay Dalkara[1], Emine Eren-Koçak[1,5,*]

1. Institute of Neurological Sciences and Psychiatry, Hacettepe University, Ankara, Turkey
2. Current address: Brain Institute, Department of Pharmacology, Vanderbilt University, Nashville, 37240-7933,TN, USA
3. Current address: Department of Psychiatry, University of Texas Southwestern, Dallas, TX, USA
4. Current address: Department of Biology, Faculty of Science, Hacettepe University, Ankara, Turkey
5. Department of Psychiatry, Faculty of Medicine, Hacettepe University, Ankara, Turkey

a These authors contributed equally to this work
*Corresponding author

*Corresponding author: Emine Eren-Kocak
Hacettepe Universitesi,
Norolojik Bilimler ve Psikiyatri Enstitusu,
 Tip Fakultesi Binasi, Asma kat,
Sihhiye, 06100 / Ankara/ Turkey

Phone number: +90 312 305 2130

Email: erenemi@hacettepe.edu.tr

Keywords: NUDT6, FGF2, FGF-AS, natural antisense transcripts, depression, S100A9, NF-κB, neurogenesis

1 **Abstract**

2 Fibroblast growth factor-2 (FGF2) is involved in the regulation of affective behavior and shows

3 antidepressant effects through Akt and ERK1/2 pathways. NUDT6 is a protein encoded from

4 the antisense strand of the FGF2 gene and its role in the regulation of affective behavior is

5 unclear. Here, we show that increasing NUDT6 expression in the hippocampus results in

6 depression-like behavior in rats without changing FGF2 levels or activating its downstream

7 effectors, Akt and ERK1/2. Instead, NUDT6 acts by inducing inflammatory signaling,

8 specifically by increasing S100A9 levels, activating NF-κB and rising microglia number along

9 with a reduction in neurogenesis. Conversely, inhibition of hippocampal NUDT6 expression by

10 shRNA results in antidepressant effects and increases neurogenesis without altering FGF2

11 levels. Together these findings suggest that NUDT6 may play a role in major depression by

12 inducing a proinflammatory state and serve as a novel therapeutic target for antidepressant

13 development. The opposite effects of NUDT6 and FGF2 on depression-like behavior may

14 serve as a mechanism to fine-tune affective behavior. Our findings open up new venues for

15 studying the differential regulation and functional interactions of sense and antisense proteins

16 in neural function and behavior as well as in neuropsychiatric disorders.

Introduction

Major depressive disorder (MDD) is a debilitating psychiatric disorder affecting 4.7% of the world population (1). Despite several treatment options, treatment resistance affecting approximately one third of patients is a major concern (2,3). Current treatment strategies for major depression lack specificity in targeting particular pathophysiological mechanisms. Therefore, better understanding of the pathophysiological mechanisms underlying MDD is necessary to develop new antidepressant interventions.

One of the leading hypotheses of major depression is the "neurotrophic hypothesis". It stems from observations of reduced growth factor levels in several brain regions of chronically stressed animals and human depressed subjects, which are reversed by antidepressant treatments (reviewed in (4,5)). In addition, increasing brain derived neurotrophic factor (BDNF) and fibroblast growth factor-2 (FGF2) levels in brain has antidepressant and anxiolytic effects in rodent models (6–9). Both FGF2 and BDNF have natural antisense transcripts (NAT), RNA molecules that are transcribed from the opposite strand of protein-coding genes, and are suggested to regulate the transcription of their sense strand (10). Supporting this, BDNF-antisense (BDNF-AS) is reported to be a discordant regulator of BDNF protein expression i.e. when BDNF-AS levels increase, BDNF protein expression decreases (11). A similar role was suggested for FGF-AS in the regulation of FGF2 expression based on initial *in vitro* studies (12–15), however several other *in vitro* studies failed to detect an interaction between FGF-AS and FGF2 expression levels (13,15–17). FGF-AS is translated into a 35 kDa protein, nudix hydrolase 6 (NUDT6), the function of which is unknown (16). Intriguingly, hippocampal NUDT6 expression was altered in response to acute stress in rats, suggesting a role for NUDT6 in the regulation of affective behavior (18,19). Although this observation points to an additional layer of complexity in the regulation of the affective behavior by neurotrophic factors, the place of NUDT6 in MDD pathophysiology has not yet been elucidated. Understanding the potential role of NUDT6 in affective behavior may therefore provide a deeper insight to mechanisms of MDD and disclose novel antidepressant therapeutic targets (6,9,20).

Here, we show that overexpression of hippocampal NUDT6 increases anxiety- and depression-like behaviors in rats. These effects were independent of FGF2 signaling, as we did not find any changes in hippocampal FGF2 expression or its downstream signaling mediated by Akt and ERK1/2. Rather, hippocampal NUDT6 overexpression increased 1) the expression of proinflammatory S100A9, an alarmin / damage associated molecular pattern (DAMP), 2) nuclear translocation (thus activation) of NF-κB-p65, a proinflammatory transcription factor and 3) number of microglia, all conforming to an inflammatory state. Moreover, NUDT6 overexpression reduced neurogenesis, possibly as a consequence of the proinflammatory changes induced. Confirming the role of NUDT6 in the pathophysiology of

depression, we found that its knockdown in the hippocampus not only had antidepressant and
anxiolytic effects, but also increased neurogenesis. Overall, our results suggest that NUDT6
can promote inflammatory signaling in the hippocampus, which may take part in decreased
neurogenesis and depression-like behavior observed on its overexpression and, reversed by
its knockdown. Our study discloses a novel mechanism in the pathophysiology of major
depression and, for the first time, underscores the functional effects of antisense proteins in
neuropsychiatric diseases. The fact that NUDT6 regulates affective behavior through a
different mechanism than its sense transcript, FGF2, whose antidepressant effects is well
known, may open up new venues for studying the functional interactions of sense and
antisense proteins not only in major depression but also in other CNS disorders.

Results

1. Hippocampal overexpression of NUDT6 induces depression- and anxiety-like behaviors

To determine whether hippocampal NUDT6 is involved in the regulation of affective behavior, first we overexpressed NUDT6 in the hippocampus by two means: 1. Repeated daily intracerebroventricular (ICV) injections of an NUDT6-expressing plasmid (NUDT6-EP) for 14 days to avoid hippocampal tissue injury and growth factor response that can result from site specific microinjection (21), 2. Bilateral intrahippocampal microinjection of NUDT6-expressing adeno-associated virus (NUDT6-AAV2) to ensure hippocampus-specific overexpression (Figure 1A). At the end of the ICV injections and one month following the microinjection of NUDT6-AAV2, we performed behavioral tests to assess depression-like behavior by sucrose preference test (SPT) and forced swim test (FST), and anxiety-like behavior by elevated plus maze test (EPM) (Figure 1A). Both repeated ICV injections of NUDT6-EP and bilateral intrahippocampal injections of NUDT6-AAV2 increased immobility time and decreased swimming time in FST, in accordance with an increase in depression-like behavior (p= 0.067 and 0.050 for ICV injections, and 0.005 and 0.003 for AAV2 injections, respectively) (Figure 1B and 1C). There was, however, no change in SPT in the NUDT6-EP or NUDT6-AAV2 groups compared to their corresponding controls treated with vehicle or Blank-AAV (p=0.501 and 0.617, respectively) (Figure S1A and S1B).

NUDT6-EP group spent less time in open arms in EPM than the vehicle-injected control group, indicating an increase in anxiety-like behavior (p=0.032) (Figure 1D). Time spent in open arms in EPM was similar in the NUDT6-AAV2 and Blank-AAV2 injected groups (p=0.89)(Figure 1E). The total distance moved in EPM was also similar between experimental and control groups in both sets of experiments (p=0.274 and 0.451, respectively) (Figure S1C and S1D).

2. Altering NUDT6 expression neither changes FGF2 levels nor interferes with FGF2 signaling pathways in the hippocampus.

We investigated whether the effects of NUDT6 on depression- and anxiety-like behavior were mediated by a change in the expression of its sense transcript, FGF2. After confirming the increase in NUDT6 protein levels in the hippocampus in both groups by immunoblotting (p=0.009 and p=0.04 in ICV NUDT6-EP, intrahippocampal NUDT6-AAV2 injection groups, respectively), we examined the corresponding hippocampal FGF2 protein levels (Figures 1F and 1G). On contrary to *in vitro* studies reporting transcriptional regulation of FGF2 by FGF-AS (12,13,15), we found no change in FGF2 protein levels by increasing

NUDT6 expression (p=0.80 and p=0.20 in ICV NUDT6-EP and intrahippocampal NUDT6-
AAV2 injection groups, respectively) (Figures 1F and 1G). FGF2 mRNA expression was also
not changed after intrahippocampal NUDT6-AAV injections (p=0.59), ruling out a possible
regulatory effect of NUDT6 on FGF2 transcription (Figure S1F).

The above findings strongly suggest that the depressogenic effect of hippocampal
NUDT6 was not mediated by a change in FGF2 levels. To rule out possible interactions at
downstream mediator levels, we studied the phosphorylation of ERK1/2 and Akt in NUDT6-
AAV2/Blank AAV2 microinjected hippocampi, molecules mediating the antidepressant actions
of FGF2 (22,23) (Figure 2A). We did not find any difference in the phosphorylation of either
Akt or ERK1/2 between the Blank-AAV2 control and NUDT6-AAV2 groups (p>0.99 and p=0.80
respectively) (Figure 2B and 2 C).

3. Increasing NUDT6 expression induces inflammatory signaling

To have an insight into the molecular mechanisms of action of NUDT6, we performed
proteomic analysis of hippocampal samples from NUDT6-AAV2 and Blank-AAV2 groups
(Figure 2D), which revealed a significant upregulation of 31 proteins and downregulation of 3
proteins in the NUDT6-AAV2 group compared to controls (p<0.05) (Figure 2E) (Supplemental
Table 1). When these 34 proteins were subjected to in silico pathway analysis using online
Panther and String softwares, we found that they are involved in neuroinflammation, cellular
growth and proliferation, organismal injury and abnormalities and mitochondrial dysfunction
(Figure 2F). FGF2 was not among the 34 proteins that were changed by hippocampal NUDT6
overexpression in line with the above immunoblotting and RT-PCR findings. However, there
was a 32-fold increase in the expression of S100 calcium binding protein A9 (S100A9), a
proinflammatory molecule mainly expressed by myeloid cells (i.e. microglia in brain),
supporting the role of inflammatory signaling in the development of NUDT6-induced
depression-like behavior (24,25) (Figure 2E).

Western blotting and RT-PCR of hippocampal samples confirmed the increase in
S100A9 mRNA and protein levels in the NUDT6–AAV group compared to Blank-AAV control
group (p=0.04 and p=0.01, respectively) (Figure 3A-C). We double-immunostained the
hippocampal sections from NUDT6-AAV and Blank-AAV groups with either anti-S100A9 &
anti-NeuN antibodies or with anti-S100A9 & anti-Iba1 antibodies to label neuronal or microglial
S100A9, respectively. We found that S100A9 was colocalized to both neurons and microglia
and its immunopositivity in both dentate granular and hilar neurons was significantly increased
in NUDT6-AAV group relative to the Blank-AAV control group (p<0.0001) (Figure 3D-E and
Figure S2A). To test whether the increase in S100A9 occurred in the same cells in which
NUDT6 was overexpressed, we double-immunostained hippocampal sections with NUDT6

&S100A9. There was a significant increase in S100A9 in NUDT6-positive cells in the NUDT6-
AAV2 group compared to Blank-AAV2 group, suggesting that the increase in S100A9 took
place downstream to NUDT6 in the same cell (Figure 3F-G).
As S100A8/A9 reportedly results in microglial cell migration *in vitro* (23), we assessed
whether NUDT6 overexpression altered the number of Iba-1 immunoreactive cells (microglia)
in hippocampus (26). The number of Iba-1 labeled microglia was increased by %87 in hilar
region of the hippocampus in NUDT6-AAV2 group compared to Blank-AAV2 group, which
indicates microglial recruitment in response to NUDT6 overexpression and further supports
the involvement of inflammatory signaling in the NUDT6's depressogenic actions (p=0.01)
(Figure 3H-I).
Previous studies report that S100A9 is associated with the activation of
proinflammatory transcription factor NF-κB (27,28). When NF-κB binding protein IκB is
phosphorylated, the p65 subunit of NF-κB is freed and translocated to the nucleus, where it
initiates transcription of several inflammatory molecules, including cytokines and chemokines
(29). Thus, nuclear translocation of NF-κB-p65 is widely used as an indicator of increased
neuroinflammatory signaling (30,31). We investigated whether NUDT6 overexpression
induced activation of NF-κB-p65 in our samples by quantifying the ratio of NF-κB-p65 positive
nuclei-to-total nuclei in immunostained hippocampal sections from NUDT6-AAV2 and blank-
AAV2 groups. We found a 2.4-fold increase in NF-κB nuclear translocation in the dentate gyrus
of NUDT6-AAV group compared to Blank-AAV injected controls (p<0.0001), indicating
activation of NF-κB by hippocampal NUDT6 overexpression (Figure 3J-K). Double labeling of
NF-κB-p65 with NUDT6 or S100A9 revealed an increased NF-κB-p65 nuclear translocation in
both NUDT6- and S100A9-positive cells in the NUDT6-AAV2 group compared to Blank-AAV2
group, respectively (p=0.0021 for hilus and p=0.031 for dentate gyrus, p<0.0001 for hilus and
p=0.0003 for dentate gyrus, respectively) (Figure 3L-P).
97

98 4. Overexpression of NUDT6 reduces hippocampal neurogenesis
Due to the putative role of neurogenesis in the regulation of affective behavior (32–34) and
involvement of NF-κB in the regulation of in hippocampal neurogenesis (24), we investigated
the effect of NUDT6 overexpression on hippocampal neurogenesis by Ki-67 immunostaining,
a marker for cell proliferation (Figure 3Q). The total number of Ki-67-positive cells was
significantly decreased in NUDT6-AAV2 group compared to Blank-AAV2 group (p=0.04)
(Figure 3R). These findings suggest that inhibition of neurogenesis may be a potential
contributor to NUDT6's mechanism of action in mediating depression-like behavior.
106

107 5. Hippocampal NUDT6 knockdown reduces depression- and anxiety-like behavior
108 and increases hippocampal neurogenesis

To determine if NUDT6 could be a potential target for antidepressant interventions, we
knocked down the expression of NUDT6 by bilateral intrahippocampal microinjections of
lentivirus expressing NUDT6-shRNA (Figure 4A). One month after the injections, we
performed the behavioral tests FST and SPT to assess depression-like behavior and EPM test
to assess anxiety-like behavior. Following behavioral experiments, the rats' hippocampi were
dissected and analyzed for NUDT6 and FGF2 protein levels, in addition to Ki-67
immunostaining to evaluate neurogenesis (Figure 4A). There was a significant decrease in
NUDT6 protein levels in the NUDT6-shRNA group compared to the non-targeting-shRNA
group confirming a successful knockdown (Figure 4B). We did not observe any difference in
FGF2 levels between groups (Figure 4B). Hippocampal NUDT6 knockdown resulted in an
increase in sucrose preference in the SPT when compared to the non-targeting-shRNA control
group (p=0.026) (Figure 4C). On the other hand, we did not observe any differences in
durations spent immobile, swimming and climbing in FST between groups (p=0.99, 0.83 and
0.72, respectively) (Figure 4D).

Experimental reduction of hippocampal NUDT6 protein levels increased total time
spent in distal half of the open arms as well as the frequency of entrance into the distal open
arms in EPM, indicating an overall anxiolytic effect (p=0.10 and 0.14, respectively) (Figure 4E-
F). Total distance traveled in the EPM was similar between groups (p=0.86), suggesting that
the difference reported above was not due to a change in locomotor activity (Figure S1E).

Finally, we evaluated the effects of hippocampal knockdown of NUDT6 on
neurogenesis by Ki-67 immunostaining, and found a significant increase in Ki-67-positive cell
number in NUDT6-shRNA group (Figure 4G, 4H). In summary knockdown of NUDT6 in the
hippocampus had antidepressant and anxiolytic effects and was associated with an increase
in neurogenesis, identifying NUDT6 as a potential target in developing novel antidepressants.

Discussion

In this study, we showed that increasing hippocampal NUDT6 levels by 33% results in depression- and anxiety-like behaviors possibly by inducing inflammatory signaling and suppressing hippocampal neurogenesis. HPLC-MS proteomics analysis showed that hippocampal NUDT6 overexpression led to increased expression of S100A9, a DAMP/alarmin, which was confirmed by immunoblotting and qRT-PCR. S100A9 intensity was found to be increased by approximately 3-fold in hilar and dentate gyrus neurons by immunohistochemical analysis. There was a 2-3 fold rise in the nuclear translocation of NF-κB in S100A9 positive cells in hilus and dentate gyrus, pertaining to the activation of inflammation. In parallel with S100A9 rise and NF-κB activation, microglial cell number was found to be increased. Plausibly microglia were recruited to the hilar region in response to chemokines released as a result of the proinflammatory state induced by NUDT6 overexpression. These findings suggest that hippocampal NUDT6 overexpression leads to depression-like behavior possibly by promoting a proinflammatory state, which is highly regarded to play an important role in the pathophysiology of depression based on several lines of evidence in the literature (24,36–40).

We did not find any changes in FGF2 mRNA or protein levels after NUDT6 overexpression in the hippocampus, on contrary to previous *in vitro* evidence mainly from cancer cells, suggesting negative regulation of FGF2 expression by FGF-AS transcript (12–15,35). This discrepancy may have resulted from completely different gene regulatory mechanisms in highly mitotic cancer cells and post-mitotic brain cells. Supporting this suggestion, Asa et al (2001) reported that overexpression of FGF-AS did not alter FGF2 expression in a pituitary derived cell line (16). In addition, we found that NUDT6 overexpression did not change phosphorylation of ERK1/2 and Akt kinases, the downstream signaling mediators of FGF2 that mediate its antidepressant effects (22,23). Thus, we conclude that the effects of hippocampal NUDT6 protein on affective behavior are not mediated through modulation of FGF2 or its downstream signaling pathways, but NUDT6 has functions of its own.

Involvement of S100A9 in stress response and depression-like behavior has also been reported by a recent study, which showed that chronic unpredictable mild stress (CUMS), a well-established depression model increased hippocampal S100A9 levels (36). In this study ICV administration of recombinant S100A9 increased depression-like behavior in mice, whereas intraperitoneal injection of S100A9 inhibitor during CUMS period blocked depressogenic effects of CUMS in mice (36). S100 protein family is comprised of several calcium-binding proteins, which form homo/heterodimers functioning by crosslinking target proteins (41). This is important in the regulation of activity of several intracellular proteins, hence, various cellular functions, including inflammation, cell proliferation and differentiation, cytoskeletal dynamics and transcription (41,42). In addition to their intracellular functions,

some S100 proteins, including S100A9 are secreted by unknown mechanisms, which then
bind to pattern recognition receptors on the cell membrane resulting in activation of NF-κB in
the target cell (26,42,43). S100A9 can also induce secretion of cytokines and chemokines in
microglial cell cultures as well as in peripheral blood monocytic cells, which upon release may
activate NF-κB in neighboring cells (44,45). Consistent with these reports, we found an
increase in nuclear translocation of NF-κB besides the increased levels of S100A9 in the
NUDT6 overexpression group. We further showed that S100A9 immunopositivity was
increased in NUDT6-positive cells and NF-κB activation was increased in both NUDT6-positive
and S100A9-positive cells. These findings suggest that NUDT6 overexpression induces
neuronal S100A9 levels and activates NF-κB. The consequent NF-κB activation could result
in an increase in transcription and release of cytokines & chemokines augmenting the
proinflammatory state. Supporting these suggestions, stress increased NF-κB activation in
both hippocampus and prefrontal cortex in mice, and ICV administration of an NF-κB inhibitor
during 4 weeks of CUMS blocked effects of CUMS on anxiety- and depression-like behavior
(24,39).
We observed an increase in microglia number in the hippocampus upon NUDT6
overexpression. This finding is consistent with previous reports showing a rise in microglia
number in animal models of depression (37,38). In addition, S100A9 has been reported to
promote microglial cell migration in microglial cell cultures (45). Recruitment of microglial cells
by S100A9 or chemokines transcribed by activated NF-κB may account for the observed
increase in microglia number following NUDT6 overexpression.

Overexpression of NUDT6 resulted in a reduction in hippocampal neurogenesis, which
suggests that the depressogenic effects of NUDT6 encompass a decrease in hippocampal
neurogenesis. This is in line with a vast body of literature reporting involvement of adult
hippocampal neurogenesis in the pathophysiology of depression (reviewed in (46)).
Hippocampal neurogenesis was reportedly decreased by both acute and chronic stress
paradigms modeling depression in rodents (47,48) (reviewed in (49)) which can be reversed
by antidepressant treatments (34,50). Moreover, effects of acute and chronic stress on
neurogenesis were reported to involve NF-κB signaling. Stress activated NF-κB signaling in
the adult hippocampus, whereas inhibition of NF-κB signaling blocked the effects of acute or
chronic stress on neurogenesis (24). Our findings of decreased neurogenesis together with
increased activation of NF-κB in NUDT6 overexpression group suggest that NUDT6 may
reduce neurogenesis by activating inflammatory pathways. In summary, NUDT6
overexpression induces proinflammatory signaling in the hippocampus, aggravated by newly
synthesized chemokines and cytokines, which then inhibits neurogenesis and recruits
microglia.

Our findings point out NUDT6 as a novel antidepressant target. We, indeed, showed
that hippocampal NUDT6 knockdown had antidepressant and anxiolytic effects. This effect
was accompanied by an increase in hippocampal neurogenesis. As neurogenesis has been
reported to be necessary for behavioral effects of the antidepressants (34), our findings
indicate that inhibition of suppressive effects of NUDT6 on neurogenesis may be involved in
antidepressant and anxiolytic effects of NUDT6 knockdown. These effects, similar to those of
NUDT6 overexpression, were not associated with changes in hippocampal FGF2 levels.

In conclusion, our results identify NUDT6 as a potential mediator of depression-like
behavior and a novel target for the antidepressant drug development. Depressogenic effect of
NUDT6 was not associated with a change in FGF2 signaling, rather it was associated with an
induction of neuroinflammatory signaling, which involves S100A9 expression, NF-κB activation
and microglial recruitment. These findings are consistent with previous reports implying a role
for inflammatory signaling in the development of depression (24,36,37,49). Our findings add
NUDT6 and S100A9 among the players in the pathophysiology of depression and reveal novel
targets for the treatment of MDD. Furthermore, this is the first *in vivo* study, to our knowledge,
which disclosed a functional role for an antisense protein in the CNS, independent of the levels
of its sense protein. These findings suggest that transcription of the same gene in the sense
and antisense direction may enable the organism to fine-tune behavioral responses as
exemplified here by NUDT6 and FGF2, both transcribed from FGF2 gene but have opposite
actions on depression-like behavior. This encouraging finding may lead to new discoveries
uncovering other protein-coding NATs that may have biological functions independent of the
regulation of their sense transcript.

As for the limitations, we performed the experiments in only male rats, therefore our
findings cannot be generalized to females. There is evidence in the literature that microglia
and neurons respond differentially to physical/psychological stressors in male animals than in
females (38,51). Therefore, possible involvement of NUDT6 in the development of depression-
like behavior in females needs to be addressed in future studies.
101

Methods and Materials

ANIMALS

Male Sprague Dawley rats, weighing 300 to 400 grams were used in this study. All animals were maintained on a 12-hour light/dark cycle with food and water available ad libitum. Animals were habituated to the housing conditions for 1 week before any manipulations. They were weighed twice weekly and monitored closely for any deterioration in their health. All the experimental procedures were approved by the Animal Experimentations Local Ethics Board of the Hacettepe University.

PLASMIDS

 pEGFP-N1 plasmid expressing full-length rat FGF-AS gene (a generous gift from Dr. Paul Murphy, Dalhousie University, Canada) was purified as endotoxin-free plasmids (Qiagen, endo-free plasmid Maxi-kit). To facilitate the in vivo delivery of the plasmids, a transfection reagent was used (in vivo-JetPEI-TM, Polyplus transfection France) according to the manufacturer's protocol. Briefly, 3 µg of plasmid was mixed with in vivo JetPEI with an N/P ratio of 6 in a total volume of 7 µl for each rat. The plasmid-in vivo JetPEI mix were prepared daily, 30 minutes before the microinjections. For the controls, we used in vivo JetPEI mix prepared without the plasmid.

VIRUS

Adeno-associated virus (AAV) serotype 2 was used to overexpress full-length rat NUDT6 in the hippocampus (pAAV-G-CMV-2A-GFP)(Abmgood, aavp6966983). A blank AAV (pAAV-G-CMV-blank) was used as a control (Abmgood, aavp002). In order to knockdown FGF-AS, lentivirus (LV) expressing shRNA targeting rat NUDT6 mRNA was injected into the hippocampus (Smart Vector Lentiviral Rat NUDT6 shRNA, Clone id: V3SVRN00_21213469, antisense sequence: AACCCTTCCCTGTGTCCG, Dharmacon). Non-targeting shRNA (with no complementarity to any known mammalian gene) expressing LV was used as control (Smart Vector Non-targeting hCMV-TurboGFP-Control, catalog# S-005000-01, Dharmacon).

SURGICAL PROCEDURES

Rats were anesthetized with ketamine (50-100 mg/kg) and xylazine (5-10mg/kg) prior to surgical procedures. For icv injections, following the application of anesthesia 26 gauge guide cannulas (Plastics One Inc., VA. USA) were implanted into the left lateral ventricle (coordinates from bregma: anteroposterior -1.4, mediolateral +1.6, dorsoventral -2.5) and anchored to the skull with dental cement for. Five to seven days after surgery, a 33 gauge injector cannula was inserted extending 1 mm below the tip of the guides. The

microinjection cannula was connected by PE-50 tubing to a Hamilton syringe mounted on a

syringe pump (World Precision Instruments, FL. USA). Rats were microinjected with either

pEGFP-N1 plasmid expressing full-length rat FGF-AS gene or in vivo JetPEI mix. Injections

were performed at a rate of 1µl/min (total volume of injection=7 µl) for 14 days.

For viral injections, either NUDT6-AAV or NUDT6-shRNA-LV were injected bilaterally

into the hippocampus (coordinates from bregma: anteroposterior -5.0, mediolateral +/-3.5,

dorsoventral -4.0) by two micro-injectors (23 gauge), which were connected by PE-50 tubing

to two Hamilton syringes mounted on a syringe pump (World Precision Instruments, FL. USA).

Injections were performed at a rate of 0.25µl/min for 4 minutes; the microinjectors were left at

the injection site for 5 minutes to allow diffusion of the virus then removed slowly in 5 minutes.

The cranial hole was filled with bone wax and the skin was sutured with 6.0 silk.

BEHAVIORAL TESTS

Behavioral tests were conducted 4 weeks after the injections of AAV or LV in order the

constructs to express and exert their functions. pEGFP-N1 plasmid expressing full-length rat

NUDT6 was injected for 14 days, and sucrose preference test and elevated plus maze test

were conducted on the 11th and 13th day of injections, respectively; whereas forced swim test

was evaluated 1 day after the last injection.

Sucrose Preference Test

The first day of the test, rats are habituated to consume sucrose by placing 2 bottles both filled

with 1% sucrose solution. Then on the next day, one of the bottles is replaced with tap water,

and 24 hours later total consumption of water and sucrose solutions is measured. For the

repeated icv NUDT6-EP or vehicle-injected groups, animals were given access to 2 bottles,

one filled with 1% sucrose and the other one filled with tap water for 2 hours after 12 hours of

water deprivation. Total water and sucrose solution consumption were measured at the end of

2 hours. Sucrose preference is calculated as the percentage of sucrose consumption to total

fluid consumption.

Elevated Plus Maze (EPM)

The rats are placed at the center of an elevated plus maze facing the closed arm. All four arms

are 55 cm high, 35 cm long and 8 cm wide, with walls surrounding the closed arms that are 51

cm high. The frequency of entrance into the center, closed arms, open arms and distal half of

open arms as well as time spent in each compartment are recorded for 5 minutes and analyzed

with Ethovision XT v.8 software (Noldus Information Technology).

Forced Swim Test (FST)

The rats are placed in transparent cylinders (diameter=26 cm, height=43 cm), filled with water at 22-24°C for 15 minutes and 5 minutes on the first and second days of the experiment, respectively. During the second day, the time spent swimming and climbing and the immobility time are recorded and then manually scored with Ethovision XT v.8 software by an investigator blinded to groups.

WESTERN BLOTTING

Following the euthanasia of the rats, the hippocampi were rapidly dissected on ice after. All samples were homogenized in RIPA buffer (NaCl, Triton-100-X, sodium deoxycholate, 1.5 M Tris Buffer [pH= 8.0], %10 SDS). Protein concentration was measured by BCA protein assay kit (Pierce) and equal amount from each sample was loaded and separated on 4-12 % Bis-Tris SDS gels (Novex, Invitrogen). The proteins were transferred to PVDF membranes (Novex, Invitrogen), which were blocked in 5% dry milk powder in TBS-T for 1 hour and then incubated in primary antibody solutions (Rabbit anti NUDT6, Proteintech 1:800; mouse anti FGF2, Millipore, 1:500; mouse anti ß-actin, Sigma, 1:7500; Rabbit anti ERK1/2, 1:1000, CST; Rabbit anti Akt 1:800, CST; Rabbit anti phospho-ERK1/2, 1:1000, CST; Rabbit anti phospho-Akt, 1:1000, CST; Rabbit anti S100A9, 1:200, CST). After treatment with the secondary antibody (anti-rabbit HRP-conjugated, 1:1000 or anti-mouse HRP-conjugated, 1:1000, Sigma), membranes were incubated in chemoluminescent (SuperSignal West Femto, Pierce) for 5 minutes and then imaged with Kodak 4000MM Image station. The band intensities were measured by Image J software (NIH). The optical intensities of all bands were normalized to that of beta-actin. To be able to compare band intensities on more than one membrane, each membrane was loaded with the same 2 samples, and the mean of the band intensities of these samples were used to normalize the intensities of the other bands on the same membrane.

IMMUNOFLUORESCENT STAINING

 For immunofluorescent staining, rats were first perfused with 4% paraformaldehyde (PFA) under deep anesthesia and their brains were collected. For each hippocampus three 40 µm thick sections that are 200 µm apart were chosen for immunolabeling. Sections were blocked in 10% NGS at room temperature (RT) and incubated in primary antibody (Anti S100A9, 1:200, CST; Anti Iba-1, 1:200, Wako; Anti NeuN, 1:200, Millipore; Anti NF-κB p65, 1:200, CST) at 4 °C overnight and in secondary antibody (Goat Anti-rabbit 1:200, Invitrogen) at RT for 1 hour. For double immunofluorescent stainings sections were blocked in 10% NGS at RT and incubated in primary antibodies simultaneously (Anti S100A9, 1:200, CST- Anti NF-κB p65, 1:200, SantaCruz; Anti NUDT6; ProteinTech, 1:200- Anti NF-κB p65, 1:200, SantaCruz; Anti NUDT6; ProteinTech, 1:200- Anti FGF2; Thermo Fisher, 1:200) at 4 °C 48 hours and in secondary antibodies simultaneously (Goat Anti-mouse 1:200, Invitrogen, Goat Anti- Rabbit

1:200, Jackson) at RT for 1 hour. There is an antigen retrieval step (80°C in citrate buffer) for

all NF-κB double staining before blocking step. For NUDT6-S100A9 double staining; sections

were blocked in %10 NGS at RT and incubated in anti NUDT6 antibody (1:200, ProteinTech)

at 4 °C 48 hours and in secondary antibody (Goat Anti- Rabbit 1:200, Jackson) at RT for 1

hour. S100A9-NeuN or S100A9-Iba1 co-stainings were performed by incubating the sections

in both primary antibodies which are labeled with fluorescent $F(ab)_2$ fragments at 4 °C for 20

hours. Images from hippocampus were taken with 25X or 40X magnification with confocal

microscope (Leica). An investigator blinded to the groups performed analyses of the labeled

sections. Upon imaging, corrected total cell fluorescence (CTCF) was calculated for S100A9

labeling in the hilar and dentate gyrus neurons, respectively. Briefly, fluorescence intensity of

S100A9 labeling was measured in NeuN positive neurons (ROI) via ImageJ from which

background fluorescence was substracted. CTCF is calculated by the following formula:

Fluorescence intensity (ROI)- (area of the selected cell x mean fluorescence of background

readings). The nuclear translocation of NF-κB-p65 was calculated by the ratio of dentate gyrus

cells positive for nuclear NF-κB staining to the total number of Hoechst positive dentate gyrus

cells (around 200 cells per image). We counted and averaged NF-κB positive nuclei in 3

sections, 200 µm apart. Iba-1 positive microglia were imaged in Z-stacks and counted and

averaged in maximum projection images obtained from three, 200 µm apart, sections. For

S100A9 and Iba-1 co-labeling, ~25 µm Z-stacks (1024x1024 pixels, ~370x370 µm2) with

isotropic voxel spacing were processed using FIJI/ImageJ 2.1.0/1.53f. for background

subtraction (Rolling ball Radius of 50 pixels and 5 pixels for red and green channel,

respectively).

Quantitative RT-PCR

Hippocampi were dissected and stored in RNAlater solution (QIAGEN) at -20 C until RNA

isolation. Total RNA was extracted from the hippocampi with RNeasy mini kit (QIAGEN) as per

manufacturer's instructions. For cDNA synthesis 500 ng of total RNA was used. cDNAs were

synthesized with random hexamer primers with RevertAid First Strand cDNA Synthesis Kit

(Thermo Fisher Scientific, K1621) according to instructions and stored at −20°C. Quantitative

RT-PCR was performed with Taqman probe-based technology. Taqman gene expression

master mix (ABI, Foster city, CA, 4369016), FAM-TAMRA labeled Taqman probes for rat FGF-

AS transcript (Assay ID: Rn00710461_m1), rat beta-actin gene (Assay ID: Rn01775763_g1),

rat FGF2 gene (Taqman probe: CGT CCA TCT TCC TTC ATA GCC AG ; forward primer: GGA

GTT GTG TCC ATC AAG; reverse primer: CAC TCT TCT GTA ACA CAC TTA) and rat

S100A9 gene gene (Taqman probe:TGC TGC GCT CCA GCT GAG ATC C; forward primer:

GCT TTG AGC AAG AAG ATG GCT; reverse primer: CCT TGT TCA GGG TGT TCA GGA)

were used. PCR was carried out in triplicates in ABI OneStep Q RT-PCR machine (ABI).

Thermal cyclic conditions were as follows: 50°C for 2 min, 95°C for 10 min followed by 40
cycles of 95°C 15 s, 60°C for 1 min. The relative expression values were calculated with ΔΔCt
method.
151

PROTEOMIC ANALYSIS
Hippocampi that were dissected 8 weeks after the injection of either NUDT6 overexpressing
AAV (n=5) or Blank AAV (n=6) were sent to Acıbadem University Department of Biochemistry
for proteomic analysis by using Liquid Chromatography Tandem Mass Spectrometry (LC-
MS/MS). The raw data obtained from LC-MS/MS was processed and compared between
groups. Principal Component Analysis was performed and proteins with significant expression
differences were determined. The network and pathway analysis were performed using online
bioinformatics websites (Panther, String). The activated or inhibited proteins were determined
by their relative activation z-scores.
161

STATISTICAL ANALYSES
Sample sizes were selected based on similar studies in the literature; no statistical analysis
was conducted to predetermine sample size. Data was analyzed by SPSS 22. Kolmogorov
Smirnov Test was used to test for the normality of the distribution. Variables that were normally
distributed were compared with student t-test, whereas those that were not normally distributed
were compared with Mann Whitney-U test. All statistical tests were conducted two-sided. Data
are presented as mean ± S.E.M.
169

ACKNOWLEDGEMENTS
This project was supported by the Scientific and Technological Research Council of Turkey,
grant number: SBAG 110S481; L'Oréal-UNESCO For Women in Science Award to EEK;
Hacettepe University Research Projects Management System, grant number: THD-2016-
11687, and Science Academy Young Scientists Award Program to EEK. We thank Dr. Paul
Murphy (Dalhousie University, Canada) for his generosity in providing the rat NUDT6
expressing plasmid, and Mesut Fırat and Sinem Yılmaz-Özcan for their expert help in technical
issues. We thank Dr Cortney Turner and Aiden Houcek for their valuable opinions on the
manuscript. Authors declare no competing financial interests. Graphical depictions were
prepared at Biorender.com.

AUTHOR CONTRIBUTIONS
Design and conception of the study: EEK ; acquisition and analysis of data: EEK, BU, FOH,
MY, ECE, ABV and KB; scientific discussions and interpretation of data: BU, EEK, TD, YA,
KB; drafting the manuscript: BU, EEK and TD

REFERENCES

1. Ferrari AJ, Somerville AJ, Baxter AJ, Norman R, Patten SB, Vos T, Whiteford HA (2013): Global variation in the prevalence and incidence of major depressive disorder: a systematic review of the epidemiological literature. *Psychol Med* 43: 471–481.

2. Nemeroff CB (2007): Prevalence and management of treatment-resistant depression. *J Clin Psychiatry* 68 Suppl 8: 17–25.

3. Trivedi MH, Rush AJ, Wisniewski SR, Nierenberg AA, Warden D, Ritz L, *et al.* (2006): Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. *Am J Psychiatry* 163: 28–40.

4. Duman RS, Monteggia LM (2006): A neurotrophic model for stress-related mood disorders. *Biol Psychiatry* 59: 1116–1127.

5. Turner CA, Watson SJ, Akil H (2012): Review The Fibroblast Growth Factor Family : Neuromodulation of Affective Behavior. *Neuron* 76: 160–174.

6. Eren-Koçak E, Turner CA, Watson SJ, Akil H (2011): Short-hairpin RNA silencing of endogenous fibroblast growth factor 2 in rat hippocampus increases anxiety behavior. *Biol Psychiatry* 69: 534–540.

7. Turner CA, Gula EL, Taylor LP, Watson SJ, Akil H (2008): Antidepressant-like effects of intracerebroventricular FGF2 in rats. *Brain Res* 1224: 63–68.

8. Hoshaw BA, Hill TI, Crowley JJ, Malberg JE, Khawaja X, Rosenzweig-Lipson S, *et al.* (2008): Antidepressant-like behavioral effects of IGF-I produced by enhanced serotonin transmission. *Eur J Pharmacol* 594: 109–116.

9. Perez JA, Clinton SM, Turner CA, Watson SJ, Akil H (2009): A new role for FGF2 as an endogenous inhibitor of anxiety. *J Neurosci Off J Soc Neurosci* 29: 6379–6387.

10. Modarresi F, Faghihi MA, Lopez-Toledano MA, Fatemi RP, Magistri M, Brothers SP, *et al.* (2012): Inhibition of natural antisense transcripts in vivo results in gene-specific transcriptional upregulation. *Nat Biotechnol* 30: 453–459.

11. Xu Y-H, Luo Y, Cao J-B, Liu Y-H, Song Y-X, Zhang X-Y, *et al.* (2022): lncRNA BDNF-AS Attenuates Propofol-Induced Apoptosis in HT22 Cells by Modulating the BDNF/TrkB Pathway. *Mol Neurobiol.* https://doi.org/10.1007/S12035-022-02757-Y

12. Baguma-Nibasheka M, Li AW, Murphy PR (2007): The fibroblast growth factor-2 antisense gene inhibits nuclear accumulation of FGF-2 and delays cell cycle progression in C6 glioma cells. *Mol Cell Endocrinol* 267: 127–136.

13. Baguma-Nibasheka M, Macfarlane LA, Murphy PR (2012): Regulation of fibroblast growth factor-2 expression and cell cycle progression by an endogenous antisense RNA. *Genes (Basel)* 3: 505–520.

14. Li AW, Murphy PR (2000): Expression of alternatively spliced FGF-2 antisense RNA transcripts in the central nervous system: regulation of FGF-2 mRNA translation. *Mol Cell Endocrinol* 162: 69–78.

15. MacFarlane L-A, Gu Y, Casson AG, Murphy PR (2010): Regulation of fibroblast growth factor-2 by an endogenous antisense RNA and by argonaute-2. *Mol Endocrinol* 24: 800–812.

16. Asa SL, Ramyar L, Murphy PR, Li AW, Ezzat S (2001): The endogenous fibroblast growth factor-2 antisense gene product regulates pituitary cell growth and hormone production. *Mol Endocrinol* 15: 589–599.

17. Baguma-Nibasheka M, Li AW, Osman MS, Geldenhuys L, Casson AG, Too CKL, Murphy PR (2005): Coexpression and regulation of the FGF-2 and FGF antisense genes in leukemic cells. *Leuk Res* 29: 423–433.

18. Frank MG, Der-Avakian A, Bland ST, Watkins LR, Maier SF (2007): Stress-induced glucocorticoids suppress the antisense molecular regulation of FGF-2 expression. *Psychoneuroendocrinology* 32: 376–384.

19. Xu J, Wang Q, Freeling M, Zhang X, Xu Y, Mao Y, *et al.* (2017): Natural antisense transcripts are significantly involved in regulation of drought stress in maize. *Nucleic Acids Res* 45: 5126–5141.

20. Elsayed M, Banasr M, Duric V, Fournier NM, Licznerski P, Duman RS (2012): Antidepressant effects of fibroblast growth factor-2 in behavioral and cellular models of depression. *Biol Psychiatry* 72: 258–265.

21. Turner CA, Eren-Koçak E, Inui EG, Watson SJ, Akil H (2016): Dysregulated fibroblast growth factor (FGF) signaling in neurological and psychiatric disorders. *Semin Cell Dev Biol* 53: 136–143.

22. Tang M-M, Lin W-J, Zhang J-T, Zhao Y-W, Li Y-C (2017): Exogenous FGF2 reverses depressive-like behaviors and restores the suppressed FGF2-ERK1/2 signaling and the impaired hippocampal neurogenesis induced by neuroinflammation. *Brain Behav Immun* 66: 322–331.

23. Wang L, Li X-X, Chen X, Qin X-Y, Kardami E, Cheng Y (2018): Antidepressant-Like Effects of Low- and High-Molecular Weight FGF-2 on Chronic Unpredictable Mild Stress Mice. *Front Mol Neurosci* 11: 377.

24. Koo JW, Russo SJ, Ferguson D, Nestler EJ, Duman RS (2010): Nuclear factor-kappaB is a critical mediator of stress-impaired neurogenesis and depressive behavior. *Proc Natl Acad Sci U S A* 107: 2669–2674.

25. Iwata M, Ota KT, Li X-Y, Sakaue F, Li N, Dutheil S, *et al.* (2016): Psychological Stress Activates the Inflammasome via Release of Adenosine Triphosphate and Stimulation of the Purinergic Type 2X7 Receptor. *Biol Psychiatry* 80: 12–22.

26. Ma L, Sun P, Zhang J-C, Zhang Q, Yao S-L (2017): Proinflammatory effects of S100A8/A9 via TLR4 and RAGE signaling pathways in BV-2 microglial cells. *Int J Mol Med* 40: 31–38.

27. Németh J, Stein I, Haag D, Riehl A, Longerich T, Horwitz E, *et al.* (2009): S100A8 and S100A9 are novel nuclear factor kappa B target genes during malignant progression of murine and human liver carcinogenesis. *Hepatology* 50: 1251–1262.

28. Vogl T, Tenbrock K, Ludwig S, Leukert N, Ehrhardt C, van Zoelen MAD, *et al.* (2007): Mrp8 and Mrp14 are endogenous activators of Toll-like receptor 4, promoting lethal, endotoxin-induced shock. *Nat Med* 13: 1042–1049.

29. Shih R-H, Wang C-Y, Yang C-M (2015): NF-kappaB Signaling Pathways in Neurological Inflammation: A Mini Review. *Front Mol Neurosci* 8: 77.

30. Liu T, Zhang L, Joo D, Sun S-C (2017): NF-κB signaling in inflammation. *Signal Transduct Target Ther* 2: 17023-.

31. Eren-Koçak E, Dalkara T (2021): Ion Channel Dysfunction and Neuroinflammation in Migraine and Depression. *Front Pharmacol* 12. https://doi.org/10.3389/fphar.2021.777607

32. Dranovsky A, Hen R (2006): Hippocampal neurogenesis: regulation by stress and antidepressants. *Biol Psychiatry* 59: 1136–1143.

33. Hill AS, Sahay A, Hen R (2015): Increasing Adult Hippocampal Neurogenesis is Sufficient to Reduce Anxiety and Depression-Like Behaviors. *Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol* 40: 2368–2378.

34. Santarelli L, Saxe M, Gross C, Surget A, Battaglia F, Dulawa S, *et al.* (2003): Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science* 301: 805–809.

35. Knee R, Li AW, Murphy PR (1997): Characterization and tissue-specific expression of the rat basic fibroblast growth factor antisense mRNA and protein. *Proc Natl Acad Sci U S A* 94: 4943–4947.

36. Gong H, Su W-J, Cao Z-Y, Lian Y-J, Peng W, Liu Y-Z, *et al.* (2018): Hippocampal Mrp8/14 signaling plays a critical role in the manifestation of depressive-like behaviors in mice. *J Neuroinflammation* 15: 252.

37. Wachholz S, Eßlinger M, Plümper J, Manitz M-P, Juckel G, Friebe A (2016): Microglia activation is associated with IFN-α induced depressive-like behavior. *Brain Behav Immun* 55: 105–113.

38. Wohleb ES, Terwilliger R, Duman CH, Duman RS (2018): Stress-Induced Neuronal Colony Stimulating Factor 1 Provokes Microglia-Mediated Neuronal Remodeling and Depressive-like Behavior. *Biol Psychiatry* 83: 38–49.

39. Gárate I, Garcia-Bueno B, Madrigal JLM, Caso JR, Alou L, Gomez-Lus ML, *et al.* (2013):

Stress-induced neuroinflammation: role of the Toll-like receptor-4 pathway. *Biol Psychiatry* 73: 32–43.

40. Bierhaus A, Wolf J, Andrassy M, Rohleder N, Humpert PM, Petrov D, *et al.* (2003): A mechanism converting psychosocial stress into mononuclear cell activation. *Proc Natl Acad Sci U S A* 100: 1920–1925.

41. Donato R (2001): S100: a multigenic family of calcium-modulated proteins of the EF-hand type with intracellular and extracellular functional roles. *Int J Biochem Cell Biol* 33: 637–668.

42. Donato R (2003): Intracellular and extracellular roles of S100 proteins. *Microsc Res Tech* 60: 540–551.

43. Chen B, Miller AL, Rebelatto M, Brewah Y, Rowe DC, Clarke L, *et al.* (2015): S100A9 induced inflammatory responses are mediated by distinct damage associated molecular patterns (DAMP) receptors in vitro and in vivo. *PLoS One* 10: e0115828.

44. Simard J-C, Cesaro A, Chapeton-Montes J, Tardif M, Antoine F, Girard D, Tessier PA (2013): S100A8 and S100A9 Induce Cytokine Expression and Regulate the NLRP3 Inflammasome via ROS-Dependent Activation of NF-κB1. *PLoS One* 8: 1–12.

45. Wu M, Xu L, Wang Y, Zhou N, Zhen F, Zhang Y, *et al.* (2018): S100A8/A9 induces microglia activation and promotes the apoptosis of oligodendrocyte precursor cells by activating the NF-κB signaling pathway. *Brain Res Bull* 143: 234–245.

46. Chooniedass-Kothari S, Emberley E, Hamedani MK, Troup S, Wang X, Czosnek A, *et al.* (2004): The steroid receptor RNA activator is the first functional RNA encoding a protein. *FEBS Lett* 566: 43–47.

47. Gascoigne DK, Cheetham SW, Cattenoz PB, Clark MB, Amaral PP, Taft RJ, *et al.* (2012): Pinstripe: a suite of programs for integrating transcriptomic and proteomic datasets identifies novel proteins and improves differentiation of protein-coding and non-coding genes. *Bioinformatics* 28: 3042–3050.

48. Stergachis AB, Haugen E, Shafer A, Fu W, Vernot B, Reynolds A, *et al.* (2013): Exonic transcription factor binding directs codon choice and affects protein evolution. *Science* 342: 1367–1372.

49. Haapakoski R, Mathieu J, Ebmeier KP, Alenius H, Kivimäki M (2015): Cumulative meta-analysis of interleukins 6 and 1β, tumour necrosis factor α and C-reactive protein in patients with major depressive disorder. *Brain Behav Immun* 49: 206–215.

50. Taupin P (2006): Neurogenesis and the effect of antidepressants. *Drug Target Insights* 1: 13–17.

51. Cao P, Chen C, Liu A, Shan Q, Zhu X, Jia C, *et al.* (2021): Early-life inflammation promotes depressive symptoms in adolescence via microglial engulfment of dendritic spines. *Neuron* 109: 2573-2589.e9.

FIGURE LEGENDS

Figure 1. NUDT6 overexpression in the hippocampus increases depression- and anxiety-like behaviors.

A. Graphical depiction of the NUDT6-EP and NUDT6-AAV experimental setup.

B. Repeated icv injections of NUDT6-EP increased time spent immobile and decreased time spent swimming in forced swim test, when compared to controls (n=7-11)

C. Intrahippocampal injection of NUDT6-AAV increased time spent immobile and decreased time spent swimming in forced swim test, when compared to Blank-AAV injected controls (n=13/group).

D. Repeated icv injections of NUDT6-EP decreased time spent in open arms in elevated plus maze (n=7-11/group).

E. Intrahippocampal injection of NUDT6-AAV did not alter time spent in open arms in elevated plus maze (n=13/group)**.**

F. Repeated icv injections of NUDT6-EP increased hippocampal NUDT6 protein levels. Hippocampal FGF2 protein levels, on the other hand, remained unchanged. Representative Western bands for NUDT6 and FGF2 of each group are shown; β-actin is used as loading control (n=5-6/group).

G. Intrahippocampal NUDT6-AAV injection increased NUDT6 mRNA expression and protein levels. FGF2 mRNA expression or protein levels in the hippocampus were not altered as detected by RT-PCR and Western blotting (n=7-8/group). Representative Western bands for NUDT6 and FGF2 of each group are shown; β-actin is used as loading control.

FGF2: fibroblast growth factor 2, NUDT6-AAV: NUDT6 expressing adeno-associated virus, NUDT6-EP: NUDT6-expressing plasmid, RT-PCR: real-time polymerase chain reaction. Significance levels were stated as follows: #p=0.067, ##p=0.05, *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001. ns denotes non-significance.

Figure 2. NUDT6 overexpression in the hippocampus did not cause any change in either FGF2 levels or phosphorylation of its downstream signaling pathways Akt and ERK1/2.

A. Graphical depiction showing FGF2 signaling pathways.

B. Representative Western blot bands for Akt, p-Akt, ERK1/2, p-ERK1/2 of each group are shown; β-actin is used as a loading control (n=7/group)

C. NUDT6 overexpression did not change phosphorylation of Akt or ERK1/2 proteins shown by relative band densities (n=7/group) (ns denotes non-significance).

D. Graphical depiction of the HPLC-Tandem MS proteomics experimental setup.

E. Proteomic analysis revealed an upregulation of 31 and a downregulation of 3 proteins in the NUDT6-AAV group (n/7/group)**.**

F. Thirteen percent (13%) of the upregulated proteins were related to inflammatory signaling whereas 1% were related to mitochondrial dysfunction and 17% were related to cellular growth & proliferation and organismal injury. The remaining 69% is related to miscellaneous pathways.

FGF2: fibroblast growth factor 2, NUDT6-AAV: NUDT6 expressing adeno-associated virus, HPLC-Tandem MS: High performance Liquid Chromatography Mass Spectrometry

Figure 3. NUDT6 overexpression in the hippocampus promotes inflammatory signaling.

A-B. NUDT6 overexpression increased S100A9 protein expression. Representative Western bands for S100A9 of each group are shown; β-actin is used as loading control (n =7/group).

C. NUDT6 overexpression increased S100A9 mRNA expression (n =7-8/group).

D. Colocalization of S100A9 with a neuronal cell marker, NeuN showed that S100A9 levels were increased in neurons. Neurons were labeled with NeuN (red) and S100A9 was labeled with green. Scale bar: 25 μm.

E. Total florescence of S100A9 in NeuN positive cells was increased in both the dentate gyrus and the hilus in the NUDT6-AAV group, when compared to the control group (n =3/group).

F-G. Colocalization of S100A9 and NUDT6 showed that S100A9 significantly increased in NUDT6-positive cells in NUDT6-AAV group compared to the control group. (n =3/group). Scale bar: 20 μm.

H-I. Number of Iba1 labeled cells in the dentate gyrus and hilus was higher in the NUDT6-AAV group than the control group (n =3/group). Scale bar: 25 μm and 5.5 μm for the inset.

J-K. Nuclear translocation of NF-κB was increased in the NUDT6-AAV group in the dentate gyrus (n =3/group). Figure insets show representative images of nuclear labeling of NF-κB (indicating translocation) in dentate gyrus. Scale bar: 25 μm and 4 μm for the inset.

L-M. Colocalization of NF-κB and NUDT6 showed that nuclear translocation of NF-κB significantly increased in NUDT6-positive cells in NUDT6-AAV group compared to the control group (n =3/group). Scale bar: 20 μm.

N-P. Colocalization of S100A9 and NF-κB showed that NF-κB activation was significantly higher in S100A9-positive cells in NUDT6-AAV group than the control group (n =3/group). Scale bar: 20 μm.

Q-R. Hippocampal Ki-67 immunofluorescent staining showed that neurogenesis was significantly lower in NUDT6-AAV group than the control group (n =3/group). Scale bar: 100 μm.

Significance levels were stated as follows: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$. ns denotes non-significance.

Figure 4. Hippocampal NUDT6 knockdown reduces depression- and anxiety-like behavior and increases hippocampal neurogenesis

A. Schematic representation of experimental setup of hippocampal NUDT6 knockdown via NUDT6-shRNA

B. Intrahippocampal NUDT6-shRNA injection decreased NUDT6 protein expression relative to the non-targeting shRNA injected group. Representative Western bands for NUDT6 and FGF2 of each group are shown; β-actin is used as loading control (n=5-6/group).

C. NUDT6 knockdown by intrahippocampal NUDT6-shRNA injection resulted in an increase in sucrose preference (n=6-7/group).

D. Knockdown of NUDT6 in the hippocampus did not change time spent immobile, climbing or swimming in the forced swim test when compared to non-targeting-shRNA control group**.**

E-F. NUDT6-shRNA group spent more time and entered more frequently into the distal open arms of elevated plus maze (n=11-13/group).

G-H. Hippocampal Ki-67 immunofluorescent staining showed that neurogenesis was significantly increased in NUDT6-shRNA group compared to the non-targeting shRNA group (n =3/group). Scale bar: 100 μm.

NUDT6-shRNA: NUDT6 short hairpin RNA. Significance levels were stated as follows: $\#p=0.10$, $\#\#p=0.14$, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$. ns denotes non-significance.

A Intrahippocampal NUDT6-shRNA or non-targeting shRNA Microinjections | Sucrose Preference Test (SPT) | Elevated Plus Maze Test (EPM) | Forced Swim Test (FST) | Hippocampal Dissection & Molecular Analyses

B
Norm to β-Actin
☐ non-targeting shRNA
■ NUDT6-shRNA
NUDT6 *
FGF2 ns
(Western blot: non-targeting shRNA, NUDT6-shRNA — NUDT6, FGF2, β-Actin)

C
Sucrose Preference (%)
non targeting shRNA / NUDT6 shRNA *

D
Time (s)
Immobility ns
Swimming ns
Climbing ns

E
Time in Distal Open Arms(s)
non targeting shRNA / NUDT6 shRNA #

F
Frequency to Enter Distal Open Arms
non targeting shRNA / NUDT6 shRNA ##

G
non-targeting shRNA | NUDT6-shRNA
Ki67 Hoechst
100μm

H
Number of Ki67 Positive Cells
non targeting shRNA / NUDT6 shRNA *