Reconstitution of a minimal ESX-5 type VII secretion system suggests a role for PPE proteins in outer

membrane transport of proteins

C. M. Bunduc[1,2,3,4], Y. Ding[1], C. Kuijl[5], T. C. Marlovits[2,3,4], W. Bitter[1,5], E.N.G. Houben[1#]

[1]Section Molecular Microbiology, Amsterdam Institute for Life and Environment (A-Life), Vrije

Universiteit Amsterdam, Amsterdam, The Netherlands

[2]Centre for Structural Systems Biology, Notkestraße 85, 22607 Hamburg, Germany

[3]Institute of Structural and Systems Biology, University Medical Center Hamburg-Eppendorf,

Notkestraße 85, 22607 Hamburg, Germany

[4]German Electron Synchrotron Centre Notkestraße 85, 22607 Hamburg, Germany

[5]Department of Medical Microbiology and Infection Control, Amsterdam UMC, Amsterdam, The

Netherlands

[#]Corresponding author. Vrije Universiteit Amsterdam, Section Molecular Microbiology, de Boelelaan

1108, 1081 HZ Amsterdam, the Netherlands. Tel.: +31 20 5983579; E-mail: e.n.g.houben@vu.nl

Keywords: ESX-5, mycobacterium, minimal system, secretion complex, type VII secretion

Abstract

Mycobacteria utilize type VII secretion systems (T7SSs) to secrete proteins across their highly

hydrophobic and diderm cell envelope. Pathogenic mycobacteria have up to five different T7SSs, called

ESX-1 to ESX-5, which are crucial for growth and virulence. Here, we use a functionally reconstituted

ESX-5 system in the avirulent species *Mycobacterium smegmatis* that lacks ESX-5, to define the role of

each *esx-5* gene in system functionality. By creating an array of gene deletions and assessing protein

levels of components and membrane complex assembly, we observed that only the five components

of the inner membrane complex are required for its assembly. However, in addition to these five core

components, active secretion also depends on both the Esx and PE/PPE substrates. Tagging the PPE

substrates followed by subcellular fractionation, surface labeling and membrane extraction showed

that these proteins localize to the mycobacterial outer membrane. This indicates that they could play a

role in secretion across this enigmatic outer barrier. These results provide a first full overview of the

role of each *esx-5* gene in T7SS functionality.

Introduction

Gram-negative bacteria have evolved an exceptional variety of specialized secretion systems to mediate protein transport across their diderm cell envelope (Costa *et al*, 2015). Bacterial pathogens usually strictly depend on these systems to complete their infection cycle. Mycobacteria have an unusual diderm cell envelope with a distinctive outer membrane, also known as the mycomembrane, which is mainly composed of mycolic acids (Dulberger *et al*, 2020). These bacteria use a specific group of homologous specialized secretion systems, called type VII secretion systems, for a diverse array of functions, such as intracellular survival, immunomodulation, as well as uptake of nutrients and metabolites (Ates *et al*, 2015; Bunduc *et al*, 2020a; Pym *et al*, 2002; Siegrist *et al*, 2009; Simeone *et al*, 2012). Pathogenic mycobacteria, such as *Mycobacterium tuberculosis*, encode up to five different T7SSs, named ESX-1 to ESX-5 (Bunduc *et al.*, 2020a). The importance of T7SS for pathogenicity is further evidenced by the lack of ESX-1 causing the attenuation of the currently used live vaccine strain *Mycobacterium bovis* BCG (Houben *et al*, 2012a; Pym *et al.*, 2002; Simeone *et al.*, 2012; van der Wel *et al*, 2007).

The mycobacterial *esx* loci contain five conserved T7SS membrane components, *i.e.* EccB, EccC, EccD and EccE (Ecc stands for <u>E</u>SX <u>c</u>onserved <u>c</u>omponent) and Mycosin or MycP. EccB is a single transmembrane domain (TMD) protein with a large periplasmic domain. EccC is a P-loop NTPase with four FtsK/SpoIIIE-like nucleotide binding domains (NBDs) and the motor protein of the complex, which has been shown to bind substrates (Champion *et al*, 2006; Rosenberg *et al*, 2015). EccD is the most hydrophobic component with eleven TMDs and EccE is anchored in the membrane via two TMDs. The four Ecc components have been shown to assemble into a large ~2 MDa machinery (Houben *et al*, 2012b) and several structures of (sub)complexes have been solved by electron microscopy (EM) (Beckham *et al*, 2017; Beckham *et al*, 2021; Famelis *et al*, 2019; Poweleit *et al*, 2019). The fifth membrane component MycP, a subtilisin-like protease with a single C-terminal TMD, interacts with and stabilizes the EccBCDE membrane complex (van Winden *et al*, 2020; van Winden *et al*, 2016). The recently solved cryo-EM structure of a full ESX-5 inner membrane complex of *M. tuberculosis* revealed the position of MycP within this complex, confirming the stabilizing role of MycP within the membrane assembly (Bunduc *et al*, 2021).

Most substrates that are secreted by mycobacterial T7SSs belong to three substrate families, *i.e.* the Esx, PE and PPE protein families. In pathogenic mycobacteria, especially the number of *pe* and *ppe* genes has expanded extensively, covering nearly 10% of the coding capacity (Cole *et al*, 1998). A specific feature of T7SS substrates is that they are secreted as folded heterodimers (Ates, 2020). Two different Esx proteins assemble into a secreted heterodimer, whereas PE proteins interact with specific

PPE proteins to form a structurally similar substrate pair. Dimerization is mediated by hydrophobic interactions resulting in a conserved four helix bundle structure. The C-terminal flexible domain of one of the secretion partner proteins additionally contains a YxxxD/E secretion signal that is important for secretion (Daleke *et al*, 2012).

Mycobacterial T7SSs also have two cytosolic components, EccA and EspG. EspG is a PPE-dedicated chaperone that binds to a hydrophobic patch on the so-called helical tip domain of these proteins (Ekiert & Cox, 2014; Korotkova *et al*, 2014). Binding of EspG to PPE substrates is required to keep PE/PPE pairs soluble in the cytosol (Ekiert & Cox, 2014; Korotkova *et al.*, 2014). EspG is additionally involved in determining system-specific substrate recognition, as swapping the EspG-binding domain between two PPE substrates resulted in rerouting of the ESX-1 substrate PPE68_1 to the ESX-5 system in *Mycobacterium marinum* (Phan *et al*, 2017). EccA is an AAA+ ATPase that has been hypothesized to disrupt the EspG/PPE interaction, leading to recycling of EspG in the cytosol, although the importance of this ATPase for the secretion process remains controversial and could be dependent on growth conditions (Ekiert & Cox, 2014; Phan *et al*, 2018).

Studying the functioning of T7SSs is impaired by the slow growth rate and virulence of pathogenic mycobacteria. Certain systems, such as ESX-3 and ESX-5, are also essential for growth, complicating genetic analyses (Ates *et al.*, 2015; Serafini *et al*, 2009; Siegrist *et al.*, 2009). Furthermore, studying the potential role of specific substrates in the secretion process is hampered by the redundancy between highly similar substrates (Ates, 2020; Gey van Pittius *et al*, 2006). *Mycobacterium smegmatis*, an avirulent and fast-growing mycobacterium, is commonly used as a model for studying mycobacterial physiology. This species possesses ESX-1, ESX-3 and ESX-4, but lacks the ESX-2 and ESX-5 systems, which are exclusively found in slow-growing mycobacteria.

Here, we exploited an ESX-5 system of *Mycobacterium xenopi* that we have previously functionally reconstituted in *M. smegmatis* (Beckham et al., 2017) to study the role of individual *esx-5* genes in system functionality. We took advantage of the limited number of substrate genes present in the introduced *esx-5* locus and defined the minimal number of genes necessary for successful assembly of the ESX-5 membrane complex and ESX-5 mediated secretion.

Results

The inter-dependence of ESX-5 components for stable expression in *M. smegmatis*

To be able to efficiently dissect the individual roles of T7SS components, we used the ESX-5 system of the slow-growing, moderately thermophilic and pathogenic mycobacterial species *M. xenopi* (Fig. 1A), which we have previously functionally reconstituted in *M. smegmatis* (Beckham *et al.*, 2017; van Winden *et al.*, 2020). As *M. smegmatis* does not have and never had a native ESX-5 system, the only ESX-5 substrates are those encoded by the introduced *esx-5* locus, *i.e.* the Esx pair EsxM/EsxN and two PE/PPE pairs that are specific for *M. xenopi* (Fig. 1A). This is in contrast to ESX-5 systems in natural hosts, which secrete multiple Esx pairs and a large number of PE and PPE substrates, of which some show redundancy (Abdallah *et al*, 2007; Ates, 2020). The limited number of substrates secreted by the reconstituted system provides an excellent opportunity to analyze their individual roles in T7SS functioning, bypassing any potential redundancy issues.

We exploited this non-essential reconstituted system to make single deletions of every *esx-5* gene that encodes for system components, *i.e.* the *eccABCDE*, *mycP* and *espG* genes. In addition, we created *esx-5* plasmids that were devoid of all *pe* and *ppe* genes (*Δpe/ppe*), lacking the two *esx* genes (*ΔesxM/N*) or devoid of all genes that encode for substrates (*Δsubstrates*) (Fig. 1A). The *Δsubstrates* and *Δpe/ppe* plasmids included the deletion of a non-conserved gene situated between the *pe/ppe* pairs, *i.e. mxen9359,* which codes for a putative SAM-methyltransferase (Fig. 1A). Finally, we also created a construct that only contained the five conserved membrane components (MC) (Fig 1A). These constructs were used to assess the effect of each gene deletion on the protein levels of other components, on the formation of the ESX-5 membrane complex and on ESX-5 mediated secretion.

First, we analyzed the impact of the various gene deletions on the expression of the other system components by SDS-PAGE and immunoblotting, using specific antisera for the different ESX-5 components. We used isolated cell envelope fractions of *M. smegmatis* bearing the different *esx-5* plasmids to detect membrane components (Fig. 1B) and whole cell lysates of the same strains to assess protein levels of the cytosolic components (Fig. 1C). Deletion of the genes coding for the cytosolic components $EspG_5$ and $EccA_5$ did not affect the protein levels of any of the five inner membrane components, *i.e.* $EccB_5$, $EccC_5$ (strep tagged), $EccD_5$, $EccE_5$ and $MycP_5$ (Fig. 1B). Furthermore, in the absence of $EccA_5$, the protein levels of $EspG_5$ were not affected and *vice versa* (Fig. 1C). Deleting the membrane components $EccB_5$, $EccC_5$ or $MycP_5$ also did not affect protein levels of the four other membrane components and the two cytosolic components. In contrast, the absence of $EccE_5$ seems to affect protein levels of $EccD_5$ and also of $EccA_5$, although the latter could be attributed to polar effects of the gene deletion, as *eccA_5* lies downstream of *eccE_5*. Knocking out *eccD_5* showed the most severe

effect on the protein levels of the other components. While EccC$_5$ and EccE$_5$ were undetectable by

immunoblotting in the absence of EccD$_5$, the protein levels of the membrane components EccB$_5$ and

MycP$_5$ were reduced. Strikingly, the two cytosolic components were also affected by the *eccD$_5$* deletion,

as EspG$_5$ could not be detected anymore and levels of EccA$_5$ were severely reduced. In conclusion,

deletion of the cytosolic components does not impact the protein levels of any of the other

components, while deleting *eccD$_5$* severely impacted the levels of all components.

Only the five membrane components are required for full assembly of the ESX-5 membrane complex

Next, we investigated which components play a role in the successful assembly of the ESX-5 inner

membrane complex. For this, we analyzed DDM-solubilized cell envelope fractions by blue native PAGE

(BN-PAGE), followed by immunoblotting using our anti-EccB$_5$ antibody or an antibody that recognized

the Strep-tag on EccC$_5$ (Fig 1D). Similar to previous observations (Beckham *et al.*, 2017), BN-PAGE

analysis of samples from *M. smegmatis* bearing the complete *esx-5* locus resulted in three main

complexes that reacted with the anti-EccB$_5$ antibody, the largest complex representing the full complex

of ~2 MDa and two subcomplexes of ~900 kDa and ~700 kDa. Staining with the anti-Strep antibody

showed a similar immunoblot pattern. Taking into consideration the influence of the detergent, lipids

and the Coomassie dye on the size estimation of membrane proteins by BN-PAGE (Wittig *et al*, 2010),

these three complexes could represent the full hexameric complex, the dimeric subcomplex and an

individual protomer, respectively. In the absence of the cytosolic components EccA$_5$ and EspG$_5$,

membrane complex assembly was not affected, as the BN-PAGE immunoblot pattern in their absence

was similar as for the WT construct (Fig. 1D). Similarly, none of the substrate deletion plasmids showed

differences in ESX-5 complexes, suggesting that substrates do not play a role in the assembly and/or

stability of the ESX-5 membrane complex. In line with this observation, a plasmid containing only the

five *esx-5* genes that code for the inner membrane components resulted in membrane complex

formation similar to the intact *esx-5* locus.

Deletion of the five membrane component genes, *i.e. eccB$_5$, eccC$_5$, eccD$_5$, eccE$_5$* and *mycP$_5$*,

showed varying effects on complex formation. In line with the observation that the *eccD$_5$* deletion

severely affected expression of the four other membrane components, no complex formation could be

observed in this condition. This suggests that EccD$_5$ is crucial for complex formation and/or stability,

which is in line with its central location within the protomers of the membrane complex (Fig. 1D). While

deletion of either *eccB$_5$, eccC$_5$* or *eccE$_5$* did not drastically affect the expression of the other membrane

components (Fig. 1B), the assembly of the membrane complex, as seen by BN-PAGE analysis, was

affected. Deletion of *eccB$_5$* affected formation of especially the full ~2 MDa complex, which is in line

with previous findings that EccB$_5$ is involved in complex multimerization (Bunduc *et al.*, 2021; Famelis

et al., 2019). In the absence of the central ATPase EccC$_5$, which gates the potential secretion pore with

its TMDs, membrane complex formation was almost completely abolished, with this construct

exhibiting only a ~500 kDa subcomplex containing EccB$_5$, which was not observed with the WT

construct. Deleting *eccE$_5$* showed a reduced amount of complexes, using both antibodies, which is in

line with our previous observation that a lower number of full complexes could be isolated under this

condition (Beckham *et al.*, 2017). In addition, the ~500 kDa complex, also present with the ΔeccC$_5$

construct, was observed in the absence of EccE$_5$ using the anti-EccB$_5$ but not the anti-Strep antibody.

Finally, while the *ΔmycP$_5$* plasmid showed no major differences in the formation of the inner membrane

complex, the ~500 kDa EccB$_5$-containing complex was also observed here. These results were similar to

previous observations in *M. marinum*, where MycP$_1$ and MycP$_5$ of the ESX-1 and ESX-5 system,

respectively, have been reported to be involved in stabilizing its corresponding T7SS membrane

complex (van Winden et al., 2016).

Taken together, optimal assembly of the ESX-5 membrane complex requires only the five

membrane components.

esx-5 locus-encoded PE/PPE proteins are required for secretion of Esx proteins

We previously observed that introduction of the plasmid that contains the complete *esx-5* locus of *M.*

xenopi in *M. smegmatis* resulted in the secretion of the locus-encoded ESX-5 substrate EsxN and also

of *M. marinum* PPE18, which was introduced by an additional plasmid containing the gene pair

pe31/ppe18 (Beckham *et al.*, 2017). This shows that the reconstituted system is functional. We

therefore tested next the impact of the various gene deletions on the ability of this system to secrete

EsxN and PPE18, the latter by again introducing the plasmid containing the *pe31/ppe18* of *M. marinum*

(PPE18 containing a C-terminal HA-tag for detection) (Fig. 2A).

While both EsxN and PPE18-HA were efficiently secreted with the WT construct, they were also

secreted in the absence of *eccA$_5$*, albeit at a reduced level (Fig. 2A). Deletion of the other cytosolic

component *espG$_5$* blocked the secretion of both PPE18-HA and EsxN (Fig. 2A). This critical role of EspG$_5$

for successful secretion of both the PE/PPE proteins and EsxM/EsxN is in line with previous findings

(Korotkova *et al.*, 2014). Additionally, PPE18-HA was not present in the bacterial pellet, suggesting that

EspG$_5$ is essential for the stability of this protein, as observed before for this and other PPE proteins

(Ekiert & Cox, 2014; Korotkova *et al.*, 2014; Phan *et al.*, 2017). In contrast, in the presence of only the

membrane components, which did not show secretion of PPE18-HA, this substrate could be detected

in the pellet fraction. The instability of PPE18 in the absence of EspG$_5$ can be rescued by deletion of

eccA$_5$ and/or the substrate-encoded *esx-5* genes. As all five membrane component deletions showed a

disruption in membrane complex assembly, it is no surprise that they also abolished the secretion of

both EsxN and PPE18-HA. Notably, deleting *eccD5* negatively affected also the protein levels of not only

PPE18-HA, which is in line with the absence of EspG$_5$ under this condition (Fig. 1C), but also of EsxN. In

the absence of EsxM/EsxN, secretion but not the stable expression of PPE18-HA was abrogated,

showing that this Esx heterodimer is required for secretion of the PE/PPE pair (Fig. 3A). Strikingly, in the

absence of PE/PPE proteins encoded by the *esx-5* locus, which did not affect expression of any ESX-5

components nor assembly of the ESX-5 membrane complex, secretion but not stable expression of both

EsxN and PPE18 was abolished (Fig. 2A). From this, we conclude that the PE/PPE pairs of the *esx-5* locus

are important for the secretion of the Esx substrates and at least one other PPE protein. This is in line

with a recent observation for ESX-1 in *M. marinum*, where secretion of EsxA is dependent on the co-

secretion of PPE68 encoded by the same gene cluster (Cronin *et al*, 2022; Damen *et al*, 2022).

To analyze the role of the individual *pe/ppe* gene pairs and of the putative methyltransferase

gene that is located between the two *pe/ppe* pairs, we created four complementation vectors

containing either the entire deleted region (whole; Fig. 1A), one of the two *pe/ppe* gene pairs with the

methyltransferase gene (*Δpe/ppe1* and *Δpe/ppe2*) or only the two *pe/ppe* pairs without the

methyltransferase gene (*Δmethylt.*). EsxN secretion could be rescued by complementation with a

plasmid containing the entire deleted region (Fig. 2B), although the observed secretion levels were

reduced when compared to the levels with the WT construct. This reduction could be due to the fact

that both introduced plasmids contain the same origin of replication, thereby reducing individual

plasmid copy numbers and the overall protein levels. The same (reduced) secretion levels were

observed when we complemented with a plasmid that contains the entire deleted region without the

gene coding for the predicted methyltransferase, suggesting that this methyltransferase gene is not

required for secretion. Methyltransferases are not often associated with *esx-5* or other T7SS loci,

suggesting that the presence of this gene could be a coincidence. However, deletion of either of the

two *pe/ppe* pairs in the complementation construct abrogated secretion of EsxN completely. This shows

that both PE/PPE pairs are required for EsxN secretion. Notably, complementing the *Δpe/ppe* plasmid

with the PE31/PPE18-HA-encoding plasmid also did not restore secretion of EsxN (Fig. 2A). This shows

that although PPE18-HA is secreted in an ESX-5 dependent manner, it cannot take over the role of the

esx-5 locus-encoded PE/PPE proteins in EsxN secretion, in line with a model in which Esx and PE/PPE

proteins are specifically paired for their co-secretion. The observation that both *esx-5*-encoded PE/PPE

proteins and EsxM/EsxN are required for PPE18-HA secretion shows that both substrate classes are

required for ESX-5 functionality.

PPE but not Esx substrates localize to the cell envelope fraction

While the interdependency of substrates for secretion can be explained by a previously proposed dual substrate recognition event by the EccC ATPase, which is required for system activation (Bunduc *et al*, 2020b; Rosenberg *et al.*, 2015), substrates could additionally be dependent on each other beyond the T7SS inner membrane complex. The dimensions of the core membrane complex dictate that it spans only the inner membrane. How protein translocation takes place beyond the inner membrane, especially across the mycobacterial outer membrane, is unknown, as there is no clear data on the identity of the outer membrane channel or components. As our reconstituted ESX-5 system is functional in *M. smegmatis*, the proteins that drive secretion over this second membrane are likely encoded by the introduced *esx-5* cluster. Within the *esx-5$_{Mxe}$* locus, the proteins that have thus far been regarded as substrates are the only proteins that could mediate translocation through the outer membrane, as all other components are localized either in the inner membrane or in the cytosol. In addition, in recent years it has become clear that various PE/PPE proteins are involved in the uptake of nutrients and localize to the cell envelope, with PPE51 as the main example (Korycka-Machala *et al*, 2020; Wang *et al*, 2020). To address whether the PPE substrates encoded by the *M. xenopi esx-5* locus could be involved in outer membrane transport of proteins, we tagged the first and the second PPE protein with a FLAG- and a Strep-tag, respectively, either at the N- or C-terminus. Additionally, to be able to detect EsxN more efficiently, we added a C-terminal HA-tag to this substrate, which has previously been shown not to interfere with the secretion of its ESX-1 equivalent EsxA (Damen *et al*, 2020). Subcellular fractionation and secretion analysis showed that a FLAG-tag at the C-terminus of the first PPE protein, hereafter called PPE$_1$, partially affected secretion of EsxN, whereas placing the tag at the N-terminus had no noticeable effect on EsxN secretion (Fig. EV 1A). The second PPE protein, hereafter called PPE$_2$, could not be detected when tagged with a Strep-tag at the N-terminus and using a Strep antibody, while placing the same tag to the C-terminus showed sufficient protein levels (Fig. EV 1A). With both constructs, EsxN was efficiently secreted to the culture supernatant (Fig. EV 1A), suggesting that PPE$_2$ with the N-terminal tag is expressed, but that the tag is cleaved off. Finally, we could also efficiently detect secreted EsxN-HA using the HA antibody.

With this information in hand, we made a final construct in which we removed the previously introduced Strep-tag at the C-terminus of EccC$_5$ and modified the substrate genes, so that they encode for PPE$_1$ with a FLAG-tag at the N-terminus, PPE$_2$ with a Strep-tag at the C-terminus and an HA-tag at the C-terminus of EsxN. We first analyzed protein secretion by *M. smegmatis* carrying this construct when grown in the presence or absence of Tween-80, a mild detergent used as anti-clumping agent for mycobacteria grown in liquid culture (Fig. 3A). EsxN was present in the culture supernatant irrespective of the presence of Tween-80. In contrast, both PPE proteins were found in the culture supernatant only when cells were grown in the presence of the detergent. In the absence of Tween-80, these proteins

were found on the cell surface instead, as they could be extracted from the cell surface using the mild
detergent Genapol X-080 (Fig 3A). This suggests that the presence of Tween-80 triggers the release of
these proteins from the cell surface into the culture medium. Notably, when the gene encoding for the
central ATPase $EccC_5$ was deleted, neither PPE proteins were secreted anymore, showing that their
secretion is dependent on a functional ESX-5 system (Fig. EV 2).

Even though we could extract the PPE proteins, a substantial amount of both PPE proteins
remained cell-associated after growth with Tween-80 and/or after Genapol X-080 treatment. To
investigate the subcellular location of these subpopulations, *M. smegmatis* cells grown in the presence
of Tween-80 were lysed and fractionated into a soluble (cytosolic) and insoluble (cell envelope) fraction
(Fig. 3B). Intriguingly, both PPE proteins localized mainly to the cell envelope fraction. Furthermore,
flow cytometry analysis using anti-FLAG, -Strep and -HA antibodies revealed that both PPE proteins
could be efficiently detected on the cell surface of cells grown with Tween-80 (Fig. 3C, Fig. EV 1B). In
contrast, EsxN could not be detected via its C-terminal HA-tag, which is in line with the observation that
this substrate is mainly secreted into the culture medium. Interestingly, after treatment of the cells with
Genapol X-080, similar levels of surface labeling could be observed as compared to cells that were not
extracted with Genapol X-080 (Fig. 3C). This suggests that a significant portion of the cell-associated
population of the two PPE proteins is firmly attached to the cell surface. To characterize the cell
envelope association of these two PPE proteins further, we subjected isolated cell envelope fractions
from Genapol X-080-treated cells to solubilization with a range of detergents, some of which have been
used previously for the characterization of mycobacterial membrane proteins (Hoffmann *et al*, 2008;
Målen *et al*, 2010). While our inner membrane protein controls $EccC_5$ and FtsH could be extracted by
the nonionic detergents DDM, beta-octylglucoside and Triton X-100, and the anionic detergent SDS,
but not by Genapol X-080 or the zwitter ionic detergent CHAPSO, PPE_1 could be solubilized, partially or
fully, by all these detergents (Fig. 3C). In contrast, PPE_2 could not be solubilized by any of these
detergents, with the exception of SDS. Together, we conclude that the majority of the two *esx-5* locus-
encoded PPE proteins of *M. xenopi* localizes to the cell envelope of *M. smegmatis*, where they are
surface exposed. While a subpopulation of the PPE proteins associates with the cell surface in a Tween-
80/Genapol X-080 sensitive manner, a substantial amount is more firmly attached to the cell surface.
The observation that the second PPE protein cannot be extracted from the cell envelope with
detergents that are able to solubilize inner membrane proteins, a characteristic that is also observed
for known or predicted integral outer membrane proteins (Heinz & Niederweis, 2000; Speer *et al*,
2015), indicates that this protein is integrated into the mycobacterial outer membrane. This protein
could therefore potentially form a channel that mediates protein transport across the second
membrane of the mycobacterial cell envelope.

Discussion

By using a functionally reconstituted ESX-5 system in *M. smegmatis*, which naturally lacks this system, we were able to study the role of each *esx-5* gene in component stability, assembly of the membrane embedded machinery and system functionality. The obtained results are summarized in Table 1.

Deletion of individual membrane components showed varying degrees of defects in membrane complex assembly, while functionality of the ESX-5 system in all these mutants was completely abolished. An *eccD$_5$* mutant showed the most striking phenotype. Here, membrane complex assembly was completely abolished, highlighting its role as a scaffold for the assembly of the inner membrane complex. This mutant also exhibited highly affected protein levels, not only for the remaining membrane components but also for cytosolic chaperones and substrates. The fact that the *esx-5* gene cluster is introduced in *M. smegmatis* that lacks ESX-5 itself makes it unlikely that these reduced levels are caused on a transcriptional level by a negative feedback loop, as has been described for the ESX-1 system (Abdallah *et al*, 2019; Bosserman *et al*, 2017; Chen *et al*, 2016). It suggests that the absence of EccD$_5$ triggers a domino-like effect on the stability of the other proteins, starting with an unassembled inner membrane machinery, which leads to degradation of the remaining membrane components and, in the absence of the membrane complex, also of the components found in the cytosol, hereby destabilizing the substrates.

Neither substrate nor chaperone mutants showed a defect in assembly of the membrane complex and complex assembly required only the five conserved membrane components. Nevertheless, conformational changes which cannot be accounted for by the methods used here could still occur. While functionality was tightly connected with the successful assembly of the membrane complex, the minimal functional T7SS ESX-5 unit requires the five membrane components, EspG$_5$, both PE/PPE pairs that are encoded by the *esx-5* locus and an Esx heterodimer. EspG$_5$ proved to be critical for successful secretion of both PE/PPE and EsxM/N proteins, similar to previous findings (Korotkova *et al.*, 2014). The system without EccA$_5$ was still functional, although to a lower extent. EccA proteins have been shown to be required for secretion in some but not all studies (Bottai *et al*, 2012; Joshi *et al*, 2012; Phan *et al.*, 2018).

In addition, we showed that secretion of the Esx heterodimers by ESX-5 is dependent on the locus encoded PE/PPE proteins. This is in line with a recent observation for ESX-1 in *Mycobacterium marinum*, where secretion of EsxA is dependent on the co-secretion of PPE68 that is encoded by the same gene cluster(Cronin *et al.*, 2022; Damen *et al.*, 2022). Notably, PPE68 was observed to be degraded on the cell surface upon export. Since PE31/PPE18 was not able to complement secretion of

the Esx pair, our results are in line with a model in which Esx and PE/PPE proteins are specifically paired

for their co-secretion. The observation that both *esx-5* encoded PE/PPE proteins and EsxM/EsxN are

required for PPE18-HA secretion shows that both substrate classes are necessary for ESX-5

functionality.

Substrate interdependence for secretion, which has been observed also in other T7SSs (Cronin

et al., 2022; Damen *et al.*, 2022), can be explained by previously proposed models in which the

membrane complex exhibits two separate recognition events, one between NBD3 of EccC and EsxB and

one mediated by the linker2 domain of EccC with PE/PPE proteins (Bunduc *et al.*, 2020b; Rosenberg *et*

al., 2015). Substrate interdependency for secretion can also be a phenomenon taking place beyond the

inner membrane. As secretion of all three heterodimeric substrates across the full cell envelope is

dependent on the reconstituted ESX-5 system in our *M. smegmatis* model organism, all components

required for secretion across the cell envelope, including the outer membrane, are likely encoded on

the *esx-5$_{xen}$* plasmid. Excluding inner membrane components and cytosolic chaperones leaves only the

substrates to be responsible for driving secretion over the outer membrane. Indeed, both PPE

substrates, but not EsxN, localize to the cell envelope fraction and are surface exposed, implying they

are embedded in the outer membrane. Furthermore, especially PPE$_2$, showed a phenotype consistent

with known mycobacterial outer membrane proteins, in that it cannot be readily solubilized by

detergents. Combined with their essential role in secretion of the Esx heterodimer and the exogenous

PE31/PPE18 pair, it suggests a new role for these specific PPE proteins in the transport of proteins

across the mycolic acid containing layer.

Based on the combined results, we propose two working models for the role of these *esx-5-*

encoded PPE proteins. In the first model, the roles of the two PPE proteins in secretion are different,

considering the differences in membrane association. Here, the first PPE protein could play a role in

activation of the inner membrane complex and co-secretion together with the EsxM/EsxN heterodimer,

similar as has been proposed for the ESX-1 dependent Esx and PE/PPE substrates (Cronin *et al.*, 2022;

Damen *et al.*, 2022) , while the second PPE protein could form the channel for protein translocation

over the outer membrane. In the second model, the two PPE proteins form the outer membrane

channel together, with PPE$_2$ creating the integral part and PPE$_1$ forming a peripheral portion of the

complex. As these PPE proteins are specific for *M. xenopi*, further studies are required to determine

whether the *esx-5*-associated PPE proteins of other mycobacterial species, such as *M. tuberculosis*,

have similar properties.

Materials and methods

Bacterial strains and growth conditions

E. coli Dh5α was grown in LB media supplemented with appropriate antibiotics at 37°C and 200 RPM. *M. smegmatis* MC^{2}155 was grown in LB media supplemented with 0.5% Tween 80 (Merck) and appropriate antibiotics at 37°C and 90 RPM. Antibiotics were used with the following concentrations: hygromycin 50 mg/l, streptomycin 30 mg/ml.

Molecular cloning

Cloning was performed using *E. coli* Dh5α and restriction enzymes from New English Biolabs. Polymerase chain reaction (PCR) products were amplified with IProof DNA polymerase from BioRad and fused in multi-product reactions with InFusion HD from TakaraBio. For construction of plasmid derivatives, genes were deleted, either completely if they were in a single operon or ~100-150 bp were left at the C-terminus, if these were in an operon with other down- and/or upstream genes. Generally, genes were deleted using the WT ESX-5$_{Mxe}$ plasmid as template using the closest unique restriction sites up- and downstream of the gene of interest and amplification of two or more PCR fragments with compatible 18 bp ends, where the gene of interest was omitted. Because for different deletions the same restriction sites were used, some primers were used for construction of multiple derivatives. For a complete overview of restriction enzymes, primers, plasmids and cloning strategies see Table EV 1, Table EV 2 and Table EV 3.

Protein secretion and immunoblot analysis

M. smegmatis MC^{2}155 with plasmids coding for ESX-5$_{Mxe}$ WT or its derivatives were grown in LB liquid medium with 0.05% Tween 80 and appropriate antibiotics at 37°C and 90 RPM until mid-log phase. Cells were sub-cultured at an OD$_{600}$ of 0.05 in LB liquid media under the same conditions for 12-16 hours until an OD$_{600}$ of 0.8-1 was reached. Cultures were spun down, supernatants were passed through 0.2 μm filters and precipitated with trichloroacetic acid (TCA), resulting in culture supernatant fractions. Cell pellets were broken via bead-beating with glass beads, resulting in whole cell lysates or bacterial pellet fractions. SDS loading buffer was added to all samples, boiled and loaded on SDS-PAGE gels (10% or 16% depending on the protein size), transferred to nitrocellulose membranes and stained with appropriate antibodies. Antibodies against EccA$_5$, EccB$_5$, EccD$_5$, EccE$_5$, EspG$_5$, FtsH, MycP$_5$, EsxN (Mtb9.9) and GroEL2 have been described previously (Houben *et al.*, 2012b); HA-, FLAG and Strep-antibodies were purchased from Thermo Scientific (2-2.2.14), Sigma-Aldrich (F-1804) and Novusbio (NBP2-41073).

Cell envelope isolation

M smegmatis MC^{2}155 with the various ESX-5$_{Mxe}$ plasmids were grown as above to an OD$_{600}$ of ~1.5. Cells were washed in PBS, resuspended in buffer (50 mM Tris-HCl, 300 mM NaCl and 10% glycerol) and broken by passing two times through a high-pressure homogenizer at 0.3 kbar (Stansted). Unbroken cells were pelleted at 5000xg. Cell envelopes were separated from the soluble fraction by ultracentrifugation at 150.000xg. After ultracentrifugation, pellets were washed and resuspended in buffer and snap-frozen in liquid nitrogen. Where stated, after culture harvesting, cells were incubated with mixing at room temperature with 0.5% Genapol X-80 for 30 minutes, to remove cell surface proteins of bacteria. Subsequently, cells were centrifuged at 5000xg, washed one time with buffer and fractionated as explained above.

BN-PAGE

For BN-PAGE analysis of membrane complexes, cell envelopes were solubilized with 0.25% DDM for one hour at 4°C. Non-solubilized material was pelleted by centrifugation at 100.000xg for 20 minutes. The resulting supernatant samples were run on 3-12% NativePage Bis-Tris Protein Gels (Invitrogen) with addition of NativePage 5% G-250 Sample Additive (Invitrogen). Gels were blotted to PVDF membrane and stained with appropriate antibodies.

Flow cytometry analysis

M. smegmatis expressing different plasmids was grown to an OD600 of 0.8 to 1.2 in LB with or without Tween-80. Bacteria were pelleted, washed with PBS with 1% bovine serum albumin (BSA; Sigma), and incubated for 2 h with antibodies recognizing the FLAG-tag (F-1804; Sigma-Aldrich), Strep-tag (NBP2-41073; Novusbio) or influenza virus hemagglutinin tag (2-2.2.14; Thermo Scientific). After washing with PBS with 1% BSA, bacteria were incubated with secondary goat anti-mouse IgG (A-21235; ThermoFischer) or secondary goat anti-rabbit IgG (A-21244; ThermoFischer) conjugated to Alexa 647 antibodies for 30 min. After washing with PBS with 1% BSA, bacteria were analyzed by flow cytometry (Attune NxT; ThermoFisher). As a control, bacteria were incubated only with secondary antibodies.

Acknowledgements

We thank Roy Ummels for technical help and advice. This work received funding by a VIDI grant (864.12.006; to C.M.B. and E.N.G.H.) from the Netherlands Organization of Scientific Research. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement no. 101030373 (to C.M.B.). This

project was supported by funds available to T.C.M. through the Behörde für Wissenschaft, Forschung

und Gleichstellung of the city of Hamburg at the Institute of Structural and Systems Biology at the

University Medical Center Hamburg–Eppendorf (UKE). The laboratory of T.C.M. is supported by DESY

(German Electron Synchrotron Center).

Conflict of interest

The authors declare no conflict of interests.

References

Abdallah AM, Gey van Pittius NC, Champion PA, Cox J, Luirink J, Vandenbroucke-Grauls CM, Appelmelk BJ, Bitter W (2007) Type VII secretion--mycobacteria show the way. *Nat Rev Microbiol* 5: 883-891

Abdallah AM, Weerdenburg EM, Guan Q, Ummels R, Borggreve S, Adroub SA, Malas TB, Naeem R, Zhang H, Otto TD *et al* (2019) Integrated transcriptomic and proteomic analysis of pathogenic mycobacteria and their esx-1 mutants reveal secretion-dependent regulation of ESX-1 substrates and WhiB6 as a transcriptional regulator. *PLoS One* 14: e0211003

Ates LS (2020) New insights into the mycobacterial PE and PPE proteins provide a framework for future research. *Mol Microbiol* 113: 4-21

Ates LS, Ummels R, Commandeur S, van de Weerd R, Sparrius M, Weerdenburg E, Alber M, Kalscheuer R, Piersma SR, Abdallah AM *et al* (2015) Essential Role of the ESX-5 Secretion System in Outer Membrane Permeability of Pathogenic Mycobacteria. *PLoS Genet* 11: e1005190

Beckham KS, Ciccarelli L, Bunduc CM, Mertens HD, Ummels R, Lugmayr W, Mayr J, Rettel M, Savitski MM, Svergun DI *et al* (2017) Structure of the mycobacterial ESX-5 type VII secretion system membrane complex by single-particle analysis. *Nat Microbiol* 2: 17047

Beckham KSH, Ritter C, Chojnowski G, Ziemianowicz DS, Mullapudi E, Rettel M, Savitski MM, Mortensen SA, Kosinski J, Wilmanns M (2021) Structure of the mycobacterial ESX-5 type VII secretion system pore complex. *Sci Adv* 7

Bosserman RE, Nguyen TT, Sanchez KG, Chirakos AE, Ferrell MJ, Thompson CR, Champion MM, Abramovitch RB, Champion PA (2017) WhiB6 regulation of ESX-1 gene expression is controlled by a negative feedback loop in <i>Mycobacterium marinum</i>. *Proceedings of the National Academy of Sciences* 114: E10772-E10781

Bottai D, Di Luca M, Majlessi L, Frigui W, Simeone R, Sayes F, Bitter W, Brennan MJ, Leclerc C, Batoni G *et al* (2012) Disruption of the ESX-5 system of Mycobacterium tuberculosis causes loss of PPE protein secretion, reduction of cell wall integrity and strong attenuation. *Mol Microbiol* 83: 1195-1209

Bunduc CM, Bitter W, Houben ENG (2020a) Structure and Function of the Mycobacterial Type VII Secretion Systems. *Annu Rev Microbiol* 74: 315-335

Bunduc CM, Fahrenkamp D, Wald J, Ummels R, Bitter W, Houben ENG, Marlovits TC (2021) Structure and dynamics of a mycobacterial type VII secretion system. *Nature* 593: 445-448

Bunduc CM, Ummels R, Bitter W, Houben ENG (2020b) Species-specific secretion of ESX-5 type VII substrates is determined by the linker 2 of EccC5. *Mol Microbiol* 114: 66-76

Champion PA, Stanley SA, Champion MM, Brown EJ, Cox JS (2006) C-terminal signal sequence promotes virulence factor secretion in Mycobacterium tuberculosis. *Science* 313: 1632-1636

Chen Z, Hu Y, Cumming BM, Lu P, Feng L, Deng J, Steyn AJ, Chen S (2016) Mycobacterial WhiB6 Differentially Regulates ESX-1 and the Dos Regulon to Modulate Granuloma Formation and Virulence in Zebrafish. *Cell Rep* 16: 2512-2524

Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon SV, Eiglmeier K, Gas S, Barry CE, 3rd *et al* (1998) Deciphering the biology of Mycobacterium tuberculosis from the complete genome sequence. *Nature* 393: 537-544

Costa TR, Felisberto-Rodrigues C, Meir A, Prevost MS, Redzej A, Trokter M, Waksman G (2015) Secretion systems in Gram-negative bacteria: structural and mechanistic insights. *Nat Rev Microbiol* 13: 343-359

Cronin RM, Ferrell MJ, Cahir CW, Champion MM, Champion PA (2022) Proteo-genetic analysis reveals clear hierarchy of ESX-1 secretion in Mycobacterium marinum. *Proc Natl Acad Sci U S A* 119: e2123100119

Daleke MH, Ummels R, Bawono P, Heringa J, Vandenbroucke-Grauls CM, Luirink J, Bitter W (2012) General secretion signal for the mycobacterial type VII secretion pathway. *Proc Natl Acad Sci U S A* 109: 11342-11347

Damen MPM, Meijers AS, Keizer EM, Piersma SR, Jiménez CR, Kuijl CP, Bitter W, Houben ENG
(2022) The ESX-1 Substrate PPE68 Has a Key Function in ESX-1-Mediated Secretion in Mycobacterium
marinum. *mBio* 13: e0281922

Damen MPM, Phan TH, Ummels R, Rubio-Canalejas A, Bitter W, Houben ENG (2020)
Modification of a PE/PPE substrate pair reroutes an Esx substrate pair from the mycobacterial ESX-1
type VII secretion system to the ESX-5 system. *J Biol Chem* 295: 5960-5969

Dulberger CL, Rubin EJ, Boutte CC (2020) The mycobacterial cell envelope - a moving target.
Nat Rev Microbiol 18: 47-59

Ekiert DC, Cox JS (2014) Structure of a PE-PPE-EspG complex from Mycobacterium tuberculosis
reveals molecular specificity of ESX protein secretion. *Proc Natl Acad Sci U S A* 111: 14758-14763

Famelis N, Rivera-Calzada A, Degliesposti G, Wingender M, Mietrach N, Skehel JM, Fernandez-
Leiro R, Bottcher B, Schlosser A, Llorca O *et al* (2019) Architecture of the mycobacterial type VII
secretion system. *Nature* 576: 321-325

Gey van Pittius NC, Sampson SL, Lee H, Kim Y, van Helden PD, Warren RM (2006) Evolution
and expansion of the Mycobacterium tuberculosis PE and PPE multigene families and their association
with the duplication of the ESAT-6 (esx) gene cluster regions. *BMC Evol Biol* 6: 95

Heinz C, Niederweis M (2000) Selective extraction and purification of a mycobacterial outer
membrane protein. *Anal Biochem* 285: 113-120

Hoffmann C, Leis A, Niederweis M, Plitzko JM, Engelhardt H (2008) Disclosure of the
mycobacterial outer membrane: cryo-electron tomography and vitreous sections reveal the lipid
bilayer structure. *Proc Natl Acad Sci U S A* 105: 3963-3967

Houben D, Demangel C, van Ingen J, Perez J, Baldeon L, Abdallah AM, Caleechurn L, Bottai D,
van Zon M, de Punder K *et al* (2012a) ESX-1-mediated translocation to the cytosol controls virulence
of mycobacteria. *Cell Microbiol* 14: 1287-1298

Houben EN, Bestebroer J, Ummels R, Wilson L, Piersma SR, Jimenez CR, Ottenhoff TH, Luirink
J, Bitter W (2012b) Composition of the type VII secretion system membrane complex. *Mol Microbiol*
86: 472-484

Joshi SA, Ball DA, Sun MG, Carlsson F, Watkins BY, Aggarwal N, McCracken JM, Huynh KK,
Brown EJ (2012) EccA1, a component of the Mycobacterium marinum ESX-1 protein virulence factor
secretion pathway, regulates mycolic acid lipid synthesis. *Chem Biol* 19: 372-380

Korotkova N, Freire D, Phan TH, Ummels R, Creekmore CC, Evans TJ, Wilmanns M, Bitter W,
Parret AH, Houben EN *et al* (2014) Structure of the Mycobacterium tuberculosis type VII secretion
system chaperone EspG5 in complex with PE25-PPE41 dimer. *Mol Microbiol* 94: 367-382

Korycka-Machala M, Pawelczyk J, Borowka P, Dziadek B, Brzostek A, Kawka M, Bekier A,
Rykowski S, Olejniczak AB, Strapagiel D *et al* (2020) PPE51 Is Involved in the Uptake of Disaccharides
by Mycobacterium tuberculosis. *Cells* 9

Målen H, Pathak S, Søfteland T, de Souza GA, Wiker HG (2010) Definition of novel cell envelope
associated proteins in Triton X-114 extracts of Mycobacterium tuberculosis H37Rv. *BMC Microbiol* 10:
132

Phan TH, Ummels R, Bitter W, Houben EN (2017) Identification of a substrate domain that
determines system specificity in mycobacterial type VII secretion systems. *Sci Rep* 7: 42704

Phan TH, van Leeuwen LM, Kuijl C, Ummels R, van Stempvoort G, Rubio-Canalejas A, Piersma
SR, Jimenez CR, van der Sar AM, Houben ENG *et al* (2018) EspH is a hypervirulence factor for
Mycobacterium marinum and essential for the secretion of the ESX-1 substrates EspE and EspF. *PLoS*
Pathog 14: e1007247

Poweleit N, Czudnochowski N, Nakagawa R, Trinidad DD, Murphy KC, Sassetti CM, Rosenberg
OS (2019) The structure of the endogenous ESX-3 secretion system. *Elife* 8: e52983

Pym AS, Brodin P, Brosch R, Huerre M, Cole ST (2002) Loss of RD1 contributed to the
attenuation of the live tuberculosis vaccines Mycobacterium bovis BCG and Mycobacterium microti.
Mol Microbiol 46: 709-717

Rosenberg OS, Dovala D, Li X, Connolly L, Bendebury A, Finer-Moore J, Holton J, Cheng Y,
Stroud RM, Cox JS (2015) Substrates Control Multimerization and Activation of the Multi-Domain
ATPase Motor of Type VII Secretion. *Cell* 161: 501-512
Serafini A, Boldrin F, Palu G, Manganelli R (2009) Characterization of a Mycobacterium
tuberculosis ESX-3 conditional mutant: essentiality and rescue by iron and zinc. *J Bacteriol* 191: 6340-
6344
Siegrist MS, Unnikrishnan M, McConnell MJ, Borowsky M, Cheng TY, Siddiqi N, Fortune SM,
Moody DB, Rubin EJ (2009) Mycobacterial Esx-3 is required for mycobactin-mediated iron acquisition.
Proc Natl Acad Sci U S A 106: 18792-18797
Simeone R, Bobard A, Lippmann J, Bitter W, Majlessi L, Brosch R, Enninga J (2012) Phagosomal
rupture by Mycobacterium tuberculosis results in toxicity and host cell death. *PLoS Pathog* 8:
e1002507
Speer A, Sun J, Danilchanka O, Meikle V, Rowland JL, Walter K, Buck BR, Pavlenok M, Holscher
C, Ehrt S *et al* (2015) Surface hydrolysis of sphingomyelin by the outer membrane protein Rv0888
supports replication of *Mycobacterium tuberculosis* in macrophages. *Mol Microbiol* 97: 881-897
van der Wel N, Hava D, Houben D, Fluitsma D, van Zon M, Pierson J, Brenner M, Peters PJ
(2007) M. tuberculosis and M. leprae translocate from the phagolysosome to the cytosol in myeloid
cells. *Cell* 129: 1287-1298
van Winden VJC, Bunduc CM, Ummels R, Bitter W, Houben ENG (2020) A Chimeric EccB-MycP
Fusion Protein is Functional and a Stable Component of the ESX-5 Type VII Secretion System
Membrane Complex. *J Mol Biol* 432: 1265-1278
van Winden VJC, Ummels R, Piersma SR, Jimenez CR, Korotkov KV, Bitter W, Houben EN (2016)
Mycosins Are Required for the Stabilization of the ESX-1 and ESX-5 Type VII Secretion Membrane
Complexes. *mBio* 7
Wang Q, Boshoff HIM, Harrison JR, Ray PC, Green SR, Wyatt PG, Barry CE, 3rd (2020) PE/PPE
proteins mediate nutrient transport across the outer membrane of Mycobacterium tuberculosis.
Science 367: 1147-1151
Wittig I, Beckhaus T, Wumaier Z, Karas M, Schagger H (2010) Mass estimation of native
proteins by blue native electrophoresis: principles and practical hints. *Mol Cell Proteomics* 9: 2149-
2161

Table 1. Overview of results.

Construct	Component/substrate protein levels				Complex formation	Secretion	
	Membrane	Cytosolic	EsxN	PPE18		EsxN	PPE18
WT	++	++	++	++	++	++	++
Δ$espG_5$	++	++	+	-	++	-	-
Δ$eccA_5$	++	++	+	+	++	+	+
Δ$eccB_5$	++	++	+	+	-	-	-
Δ$eccC_5$	+	++	+	+	-	-	-
Δ$eccD_5$	-	+/-	-	+/-	-	-	-
Δ$eccE_5$	+	+/-	+	+	+	-	-
Δ$mycP_5$	++	++	+	+	+	-	-
MC	++	n.a.	n.a.	+	++	n.a.	-
Δ$esxM/N$	++	++	n.a.	+	++	n.a.	-
Δpe/ppe	++	++	+	+	++	-	-
Δ$substrates$	++	++	n.a.	+	++	n.a.	-

568   -, entirely affected; +/-, significantly affected; +, slightly affected; ++, not affected; n.a, not applicable

 Figure legends

Figure 1. Protein expression and membrane complex formation for ESX-5 mutants.

(A) Genetic organization of the *esx-5* locus of *M. xenopi* RIVM700367 and several of its derivatives.

Depicted are the complete gene cluster, the cluster derivatives, in which more than one gene was

deleted, and the gene organization for the *pe/ppe* region complementation plasmid. (B) SDS-PAGE

and immunoblot analysis of cell envelope fractions of *M. smegmatis* carrying the ESX-5$_{Mxe}$ plasmid or

its derivatives using antibodies reactive against the four membrane components EccB$_5$, EccD$_5$, EccE$_5$

and MycP$_5$ or antibodies directed against a Strep-tag linked to the membrane component EccC$_5$.

Antibodies against the unrelated membrane protein FtsH were used as loading control. Notably,

MycP$_5$ is processed in one of the loops of its protease domain (van Winden *et al.*, 2016), resulting in

two immuno-reactive bands (arrow heads). The high hydrophobicity of EccD$_5$ compromises its

biochemical analysis, resulting in its variable detection by SDS-PAGE and immunoblotting. (C) SDS-

PAGE and immunoblot analysis of whole cell lysates of *M. smegmatis* carrying the ESX-5$_{Mxe}$ plasmid or

its derivatives using antibodies reactive against the cytosolic ESX-5 components EccA$_5$ and EspG$_5$.

Antibodies against GroEL2 were used as loading control. (D) Model of the ESX-5 system depicting all

components and substrates (left) and BN-PAGE and immunoblot analysis of DDM-solubilized cell

envelope fractions of *M. smegmatis* carrying the ESX-5$_{Mxe}$ plasmid or its derivatives using antibodies

against EccB$_5$ or the Strep-tag introduced on EccC$_5$ (right). Asterisks denote the ~500 kDa subcomplex

seen in the *eccC$_5$, eccE$_5$* and *mycP$_5$* mutants. MOM, mycobacterial outer membrane; IM, inner

membrane.

Figure 2. Secretion analysis of *M. smegmatis* carrying the ESX-5$_{Mxe}$ plasmid or its derivatives.

(A) SDS-PAGE and immunoblot analysis of secreted fractions (culture supernatants) and whole cell

lysates (bacterial pellets) using antibodies against EsxN. In addition, an HA-antibody was used to

detect PPE18-HA that was encoded together with PE31, its partner protein, on a separate plasmid.

Antibodies against GroEL2 were used as lysis and loading control for the whole cell lysates and

antibodies against Apa were used as loading control for the secreted fraction. (B) Secretion analysis

of *M. smegmatis* carrying the ESX-5$_{Mxe}$ Δ*pe/ppe* plasmid and various *pe/ppe* complementation vectors.

The used bacterial fractions and antibodies are the same as under A.

Figure 3. Subcellular localization of the tagged PPE substrates of the *esx-5$_{Mxe}$* locus expressed in *M.*

smegmatis

(A) Secretion analysis of *M. smegmatis* carrying the ESX-5$_{Mxe}$ plasmid expressing tagged substrates.

SDS-PAGE and immunoblot analysis of secreted fractions (culture supernatants) and whole cell lysates

(bacterial pellets) of *M. smegmatis* carrying the ESX-5$_{Mxe}$ plasmid encoding the first PPE substrate

(PPE$_1$) with a FLAG-tag at the N-terminus, the second PPE protein (PPE$_2$) with a Strep-tag at the C-

terminus and EsxN with an HA-tag at the C-terminus. Antibodies against GroEL2 were used as lysis

control. Cells were grown in the presence or absence of Tween-80, after which they were harvested

and half was treated with Genapol X-080 to extract surface proteins. P, bacterial pellet; S, culture

supernatant; GP, Genapol pellet; GS, Genapol supernatant. Background bands are marked with

asterisks. (B) Cells expressing the *M. xenopi esx-5* locus encoding substrates with tags were grown in

the presence of 0.05% Tween 80, after which cells were fractionated into total (T), cytosol (Cyt) and

cell envelope (CE) fractions. Fractions were loaded in equal amounts. Antibodies against FtsH (inner

membrane protein) and GroEL2 (cytosolic protein) were used as fractionation control. (C) Surface

exposure of substrates on whole cells was measured by flow cytometry of cells grown in the presence

of Tween-80 without (top) or with (bottom) surface protein extraction with Genapol X-080 after

culture harvesting. (D) SDS-PAGE and immunoblot or Coomassie brilliant blue (CBB) stain analysis of

cell envelope fractions isolated from cells treated with Genapol X-080 and solubilized with the

mentioned detergents. Concentrations for the detergents were: 0.25% DDM, 1% n-Octyl-β-D-

Glucopyranoside, 2% Triton X-100, 2% SDS, 0.5% Genapol X-080 and 5% CHAPSO. Fractions were

loaded in equal amounts. Antibodies against EccB$_5$ and FtsH were used as inner membrane controls.

S, soluble fraction; P, non-soluble fraction.

Expanded View Figure 1. Subcellular fractionation of *M. smegmatis* carrying the *esx-5$_{Mxe}$* plasmid

expressing tagged ESX-5 substrates.

(A) Cells expressing the *M. xenopi esx-5* locus encoding substrates with tags at the N or C- termini

were grown in the presence of 0.05% Tween 80, after which supernatants (S) were separated from

bacterial cells. The cells were subsequently fractionated into total (T), cytosol (Cyt) and cell envelope

(CE) fractions. Fractions were loaded in a ratio of 2:1:1:1 for S:T:Cyt:CE. Antibody against FtsH (inner

membrane protein) was used as fractionation control. (B) Surface exposure of substrates on whole

cells was measured by flow cytometry of cells grown in the presence of Tween-80.

Expanded View Figure 2. Secretion analysis of the *esx-5*-encoded PPE proteins by *M. smegmatis*

carrying the *esx-5$_{Mxe}$* Δ*eccC$_5$* plasmid. SDS-PAGE and immunoblot analysis of secreted fractions (S) and

whole cell lysates (P) of *M. smegmatis* carrying the ESX-5$_{Mxe}$ plasmid encoding the first PPE substrate

(PPE$_1$) with a FLAG-tag at the N-terminus, the second PPE protein (PPE$_2$) with a Strep-tag at the C-

terminus and EsxN with a HA-tag at the C-terminus and the same plasmid with *eccC₅* deleted.

Antibodies against GroEL2 were used as lysis control. Cells were grown in the presence of Tween-80.

Expanded View Table 1. List of plasmids used in this study

Expanded View Table 2. Primers used in this study for molecular cloning

Expanded View Table 3. Overview of molecular cloning strategies for generating the plasmids shown

in Table EV1 and using primers shown in Table EV2.

Figure 1

Figure 2

A

Bacterial pellets | Culture supernatants

kDa — WT, ΔespG₅, ΔeccA₅, ΔeccB₅, ΔeccC₅, ΔeccD₅, ΔeccE₅, ΔmycP₅, MC, ΔesxM/N, Δpe/ppe, Δsubstrates

GroEL2

HA (PPE18)

EsxN

Apa

1 2 3 4 5 6 7 8 9 10 11 12   13 14 15 16 17 18 19 20 21 22 21 22

B

Bacterial pellets | Culture supernatants

Δpe/ppe | Δpe/ppe

kDa — WT, -, whole, Δpe/ppe₁, Δmethylt., Δpe/ppe₂

GroEL2

EsxN

Apa

1 2 3 4 5 6   7 8 9 10 11 12

Figure 3

Expanded View Figure 1

Expanded View Figure 2

Expanded View Table 1: List of plasmids used in this study.

Nr.	Construct	Source
1.	pMV *esx-5*$_{Mxe}$	Beckham *et al.*, 2017
2.	pMV *esx-5*$_{Mxe}$ Δ*eccB*$_5$	this study
3.	pMV *esx-5*$_{Mxe}$ Δ*eccC*$_5$	this study
4.	pMV *esx-5*$_{Mxe}$ Δ*pe/ppe*	this study
5.	pMV *esx-5*$_{Mxe}$ Δ*esxM/N*	this study
6.	pMV *esx-5*$_{Mxe}$ Δ*substrates*	this study
7.	pMV *esx-5*$_{Mxe}$ Δ*espG*$_5$	this study
8.	pMV *esx-5*$_{Mxe}$ Δ*eccD*$_5$	this study
9.	pMV *esx-5*$_{Mxe}$ Δ*mycP*$_5$	van Winden *et al.*, 2020
10.	pMV *esx-5*$_{Mxe}$ Δ*eccE*$_5$	this study
11.	pMV *esx-5*$_{Mxe}$ Δ*eccA*$_5$	this study
12.	pMV *esx-5*$_{Mxe}$ membrane components (MC)	this study
13.	pSMT3 *pe/ppe* whole	this study
14.	pSMT3 *pe/ppe* Δ*pe/ppe*$_1$	this study
15.	pSMT3 *pe/ppe* Δmethyltransferase	this study
16.	pSMT3 *pe/ppe* Δ*pe/ppe*$_2$	this study
17.	pMV *esx-5*$_{Mxe}$ *ppe*$_1$ FLAG N-term	this study
18.	pMV *esx-5*$_{Mxe}$ *ppe*$_1$ FLAG C-term	this study
19.	pMV *esx-5*$_{Mxe}$ *ppe*$_2$ Strep N-term	this study
20.	pMV *esx-5*$_{Mxe}$ *ppe*$_2$ Strep C-term	this study
21.	pMV *esx-5*$_{Mxe}$ *esxN-HA*	this study
22.	pMV *esx-5*$_{Mxe}$ *ppe*$_1$ FLAG N-term, *ppe*$_2$ Strep C-term, *esxN-HA*, No Strep on *eccC*$_5$	this study
23.	pMV *esx-5*$_{Mxe}$ *ppe*$_1$ FLAG N-term, *ppe*$_2$ Strep C-term, *esxN-HA*, Δ*eccC*$_5$	this study

Expanded View Table 2: List of primers used in this study.

Nr.	Primer name	Primer sequence 3' -5'
1.	ESX-5 xen del.EccB 1 FW	TGTTTGCTTGTCGGACACCT
2.	ESX-5 xen del.EccB 1 RV	CTATTTGAGTCACGATCCCCGTTC
3.	ESX-5 xen del.EccB 2 FW	GATCGTGACTCAAATAGGCACCATGGGTGGCACTG
4.	ESX-5 xen del.EccB 2 RV	TCCCATTGCGGGTCTTCGAT
5.	ESX-5 xen del.EccC FW	GTACGCGAAGAACCACTTAATTAATGC
6.	ESX-5 xen del.EccC RV	GAGGTGCCTGCCTTAAGTCACTTCGGTACCGCCAATTC
7.	Del EspG xen FW	GGCGGGGGCTTCTCTAGATG
8.	Del EsxM/N xen 1 rv	TGCGGTTGGTTGTTACTGAATATCTCCTTAGCTCGAAATCGGCTG
9.	Del EsxM/N xen 2 fw	TAACAACCAACCGCAGGCGC
10.	Del EsxM/N xen 2 rv	CATCTCGACCGAATACGTAAGTGAAAG
11.	Xen ESX-5 del sub. INF. FW	CGAGAAGTAAAGCTCTTAAGTAACAACCAACCGCAGGCGCG
12.	Del EspG xen FW	GGCGGGGGCTTCTCTAGATG
13.	Del EspG xen RV	CATCTCGACCGAATACgtaGGCTGCTCCCATGTTGGCGTG
14.	Del EccE clean 1 FW	GATCCTGCGCGGCCGCTCAT
15.	Del EccE clean 1 RV	CAGCCGTGGGCGTCGTCACGTCATGGCTGCCTCCCTTGTCG
16.	Del EccE clean 2 FW	CGTGACGACGCCCACGGCTGGTG
17.	Del EccE clean 2 RV	CGTCGACATCGATAAGCTTTGG
18.	Del EccA xen FW	GCCATCTGCCGGACGCGTG
19.	Del EccA xen RV	GTCGACATCGATAAGCTTCATCGGGCCGCCGCCGGG
20.	Xen whole PE/PPE inf FW	GAGGAATCACGCTAGCGATGTCGTTCGTGACC
21.	Xen whole PE/PPE inf RV	AGATATCCATGGATCCTAACCGGCGTAGACC
22.	Xen tag substr 1 FW	CGAGAAGTAAAGCTCTTAAGGCAGGCAC
23.	Xen 1st PPE flag C 1 RV	TGTCGTCATCGTCTTTGTAGTCGCCGGCGAACGGTGGGCGTG
24.	Xen 1st PPE Flag C 2 FW	ACAAAGACGATGACGACAAGTAGCAACACGCGAAGCGCCG
25.	Xen tag substr 2 RV	TGCATCTCGACCGAATACGTAAGTGAAA
26.	Xen 2nd PPE Strep C 1 RV	TTCAAATTGGGGATGCGACCATGATCCTCCACCGGCGTAGACCGGGCG
27.	Xen 2nd PPE Strep C 2 FW	TCGCATCCCCAATTTGAAAAATAGGAAAGGGGGCCAGCCG
28.	Xen EsxN-HA 1 RV	TAATCAGGAACATCATACGGATAGGCCCAGCTGGAGCCGAC
29.	Xen EsxN-HA 2 FW	CGTATGATGTTCCTGATTATGCTTAACAACCAACCGCAGGCGC
30.	Xen 1st PPE flag Nterm 1 RV	CGAACCTCCCGAACCTCCCTTGTCGTCATCGTCTTTGTAGTCCATTAGTACTCTCCTTAGCCCGCCG
31.	Xen 1st PPE Flag Nterm 2 FW	GGAGGTTCGGGAGGTTCGGTGCTGGACTTCGGGGCATTTC
32.	Xen 2nd PPE Strep Nterm 1 RV	CGAACCTCCCGAACCTCCTTTTTCAAATTGGGGATGCGACCACATCTAGAGAAGCCCCCGCC
33.	Xen 2nd PPE Strep Nterm 2 FW	GGAGGTTCGGGAGGTTCGTTGGATTTCGGGCTGTTTCCG
34.	Rem Str EccC5xen FW	CCGACCTGCGCCGGTTTAAATAGCTTAAGGCAGGCACCTCTTGAG
35.	Rem Str EccC5xen RV	CTCAAGAGGTGCCTGCCTTAAGCTATTTAAACCGGCGCAGGTCGG

Expanded View Table 3: Overview of molecular cloning strategies for generating the plasmids shown in Expanded View Table 1 and using primers shown in Expanded View Table 2.

Nr.	Construct	Starting backbone	Restriction sites used to cut the backbone	Primers used	PCR product size	Comments
2.	pMV esx-5_Mxe ΔeccB_5	pMV esx-5_Mxe	EcoRV, SexAI	1, 2	330 bp	Restricted backbone and the two PCR products were ligated with InFusion HD.
				3, 4	1140 bp	
3.	pMV esx-5_Mxe ΔeccC_5	pMV esx-5_Mxe	PacI, AflII	5, 6	2174 bp	Restricted backbone and the PCR product were ligated with InFusion HD.
4.	pMV esx-5_Mxe Δpe/ppe	pMV esx-5_Mxe	AflII, XbaI			Restriction removes pe/ppe genes. Resulting backbone was recircularized.
5.	pMV esx-5_Mxe ΔesxM/N	pMV esx-5_Mxe	Xbai, SnaBI	7, 8	1356 bp	Restricted backbone and the two PCR products were ligated in InFusion HD.
				9, 10	1061 bp	
6.	pMV esx-5_Mxe Δsubstr.	pMV esx-5_Mxe	AflII, SnaBI	11, 10	1082 bp	Restricted backbone and the PCR product were ligated with InFusion HD.
7.	pMV esx-5_Mxe ΔespG_5	pMV esx-5_Mxe	XbaI, SnaBI	12, 13	2053 bp	Restricted backbone and the PCR product were ligated with InFusion HD.
8.	pMV ESX-5xen ΔeccD_5	pMV esx-5_Mxe	SnaBI, NdeI			Restriction removes eccD_5. Resulting backbone was recircularized.
10.	pMV ESX-5xen ΔeccE_5	pMV esx-5_Mxe	NdeI, HindIII	14, 15	2068 bp	Restricted backbone and the two PCR products were ligated with InFusion HD.
				16, 17	2607 bp	
11.	pMV ESX-5xen ΔEccA5	pMV esx-5_Mxe	MluI, HindIII	18, 19	1129 bp	Restricted backbone and the PCR product were ligated with InFusion HD.
12.	pMV ESX-5xen MC	pMV esx-5_Mxe ΔEccA5	AflII, SnaBI			Restriction removes all substrate genes and espG_5. Resulting backbone was recircularized.
13.	pSMT3 pe/ppe whole	pSMT3 empty		20, 21	4583 bp	Backbone and PCR product were ligated with InfusionHD.
14.	pSMT3 pe/ppe Δpe/ppe_1	pSMT3 pe/ppe whole	NheI			Restriction removes 1st pe/ppe genes. Resulting backbone was recircularized.
15.	pSMT3 pe/ppeΔmethyltransf.	pSMT3 pe/ppe whole	NheI, BamHI			Partial restriction to delete methyltransferase. Resulting backbone was recircularized.
16.	pSMT3 pe/ppe Δpe/ppe_2	pSMT3 pe/ppe whole	BamHI			Restriction remove 2nd pe/ppe genes. Resulting backbone was recircularized.
17.	pMV esx-5_Mxe ppe_1 FLAG N-term	pMV esx-5_Mxe	AflII, SnaBI	22, 30	688 bp	Restricted backbone and the two PCR products were ligated with InFusion HD.
				31, 25	6035 bp	
18.	pMV esx-5_Mxe ppe_1 FLAG C-term	pMV esx-5_Mxe	AflII, SnaBI	22, 23	1934 bp	Restricted backbone and the two PCR products were ligated with InFusion HD.
				24, 25	4769 bp	
19.	pMV esx-5_Mxe ppe_2 Strep N-term	pMV esx-5_Mxe	AflII, SnaBI	22, 32	3716 bp	Restricted backbone and the two PCR products were ligated with InFusion HD.
				33, 25	3005 bp	
20.	pMV esx-5_Mxe ppe_2 Strep C-term	pMV esx-5_Mxe	AflII, SnaBI	22, 26	4910 bp	Restricted backbone and the two PCR products were ligated with InFusion HD.
				27, 25	1802 bp	
21.	pMV esx-5_Mxe esxN-HA	pMV esx-5_Mxe	AflII, SnaBI	22, 28	5621 bp	Restricted backbone and the two PCR products were ligated with InFusion HD.
				29, 25	1802 bp	
22.	pMV esx-5_Mxe ppe_1 FLAG N-term, ppe_2 Strep C-term, esxN-HA, no Strep on eccC_5.	pMV esx-5_Mxe ppe_1 Flag N-term	AflII, SnaBI	22, 26	4910 bp	Restricted backbone and the two PCR products were ligated with InFusion HD into an intermediate plasmid.
		pMV esx-5_Mxe esxN-HA	AflII, SnaBI	27, 25	1802 bp	
		Intermediate plasmid	DraI, AflII	34, 35		Restricted intermediate plasmid was religated with InFusion HD with the overlapping primers to swap the TwinStrep at the C-terminus of eccC_5 with a stop codon.
23.	pMV esx-5_Mxe ppe_1 FLAG N-term, ppe_2 Strep C-term, esxN-HA, ΔeccC_5	pMV esx-5_Mxe ppe_1 FLAG N-term, ppe_2 Strep C-term, esxN-HA, No Strep on eccC_5.	PacI, AflII	5, 6	2174 bp	Restricted backbone and the PCR product were ligated with InFusion HD.