The effects of *de novo* mutation on gene expression and the consequences for fitness in *Chlamydomonas reinhardtii*

Running Title: Transcriptomic effects of *de novo* mutation and the fitness consequences

Eniolaye J. Balogun[1,2] and Rob W. Ness[1]

1. Department of Biology, William G. Davis Building, University of Toronto, Mississauga, L5L-1C6, Canada
2. Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, M5S-3B2, Canada

Keywords: Gene expression, mutation, evolution, *Chlamydomonas reinhardtii*

Abstract

Mutation is the ultimate source of genetic variation, the bedrock of evolution. Yet, predicting the consequences of new mutations remains a challenge in biology. Gene expression provides a potential link between a genotype and its phenotype. But the variation in gene expression created by *de novo* mutation and the fitness consequences of mutational changes to expression remain relatively unexplored. Here, we investigate the effects of >2600 *de novo* mutations on gene expression across the transcriptome of 28 mutation accumulation lines derived from two independent wild-type genotypes of the green algae *Chlamydomonas reinhardtii*. We observed that the amount of genetic variance in gene expression created by mutation (V_m) was similar to the variance that mutation generates in typical polygenic phenotypic traits and approximately 15-fold the variance seen in the limited species where V_m in gene expression has been estimated. Despite the clear effect of mutation on expression, we did not observe a simple additive effect of mutation on expression change, with no linear correlation between the total expression change and mutation count of individual MA lines. We therefore inferred the distribution of expression effects of new mutations to connect the number of mutations to the number of differentially expressed genes (DEGs). Our inferred DEE is highly L-shaped with 95% of mutations causing 0-1 DEG while the remaining 5% are spread over a long tail of large effect mutations that cause multiple genes to change expression. The distribution is consistent with many *cis*-acting mutation targets that affect the expression of only one gene and a large target of *trans*-acting targets that have the potential to affect tens or hundreds of genes. Further evidence for *cis*-acting mutations can be seen in the overabundance of mutations in or near differentially expressed genes. Supporting evidence for *trans*-acting mutations comes from a 15:1 ratio of DEGs to mutations and the clusters of DEGs in the co-expression network, indicative of shared regulatory architecture. Lastly, we show that there is a negative correlation with the extent of expression divergence from the ancestor and fitness, providing direct evidence of the deleterious effects of perturbing gene expression.

Introduction

Mutations provide both the raw variation for evolution through adaptation and contribute towards disease and aging. Unfortunately, a challenge in predicting the effects of mutations is linking the genotypic consequences of a mutation to phenotypic change. Gene expression provides a potentially useful link where we can observe changes in the regulation of genes due to mutation. Moreover, there is a longstanding hypothesis that evolution may proceed primarily through alteration in gene expression rather than changes in protein structure (King and Wilson 1975). For this to be true, mutations altering

expression must either be more common, have less harmful effects or be more beneficial than protein-

coding mutations. Though there are now numerous examples of regulatory changes driving both adaptive

changes (e.g. Abzhanov et al. 2006) and harmful diseases (e.g. Kleinjan and van Heyningen 2005), we

still lack a general description of how variation in gene expression originates from *de novo* mutation and

the distribution of effects of these mutations on fitness.

Comparative research has revealed numerous patterns in gene expression variation that provide insights

into the impact of mutation on gene regulation (reviewed in Hill et al. 2021). In general, essential genes,

genes typically with high expression and those that are central in molecular interaction networks tend to

display reduced expression variation within and between species (Jeong et al. 2001; Pál et al. 2001; Yu et

al. 2004; Hahn and Kern 2005; He and Zhang 2006; Liao et al. 2006; Yu et al. 2007; Zotenko et al. 2008).

An obvious explanation is that stabilizing selection on expression level removes deleterious variants and

reduces the overall segregating expression variation. However, the distribution of mutational effects on

expression (DEE) may also influence the extent of variation in different gene classes if mechanisms have

evolved that buffer essential genes from perturbations by mutation. It has been observed that within

species variation is driven by *trans*-regulatory variants, defined by regulatory mutations that affect distal

genes. In contrast, *cis*-regulatory mutations, which affect proximal genes, underlie between-species

variation (Wittkopp et al. 2004; Wittkopp et al. 2008; Graze et al. 2009; Signor and Nuzhdin 2018). This

trend might be explained if *trans*-regulatory mutations are more common, due to a larger mutation target.

The more genes affected, the increasing likelihood of reducing fitness (Zande et al. 2022). With this in

mind, we may also expect *trans*-regulatory variants to be kept at low frequencies and thus not contribute

to between species divergence. Critically, the above explanations all depend on the rate, target size,

expression effect and fitness impact of regulatory mutations, highlighting the importance of studying the

effect of mutation on expression.

Some of the most detailed information on how *de novo* mutation affects gene expression comes from

experiments using single gene models where the effect of mutations on a reporter gene can be studied in

detail. For example, in *S. cerevisiae, cis*-acting mutations in the *TDH3* promoter were shown to generally

reduce expression and have a greater effect on expression than *trans*-acting mutations, which did not

show an asymmetry in the direction of effect (Metzger et al. 2016). However, although *trans*-acting

mutations were relatively weak, they were 265 times more likely to occur than *cis*-acting mutations,

implying a much larger mutational target (Gruber et al. 2012; Metzger et al. 2016). Despite the valuable

insights gained from such precise experiments, these systems can not capture the diverse mechanisms that

may govern the effects of natural mutations on expression throughout the transcriptome. For example,

some genes have redundant promoters and regulatory proteins that maintain expression in the presence of

mutational perturbations (reviewed by Payne and Wagner 2015; Signor and Nuzhdin 2018). Network

feedback may also serve to mute the effects of mutation on expression of genes, but such effects would

likely be most evident when genes native to the genome in question are investigated. Additionally, there

is evidence that *trans*-acting eQTLs may be clustered in hotspots, affecting thousands of genes (Albert et

al. 2018), which are not observable in single gene model systems. There is therefore good reason to

investigate the effect of *de novo* mutation on the transcriptome.

Investigating spontaneous mutations has been constrained by the rarity with which new mutations arise.

Mutation accumulation (MA) experiments overcome this limitation by evolving lines under minimal

selection, therefore facilitating the build-up of spontaneous changes irrespective of their fitness (Muller

1928). MA has been used extensively in the estimation of the rate, spectrum, and fitness effects of

mutations. However, there are surprisingly few studies investigating the expression changes in MA lines

in a small number of model organisms including *S. cerevisiae*, *D. melanogaster* and *C. elegans* (Denver

et al. 2005; Rifkin et al. 2005; Landry et al. 2007; McGuigan et al. 2014; Huang et al. 2016; Zalts and

Yanai 2017). A key parameter in measuring how mutation affects gene expression is the contribution of

mutation to expression variation per generation *i.e.,* mutational variance (V_m) (Lynch and Walsh 1998).

From the studies formerly mentioned, the mutational variance of *C. elegans* (Denver et al. 2005) was up

to 9-fold higher than that of *D. melanogaster* and (Rifkin et al. 2005) and yeast (Landry et al. 2007). MA

studies have also shown that mutation introduced far more variation in expression than would be expected

based on standing expression variation in natural populations, suggesting strong stabilizing selection in

nature (Lemos et al. 2004; Denver et al. 2005; Rifkin et al. 2005; Huang et al. 2016). We have also

learned that the expression of genes likely involved in fitness-related traits and pivotal developmental

stages are highly constrained (Zhang and Li 2004; Rifkin et al. 2005; Zalts and Yanai 2017). Moreover,

there is evidence of an inverse correlation between V_m and expression level (Rifkin et al. 2005). Despite

these breakthroughs (Hine et al. 2018; Hodgins-Davis et al. 2019), our current knowledge of the global

effects of mutations across the transcriptome is restricted to a small number of organisms with sometimes

limited technologies and often in the absence of information about the underlying genomic mutations.

Here, we investigate the effect of *de novo* mutation on gene expression and fitness in *Chlamydomonas*

reinhardtii. This haploid green alga is a longstanding model for numerous aspects of cell, molecular, and

genome biology (Harris 2001; Grossman et al. 2003) and has emerged as a model for studying the

evolution of mutation in eukaryotes. Over the past decade, numerous studies have explored mutation rate

(Ness et al. 2012; Sung et al. 2012; Ness et al. 2015; Ness et al. 2016; Flynn et al. 2018), base spectrum,

and the predictors of mutational variation across the genome of *C. reinhardtii* and related species (Ness et

al. 2015; López-Cortegano et al. 2021). We also know that on average mutations in *C. reinhardtii* are

deleterious, but evidence from the genetic mapping of mutational effects demonstrates a significant

fraction of spontaneous mutations are beneficial (Morgan et al. 2014; Böndel et al. 2019). However, in

this system, it remains unexplored how the accumulation of mutations has altered patterns of gene

expression. Here, we sequenced the transcriptomes of 28 well-studied MA lines derived from two

independent wild-type genotypes with fully characterized mutations and known fitness (Morgan et al.

2014; Ness et al. 2015; Böndel et al. 2019). Each MA line was grown for 800 – 1100 generations and

carries approx. 76 mutations. Integrating new transcriptome data with detailed characterization of these

lines, we address the following questions: (1) How much does *de novo* mutation influence the direction

and overall variation in gene expression? (2) What is the distribution of mutational effects on expression?

(3) What are the properties of genes that are more or less robust to gene regulation perturbation by

mutation (4) Do MA lines with more transcriptome divergence from mutation also suffer greater fitness

declines?

Methods

Sample collection and growth

We examined gene expression in 28 mutation accumulation lines derived from two wild-type *C.*

reinhardtii strains: CC-2391 (n=13) and CC-2344 (n=15) (Morgan et al. 2014; Ness et al. 2015; Böndel et

al. 2019). These MA lines have been intensively studied, including the characterization of the location of

each mutation; the relative fitness of each MA line with respect to their unmutated ancestral line; the

mutation rate of each line; and the distribution of fitness effects of all mutations. The MA lines were

grown for 800-1100 generations and carry a mean of 76.2 mutations in CC-2344 and 112.1 mutations in

CC-2931. To measure expression change from *de novo* mutation, we grew three replicates of each MA

line and their unmutated ancestors synchronously on Bold's agar medium at 25ºC under standard,

uniform lab conditions. Each replicate was grown until the density of colonies was sufficient for RNA

isolation, then frozen at -80ºC until extraction. RNA extraction was conducted via the Maxwell RSC 48

Instrument and assessed for degradation, purity, and quantity using gel electrophoresis and a Qubit

fluorometer. mRNA isolation and library preparation were conducted with the NEB mRNA stranded

library preparation kit and Illumina NovaSeq preparation. All samples were sequenced using Illumina

100bp paired-end sequencing with at least 20 M high-quality reads per sample. Sequencing and library

preparation were performed by Genome Quebec.

Sequence read filtering and quality control

Sequence quality of raw reads was assessed with FastQC (*v0.11.9*) (Andrews 2010) and aligned using

STAR aligner (*v2.7.8a*) (Dobin et al. 2013). We used the *C. reinhardtii* v6.0 reference genome with the

mt+ mating type allele, and organelle genomes included (Craig et al. 2022). STAR alignment was

conducted in a two-pass mode, where the first pass identified possible splice junctions (SJ), and the

second pass used the reference gene annotation in conjunction with the SJ output. Parameters of both

alignments included "--alignIntronMax 5000" to exclude known false large introns identified in other

STAR-mapped BAM files and "--outFilterMismatchNoverLmax 0.2", which allowed 0.2 x 100 bp

mismatches for each single read to account for *C. reinhardtii's* high genetic diversity (~3%). The

resulting BAM files were sorted and indexed with samtools (Li et al. 2009). Sequence coverage and

mapping quality was measured with bamqc and rnaseq from the Qualimap quality control package

(*v2.2.2a*) (Okonechnikov et al. 2016). Lastly, transcript quantification was done with featurecounts

(*v2.0.0*) (Liao et al. 2013).

Differentially expressed gene (DEG) calling

Read count normalization was conducted by DESeq2 (*v1.30.1*) (Love et al. 2014) using the median of

ratios method to account for differing sequencing depth across samples. Low read counts were filtered by

DESeq2 using the mean of normalized read counts across all samples within a strain (*i.e.,* ancestors and

replicates) as a filter statistic that optimizes statistically significant differentially expressed gene calling.

We retained ~ 95% of all genes across samples. We found a high expression correlation between the

biological replicates of all samples using FPKM (fragments per kilobase of exon per million) normalized

read counts of each gene (Pearson's $R > 0.96$, $p < 1.0$ x 10^{-4}). Of the 90 replicates, only two had a lower

correlation with their respective replicates (CC-2344 L6 replicate 2, Pearson's $R \sim 0.78$, $p < 1.0$ x 10^{-4};

CC-2931 L3 replicate 3, Pearson's $R \sim 0.80$, $p < 1.0$ x 10^{-4}). These replicates were removed from all

analysis dependent on averaged normalized read counts, but all replicates were kept for calling

differentially expressed genes (DEGs) given DESeq2 accounts for variance among replicates. DEGs in

each MA line were identified with respect to its unmutated ancestral line with DESeq2. DESeq2 uses the

Wald's test, $p_{adj} < 0.05$ for DEG calling, and the Benjamini-Hochberg method to account for multiple

testing (*FDR < 0.05*). Across all MA lines, we identified 23,557 DEGs in CC-2344 (mean ~1571) and

15,829 DEGs in CCC-2391 (mean ~1218).

Quantifying expression variance introduced by de novo mutation

To estimate the mutational variance, we used a generalized linear mixed model with a Poisson distribution to find the between line variation across MA lines.

$$\ln(Y_{ijk}) = \beta_1 \text{generation}_{ij} + u_{1i}\sqrt{generation}_{ij} + b_k + \ln(\frac{1}{size\ factor\ i}) + \varepsilon_{ijk} \qquad (1)$$

The model was fitted for each gene separately, where the between line variance per generation is given by the variance of u_{1i} ($2\text{Var}(u_{1i}) \sim$ mutational variance, V_m) (Lynch and Walsh 1998). The MA line was treated as a random effect $u_{1i} \sim N(0, \sigma^2_u)$ to account for per line variation given the MA lines evolved independent of each other. Since we know the total number of mutations in each line, we were able to estimate the increase in expression variation per *de novo* mutation (V_{mm}) by substituting the number of generations with the mutation count in equation 1. In our model, Y_{ijk} is the raw read count of gene j, line i and replicate k, which is normalized by a size factor $\ln(\frac{1}{size\ factor})$ to consider the varying sequencing depths across samples. Though we found a strong correlation between replicates, we also included an observation level random effect $b_k \sim N(0, \sigma^2_b)$ to account for overdispersion in our data. To estimate the mutational heritability $h^2_m = \frac{Vm}{Ve}$, the environmental variance (V_e) for each gene is given by the residual variance of the random slope $\sim \sigma^2_\varepsilon$. For subsequent analysis, all variance estimates were $\log_2$ transformed to facilitate comparison with previous studies.

To test whether lines with more mutations demonstrated larger expression divergence from the unmutated ancestor, we quantified the $\log_2$-fold change of gene expression for each gene in each MA line relative to its respective unmutated ancestor. We then summed these expression changes for all significant DEGs ($p_{adj} < 0.05$) for each MA line and plotted it against the mutation count of the respective line to get an overall measure of divergence (Fig 1). The DEGs in each MA line were divided into positive and negative expression change bins to consider any bias in the direction of change and avoid the cancelling out of expression changes. We found no correlation between expression divergence and mutation count and found similar results after plotting the sum of absolute fold change of DEGs per MA line against mutation count (Fig S1).

Inference of the distribution of expression effects (DEE) of novel mutations

The number of DEGs generated by mutation is a function of the distribution of expression effects of new mutations. Given that many mutations likely have no effect on expression, while other mutations may affect many genes (reviewed in Emerson and Li 2010), we inferred the distribution of expression effects

of mutations (DEE) by fitting a model to the number of DEGs caused by mutation in each of the 28 MA

lines. We first fit a model where the number of DEGs caused by a mutation was drawn from a gamma

distribution:

$$X \sim Gam(\theta, k),$$

where X is the number of DEGs and θ and k are the parameters for the shape and scale respectively. We

also reasoned that if many mutations caused 0 DEGs a better fitting model may allow a fraction of

mutations to cause 0 DEGs while other mutations caused 1 or more DEGs drawn from a gamma

distribution shifted by 1. To find the parameters that best optimize the likelihood of seeing all 28

mutations-DEG counts in a given distribution, we use the SciPy global minima tool dual annealing, which

was amended to find the global maximum instead of the minimum (Virtanen et al. 2020). The tool was

run for one million iterations under varying temperatures to get a good estimate of the optimum. For the

modified distribution, we split the bounds of no-effect mutations into 0.2 intervals and ran the model for

each interval. The model of best fit was determined through Akaike and Bayesian Information criteria.

Correlates of expression change

To test whether expression change depends on transcription level in the ancestral genotype, we

categorized the genes into 10 percentile bins according to the FPKM (fragments per kilobase of exon per

million)-normalized read counts of the ancestral line. The lowest bin represents genes with the lowest

read counts and those in the highest percentile bins consist of high expression genes. Each bin contains

1745 and 1736 genes for CC-2344 and CC-2931 respectively. Next, we plotted the distribution of

expression change within each percentile bin by taking the median $\log_2$ fold change of each gene across

all MA lines within a strain.

Using genes in the metabolic network (iCre1355), we tested whether the position of genes in the network,

specifically their connectedness to other genes predicted their robustness to mutational perturbations. The

degree centrality (*i.e.,* indegree, outdegree and betweenness centrality) of each gene was estimated (Imam

et al. 2015) from the iCre1355 network after conversion to a gene centric network (Chaiboonchoe et al.

2016). iCre1355 was constructed using the *C. reinhardtii* v5 gene annotation with 1355 metabolic genes,

and after liftover to the v6 annotation, 1304 metabolic genes remained.

We also tested whether genes central in the co-expression network were buffered against expression

change due to mutation. We identified hub genes in the Strenkert et al (2019) co-expression network that

measured the expression of the transcriptome at two hour intervals over a 24-hr period. Using the

log$_2$(FPKM + 1) normalized expression data retrieved from the Strenkert et al. (2019) supplementary, we

generated an adjacency matrix with a soft threshold of 2 using WGCNA (Langfelder and Horvath 2008;

Langfelder and Horvath 2012). The adjacency matrix was subsequently transformed into a topological

overlap matrix which was used to cluster coexpressed genes using hierarchical clustering analysis. Of the

33 resultant clusters, those that were correlated by 0.75 were merged to provide a total of 13 distinct

clusters. 99.9% of genes were placed within these 13 co-expression modules. Hub genes were identified

as genes with the top 5% intramodular connectivity per module. There were 890 hub genes based on the

v5 gene annotation, which translated to 774 hub genes in v6. We measured the relationship between

expression change and intramodular/total connectivity using multiple regression and compared the

median expression change in hub genes against non hub genes.

Evidence of cis- and trans-acting mutations

To identify putative *cis* mutations, we measured the number of mutations located within 100 bp of the

nearest DEG. Given the possibility of random chance neighboring mutation-DEG pairs, we use a

permutation test to identify the presence of significant mutation-DEG clustering. We compared the

number of nearby mutations to *n* observed DEGs per MA line to a null distribution generated by 10,000

iterations of nearby mutations to *n* randomly selected DEGs. The *p* value, the fraction of simulated DEGs

with a higher number of mutation-DEG clustering than our real dataset, was subsequently compared to α

= 0.05 to test for the presence of potential *cis* mutations.

To group DEGs that might arise from mutations in common regulatory elements, we considered whether

DEGs were clustered in the *Chlamydomonas* co-expression network (Strenkert et al. 2019). To test for

excess clustering, we calculated the shortest path in a network between DEGs in the same MA line using

the multi_source_dijkstra function in the NetworkX python package (Hagberg et al. 2008). Paths between

genes (*i.e.,* edges) in the co-expression network represent correlations in gene expression. Since genes can

be connected through hubs, we measured total connectedness by converting each correlation coefficient

(weight of the path) to path length using the equation: distance = correlation coefficient^{-1}. We then took

the sum of all paths between two genes to get the total distance between two DEGs in the coexpression

network. The shortest total paths represent the strongest correlations. We used a permutation test to

compare the distribution of shortest path lengths between observed DEGs for each MA line against a null

distribution generated from the shortest path lengths of *n* random genes, where *n* is the number of DEGs

in a given MA line. Lines with signs of significant clustering had shorter distances than their simulated

counterparts. A *p* value was assigned as the fraction of trials where simulated DEGs had equal or shorter

path lengths than the observed data. The function multi_source_dijkstra from NetworkX was amended to

exclude unconnected DEGs from the analysis.

Fitness of mutation accumulation lines

The fitness of all 28 MA lines were previously characterized in Morgan et al (2014). Fitness was

measured as the mean of maximum growth rate from two independent assays in Bold's medium. To

assess fitness change, all fitness measures were scaled relative to the growth rate of the unmutated

ancestor. To test for an association of fitness change with expression change, we performed a linear

regression of relative fitness against the number of DEGs per line. We performed a similar regression

using absolute fold expression change which resulted in qualitatively similar results (Fig S6).

The code used for the analysis can be found at https://github.com/enibalogun/The-effect-of-mutation-on-

gene-expression.git and the raw data at the European Genome-Phenome Archive

[PRJEB51708/ERP136361].

Results and Discussion

Mutational variance

In this study, we examined how >2600 mutations across 28 MA lines have altered gene expression in the

transcriptome of *C. reinhardtii*. On average each MA line had 1407 differentially expressed genes

(DEGs), with more DEGs in MA lines of strain CC-2344 (mean 1571 DEGs/MA line) than CC-2931

(mean 1218 DEGs/MA line). We first estimated mutational variance (V_m) (Lynch and Walsh 1998),

which describes the change in expression variation introduced into each gene in each generation due to

mutation. Measured as a $\log_2$-fold change in RNASeq read count, we found that the median V_m across

genes was 8.2×10^{-4} and 6.1×10^{-4} for CC-2344 and CC-2931 respectively. Similar to the mean number

of DEGs, the V_m per gene was higher in MA lines derived from CC-2344. The 1.3-fold elevation in CC-

2344 was unexpected given that it has a lower mutation rate than CC-2931 (Ness et al. 2015). Our

estimates of V_m for gene expression were on average 1.8✕ that of *C. elegans* (Denver et al. 2005;

Hodgins-Davis et al. 2019) and 6 - 15✕ that of *S. cerevisiae* (Landry et al. 2007; Hodgins-Davis et al.

2019). However, it is difficult to directly compare V_m between different organisms due to numerous

factors including varying experimental designs, inclusion of all genes versus only DEGs, and

technological differences between microarray and RNASeq. Even including these estimates, it is very

clear from the within and between species variation in V_m that there is a compelling reason for similar

studies in a much broader array of organisms.

Our estimates of V_m also permit calculation of mutational heritability ($h^2_m \sim \frac{vm}{ve}$, where V_e is the

environmental variance) which is useful in comparing how much variation mutation can generate across

different traits. We observed a median mutational heritability of per gene expression as 1.4×10^{-3} across

the 28 MA lines from two strains. A recent review by Conradsen et al (2022) showed that h^2_m ranged from

2.5×10^{-5} to 1.0×10^{-2} across numerous traits and taxa. Therefore, our estimate of h^2_m for gene expression

fell in the higher end of this range, but is broadly comparable to the amount of variation generated by

mutation in other largely polygenic traits. Notably, microbes and transcriptomic traits were excluded from

the review, but the fact that the h^2_m of gene expression in *C. reinhardtii* is similar to that of other traits

implies that mutation can create a substantial amount of variation for natural selection to act upon. As *C.*

reinhardtii is haploid, the effects of mutations can not be masked by heterozygosity, which may

contribute to higher estimates of h^2_m. One might assume that the mutational target size of a single gene's

expression is relatively low compared to complex life history traits or fitness, but if there are numerous

trans-acting effects caused by regulatory network feedback this may not be the case. The P_{TDH3}-YFP

reporter gene in yeast was estimated to have a *trans* mutational target size of 118 Kbp (Metzger et al.

2016). Assuming the *C. reinhardtii* single base mutation rate of 1.2×10^{-9} and 900 generations (Ness et al.

2015), this target size implies that 1 in 7.7 genes should experience a regulatory mutation during the MA

experiment, and some of these mutations may affect multiple genes.

Mutational variance is often scaled per generation, partly to facilitate estimates of trait evolution over

time, but also because the number of mutations causing the change in each line is often unknown. As

mutations are the cause of expression change and the expected number of mutations per generation varies

between individuals and across species, we may gain insight from per-mutation estimates of mutational

variance. Here, we were able to use mutation counts for each line to estimate expression change per

mutation (V_{mm}) using generalized mixed-effect linear modelling. Across genes, we found a 4.9⬚

difference in median V_{mm} of the strains. These values are both substantially higher than the V_m per

generation, which is predicted given that we only expect ~1 mutation per 10 generations in *C. reinhardtii*.

However, while the direction of change between mutation variance scaled per generation and per

mutation is expected, the magnitude is much larger than expected based on the number of mutations per

generation (V_{mm} is 243⬚ V_m). This discrepancy may in part be due to the evolution of mutator strains

among the MA lines, with 3.5⬚ - 8⬚ higher mutation rates than other MA lines from the same strain.

When we excluded these two mutators from our model, V_{mm} estimates dropped by more than half (CC-

2344 V_{mm} = 0.15; CC-2931 V_{mm} = 0.028), but not enough to explain the discrepancy between V_m and V_{mm}.

Taken together, the lack of congruence between the mutation rate of the two strains and the amount of

mutation variance for expression, as well as the relatively larger V_{mm} to V_m estimates suggests the

relationship between mutation count and the extent of expression divergence is complex, which we

explore more below.

Distribution of expression effects of mutation

Contrary to the simple prediction that more mutations would introduce greater changes in expression, we

found no linear correlation between total absolute $\log_2$-fold expression change and the number of

mutations in a MA line (Fig S1, R^2 = 0.058, p = 0.421). In fact, some MA lines with very few mutations

(CC-2344) had the most DEGs and total change in expression (Fig 1). To investigate whether there was a

bias in the direction of expression change, we separated genes into up- and down-regulated categories.

Lines that experienced large expression changes tended to accumulate a similar amount of expression

change in both directions (Pearson's R = 0.826, p = 6.03 x 10^{-8}). To ensure that the pattern between

mutation count and expression change was not driven by large fold-changes in genes with ancestrally low

expression, we also considered the relationship between mutation and read count differences (MA -

ancestral read count) but similarly found no correlation (Fig S2).

Fig 1. No correlation between total expression change and mutation count. Plot of the total expression change of

differentially expressed genes in each MA line against mutation count. Genes with positive and negative $\log_2$ fold

changes were summed separately and their total expression change was represented as points above and below the

black horizontal line. The dashed line represents the average directional rate of 2-fold expression change per

mutation. There is no correlation between expression change in either direction and the number of mutations

(positive: Pearson's $R = 0.1$, $p = 0.625$; negative: Pearson's $R = 0.08$, $p = 0.702$).

One explanation for the weak linear correlation that we see between expression change and mutation

count is that the underlying distribution of expression effects of mutation (DEE) includes a large number

of mutations with little or no effect and a long tail of rare mutations that drive large changes in

expression. We therefore attempted to infer the DEE as a gamma distribution that would describe the

number of DEGs seen in each of the 28 MA lines as a function of the number of mutations they each

harbour. We fit three models, one gamma distribution and two shifted gamma's with a preset fraction of

mutations that introduced 0 DEGs or 0 and 1 DEG. Of the three models, the best fitting distribution was

strongly right skewed with 95% of mutations causing 0-1 DEGs. According to this model with a discrete

bin of zero-effect mutations plus a gamma, 42% of mutations caused no change in expression, 53%

caused 1 DEG and 5% caused 2 or more genes to alter in expression (Fig 2). The long tail of high effect

mutations meant that the overall mean number of DEGs per mutation is 14.3. Although we likely do not

have enough data to precisely infer the shape of the DEE for mutations of large effect (those causing 2 or

more DEGs), the three models reach a consensus. Many mutations have no effect on expression, a

fraction of mutations affect a single gene, while the long tail of large effect mutations have the potential

to create many DEGs. This model fits with what we know about the effects of *cis* and *trans* regulatory

mutations with *cis* mutations affecting only local genes, while *trans*-acting mutations have the potential to

alter the expression of many genes through pleiotropy (Wittkopp 2023). The existence of a small fraction

of large effect mutation also potentially explains why some MA lines with very few mutations have very

divergent transcriptomes while others with more mutations have relatively few DEGs. In fact, when we

simulate the expected number of DEGs seen in 28 MA lines with the observed mutation counts, the

predicted number of DEGs per line is very similar to observed data (1321 DEGs/MA line predicted, 1394

DEGs/MA line observed).

Fig 2. Distribution of expression effects of *de novo* mutations. The inferred distributions describe the number of differentially expressed genes caused by a randomly selected *de novo* mutation. Three distributions were used to fit the mutation and DEG counts of all 28 MA lines. Of the three distributions, the best fitting model (green) included a category of mutations that cause no differential expression (0 DEG) and a gamma distribution shifted by one. The mean number of DEGs per mutation was 14.3 but 95% of mutations were estimated to change the expression of zero or one gene, while the remaining 5% caused 2 or more (0-DEG category =0.42, gamma shape = 0.0118, gamma scale = 2005)

An alternative explanation to our modelled DEE is that the expression of many genes are affected by more than one mutation. Each gene's expression is the sum of multiple up- and down-regulating mutations creating the possibility of compensatory mutations that would mask the individual effects of mutations. However, under this compensatory model there should be an increase in both the mean and standard deviation of expression change with increasing mutations per line - which is not what we observed.

Correlates of expression change

It is often hypothesized that genes with high expression are more important to fitness and therefore subject to stronger selective constraints. Indeed, in numerous species there is a negative correlation between expression level and the rate of protein sequence evolution (Pál et al. 2001; Pál et al. 2003;

Bloom and Adami 2004; Marais et al. 2004; Rocha and Danchin 2004; Zhang and Li 2004). We find a

similar pattern in *C. reinhardtii* with high expression genes displaying reduced sequence divergence

relative to *C. incerta* (0.12 for $K_a/K_{s \text{ high expression genes}}$ < 0.33 for $K_a/K_{s \text{ low expression genes}}$, Brunner Munzel test, *p*

= 5.41 x 10^{-104}). It is also reasonable to posit that genes with high expression may be critical to the

transcriptome and may have evolved robust expression regulation, making them less susceptible to

mutational or environmental perturbation. Contrary to this prediction we see relatively little evidence that

ancestral expression level predicts the extent of expression change due to mutation (Fig S3). Genes with

higher ancestral expression in fact show a slight elevation in fold expression change from the ancestor.

On the other hand, we find a clear correlation that higher expression genes have higher V_m and and h^2_m;

but this reflects the expected mean-variance correlation when V_m is estimated from read counts, such that

genes with more reads have more scope to vary in absolute terms. Relatively few other studies have

examined the relationship of expression level and susceptibility to mutation. Similar to our findings, there

was an overall weak positive relationship of h^2_m and expression observed in *D. melanogaster* (Fig S4), but

that was slightly negative when developmental stages were separated (Rifkin et al. 2005). So while our

results support the finding that protein evolution is slower in high expression genes, we find no evidence

that the mutational effects are buffered in these high expression genes.

In addition to expression level, the interactions among genes and their relative regulation can be

summarized in interaction networks. The centrality-lethality rule describes the concept that highly

connected genes in an interaction network are more likely to be critical to fitness (Jeong et al. 2001; Yu et

al. 2004; Hahn and Kern 2005; He and Zhang 2006; Yu et al. 2007; Zotenko et al. 2008). There is clear

support for this idea from experimental and comparative genomics (Pál et al. 2001; Rocha and Danchin

2004; Subramanian and Kumar 2004; Zhang and He 2005; Liao et al. 2006; Zhang and Yang 2015).

Extending this idea, it is reasonable that the expression level of central genes is also subject to stronger

stabilizing selection, and that mechanisms may have evolved to protect central genes from mutations that

alter their expression. Using a reconstruction of *C. reinharditii's* metabolic network, iCre1355 (Imam et

al. 2015), we tested whether highly-connected metabolic genes (*i.e.*, degree and betweenness centrality)

underwent less expression change due to mutation. Contrary to our prediction, we found that the genes

with the highest degree (those with the most connections to other genes) had a 25% higher median

absolute-fold expression change than the lowest degree genes (low degree genes = 0.35, high degree

genes = 0.425, Brunner Munzel test, *p* < 0.03). We also noted that the subset of genes included in the

metabolic network had higher expression than the genome average and therefore tended to have higher V_m

because of the same mean-variance correlation of V_m and expression noted above. When we look for

altered patterns of expression change in the regulatory network, we find a similarly significant positive

correlation of connectivity (*i.e.,* intramodular connectivity) in a regulatory module with the median

expression change of a gene although the amount of variation explained is relatively low ($R^2 = 0.04$, $p =$

0.001). The positive relationship between connectivity and expression divergence does not support the

idea that mechanisms have evolved to buffer these genes from mutational effects, but nor does it rule it

out. Given the interconnectedness of hub genes, it may be expected that they would be more prone to

perturbation and the fact that their mutation vulnerability is quite similar to other genes could be a

consequence of these buffering mechanisms. No previous mutation accumulation studies have examined

the relationship between network topology and mutational effects on expression. However, in *D.*

melanogaster (Lemos et al. 2004; Lemos et al. 2005) reduced divergence and polymorphism of

expression level was seen in highly connected proteins, and if this trend holds in *Chlamydomonas* it

would imply that there is stronger stabilizing selection removing the incoming variation from mutation.

Evidence of cis- and trans-acting mutations

The inferred DEE suggested a large fraction of mutations that cause only 1 or a small number of DEGs,

which could represent mutations in *cis*-acting regulatory regions that affect local genes. We therefore

sought to test for an over-representation of mutations near DEGs, under the simplifying assumption that

cis-mutations are near the genes they affect. To generate a null distribution of the expected distance of a

DEG to its nearest mutation, we conducted 10,000 iterations where the relative positions of mutations and

DEGs were randomized. To reflect the non-uniform distribution of mutations, we retained their position

in the genome and randomly assigned genes as DEGs. When we compared the distribution created with

observed DEGs to the null, we found an enrichment of DEGs with mutations within 100 bp of the gene

flanks in 12 of 28 MA lines (permutation test, $p < 0.05$; Fig S5) and an additional six lines with

marginally significant *p* values (permutation test, $0.05 < p < 0.1$). In Fig 3, we display a representative

sample (CC-2344 L14) showing evidence of an over-abundance of mutations near DEGs (red) relative to

the expectation if there was no relationship (blue). We also found several mutations within DEGs, which

were twice as likely to be found in UTRs as would be expected based on the proportion of UTR

sequences in the genome. This may be expected given that UTRs are enriched with promoters/enhancers

and are known to regulate RNA transcription, stability, and translation (Steri et al. 2018). We also noticed

that MA lines with many DEGs were less likely to show the enrichment of nearby mutations. This may be

due to the presence of one or more *trans*-acting mutations with multiple regulatory targets that masked the

signals of any *cis*-acting mutations that only affect one gene. Though previous experimental results

suggest that *cis*-acting mutations tend to down-regulate genes (Metzger et al. 2016), we did not see a bias

in the direction of expression change in DEGs that were <100 bp from a mutation. From single gene

reporter systems (Metzger et al. 2016), we have estimates that the *trans*-acting mutation target is 385-fold

larger than the *cis*-target for that single reporter gene. However, if *cis* acting mutations are gene-specific,

and *trans*-acting mutations are pleiotropic, then the full DEE of all genes might have a larger overall

proportion of target sites for *cis*-acting mutations and could reconcile why we see a relatively clear signal

of mutations near genes.

Fig 3. Evidence of *cis*-regulatory mutations. Distribution of distance between a DEG and its nearest mutation

for observed (red) and simulated data (blue). The observed distribution is one representative MA line and the

expected distribution is based on 10,000 trials where the positions of DEGs in the genome are randomized. The

excess of DEGs co-localizing with mutations is apparent in the excess of observed data on the left of the

distribution. Mutations found within the coding or non-coding regions (UTRs and introns) were assigned a distance

of 1.

Although the majority of the inferred DEE is small effect mutations, the long tail of mutations causing

multiple DEGs may indicate the presence of *trans*-acting sites with pleiotropic effects. It is clear from the

ratio of DEGs to mutations (mean 15.7 DEGs/mutation) that there are many mutations that affect multiple

genes, especially given that many mutations will have little or no effect on expression. We also found that

DEGs tended to be clustered near one another in the *Chlamydomonas* co-expression network (Strenkert et

al. 2019). The shortest path between pairs of DEGs from the same MA line was less than the shortest path

length between random pairs of genes in 14 out of 28 MA lines (Permutations test, $p < 0.05$). Fig 4 is a

representative plot of one MA line (CC-2344 L1), where the signal of co-expression network clustering

can be observed as an excess of short paths between DEGs. A similar pattern was seen in *C. elegans*

where gene expression changes from mutation accumulation clustered in the network, and were

interpreted as evidence of *trans*-acting mutations. It has been shown in *S. cerevisiae* that the topology of

the co-expression network can explain the pleiotropic effects of *trans*-acting mutations, in line with what

we observe. Significant clustering of DEGs could be the consequence of mutation in a common *trans*-

acting factor that controls expression in a given co-expression module or regulatory feed backs that result

from alterations to the expression or structure of related genes.

Fig 4. Clustering of DEGs in the co-expression network. Overlaid distributions of the shortest path lengths

between observed DEGs (red) and the null (blue) expectation. The null distribution was created by calculating the

shortest paths among a random sample of genes from the co-expression network. The figure above is a

representative plot of a single MA line demonstrating an excess of observed short paths among DEGs at the left of

the plot indicative of clustering (Permutations test, $p < 0.05$). 14 of 28 MA lines had a significant excess of clustered

DEGs.

Fitness consequences of expression change

It has been shown that expression is often under stabilizing selection (Denver et al. 2005; Rifkin et al.

2005; Huang et al. 2016), thus we would expect MA lines with more total expression change to equally

experience more fitness decline. Consistent with this prediction, we found a negative correlation between

the number of DEGs per MA line and fitness (Fig 5, $R^2 = 0.233$, $p = 0.032$), and a similar relationship
between the sum of absolute log$_2$-fold change and fitness (Fig S6; $R^2 = 0.2$, $p = 0.034$). It is worth noting
that the correlation was dependent on a single influential data point with high expression divergence and
low fitness at the bottom right of Fig 5 (excluding this data point, $R^2 = 0.183$, $p = 0.201$). However, there
is nothing otherwise unusual about this particular data that would justify its exclusion, so while the
relationship is not strong, more replication would be required to determine if the relationship between
fitness and expression is real. The association between declining fitness and increasing number of DEGs
implies the deleterious nature of large changes in the transcriptome due to mutation. Zande et al (2022)
had similar findings in yeast, where *trans* mutations had negative fitness effects due to pleiotropy. It is
also worth noting that these same lines have previously been analyzed, where it was seen that there was a
negative relationship between the number of mutations and fitness (Kraemer et al. 2017; Böndel et al.
2019; Böndel et al. 2022), but given that we did not seen a relationship between the amount of expression
divergence and mutation count the contribution of some deleterious mutations may have been to non-
expression related traits while others influenced expression to cause an overall decline in fitness. In
combination with our results which describe the effects of a random set of *de novo* mutations, evidence
suggests that *C. reinhardtii* is vulnerable to harmful mutations that disrupt the regulation of gene
expression.

Fig 5. Evidence of fitness decline in lines with more differentially expressed genes. Linear regression modelling

the negative relationship between the number of significant differentially expressed genes in each MA line and its

fitness relative to the unmutated ancestor ($R^2 = 0.233$, $p = 0.032$).

Taken together we see clear evidence that mutation can generate a substantial amount of genetic variation

in gene expression and this new variation is important for organismal performance. For example, it has

been a long-held belief that the contribution of mutation to genetic variance is less than 1% of the

environmental variance. But, we found that mutation may contribute significantly more to variation per

generation than formerly considered, almost 24%. This is a considerable amount of mutational variation

that is heritable, which may change the perception of the raw materials available for evolution via

selection to act upon. Using mutation accumulation lines allows us to observe the cumulative effect of

mutations from the spectrum of naturally occurring mutations. Our estimated DEE indicates that there is a

substantial mutational target for large effect mutations that pleiotropically generate expression changes

across the transcriptome. There are some lines harbouring tens of mutations showing thousands of

differentially expressed genes. It seems likely that these pleiotropic mutations are driven not only by

mutations in *trans*-acting regulators like transcription factors, but also by alterations in the balance of the

metabolism of the cell and feedbacks in the regulatory network. Despite this, we see no evidence for

mutational robustness in genes central to the metabolic and co-expression network. A limitation of our

data is that we can not link the expression changes to individual mutations and therefore can not assess

the nature of these large effect mutations. Therefore, the possibility of compensatory mutations means

that some expression change may be masked by subsequent mutations. With current technology it is not

cost-effective to map *trans*-acting mutations to their expression effects but advances in single cell

technologies may allow for such resolution in the near future.

Acknowledgments

531 We would like to thank Dr. Peter Keightley and anonymous reviewers for helpful comments on our paper.

532 We would also like to thank Dr. Jarrod Hadfield for invaluable help in estimating mutational variance.

533 This work was supported by a Natural Sciences and Engineering Research Council (NSERC) Discovery

534 grant (RGPIN/06331-2016), the Canadian Foundation for Innovation John R. Evans Leaders fund

535 (35591) to R.W.N. and an NSERC graduate scholarship to EJB.

Literature cited

Abzhanov A, Kuo WP, Hartmann C, Grant BR, Grant PR, Tabin CJ. 2006. The calmodulin pathway and
evolution of elongated beak morphology in Darwin's finches. *Nature* 442:563–567.

Albert FW, Bloom JS, Siegel J, Day L, Kruglyak L. 2018. Genetics of *trans*-regulatory variation in gene
expression. *Elife* [Internet] 7. Available from: http://dx.doi.org/10.7554/eLife.35471

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.

Bloom JD, Adami C. 2004. Evolutionary rate depends on number of protein-protein interactions
independently of gene expression level: response. *BMC Evol. Biol.* 4:14.

Böndel KB, Kraemer SA, Samuels T, McClean D, Lachapelle J, Ness RW, Colegrave N, Keightley PD.
2019. Inferring the distribution of fitness effects of spontaneous mutations in *Chlamydomonas*
reinhardtii. *PLoS Biol.* 17:e3000192.

Böndel KB, Samuels T, Craig RJ, Ness RW, Colegrave N, Keightley PD. 2022. The distribution of fitness
effects of spontaneous mutations in *Chlamydomonas reinhardtii* inferred using frequency changes
under experimental evolution. *PLoS Genet.* 18:e1009840.

Chaiboonchoe A, Ghamsari L, Dohai B, Ng P, Khraiwesh B, Jaiswal A, Jijakli K, Koussa J, Nelson DR,
Cai H, et al. 2016. Systems level analysis of the *Chlamydomonas reinhardtii* metabolic network
reveals variability in evolutionary co-conservation. *Mol. Biosyst.* 12:2394–2407.

Conradsen C, Blows MW, McGuigan K. 2022. Causes of variability in estimates of mutational variance
from mutation accumulation experiments. *Genetics* [Internet] 221. Available from:
http://dx.doi.org/10.1093/genetics/iyac060

Craig RJ, Gallaher SD, Shu S, Salomé P, Jenkins JW, Blaby-Haas CE, Purvine SO, O'Donnell S, Barry
K, Grimwood J, et al. 2022. The *Chlamydomonas* Genome Project, version 6: reference assemblies
for mating type plus and minus strains reveal extensive structural mutation in the laboratory. *Plant*
Cell [Internet]. Available from: http://dx.doi.org/10.1093/plcell/koac347

Denver DR, Morris K, Streelman JT, Kim SK, Lynch M, Thomas WK. 2005. The transcriptional
consequences of mutation and natural selection in *Caenorhabditis elegans*. *Nat. Genet.* 37:544–548.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15–21.

Emerson JJ, Li W-H. 2010. The genetic basis of evolutionary change in gene expression levels. *Philos.*
Trans. R. Soc. Lond. B Biol. Sci. 365:2581–2590.

Flynn JM, Lower SE, Barbash DA, Clark AG. 2018. Rates and Patterns of Mutation in Tandem
Repetitive DNA in Six Independent Lineages of *Chlamydomonas reinhardtii*. *Genome Biol. Evol.*
10:1673–1686.

Graze RM, McIntyre LM, Main BJ, Wayne ML, Nuzhdin SV. 2009. Regulatory divergence in
Drosophila melanogaster and *D. simulans*, a genomewide analysis of allele-specific expression.
Genetics 183:547–561, 1SI – 21SI.

Grossman AR, Harris EE, Hauser C, Lefebvre PA, Martinez D, Rokhsar D, Shrager J, Silflow CD, Stern
D, Vallon O, et al. 2003. *Chlamydomonas reinhardtii* at the crossroads of genomics. *Eukaryot. Cell*
2:1137–1150.

Gruber JD, Vogel K, Kalay G, Wittkopp PJ. 2012. Contrasting properties of gene-specific regulatory,
coding, and copy number mutations in *Saccharomyces cerevisiae*: frequency, effects, and
dominance. *PLoS Genet.* 8:e1002497.

Hagberg A, Swart P, S Chult D. 2008. Exploring network structure, dynamics, and function using
networkx. Los Alamos National Lab. (LANL), Los Alamos, NM (United States) Available from:
https://www.osti.gov/biblio/960616

Hahn MW, Kern AD. 2005. Comparative genomics of centrality and essentiality in three eukaryotic
protein-interaction networks. *Mol. Biol. Evol.* 22:803–806.

Harris EH. 2001. *Chlamydomonas* as a model organism. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*
52:363–406.

He X, Zhang J. 2006. Why do hubs tend to be essential in protein networks? *PLoS Genet.* 2:e88.

Hill MS, Vande Zande P, Wittkopp PJ. 2021. Molecular and evolutionary processes generating variation
in gene expression. *Nat. Rev. Genet.* 22:203–215.

Hine E, Runcie DE, McGuigan K, Blows MW. 2018. Uneven Distribution of Mutational Variance Across
the Transcriptome of *Drosophila serrata* Revealed by High-Dimensional Analysis of Gene
Expression. *Genetics* 209:1319–1328.

Hodgins-Davis A, Duveau F, Walker EA, Wittkopp PJ. 2019. Empirical measures of mutational effects
define neutral models of regulatory evolution in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. U.*
S. A. 116:21085–21093.

Huang W, Lyman RF, Lyman RA, Carbone MA, Harbison ST, Magwire MM, Mackay TF. 2016.
Correction: Spontaneous mutations and the origin and maintenance of quantitative genetic variation.
Elife [Internet] 5. Available from: http://dx.doi.org/10.7554/eLife.22300

Imam S, Schäuble S, Valenzuela J, López García de Lomana A, Carter W, Price ND, Baliga NS. 2015. A

refined genome-scale reconstruction of *Chlamydomonas* metabolism provides a platform for
systems-level analyses. *Plant J.* 84:1239–1256.

Jeong H, Mason SP, Barabási AL, Oltvai ZN. 2001. Lethality and centrality in protein networks. *Nature*
411:41–42.

King MC, Wilson AC. 1975. Evolution at two levels in humans and chimpanzees. *Science* 188:107–116.

Kleinjan DA, van Heyningen V. 2005. Long-range control of gene expression: emerging mechanisms and
disruption in disease. *Am. J. Hum. Genet.* 76:8–32.

Kraemer SA, Böndel KB, Ness RW, Keightley PD, Colegrave N. 2017. Fitness change in relation to
mutation number in spontaneous mutation accumulation lines of *Chlamydomonas reinhardtii.*
Evolution 71:2918–2929.

Landry CR, Lemos B, Rifkin SA, Dickinson WJ, Hartl DL. 2007. Genetic properties influencing the
evolvability of gene expression. *Science* 317:118–121.

Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network analysis. *BMC*
Bioinformatics 9:559.

Langfelder P, Horvath S. 2012. Fast R Functions for Robust Correlations and Hierarchical Clustering. *J.*
Stat. Softw. [Internet] 46. Available from: https://www.ncbi.nlm.nih.gov/pubmed/23050260

Lemos B, Bettencourt BR, Meiklejohn CD, Hartl DL. 2005. Evolution of proteins and gene expression
levels are coupled in *Drosophila* and are independently associated with mRNA abundance, protein
length, and number of protein-protein interactions. *Mol. Biol. Evol.* 22:1345–1354.

Lemos B, Meiklejohn CD, Hartl DL. 2004. Regulatory evolution across the protein interaction network.
Nat. Genet. 36:1059–1060.

Liao B-Y, Scott NM, Zhang J. 2006. Impacts of gene essentiality, expression pattern, and gene
compactness on the evolutionary rate of mammalian proteins. *Mol. Biol. Evol.* 23:2072–2080.

Liao Y, Smyth GK, Shi W. 2013. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* 30:923–930.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000
Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and
SAMtools. *Bioinformatics* 25:2078–2079.

López-Cortegano E, Craig RJ, Chebib J, Samuels T, Morgan AD, Kraemer SA, Böndel KB, Ness RW,
Colegrave N, Keightley PD. 2021. *De Novo* Mutation Rate Variation and Its Determinants in
Chlamydomonas. Mol. Biol. Evol. 38:3709–3723.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol.* 15:550.

Lynch M, Walsh B. 1998. Genetics and analysis of quantitative traits. Available from:
http://www.invemar.org.co/redcostera1/invemar/docs/RinconLiterario/2011/febrero/AG_8.pdf

Marais G, Domazet-Lošo T, Tautz D, Charlesworth B. 2004. Correlated Evolution of Synonymous and
Nonsynonymous Sites in Drosophila. *J. Mol. Evol.* 59:771–779.

McGuigan K, Collet JM, McGraw EA, Ye YH, Allen SL, Chenoweth SF, Blows MW. 2014. The nature
and extent of mutational pleiotropy in gene expression of male *Drosophila serrata*. *Genetics*
196:911–921.

Metzger BPH, Duveau F, Yuan DC, Tryban S, Yang B, Wittkopp PJ. 2016. Contrasting Frequencies and
Effects of *cis*- and *trans*-Regulatory Mutations Affecting Gene Expression. *Mol. Biol. Evol.*
33:1131–1146.

Morgan AD, Ness RW, Keightley PD, Colegrave N. 2014. Spontaneous mutation accumulation in
multiple strains of the green alga, *Chlamydomonas reinhardtii*. *Evolution* 68:2589–2602.

Muller HJ. 1928. The Measurement of Gene Mutation Rate in *Drosophila*, Its High Variability, and Its
Dependence upon Temperature. *Genetics* 13:279–357.

Ness RW, Kraemer SA, Colegrave N, Keightley PD. 2016. Direct Estimate of the Spontaneous Mutation
Rate Uncovers the Effects of Drift and Recombination in the *Chlamydomonas reinhardtii* Plastid
Genome. *Mol. Biol. Evol.* 33:800–808.

Ness RW, Morgan AD, Colegrave N, Keightley PD. 2012. Estimate of the spontaneous mutation rate in
Chlamydomonas reinhardtii. *Genetics* 192:1447–1454.

Ness RW, Morgan AD, Vasanthakrishnan RB, Colegrave N, Keightley PD. 2015. Extensive *de novo*
mutation rate variation between individuals and across the genome of *Chlamydomonas reinhardtii*.
Genome Res. 25:1739–1749.

Okonechnikov K, Conesa A, García-Alcalde F. 2016. Qualimap 2: advanced multi-sample quality control
for high-throughput sequencing data. *Bioinformatics* 32:292–294.

Pál C, Papp B, Hurst LD. 2001. Highly expressed genes in yeast evolve slowly. *Genetics* 158:927–931.

Pál C, Papp B, Hurst LD. 2003. Genomic function: Rate of evolution and gene dispensability. *Nature*
421:496–497; discussion 497–498.

Payne JL, Wagner A. 2015. Mechanisms of mutational robustness in transcriptional regulation. *Front.*
Genet. 6:322.

Rifkin SA, Houle D, Kim J, White KP. 2005. A mutation accumulation assay reveals a broad capacity for
rapid evolution of gene expression. *Nature* 438:220–223.

Rocha EPC, Danchin A. 2004. An analysis of determinants of amino acids substitution rates in bacterial
proteins. *Mol. Biol. Evol.* 21:108–116.

Signor SA, Nuzhdin SV. 2018. The Evolution of Gene Expression in *cis* and *trans*. *Trends Genet.*
34:532–544.

Steri M, Idda ML, Whalen MB, Orrù V. 2018. Genetic variants in mRNA untranslated regions. *Wiley*
Interdiscip. Rev. RNA 9:e1474.

Strenkert D, Schmollinger S, Gallaher SD, Salomé PA, Purvine SO, Nicora CD, Mettler-Altmann T,
Soubeyrand E, Weber APM, Lipton MS, et al. 2019. Multiomics resolution of molecular events
during a day in the life of *Chlamydomonas*. *Proceedings of the National Academy of Sciences*
116:2374–2383.

Subramanian S, Kumar S. 2004. Gene expression intensity shapes evolutionary rates of the proteins
encoded by the vertebrate genome. *Genetics* 168:373–381.

Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M. 2012. Drift-barrier hypothesis and mutation-rate
evolution. *Proc. Natl. Acad. Sci. U. S. A.* 109:18488–18492.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P,
Weckesser W, Bright J, et al. 2020. SciPy 1.0: fundamental algorithms for scientific computing in
Python. *Nat. Methods* 17:261–272.

Wittkopp PJ. 2023. Contributions of mutation and selection to regulatory variation: lessons from the
Saccharomyces cerevisiae TDH3 gene. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 378:20220057.

Wittkopp PJ, Haerum BK, Clark AG. 2004. Evolutionary changes in *cis* and *trans* gene regulation.
Nature 430:85–88.

Wittkopp PJ, Haerum BK, Clark AG. 2008. Regulatory changes underlying expression differences within
and between *Drosophila* species. *Nat. Genet.* 40:346–350.

Yu H, Greenbaum D, Xin Lu H, Zhu X, Gerstein M. 2004. Genomic analysis of essentiality within
protein networks. *Trends Genet.* 20:227–231.

Yu H, Kim PM, Sprecher E, Trifonov V, Gerstein M. 2007. The importance of bottlenecks in protein
networks: correlation with gene essentiality and expression dynamics. *PLoS Comput. Biol.* 3:e59.

Zalts H, Yanai I. 2017. Developmental constraints shape the evolution of the nematode mid-
developmental transition. *Nat Ecol Evol* 1:113.

Zande PV, Hill MS, Wittkopp PJ. 2022. Pleiotropic effects of *trans*-regulatory mutations on fitness and
gene expression. *Science* 377:105–109.

Zhang J, He X. 2005. Significant impact of protein dispensability on the instantaneous rate of protein
evolution. *Mol. Biol. Evol.* 22:1147–1155.

Zhang J, Yang J-R. 2015. Determinants of the rate of protein sequence evolution. *Nat. Rev. Genet.*
16:409–420.

Zhang L, Li W-H. 2004. Mammalian housekeeping genes evolve more slowly than tissue-specific genes.
Mol. Biol. Evol. 21:236–239.

Zotenko E, Mestre J, O'Leary DP, Przytycka TM. 2008. Why do hubs in the yeast protein interaction
network tend to be essential: reexamining the connection between the network topology and
essentiality. *PLoS Comput. Biol.* 4:e1000140.

Supplementary materials

Fig S1: Scatterplot of the sum of the absolute log$_2$ fold change by mutation count. Expression change
is measured as the sum of absolute log$_2$ fold change across all genes in an MA line and is plotted against
the number of mutations in that MA line. There was no significant correlation found between the two
variables $(R^2 = 0.058, p = 0.421)$.

Fig S2. No correlation between the sum of FPKM differences and mutation count. Plot of the $\log_2$
transformed summed FPKM-normalized read count differences between MA lines and their respective
unmutated ancestor. Only differentially expressed genes with $p_{adj} < 0.05$ were considered. Each MA line
is shown twice with the same mutation count, points above the y=0 line represent upregulated genes and
below the line are down-regulated genes. The dashed line represents the average rate at which expression
changes per mutation. We did not find a correlation between the non-transformed differences in
expression and mutation count (positive expression change ~ Pearson $R = -0.0825$, $p = 0.676$; negative
expression change ~ Pearson $R = -0.0607$, $p = 0.759$).

Fig S3. Ancestral expression level does not predict absolute expression change. Split-violin plot showing the distribution of genes in each FPKM-normalized ancestral expression bin ($n_{CC\text{-}2344}$ = 1745, $n_{CC\text{-}2931}$ = 1737). Each measurement within the distribution represents the median absolute expression change across all 28 MA lines per gene. There is no relationship between the ancestral expression level of a gene and its respective degree of expression change. The median expression change decreases from the 10[th] to 50/60[th] percentile expression bin and increases thereafter.

Fig S4. Slight positive correlation between mutational heritability and expression level. Scatterplot of the FPKM normalized gene expression counts of the ancestral line of each strain against its respective mutational heritability. The mutational heritability was estimated using the per generation mutational variance. There is a weak positive relationship for CC-2344 (Pearson's $R = 0.20$, $p = 1.22$ x 10^{-162}) and CC-2931 (Pearson's $R = 0.06$, $p = 7.59$ x 10^{-14}).

Fig S5. Overrepresentation of mutations within 100 bp of differentially expressed genes. Evidence of *cis*-regulatory mutations was inferred from an over-representation of mutations within 100 bp of

differentially expressed genes (DEGs). The observed distance between mutations and their nearest DEGs
was compared to the expected distribution created by randomizing the DEGs across the genome over
10000 iterations. The p values represent the fraction of trials where the cases of simulated DEGs co-
localizing with mutations was greater than those with observed DEGs. The distribution of p values is
heavily right-skewed, showing the presence of an excess of mutations in or near DEGs in most MA lines.

Fig S6. A negative correlation between relative fitness and total absolute sum of log$_2$ fold change.
Linear regression modelling the relationship between the total sum of absolute log$_2$ fold change in
differentially expressed genes and its fitness relative to the unmutated ancestor ($R^2 = 0.23$, $p = 0.034$).