No evidence for associations between brood size, gut microbiome diversity and survival in great tit (*Parus major*) nestlings

Liukkonen[1*], M., Hukkanen[2,3] M., Cossin-Sevrin[2], N., Stier[2,4,5], A., Vesterinen[2], E., Grond[6], K. & Ruuskanen[1,2], S.

[1] Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland
[2] Department of Biology, University of Turku, Turku, Finland
[3] Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland
[4] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622, Lyon, France
[5] Institut Pluridisciplinaire Hubert Curien, UMR7178, Université de Strasbourg, CNRS, Strasbourg, France
[6] Department of Biological Sciences, University of Alaska Anchorage, Anchorage, AK 99508, USA

* Corresponding author: Martta Liukkonen
email: martta.liukkonen@gmail.com or martta.a.liukkonen@jyu.fi

Keywords: avian microbiome, brood size, gut microbiome, *Parus major*, 16S rRNA gene

ABSTRACT

Background

The gut microbiome forms at an early stage, yet data on the environmental factors influencing the development of wild avian microbiomes is limited. As the gut microbiome is a vital part of organismal health, it is important to understand how it may connect to host performance. The early studies with wild gut microbiome have shown that the rearing environment may be of importance in gut microbiome formation, yet the results vary across taxa, and the effects of specific environmental factors have not been characterized. Here, wild great tit (*Parus major*) broods were manipulated to either reduce or enlarge the original brood soon after hatching. We investigated if brood size was associated with nestling bacterial gut microbiome, and whether gut microbiome diversity predicted survival. Fecal samples were collected at mid-nestling stage and sequenced with the 16S rRNA gene amplicon sequencing, and nestling growth and survival were measured.

Results

Gut microbiome diversity showed high variation between individuals, but this variation was not significantly explained by brood size or body mass. Additionally, we did not find a significant effect of brood size on body mass or gut microbiome composition. We also demonstrated that early handling had no impact on nestling performance or gut microbiome. Furthermore, we found no significant association between gut microbiome diversity and short-term (survival to fledging) or mid-term (apparent juvenile) survival.

Conclusions

We found no clear association between early-life environment, offspring condition

and gut microbiome. This suggests that brood size is not a significantly contributing

factor to great tit nestling condition, and that other environmental and genetic

factors may be more strongly linked to offspring condition and gut microbiome.

Future studies should expand into other early-life environmental factors e.g., diet

composition and quality, and parental influences.

INTRODUCTION

The digestive tract hosts a large community of different microorganisms (i.e., gut

microbiome) and is known to be a fundamental part of organismal health and a

powerful proximate mechanism affecting host performance [1-2]. The gut

microbiome has been studied across a wide range of animal taxa e.g., humans [3-5],

fish [6], and economically important species such as poultry [7], and data from wild

populations is slowly increasing [8]. Generally, a more diverse gut microbiome is

considered beneficial for individual health [9], but there are also community

structure effects that define the functionality [10]. For example, laboratory-bred mice

with a less diverse gut microbiome have a substantially lower chance of surviving an

influenza infection compared to their wild counterparts unless receiving a gut

microbiota transplant from their wild counterparts [11-12]. Moreover, gut

microbiome had been linked to host fitness and survival in the Seychelles warbler

(*Acrocephalus sechellensis*). Individuals that harbored opportunistic pathogens, i.e.,

microbes that usually do not cause disease in healthy individuals, but may become

harmful in individuals that are immunocompromised, in their gut microbiome showed higher mortality [13-14]. Therefore, understanding how gut microbiome affects fitness within and between individuals is necessary for not only understanding species survival but also evolution [15-17].

Gut microbiome forms at a young age and remains somewhat stable in adulthood as found for example in laboratory bred mice [18-20]. Disruption in the gut microbiome that leads to a microbiome imbalance at a young age could result in both short-term and long-term changes in the gut microbiome [21-22]. Of the environmental effects, diet [23], including e.g., macronutrient balance (carbohydrates, fats, amino acids) [3, 24] has been concluded to be major determinants of rat and mouse gut microbiome, and this effect has recently been seen in avian models as well [25-28]. Moreover, macronutrient balance has been linked to intestinal microbiome composition [3, 24] and the functioning of individual immune response [29-30]. However, as a large part of the prior research has focused strictly on humans or species living in controlled environments in which environmental effects on both the microbiome and host are sidelined [31-32], many species, including most birds [8], have only started to attract attention [33].

The mechanisms of bacterial colonization of the bird gut are somewhat unique as avian life-histories differ significantly from those of e.g., mammals [34]. In mammals, the offspring are exposed to bacterial colonization during vaginal birth [35] and lactation [36-37], whereas bird hatchlings are exposed to bacteria first upon hatching

[20, 38]. Few studies have investigated the possibility of bacterial colonization *in ovo*,

but results are still lacking [39]. Genetics [40-42] as well as the post-hatch

environment [20, 43-46] have a significant effect on the formation of the avian gut

microbiome. Once hatched, most altricial birds feed their young, which exposes the

hatchlings to various bacteria that originate from the parents i.e., via vertical

transmission [47]. It has also been shown that environmental factors are major

contributors in the formation of gut microbiome [48-51], one of these being the

rearing environment in the nest [44].

As early-life environment is connected to the establishment of gut microbiome,

brood size may affect gut microbiome [52]. Brood size is often associated with

parents' performance and ability to feed their young [53], and the trade-off between

offspring quality and quantity has been studied widely [54-55]. Food quantity per

nestling can decrease in enlarged broods, as parents may not be able to fully

compensate for the additional amount of food an enlarged brood requires [56-57].

For example, in great tits (*Parus major*) it has been shown that nestlings from reduced

broods may have a higher body mass [58] and tend to survive better [59].

Importantly, great tit nestling body mass has been connected to gut microbiome

diversity and composition: body mass positively correlates with gut microbiome

richness [52]. This could imply that good physiological condition and high food

availability would allow the host to have a diverse gut microbiome that promotes a

healthy gut.

Alterations in early-life gut microbiome could have long-term consequences on

individual performance [60], yet such effects have rarely been studied in wild

organisms. In wild birds, some bacterial taxa have been linked to better survival. For

example, a high abundance of bacteria in the order *Lactobacillales* of the phylum

Firmicutes is related to higher individual fitness in Seychelles warblers [14] and great

tits [61]. These bacteria are also known for the benefits for bird health in

economically important species such as poultry, in which *Lactobacilli* are used as

probiotics to boost immune functioning [62]. Besides *Lactobacillales,* gut bacteria

belonging to other genera such as *Clostridium* and *Streptococcus* are important for the

degradation of non-starch polysaccharides and for the synthesis of essential

molecules such as the short-chain fatty acids [63-64]. Short-chain fatty acids are

important in host energy metabolism [65] and therefore crucial for performance.

Changes in nestling's early-life gut microbiome could affect such key physiological

processes that could influence for example nestling body mass, which is tightly

linked to survival to fledging [58-59]. Because the gut microbiome establishes at a

young age and is less plastic later in life [18-20], gut microbiome and changes to its

richness can have long-term effects on juvenile and adult survival [21-22]. For

example, antibiotic treatment at infancy can affect the expression of genes involved

in immune system functioning and lead to long-term effects on host metabolism [20].

Moreover, changes in the rearing environment can affect individual physiology and

these effects can carry over to later stages of an individual's life such as survival to

fledging and lifetime reproductive success [66].

Here, we use an experimental approach to investigate whether brood size

manipulation influenced wild great tit nestlings' bacterial gut microbiome diversity

on day 7 post-hatch. We also investigated whether brood size influenced nestling

body mass on day 7 or on day 14 post-hatch, and if the gut microbiome predicts

short-term (i.e., survival to fledging) and mid-term (i.e., apparent juvenile) survival.

The great tit is a well-studied species in the fields of ecology and evolution, and it is

easy to monitor in the wild due to its habit of breeding in nest boxes. Great tit

nestlings' gut microbiome undergoes profound shifts during early life [52], and it

has been linked to nestling natal body mass and body size [52, 61], yet studies

focusing on gut microbiome associations with survival are still scarce. Here, we

manipulated wild great tit broods by reducing or enlarging the original brood size in

order to analyze if this affected the gut microbiome. In large broods, nestlings need

to compete for their food more [67-68], and the lower food availability could result in

a lower gut microbiome diversity. This might impair nestling body mass and fitness

prospects [13, 52]. We used a partial cross-fostering design that enabled us to

disentangle the relative contributions of genetic background, early maternal effects,

and rearing environment such as parents, nest and nestmates on gut microbiome.

Furthermore, we used an unmanipulated control group in which no nestling was

cross-fostered to control for the possible effects of moving the nestlings between

nests. For example, early human handling such as marking and weighing at day 2

post-hatch could influence gut microbiome later on. We hypothesized that 1) in

reduced broods nestlings would have a higher body mass, 2) in reduced broods

nestling gut microbiome would be more diverse than in enlarged broods, and 3)

higher gut microbiome diversity on day 7 post-hatch would increase survival to

fledging and potentially reflect apparent juvenile survival. Such knowledge could

provide new information about gut microbiome in wild passerine bird population

and how the early-life environment may associate with nestling gut microbiome,

body mass, and short-term and mid-term survival.

METHODS

Study area and species

The great tit is a small passerine bird, which breeds in secondary holes and artificial nest-boxes, making it a suitable model species. Great tits breed throughout Europe and inhabit parts of Northern Africa and Asia as well, and the breeding areas differ in environment and diet [69]. In Finland the great tit is a common species with an estimate of 1.5 to 2 million breeding pairs. They lay 6 to 12 eggs between April and May and the female incubates the eggs for 12 – 15 days. The nestlings fledge approximately 16 to 21 days after hatching. The study was conducted during the breeding season (May-July 2020) on Ruissalo island (60° 25' 59.99" N 22° 09' 60.00" E). Ruissalo island habitat is a mostly temperate deciduous forest and meadows, and some areas have small patches of coniferous trees.

Brood size manipulation experiment

Nest boxes were first monitored weekly and later daily when clutches were close to the estimated hatching date. Brood size manipulation took place on day 2 after hatching. Increases in great tit brood size can lead to lowered weight in both the nestlings and adults [70-75], and our decision on the number (i.e., +2 or -2) of manipulated nestlings (i.e,. +2 or -2) followed the cited studies. We had four treatment groups (see Fig. 1): in the 'enlarged group (henceforward called E)', we increased the brood size by two individuals that were taken from a 'reduced brood'. Correspondingly, in the 'reduced group (henceforward called R)', we decreased the

brood size by two individuals, that were added to the enlarged broods. In the

'control group (henceforward called C)', we swapped nestlings between nests but

did not change the brood size. And lastly, in the 'unmanipulated control group

(henceforward called COU)', we only weighed and collected fecal samples on day 7

but did not move the nestlings between nests. We also moved nestlings between the

reduced nests to ensure that all nests except for COU had both original and fostered

nestlings. Control nests were used to control for potential cross-fostering effects

unrelated to brood size. Additionally, in the unmanipulated control group nestlings

were not moved or weighed on day 2 in order to control for any handling effects *per*

se. This study design enabled us to test the potential impacts of handling nestlings

and swapping the nest early after hatching. We aimed to move approximately half of

the chicks in the manipulated nests, so that the number of original and the fostered

nestlings would be the same in each nest after manipulation.

Before they were moved, nestlings were weighed using a digital scale with a

precision of 0.1 g and identified by clipping selected toenails. We aimed to

add/remove nestlings that were of similar weight to avoid changing the sibling

hierarchy in the brood. The moving procedure was performed as quickly as possible

to minimize the risk of stress and the nestlings were kept in a warmed box during

transportation. For each pair of nests in the brood size manipulation experiment, we

selected nests that had a similar hatching date. In case of uneven number of nests

hatching within a day, one or three nest(s) was/were allocated to the COU group. To

avoid potential bias from hatching date, we allocated nests in any given day evenly

to each treatment. We also checked that the treatments had an equal brood size on

average i.e., we did not want to only reduce the larger clutches and enlarge the

smaller clutches. These is also a significant bias towards COU nests being later in the

season on average (Table 1).

Table 1. (A) Brood size before and after manipulation, (B) Hatching date across
treatments.

A	Brood size	Before manipulation (mean±s.d.)	After manipulation (mean±s.d.)
	Enlarged broods (E)	7.700±1.261	9.650±1.309
	Reduced broods (R)	8.375±1.637	6.375±1.637
	Control broods (C)	7.565±1.805	7.565±1.805
	Unmanipulated broods (COU)	7.810±2.112	na
	ANOVA	F3=0.987, p=0.403	

B	Hatching date	mean±s.d.
	Enlarged broods (E)	58.60±5.77
	Reduced broods (R)	59.83±6.41
	Control broods (C)	58.74±5.34
	Unmanipulated broods (COU)	63.81±4.79

B Tukey's *post-hoc* for between-group comparisons
Average hatching date

	ANOVA	F3=3.964, p=0.011 *			
Contrasts	Estimate	s.e.	t.ratio	p	
COU-C	5.070	1.70	2.983	0.019	*
COU-E	5.210	1.76	2.961	0.020	*
COU-R	3.976	1.68	2.363	0.092	
C-E	0.139	1.72	0.081	0.100	
C-R	-1.094	1.64	-0.666	0.910	
E-R	-1.233	1.70	-0.723	0.888	

222
223 (E) enlarged brood size, (R) reduced brood size, (C) control brood size, (COU) unmanipulated control brood size. Brood size
224 was successfully either reduced or enlarged by two chicks.

Figure 1. Brood size manipulation experiment schematic diagram. 2-day-old nestlings were moved between nestboxes to enlarge or to reduce original brood size (an example with brood size of seven is given). Some nests were kept as control nests (nestlings were moved but brood size remained the same) and some were kept as unmanipulated control nests (nestlings were not moved at all to test whether early-life handling affects gut microbiome). The original brood size varied between nests.

Fecal sample collection

To study the effects that brood size may have on the nestling gut microbiome and its links to individual nestling body mass, survival to fledging and apparent juvenile survival, we used a subset of data from a larger experiment (Cossin-Sevrin et al., *unpublished data*). In this subset, we use individuals from which fecal samples were collected on day 7 after hatching and analyzed for microbiome diversity and composition (C=23 nestlings/15 nests, COU=22/13, E=23/15, R=24/16) We aimed to collect two samples (one from original and one from foster nestlings) per nest. Fecal samples from the nestlings were collected gently by stimulating the cloaca with the collection tube. Samples were collected straight into a sterile 1.5 ml Eppendorf tube

to avoid possible contamination of the sample. At time of sampling, each nestling

was weighed (0.1g), and the nestlings were ringed for individual identification using

aluminum bands. The samples were stored in cool bags onsite and afterwards

moved to a -80 °C freezer for storage until DNA extraction.

Apparent juvenile survival

We monitored all study nests until fledging to measure short-term survival. On day

14 post-hatch, the sampled nestlings were weighed, and wing-length was measured

to detect if the manipulation had any effects on nestling growth. Nests were

subsequently monitored for fledging success. Additionally, we monitored our study

population for apparent juvenile survival (i.e., mid-term survival) after the breeding

season (i.e., approximately 3 months after fledging) to assess the association between

gut microbiome and post-fledging survival. We captured juvenile great tits by mist

netting during the autumn-winter 2020 at six different feeding stations that had a

continuous supply of sunflower seeds and suet blocks. Feeding stations were located

within the previously mentioned nest box population areas. For each site mist

netting with playback was conducted on three separate days during October-

November 2020 for three hours at a time, leading to a total of 69 hours of mist

netting. A total of 88 individuals from the brood size manipulation experiment were

caught, and the caught juvenile great tits were weighed, and wing length was

measured. Our catching method provides an estimate of post-fledging survival yet,

it could be slightly biased based by dispersal. In a previous study in our population

[76], none of the birds ringed as nestlings were recaptured outside the study area,

suggesting that dispersal is likely limited.

DNA extraction and sequencing

We chose two samples per nest for DNA extraction, yet in such a way that both

fledged and not-fledged nestlings would be included in the dataset. DNA was

extracted from nestling fecal samples using the Qiagen QIAamp PowerFecal Pro

DNA Kit (Qiagen; Germany) following the manufacturer's protocols. Additionally,

we included negative (RNAse and DNAse free ddH_2O) controls to control for

contamination during DNA extraction and additional controls to confirm successful

amplification during PCR. A short fragment of hypervariable V4 region in the 16S

rRNA gene was amplified using the purified DNA samples as template with the

following primers: 515F_Parada (5' - GTGYCAGCMGCCGCGGTAA -3') and

806R_Apprill (5' - GGACTACNVGGGTWTCTAAT - 3') [77-78]. PCRs were

performed in a total volume of 12 µL using MyTaq RedMix DNA polymerase

(Meridian Bioscience; Cincinnati, OH, USA). The PCR cycling conditions were as

follows: first, an initial denaturation at 95 °C for 3 minutes followed by 30 cycles of

95 °C for 45 seconds, 55 °C for 60 seconds, and 72 °C for 90 seconds, and finished

with a 10-minute extension at 72 °C. After the first round of PCR, a second round

was conducted to apply barcodes for sample identification [79]. For this, PCR cycling

conditions were as follows: first, an initial denaturation at 95 °C for 4 minutes

followed by 18 cycles of 98 °C for 20 seconds, 60 °C for 15 seconds, and 72 °C for 30

seconds, and finished with a 3-minute extension at 72 °C. We performed replicate

PCR reactions to control for errors during the amplification. Further on, the PCR

products were measured for DNA concentration with Quant-IT PicoGreen dsDNA

Assay Kit (ThermoFischer Scientific; Waltham, MA, USA) and for quality with

TapeStation 4200 (Agilent; Santa Clara, CA, USA). The samples from each of the PCR

replicates were pooled equimolarly creating two separate pools and purified using

NucleoMag NGS Clean-up and Size Select beads (Macherey-Nagel; Düren,

Germany). Finally, pooled samples were sequenced (2 x 300 bp) on the Illumina

MiSeq platform (San Diego, CA, USA) at the Finnish Functional Genomic Center at

the University of Turku (Turku, Finland).

Sequence processing

All statistical analyses were performed with R (v. 4.11.0; R Development Core Team

2021) unless otherwise stated. The demultiplexed Illumina sequence data was first

processed with Cutadapt version 2.7 [80] to remove locus-specific primers from both

R1 and R2 reads. Then, the DADA2 pipeline (v. 1.24.0; [81]) was used to filter the

reads based on quality, merge the paired-end (R1 and R2) reads, to define the single

DNA sequences i.e., Amplicon Sequence Variants (henceforward ASV), and to

construct a 'seqtab'. Seqtab is a matrix also known as otutable or readtable: ASVs in

columns, samples in rows, number of reads in each cell, using default parameter

settings. In total, our seqtab consisted of 6,929,537 high-quality reads. Reads were

assigned to taxa against the SILVA v132 reference database [82] resulting in 8658

ASVs. To control for contamination, negative DNA extraction and PCR controls

were used to identify contaminants (60 ASVs) using the decontam package (v. 1.12;

[83]) and all were removed from the dataset. Sequencing runs (replicate PCR's) were

merged using the phyloseq package (v. 1.32.0) and non-bacterial sequences (mainly

Chlorophyta) were removed from the data as they were not of interest in this study

resulting in a total of 6566 ASVs (a total of 4,107,028 high-quality reads in all

samples; mean per sample: 15,155.085; mean range per sample: 0-97,264). Singleton

reads were removed from the dataset by the DADA2 pipeline. Data was further

analyzed with the phyloseq package (v. 1.32.0; [84]), and the microbiome package (v.

1.18.0; [85]) and visualized with the ggplot2 package (v. 3.3.6; [86]).

The final dataset contained 92 samples from great tit nestlings resulting in a total of

3,161,696 reads (mean per sample: 34,366.261; mean range per sample 108 – 189,300

reads), which belonged to 6,505 ASVs. The dataset was then rarefied for alpha

diversity analyses at a depth of 5000, as this was where the rarefaction curves

plateaued (see Supplementary file 2). The rarefied dataset contained 4,791 ASVs in

88 samples. For beta diversity, the unrarefied dataset was used after confirming that

the beta diversity statistics were quantitatively similar for the rarefied and

unrarefied datasets. Bacterial relative abundances were summarized at the phylum

and genus level and plotted based on relative abundance for all phyla and genera. A

Newick format phylogenetic tree with the UPGMA algorithm to cluster treatment

groups together was used to visualize sample relatedness (see Supplementary file 3)

and was constructed using the DECIPHER (v. 2.24.0; [87]), phangorn (v. 2.8.1; [88]),

and visualized with ape (v. 5.6-2; [89]), and ggtree (v. 3.4.0; [90]) packages.

Statistical analyses

Nestling body mass

First, to analyze whether brood size manipulation affected nestling body mass in the C, E, and R treatment groups, we ran two linear mixed-effects models with the *lme4* package (v. 1.1-29; [91]). In these models we used either body mass on day 7 or 14 as the dependent variable and brood size manipulation treatment, hatching date, body mass on day 2 post-hatch and original brood size as predicting variables. Hatching date is used as a predicting variable because it is known to affect nestling body mass during the breeding season [92] and there were differences in hatching date between the COU and other treatment groups (see Table 1). We included the interaction between original brood size and brood size manipulation treatment in both models as the effect of manipulation may depend on the original brood size. For example, there could be stronger effect of enlargement in already large broods. Nest of origin and nest of rearing were used as random intercepts to control for the non-independence of nestlings sharing the same original or foster nests. Here, we did not include the COU group in the analysis because we wanted to measure the effects of treatment on nestling body mass, and only enlarged, reduced or control broods' nestlings were moved between nests.

Second, to analyze whether the actual brood size affected nestling body mass, we ran two models where we used it as a continuous dependent variable to explain body mass either on day 7 or on day 14 post-hatch. Hatching date and body mass on day 2 post-hatch were used as predicting variables and nest of origin and nest of rearing as random intercepts to control for the non-independency of samples. We

included the interaction between manipulated brood size and hatching date in the

models because the effect of brood size may depend on the hatching date. For

example, hatching date can reflect environmental conditions and large broods may

perform poorly late in the season due to poorer food availability. The COU group

was initially excluded from this model to see which of the two random effects, nest

of origin or nest of rearing, explained a larger portion of variation in the treatment

groups. In the COU group, nest of origin and nest or rearing were the same, which

meant we could not include both random effects in models where all treatment

groups were present due to the model failing to converge. Nest of origin explained

more of the variation in the first model (see Supplementary file 4) and therefore, we

used it in the full models with all treatment groups: C, COU, E and R. In these

models, nestling body mass either on day 7 and or on day 14 post-hatch was used as

a dependent variable and manipulated brood size as the explanatory variable.

Hatching date and body mass on day 2 post-hatch were set as predicting variables.

Nest of rearing was used as a random intercept to control for the non-independence

of nestlings sharing the same foster nests. The significance of factors included in the

models were tested using the F-test ratios in analysis of variance (Type III ANOVA).

Alpha diversity

For alpha diversity analyses, which measures within-sample species diversity, we

ran two linear mixed-effects models with the *lme4* package (v. 1.1-29; [91]) to

measure if either brood size manipulation or manipulated brood size as a continuous

variable were associated with gut microbiome diversity. We used two alpha

diversity metrics: the Shannon Diversity Index, which measures the number of

bacterial ASVs and their abundance evenness within a sample, and Chao1 Richness,

which is an estimation of the number of different bacterial ASVs in a sample. Both

metrics were used to check if alpha diversity results were consistent across different

metrics. Each diversity index was used as the dependent variable at a time and either

brood size manipulation treatment or manipulated brood size as a predicting

variable. In both models we included original brood size, weight on day 7 post-hatch

and hatching date as covariates.   We included interaction between brood size

manipulation treatment and original brood size as there could be a stronger effect of

enlargement in initially large broods. We also included interaction between

manipulated brood size and weight on day 7 post-hatch because effect of brood size

on microbiome may depend on nestling weight. We also tested whether alpha

diversity predicted weight on day 7 post-hatch, as weight and gut microbiome

diversity have been connected in previous studies. In this analysis we used weight

on day 7 post-hatch as the dependent variable and alpha diversity (Shannon

Diversity Index and Chao1 Richness), treatment and hatching date as predicting

variables and nest of rearing as the random effect. In these sets of models, we first

excluded the COU group to see which of the two random effects, nest of origin or

nest of rearing, explained a larger proportion of variation in the treatment groups.

Nest of rearing explained more of the variation in this model (see Supplementary file

4) and therefore, we used it in the full model with all treatment groups: C, COU, E

and R. The significance of factors included in the models were tested using the F-test

ratios in analysis of variance (ANOVA).

Short-term survival

To explore whether alpha diversity associated with survival to fledging (i.e., short-term survival) and with apparent juvenile survival in Autumn 2020 (i.e., mid-term survival), we used generalized linear models with binomial model (v. 1.1-29; lme4 package, [91]), and then tested the significance of factors with type 2 ANOVA from the car package (v. 3.0-13; [93]). Type 2 ANOVA was used because the model did not contain interaction between predicting and there was no order between covariates, as they could not be ranked. Survival to fledging and recapture in Autumn 2020 were used as the binomial response variable (yes-no) in each model. Alpha diversity (Shannon Diversity Index and Chao1 Richness) was the main predicting variable, and weight on day 7 post-hatch (same time as sampling the fecal gut microbiome), hatching date and manipulated brood size were included as covariates in the model. We did not include brood size manipulation treatment in the survival models as not enough birds from each treatment group were recorded for fledging and juvenile survival. Moreover, we excluded random effects from this model as the model failed to converge. 65 nestlings fledged successfully, while 8 nestlings were found dead in nest boxes. For 15 nestlings we had no fledging record, so these were excluded from the survival to fledging analysis. In apparent juvenile survival, 19 birds out of 92 (with data on microbiome diversity) were recaptured as juveniles. For all analyses, the R package car (v. 3.0-13; [93]) was used to test Variance Inflation Factors (VIFs) and the package DHARMa (v. 0.4.5; [94]) to test model diagnostics for linear mixed-effects and generalized linear models.

Beta diversity

For visualizing beta diversity, i.e., the similarity or dissimilarity between the

treatment group gut microbiomes, non-metric multidimensional scaling (NMDS)

was used with three distance matrices: Bray-Curtis [95], weighted UniFrac, and

unweighted UniFrac [96]. Permutational multivariate analysis of variance

(PERMANOVA) using the Euclidean distance matrix and 9999 permutations was

tested with the R package vegan (*adonis2* function; v. 2.6-2; [97]) to investigate if any

variables affected to the variation in gut microbiome composition. Nest of rearing

was set as a blocking factor in the PERMANOVA to control for the non-desirable

effects of the repeated sampling of foster siblings. The test for homogeneity of

multivariate dispersions was used to measure the homogeneity of group dispersion

values. We used the phyloseq package (v. 1.32.0; [84]) to run a differential

abundance analysis with a significance cut-off $p < 0.01$ to test the differential

abundance of ASVs between the treatment groups.

442
RESULTS

The effects of brood size manipulation on nestling body mass

Brood size manipulation did not significantly affect nestling body mass on day 7

post-hatch (ANOVA: $F_{2, 25.832} = 0.441$, p=0.648; see Supplementary file 5). Moreover,

there was no significant interaction between brood size manipulation and original

brood size (ANOVA: $F_{2, 24.610} = 0.678$, p=0.517; see Supplementary file 5). On day 14

post-hatch, brood size manipulation did not significantly affect nestling body mass

(ANOVA: $F_{2,\ 24.335}=0.831$, p=0.448; see Supplementary file 5). However, body mass

increased with increasing hatching date (ANOVA: $F_{1\ 24.070}=13.367$, p=0.001; see

Supplementary file 5). Next, we did not find any significant associations between

manipulated brood size and nestling body mass (ANOVA for weight on day 7: $F_{1,}$

$_{35.149}=1.777$, p=0.191; ANOVA for weight on day 14: $F_{1,\ 29.491}=2.156$, p=0.153; see

Supplementary file 6). Nest of origin explained a larger proportion of the variation in

weight than the nest of rearing on both day 7 (nest of origin 41.1 % and nest of

rearing 24.4 %) and day 14 (nest of origin 65.5 % and nest of rearing 21.9 %) post-

hatch, but this result was not statistically significant (Pr > $\chi 2 = 1$) (see Supplementary

file 4).

Alpha diversity

As 7-day-old nestlings, most bacterial taxa belonged to the phyla *Proteobacteria*,

Firmicutes, and *Actinobacteria* (Fig. 2).

Figure 2. Bacterial relative abundances on Phylum level across the four treatment groups. Each bar represents an individual sample. Treatment groups are control (C), unmanipulated control (COU), enlarged (E), and reduced (R). N=88 samples divided into treatment groups as follows: C=23, COU=21, E=20, R=24. Phyla with less that 10 % in relative abundance is collapsed into the category "< 10 % abundance."

Brood size manipulation did not significantly influence alpha diversity (Shannon Diversity Index) (ANOVA: $F_{3,\ 47.488}$=1.026, p=0.390, Fig. 3; see Supplementary file 7).

Moreover, original brood size (ANOVA: $F_{1,\ 50.269}$=0.388, p=0.536; see Supplementary file 7), weight on day 7 post-hatch (ANOVA: $F_{1,\ 80.551}$=0.003, p=0.959; see Supplementary file 7), and hatching date (ANOVA: $F_{1,\ 50.276}$=1.073, p=0.305; see Supplementary file 7) did not significantly associate with alpha diversity. There was no significant interaction between brood size manipulation and original brood size (ANOVA: $F_{3,\ 48.053}$=0.126, p=0.944; see Supplementary file 7). Results for Chao1 Richness were quantitatively similar: brood size manipulation did not affect alpha

diversity (ANOVA: $F_{3,\ 45.936}=0.358$ p=0.784, Fig. 3; see Supplementary file 7). Nest of

rearing explained a larger proportion of the observed variance in alpha diversity

(27.7 %) than nest of origin (10.8 %), but the result was not statistically significant (Pr

$> \chi2 = 1$) (see Supplementary file 4).

Figure 3. The gut microbiome alpha diversity of 7-day-old great tit nestlings
across the four treatment groups visualized with two diversity metrics: A)
Shannon Diversity Index and B) Chao1 Richness. The black dots represent each
observation within a treatment group. The whiskers represent 95 % confidence
intervals. Treatment groups are control (C), unmanipulated control (COU), enlarged
(E), and reduced (R). N=88 samples divided into treatment groups as follows: C=23,
COU=21, E=20, R=24.

Next, we tested whether the manipulated brood size as a continuous variable was

associated with alpha diversity (Shannon Diversity Index), but found no significant

association (ANOVA: $F_{1, 63.001} < 0.001$, p=0.984; see Supplementary file 8) in this

analysis either. Weight on day 7 post-hatch (ANOVA: $F_{1, 82.840} = 0.015$, p=0.903; see

Supplementary file 8) and hatching date (ANOVA: $F_{1, 59.734} = 0.137$, p=0.713; see

Supplementary file 8) did not correlate with alpha diversity in this model either.

There was no significant interaction between manipulated brood size and weight on

day 7 post-hatch (ANOVA: $F_{1, 82.702} < 0.000$, p=0.998; see Supplementary file 8). Results

for Chao1 Richness were quantitatively similar (ANOVA: $F_{1, 65.064} = 0.246$, p=0.622; see

Supplementary file 8): manipulated brood size did not affect alpha diversity, and

neither did weight on day 7 post-hatch (ANOVA: $F_{1\ 83.513} = 0.690$, p=0.409; see

Supplementary file 8) nor hatching date (ANOVA: $F_{1\ 57.110} = 1.133$, p=0.292; see

Supplementary file 8).

507
Alpha diversity and short/mid-term survival

Next, we explored whether alpha diversity (Shannon Diversity Index and Chao1

Richness) contributed to predicting short/mid-term survival (survival to fledging

and apparent juvenile survival). Survival to fledging was not predicted by alpha

diversity (Shannon Diversity Index: $\chi^2 = 0.010$, df=1, p=0.923; see Supplementary files

9 and 10), manipulated brood size ($\chi^2 = 0.090$, df=1, p=0.764; see Supplementary file

9), weight on day 7 post-hatch ($\chi^2 = 0.388$, df=1, p=0.533; see Supplementary file 9) or

hatching date ($\chi^2 = 0.438$, df=1, p=0.508; see Supplementary file 9).

Apparent juvenile survival was not significantly associated with alpha diversity

(Shannon Diversity Index: $\chi2$=1.916, df=1, p=0.166; see Supplementary files 9 and

10). Moreover, there was no significant interaction between alpha diversity and

manipulated brood size ($\chi2$= 1.268, df=1, p=0.260; see Supplementary file 9).

However, apparent juvenile survival was negatively associated with hatching date

($\chi2$=4.654, df=1, p=0.031; see Supplementary file 9). Additional analyses to check for

the consistency of results were tested the following way: survival to fledging with

nestlings from the COU group removed and apparent juvenile survival without the

nestlings with no recorded survival for fledging (see methods). These results were

quantitatively similar as in the whole dataset for both Shannon Diversity Index

(survival to fledging: $\chi2$= 2.285, df=1, p= 0.131; apparent juvenile survival: $\chi2$=1.515,

df=1, p=0.218; see Supplementary file 11) and Chao1 Richness (survival to fledging:

$\chi2$=0.665, df=1, p=0.415; apparent juvenile survival: $\chi2$=2.654, df=1, p=0.103; see

Supplementary file 11).

Beta diversity

Non-metric multidimensional scaling (NMDS) using weighted and unweighted

UniFrac and Bray-Curtis dissimilarity did not show clear clustering of samples based

on brood size manipulation treatment (Fig. 5; see Supplementary file 3). The test for

homogeneity of multivariate dispersions supported the visual assessment of the

NMDS (Betadispersion$_{9999 \text{ permutations}}$: $F_{3, 0.069}$= 0.650, p<0.001; see Supplementary file

12). Pairwise PERMANOVA further indicated that the treatment (PERMANOVA:

R^2=0.061, F=1.951, p=0.278; see Supplementary file 12), weight on day 7 post-hatch

(PERMANOVA: R^2=0.015, F=1.387, p=0.091) or hatching date (PERMANOVA:

R^2=0.0232, F=2.214, p=0.993) did not significantly contribute to the variation in gut

microbiome composition between the treatment groups. Differential analysis of ASV

abundance between the treatment groups showed that there is variation in taxa

abundance. E group showed higher taxa abundance when compared to COU and C

groups and was slightly higher than the R group. C and COU groups were generally

lower in taxa abundance than R and E groups, and COU group showed lower

abundance than the other groups in each comparison (see Supplementary file 13).

DISCUSSION

In this study, we investigated the associations between great tit nestling gut

microbiome, brood size, and nestling body mass by experimentally manipulating

wild great tit broods to either reduce or enlarge the original brood size. The results

show that even though there was individual variation in the nestling gut

microbiome (Fig. 2), brood size did not significantly contribute to gut microbiome

diversity. Neither did gut microbiome diversity explain short-term (survival to

nestling) nor mid-term (apparent juvenile) survival. Body mass was also not

significantly affected by brood size manipulation. The COU group that functioned as

a control for moving and handling effects, did not differ in this respect from the

other groups. This suggests that human contact or handling nestlings 2 days post-

hatch did not influence nestling gut microbiome or body mass. The partial cross-

fostering design enabled us to disentangle the relative contributions of rearing

environment (i.e., parents, nest and nestmates) from genetic, prenatal such as

maternal allocation to egg, and early post-natal effects such as feeding up to day 2.

Nest of rearing seemed to explain more of the variation in nestling gut microbiome

diversity than the nest of origin (although not statistically significant), which follows

previous studies. Contrastingly, nest of origin seemed to be a stronger contributor

than the nest of rearing on nestling body mass on day 7 and day 14 post-hatch. This

result was also not statistically significant.

Brood size manipulation and nestling body mass

First, we explored whether brood size was associated with nestling body mass, as

such changes may explain the underlying patterns in gut microbiome [52]. Against

our hypothesis, we found no significant association between nestling body mass and

brood size: neither reduction nor enlargement of the broods resulted in significant

body mass differences in the nestlings on day 7 and day 14 post-hatch. While the

result is supported by some studies in which associations between nestling body

mass and brood size have been tested [61, 98], the majority of the literature shows

that brood size negatively correlates with nestling body mass: in larger broods

nestlings are generally of lower mass [52-53, 57, 67, 99-104].

There are a few possible explanations why brood size manipulation did not affect

nestling body mass. Firstly, if environmental conditions were good, parents may

have been able to provide enough food even for the enlarged nests and thus,

variance in brood size may not result in differences in nestling body mass between

reduced and enlarged nests.   In that case the number of nestlings transferred

between enlarged and reduced nests should probably have been larger to create

differences in nestling body mass between the two treatments. Still, we think that the

decision to transfer +2/-2 was reasonable since it was based on extensive evidence

from previous studies [103]. Secondly, it could be that the enlarged brood size

negatively influences some other physiological traits while body mass was retained

at the expense of these other traits e.g., immune system functioning [105-106].

Moreover, our analysis showed that hatching date had a significant effect on nestling

body mass: nestlings that hatched later in the season were of lower weight. This

could be a result of changes in the food items that great tits use, changes in

temperature conditions or in parental investment during the breeding season. As the

season progresses, the abundance of insect taxa varies, and this can result in changes

in nutrient rich food [103, 107]. For example, great tits can select certain lepidopteran

larvae that vary in their abundance during the great tit breeding season [108].

Thirdly, it could be that the change in brood size was influencing the parents'

condition instead of the nestlings [109-110]. In enlarged broods, parents are required

to forage more which can lead to higher energy expenditure and increased stress

levels in parents [72-73, 109].

Brood size manipulation and gut microbiome

We found large inter-individual differences in gut microbiome diversity, yet this

variation was not explained by brood size or nestling body mass. It is possible that

brood size did not result in differences in food intake. For example, parents were

likely able to provide an equivalent amount of food, given that body mass was not

significantly affected by the brood size manipulation. Therefore, brood size

manipulation did not affect gut microbiome diversity through differences in nutrient

uptake. Alternatively, in this study, fecal sampling took place 5 days after the initial

brood size manipulation (day 2 post-hatch). It could be that sampling on a later date

or at multiple timepoints [61, 111] would have led to different results. Firstly, the

time interval may not have been long enough to detect effects of the brood size

manipulation. Secondly, it has been shown in previous studies that the nestling gut

microbiome undergoes profound shifts at the nestling stage: overall gut microbiome

diversity decreases but relative abundance in some taxa increases [52]. We suggest

that fecal samples could be collected on multiple days post-hatch to understand the

potential day to day changes in the nestling gut microbiome.

Our results suggest that the variance in gut microbiome is a result of other factors

than those linked to brood size. Firstly, one of these factors could be diet (i.e., food

quality) which has gained attention in gut microbiome studies during the past years

[25, 27, 112-115]. The overall diversity in gut microbiome could be explained by

adaptive phenotypic plasticity because it is sensitive to changes in the environment

e.g., changes in diet [116-117]. The food provided by the parents can vary between

broods in different environments [118], and this variation in diet can lead to

differences in gut microbiome diversity [114-119]. For example, abundance in certain

dietary items such as insects or larvae can result in lower gut microbiome diversity

than other dietary items [113-116]. As great tits have been reported to adapt their

diet along the breeding season due to changes in insect taxa frequency [103, 107] this

could affect the between-nestling and between-nest gut microbiome diversity.

However, using wild bird populations in gut microbiome studies limits the ability to

control the consumed dietary items because parents may use variable food resources

and there can be variance in dietary between sexes and even individuals. Visual

assessment of dietary items [116] and metabarcoding could be of use here as they

enable the identification of food items on genus and even species level from e.g.,

fecal samples [119].

Secondly, breeding habitat may lead to differences in gut microbiome diversity

[120]: adult birds living in deciduous forests have shown to harbor different gut

microbiome diversity than their counterparts living in open forested hay meadows.

Here, we used a cross-fostering design to study if the rearing environment

contributed to the variation in gut microbiome diversity: Our study indicated that

the nest of rearing seemed to explain more of the gut microbiome variation than the

nest of origin (although not significant), which follows some previous results [43-44,

52]. For example, a study with great and blue tit (*Cyanistes caerulaeus*) nestlings

showed that the nest of rearing contributed more to the gut microbiome than the

nest of origin [43], and another study with the brown-headed cowbird (*Molothrus*

ater) concluded that the sampling locality had a significant contribution to the gut

microbiome [44]. Teyssier et al. [52] conducted cross-fostering at day 8 post-hatch in

great tits and found that the nest of rearing influenced the gut microbiome more

than the nest of origin. Additionally, parents can pass down their bill and feather

microbiome through vertical transmission, which could influence nestling gut

microbiome [20].

Results from beta diversity analysis were similar to that of alpha diversity: brood

size manipulation did not contribute to the variation in gut microbiome composition.

Overall, variation in gut microbiome composition could be a result of different

genetic and environmental contributors. Firstly, great tit nestling gut microbiome

composition could be explained by underlying genetic effects that we did not

measure in this study. Phylosymbiosis i.e., the matching of gut microbiome

composition to host genetic structure, could be explained by underlying genetics

that may translate into physiological differences that affect the gut microbiome e.g.,

founder effects or genetic drift [121]. Davies et al. [14] found that MHC genes

correlate with gut microbiome composition: the expression of specific alleles in the

MHC genes was connected to the abundance of specific bacterial taxa such as

Lactobacillales and *Bacteroidales* that influenced host health. In a study by Benskin et

al. [41] captive zebra finches (*Taeniopygia guttata*) showed significant variation in gut

microbiome composition between individuals even though their diet and housing

conditions were standardized. The study suggested that individual homeostatic

mechanisms linking to naturally occurring differences in individual gut microbiome

could be why gut microbiome composition varied even with standardized housing

conditions [41]. Secondly, gut microbiome composition could have been affected by

the same environmental effects that may have linked to the variation in gut

microbiome diversity: diet and feeding behavior [115-116].

Differential analysis of ASV abundance showed variation in differential abundance

of taxa between the treatment groups. However, several ASVs were not

taxonomically assigned beyond family level making it difficult to draw conclusions

about the significance of these results. All treatment groups had taxa belonging to

the order *Firmicutes, Proteobacteria* and *Actinobacteria*, which was to be expected

because they are usually the most core phyla in passerine gut microbiomes [33].

Nestlings belonging to E, R or C group showed higher taxa abundance than the COU

group in each comparison. This result could be a result of the COU nestlings

generally hatching later in the season and potentially having a less diverse diet [103,

107]. Of the E, R and C groups, C group was less abundant than E and R groups.

Both E and R group showed high taxa abundance, which is interesting because we

hypothesized that nestlings belonging to the E group would potentially experience

less parental investment per nestling and have lower gut microbiome diversity and

therefore, be less abundant [56-57, 67-68]. We did not observe differential abundance

in e.g., the order *Lactobacillales* which would have been of interest, because the order

hosts taxa that are beneficial for gut microbiome health [14, 62]. The genus

Staphylococcus was differentially abundant in the E group, but not in the other

groups. *Staphylococcus* is a gram-positive genus of bacteria and known to cause

infections in its host species [122]. Curiously, the COU group was differentially

abundant in the genus *Dietzia*, which is a human pathogen [123].

Gut microbiome and short-term and mid-term survival

Our results showed that gut microbiome diversity and brood size were not

significantly associated with short-term (survival to fledging) or mid-term (apparent

juvenile) survival. However, while a more diverse gut microbiome is considered a

possible indicator of a healthy gut microbiome, the effects of the gut microbiome on

the host health may often be more complex and related to specific taxa [9-10]. For

example, Worsley et al. [13] did not find a correlation between body condition and

gut microbiome diversity, yet they found that specific taxa in the gut microbiome

linked with individual body condition and survival. Not only environment, but also

genetic background of the individual may contribute to gut microbiome and

survival. In a study by Davies et al. [14], *Ase-ua4* allele of the MHC genes was linked

to lower gut microbiome diversity and it was suspected that the variation in the

MHC genes could affect the sensitivity to pathogens that could lead to variation in

gut microbiome diversity and eventually, host survival.

To gain a better understanding of gut microbiome diversity and the contribution of

different taxa to host survival, functional analyses of the gut microbiome should be

included in gut microbiome studies. Different bacterial taxa can have similar

functions in the gut microbiome [5, 124] and therefore, the absence of some taxa may

be covered by other functionally similar taxa, resulting in a gut microbiome that is

functionally more stable [125]. Similarity in functions may also contribute to host's

local adaptation e.g., to the changes in the host's early-life environment [124]:

changes in brood size or dietary items could result in variation in the gut

microbiome diversity, yet there may be no effects on host body condition.

The lack of association between brood size, nestling size and survival contrasts with

previous studies, but it should be noted that the majority of previous studies have

been done with adult birds and not nestlings. Because nestling gut microbiome is

still quite flexible compared to that of the adults [20], it is possible that our

experiment did not result in a strong enough effect on the gut microbiome. In future

studies, it would be important to study the parents as well as it could be more likely

to find an association between adult microbiome and fitness than with nestling gut

microbiome and survival. Also, our sample size in the survival analyses was small,

and it is hard to determine if the result was affected by the sample size. Firstly,

nestling survival is often found to correlate with brood size and more specifically,

with fledging mass and in particular, the ability to forage for food [61, 126]. Intra-

brood competition may explain survival to fledging, as competition between

nestlings can limit food availability and thus, leading to lower nestling body

condition [68, 127]. A study with blackbirds (*Turdus merula*) showed that nestling

body mass explained juvenile survival [128], and similar results have been shown

with great tits and collared flycatchers (*Ficedula albicollis*; [31]). Contrastingly,

Ringsby et al. [129] observed that in house sparrows (*Passer domesticus*) juvenile

survival was independent of nestling mass and brood size. Moreover, natal body

mass is often positively correlated with survival to fledging and juvenile survival as

heavier nestlings are more likely to be recruited [92, 130-131], yet we failed to

demonstrate this in our study. Hatching date is also often positively correlated with

fledging success [132] yet we did not find this association in our study, but instead

found a significant association between hatching date and apparent juvenile

survival.

CONCLUSIONS

Offspring condition can be affected by the early-life environment and early-life gut microbiome, thus highlighting the importance of understanding how changes in the rearing environment affect individual body mass and survival. Even though our results showed between-individual variation in nestling gut microbiome diversity, we did not find a significant link between brood size and nestling gut microbiome. Moreover, we did not find a significant association between nestling gut microbiome diversity and short-term or mid-term survival. This suggests that other environmental factors (e.g., diet quality) may contribute more to variation in nestling gut microbiome. Further research is needed to uncover the environmental factors that contribute to nestling gut microbiome in wild bird populations, and how gut microbiome may be linked to nestling survival. Gut microbiome can adapt faster to environmental changes than the host, which makes it important to understand the causes of inter-individual variation in microbiome, and how variation in microbiome possibly mediate adaptation to environmental changes.

DECLARATIONS

Ethics statement

All animal work was conducted under relevant national and international guidelines and legislation. The animal work was licensed by the environmental and ethical committee of Varsinais-Suomi (environmental permit license number

VARELY/890/2020; animal ethics permit number ESAVI/5454/2020). The birds

were ringed from their nest-sites by permission from the Finnish Ringing Centre to

SR (Finnish Museum of Natural History).

Consent for publication

Not applicable

Availability of data and material

Sequence files are available at NCBI database (BioProject PRJNA877058). Metadata

and R scripts used to run the analyses are available in the Github Repository,

https://github.com/marannli/bsm_analyses.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by the Emil Aaltonen Foundation. NCS was supported by

EDUFI Fellowship (Opetushallitus). AS was funded by the Turku Collegium for

Science and Medicine, who contributed to fund the field study. AS acknowledges

funding from the European Commission Marie Sklodowska-Curie Postdoctoral

Fellowship (#894963) at the time of writing.

Author contribution

The idea for this study was by AS, SR and NCS. Sample collection was done by NCS, MH, AS and SR. Laboratory analyses were done by ML with assistance from SR, KG and EV. Sequence processing and statistical analyses were done by ML with assistance from KG, EV (bioinformatics) and SR (statistics). Manuscript was written by ML and all authors commented and approved the manuscript.

Acknowledgements

We would like to thank Charli Davies, Petri Papponen, Sari Viinikainen and Elina Virtanen at the University of Jyväskylä for their help with bioinformatics (CD) laboratory optimizations (PP, SV and EV), as well as Jorma Nurmi, Robin Cristofari, Natacha Garcin, Toni Laaksonen at the University of Turku and volunteer bird ringers for their help in the field.

REFERENCES

[1] Kinross JM, Darzi AW & Nicholson JK. Gut microbiome-host interactions in

health and disease. Genome medicine. 2011; 3:3:1-12.

[2] van Dongen WF, White J, Brandl HB, Moodley Y, Merkling T, Leclaire S, Wagner

RH. (2013). Age-related differences in the cloacal microbiota of a wild bird species.

BMC ecology. 2013; 13:1:1-12.

[3] David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE,

Turnbaugh PJ et al. Diet rapidly and reproducibly alters the human gut microbiome.

Nature. 2014; 505:7484:559-563.

[4] Sánchez B, Delgado S, Blanco-Míguez A, Lourenço A, Gueimonde M & Margolles

A. Probiotics, gut microbiota, and their influence on host health and disease.

Molecular nutrition & food research. 2017; 61:1:1600240.

[5] Heintz-Buschart A & Wilmes P. Human gut microbiome: function matters.

Trends in microbiology. 2018; 26:7:563-574.

[6] Egerton S, Culloty S, Whooley J, Stanton C & Ross RP. The gut microbiota of

marine fish. Frontiers in microbiology. 2018; 9:873.

[7] Pan D & Yu Z. Intestinal microbiome of poultry and its interaction with host and

diet. Gut microbes. 2014; 5:1:108-119.

[8] Hird SM. Evolutionary biology needs wild microbiomes. Frontiers in

Microbiology 2017;8.

[9] Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability

and resilience of the human gut microbiota. Nature. 2012; Sep;489:7415:220-30.

[10] Zaneveld JR, McMinds R & Vega Thurber R. Stress and stability: applying the

Anna Karenina principle to animal microbiomes. Nature microbiology. 2017; 2:9: 1-8.

[11] Rosshart SP, Vassallo BG, Angeletti D, Hutchinson DS, Morgan AP, Takeda K,

Rehermann B. Wild mouse gut microbiota promotes host fitness and improves

disease resistance. Cell. 2017a; 171:5:1015-1028.

[12] Rosshart SP, Vassallo BG, Angeletti D, Hutchinson DS, Morgan AP, Takeda K,

Rehermann B. Wild mouse gut microbiome protects laboratory mice against lethal

influenza virus infection and colorectal cancer. The Journal of Immunology. 2017b;

198.

[13] Worsley SF, Davies CS, Mannarelli ME, Hutchings MI, Komdeur J, Burke T &

Richardson DS. Gut microbiome composition, not alpha diversity, is associated with

survival in a natural vertebrate population. Animal microbiome. 2021; 3:1:1-18.

[14] Davies CS, Worsley SF, Maher KH, Komdeur J, Burke T, Dugdale HL &

Richardson DS. Immunogenetic variation shapes the gut microbiome in a natural

vertebrate population. Microbiome. 2022; 10:1:1-22.

[15] Zilber-Rosenberg I & Rosenberg E. Role of microorganisms in the evolution of

animals and plants: the hologenome theory of evolution. Fems Microbiology

Reviews. 2008; 32:723-735.

[16] Round JL & Mazmanian SK. The gut microbiota shapes intestinal immune

responses during health and disease. Nature Reviews Immunology. 2009; 9:313-323.

[17] Kopac SM & Klassen JL. Can they make it on their own? Hosts, microbiomes,

and the holobiont niche. Frontiers in Microbiology. 2016; 7:6.

[18] Hansen CHF, Nielsen DS, Kverka M, Zakostelska Z, Klimesova K, Hudcovic T,

Hansen AK. Patterns of early gut colonization shape future immune responses of the

host. PloS one. 2012; 7:3:e34043.

[19] Cox LM, Yamanishi S, Sohn J, Alekseyenko AV, Leung JM, Cho I, Blaser MJ et

al. Altering the intestinal microbiota during a critical developmental window has

lasting metabolic consequences. Cell. 2014; 158:4:705-721.

[20] Grond K, Lanctot RB, Jumpponen A, Sandercock BK. Recruitment and

establishment of the gut microbiome in arctic shorebirds. FEMS Microbiology

Ecology. 2017; 93:9.

[21] Knutie SA, Shea LA, Kupselaitis M, Wilkinson CL, Kohl KD & Rohr JR. Early-

life diet affects host microbiota and later-life defenses against parasites in frogs.

Integrative and Comparative Biology. 2017; 57:4:732-742.

[22] Videvall E, Song SJ, Bensch HM, Strandh M, Engelbrecht A, Serfontein N,

Cornwallis CK et al. Early-life gut dysbiosis linked to juvenile mortality in ostriches.

Microbiome. 2020; 8:1:1-13.

[23] Carmody RN, Gerber GK, Luevano Jr JM, Gatti DM, Somes L, Svenson KL &

Turnbaugh PJ. Diet dominates host genotype in shaping the murine gut microbiota.

Cell host & microbe. 2015; 17:1:72-84.

[24] Hooper LV, Littman DR & Macpherson AJ. Interactions between the microbiota

and the immune system. Science. 2012; 336:6086:1268-1273.

[25] Teyssier A, Matthysen E, Hudin NS, De Neve L, White J & Lens L. Diet

contributes to urban-induced alterations in gut microbiota: experimental evidence

from a wild passerine. Proceedings of the Royal Society B. 2020; 287:1920:20192182

[26] Bodawatta KH, Freiberga I, Puzejova K, Sam K, Poulsen M & Jønsson KA.

Flexibility and resilience of great tit (Parus major) gut microbiomes to changing

diets. Animal microbiome. 2021a;3:1:1-14.

[27] Góngora E, Elliott KH & Whyte L. Gut microbiome is affected by inter-sexual

and inter-seasonal variation in diet for thick-billed murres (Uria lomvia). Scientific

reports. 2021; 11:1:1-12.

[28] Schmiedová L, Tomášek O, Pinkasová H, Albrecht T & Kreisinger J. Variation in

diet composition and its relation to gut microbiota in a passerine bird. Scientific

reports. 2022; 12:1:1-13.

[29] Datta FU, Nolana JV, Rowe JB & Gray GD. Protein supplementation improves

the performance of parasitised sheep fed a straw-based diet. International Journal for

Parasitology. 1998; 28:8:1269-1278.

[30] Cotter SC, Simpson SJ, Raubenheimer D & Wilson K. Macronutrient balance

mediates trade-offs between immune function and life history traits. Functional

Ecology. 2011; 25:1:186-198.

[31] Linden M, Gustafsson L & Part T. Selection on fledging mass in the collared

flycatcher and the great tit. Ecology. 1992; 73:1:336-343.

[31] Spor A, Koren O & Ley R. Unravelling the effects of the environment and host

genotype on the gut microbiome. Nature Reviews Microbiology. 2011; 9:4:279-290

[33] Bodawatta KH, Hird SM, Grond K, Poulsen M & Jønsson KA. Avian gut

microbiomes taking flight. Trends in microbiology. 2021c.

[34] Waite DW & Taylor M. Exploring the avian gut microbiota: current trends and

future directions. Frontiers in microbiology. 2015; 6:673.

[35] Palmer C, Bik EM, DiGiulio DB, Relman DA & Brown PO. Development of the

human infant intestinal microbiota. PLoS biol. 2007; 5:7:e177.

[36] Chen X, Xu J, Ren E, Su Y & Zhu W. Co-occurrence of early gut colonization in

neonatal piglets with microbiota in the maternal and surrounding delivery

environments. Anaerobe. 2018; 49:30-40.

[37] Kalbermatter C, Fernandez Trigo N, Christensen S & Ganal-Vonarburg SC.

Maternal microbiota, early life colonization and breast milk drive immune

development in the newborn. Frontiers in immunology. 2021; 12:1768.

[38] Kohl KD. Diversity and function of the avian gut microbiota. Journal of

Comparative Physiology B. 2012; 182:5:591-602.

[39] Trevelline BK, MacLeod KJ, Knutie SA, Langkilde T, Kohl KD. In ovo microbial

communities: a potential mechanism for the initial acquisition of gut microbiota

among oviparous birds and lizards. Biology Letters. 2018; 14:7:4

[40] Ruiz-Rodriguez M, Lucas FS, Heeb P & Soler JJ. Differences in intestinal

microbiota between avian brood parasites and their hosts. Biological Journal of the

Linnean Society. 2009; 96:2:406-414.

[41] Benskin CMH, Rhodes G, Pickup RW, Wilson K & Hartley IR. Diversity and

temporal stability of bacterial communities in a model passerine bird, the zebra

finch. Molecular ecology. 2010;19.24:5531-5544.

[42] Lee CY, Peralta-Sánchez JM, Martínez-Bueno M, Møller AP, Rabelo-Ruiz M,

Zamora-Muñoz C & Soler JJ. The gut microbiota of brood parasite and host nestlings

reared within the same environment: disentangling genetic and environmental

effects. The ISME Journal. 2020; 14:11:2691-2702.

[43] Lucas FS & Heeb P. Environmental factors shape cloacal bacterial assemblages

in great tit Parus major and blue tit P. caeruleus nestlings. *Journal of Avian Biology*.

2005; *36*:6:510-516.

[44] Hird SM, Carstens BC, Cardiff S, Diffmann DL & Brumfield RT. Sampling

locality is more detectable than taxonomy or ecology in the gut microbiota of the

brood-parasitic Brown-headed Cowbird (Molothrus ater). Peerj 2014.

[45] van Veelen HPJ, Salles JF & Tieleman BI. Multi-level comparisons of cloacal,

skin, feather and nest-associated microbiota suggest considerable influence of

horizontal acquisition on the microbiota assembly of sympatric woodlarks and

skylarks. Microbiome. 2017; 5:1:1-17.

[46] Devaynes A, Antunes A, Bedford A & Ashton P. Bacterial species richness at

three stages of the breeding season in Cyanistes caeruleus (blue tit). Acta Oecologica.

2018; 92:123-130.

[47] Ding J, Dai R, Yang L, He C, Xu K, Liu S et al. Inheritance and establishment of

gut microbiota in chickens. Frontiers in Microbiology. 2017; 8:1967: doi:

10.3389/fmicb.2017.01967

[48] Preest MR, Folk DG & Beuchat CA. Decomposition of nitrogenous compounds

by intestinal bacteria in hummingbirds. The Auk. 2003; 120:4:1091-1101.

[49] Godoy-Vitorino F, Leal SJ, Díaz WA, Rosales J, Goldfarb KC, García-Amado

MA, Domínguez-Bello MG et al. Differences in crop bacterial community structure

between hoatzins from different geographical locations. Research in Microbiology.

2012; 163:3:211-220.

[50] Mendoza MLZ, Roggenbuck M, Vargas KM, Hansen LH, Brunak S, Gilbert MTP

& Sicheritz-Pontén T. Protective role of the vulture facial skin and gut microbiomes

aid adaptation to scavenging. Acta Veterinaria Scandinavica. 2018; 60:1:1-19.

1000

1001 [51] Fuirst M, Veit RR, Hahn M, Dheilly N & Thorne LH. Effects of urbanization on

1002 the foraging ecology and microbiota of the generalist seabird Larus argentatus. PLoS

1003 One. 2018; 13:12:e0209200.

1004

1005   [52] Teyssier A, Lens L, Matthysen E & White J. Dynamics of gut microbiota

1006   diversity during the early development of an avian host: evidence from a cross-foster

1007   experiment. Frontiers in microbiology. 2018; 9:1524

1008

1009   [53] Pettifor RA, Perrins CM & McCleery RH. Individual optimization of clutch size

1010   in great tits. Nature. 1988; 336:6195:160-162.

1011

1012   [54] Wright J, Both C, Cotton PA & Bryant D. Quality vs. quantity: energetic and

1013   nutritional trade-offs in parental provisioning strategies. Journal of Animal Ecology.

1014   1998; 620-634.

1015

1016   [55] Badyaev AV & Ghalambor CK. Evolution of life histories along elevational

1017   gradients: Trade-off between parental care and fecundity. Ecology. 2001; 82:10:2948-

1018   2960.

1019

1020   [56] Naef-Daenzer B & Keller LF. The foraging performance of great and blue tits

1021   (Parus major and P. caeruleus) in relation to caterpillar development, and its

1022   consequences for nestling growth and fledging weight. Journal of Animal Ecology.

1023   1999, 68: 708–718.

1024

1025   [57] de Kogel CH. Long-term effects of brood size manipulation on morphological

1026   development and sex-specific mortality of offspring. Journal of Animal Ecology.

1027   1997; 167-178.

1028

1029   [58] Pettifor RA, Perrins CM & McCleery RH. The individual optimization of fitness:

1030   variation in reproductive output, including clutch size, mean nestling mass and

1031   offspring recruitment, in manipulated broods of great tits Parus major. Journal of

1032   Animal Ecology. 2001; 70:1:62-79.

1033

1034   [59] Smith HG, Källander H & Nilsson JA. The trade-off between offspring number

1035   and quality in the great tit Parus major. The Journal of Animal Ecology. 1989; 383-

1036   401.

1037

1038   [60] Simon K, Verwoolde MB, Zhang J, Smidt H, de Vries Reilingh G, Kemp B &

1039   Lammers A. Long-term effects of early life microbiota disturbance on adaptive

1040   immunity in laying hens. Poultry science. 2016; 95:7:1543-1554.

1041

1042   [61] Davidson GL, Somers SE, Wiley N, Johnson CN, Reichert MS, Ross RP, Quinn JL

1043   et al. A time-lagged association between the gut microbiome, nestling weight and

1044   nestling survival in wild great tits. Journal of Animal Ecology. 2021; 90:4:989-1003.

1045

1046   [62] Kabir SM. The role of probiotics in the poultry industry. International Journal of

1047   Molecular Sciences. 2009; *10*:8:3531-3546.

1048

1049   [63] Maki JJ, Klima CL, Sylte MJ & Looft T. The microbial pecking order: utilization

1050   of intestinal microbiota for poultry health. *Microorganisms.* 2019; *7*:10:376.

1051

1052   [64] Du X, Xiang Y, Lou F, Tu P, Zhang X, Hu X, Xiao Y et al. Microbial community

1053   and short-chain fatty acid mapping in the intestinal tract of quail. Animals. 2020;

1054   10:6:1006.

1055

1056   [65] Den Besten G, Van Eunen K, Groen AK, Venema K, Reijngoud DJ & Bakker BM.

1057   The role of short-chain fatty acids in the interplay between diet, gut microbiota, and

1058   host energy metabolism. Journal of lipid research. 2013; 54:9:2325-2340.

1059

1060   [66] Harrison XA, Blount JD, Inger R, Norris DR & Bearhop S. Carry-over effects as

1061   drivers of fitness differences in animals. Journal of Animal Ecology. 2011; 80:1, 4-18.

1062

1063   [67] Leonard ML, Horn AG, Gozna A & Ramen S. Brood size and begging intensity

1064   in nestling birds. Behavioral Ecology. 2000; 11:2:196-201.

1065

1066   [68] Nicolaus M, Michler SP, Ubels R, Van Der Velde M, Komdeur J, Both C &

1067   Tinbergen JM. Sex-specific effects of altered competition on nestling growth and

1068 survival: an experimental manipulation of brood size and sex ratio. Journal of

1069 Animal Ecology. 2009; 78:2:414-426.

1070

1071 [69] Perrins CM. Population fluctuations and clutch size inthe Great tit, Parus major.

1072 Journal of Animal Ecology. 1965; 34:601–647

1073

1074 [70] Smith HG, Källander H & Nilsson JÅ. Effect of experimentally altered brood size

1075 on frequency and timing of second clutches in the great tit. The Auk. 1987; 104:4:700-

1076 706.

1077

1078 [71] Hõrak P, Tegelmann L, Ots I & Møller AP. Immune function and survival of

1079 great tit nestlings in relation to growth conditions. Oecologia. 1999; 121:3:316-322.

1080

1081 [72] Sanz JJ & Tinbergen JM. Energy expenditure, nestling age, and brood size: an

1082 experimental study of parental behavior in the great tit Parus major. Behavioral

1083 Ecology. 1999; 10:5:598-606.

1084

1085 [73] Tinbergen JM & Verhulst S. A fixed energetic ceiling to parental effort in the

1086 great tit? Journal of Animal Ecology. 2000; 69:2:323-334.

1087

1088   [74] Neuenschwander S, Brinkhof MW, Kölliker M & Richner H. Brood size, sibling

1089   competition, and the cost of begging in great tits (Parus major). Behavioral Ecology.

1090   2003; 14:4:457-462.

1091

1092   [75] Tinbergen JM & Sanz JJ. Strong evidence for selection for larger brood size in a

1093   great tit population. Behavioral Ecology. 2004; 15:4:525-533.

1094

1095   [76] Cossin-Sevrin N, Hsu BY, Marciau C, Viblanc VA, Ruuskanen S & Stier A. Effect

1096   of prenatal glucocorticoids and thyroid hormones on developmental plasticity of

1097   mitochondrial aerobic metabolism, growth and survival: an experimental test in wild

1098   great tits. Journal of Experimental Biology. 2022; 225:9:jeb243414.

1099

1100   [77] Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N,

1101   Knight R. et al. Ultra-high-throughput microbial community analysis on the Illumina

1102   HiSeq and MiSeq platforms. The ISME journal. 2012; 6:8:1621-1624.

1103

1104   [78] Parada AE, Needham DM & Fuhrman JA. Every base matters: assessing small

1105   subunit rRNA primers for marine microbiomes with mock communities, time series

1106   and global field samples. Environmental Microbiology. 2016; 18:5:1403–1414

1107

1108 [79] de Muinck EJ, Trosvik P, Gilfillan GD, Hov JR & Sundaram AY. A novel ultra

1109 high-throughput 16S rRNA gene amplicon sequencing library preparation method

1110 for the Illumina HiSeq platform. Microbiome. 2017; 5:1:1-15.

1111

1112 [80] Martin M. Cutadapt removes adapter sequences from high-throughput

1113 sequencing reads. *EMBnet. Journal.* 2011; *17*:1:10-12.

1114

1115 [81] Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA & Holmes SP.

1116 DADA2: high-resolution sample inference from Illumina amplicon data. Nature

1117 methods. 2016; 13:7:581-583.

1118

1119 [82] Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Glöckner FO et al.

1120 The SILVA ribosomal RNA gene database project: improved data processing and

1121 web-based tools. Nucleic acids research. 2012; 41:D1:D590-D596.

1122

1123 [83] Davis NM, Proctor DM, Holmes SP, Relman DA & Callahan BJ. Simple

1124 statistical identification and removal of contaminant sequences in marker-gene and

1125 metagenomics data. Microbiome. 2018; 2018; 6:1:1-14.

1126

1127 [84] McMurdie PJ & Holmes S. phyloseq: an R package for reproducible interactive

1128 analysis and graphics of microbiome census data. PloS one. 2013; 8:4:e61217.

1129

1130 [85] Lahti L & Sudarshan Shetty et al. Tools for microbiome analysis in R.

1131 2017;<http://microbiome.github.com/microbiome>

1132

1133 [86] Wickham H. Data analysis. In *ggplot2* (pp. 189-201). 2016; Springer, Cham.

1134

1135 [87] Wright ES. Using DECIPHER v2. 0 to analyze big biological sequence data in R.

1136 R Journal. 2016; 8:1.

1137

1138 [88] Schliep KP. phangorn: phylogenetic analysis in R. Bioinformatics. 2011; 27:4:

1139 592-593.

1140

1141 [89] Paradis E, Blomberg S, Bolker B, Brown J, Claude J, Cuong HS, Didier G et al.

1142 Package 'ape'. Analyses of phylogenetics and evolution, version. 2019; *2*(4).

1143

1144 [90] Yu G, Smith DK, Zhu H, Guan Y & Lam TTY. ggtree: an R package for

1145 visualization and annotation of phylogenetic trees with their covariates and other

1146 associated data. Methods in Ecology and Evolution. 2017; 8:1:28-36.

1147

1148 [91] Bates D, Maechler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models

1149 Using lme4. Journal of Statistical Software. 2015; 67(1), 1-48;doi:10.18637/jss.v067.i01.

1150

1151 [92] Monrós JS, Belda EJ, Barba E. Post-fledging survival of individual great tits: the

1152 effect of hatching date and fledging mass. Oikos. 2002; Dec;99:3:481-8.

1153

1154 [93] Fox J, Weisberg S, Adler D, Bates D, Baud-Bovy G, Ellison S, Monette G et al.

1155 Package 'car'. Vienna: R Foundation for Statistical Computing. 2012; 16.

1156

1157 [94] Hartig F & Hartig MF. Package 'DHARMa'. *R package*. 2017.

1158

1159 [95] Bray JR & Curtis JT. An ordination of the upland forest communities of southern

1160 Wisconsin. Ecological monographs.1957; 27:4:326-349.

1161

1162 [96] Lozupone C & Knight R. UniFrac: a new phylogenetic method for comparing

1163 microbial communities. Applied and environmental microbiology. 2005; 71:12:8228-

1164 8235.

1165

1166 [97] Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara RB, Oksanen

1167 MJ etl al. Package 'vegan'. Community ecology package, version. 2013; 2:9:1-295.

1168

1169 [98] Moreno J, Cowie RJ, Sanz JJ & Williams RS. Differential response by males and

1170 females to brood manipulations in the pied flycatcher: energy expenditure and

1171 nestling diet. Journal of Animal Ecology. 1995; 721-732.

1172

1173   [99] Askenmo C. Effects of addition and removal of nestlings on nestling weight,

1174   nestling survival, and female weight loss in the Pied Flycatcher Ficedula hypoleuca

1175   (Pallas). Ornis Scandinavica. 1977;1-8.

1176

1177   [100] Nur N. The consequences of brood size for breeding blue tits II. Nestling

1178   weight, offspring survival and optimal brood size. The Journal of Animal Ecology.

1179   1984; 497-517.

1180

1181   [101] Sanz JJ. Clutch size manipulation in the pied flycatcher: effects on nestling

1182   growth, parental care and moult. Journal of Avian Biology. 1998; 157-162.

1183

1184   [102] Burness GP, McClelland GB, Wardrop SL & Hochachka PW. Effect of brood

1185   size manipulation on offspring physiology: an experiment with passerine birds.

1186   Journal of Experimental Biology. 2000; 203:22:3513-3520.

1187

1188   [103] Naef-Daenzer L, Naef-Daenzer B & Nager RG. Prey selection and foraging

1189   performance of breeding Great Tits Parus major in relation to food availability.

1190   Journal of Avian Biology. 2000; 31:2:206-214.

1191

1192   [104] Rytkönen S & Orell M. Great tits, Parus major, lay too many

1193   eggs:experimental evidence in mid-boreal habitats. Oikos. 2001; 93: 439 – 450

1194

1195   [105] van der Most PJ, de Jong B, Parmentier HK & Verhulst S. Trade-off between

1196   growth and immune function: a meta-analysis of selection experiments. Functional

1197   Ecology. 2011; 25:1:74-80.

1198

1199   [106] Rauw WM. Immune response from a resource allocation perspective. Frontiers

1200   in genetics. 2012; 3:267.

1201

1202   [107] Naef-Daenzer B & Keller LF. The foraging performance of great and blue tits

1203   (Parus major and P. caeruleus) in relation to caterpillar development, and its

1204   consequences for nestling growth and fledging weight. Journal of Animal Ecology.

1205   1999; 6:84:708-718.

1206

1207   [108] Barba E & Gil-Delgado JA. Seasonal variation in nestling diet of the great tit

1208   Parus major in orange groves in eastern Spain. Ornis scandinavica. 1990; 296-298.

1209

1210   [109] Tinbergen JM & Dietz MW. Parental energy expenditure during brood rearing

1211   in the great tit (Parus major) in relation to body mass, temperature, food availability

1212   and clutch size. Functional ecology. 1994; 563-572.

1213

1214   [110] Ilmonen P, Hasselquist D, Langefors Å & Wiehn J. Stress, immunocompetence

1215   and leukocyte profiles of pied flycatchers in relation to brood size manipulation.

1216   Oecologia. 2003; 136:1:148-154.

1217

[111] Kohl KD, Brun A, Caviedes-Vidal E & Karasov WH. (2019). Age-related changes in the gut microbiota of wild House Sparrow nestlings. Ibis. 2019; 161:1:184-191.

[112] Waite DW & Taylor MW. Characterizing the avian gut microbiota: membership, driving influences, and potential function. Frontiers in microbiology. 2014; 5:223.

[113] Davidson GL, Wiley N, Cooke AC, Johnson CN, Fouhy F, Reichert MS, Quinn JL et al. Diet induces parallel changes to the gut microbiota and problemsolving performance in a wild bird. Scientific reports. 2020; 10:1:1-13.

[114] Murray MH, Lankau EW, Kidd AD, Welch, CN, Ellison T, Adams HC, Hernandez SM et al. Gut microbiome shifts with urbanization and potentially facilitates a zoonotic pathogen in a wading bird. PLoS One. 2020; 15:3:e0220926.

[115] Bodawatta KH, Koane B, Maiah G, Sam K, Poulsen M & Jønsson KA. Species-specific but not phylosymbiotic gut microbiomes of New Guinean passerine birds are shaped by diet and flight-associated gut modifications. Proceedings of the Royal Society B. 2021b;288:1949:20210446.

1238

1239 [116] Bodawatta KH, Klečková I, Klečka J, Pužejová K, Koane B, Poulsen M, Sam K

1240 et al. Specific gut bacterial responses to natural diets of tropical birds. Scientific

1241 reports. 2022; 12:1:1-15.

1242

1243 [117] Alberdi A, Aizpurua O, Bohmann K, Zepeda-Mendoza ML & Gilbert MTP. Do

1244 vertebrate gut metagenomes confer rapid ecological adaptation? Trends in Ecology

1245 & Evolution 2016; 31:9:689-699.

1246

1247 [118] Wilkin TA, King LE & Sheldon BC. Habitat quality, nestling diet, and

1248 provisioning behaviour in great tits Parus major. Journal of Avian Biology. 2009;

1249 40:2:135-145.

1250

1251 [119] Rytkönen S, Vesterinen EJ, Westerduin C, Leviäkangas T, Vatka E, Mutanen M,

1252 Orell M et al. From feces to data: A metabarcoding method for analyzing consumed

1253 and available prey in a bird-insect food web. Ecology and evolution. 2019; 9:1:631-

1254 639.

1255

1256 [120] Drobniak SM, Cichoń M, Janas K, Barczyk J, Gustafsson L &

1257 Zagalska-Neubauer M. Habitat shapes diversity of gut microbiomes in a wild

1258 population of blue tits Cyanistes caeruleus. Journal of Avian Biology. 2022;2:e02829.

1259

1260   [121] Fleischer R, Risely A, Hoeck PE, Keller LF & Sommer S. Mechanisms governing

1261   avian phylosymbiosis: Genetic dissimilarity based on neutral and MHC regions

1262   exhibits little relationship with gut microbiome distributions of Galápagos

1263   mockingbirds. Ecology and evolution. 2020; 10:23:13345-13354.

1264

1265   [122] Kloos WE. Natural populations of the genus Staphylococcus. Annual Reviews

1266   in Microbiology. 1980; 34:1:559-592.

1267

1268   [123] Gharibzahedi SMT, Razavi SH & Mousavi SM. Characterization of bacteria of

1269   the genus Dietzia: an updated review. Annals of microbiology. 2014; 64:1:1-11.

1270

1271   [124] Hanning I, Diaz-Sanchez S. The functionality of the gastrointestinal

1272   microbiome in non-human animals. Microbiome. 2015; 3:5: doi: 10.1186/s40168-015-

1273   0113-6

1274

1275   [125] Doak DF, Bigger D, Harding EK, Marvier MA, O'malley RE & Thomson D. The

1276   statistical inevitability of stability-diversity relationships in community ecology. The

1277   American Naturalist. 1998; 151:3:264-276.

1278

1279   [126] Naef-Daenzer B, Widmer F & Nuber M. Differential post-fledging survival of

1280   great and coal tits in relation to their condition and fledging date. Journal of Animal

1281   ecology. 2001; 70:5:730-738.

1282

[127] Nilsson JA & Svensson M. Sibling competition affects nestling growth

strategies in marsh tits. Journal of Animal Ecology. 1996; 825-836.

1285

[128] Magrath RD. Nestling weight and juvenile survival in the blackbird, Turdus

merula. The Journal of Animal Ecology. 1991; 335-351.

1288

[129] Ringsby TH, Sæther BE & Solberg EJ. Factors affecting juvenile survival in

house sparrow Passer domesticus. Journal of Avian Biology. 1998; 241-247.

1291

[130] Both C, Visser ME & Verboven N. Density–dependent recruitment rates in

great tits: the importance of being heavier. Proceedings of the Royal Society of

London. Series B: Biological Sciences. 1999; 266:1418:465-469.

1295

[131] Cleasby IR, Nakagawa S, Gillespie DO & Burke T. The influence of sex and

body size on nestling survival and recruitment in the house sparrow. Biological

Journal of the Linnean Society. 2010; 101:3:680-688.

1299

[132] Tinbergen JM & Boerlijst MC. Nestling weight and survival in individual great

tits (Parus major). The Journal of Animal Ecology. 1990; 1113-1127.

1302

SUPPLEMENTARY INFORMATION

All supplementary files are provided in one word (.docx) document with each

section titled as follows:

Supplementary file 1. Brood size before and after manipulation: brood sizes

between treatment groups were tested with a linear model to see if the differences

were statistically significant. **Supplementary file 2. Rarefaction curves for the**

unrarefied dataset. Species (ASVs) plateaued at about 5000 reads which was used as

the rarefying depth. **Supplementary file 3.** Phylogenetic tree using the Newick-

format. The tree describes the dissimilarity among the treatment groups. Each tip

represents an individual sample, and each tip is colored and shaped based on

treatment. Treatment groups are clustered using the UPGMA algorithm.

Supplemental information 4. (A) Linear mixed effects model for gut microbiome

diversity (Shannon Diversity Index and Chao1 Richness) and brood size

manipulation treatment. (B) Linear mixed effects model for GM diversity (Shannon

Diversity Index and Chao1 Richness) and manipulated brood size. **Supplementary**

file 5. A linear mixed effects model investigating the effects of brood size

manipulation on nestling body mass on day 7 and day 14 post-hatch.

Supplementary file 6. A linear mixed effects model investigating the effects of

manipulated brood size on nestling body mass on day 7 and day 14 post-hatch.

Supplementary file 7. A linear mixed effects model investigating the associations

between alpha diversity (Shannon Diversity Index and Chao1 Richness) and brood

size manipulation. **Supplementary file 8.** A linear mixed effects model investigating

the association between alpha diversity (Shannon Diversity Index and Chao1

Richness) and manipulated brood size. **Supplementary file 9.** A generalized linear

model exploration into alpha diversity's (Shannon Diversity Index and Chao1

Richness) association with short-term (survival to fledging) and mid-term (apparent

juvenile) survival. **Supplementary file 11.** Generalized linear model to measure the

association between alpha diversity (Shannon Diversity Index and Chao1 Richness)

survival to fledging and apparent juvenile survival. **Supplementary file 11.**

Generalized linear model to measure the association between alpha diversity

(Shannon Diversity Index and Chao1 Richness) survival to fledging and apparent

juvenile survival. **Supplementary file 12.** Ordination of the gut microbial

communities.

1337

Before manipulation

After manipulation

E

Enlarged nest gained four nestlings from reduced nest. Additionally, two nestlings from the original enlarged nest were moved to the reduced nest.

R

Reduced nest lost two nestlings when four of the nestlings were moved to the enlarged nests and only two nestlings were moved back to the reduced nest.

Control

C

Moved across broods but original and manipulated brood sizes were identical.

Unmanipulated control   **COU**

Not moved at all, but only sampled for a fecal sample at day 7 post-hatch.