Asgard ESCRT-IIIs assemble into helical filaments in the presence of

DNA and remodel eukaryotic membranes

Nataly Melnikov[1,2], Dikla Nachmias[1,2], Gal Halbi[3], Alexander Upcher[4], Ran Zalk[4], Ann

Bernheim-Groswasser[3,4], Natalie Elia[1,2]

1 Department of Life Sciences, Ben-Gurion University of the Negev, Beer Sheva 84105,

Israel

2 National Institute for Biotechnology in the Negev (NIBN), Ben-Gurion University of the

Negev, Beer Sheva 84105, Israel

3 Department of Chemical Engineering, Ben-Gurion University of the Negev, Beer

Sheva 84105, Israel

4 Ilse Katz Institute for Nanoscale Science and Technology, Ben Gurion University of

the Negev, Beer Sheva 84105, Israel

* Corresponding author:

Natalie Elia

Phone: 972-8-6428735

Email: elianat@post.bgu.ac.il

ABSTRACT

The ESCRT machinery mediate membrane remodeling in numerous processes in cells

including cell division and nuclear membrane reformation. The identification of ESCRT

homologs in Asgard archaea, currently considered the closest ancestor of eukaryotes,

suggests a role for ESCRTs in the membrane remodeling processes that occurred

during eukaryogenesis. Yet, the function of these distant ESCRT homologs is mostly

unresolved. Here we show that Asgard ESCRT-III proteins self-assemble into homo-

and hetero- helical tubes, a hallmark of the eukaryotic ESCRT system. Asgard ESCRT-

III tube assembly was facilitated in the presence of DNA and inhibited by DNAase.

Notably, Asgard ESCRT-III filaments remodeled eukaryotic-like membrane vesicles,

also in the presence of DNA, indicating an ancient role for the ESCRT complex in

membrane remodeling, that may involve DNA binding. The ability of Asgard archaeal

ESCRTs to remodel eukaryotic-like membranes, places them at the junction between

prokaryotes and eukaryotes, substantiating a role for ESCRTs in eukaryogenesis.

INTRODUCTION

The Asgard superphyla, manifest a unique branch that, according to current

evolutionary theories, is considered to be the closest prokaryotic relative of eukaryotes.

All Asgard archaeal species were found to encode for Eukaryotic Signature Proteins

(ESPs), that carry cellular functions that until recently were considered as exclusive to

eukaryotes (1-3). Thus, Asgard proteins potentially hold the core functions of ancient

eukaryotic cellular machines. Of particular interest, are ESPs with membrane

remodelling capabilities, as these proteins may contribute to the establishment of

cellular compartmentalization during eukaryogenesis. Indeed, Asgard archaea encode

for several eukaryotic membrane remodelling machineries including the SNARE

complex, BAR domains proteins and the ESCRT machinery (1, 3, 4). Among these

machineries, ESCRTs are the most conserved and are encoded by all Asgard species

discovered up to date, suggesting that they carry essential functions in these unique

prokaryotes (5, 6).

The ESCRT machinery constitute one of the most robust and versatile cellular

apparatus for membrane constriction and fission. Proteins of the machinery are

encoded by all domains of life and were shown to mediate membrane remodelling in a

wide range of eukaryotic cellular membranes including plasma membrane, nuclear

membrane and endocytic membranes (7, 8). Moreover, ESCRT-mediated processes

span a large spectrum of length scales, from less than 100 nm for vesicle release to up

to 1 µm for cell division. How the ESCRT complex mediate membrane remodelling in

this large landscape of membrane sources and length scales is mostly unknown (9).

The eukaryotic ESCRT system is composed of five subfamilies, i.e. ESCRT 0-III and the

AAA-ATPase VPS4. Within this complex, ESCRT-III (named CHMPs in animal cells)

and VPS4 manifest the minimal unit required for driving membrane fission (9, 10).

According to current models, membrane constriction and scission are mediated by

polymerization of ESCRT-III proteins into helical hetero-filaments and their remodelling

by VPS4 (10-14). Indeed, ESCRT-III proteins have been shown organize in helical

filaments and tubes both in vitro and in cells (14-19). However, the large number of

proteins within the ESCRT-III subfamily —twelve in humans and eight in yeast - have

challenged mechanistic studies of the basic function of the machine.

Reduced, simplified ESCRT-III systems have been identified in prokaryotes including in

bacteria and archaea (named Vipp1 and PspA in bacteria and CdvB in archaea) (7, 20,

21). While the bacterial homologs were shown to remodel membranes in vitro, no VPS4

was identified in this system so far, and whether these distant homologs can sever

membranes has not been clarified. ESCRTs encoded by TACK archaea were shown to

participate in cell division, suggesting functional similarities between the eukaryotic and

archaeal ESCRT systems (22-25). Within the archaeal domain the Asgard ESCRT

system is considerably closer to the eukaryotic system. While all other archaea encode

only for homologs of the ESCRT-III/VPS4 module, Asgard archaea encode for

homologs of the complete ESCRT system (ESCRT-I-III and VPS4) (6, 7). Additionally,

the sequence of Asgard ESCRT-III/VPS4 proteins is more closely related to those of

eukaryotes than to other archaeal ESCRT systems (the CDV system) (5, 20). Lastly,

VPS4 homologs of Asgard archaea were shown to functionally interact with eukaryotic

ESCRTs in both yeast and mammalian cells (5, 26). Notably, the Asgard ESCRT-III

subfamily is substantially reduced compared to eukaryotes with only two ESCRT-III

proteins (named here CHMP1-3 and CHMP4-7) encoded in some species including

Loki and Heimdall archaeota (1, 5, 7, 26). Hence, the Asgard ESCRT system

constitutes a minimalist ESCRT machinery that may hold the core capabilities of

eukaryotic ESCRTs and could have been involved in membrane remodelling processes

that occurred during eukaryogenesis. Yet, the function, organization, and properties of

Asgard ESCRT-III complexes as well as their ability to remodel membranes are mostly

unknown.

In this work, we provide the first evidence that Asgard ESCRT-III complexes share

functional similarities with eukaryotic ESCRTs. Using purified ESCRT-III homologs

encoded by the most abundant Asgard phyla, Lokiarchaeota (Loki), we show that the

Asgard ESCRT-III protein CHMP4-7 self-assemble into homo- and hetero- helical tubes

that resemble those of eukaryotes (15, 18, 19, 27). The proportion of helical tubes was

dramatically enriched in the presence of short DNA oligos, while no filaments could be

observed upon DNAse treatment. Notably, Loki ESCRT filaments, formed under any of

these conditions, were able to bind and deform small unilamellar vesicles (SUVs)

comprised of eukaryotic-like synthetic phospholipids. Hence, Loki ESCRT-III assemble

into helical filaments that are affected by oligonucleotides and are capable of

remodelling eukaryotic-like membranes.

RESULTS

Eukaryotic ESCRT-III proteins polymerize into helical filaments - a hallmark of ESCRT-

III organization that is thought to be essential for their function in membrane remodeling

(8, 28). Polymerization is mediated by the snf7 domain that is conserved in both

archaea and eukaryotes, but the polymerization properties of Asgard ESCRT-III has not

been examined (5, 20, 26). To investigate the assembly properties of Asgard ESCRT-III

proteins, we purified recombinant versions of the ESCRT-III proteins CHMP4-7 and

CHMP 1-3, encoded by the Loki GC14_75 strain (KKK44605.1 and KKK42122.1,

respectively). Using our assay conditions (see material and methods), full length CHMP

4-7 self-assembled into filaments (Fig 1). No filaments were observed under the same

conditions using full length CHMP1-3 (sup Fig. 1A). In negative stain TEM images,

CHMP 4-7 was predominantly seen in either long, thin filaments or rod-shaped tubes

(Fig 1A, see arrows). A pattern of parallel, periodic stripes was detected in the CHMP 4-

7 tube structures by cryo-EM, supporting a helical tube organization (Fig 1B, C). The

outer diameter of the CHMP4-7 tubes was variable, ranging between 35 to 55 nm

(averaged diameter 44nm ± 30.9). Tube length was typically between 300 - 450 nm (Fig

1D). The overall shape of CHMP4-7 helical tubes resembled the helical tubes previously

described for eukaryotic or bacterial ESCRT-III filaments (15, 18, 19, 29). The diameter

of Loki CHMP 4-7 helical tubes was slightly larger than the diameters measured for

human CHMP1B (~25nm) or the bacterial ESCRT III homologs PspA (~21 nm), but was

in the range of diameters measured for human CHMP2A-CHMP3 (38-43 nm) (18, 19,

29). Collectively, these data indicate that Loki CHMP 4-7 spontaneously self-assemble

into the typical helical tube organization described for the ESCRT-III complex.

Next, we examined whether, similar to their human homologs, Loki ESCRT-IIIs can

assemble into co-polymers (18, 19). To this end, full lengths Loki CHMP4-7 and CHMP

1-3 were incubated at different ratios. While no filaments were seen at 1:1 CHMP4-7-

CHMP1-3 ratio, helical tubes could be readily observed at 2:1, 4:1 and 6:1 ratios (Fig 2

and Sup Fig 1B). Notably, CHMP 4-7-CHMP 1-3 helical tubes exhibited different

properties than those observed for the CHMP 4-7 homo-polymer. First, their outer

diameter was significantly smaller and less diverse (CHMP 4-7-CHMP 1-3 2:1 ratio,

31±7.7) (Fig. 2B, left panel). Second, helical tubes formed in the presence of both

CHMP4-7 and CHMP1-3 were significantly longer than those formed by the CHMP4-7

homo-polymer (>X1.5 folds increase for 2:1 and 4:1 ratios) (Fig. 2B, right panel). Third,

the morphology of the CHMP4-7-CHMP1-3 helical tube was slightly different: tubes

appeared to be more curved and the periodic stripes, which could only be observed by

cryo-EM for CHMP 4-7 homo-polymers, were readily detected in negative stain images

of the co-polymer, suggesting that packing of the helical tube is less tight in the

presence of CHMP1-3 (Fig 2A). These phenotypic changes were not induced upon

substituting CHMP 1-3 with purified GFP, suggesting that they are specifically

contributed by CHMP 1-3 (Sup. Fig 1C). We therefore concluded that Loki ESCRT-III

proteins assemble into helical tubes composed of homo- and co-polymers with the co-

polymers forming longer, narrower, and more homogenous tubes than the homo-

polymer. These findings are in line with previous observations obtained for human

ESCRT-III CHMP1B homo- and co-polymers (in the presence of IST1), which also

formed a narrower, more uniform tubes in the co-polymer composition (30). The

realization that both human and Asgard ESCRT-III proteins assemble into homo-and

co- polymers with the latter adopting a narrower more uniform tube shape, strongly

suggests that this is a basic, conserved property of the ESCRT-III system.

Recent work from our laboratory showed that purified Loki CHMP4-7 binds short DNA

oligonucleotides (26). Human CHMP1B, exhibited similar properties and a recently

solved cryo-EM structure of human CHMP1B/IST1 filaments demonstrated direct

interactions between the ESCRT-III filament and nucleic acids (26, 31). We, therefore,

set to examine the effect of short DNA oligonucleotides on the self-assembly properties

of Asgard ESCRT-III filaments. A significant increase in helical tube assemblies was

observed for the co-polymer in the presence of ssDNA, in both negative stain- and cryo-

TEM images (Fig 3A-B and Sup Fig 2A). More helical tubes were also observed upon

addition of dsDNA, but to a lower extent. No filaments were observed when incubating

CHMP1-3 with DNA, indicating that that observed increase in helical tubes do not result

from self-assembly of CHMP1-3 into homo-polymers (Sup Fig. 2D). Addition of ssDNA

to CHMP4-7 homo-polymers also led to a dramatic increase in helical tubes, indicating

that the effect of DNA is not restricted to the co-polymer (Fig. 3C, Sup Fig 2B). The

presence of DNA also affected the properties of the helical tubes. CHMP 4-7-CHMP1-3

helical tubes formed in the presence of either ds- or ss- DNA exhibited larger and more

variable diameters (43.9nm ±58 for CHMP4-7-CHMP1-3 2:1 ssDNA vs CHMP4-7-

CHMP1-3 2:1 31nm ±7.7), resembling the diameter measured for the CHMP 4-7 homo-

polymer (CHMP4-7 44nm ±30.9) (Fig. 3D). Moreover, filaments were less curved, and

the periodic stripes observed in negative stain images in the co-polymer could not be

observed (Fig 3A). Addition of DNA did not affect the characteristics length or diameter

of tubes formed by CHMP 4-7 homo-polymers (Fig. 3D-E). It, therefore, appears that

DNA aids the organization of Loki-ESCRT-III proteins into helical tubes and counteract,

at least to some extent, the effect of CHMP1-3 on tube characteristics.

To further investigate the possible effect of DNA on Loki ESCRT-III polymerization, we

performed the self-assembly reaction in the presence of DNase. Surprisingly, filament

formation was completely abolished for both homo- and co-polymers in the presence of

DNase (Fig 3F and Sup Fig. 2E-F). Additionally, no filaments were observed upon

incubation with DNAse post self-assembly, suggesting that filament stability and/or

integrity are dependent on DNA. Altogether, these findings strongly suggest that DNA is

an essential component of Loki ESCRT-III filamentous organization.

Eukaryotic ESCRT-IIIs were shown to bind and deform membranes *in vitro* and in cells

(28). To determine whether ESCRTs encoded by Asgard archaea carry the ability to

remodel membranes, we incubated Loki ESCRT-III pre-assembled filaments with small

unilamellar vesicles (SUVs, ~100 nm) composed of negatively charged eukaryotic-like

phospholipids (1:1, PC:PS). Under these conditions, ESCRT-III filaments, assembled

under any of the conditions described above (including in the presence of DNA), were

found to intimately interact with SUVs (Fig. 4A, B and Sup Fig. 3). We identified four

main types of ESCRT-III-SUV interactions: (1) Attachment of the SUV to the outside

surface of the tube, which was associated with flattening of the vesicle at the filament-

vesicle interface; (2) Docking of the SUV at the tip of the helical tube; (3) Invagination of

the SUV to the interior of the ESCRT-III tube; and (4) Dramatic remodeling of both the

SUV and the ESCRT-III tube (see examples in Fig. 4C, left panel). Combinations of

several types of interactions on the same filament were also observed, indicating that

these interactions are not mutually exclusive. Moreover, a docked SUV that invaginates

the helical tube was observed occasionally (Fig. 4B, left panel and sup movies 1, 2),

suggesting that these two types of interactions represent successive steps.

Invaginations of more than one vesicle to the same tube could be observed, in cryo-

TEM and cryo-tomography reconstructions, giving rise to the formation of a complex

membrane system with several vesicles internalized into one another (Fig 4B, and sup

movies 1, 2). In these complex organizations, which were often accompanied by a

considerable deformation of both the filament and the vesicles, the characteristics

periodic stripes of the ESCRT-III filament could often be detected at the exterior of the

filament-vesicle complex, suggesting that the observed morphologies result from

internalization of vesicles into the ESCRT-III helical tube followed by deformation of the

tube. Loki ESCRT III filaments also interacted with neutrally charged SUVs (100% PC)

but to a much lower extent, exhibiting predominantly binding of the SUV on the outside

of the helical tube (Sup Fig. 4). Therefore, Loki ESCRT-III filaments associate with

eukaryotic-like membrane vesicles, and deform negatively charged vesicles via the

interior of the tube (see proposed model in Fig. 4D).

Last, we asked whether the composition of ESCRT-III helical tubes, described here,

affect their capabilities to remodel membranes. Because vesicle-filaments interactions

were identified under all conditions, we analyzed the proportion of the four types of

filament-vesicle interaction, we identified, in different ESCRT-III compositions. No

difference was observed in the distributions of homo- and co- polymers, suggesting that

the composition of the filament does not significantly affect membrane indications. The

presence of DNA affected the distributions of ESCRT-III-SUV interactions in both the

homo- and co- polymers, as follows (Fig. 4C): (1) The association of vesicles with the

outside surface of the helical tube was considerably enriched (outside, type 1, No DNA,

<15%; with DNA, >35%). (2) Interactions with the tip of the filament (docking, type 2)

and invagination of the vesicle (inside, type 3) were significantly enriched. (3) The

extent of complete remodeling events (remodeling, type 4) was dramatically reduced

(with DNA, ~5%; no DNA ~80%). Therefore, homo- and co- Loki ESCRT-III filaments

are capable of interacting and remodeling lipid membranes, both in the presence and

absence of DNA, but the extent of remodeling appear to be attenuated by the presence

of DNA.

DISCUSSION

In this work, we provide the first evidence to show that, similar to their eukaryotic

homologs, Asgard ESCRT-III proteins assemble into helical filaments that remodel

membranes. These findings confirm that ESCRT homologs encoded by Asgard archaea

are functionally related to their eukaryotic homologs. We further provide data to support

prominent role for DNA in the Asgard ESCRT-III system. Finally, by using SUVs

composed of eukaryotic phospholipids, which are fundamentally different from archaeal

lipids, we demonstrate that the Asgard ESCRT-III could, in-principle, contribute to the

membrane remodeling processes that occurred during eukaryogenesis and gave rise to

the complex phospholipids-based endomembrane system of eukaryotes.

Our data point to similarities between the human and Asgard ESCRT-III systems. First,

the nature of self-assembly appear to be conserved. Similar to the human CHMP1B-

IST1 complex, Loki CHMP4-7 was found to polymerize alone or in the presence of

CHMP 1-3, with the homo-polymer exhibiting a wider, more variable diameter compared

to the co-polymer (30). What controls the diameter of the ESCRT-III helical tube is

currently unknown. Our findings that, similar to their human homologs, Asgard ESCRT-

III homo-filaments can adopt a wide range of diameters that can be tuned by the

addition of another ESCRT-III components, strongly suggest that this is a conserved

feature of the ESCRT-III system. Therefore, a basic feature of the ESCRT-III system

may be their ability to assemble into flexible homo filaments that can be tuned to adopt

different shapes by incorporating additional ESCRT-III proteins into the polymer. Such a

model can explain how the ESCRT-III system execute the large repertoire of membrane

remodeling processes in cells, which greatly vary in scales and topologies (7, 8).

The topology of membrane binding, described here for Loki ESCRT-IIIs, resemble the

topology reported for the human CHMP1B-IST1 polymer, i.e. bindings via the interior of

the helical tube (30). This topology is reversed to the conventional topology attributed to

the ESCRT complex and that was described for the CHMP2A-CHMP3 co-polymer (19,
27), which is binding and deforming membranes via the external surface of the tube.
Reversed membrane binding topology was also recently reported for the bacterial
ESCRT-III homologs Vipp1 and PspA (29, 32). Whether this topology represent the
preferred topology of ancient ESCRTs or is a consequence of the experimental, in vitro,
system is yet to be determined. Regardless, these similarities suggest that the ESCRT-
III membrane remodeling properties reported here are a common property of the
ESCRT system, conserved in all domains of life.
286
Our results highlight DNA as an essential Loki ESCRT-III partner. Organization of both
homo- and co- ESCRT-III polymers into helical tubes was enhanced in the presence of
short DNA oligonucleotides and helical tube formation was completely blocked upon
addition of DNase. Whether DNA has a role in stabilizing Loki ESCRT-III filaments or in
facilitating their nucleation is still unknown. Notably, human CHMP1B-IST1 filaments
were shown to bind DNA in the inner side of the tube (31). Additionally, nucleotides
were recently resolved in cryo-EM structures of the bacterial ESCRT-III Vipp1 and were
suggested to play a role in helical tube stabilization (32). It is therefore tempting to
speculate that nucleotide binding is an additional core property of the ESCRT-III
machine that is conserved in all domains of life.
297
Our data show that Loki ESCRT-III filaments that polymerized in the presence of DNA
were able to bind and deform membrane vesicles. Yet, the extent of membrane
deformation that occurred inside the tube was attenuated, while binding on the exterior
of the ESCRT-III tube was enhanced. These results are consistent with competition of
DNA and membranes over the same binding sites, located at the interior of the tube, as
recently suggested for human CHMP1B (31). Alternatively, the presence of DNA may
change the biophysical properties of the tube, thereby resisting the deformation of the
tube needed to facilitate membrane remodeling from within. Future experiments
allowing direct visualization of ESCRT-III proteins, DNA and the membrane at high
resolution are needed in order to examine these models and to resolve the intimate
relationship between these three partners.

Overall, the Loki ESCRT-III filaments characterized here, share more similar

characteristics to human CHMP1B-IST1 filaments than to human CHMP2A-CHMP3 or

yeast snf7 (human CHMP4B) filaments. Filaments of both Asgard ESCRT-III and

human CHMP1B, were shown to assemble into homo- and co- polymers with the latter

being more constricted; both filaments were reported to bind DNA in vitro; and both

deformed membranes via binding to the inner side of the filamentous tube. Consistently,

we recently showed that both Loki CHMP4-7 and human CHMP1B associate with

chromatin when expressed in mammalian cells (26). We therefore suggest that Asgard

ESCRT-IIIs are functionally most similar to CHMP1B in the eukaryotic ESCRT-III

system.

The biological function of Asgard ESCRTs has yet to be defined. The findings that Loki

ESCRT-III filaments can bind both DNA and membranes raises a few attractive

possibilities. First, given the established role of ESCRTs in cell division in TACK

archaea and in eukaryotes, it is possible that the ESCRT machinery manifest a basic

cell division machinery capable of orchestrating DNA segregation and membrane

splitting (22-25). In this regard, the ability of DNA to retain extensive membrane

deformation may represent a biological mechanism to avoid premature membrane

constriction during cell division. Second, the ESCRT complex may have been involved

in nuclear membrane formation during eukaryogenesis, by bringing together

membranes and DNA and facilitating DNA encapsulation by membranes. In this

respect, mammalian ESCRT-III proteins have been shown to mediate sealing of the

nuclear membrane post cell division and to regulate the attachment of heterochromatin

to the newly formed nuclear envelop (33-35). Moreover, we recently reported that Loki

CHMP 4-7 is targeted to the nucleus when over expressed in eukaryotic cells (26).

Finally, the membrane binding experiments performed here were done using

eukaryotic-like phospholipids, which are fundamentally different from archaeal lipids but

may resemble the lipid composition of the newly formed nuclear membrane. Therefore,

the ESCRT-III system encoded by Asgard archaea could have contributed to the

membrane remodeling that occurred during the formation of the phospholipids-based
nuclear membrane.
341
MATERIALS AND METHODS
Cloning
Open reading frames (ORFs) from Loki MAG sp. GC14_75, encoding for CHMP1-3 and
CHMP4-7 genes were obtained from NCBI (accession numbers KKK42122.1,
KKK44605.1, respectively) were amplified by PCR and sub-cloned into pSH21 vector
subsequently to the TEV cleavage tag that followed an N terminal polyhistidine and
MBP protein tags (a kind gift from Eyal Gur, BGU, Israel)(36). Plasmids were verified by
sequencing.
350
Protein Expression and Purification
pSH21 plasmids containing Loki CHMP1-3 or CHMP4-7 were expressed Escherichia
coli BL21 strain and incubated in LB 50 µg/mL Ampicillin at 37°C until reaching O.D. of
0.6. Then, induction with 0.5 mM IPTG was performed and the cultures were grown at
30°C for 3-4 hours. Cells were then resuspended and lysed by sonication in buffer A (25
mM Hepes pH= 8, 500 mM NaCl, 10% glycerol, 15 mM 2-mercaptoethanol, 10 mM
imidazole) supplemented with protease inhibitor cocktail (cOmplete ultra-tablets, EDTA-
free, Roche) and 500 µg/ml DNase I (10104159001; Roche Diagnosis GmbH, Manheim,
Germany), and the supernatant was clarified by centrifugation (25,000 g, 40 mins, 4°C).
For CHMP4-7 purification, supernatant was subjected to HisPur™ Ni-NTA Resin
(88222; Thermo Fisher Scientific, Weltham, MA) column and eluted in buffer B (buffer A
supplemented with 50 mM imidazole). Protein tags were then removed using a TEV
protease (1 mg/10 mg protein) in buffer C (50mM Tris-HCL pH=7.4, 10% glycerol, 500
mM NaCl, 15 mM 2-mercaptoethanol). Finally, a second Ni-NTA step was performed
using buffer C and the flow-through was collected.
For CHMP1-3 purification, supernatant was subjected to Ni-NTA column and eluted in
buffer B. Samples were then subjected to amylose resin (E8021; NEB, Frankfurt,
Germany) in buffer D (20mM Tris pH=7.4, 500 mM NaCl, 10mM β-Mercaptoethanol,
10% Glycerol) and eluted in buffer D supplemented with 10mM maltose. Protein tags

were then removed using a TEV protease (1 mg/10 mg protein) in buffer C (50mM Tris-

HCL pH=7.4, 10% glycerol, 500 mM NaCl, 15 mM 2-mercaptoethanol). Finally, a

second Ni-NTA step was performed using buffer C and the flow-through was collected.

Purified proteins were verified my mass spectrometry.

For polymerization, proteins were incubated in buffer E (25mM Tris pH=7.6, 50 mM KCl)

for 5 hours at RT at the indicated conditions. For DNA experiments,

ATCCACCTGTACATCAACTCGCCCGGCGGCTCGATCAGCG (40 bases probe), ss or

ds were used at the indicted concentrations. DNase I (10104159001; Roche Diagnosis

GmbH, Manheim, Germany) was added at 20mg/ml concentrations when indicated.

Preparation of small unilamellar vesicles (SUVs)

Chloroform Lipid solutions (DOPC or DOPS) were purchased from Avanti Polar Lipids

Inc. (Albaster, AL, USA catalog number 850375, 840035). Lipid mixtures of

DOPC/DOPS 1:1 molar ratio and pure DOPC were prepared at a total concentration of

1 mg/ml in Tris 20 mM, KCl 50 mM, pH 7.8 were prepared. Each lipid suspension was

extruded 20 times via a 100 nm polycarbonate membrane using a mini-extruder (Avanti

Polar Lipids, Alabaster, AL, USA). In the final stage, the vesicles were mixed with the

Asgard Loki ESCRT-III filaments. All experiments were done with 8 µM ESCRT-III and

124 µM vesicles in a buffer of Tris 25 mM, KCl 50 mM, at pH 7.6.

Negative Stain grid preparation and imaging

The negative stain samples for TEM analysis were prepared in the following way: 300

mesh copper grids (Ted Pella, Prod No. 01813-F) were glow discharged to enhance

hydrophilicity of their surface. Next, 2.5 uL of the sample was applied on to the grid and

the excess liquid was blotted with filter paper after 1 minute. The grid was dried in air for

1 minute, following by applying 5 uL of uranyl acetate 2% solution (SPI CAS# 6159-44-

0) for negative staining to increase the sample contrast. Next, the grid was blotted ones

more to remove the excess uranyl acetate. Finally, the grid was dried in air before

insertion into the microscope. The imaging of the samples was performed with Thermo

Fisher Scientific Talos F200C transmission electron microscope operating at 200 kV.

The images were taken with Ceta 16M CMOS camera.

Cryo-electron microscopy

Sample preparation: 3 μL of proteins / proteins-phospholipids mix samples were gently

deposited on glow discharged Quantifoil R 1.2/1.3 holey carbon grids (Quantifoil Micro

Tools GmbH, Germany). Samples were manually blotted for four seconds at room

temperature and vitrified by rapidly plunging into liquid ethane using a home-built

plunging apparatus. The frozen samples were stored in liquid nitrogen until imaging.

Micrographs acquisition: Samples were loaded under cryogenic conditions and

imaged in low dose mode on a FEI Tecnai F30 Polara microscope (FEI, Eindhoven)

operated at 300 kV. Micrographs were collected using SerialEM, at a calibrated pixel

size of 1.1 Å by a K2 Summit direct electron detector fitted behind an energy filter

(Gatan Quantum GIF) set to ±10 eV around zero-loss peak with total electron dose of

80 ē/$Å^2$. Each dose-fractionated movie had 50 frames, micrographs sums were aligned

in SerialEM.

Cryo-tomography. Micrographs were acquired with a FEI Tecnai F30 Polara

microscope (FEI, Eindhoven) operated at 300 kV, using a K2 Summit direct electron

detector fitted behind an energy filter (Gatan Quantum GIF), set to ±10 eV slit around

the zero-loss peak. Pixel size at the sample plane was 2.3 Å. The camera was operated

in counting mode at a dose rate of 6−7 ē/pixels/sec. Tilt series of 41 exposures from

−21° to +60° and back to -60° in 3° intervals and a total dose of ∼ 120 ē /$Å^2$ were

collected with Serial-EM. Tilt series were aligned, and tomograms were reconstructed

using eTomo (IMOD4.11 package) (37). Movies were generated in imageJ.

Measurements and statistical analysis

Filament length and diameter measurements were performed in Velox software version

3.5 (Thermo Fisher Scientific, Inc.).

Statistical analysis was performed in Graph Pad Prism version 9.00 (La Jolla, CA,

USA). Unless specified otherwise, comparison between two groups was calculated by

one way ANOVA * p- value ≤ 0.1, **p- value ≤ 0.01, ***p- value ≤ 0.001, ****p-

value ≤ 0.0001.

ACKNOWLEDMENTS

We thank the laboratories of Eyal Gur (BGU) and Eyal Arbely (BGU) for their practical

advice on proteins expression and purifications and for kindly sharing their reagents and

equipment with us. We also thank Itzik Mizrahi (BGU) and Ori Avinoam (WIS) for

constructive feedback on the project. Last, we thank all members of the Elia lab for

critical feedback throughout the project. The Elia laboratory is funded by the Israeli

Science Foundation (ISF) Grant no. 1323/18. ABG is grateful to the Israel Science

Foundation for financial support (grant 2101/20).

AUTHOR CONTRIBUTIONS

NE and DN conceptualized the project. NM performed and analyzed all experiments.

GH and AB, were in charge of experiments using SUVs. AU acquired negative stain

TEM images. RZ acquired and analyzed cryo-EM data. NE, DN, AB and RZ were

involved in experimental design and data analysis. NE wrote the manuscript. All authors

read and revised the manuscript.

Competing Interest Statement: We declare no competing interest

REFERENCES

1.   K. Zaremba-Niedzwiedzka *et al.*, Asgard archaea illuminate the origin of eukaryotic
cellular complexity. *Nature* **541**, 353-358 (2017).

2.   F. MacLeod, G. S. Kindler, H. L. Wong, R. Chen, B. P. Burns, Asgard archaea: Diversity,
function, and evolutionary implications in a range of microbiomes. *AIMS Microbiol* **5**, 48-
61 (2019).

3.   A. Spang *et al.*, Complex archaea that bridge the gap between prokaryotes and
eukaryotes. *Nature* **521**, 173-179 (2015).

4.   E. Neveu, D. Khalifeh, N. Salamin, D. Fasshauer, Prototypic SNARE Proteins Are Encoded
in the Genomes of Heimdallarchaeota, Potentially Bridging the Gap between the
Prokaryotes and Eukaryotes. *Curr Biol* **30**, 2468-2480 e2465 (2020).

5.   Z. Lu *et al.*, Coevolution of Eukaryote-like Vps4 and ESCRT-III Subunits in the Asgard
Archaea. *mBio* **11** (2020).

6.   T. Hatano *et al.*, Asgard archaea shed light on the evolutionary origins of the eukaryotic
ubiquitin-ESCRT machinery. *Nat Commun* **13**, 3398 (2022).

7.   Y. Caspi, C. Dekker, Dividing the Archaeal Way: The Ancient Cdv Cell-Division Machinery.
Front Microbiol **9**, 174 (2018).

8.   J. H. Hurley, ESCRTs are everywhere. *EMBO J* **34**, 2398-2407 (2015).

9.   Y. A. M. Alonso, S. M. Migliano, D. Teis, ESCRT-III and Vps4: a dynamic multipurpose tool
for membrane budding and scission. *FEBS J* **283**, 3288-3302 (2016).

10.   J. McCullough, L. A. Colf, W. I. Sundquist, Membrane fission reactions of the mammalian
ESCRT pathway. *Annu Rev Biochem* **82**, 663-692 (2013).

11.   C. Caillat, S. Maity, N. Miguet, W. H. Roos, W. Weissenhorn, The role of VPS4 in ESCRT-III
polymer remodeling. *Biochem Soc Trans* **47**, 441-448 (2019).

12.   S. Maity *et al.*, VPS4 triggers constriction and cleavage of ESCRT-III helical filaments. *Sci
Adv* **5**, eaau7198 (2019).

13.   B. E. Mierzwa *et al.*, Dynamic subunit turnover in ESCRT-III assemblies is regulated by
Vps4 to mediate membrane remodelling during cytokinesis. *Nat Cell Biol* **19**, 787-798
(2017).

14.   A. K. Pfitzner *et al.*, An ESCRT-III Polymerization Sequence Drives Membrane
Deformation and Fission. *Cell* **182**, 1140-1155 e1118 (2020).

15.   W. M. Henne, N. J. Buchkovich, Y. Zhao, S. D. Emr, The endosomal sorting complex
ESCRT-II mediates the assembly and architecture of ESCRT-III helices. *Cell* **151**, 356-371
(2012).

16.   I. Goliand *et al.*, Resolving ESCRT-III Spirals at the Intercellular Bridge of Dividing Cells
Using 3D STORM. *Cell Rep* **24**, 1756-1764 (2018).

17.   J. Guizetti *et al.*, Cortical constriction during abscission involves helices of ESCRT-III-
dependent filaments. *Science* **331**, 1616-1620 (2011).

18.   J. McCullough *et al.*, Structure and membrane remodeling activity of ESCRT-III helical
polymers. *Science* **350**, 1548-1551 (2015).

19.   S. Lata *et al.*, Helical structures of ESCRT-III are disassembled by VPS4. *Science* **321**,
1354-1357 (2008).

20.   B. P. Frohn, T. Hartel, J. Cox, P. Schwille, Tracing back variations in archaeal ESCRT-based
cell division to protein domain architectures. *PLoS One* **17**, e0266395 (2022).

21. J. Liu *et al.*, Bacterial Vipp1 and PspA are members of the ancient ESCRT-III membrane-
remodeling superfamily. *Cell* **184**, 3660-3673 e3618 (2021).
22. A. C. Lindas, E. A. Karlsson, M. T. Lindgren, T. J. Ettema, R. Bernander, A unique cell
division machinery in the Archaea. *Proc Natl Acad Sci U S A* **105**, 18942-18946 (2008).
23. R. Y. Samson, T. Obita, S. M. Freund, R. L. Williams, S. D. Bell, A role for the ESCRT
system in cell division in archaea. *Science* **322**, 1710-1713 (2008).
24. R. Y. Samson *et al.*, Molecular and structural basis of ESCRT-III recruitment to
membranes during archaeal cell division. *Mol Cell* **41**, 186-196 (2011).
25. G. Tarrason Risa *et al.*, The proteasome controls ESCRT-III-mediated cell division in an
archaeon. *Science* **369** (2020).
26. D. Nachmias *et al.* (2022) Asgard ESCRT-III and VPS4 reveal evolutionary conserved
chromatin binding properties of the ESCRT machinery. in *bioRxiv*
27. K. Azad *et al.* (2022) Structural basis of CHMP2A-CHMP3 ESCRT-III polymer assembly and
membrane cleavage. in *bioRxiv*.
28. Y. Olmos, The ESCRT Machinery: Remodeling, Repairing, and Sealing Membranes.
Membranes (Basel) **12** (2022).
29. B. Junglas *et al.*, PspA adopts an ESCRT-III-like fold and remodels bacterial membranes.
Cell **184**, 3674-3688 e3618 (2021).
30. H. C. Nguyen *et al.*, Membrane constriction and thinning by sequential ESCRT-III
polymerization. *Nat Struct Mol Biol* **27**, 392-399 (2020).
31. N. Talledge *et al.* (2018) The ESCRT-III proteins IST1 and CHMP1B assemble
around nucleic acids. in *bioRxiv*.
32. T. K. Gupta *et al.*, Structural basis for VIPP1 oligomerization and maintenance of
thylakoid membrane integrity. *Cell* **184**, 3643-3659 e3623 (2021).
33. G. H. Pieper, S. Sprenger, D. Teis, S. Oliferenko, ESCRT-III/Vps4 Controls
Heterochromatin-Nuclear Envelope Attachments. *Dev Cell* **53**, 27-41 e26 (2020).
34. Y. Olmos, L. Hodgson, J. Mantell, P. Verkade, J. G. Carlton, ESCRT-III controls nuclear
envelope reformation. *Nature* **522**, 236-239 (2015).
35. M. Vietri *et al.*, Spastin and ESCRT-III coordinate mitotic spindle disassembly and nuclear
envelope sealing. *Nature* **522**, 231-235 (2015).
36. Y. Elharar *et al.*, Survival of mycobacteria depends on proteasome-mediated amino acid
recycling under nutrient limitation. *EMBO J* **33**, 1802-1814 (2014).
37. D. N. Mastronarde, Automated electron microscope tomography using robust prediction
of specimen movements. *J Struct Biol* **152**, 36-51 (2005).

Fig 1. Loki CHMP4-7 protein self-assemble into filaments and helical tubes.

A. Negative- stain TEM micrographs of Loki CHMP4-7 homo-filaments. Representative

zoomed-out (left, scale = 200 nm) and zoomed- in (middle and right, scale = 100 nm)

images are shown. Orange arrows indicate filaments, red arrow indicates a helical tube.

B. Representative Cryo-EM micrographs of Loki CHMP4-7 homo-polymers. Left,

zoomed-out image (Scale = 100 nm), middle and right, zoomed-in images (Scale = 50

nm). Box in middle panel specify the zoomed-in region shown to the right. **C.** A

representative 2D class average of 502 particles extracted from CHMP 4-7 projections.

Scale = 100 Å. **D.** Measurements of Loki CHMP4-7 helical diameters (left, n = 37) and

length (right, n = 36) obtained from negative stain TEM images acquired from two

independent experiments. Averaged CHMP 4-7 helical tube diameter 44nm±30.9.

Fig 2. Loki CHMP4-7 and CHMP1-3 co-polymers assemble into helical tubes exhibiting different properties from CHMP 4-7 homo-polymers.

A. Representative Negative- stain TEM images of Loki CHMP4-7 homo-polymers (upper panels) and CHMP4-7-CHMP1-3 co-polymers (bottom panels, 2:1 molar ratio). Zoomed- out (left, scale = 1 µm) and zoomed- in (right, scale = 50 nm) images are shown. Yellow asterisks indicate helical tubes. **B**. Measurements of CHMP4-7-CHMP1-3 hetero-filaments diameters (left) and length (right) at the indicated molar ratios obtained from negative stain TEM images. Averaged diameters (nm): 2:1 ratio, 31±7.7; 4:1 ratio 28±1.6, 6:1 ratio 32±3.8. n = 25 for each condition.

Fig 3. Organization into helical tubes is increased in the presence of DNA

A. Negative- stain TEM micrographs of CHMP 4-7-CHMP 1-3 co- polymers (2:1 molar ratio) in the presence of 4.6 µM 40 bases oligonucleotides (ssDNA or dsDNA, as indicated). Representative zoomed- out (left, Scale = 1 µm) and zoomed- in (right, Scale = 50 nm) images are shown. Helical tubes are colored in yellow in zoomed out images to highlight the observed increase in tubes (raw images are provided in Sup Fig 2A). **B.** Representative Cryo-EM images of CHMP 4-7– CHMP 1-3 co-polymers (molar ratio, 4:1) in the presence of ssDNA. Scale = 200 nm. **C.** Negative- stain TEM micrographs of CHMP 4-7 homo-polymers in the presence of 4.6 µM 40 bases ssDNA. Representative zoomed- out (left, Scale = 1 µm) and zoomed- in (right, Scale = 50 nm) images are shown. Helical tubes are colored in yellow in zoomed out images to highlight the observed increase in tubes (raw images are provided in Sup Fig 2B). **D-E** Diameter (**D**) and length (**E**) measurements of helical tubes formed by homo- and the co-polymers (2:1 ratio) in the presence of nucleic acids obtained from negative stain TEM images. Averaged diameters (nm): CHMP4-7 ssDNA 43.9±33, n=21; CHMP4-7-CHMP1-3 2:1 dsDNA 51.9±94, n=44; CHMP4-7-CHMP1-3 2:1 ssDNA 43.9±58, n=48). **F.** Representative Negative- stain TEM images of co-polymers (molar ratio 2:1) upon addition of 20 mg/ml DNase I during (middle panel) or after (right panel) polymerization. Helical tubes are colored in yellow. Scale =200 nm. Data was reproduced in four independent experiments.

A CHMP 4-7 - CHMP 1-3 ssDNA + SUVs

B CHMP 4-7 - CHMP 1-3 + SUVs

ssDNA   No DNA

C

D

Fig 4. Loki ESCRT-III filaments induce membrane remodeling

A-B. Representative Negative- stain TEM (**A**) and Cryo-EM images (**B**) of CHMP4-7-CHMP1-3 (molar ratio, 4:1) polymerized in the presence of ssDNA (10 μM) and incubated with SUVs (DOPC:DOPS, 1:1 ratio). Scale = 100 nm. Arrows in **B** specify internalized vesicles. Reconstructed tomograms of filaments collected from the same grid are provided in Sup. movies 1 and 2. **C**. Percentages of the different types of vesicle-tube interactions observed under the specified conditions. Left panel, represented negative-stain images of the four types of interactions quantified. Outside, attachment of SUV to the outside of the helical tube (type 1); Docking, binding of the vesicle to the tip of the helical tube (type 2); Inside, invagination of the SUV to the helical tube (type 3); Remodeling, deformation of both the vesicles and the ESCRT-III helical tube (type 4). Data was quantified from negative stain imaged obtained in 3 independent experiments (n = CHMP4-7 ssDNA, 124; CHMP4-7-CHMP1-3 ssDNA, 159; CHMP4-7, 197; CHMP4-7-CHMP1-3, 161). Scale = 100 nm. Chi square statistical analysis show a significant difference in the distributions of CHMP4-7 ssDNA vs CHMP4-7 (****p- value ≤ 0.0001), CHMP4-7-CHMP1-3 ssDNA vs CHMP4-7-CHMP1-3 (****p- value ≤ 0.0001), and CHMP4-7 ssDNA vs CHMP4-7-CHMP1-3 ssDNA (****p-value ≤ 0.0001). No significance was found in the distributions of CHMP4-7 vs CHMP4-7-CHMP1-3. **D**. A suggested model for Loki ESCRT-III mediated vesicle remodeling. Vesicles dock at the tip of the ESCRT-III tube. Docking allows the vesicle to directly interact with the interior of the tube, leading to deformation and invagination of the vesicle by the ESCRT-III filament. This process occurs repeatedly leading to the formation of a complex membrane network inside the helical tube, which pushes the ESCRT-III tube from the inside and ultimately causing deformation and remodeling of the tube. This stage is inhibited in the presence of DNA, resulting the accumulation of ESCRT-III-SUV interactions in all the stages that preceded this complete remodeling stage.