The Asgard archaeal ESCRT-III system forms helical filaments and remodels eukaryotic membranes, shedding light on the emergence of eukaryogenesis

Nataly Melnikov[1,2], Benedikt Junglas[3$], Gal Halbi[4$], Dikla Nachmias[1,2], Erez Zerbib[1,2], Alexander Upcher[5], Ran Zalk[5], Carsten Sachse[3,6*], Ann Bernheim-Groswasser[4,5], Natalie Elia[1,2*]

Department of Life Sciences, Ben-Gurion University of the Negev, Beer Sheva 84105, Israel

National Institute for Biotechnology in the Negev (NIBN), Ben-Gurion University of the Negev, Beer Sheva 84105, Israel

Ernst-Ruska Centre for Microscopy and Spectroscopy with Electrons, ER-C-3/Structural Biology, Forschungszentrum Jülich, 52425 Jülich, Germany

Department of Chemical Engineering, Ben-Gurion University of the Negev, Beer Sheva 84105, Israel

Ilse Katz Institute for Nanoscale Science and Technology, Ben Gurion University of the Negev, Beer Sheva 84105, Israel

Department of Biology, Heinrich Heine University, Universitätsstr. 1, 40225 Düsseldorf, Germany

$ these authors contributed equally to this work.

* Corresponding authors:

Structural analysis:

Carsten Sachse

Phone: +49 2461/61-2030

Email: c.sachse@fz-juelich.de

All other subjects:

Natalie Elia

Phone: 972-8-6428735

Email: elianat@post.bgu.ac.il

ABSTRACT

The ESCRT machinery mediates membrane remodeling in numerous processes in cells

including cell division and nuclear membrane reformation. The identification of ESCRT

homologs in Asgard archaea, currently considered the closest ancestor of eukaryotes,

implies a role for ESCRTs in the membrane remodeling processes that occurred during

eukaryogenesis. Yet, the function of these distant ESCRT homologs is mostly

unresolved. Here we show that Asgard ESCRT-III proteins of the Lokiarcheota self-

assemble into helical filaments, a hallmark of the ESCRT system. We determined the

cryo-EM structure of the filaments at 3.6 Å resolution and found that they share features

of bacterial and eukaryotic ESCRT-III assemblies. Markedly, Asgard ESCRT-III

filaments bound and deformed eukaryotic-like membrane vesicles. Oligonucleotides

facilitated the assembly of ESCRT-III filaments and tuned the extent of membrane

remodeling. The ability of Asgard archaeal ESCRTs to remodel eukaryotic-like

membranes, which are fundamentally different from archaeal membranes, and the

structural properties of these proteins places them at the junction between prokaryotes

and eukaryotes, substantiating a role for ESCRTs in eukaryogenesis.

GRAPHICAL ABSTRACT

INTRODUCTION

The Asgard superphyla, manifest a unique branch that, according to current
evolutionary theories, is considered to be the closest prokaryotic relative of eukaryotes.
All Asgard archaeal species were found to encode for Eukaryotic Signature Proteins
(ESPs), that carry cellular functions that until recently were considered as exclusive to
eukaryotes (1-3). Thus, Asgard proteins potentially hold the core functions of ancient
eukaryotic cellular machines. Of particular interest, are ESPs with membrane
remodelling capabilities, as these proteins may contribute to the establishment of
cellular compartmentalization during eukaryogenesis. Indeed, Asgard archaea encode
for several eukaryotic membrane remodelling machineries including the SNARE
complex, BAR domains proteins and the ESCRT machinery (1, 3, 4). Among these
machineries, ESCRTs are the most conserved and are encoded by all Asgard species
discovered up to date, suggesting that they carry essential functions in these unique
prokaryotes (5, 6).

The ESCRT machinery constitute one of the most robust and versatile cellular apparatus for membrane constriction and fission. Proteins of the machinery are encoded by all domains of life and were shown to mediate membrane remodelling in a wide range of eukaryotic cellular membranes including plasma membrane, nuclear membrane and endocytic membranes (7, 8). Moreover, ESCRT-mediated processes span a large spectrum of length scales, from less than 100 nm for vesicle release to up to 1 μm for cell division. How the ESCRT complex mediate membrane remodelling in this large landscape of membrane sources and length scales is mostly unknown (9).

The eukaryotic ESCRT system is composed of five subfamilies, i.e. ESCRT 0-III and the AAA-ATPase VPS4. Within this complex, ESCRT-III (named CHMPs in animal cells) and VPS4 manifest the minimal unit required for driving membrane fission (9, 10). According to current models, membrane constriction and scission are mediated by polymerization of ESCRT-III proteins into helical hetero-filaments and their remodelling by VPS4, seen in vitro and in cells (10-19). However, the large number of proteins within the ESCRT-III subfamily —twelve in humans and eight in yeast - have challenged mechanistic studies of the basic function of the machine.

Reduced, simplified ESCRT-III systems have been identified in prokaryotes including in bacteria and archaea (named Vipp1 and PspA in bacteria and CdvB in archaea) (7, 20, 21). While the bacterial homologs were shown to remodel membranes, no VPS4 was identified in this system so far. ESCRTs encoded by TACK archaea were shown to participate in cell division, suggesting functional similarities between the eukaryotic and archaeal ESCRT systems (22-25). Within the archaeal domain the Asgard ESCRT system is considerably closer to the eukaryotic system. While all other archaea encode only for ESCRT-III and VPS4 homologs, Asgard archaea encode for homologs of the complete ESCRT system (ESCRT-I-III and VPS4) (6, 7). Additionally, the sequence of Asgard ESCRT-III/VPS4 proteins is more closely related to those of eukaryotes than to other archaeal ESCRT systems (named CDV) (5, 20). Lastly, VPS4 homologs of Asgard archaea were shown to functionally interact with eukaryotic ESCRTs in both yeast and mammalian cells (5, 26). Notably, the Asgard ESCRT-III subfamily is

substantially reduced compared to eukaryotes with only two ESCRT-III proteins (named

here CHMP1-3 and CHMP4-7) encoded in some species including Loki and Heimdall

archaeota (1, 5, 7, 26). Hence, the Asgard ESCRT system may constitute a more basic

version of the ESCRT machinery, which holds the core capabilities of eukaryotic

ESCRTs, and could have been involved in membrane remodelling processes that

occurred during eukaryogenesis. Yet, the function, organization, and properties of

Asgard ESCRT-III complexes as well as their ability to remodel membranes are mostly

unknown.

In this work, we provide the first evidence that Asgard ESCRT-III complexes share

functional similarities with eukaryotic ESCRTs. Using purified ESCRT-III homologs

encoded by the most abundant Asgard phyla, Lokiarchaeota (Loki), we show that the

Asgard ESCRT-III protein CHMP4-7 can self-assemble, on its own and in the presence

of CHMP1-3, into helical tubes that resemble those of eukaryotes (15, 18, 19, 27). The

frequency of helical tubes was increased in the presence of short ssDNA oligos, while

no filaments could be observed upon DNAse treatment. Moreover, we determined the

cryo-EM structure of helical CHMP 4-7 tubes in the presence of CHMP 1-3 and DNA at

a resolution of 3.6 Å. The 36 nm wide tubular assemblies are built from protomers

adopting the canonical ESCRT-III fold while they share characteristic intersubunit

contacts with bacterial and eukaryotic ESCRT-III polymers. Finally, we found that Loki

ESCRT filaments, formed under any of these conditions, are able to bind and deform

small unilamellar vesicles (SUVs) comprised of eukaryotic-like synthetic phospholipids.

Hence, Loki ESCRT-III assemble into helical filaments that are affected by

oligonucleotides and are capable of remodelling eukaryotic-like membranes.

RESULTS

Current phylogenic models suggests that eukaryotes integrate from the archaeal Asgard

clade (Fig. 1A). ESCRT-III proteins encoded in eukaryotes polymerize into helical

filaments - a hallmark of ESCRT-III organization that is thought to be essential for their

function in membrane remodeling (8, 28). Polymerization is mediated by the snf7

domain that is conserved in both archaea and eukaryotes, but the polymerization

properties of their ancestor Asgard ESCRT-III homologs has not been examined (5, 20,

26). To investigate the assembly properties of Asgard ESCRT-III proteins, we purified

recombinant versions of the ESCRT-III proteins CHMP4-7 and CHMP1-3, encoded by

the Loki GC14_75 strain (KKK44605.1 and KKK42122.1, respectively). Purified proteins

were then subjected to polymerization reaction (see methods) and visualized by

negative-stain TEM and cryo-EM. Recombinant full-length CHMP4-7 spontaneously

self-assembled into helical filaments (Fig. 1B), while no evidence for polymerization

could be detected for purified CHMP1-3 using a range of experimental conditions (Fig.

S1). The overall shape of CHMP4-7 filaments, seen by negative-stain EM, resembled

those previously described for eukaryotic or bacterial ESCRT-III filaments (15, 18, 19,

29). CHMP4-7 polymers were predominantly seen as long, thin and curved filaments or

as rod-shaped tubes (Fig. 1B, see arrows and Fig. 1C). A Moiré pattern of parallel,

periodic stripes was detected in the CHMP4-7 tube structures by cryo-EM, supporting a

helical tube organization (Fig. 1C, D). The outer diameter of the CHMP4-7 tubes was

variable, ranging between 35 to 55 nm (averaged diameter 44 ± 30.9 nm). Tube length

was typically between 300 - 450 nm (Fig. 1E). The diameter of Loki CHMP4-7 helical

tubes was slightly larger than the diameters measured for human CHMP1B (~25nm) or

the bacterial ESCRT-III homologs PspA (~21 nm), but was in the range of diameters

measured for human CHMP2A-CHMP3 (38 - 43 nm) (18, 19, 29). Collectively, these

data indicate that Loki CHMP 4-7 spontaneously self-assemble into the typical helical

tube organization described for the ESCRT-III complex.

Next, we examined the effect of CHMP1-3, which did not polymerize on its own, on the

ESCRT-III filaments. To this end, full lengths Loki CHMP4-7 and CHMP1-3 were co-

incubated at different ratios and subjected to polymerization reaction. While no

polymers were seen at 1:1 CHMP4-7-CHMP1-3 ratio, helical tubes could be readily

observed at 2:1, 4:1 and 6:1 ratios (Fig. 2 and S2A). Notably, helical tubes assembled in

the presence of both CHMP4-7 and CHMP1-3 were morphologically distinct from helical

tubes assembled using only CHMP4-7. First, their outer diameter was significantly

smaller and less diverse (28 ± 1.6 nm vs 44 ± 30.9 nm for CHMP4-7/CHMP1-3 4:1 ratio

174 and CHMP4-7 alone, respectively) (Fig. 2B, left panel). Second, helical tubes formed in

175 the presence of both CHMP4-7 and CHMP1-3 were significantly longer than those

176 formed by the CHMP4-7 homo-polymer (>X1.5 folds increase for 2:1 and 4:1 ratios)

177 (Fig. 2B, right panel). Third, the morphology of the CHMP4-7/CHMP1-3 helical tube was

178 slightly different: tubes appeared to be more curved and periodic stripes, were readily

179 detected in negative stain images, suggesting that packing of the helical tube is less

180 tight in the presence of CHMP1-3 (Fig. 2A). Additionally, unpacked filaments,

181 resembling a telephone cord were observed, further supporting a continuous helical

182 tube assembly (Fig. S2B). The phenotypic changes observed in the presence of

183 CHMP1-3 were not induced upon substituting CHMP1-3 with purified GFP or by

184 increasing the concentration of CHMP4-7, suggesting that they are specifically

185 contributed by CHMP1-3 (Fig. S2C). Therefore, the Loki ESCRT-III protein CHMP4-7

186 self-assemble into helical filaments, with CHMP1-3 affecting the properties of these

187 filaments. These findings are in line with previous observations obtained for human

188 ESCRT-III CHMP1B helical filaments, which also formed a narrower, more uniform

189 tubes in the presence of IST1(30). The realization that both human and Asgard ESCRT-

190 III proteins assemble into helical filaments that can adopt a narrower, more uniform tube

191 shape in the presence an additional subunit, suggests that this is a conserved property

192 of the ESCRT-III system.

194 Recent work from our laboratory showed that purified Loki CHMP4-7 binds short DNA

195 oligonucleotides (26). Human CHMP1B, exhibited similar properties and a recently

196 solved cryo-EM structure of human CHMP1B/IST1 filaments demonstrated direct

197 interactions between the ESCRT-III filament and nucleic acids (26, 31). We, therefore,

198 set to examine the effect of short DNA oligonucleotides on the self-assembly properties

199 of Asgard ESCRT-III filaments. A five-fold increase in the occurrence of helical tubes

200 was observed for CHMP4-7/CHMP1-3 helical tubes that were assembled in the

201 presence of ssDNA, in negative stain TEM images (Fig. 3A-C and Fig. S3A). Addition of

202 dsDNA, comprising the same sequence, did not induce this effect (Fig. S3B). Incubating

203 CHMP1-3 with ssDNA, did not induce polymerization, further supporting the inability of

204 CHMP1-3 to self-assemble in the absence of CHMP4-7 (Fig. S3C). The presence of

 7

DNA affected the properties of CHMP4-7/CHMP1-3 helical tubes. Helical tubes formed

in the presence of ssDNA were wider and slightly longer (averaged diameters: 40±4 nm

vs. 29.4±4.6 nm; averaged lengths: 777.5 ±210 nm vs. 535 ±119 nm for helical tubes

polymerized with and without ssDNA, respectively) (Fig. 3D). Moreover, filaments were

less curved, and the periodic stripes observed in negative stain images could not be

observed (Fig. 3A). Addition of ssDNA also increased the frequency of helical tubes

formed by CHMP4-7 alone, but to a lesser extent (three folds), without affecting their

characteristic length or diameter (Fig. 3B-D and Fig. S3D). Helical tube formation was

completely abolished for both CHMP 4-7/CHMP1-3 and CHMP4-7 filaments in the

presence of DNase, resulting aggregates formation, further supporting a role for DNA in

ESCRT-III helical tube assembly (Fig. 3E and Fig. S3E).

Given the higher frequency of ESCRT-III helical filaments obtained for CHMP4-

7/CHMP1-3 filaments in the presence of ssDNA, we set out to resolve the cryo-EM

structure of the filaments. In our cryo-EM micrographs, we found two types of filaments:

straight tube/rod-like polymers with variable diameters and thinner highly curved

polymers (as described above) (Fig. 4A, Fig. S4A). Firstly, the curved filaments had

diameters of 6 to 15 nm and showed distinct features in their 2D class averages such as

saw-tooth-like spikes emanating sideways from the filament including, in some cases, a

central smooth line (Fig. S4B). Nevertheless, we were unable to generate reliable 3D

reconstructions, likely due to their variable curvature and remaining heterogeneity.

Secondly, the rod-shaped filaments had diameters ranging from 30 to over 45 nm. The

2D class averages showed well-defined spikes at the filament rims and a Christmas-

tree like pattern (Fig. 4B), overall closely resembling PspA rods (29). We were able to

determine the structure of 36 nm PspA-like rods at a global resolution of 3.6 Å and a

local resolution ranging from 3.0 Å at the inner rod wall to 6.0 Å at the outer tube wall

(Fig. 4C-E). The rods are arranged by helical symmetry with 1.05 Å rise and 173.9 °

rotation. In the lumen of the rods, weak cylindrical density was found close to the inner

wall of the rods at approx. 13 nm radial position (Fig. S4C). We relied on the strong

density to build a model for Loki CHMP4-7 α1-5 into the density (Fig. 5A-B), while no

density was found for helices α0 and α6. Despite the presence of CHMP1-3 and ssDNA

in the sample and the described effects on frequency and morphology of the
assemblies, we could not identify any corresponding density in the cryo-EM map.
CHMP1-3 is, therefore, not a regular member of the symmetric rod assemblies while it
may be part of the curved filaments, involved in nucleation or occasionally bound to the
rod assemblies. The structured elements of CHMP 4-7 adopt the canonical ESCRT-III
fold: α1-3 form an extended hairpin, connected to α4 via a short linker (E117-P120 Fig.
5B inset). Helix α4 is separated in two parts by a short loop at G146 and connected to
α5 by an extended linker (I154-E159). The monomers are arranged in a j+4 repeat,
where α5 of chain j packs perpendicularly against the tip of the hairpin of j+4 (Fig. 5C),
forming the typical ESCRT-III assembly motif known from IST1B/CHMP1B (18). In the
assembly, helices α1-4 make up the wall of the rods, while α5 together with the tip of
the α1+2 hairpin forms the spikes on the outside of the wall (Fig. S4D and 5D).
Consequently, although unresolved, α6 is likely located outside the rods while α0 points
into the lumen perhaps as part of the weak density found inside the rods. The
electrostatic surface of the rod assembly reveals a distinct polarity with one negatively
and one positively charged end (Fig. S4E). The same interfaces likely also stabilize the
layers of the rod assembly by electrostatic interactions.
253
Loki CHMP4-7 adopts the canonical ESCRT-III fold in the open form and the assembly
resembles typical features of ESCRT-III tubes including the conserved motif between
α1/2-α5. Comparing the CHMP4-7 rods with ESCRT-III tube-like assemblies of other
domains of life, we found that their architecture is most similar to the bacterial
assemblies that have a α5-hairpin structure emanating from the tubular wall and giving
rise to the characteristic spikes in the projections (Fig. 5E top). In contrast, in eukaryotic
assemblies such as IST1 and CHMP1B the α5-hairpin structure does not protrude from
the assembly, thereby forming smooth tubes. In CHMP4-7 rods of Asgard, α5 only
interacts with a single hairpin at the bottom end, which is a common feature in
eukaryotic ESCRT-III assemblies, while in bacteria top and bottom ends of α5 are
interacting with two hairpins connecting two layers of the assembly (Fig. 5E bottom).
Additionally, on the monomer level, CHMP4-7 more closely resembles the eukaryotic
ESCRT-III proteins: firstly, the bacterial ESCRT-IIIs have more extended α0 and α2+3

helices (corresponding to aa 10-20, aa 108-123 in the consensus sequence Fig. S4F).

Secondly, the linker between α3 and α4 is longer and contains more helix breaking

residues in the bacterial ESCRT-IIIs (i.e., G156, G157 and G159 in Vipp1). Therefore,

the Loki CHMP4-7 assembly more closely resembles the bacterial configuration with the

characteristic spike while the Loki CHMP4-7 monomer adopts a conformation closer to

previously determined eukaryotic structures including the α1/2-α5 interaction motif.

Eukaryotic and bacterial ESCRT-IIIs were shown to bind and deform membranes *in*

vitro and in cells (*21, 28, 29, 32*). To determine whether ESCRTs encoded by Asgard

archaea carry the ability to remodel membranes, we incubated Loki ESCRT-III pre-

assembled filaments with small unilamellar vesicles (SUVs, ~100 nm) composed of

negatively charged eukaryotic-like phospholipids (1:1, PC:PS). Under these conditions,

ESCRT-III filaments, were found to intimately interact with SUVs under any of the

conditions tested (including in the presence of ssDNA) (Fig. 6A, B and Fig. S5). We

identified four main types of ESCRT-III-SUV interactions: (1) Attachment of the SUV to

the outside surface of the tube, which was associated with flattening of the vesicle at the

filament-vesicle interface; (2) Docking of the SUV at the tip of the helical tube; (3)

Invagination of the SUV to the interior of the ESCRT-III tube; and (4) Dramatic

remodeling of both the SUV and the ESCRT-III tube (see examples in Fig. 6C, left

panel). Combinations of several types of interactions on the same filament were also

observed, indicating that these interactions are not mutually exclusive. In addition, we

occasionally observed a docked SUV invaginating into the helical tube (Fig. 6B, left

panel, Fig. 6D and sup movies 1, 2), suggesting that these two types of interactions

represent successive steps. Invaginations of more than one vesicle to the same tube

could be observed, giving rise to the formation of a complex membrane system with

several vesicles internalized into one another (Fig 6B). The characteristic periodic

stripes of the ESCRT-III filament could often be detected at the exterior of the filament-

vesicle complex, suggesting that the observed morphologies result from internalization

of vesicles into the ESCRT-III helical tube followed by deformation of the tube. Loki

ESCRT III filaments also interacted with neutrally charged SUVs (100% PC) but to a

lesser extent, (27% vs. 15% of the total SUVs interacted with ESCRT-III tubes upon

incubation with PC:PS vs PC vesicles, respectively), predominantly exhibiting

attachment of the SUV to the exterior of the helical tube (Fig. S6). Therefore, Loki

ESCRT-III filaments associate with membrane vesicles and deform negatively charged

vesicles via the interior of the tube (see proposed model in Fig. 6E).

Last, we asked whether the composition of ESCRT-III helical tubes, described here,

affect their capabilities to remodel membranes. To this end, we quantified the

frequencies of the different types of filament-vesicle interactions we defined, at the

various assembly conditions. No difference in frequencies were observed for helical

filaments that assembled in the presence or absence of CHMP1-3 (CHMP4-7/CHMP1-3

vs CHMP4-7), suggesting that CHMP1-3 does not significantly contribute to membrane

interactions. The presence of DNA, however, affected the distributions of ESCRT-III-

SUV interactions, as follows (Fig. 6C): First, the association of vesicles with the outside

surface of the helical tube (outside, type 1) was considerably enriched (No DNA, <15%;

with DNA, >35%). Second, interactions with the tip of the filament (docking, type 2) and

invagination of the vesicle (inside, type 3) were significantly enriched. Finally, the most

significant difference was the abundance of the complete remodeling events

(remodeling, type 4) (no DNA ~80%, with DNA, ~5%;), suggesting that the presence of

DNA inhibits this terminal tube-vesicle interaction stage. The effect of DNA on Loki-

ESCRT-III induced membrane remodeling appeared to be irreversible, as addition of

DNase to pre-established tube-SUV complexes did not lead to extensive membrane

remodeling (Fig. S5C). Therefore, Loki ESCRT-III filaments are capable of interacting

and remodeling lipid membranes, both in the presence and absence of DNA, but the

extent of remodeling appear to be attenuated by the presence of DNA.

DISCUSSION

In this work, we provide the first evidence to show that, Asgard ESCRT-III proteins

assemble into helical tubes that resemble those of their eukaryotic homologs and

remodel membranes. These findings confirm that ESCRT homologs encoded by Asgard

archaea are functionally related to their eukaryotic homologs. We further provide data to

support a role for DNA in the assembly of Asgard ESCRT-III filaments. Finally, by using

SUVs composed of eukaryotic phospholipids, which are fundamentally different from

archaeal lipids, we demonstrate that the Asgard ESCRT-III could, in-principle, contribute

to the membrane remodeling processes that occurred during eukaryogenesis and gave

rise to the complex phospholipids-based endomembrane system of eukaryotes.

Our data point to similarities between the human and Asgard ESCRT-III systems.

Firstly, the nature of self-assembly appear to be conserved. Similar to the human

CHMP1B/IST1 complex, Loki CHMP4-7 was found to polymerize alone or in the

presence of CHMP1-3, with the former exhibiting a wider, more variable diameter (30).

What controls the diameter of the ESCRT-III helical tube is currently unknown. Notably,

while IST1 was found to form a layer on the outside of the CHMP1B tube, CHMP1-3

could not be detected in our cryo-EM structure suggesting that it obtains a different

mode of operation. Regardless of the mechanism, the realization that both human and

Asgard ESCRT-III filaments can adopt a wide range of diameters that can be tuned by

the addition of another ESCRT-III components, strongly suggest that this is a conserved

feature of the ESCRT-III system. Therefore, a basic feature of the ESCRT-III system

may be their ability to assemble into adaptable filaments that can be tuned, by

additional ESCRT-III proteins, to adopt specific shapes. Such a model can explain how

the ESCRT-III system executes the large repertoire of membrane remodeling processes

in cells, which greatly vary in scales and topologies (7, 8).

The determined cryo-EM structure of CHMP4-7 helical filaments reveals the canonical

ESCRT-III fold made of 6 α-helices including the conserved α1/2-α5 packing equally

present in Loki archaea. For eukaryotes and Loki archaea, we detected a number of

residue omissions in helices α2+α3 and the linker between α3 and α4, whose

associated properties appear to be dispensable in the context of ESCRT evolution and

may be responsible for functional diversification of different isoforms. The basic

structural arrangement of the molecule in the CHMP4-7 assembly resembles that of

previously determined bacterial ESCRT-III proteins PspA and Vipp1 (29, 32). In both

cases, the hairpin formed by α1/α2 emanates from the assembly giving rise to

characteristic spikes in the EM images. Another feature that is closer to the determined
eukaryotic assemblies CHMP1B and IST1 is that helix α5 contacts only a single α1/2
hairpin as opposed to contacting two α1/2 hairpins described for bacterial ESCRT-III
proteins (18). This feature makes eukaryotic as well as Loki rods less tightly packed
potentially offering additional access for binding partners and thus opening up the
possibility to functional modulation and regulation of membrane remodeling by other
isoforms of the ESCRT-III family.
367
The topology of membrane binding, described here for Loki ESCRT-IIIs, involves
invagination of the membrane by the ESCRT-III tube. Although the conventional
topology attributed to the ESCRT complex is reversed, as described for the CHMP2A-
CHMP3 co-polymer (19, 27), membrane binding via the interior of the tube has been
reported for the human CHMP1B/IST1 polymer (30), and for the bacterial ESCRT-III
homologs Vipp1 and PspA (29, 32). Consistently, in the cryo-EM structure of Loki
ESCRT-III the N terminus, which is associated with membrane binding, is facing the
interior of the helical tube. Whether internal membrane binding topology represents the
preferred topology of ancient ESCRTs or is a consequence of the, in vitro, experimental
system is yet to be determined. Regardless, these similarities suggest that the ESCRT-
III membrane remodeling properties reported here are a common property of the
ESCRT system in all domains of life.
380
Our results highlight DNA as an essential Loki ESCRT-III partner. Self-assembly of the
Loki ESCRT-III polymers into high-ordered helical tubes was enhanced in the presence
of short ssDNA oligonucleotides, and helical tubes could not be found in the presence of
DNase. While the determined cryo-EM structure of Loki ESCRT-III, reported here, does
not reveal any density of DNA, a cryo-EM structure of CHMP1B/IST1 filaments shows
the presence of DNA in the inner side of the tube and nucleotides were detected in the
cryo-EM structure of the bacterial ESCRT-III Vipp1 (31, 32). Perhaps the DNA reside in
the lumen of the rods seen in the unresolved weak cylindrical density of the Loki
ESCRT-III cryo-EM structure.
390

Notably, in the ESCRT homologous archaeal CDV system, CDVA, was reported to bind

DNA. The exact role of DNA in ESCRT-III filaments assembly is yet to be determined

but the accumulating evidence for nucleotide binding in prokaryotic and eukaryotic

ESCRTs supports the notion that oligonucleotides binding is another property of the

ESCRT-III machine that is conserved in all domains of life.

The mechanisms by which ssDNA affects the extent of membrane remodeling are

unclear. The presence of ssDNA may change the biophysical properties of the tube,

thereby resisting the deformation of the tube needed to facilitate membrane remodeling

from within. Alternatively, ssDNA and membranes may compete over the same binding

sites. Recent results from our laboratory showed that removal of a positively charged N

terminal domain, conserved in Asgard CHMP4-7 proteins, from Loki CHMP4-7 abolish

DNA binding (26). Indeed, the N terminal region of CHMP4-7 is facing the inner side of

the tube in the cryo-EM structure (26). A plausible scenario, therefore, is that both DNA

and membrane binding are mediated via the positively charged N terminal patch located

at the interior of the tube. As a result, the presence of ssDNA will reduce the

interactions of the tube with the membrane, resulting in a reduced membrane

remodeling. Consistent with this model, SUV invaginations were barely detected using

neutrally charged SUVs. Future experiments allowing direct visualization of ESCRT-III

proteins, DNA, and the membrane at high resolution are needed to examine these

models and resolve the intimate relationship between these three partners.

Overall, the tubular Loki ESCRT-III filaments characterized here, share more similar

characteristics to bacterial PspA and Vipp1 and human CHMP1B/IST1 filaments than

eukaryotic Vps24 or Snf-7 filaments. Filaments of both Asgard ESCRT-III and human

CHMP1B, were shown to assemble into helical polymers that exhibit a constricted, more

uniform diameter in the presence of an additional ESCRT-III (CHMP1-3 and IST1,

respectively); both filaments were reported to bind DNA in vitro; and both deformed

membranes via binding to the inner side of the filamentous tube. Consistently, we

recently showed that both Loki CHMP4-7 and human CHMP1B associate with

chromatin when expressed in mammalian cells (26). We therefore suggest that Asgard

ESCRT-IIIs are functionally more similar to CHMP1B in the eukaryotic ESCRT-III

system. The findings that the overall structure of the Loki ESCRT-III helical tube

resemble that of the bacterial PspA while the intermolecular interactions between the

ESCRT-III helices are more similar to the ones found in CHMP1B, support the notion

that the Asgard ESCRT system is an intermediate between the prokaryotic and

eukaryotic systems.

The biological function of Asgard ESCRTs has yet to be defined. The findings that Loki

ESCRT-III filaments can bind both DNA and membranes raises a few attractive

possibilities. First, given the established role of ESCRTs in cell division in TACK

archaea and in eukaryotes, it is possible that the ESCRT machinery manifest a basic

cell division machinery capable of orchestrating DNA segregation and membrane

splitting (22-25). In this regard, the ability of DNA to retain extensive membrane

deformation may represent a biological mechanism to avoid premature membrane

constriction during cell division. Second, the ESCRT complex may have been involved

in nuclear membrane formation during eukaryogenesis, by bringing together

membranes and DNA and facilitating DNA encapsulation by membranes. In this

respect, mammalian ESCRT-III proteins have been shown to mediate sealing of the

nuclear membrane post cell division and to regulate the attachment of heterochromatin

to the newly formed nuclear envelope (33-35). Moreover, we recently reported that Loki

CHMP4-7 is targeted to the nucleus when over expressed in eukaryotic cells (26).

Finally, the membrane binding experiments performed here were done using

eukaryotic-like phospholipids, which are fundamentally different from archaeal lipids but

may resemble the lipid composition of the newly formed nuclear membrane. Therefore,

the ESCRT-III system encoded by Asgard archaea could have contributed to the

membrane remodeling that occurred during the formation of the phospholipids-based

nuclear membrane.

MATERIALS AND METHODS

Cloning

Open reading frames (ORFs) from Loki MAG sp. GC14_75, encoding for CHMP1-3 and CHMP4-7 genes were obtained from NCBI (accession numbers KKK42122.1, KKK44605.1, respectively) were amplified by PCR and sub-cloned into pSH21 vector subsequently to the TEV cleavage tag that followed an N terminal polyhistidine and MBP protein tags (a kind gift from Eyal Gur, BGU, Israel) (36). Plasmids were verified by sequencing.

Protein Expression and Purification

pSH21 plasmids containing Loki CHMP1-3 or CHMP4-7 were expressed Escherichia coli BL21 strain and incubated in LB 50 µg/mL Ampicillin at 37°C until reaching O.D. of 0.6. Then, induction with 0.5 mM IPTG was performed and the cultures were grown at 30°C for 3-4 hours. Cells were then resuspended and lysed by sonication in buffer A (25 mM Hepes pH= 8, 500 mM NaCl, 10% glycerol, 15 mM 2-mercaptoethanol, 10 mM imidazole) supplemented with protease inhibitor cocktail (cOmplete ultra-tablets, EDTA-free, Roche) and 500 µg/ml DNase I (10104159001; Roche Diagnosis GmbH, Manheim, Germany), and the supernatant was clarified by centrifugation (25,000 g, 40 mins, 4°C). For CHMP4-7 purification, supernatant was subjected to HisPur™ Ni-NTA Resin (88222; Thermo Fisher Scientific, Weltham, MA) column and eluted in buffer B (buffer A supplemented with 50 mM imidazole). Protein tags were then removed using a TEV protease (1 mg/10 mg protein) in buffer C (50mM Tris-HCL pH=7.4, 10% glycerol, 500 mM NaCl, 15 mM 2-mercaptoethanol). Finally, a second Ni-NTA step was performed using buffer C and the flow-through was collected.

For CHMP1-3 purification, supernatant was subjected to Ni-NTA column and eluted in buffer B. Samples were then subjected to amylose resin (E8021; NEB, Frankfurt, Germany) in buffer D (20mM Tris pH=7.4, 500 mM NaCl, 10mM β-Mercaptoethanol, 10% Glycerol) and eluted in buffer D supplemented with 10mM maltose. Protein tags were then removed using a TEV protease (1 mg/10 mg protein) in buffer C (50mM Tris-HCL pH=7.4, 10% glycerol, 500 mM NaCl, 15 mM 2-mercaptoethanol). Finally, a

second Ni-NTA step was performed using buffer C and the flow-through was collected.

Purified proteins were verified my mass spectrometry.

For polymerization, proteins were incubated in buffer E (25mM Tris pH=7.6, 50 mM KCl)

for 5 hours at RT at the indicated conditions. For DNA experiments, buffer E was

supplemented with ATCCACCTGTACATCAACTCGCCCGGCGGCTCGATCAGCG (40

bases probe, 10 µM), ss or ds at the indicted concentrations. DNase I (10104159001;

Roche Diagnosis GmbH, Manheim, Germany) was added at 20 mg/ml concentrations

when indicated. Homo polymerization of CHMP1-3 was examined at a range of pH (5.0-

8.8), different salt concentrations (0-500 mM NaCl). Polymerization could not be

observed under any of these conditions (Fig. S1).

Preparation of small unilamellar vesicles (SUVs)

Chloroform Lipid solutions (DOPC or DOPS) were purchased from Avanti Polar Lipids

Inc. (Albaster, AL, USA catalog number 850375, 840035). Lipid mixtures of

DOPC/DOPS 1:1 molar ratio and pure DOPC were prepared at a total concentration of

1 mg/ml in Tris 20 mM, KCl 50 mM, pH 7.8 were prepared. Each lipid suspension was

extruded 20 times via a 100 nm polycarbonate membrane using a mini-extruder (Avanti

Polar Lipids, Alabaster, AL, USA). In the final stage, the vesicles were mixed with the

Asgard Loki ESCRT-III filaments. All experiments were done with 8 µM ESCRT-III and

124 µM vesicles in a buffer of Tris 25 mM, KCl 50 mM, at pH 7.6. DNase was added to

the indicated conditions at 20mg/ml concentrations for one hour post vesicle-ESCRT-III

incubation.

Negative stain grid preparation and imaging

The negative stain samples for TEM analysis were prepared in the following way: 300

mesh copper grids (Ted Pella, Prod No. 01813-F) were glow discharged to enhance

hydrophilicity of their surface. Next, 2.5 µL of the sample was applied on to the grid and

the excess liquid was blotted with filter paper after 1 minute. The grid was dried in air for

1 minute, following by applying 5 µL of uranyl acetate 2% solution (SPI CAS# 6159-44-

0) for negative staining to increase the sample contrast. Next, the grid was blotted once

more to remove the excess uranyl acetate. Finally, the grid was dried in air before

insertion into the microscope. Imaging of the samples was performed with Thermo

Fisher Scientific Talos F200C transmission electron microscope operating at 200 kV.

The images were taken with Ceta 16M CMOS camera.

Cryo-electron microscopy

Sample preparation: 3 µL of proteins / proteins-phospholipids mix samples were gently

deposited on glow discharged Quantifoil R 1.2/1.3 holey carbon grids (Quantifoil Micro

Tools GmbH, Germany). Samples were manually blotted for four seconds at room

temperature and vitrified by rapidly plunging into liquid ethane using a home-built

plunging apparatus. The frozen samples were stored in liquid nitrogen until imaging.

Micrograph acquisition. For an initial data set, samples were loaded under cryogenic

conditions and imaged in low dose mode on either a FEI Tecnai F30 Polara microscope

(FEI, Eindhoven) operated at 300 kV or a Glacios (Thermo-Fisher Scientific) operated at

200 kV. On the Polara, micrographs were loaded under cryogenic conditions and

imaged in low dose mode on a FEI Tecnai F30 Polara microscope (FEI, Eindhoven)

operated at 300 kV. Micrographs were collected using SerialEM, at a calibrated pixel

size of 1.1 Å by a K2 Summit direct electron detector fitted behind an energy filter

(Gatan Quantum GIF) set to ±10 eV around zero-loss peak with total electron dose of

80 ē/Å2. Each dose-fractionated movie had 50 frames, micrographs sums were aligned

in SerialEM. On the Glacios, micrographs were collected using Tomo (Single

micrograph) at a calibrated pixel size of 0.87 Å by a Falcon 4i direct electron detector

fitted behind a Selectris X energy filter set to ±5 eV around zero-loss peak with total

electron dose of 30 e$^-$/Å2. The final dataset used for was collected on 300 kV Titan

Krios G4 (ThermoFisher Scientific) electron microscope equipped with a Biocontinuum

K3 (Gatan) detector operated by EPU (ThermoFisher Scientific) (see details Table 1).

Dose-fractionated movie frames (30 frames per movie) were acquired at a nominal

defocus of 1.5 to 3 µm with a calibrated pixel size of 1.36 Å and a total dose of 30 e$^-$/Å^2.

Helical reconstruction. For the initial dataset, movie frames were gain-corrected,

dose-weighted, and aligned using cryoSPARC Live (37). Initial 2D classes were

produced using the auto picker implemented in cryoSPARC Live. The following image

processing steps were performed using cryoSPARC. The classes with most visible

details were used as templates for the filament tracer (50 Å spacing, 50 Å low-pass

filter) The resulting filament segments were extracted with 1000 px box size (Fourier

cropped to 300 px) and subjected to multiple rounds of 2D classification. The remaining

segments were sorted by rod diameter and 2D class averages of the most abundant

diameter for initial symmetry guesses in PyHI (38). Initial symmetry estimates were

validated by helical refinement in cryoSPARC and selection of the helical symmetry

parameter yielding a reconstruction with typical ESCRT-III features and the best

resolution. For the main dataset, movie frames were gain-corrected, dose-weighted,

and aligned using cryoSPARC Live (37). Initial 2D classes were produced using the

auto picker implemented in cryoSPARC Live. The following image processing steps

were performed using cryoSPARC. The classes were separated by filament type (PspA-

like rods and curved filaments). Classes with most visible detail were used as templates

for the template picker (50 Å spacing). The resulting curved filament segments were

extracted with 400 px box size (Fourier cropped to 200 px) and subjected to multiple

rounds of 2D classification. The remaining segments were sorted by curvature. Only the

least curved segments were kept. No regular helical symmetry could be identified for

these filaments, and initial attempts with *ab initio* reconstruction did not yield consistent

maps. The PspA-like rods were extracted with 800 px box size (Fourier cropped to 200

px) and subjected to multiple rounds of 2D classification. The remaining segments were

sorted by diameter. The previously determined helical symmetry was used for an initial

reconstruction of the most abundant rods (after re-extraction with 600 px box size,

Fourier cropped to 300 px). The corresponding segments were classified by

heterogeneous refinement and followed by 3D classifications using the initial helical

reconstructions as templates. The resulting helical reconstructions were subjected to

multiple rounds of helical refinement including the symmetry search option. For the final

polishing, the segments were re-extracted at 600 px without Fourier cropping and

subjected to helical refinement including non-uniform refinement. Higher-order

aberrations were corrected using global and local CTF-refinement. The local resolution

distribution and local filtering for the resulting map was performed using cryoSPARC (as

shown in Fig 4E). The resolution of the final reconstruction was determined by Fourier

shell correlation (auto-masked, FSC=0.143).

Cryo-EM map interpretation and model building. The handedness of the maps was

determined by rigid-body fitting the hairpin of an Alphafold2 (*39, 40*) prediction of Loki

CHMP4-7 into the final map using ChimeraX (*41, 42*) and flipped accordingly.

Predictions of Loki CHMP 1-3 and CHMP 4-7 monomers were flexibly MDFF fitted to

the 3D reconstruction using ISOLDE (*43*). Only the CHMP4-7 model fitted the

secondary structure and large residue pattern present in the map. The CHMP1-3 model

was discarded. The CHMP4-7 monomer model was truncated to the parts covered by

the map (aa 15-172) and manually refined in ISOLDE. Subsequently, the model was

subjected to auto-refinement with *phenix.real_space_refine* (*44*). The auto-refined

model was checked/adjusted manually in Coot (*45*). The respective helical symmetry

was applied to the model to create assemblies of 60 monomers in ChimeraX and the

assembly structure was subjected to another cycle of auto-refinement (with NCS

restraints and refinement) using *phenix.real_space_refine* (*44*). After the final

inspection, the models were validated in *phenix.validation_cryoem* (*46*) */Molprobity* (*47*).

Cryo-tomography. Micrographs were acquired using the same setups as of the cryo-

EM micrographs. Pixel size at the sample plane on the Polara was 2.3 Å, and 1.9 Å on

the Glacios. On the Polara, the camera was operated in counting mode at a dose rate of

6−7 ē/pixels/sec, tilt series of 41 exposures from −21° to +60° and back to -60° in 3°

intervals and a total dose of ∼ 120 ē /Å^2 were collected with Serial-EM. On the Glacios,

the camera was operated in counting mode at a dose rate of 2−3 ē/pixels/sec. Fiducials-

free, dose-symmetric tilt series of 61 exposures from −60° to +60° in 2° intervals and a

total dose of ∼120 ē /Å^2 were collected with Tomo. Tilt series were aligned, and

tomograms were reconstructed using eTomo (IMOD4.11 package) (*48*). Movies were

generated in imageJ.

Measurements and statistical analysis

Filament length and diameter measurements were performed in Velox software version

3.5 (Thermo Fisher Scientific, Inc.).

Statistical analysis was performed in Graph Pad Prism version 9.00 (La Jolla, CA,
USA). Normality was tested using the D'Agostino-Pearson test. For three or more
groups, statistical significance was determined by one-way analysis of variance
(ANOVA) Kruskal-Wallis test followed by Dunn's post hoc test or by one-way analysis of
variance (ANOVA) followed by Tukey post hoc test, for multiple comparisons. Pearson
chi-squared test was used to determine the difference in the distribution of the
assembled polymers. P values in all figures are as follows: * p- value ≤ 0.1, **p- value
≤ 0.01, ***p- value ≤ 0.001, ****p- value ≤ 0.0001. Bars in all plots indicate mean values
and standard deviation of the mean.
615

616 ACKNOWLEDMENTS
617
We thank the laboratories of Eyal Gur (BGU) and Eyal Arbely (BGU) for their practical
advice on proteins expression and purifications and for kindly sharing their reagents and
equipment with us. We also thank Itzik Mizrahi (BGU) and Ori Avinoam (WIS) for
constructive feedback on the project. Last, we thank all members of the Elia lab for
critical feedback throughout the project. The Elia laboratory is funded by the Israeli
Science Foundation (ISF) Grant no. 1323/18. ABG is grateful to the Israel Science
Foundation for financial support (grant 2101/20). The authors gratefully acknowledge
the electron microscopy training, imaging, and access time granted by the life science
EM facility of the Ernst-Ruska Centre at Forschungszentrum Jülich. The authors
gratefully acknowledge the computing time granted by the JARA Vergabegremium and
provided on the JARA Partition part of the supercomputer JURECA at
Forschungszentrum Jülich (49).

630
631 AUTHOR CONTRIBUTIONS
632
NE and DN conceptualized the project. NM performed and analyzed all experiments.
GH and AB, were in charge of experiments using SUVs. AU acquired negative stain
TEM images. RZ, BJ and CS acquired and analyzed cryo-EM data. BJ and CS built and
refined the atomic models. NE, DN, AB and RZ were involved in experimental design

and data analysis. NE, BJ and CS wrote the manuscript. All authors read and revised

the manuscript.

Competing Interest Statement: We declare no competing interests.

REFERENCES

1. K. Zaremba-Niedzwiedzka *et al.*, Asgard archaea illuminate the origin of eukaryotic cellular complexity. *Nature* **541**, 353-358 (2017).
2. F. MacLeod, G. S. Kindler, H. L. Wong, R. Chen, B. P. Burns, Asgard archaea: Diversity, function, and evolutionary implications in a range of microbiomes. *AIMS Microbiol* **5**, 48-61 (2019).
3. A. Spang *et al.*, Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature* **521**, 173-179 (2015).
4. E. Neveu, D. Khalifeh, N. Salamin, D. Fasshauer, Prototypic SNARE Proteins Are Encoded in the Genomes of Heimdallarchaeota, Potentially Bridging the Gap between the Prokaryotes and Eukaryotes. *Curr Biol* **30**, 2468-2480 e2465 (2020).
5. Z. Lu *et al.*, Coevolution of Eukaryote-like Vps4 and ESCRT-III Subunits in the Asgard Archaea. *mBio* **11**, (2020).
6. T. Hatano *et al.*, Asgard archaea shed light on the evolutionary origins of the eukaryotic ubiquitin-ESCRT machinery. *Nat Commun* **13**, 3398 (2022).
7. Y. Caspi, C. Dekker, Dividing the Archaeal Way: The Ancient Cdv Cell-Division Machinery. *Front Microbiol* **9**, 174 (2018).
8. J. H. Hurley, ESCRTs are everywhere. *EMBO J* **34**, 2398-2407 (2015).
9. Y. A. M. Alonso, S. M. Migliano, D. Teis, ESCRT-III and Vps4: a dynamic multipurpose tool for membrane budding and scission. *FEBS J* **283**, 3288-3302 (2016).
10. J. McCullough, L. A. Colf, W. I. Sundquist, Membrane fission reactions of the mammalian ESCRT pathway. *Annu Rev Biochem* **82**, 663-692 (2013).
11. C. Caillat, S. Maity, N. Miguet, W. H. Roos, W. Weissenhorn, The role of VPS4 in ESCRT-III polymer remodeling. *Biochem Soc Trans* **47**, 441-448 (2019).
12. S. Maity *et al.*, VPS4 triggers constriction and cleavage of ESCRT-III helical filaments. *Sci Adv* **5**, eaau7198 (2019).
13. B. E. Mierzwa *et al.*, Dynamic subunit turnover in ESCRT-III assemblies is regulated by Vps4 to mediate membrane remodelling during cytokinesis. *Nat Cell Biol* **19**, 787-798 (2017).
14. A. K. Pfitzner *et al.*, An ESCRT-III Polymerization Sequence Drives Membrane Deformation and Fission. *Cell* **182**, 1140-1155 e1118 (2020).
15. W. M. Henne, N. J. Buchkovich, Y. Zhao, S. D. Emr, The endosomal sorting complex ESCRT-II mediates the assembly and architecture of ESCRT-III helices. *Cell* **151**, 356-371 (2012).
16. I. Goliand *et al.*, Resolving ESCRT-III Spirals at the Intercellular Bridge of Dividing Cells Using 3D STORM. *Cell Rep* **24**, 1756-1764 (2018).
17. J. Guizetti *et al.*, Cortical constriction during abscission involves helices of ESCRT-III-dependent filaments. *Science* **331**, 1616-1620 (2011).
18. J. McCullough *et al.*, Structure and membrane remodeling activity of ESCRT-III helical polymers. *Science* **350**, 1548-1551 (2015).
19. S. Lata *et al.*, Helical structures of ESCRT-III are disassembled by VPS4. *Science* **321**, 1354-1357 (2008).

20. B. P. Frohn, T. Hartel, J. Cox, P. Schwille, Tracing back variations in archaeal ESCRT-based
cell division to protein domain architectures. *PLoS One* **17**, e0266395 (2022).
21. J. Liu *et al.*, Bacterial Vipp1 and PspA are members of the ancient ESCRT-III membrane-
remodeling superfamily. *Cell* **184**, 3660-3673 e3618 (2021).
22. A. C. Lindas, E. A. Karlsson, M. T. Lindgren, T. J. Ettema, R. Bernander, A unique cell
division machinery in the Archaea. *Proc Natl Acad Sci U S A* **105**, 18942-18946 (2008).
23. R. Y. Samson, T. Obita, S. M. Freund, R. L. Williams, S. D. Bell, A role for the ESCRT
system in cell division in archaea. *Science* **322**, 1710-1713 (2008).
24. R. Y. Samson *et al.*, Molecular and structural basis of ESCRT-III recruitment to
membranes during archaeal cell division. *Mol Cell* **41**, 186-196 (2011).
25. G. Tarrason Risa *et al.*, The proteasome controls ESCRT-III-mediated cell division in an
archaeon. *Science* **369**, (2020).
26. D. Nachmias *et al.*, Asgard ESCRT-III and VPS4 reveal conserved chromatin binding
properties of the ESCRT machinery. *ISME J*, (2022).
27. K. Azad *et al.*, in *bioRxiv*. (2022).
28. Y. Olmos, The ESCRT Machinery: Remodeling, Repairing, and Sealing Membranes.
Membranes (Basel) **12**, (2022).
29. B. Junglas *et al.*, PspA adopts an ESCRT-III-like fold and remodels bacterial membranes.
Cell **184**, 3674-3688 e3618 (2021).
30. H. C. Nguyen *et al.*, Membrane constriction and thinning by sequential ESCRT-III
polymerization. *Nat Struct Mol Biol* **27**, 392-399 (2020).
31. N. Talledge *et al.*, in *bioRxiv*. (2018).
32. T. K. Gupta *et al.*, Structural basis for VIPP1 oligomerization and maintenance of
thylakoid membrane integrity. *Cell* **184**, 3643-3659 e3623 (2021).
33. G. H. Pieper, S. Sprenger, D. Teis, S. Oliferenko, ESCRT-III/Vps4 Controls
Heterochromatin-Nuclear Envelope Attachments. *Dev Cell* **53**, 27-41 e26 (2020).
34. Y. Olmos, L. Hodgson, J. Mantell, P. Verkade, J. G. Carlton, ESCRT-III controls nuclear
envelope reformation. *Nature* **522**, 236-239 (2015).
35. M. Vietri *et al.*, Spastin and ESCRT-III coordinate mitotic spindle disassembly and nuclear
envelope sealing. *Nature* **522**, 231-235 (2015).
36. Y. Elharar *et al.*, Survival of mycobacteria depends on proteasome-mediated amino acid
recycling under nutrient limitation. *EMBO J* **33**, 1802-1814 (2014).
37. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).
38. X. Zhang, Python-based Helix Indexer: A graphical user interface program for finding
symmetry of helical assembly through Fourier-Bessel indexing of electron microscopic
data. *Protein Sci* **31**, 107-117 (2022).
39. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583-589 (2021).
40. M. Mirdita *et al.*, ColabFold: making protein folding accessible to all. *Nat Methods* **19**,
679-682 (2022).
41. T. D. Goddard *et al.*, UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. *Protein Sci* **27**, 14-25 (2018).

42.   E. F. Pettersen *et al.*, UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. *Protein Sci* **30**, 70-82 (2021).

43.   T. I. Croll, ISOLDE: a physically realistic environment for model building into low-
resolution electron-density maps. *Acta Crystallogr D Struct Biol* **74**, 519-530 (2018).

44.   P. V. Afonine *et al.*, Real-space refinement in PHENIX for cryo-EM and crystallography.
Acta Crystallogr D Struct Biol **74**, 531-544 (2018).

45.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta*
Crystallogr D Biol Crystallogr **66**, 486-501 (2010).

46.   P. V. Afonine *et al.*, New tools for the analysis and validation of cryo-EM maps and
atomic models. *Acta Crystallogr D Struct Biol* **74**, 814-840 (2018).

47.   C. J. Williams *et al.*, MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Sci* **27**, 293-315 (2018).

48.   D. N. Mastronarde, Automated electron microscope tomography using robust prediction
of specimen movements. *J Struct Biol* **152**, 36-51 (2005).

49.   T. P., JURECA: Data Centric and Booster Modules implementing the Modular
Supercomputing Architecture at Jülich Supercomputing Centre. *Journal of large-scale*
reseach facilities JLSRF **7**, 182-184 (2021).

50.   A. M. Waterhouse, J. B. Procter, D. M. Martin, M. Clamp, G. J. Barton, Jalview Version 2--
a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189-
1191 (2009).

51.   P. Di Tommaso *et al.*, T-Coffee: a web server for the multiple sequence alignment of
protein and RNA sequences using structural information and homology extension.
Nucleic Acids Res **39**, W13-17 (2011).

792 **Table 1. Data collection, image processing, and model refinement.**

Movies	12,751
Magnification	63 kx
Voltage (kV)	300
Total dose (e⁻/ Å²)	30
Defocus range (μm)	1.5 to 3
Physical pixel size (Å)	1.36
Detector	Gatan K3
Symmetry imposed	1.05 Å rise 173.9 ° rotation C1
Final no. of segments (ASUs)	228,688 (11,205,712)
Global map resolution (Å, FSC = 0.143)	3.55
Local map resolution range (Å, FSC = 0.5)	3.0 -6.0
Initial model used (PDB code)	AF2 prediction
Model refinement	
Model resolution	3.5
CC mask	0.76
CC box	0.29
CC peaks	-0.19
CC volume	0.78
CC ligands	-
Map sharpening B-factor (Å²)	-76
Model composition	
Nonhydrogen atoms	1251
Protein	1251
RMSDs	
Bond lengths (Å)	0.007
Bond angles (°)	0.909
Validation	
MolProbity score	1.34
Clashscore	3.57
Rotamer outliers (%)	0.74
Ramachandran plot	
Favored (%)	96.79
Allowed (%)	3.21
Disallowed (%)	0.0
Deposition IDs	
EMDB	50583
PDB	9FN1

Fig 1. Loki CHMP4-7 self-assembles into helical tubes.

A. Schematic representation of the currently suggested models for the evolutionary relationships between Bacteria, Archaea, and Eukarya that integrate the archaeal Asgard clade. **B**. Negative-stain TEM micrographs of Loki CHMP4-7 homo-polymers. Representative zoomed-out (left, scale = 200 nm) and zoomed-in (middle and right, scale = 100 nm) images are shown. Orange arrows indicate proto-filaments, yellow arrow indicates a helical tube. **C.** Representative Cryo-EM micrographs of Loki CHMP4-7 homo-polymers. Left, scale = 100 nm, middle and right, scale = 50 nm. An enlargement of the area depicted in black box in middle panel is show in right panel. **D.** A representative 2D class average of 502 particles extracted from CHMP4-7 projections. Scale = 100 Å. **E.** Frequency histograms of Loki CHMP4-7 helical diameters (left, n = 37) and length (right, n = 36) obtained from negative stain TEM images acquired from two independent experiments. Averaged CHMP4-7 helical tube diameter 44 ± 30.9 nm.

A CHMP 4-7

CHMP 4-7 - CHMP 1-3

B

Fig 2. CHMP1-3 induces morphological changes in Loki ESCRT-III helical tubes

A. Representative negative-stain TEM images of Loki CHMP4-7 homo-polymers (upper panels) and CHMP4-7-CHMP1-3 co-polymers (bottom panels, 4:1 molar ratio). Zoomed-out (left, scale = 1 µm) and zoomed-in (right, scale = 50 nm) images are shown. Yellow arrows indicate helical tubes. Note, that the helical tube in co-polymers is more curved and displays periodic stripes that are not seen in the homo-polymers by negative staining **B**. Diameter (left) and length (right) measurements of CHMP4-7-CHMP1-3 helical-tubes that were assembled at the indicated molar ratios. Measurements were obtained from negative stain TEM images. Averaged diameters (nm): 2:1 ratio, 31 ± 7.7, n=25; 4:1 ratio 28 ± 1.6, n=26, 6:1 ratio 32 ± 3.8, n=23. One-way analysis of variance (ANOVA) Kruskal-Wallis test followed by Dunn's post hoc test was performed (see methods). Averaged lengths (nm): 2:1 ratio 641 ± 173, n=25; 4:1 ratio 594 ± 158, n=27; 6:1 ratio 492 ± 109, n=25

Fig 3. DNA facilitates Loki ESCRT-III helical-tube assembly.

A-D Loki ESCRT-III proteins CHMP4-7 and CHMP1-3 (4:1 molar ratio) (**A**) or CHMP4-7 alone (**B**) were subjected to plomerization reaction (see methods) in the absence or presence of 40 bases oligonucleotides (ssDNA or dsDNA, as indicated), and documented using negative-stain TEM. Representative zoomed-out (left, Scale = 1 μm) and zoomed-in (right, Scale = 50 nm) images are shown in **A-B** and in Fig. S3A-B. Helical tubes are colored in yellow in zoomed-out images (raw images, Fig. S3A). **C.** Helical tubes density at various conditions. Measurements were performed on an area >565 μm^2 from each experiment. Data from two independent experiments are presented for each condition. **D.** Diameters of CHMP4-7 and CHMP4-7/CHMP1-3 helical tubes assembled in the presence of oligonucleotides. Averaged diameters (nm): CHMP4-7/CHMP1-3 29.4 ±4.6, n=27; CHMP4-7/CHMP1-3 ssDNA 40±4.1, n=23; CHMP4-7/CHMP1-3 dsDNA 48±11.5, n=37; CHMP4-7 43±5, n=23; CHMP4-7 ssDNA 44 ± 5.7, n = 22. **E**. Loki ESCRT-III proteins CHMP4-7 and CHMP1-3 (4:1 molar ratio) (left) or CHMP4-7 alone (right) were assembled under control conditions, or in the presence of 20 mg/ml DNase, as indicated. Helical tubes are colored in yellow (raw images, Fig. S3E). Scale = 1 μm. Data was reproduced in four independent experiments. Statistics in C and D was performed using one-way analysis of variance (ANOVA) (see methods).

Fig 4. Loki CHMP4-7 forms PspA-like rod structures.

A. Example cryo-EM micrograph of Loki CHMP4-7 showing straight rod structures with

different diameters. **B.** Set of 2D class averages of Loki CHMP4-7 rods with the typical

Christmas tree pattern of PspA rods. **C.** Side (left) and top (right) view of cryo-EM

density map of the rod structure with an inner and diameter of 26 and 36 nm,

respectively. **D.** Fourier shell correlation of the cryo-EM density map indicating a

nominal resolution at 3.6 Å according to the FSC 0.143 cutoff. **E.** Estimates of local

resolution mapped in color code (blue=3.0 and red 6.0 Å resolution, respectively) on the

cryo-EM density.

Fig 5. Loki CHMP4-7 rods are built from monomers sharing typical ESCRT-III

features.

A. Topology plot of the Loki CHMP4-7 structure with color-coded helices: α0 white, α1

red, α2+3 violet, α4 blue, α5 cyan, α6 white. **B.** Segmented density including modeled

atomic Loki CHMP4-7 monomer structure (inset visible side-chains aa 104-134) found

in 36 nm rods showing the ESCRT-III fold. **C.** Organization of the CHMP 4-7 monomers

in the helical assembly: α5 of chain j interacts with the hairpin of chain j+4. **D.** Cryo-EM

map of Loki CHMP4-7 rod with isosurface colored CHMP 4-7 α-helices and enlarged

view of the cryo-EM map with the fitted model showing the interactions in the tube wall

and helix α5 interactions in the periphery. **E.** Comparison of tubular ESCRT-III

assemblies from bacteria, Asgard archaea and eukaryotes. PspA: 7abk, Vipp1: 7o3w,

IST1: 6tza, CHMP1B: 6tz9, CHMP2A-CHMP3: 7zch. Color-coded helices: α0 green, α1

red, α2+3 violet, α4 blue, α5 cyan.

A CHMP 4-7 - CHMP 1-3 ssDNA + SUVs

B CHMP 4-7 - CHMP 1-3 + SUVs
ssDNA   No DNA

C
Outside   Docking
Inside   Remodeling

D
slice 111
slice 136

E

Fig 6. Loki ESCRT-III filaments deform eukaryotic-like membrane vesicles

A-B. Representative negative-stain TEM (**A**) and cryo-EM (**B**) images of CHMP4-7-CHMP1-3 helical tubes (molar ratio, 4:1) assembled in the presence of ssDNA (10 µM), and incubated with SUVs (DOPC:DOPS, 1:1 ratio). Scale = 100 nm. Arrows in **B** specify internalized vesicles. **C**. Percentages of the different types of vesicle-tube interactions observed under the specified conditions. Left panel, representative negative-stain images of the four types of interactions quantified. Outside (type 1); Docking (type 2); Inside (type 3); Remodeling (type 4). Data was quantified from negative stain images obtained from at least two independent experiments for each condition (n = CHMP4-7 ssDNA, 124; CHMP4-7-CHMP1-3 ssDNA, 159; CHMP4-7, 197; CHMP4-7-CHMP1-3, 161). Scale = 100 nm. Pearson chi-squared analysis show a significant difference in the distributions of homo- and co-polymers assembled in the presence vs absence of DNA (p- value ≤ 0.0001 for all pairs). No significance was found in the distributions obtained for ESCRT-III filaments assembled in the presence or absence of CHMP 1-3 that were subjected to similar treatments. Scale =1 µm. **D**. Selected slices of reconstructed tomograms of filaments-SUV complexes showing the membrane (yellow arrows) and ESCRT-III filament (white arrows). The complete series is provided in Sup. movies 1. **E.** A suggested model for Loki ESCRT-III mediated vesicle remodeling. Vesicles docked at the tip of the ESCRT-III tube directly interact with the interior of the tube, leading to deformation and invagination of the vesicle by the ESCRT-III filament. This process occurs repeatedly, leading to the formation of a complex membrane network inside the helical tube, which ultimately causes deformation and remodeling of the ESCRT-III tube. Tube remodeling is inhibited in the presence of DNA, resulting in the accumulation of ESCRT-III-SUV interactions in all the stages that precede this stage.

Fig S1: Loki CHMP1-3 do not self-assemble into high-ordered structures. (**A-C**)

representative negative stain TEM images of full-length Loki CHMP1-3 that was

subjected to polymerization reaction at different buffers. (**A**) buffer used for CHMP4-7

self-assembly: 25 mM Tris pH=7.6, 50 mM KCl). (**B**) Using acidic (50 mM Sodium

Acetate pH=5, 50 mM KCl) or basic (25 mM Tris pH=8.8 50 mM KCl buffers). (**C**) at

different ionic strengths (high salt: 25 mM Tris pH=7.6, 200 mM KCl; No salt: 25 mM

Tris pH=7.6. A final concentration of 10.88 µM CHMP1-3 was used for all buffer

conditions. No filaments were observed under any of the examined conditions.

A CHMP 4-7 - CHMP 1-3 (2:1)

CHMP 4-7 - CHMP 1-3 (6:1)

B CHMP 4-7 - CHMP 1-3 ssDNA

C CHMP 4 -7   CHMP 4-7 - GFP

Fig. S2: Characterization of Loki ESCRT-III filaments. A. Representative negative

stain TEM images of Loki ESCRT-III filaments assembled using CHMP4-7 and CHMP1-

3 at molar ratios 2:1 (upper panel) and 6:1 (bottom panel). Zoomed-out (left, Scale = 1

µm); zoomed-in (right, Scale = 50 nm). Yellow arrows indicate helical tubes.

Corresponds to measurements in Fig. 2B. **B.** Negative stain TEM images of loose Loki

Loki ESCRT-III filaments assembled in the presence of CHMP4-7 – CHMP1-3 (2:1

molar ratio, ssDNA) resembling a telephone cord, supporting that the ESCRT-III tube is

composed of a packed helical filament. Scale = 200 nm **C.** Representative negative

stain TEM images of filaments assembled in the presence of Loki CHMP 4-7 at a total

concentration that equals the concentration of both CHMP4-7 - CHMP1-3 at 2:1 ratio

(left, final concentration of 16 µM) and in the presence of both CHMP4-7 and purified

recombinant GFP (right panel), reaching the same final concentrations (CHMP 4-7

10.67 µM, GFP 5.4 µM). Scale = 200 nm. No effect on filament formation was identified

under these assembly conditions. Data in all panels was reproduced in at least two

independent experiments.

Fig. S3: Helical tube formation by Loki ESCRT-III in the presence of nucleic acids.

A. Raw data of the representative zoomed-out negative stain TEM images presented in

Fig. 3A. Scale = 1 μm. **B.** CHMP4-7 and CHMP1-3 (4:1 molar ratio) were subjected to

polymerization reaction (see methods) in the presence of 40 bases dsDNA, and

documented using negative-stain TEM. Scale zoomed out, 1 μm; zoomed in, 50 nm **C.**

Representative zoomed-out negative stain TEM images of assembly reactions using

Loki CHMP 1–3 with or without ssDNA (40 bases probe) showing that addition of

ssDNA does not induce polymerization of CHMP 1-3. Scale = 200 nm. **D.** Length

measurements of helical tubes formed under the indicated conditions (corresponds to

Fig. 3). Measurements obtained from negative stain TEM images of samples produced

in at least two independent experiments for each condition. Averaged lengths (nm):

CHMP4-7/CHMP1-3 534.8 ±118.8, n=25; CHMP4-7/CHMP1-3 ssDNA 777.5 ±210.6,

n=30; CHMP4-7/CHMP1-3 dsDNA 599±218.2, n=45; CHMP4-7 364±67.8, n=23;

CHMP4-7 ssDNA 363 ± 96.1, n = 21. **E.** Raw data of the representative zoomed-out

negative stain TEM images presented in Fig. 3E.

 Fig. S4: Curved Filaments and ESCRT-III comparison.

 A. Cryo-EM micrograph of CHMP 4-7/CHMP 1-3 (4:1 molar ratio, assembled in the

 presence of ssDNA) showing curved filaments and PspA-like rods. **B.** 2D class

 averages of curved filaments. **C.** z and xy slice (averaged over 10 px) of the Loki

 CHMP4-7 rods showing tube-like fuzzy density in the lumen of the rods. **D.** Top view of

 the density map and the fitted polymer model. **E.** Electrostatic surface of helical

assembly reveals a negatively and a positively charged end indicating a distinct polarity

of the rod. **F.** Sequence alignment of ESCRT-III proteins with known structures

(truncated to α5) in comparison with Loki CHMP 1-3 and 4-7. The alignment is

superimposed with the consensus secondary structures derived from the deposited

models. Residues with similarity >30% are colored according to their properties

(ClustalX color scheme). Sequences were aligned using the T-Coffee webserver and

visualized with JalView *(50, 51)*.

A CHMP 4-7 - CHMP 1-3 + DOPC : DOPS

CHMP 4-7 ssDNA + DOPC : DOPS

CHMP 4-7 + DOPC : DOPS

B DOPC : DOPS

C

Fig. S5: Interactions of Loki ESCRT-III filaments with negatively charged SUVs (DOPC:DOPS 1:1). A. Representative negative stain TEM images of Loki ESCRT-III filaments that were assembled at the indicated conditions and incubated for 1hour with DOPC:DOPS SUVs. Zoom out images are shown on left panels. Arrows indicate interactions between SUVs and filaments. Corresponds to quantification shown in Fig. 6C. Scale left panels = 500 nm, middle and right panels =200 nm. **B.** A representative negative stain TEM image of DOPC:DOPS SUVs prior to incubation with Loki ESCRT-III. (Scale = 500 nm). **C.** Effect of DNase on pre-assembled ESCRT-III-SUV complexes. ESCRT-III-SUV complexes were generated as in A and were subjected (or not) to DNase treatment and the distribution of the different vesicle-tube interactions observed under the specified conditions was quantified. Note that DNase was not able to disrupt pre-assembled ESCRT-III-SUV interactions (categories are as in Fig. 6C).

A CHMP 4-7 -CHMP 1-3 ssDNA + DOPC

CHMP 4-7 -CHMP 1-3 + DOPC

CHMP 4-7 + DOPC

B DOPC

C

Fig. S6: Interactions of Loki ESCRT-III filaments with DOPC SUVs. A.

Representative negative stain TEM images of Loki ESCRT-III filaments that were

assembled at the indicated conditions and incubated for 1hour with DOPC SUVs. Zoom

out images are shown on left panels. Arrows indicate interactions between SUVs and

filaments. Scale: left panels = 500 nm, middle and right panels =200 nm. Note that the

most prominent interaction is adhesion of the vesicle to the outside of the tube. **B.**

representative negative stain TEM image of DOPC SUVs prior to incubation with Loki

ESCRT-III. (Scale = 500 nm). **C.** Percentages of interactions between Loki CHMP 4-7 –

CHMP 1-3 (4:1) filaments with SUVs comprised of DOPC or DOPC-DOPS (1:1),

showing preference to DOPS containing vesicles.

48