Characterization of TelE, an LXG effector of *Streptococcus*

***gallolyticus*, antagonized by a non-canonical immunity protein**

Wooi Keong Teh[1], Yichen Ding[1,2], Francesca Gubellini[3], Alain Filloux[1,4], Claire

Poyart[5], Shaynoor Dramsi[6,7,*], Michael Givskov[1,8,*]

[1] Singapore Centre for Environmental Life Sciences Engineering, Nanyang

Technological University, Singapore

[2] Present address: Temasek Life Sciences Laboratory, Singapore

[3] Institut Pasteur, Unité de Microbiologie Structurale, Paris, France

[4] MRC Centre for Molecular Bacteriology and Infection, Department of Life Sciences,

Imperial College London, London SW7 2AZ, UK

[5] Université de Paris, Assistance Publique Hôpitaux de Paris, Service de

Bactériologie, Centre National de Référence des Streptocoques, Groupe

Hospitalier Paris Centre site Cochin, Paris, France

[6] Institut Pasteur, Université Paris Cité, CNRS UMR6047, Biology of Gram-positive

Pathogens Unit, Paris 75015 France.

[7] Centre National de la Recherche Scientifique (CNRS) UMR2001, Paris, France

[8] Costerton Biofilm Centre, Department of Immunology and Microbiology, University

of Copenhagen, Denmark

* Correspondence:

Shaynoor Dramsi, shaynoor.dramsi@pasteur.fr;

Michael Givskov, mgivskov@sund.ku.dk

Abstract

Streptococcus gallolyticus subsp. *gallolyticus* (*Sgg*) is an opportunistic bacterial pathogen strongly associated with colorectal cancer. Here, through comparative genomics analysis, we demonstrated that the genetic locus encoding the Type VIIb Secretion System (T7SSb) machinery is uniquely present in *Sgg* in two different arrangements. *Sgg* UCN34 carrying the most prevalent T7SSb genetic arrangement was chosen as the reference strain. To identify the effectors secreted by T7SSb, we inactivated the *essC* gene encoding the motor of this machinery. Comparison of the proteins secreted by Sgg UCN34 WT and its isogenic Δ*essC* mutant revealed six T7SSb effector proteins, including an LXG-family toxin named herein TelE displaying pore-forming activity. Seven homologs of TelE harboring a conserved glycine zipper motif at the C-terminus were identified in *Sgg* isolates. Scanning mutagenesis of this motif showed that the glycine residue at position 470 was crucial for TelE pore-forming activity. Unlike other pore-forming toxins commonly antagonized by a membrane protein, TelE activity was antagonized by a cytosolic protein TipE belonging to the DUF5085 family. Finally, we demonstrated that purified TelE can permeabilize *E. coli* cells rapidly when introduced extracellularly. Overall, we report herein a unique *Sgg* T7SSb effector exhibiting a pore-forming activity against non-immune bacteria.

Author summary

In this study, 38 clinical isolates of *Streptococcus gallolyticus* subsp*. gallolyticus* (*Sgg*) were sequenced and a genetic locus encoding the Type VIIb secretion system (T7SSb) was found conserved in all 38 genomes. This locus absent from the closely related *S. gallolyticus* subsp. *pasteurianus*, constitutes a pathogenicity island. T7SSb is a specialized machinery involved in the secretion of multiple effectors that have been implicated in bacterial virulence, interbacterial killing or host immunity modulation. Here, we report that the model organism *Sgg* strain UCN34 secretes six T7SSb effectors. One of the six effectors named TelE displayed a strong toxicity when expressed in Gram-negative bacteria. Our results indicate that TelE is a pore forming toxin whose activity can be antagonized by a non-canonical immunity protein named TipE. Overall, we report a unique toxin-immunity protein pair and our data expand the range of effectors secreted through T7SSb. As shown previously for gallocin, a bacteriocin giving a selective advantage to *Sgg* in the tumoral microenvironment, we propose that T7SSb effectors including TelE may help *Sgg* to alter the host microbiota contributing to colon colonization.

Keywords: *Streptococcus gallolyticus*; Type VIIb secretion system; pore-forming toxin; LXG family toxin; glycine zipper

Introduction

Streptococcus gallolyticus subsp*. gallolyticus* (*Sgg*) is a commensal bacterium in the rumen of herbivores*.* The first complete genome of strain UCN34 provided clear insights for its adaptation to the rumen (Rusniok et al., 2010)*. Sgg* is also an opportunistic pathogen causing septicemia and endocarditis in the elderly who often have concomitant colon tumours (Corredoira et al., 2014, Corredoira-Sanchez et al., 2012, Corredoira et al., 2015, Kim et al., 2010, Kwong et al., 2018, Wending et al., 2006, Boleij et al., 2011, Zhang et al., 2015). *Sgg* was formerly known as *S. bovis* biotype I belonging to the large *S. bovis / equinus* complex, which is comprised of several opportunistic pathogens and non-pathogenic bacteria. In this complex, *Sgg* is assigned into a clade with two other closely related subspecies named *S. gallolyticus* subsp. *pasteurianus* (referred to hereafter as *Sgp*) and *S. gallolyticus* subsp. *macedonicus* (referred to hereafter as *Sgm*) (Dekker and Lau, 2016, Schlegel et al., 2003). In general, *Sgp* is considered less pathogenic than *Sgg*, mainly causing neonatal meningitis (Onoyama et al., 2009, Chen et al., 2020), whereas *Sgm* is generally recognized as a safe non-pathogenic strain (Georgalaki et al., 2002, Maragkoudakis et al., 2009).

The Type VIIb Secretion System (T7SSb) is a specialized secretion system exploited by Firmicutes to secrete various substrates or effectors involved in host immune system modulation or bacterial competition (Cao et al., 2016, Anderson et al., 2017, Whitney et al., 2017, Anderson et al., 2011, Unnikrishnan et al., 2017). T7SSb is distantly related to the Type VII secretion system originally discovered in *Mycobacterium tuberculosis.* These two evolutionarily related secretion systems have

two components in common; a main FtsK/SpoEIII ATPase (EssC/EccC) constituting

the motor of the secretion machinery, and a secreted effector named EsxA/ESAT6

belonging to the WXG100 family (Abdallah et al., 2007). In Firmicutes, the T7SSb core

machinery is composed of 4 additional proteins namely EssA, EsaA, EsaB and EssB.

T7SSb is involved in the secretion of a variety of effectors including (i) WXG100

proteins of approximately 100 amino acids (aa) with a WXG motif in the center such

as EsxA and EsxB (Burts et al., 2005); (ii) proteins of approximately 100-aa without a

WXG motif (named WXG100-like proteins) such as EsxC and EsxD (Anderson et al.,

2013, Burts et al., 2008); (iii) protein with an N-terminal domain COG5444 such as

EsaD/EssD (Cao et al., 2016, Ohr et al., 2017); and (iv) polymorphic toxins containing

a N-terminal domain of about 200 aa with an LXG motif (also known as LXG family

proteins) such as TelA, TelB, TelC, TspA and EsxX (Ulhuq et al., 2020, Whitney et al.,

2017, Dai et al., 2017, Holberger et al., 2012). Previous studies revealed that these

effectors function as a NADase toxin (TelB) (Whitney et al., 2017), a lipid II

phosphatase (TelC) (Whitney et al., 2017), nucleases (EsaD, YobL,, YxiD) (Cao et al.,

2016, Ohr et al., 2017, Holberger et al., 2012), or a membrane depolarizing toxin (TspA)

(Ulhuq et al., 2020).

Importantly, T7SSb of *Sgg* strain TX20005 was previously shown to contribute to the

development of colon tumors in a murine model (Taylor et al., 2021). Here, using

comparative genomics, we identified that the T7SSb locus residing in a 26-kb genomic

island is uniquely present in *Sgg* but neither in *Sgp* nor in *Sgm*. Subsequently, we

found that six proteins were secreted through the T7SSb machinery of the reference

strain UCN34, including the WXG100 effector EsxA, two hypothetical proteins, and

three LXG family proteins including two TelC homologs. One of the unknown LXG

family protein was characterized herein and named TelE. We show that TelE is a pore-

forming protein whose activity is antagonized by a non-canonical immunity protein.

Results

A T7SSb locus specific to *Sgg*. In order to identify the specific pathogenic trait(s) of *Sgg*, comparative genomics was performed on newly sequenced 13 *Sgp* and 38 *Sgg* clinical isolates obtained from the National Reference Center of Streptococci in France. Phylogenetic analysis of these newly sequenced genomes, together with the complete genomes of *Sgg* and *Sgp* publicly available on NCBI GenBank assigned *Sgg* and *Sgp* into two distinct clades, confirming their correct classification **(Figure S1)**. BLASTn-based comparative genomic analysis of the reference genome UCN34 together with the representative *Sgg* and *Sgp* genomes uncovered a genomic island unique to *Sgg* **(Figure 1A)**. This genomic island contains the six genes (*esxA*, *essA*, *esaB*, *essB*, *essC* and *esaA*) encoding the core components of the Type VIIb Secretion System (T7SSb) machinery **(Figure 1A and 1B)**.

An extended protein homology search of each T7SSb core components across all the newly sequenced genomes confirmed the conservation of T7SSb in *Sgg* genomes and its absence in *Sgp* or in *Sgm* **(Figure 1C)**. A closer inspection of the T7SSb genetic locus uncovered two different gene arrangements in the *Sgg* isolates sequenced so far. In a vast majority of *Sgg* isolates, the T7SSb locus is arranged in the order of *esxA/essA-esaB-essB-essC-esaA*, as in our reference strain UCN34. Notably, *esxA* is transcribed in the opposite direction from the other five genes *essA-esaB-essB-essC-esaA,* a unique feature distinct from other Firmicutes. In contrast, in a small minority of *Sgg* isolates including the US isolate TX20005, the T7SSb genes are arranged in a different order of *esxA-esaA-essA-esaB-essB-essC* with an evident shuffling of *esaA*, and the six genes transcribed in the same direction **(Figure 1D)**.

***Sgg* UCN34 secretes 6 T7SSb effectors.** In order to identify the T7SSb effectors, we inactivated the T7SSb machinery, through in-frame deletion of *essC* gene encoding the main ATPase, in the strain UCN34. We next compared the secretome of the UCN34 WT strain to that of the Δ*essC* mutant. This approach identified 6 proteins that were differentially expressed **(Figure 2A, Figure S2A)** including the ubiquitous WXG100 effector (EsxA/Gallo_0553) that was found 4-fold less abundant in the Δ*essC* mutant, confirming that the Δ*essC* mutant is a *bona fide* T7SSb-deficient strain. In addition, five proteins namely Gallo_0559, Gallo_0560, Gallo_0562, Gallo_1068 and Gallo_1574 were detected only in the secretome of the UCN34 WT but not in the Δ*essC* mutant.

Prediction of the putative function of these effectors by *in silico* SMART analysis revealed that Gallo_0559 contains a domain of unknown function DUF5082, Gallo_0560 as a TIGR04197 family protein, whereas Gallo_0562, Gallo_1068 and Gallo_1574 contain the prototypical N-terminal LXG domain (pfam04740) found in other T7SSb effectors **(Figure 2B)**. Protein homology search indicated that Gallo_1068 and Gallo_1574 are homologous (>98% coverage, >40% similarity) to the *Streptococcus intermedius* TelC effector that functions as a lipid II phosphatase (Whitney et al., 2017). Similar to TelC, both Gallo_1068 and Gallo_1574 contain an intact C-terminal aspartate-rich motif critical for TelC enzymatic activity. We thus propose to rename Gallo_1068 and Gallo_1574 as TelC1 and TelC2, respectively **(Figure 2B)**. The third LXG-containing effector (Gallo_0562) is a protein of 514 amino acids sharing no similarities with known proteins that we propose to rename TelE in accordance with the gene nomenclature (<u>T</u>oxin exported by <u>E</u>sx with <u>L</u>XG domain)

proposed for *S. intermedius* (Whitney et al., 2017). Of note, 4 out of 6 proteins

identified in the secretome of UCN34 WT are encoded in the T7SSb locus and three

are encoded by genes likely transcribed as an operon, namely *gallo_0559*, *gallo_0560*

and *telE (gallo_0562)*.

TelE is toxic when overexpressed in Gram-negative bacteria. LXG family proteins

are polymorphic toxins implicated in interbacterial competition (Whitney et al., 2017).

However, we failed to observe any role of *Sgg* T7SSb in interbacterial competition

under standard laboratory conditions. We reasoned that this could be due to the low

expression of T7SSb effectors as reflected by the low total peptide counts in the

proteomics data **(Figure S2A)** and decided to test their potential antibacterial activity

through heterologous and inducible expression in the Gram-negative *Escherichia coli*

K-12. As shown in **Figure S2B**, overexpression of *telE*, but none of the other T7SSb

effectors, caused a strong toxicity in *E. coli* or in another Gram-negative bacterium

Pseudomonas aeruginosa, but not in the Gram-positive bacteria *Enterococcus*

faecalis, *Streptococcus agalactiae* and *Sgm* **(Figure 2C)**. Since TelC was shown

previously to be toxic to Gram-positive bacteria (Whitney et al., 2017), TelC1 and

TelC2 were then expressed in *Sgm* and *E. faecalis.* Although the survival of *Sgm* or

E. faecalis was not affected upon induced expression of TelC1 or TelC2, we observed

an altered colony morphology in *Sgm* **(Figure S2C)**. Of note, the lack of activity of

TelC1, TelC2 and TelE in Gram-positive bacteria could be due to the lower production

of these effectors in these cells by unknown mechanisms (data not shown).

TelE is encoded in the vicinity of the T7SSb core machinery locus (**Figure 3A**). This

prompted us to examine the prevalence and the conservation of TelE among the *Sgg*

clinical isolates. As shown in **Figure S3,** TelE homologs were found in over 90% of

Sgg isolates. Based on their primary amino acid sequences, TelE homologs can be

categorized into 7 variants named TelE1 (initial TelE found in UCN34) through TelE7.

Notably, multiple copies of TelE homologs can be found in a few *Sgg* genomes. Some

of these TelE homologs are N-terminally truncated, resembling the orphan modules

commonly seen in polymorphic toxins (Jamet and Nassif, 2015) **(Figure S3).** To

assess the antimicrobial activity of these TelE homologs, each TelE variant was

expressed in *E. coli.* As shown in **Figure 3B,** all the TelE variants exhibited a strong

toxicity upon induction in *E. coli*. Interestingly, a multiple sequence alignment of these

seven TelE variants uncovered a highly conserved sequence motif at the C-terminus

(**Figure S4**). This motif, rich in small hydrophobic amino acids alternating with

conserved glycine residues, is similar to the glycine zipper motif (GxxxGxxxG) found

in many membrane proteins (Kim et al., 2005) **(Figure 3C)**. Three highly conserved

glycine residue were identified at positions 458, 470 and 474, and three less-

conserved glycine residues were found at positions 466, 478 and 480 **(Figure 3C and**

3D). Single replacement of each glycine residue to a slightly bulkier valine residue

demonstrated that those at positions 466, 470 and 474 were key for TelE activity in *E.*

coli **(Figure 3E)**, with the replacement of the glycine residue at position 470 showing

the strongest impact on TelE activity (**Figure 3E**). Together, these data showed that

nearly all the *Sgg* isolates sequenced as to date encode a functional TelE homolog

whose primary sequence varies from one isolate to another, but all 7 TelE variants

contain a conserved glycine zipper motif at the C-terminus essential for TelE

antimicrobial activity.

TelE is a pore-forming toxin. Structural modelling of TelE did not show any similarities with known proteins but indicated that TelE secondary structures are rich in alpha-helices **(Figure 4A).** Alpha-helical structures with glycine zipper motifs are the two main characteristics of membrane proteins with a high propensity to form homo-oligomeric ion channels (Kim et al., 2005), suggesting that TelE might have a pore-forming activity. To investigate this possibility, we used propidium iodide, a large molecule (668 Da) that is conventionally used to assess the presence of membrane pores or membrane disruption in bacterial cells (LaCourse et al., 2018, Fridman et al., 2020). Propidium iodide can only enter the cellular cytoplasmic space through membrane lesions or large membrane pores to subsequently bind to intracellular DNA and emit a red fluorescence signal. Using time-lapse fluorescence microscopy, we observed a gradual increase of red fluorescence signal in *E. coli* cells expressing TelE, but not in control cells carrying the empty vector **(Figure 4B)**, indicating that TelE expression resulted in an influx of propidium iodide into the cells. Intriguingly, this influx was preceded immediately by shrinkage of the bacterial cell visible by phase contrast microscopy **(Figure 4C)**. Similar morphological changes have been observed on cells experiencing water loss or cytoplasmic content leakage (Sutterlin et al., 2016, Shi et al., 2020), which can be due to the excessive presence of membrane pores. Our microscopy observations thus indicate that TelE can form membrane pores in *E. coli* cells.

Next, we probed the subcellular localization of TelE in *E. coli* by using TelE with an hexahistidyl tag either at the N- or C-terminus. Notably, we found that TelE with a His-tag at the C terminus (TelE-His) is significantly less active in intoxicating *E. coli* as compared to the native or N-terminally tagged TelE. Since TelE-His was still functional

when overexpressed in strain BL21(DE3) **(Figure S5A-C)**, we proceeded with this less

toxic TelE-His as a means to monitor TelE subcellular localization. As shown in **Figure**

4D, TelE-His was predominantly found in the inner and outer membrane of *E. coli*. On

the basis that TelE localizes in the membrane and its expression induces cellular

uptake of propidium iodide from the extracellular medium, we propose that TelE is a

pore-forming toxin. We next investigated the subcellular localization of the inactive

TelE$_{G470V}$ mutant by adding a hexahistidyl tag at the C-terminus (TelE$_{G470V}$-His). As

shown in **Figure 4D**, this construct was stably expressed in *E. coli* cells and was able

to localize in the membrane as the wildtype TelE-His, indicating that the loss of activity

of the inactive TelE$_{G470V}$ variant was not due to protein instability or protein

delocalization in the cells.

The cognate immunity protein TipE antagonizes TelE toxicity. The activity of LXG

family toxins such as TelA, TelB and TelC can be antagonized by cognate immunity

proteins (Whitney et al., 2017). To identify the immunity protein able to counteract TelE

toxicity, each gene located downstream from *telE* was cloned and co-expressed in *E.*

coli cells using a second compatible plasmid (pJN105). We found that only co-

expression of the third gene downstream from *telE* (*gallo_0565*) rescued *E. coli* from

TelE intoxication **(Figure 2A, 5A)**. This gene encodes a 151 amino acids-long protein

renamed herein as TipE (<u>T</u>el <u>i</u>mmunity <u>p</u>rotein), which harbor a domain of unknown

function DUF5085. An additional DUF5085 domain-containing protein (Gallo_0570),

sharing ~30% protein sequence similarity with TipE, was found five genes downstream

of TipE **(Figure 2A)**. However, co-expression of this protein did not protect *E. coli* from

TelE intoxication, suggesting a specific interaction between TipE and TelE **(Figure**

5A).

To decipher how TipE confers immunity, we constructed a recombinant TipE tagged

with a 9-aa hemagglutinin tag (HA tag) at the N-terminus for the detection of the

recombinant TipE using a specific anti-HA monoclonal antibody. We showed that the

recombinant HA-TipE was functional and able to rescue *E. coli* from TelE intoxication

(Figure S5B). Subcellular fractionation experiments indicates that TipE is

predominantly a soluble protein **(Figure 5B)**. To determine whether TipE is a

periplasmic or a cytosolic protein, we performed a separate extraction of periplasmic

proteins but did not detect TipE in this fraction (**Figure 5B**). Thus, we conclude that

TipE is a cytosolic protein. In contrast, when HA-TipE was co-expressed with TelE-

His, these two proteins were mostly localized in the insoluble fraction **(Figure S5C)**,

suggesting that TipE interacts with TelE to counteract its function.

To demonstrate that TipE interacts with TelE, a protein pulldown experiment was

carried out using nickel-immobilized resin. Incubation of TelE-bound resin with cellular

lysate containing HA-TipE captured a significant amount of TipE, which was detected

through immunoblot using anti-HA specific antibodies following TelE elution (**Figure

5C**). Importantly, TipE was not detected in similar amounts when the cellular lysate

was incubated with free nickel-immobilized resin (**Figure 5C**), indicating that TipE

indeed interacts with TelE. To monitor in more detail the impact of TipE on TelE in live

cells, we constructed a chimeric TelE variant C-terminally fused with superfolder green

fluorescent protein (TelE-sfGFP). This TelE chimera protein, upon excitation, allows

direct visualization of TelE in live cells using fluorescence microscopy. We first showed

that TelE-sfGFP is still toxic in *E. coli* **(Figure S5A)** and is able to cause cellular influx

of propidium iodide when expressed **(Figure 5D)**. As shown in **Figure 5D**, sfGFP was

found uniformly distributed throughout the cytoplasm in its native form **(Figure 5D)**.

However, in *E. coli* cells expressing TelE-sfGFP, TelE was found in small foci

distributed around the cell body **(Figure 5D)**. In contrast, in *E. coli* cells co-expressing

TelE-sfGFP and TipE, the GFP signal was mainly observed in the cytoplasm of these

cells **(Figure 5D)**, suggesting that TelE cellular localization was altered in the presence

of the immunity protein TipE. Taken together, these data indicate that TipE

antagonizes TelE activity by disrupting TelE cellular localization through protein-

protein interaction.

Purified recombinant TelE protein is able to permeabilize live *E. coli* cells. Since

TelE is a T7SSb effector found in the supernatant of *Sgg* UCN34, we wondered

whether TelE was able to induce pore-formation on *E. coli* cells when present in the

external milieu. To purify TelE, we turned to the TelE variant appended with 9-aa long

HA tag at the C-terminus (TelE-HA) that exhibits drastic reduction of toxicity in *E. coli*.

This allowed the purification of sufficient amount of TelE from the membrane fraction.

We next introduced the purified TelE onto *E. coli* cells labelled with the fluorescent dye

carboxyfluorescein diacetate succinimidyl ester (CFSE). This cell-tracker dye is

fluorescent only intracellularly upon cleavage by a non-specific esterase. It is long

lasting and well retained dye within labeled cells. TelE addition to fluorescent gut-

adapted *E. coli* pks+ induced a rapid leakage of the fluorescent dye in 10 min at 37°C

(**Figure 6**). A similar effect can be seen when the cells were treated with membrane-

permeabilizing detergent such as Triton X-100 or Tween 20 (**Figure 6A lower panel**

and 6B). Importantly, detergent octyl glucoside used for TelE purification had little

effect on *E. coli* permeabilization. We further confirmed that the leakage of fluorescent

dye from TelE-treated *E. coli* cells was not due to the detergent octyl glucoside by

incubating the cells with TelE depleted of detergent either by beta-cyclodextrins (TelE1)

or Bio-Beads (TelE2 and TelE3) (**Figure 6C**). Taken together, we conclude that TelE

is able to permeabilize live *E. coli* cells when added in the external milieu.

Discussion

In this study, we show that a T7SSb cluster is a putative pathogenicity island conserved in all the clinical isolates of *Streptococcus gallolyticus subsp. gallolyticus* (*Sgg*). Interestingly, two different T7SSb genetic arrangements can be found in *Sgg* clinical isolates. The most prevalent T7SS arrangement resembles those found in *Streptococcus intermedius*, whereas the T7SS of a small minority of *Sgg* isolates is identical to those of *Staphylococcus aureus* (Cao et al., 2016, Whitney et al., 2017) **(Figure 1D)**. Six T7SSb effectors were identified in strain UCN34, including the prototypical EsxA/ESAT-6, and three LXG-containing proteins. Amongst the three LXG effectors, two encoded outside of the T7SSb locus are homologs of TelC, a lipid II phosphatase shown to inhibit peptidoglycan biosynthesis in Gram-positive bacteria (Whitney et al., 2017, Klein et al., 2018). The third LXG effector of unknown function is unique to *Sgg* and is encoded in vicinity of the locus encoding T7SSb machinery. Here, we showed that TelE exhibits pore-forming activity and its activity can be neutralized by a cytoplasmic protein TipE.

TipE is an atypical immunity protein. In other pore-forming toxins similarly containing glycine zipper motifs such as CdzC/D from *Caulobacter crescentus* and Tse4 from *Pseudomonas aeruginosa*, the cognate immunity proteins are multi-pass transmembrane proteins residing predominantly in the inner membrane (Garcia-Bayona et al., 2017, LaCourse et al., 2018). Similarly, an inner membrane protein protects the *E. coli* cells expressing the *S. aureus* membrane depolarizing T7SSb toxin TspA (Ulhuq et al., 2020). Remarkably, the six-pass transmembrane protein Gallo_0563 encoded by the gene located immediately downstream of *telE* did not

protect *E. coli* from TelE intoxication **(Figure 5A, 5B)**, and our data showed that only

TipE is able to block TelE activity through direct interaction and sequestration. Notably,

TipE is a DUF5085 family protein predicted related to T7SS in the Pfam database.

Based on our data, we propose herein that DUF5085 family proteins could serve as

immunity proteins to T7SSb toxins.

Interestingly, a diverse pool of TelE variants (TelE to TelE7) exist in the genomes of

Sgg isolates, all harboring the conserved C-terminal glycine zipper motif essential for

TelE activity. Glycine zipper motif is a common feature in membrane proteins

displaying pore-forming activity. In addition to the aforementioned CdzC/D and Tse4,

glycine zipper motif is also found in the vacuolating toxin VacA from *Helicobacter pylori*,

the prion protein (PrP) and the amyloid beta peptide (Aβ) (Kim et al., 2005), many of

which display pathological roles. For instance, VacA is involved in gastric ulceration

caused by *H. pylori*, whereas PrP and Aβ are associated with prion and Alzheimer's

disease, respectively[31]. In line with previous observations (LaCourse et al., 2018, Kim

et al., 2005, McClain et al., 2003), mutation of the key glycine residue in this motif

abolishes TelE activity, yet did not alter TelE membrane localization. Since the glycine

residues of the motif are usually positioned in the inner part of membrane pores, we

speculate that change of the small glycine to a bulkier valine residue may affect the

oligomerization and/or pore opening of TelE, as demonstrated for other glycine zipper

toxins (Kim et al., 2005).

In addition to TelE, we discovered two additional yet to be characterized T7SSb

effectors namely Gallo_0559 and Gallo_0560 located immediately upstream of TelE

and probably encoded in the same operon. Bioinformatic analysis showed that

Gallo_0559 is a DUF5082 family protein whereas Gallo_0560 is a TIGR01497 family

protein. Proteins belonging to these two families were previously identified in *S.*

intermedius as WxgA and WxgC, respectively (Whitney et al., 2017). WxgA and WxgC

are chaperone proteins essential for the secretion of LXG family proteins (Tel) in *S.*

intermedius (Whitney et al., 2017). These proteins were suggested as WXG100-like

proteins (Whitney et al., 2017), but WxgA and another chaperone protein WxgB are

40% longer than the canonical WXG100 proteins. These chaperones have not been

detected in the *S. intermedius* bacterial supernatant and were henceforth

hypothesized to disassociate from Tel proteins prior to or during secretion (Whitney et

al., 2017). Here we found that these probable Wxg-like chaperones were detected in

the supernatant of *Sgg* WT but not in the Δ*essC* mutant, indicating that the secretion

of these proteins is T7SSb-dependent, and that they may remain associated with LXG

toxins post secretion. Further investigation on the potential interaction of these two

proteins with TelE may provide insight into T7SSb effectors secretion and delivery.

The three LXG family toxins identified in this study expands our knowledge on the

antibacterial arsenal possessed by *Sgg*. As demonstrated here, TelE is a toxin able to

permeabilize Gram-negative bacteria, whereas the two other LXG family toxins TelC1

and TelC2 homologous to lipid II phosphatase TelC probably targets Gram-positive

bacteria (Whitney et al., 2017). It is noteworthy that *Sgg* also produces bacteriocins

such as gallocin exhibiting antimicrobial activity against gut commensals such as *E.*

faecalis (Aymeric et al., 2018), which together with the newly identified Tel toxins, are

likely the arsenal essential for *Sgg* survival in the highly dynamic and populated human

gut environment.

Finally, our findings may have further implications in understanding the pathogenicity of *Sgg*. Indeed, it was shown in *Sgg* strain TX20005 that T7SSb is a key component contributing to *Sgg* gastrointestinal tract colonization and cancer-promoting capability (Taylor et al., 2021). Of note, TX20005 belongs to the small minority of *Sgg* isolates displaying a different T7SSb gene locus arrangement as compared to UCN34, and the impact of these differences on the pathogenic outcome is currently unclear. The contribution of pathogenicity associated islands in bacterial virulence is well established. For instance, the CagA toxin, VacA toxin and Type IV secretion system encoded in the *Helicobacter pylori* pathogenicity island (Franco et al., 2008, Wen and Moss, 2009), or the colibactin synthase genes encoded in the *E. coli* polyketide synthase (*pks*) genotoxic island (Arthur et al., 2012, Cuevas-Ramos et al., 2010, Pleguezuelos-Manzano et al., 2020) are key to the progression of gastric cancer and colorectal cancer, respectively. Of particular interest is VacA that also contains a glycine zipper motif essential for its function causing ulceration and gastric lesions in murine model (Telford et al., 1994). Whether TelE could also alter host cells is a perspective for future studies.

Methods

Bacterial culture conditions

All bacterial strains and plasmids used in this study are listed in **Table S1**. Cultures of *E. coli* and *P. aeruginosa* strains were prepared from single colonies in lysogeny broth (LB) (BD) and incubated overnight at 37°C, 200 rpm, whereas cultures of *E. faecalis*, *S. agalactiae* and *S. gallolyticus* strains were prepared from single colonies in brain heart infusion broth (BHI) (Acumedia) and incubated statically overnight at 37°C. When the use of growth medium with minimal amount of peptides were necessary, overnight culture of *Sgg* were prepared from single colonies in M9YEG broth (1X M9 minimal salts supplemented with 0.5% yeast extract and 1.0% of glucose) (Teh et al., 2019) and incubated statically at 37°C. When appropriate, antibiotics were supplemented at the following concentrations: kanamycin, 50 µg/mL for Gram-negative bacteria and 500 µg/mL for Gram-positive bacteria; gentamicin, 20 µg/mL for *E. coli*, and 60 µg/mL for *P. aeruginosa*; erythromycin, 2 µg/mL for Gram-positive bacteria.

Genetic techniques

Standard molecular cloning procedures as previously described (Green et al., 2012) were followed to construct plasmids for gene deletion and gene expression. All primers used in this study were synthesized by Integrated DNA Technology and listed in **Table S1**. The enzymes Q5 DNA polymerase and restriction enzymes were obtained from New England Biolabs, whereas T4 ligase was obtained from Thermo Fisher Scientific. All genetic constructs were delivered into chemically transformed *E. coli* DH5α prepared as described (Inoue et al., 1990) and verified by Sanger sequencing

performed at 1st BASE. For gene expression plasmids construction, the

corresponding open reading frames with the native ribosomal binding sites were

cloned downstream of inducible promoters such as P_{tetO} (anhydrotetracycline-

inducible) and P_{BAD} (arabinose-inducible).

To construct plasmids for gene deletion, two ~500bp DNA fragments corresponding

to the upstream and downstream region of the gene of interest were first PCR

amplified. The resulting two DNA fragments were subsequently spliced by overlap

extension PCR and ligated into pG1 plasmid at *Sma*I restriction site. The constructed

plasmids were transformed into *E. coli* and later transformed into *S. gallolyticus* as

described in the transformation section below. Subsequently, *Sgg* in-frame

chromosomal deletion mutants were generated by allelic exchange as previously

described (Danne et al., 2013).

To generate plasmids encoding TelE with a specific residue substitution, *telE* was

PCR-amplified into two fragments corresponding to the 5' and 3' regions by using

primers with the desired mutation incorporated. These two fragments were spliced into

a full-length *telE* open reading frame by overlap extension PCR and ligated into

pTCV*erm*-ptetO at the *Bam*HI/*Sph*I restriction site.

Plasmid encoding TelE with the C-terminus fused with superfolder GFP was

constructed by splicing the PCR amplicons corresponding to the *telE* open reading

frame without a stop codon and the superfolder GFP open reading frame that has

been codon-optimized for *Sgg* (gene synthesized at GenScript).

Transformation

E. coli DH5α was chemically transformed as described in (Inoue et al., 1990) whereas *P. aeruginosa* was transformed by electroporation as previously described (Smith and Iglewski, 1989). *E. faecalis* and *S. agalactiae* were transformed by electroporation as previously described (Bae et al., 2002, Dunny et al., 1991, Biswas et al., 1993). *S. agalactiae* carrying the respective plasmids were used for conjugal transfer into *Sgm* as described previously (Danne et al., 2013). *Sgg* was naturally transformed using a newly established method, by supplementing the log-phase culture grown in M9YEG with 10 µM of the competence inducing peptides (ITGWWGL) outlined in the previous study (Morrison et al., 2013), at least 5 µg of plasmid DNA, 10 µM of magnesium chloride, 10 µM of magnesium sulfate and 1 µM of ferrous (II) chloride. The mixture was incubated for an hour at 37°C, followed by three hours at 30°C, before being plated onto BHI agar supplemented with erythromycin.

Genomic DNA isolation

S. gallolyticus genomic DNA was extracted using either Wizard Genomic DNA Purification Kit or Wizard SV Genomic DNA Purification Kit (Promega). The extraction was performed according to the manufacturer's instructions, except that 20U to 50U of mutanolysin (Sigma-aldrich) was added at the lysozyme treatment step. When necessary, an additional phenol:chloroform clean up step was performed to remove the excessive impurities in the samples.

Genomic DNA sequencing and assembly

High integrity DNA were sequenced to at least a coverage of 100X on the Illumina Miseq platform (reagent kit v3, 2 x 300bp) at the in-house next-generation sequencing

facility in the Singapore Centre for Environmental Life Sciences Engineering. The

paired-end sequencing reads were trimmed and assembled *de novo* into contigs on

the CLC Genomics Workbench 10.0 (Qiagen). These assembled contigs were

subsequently uploaded to the RAST server (Aziz et al., 2008, Brettin et al., 2015,

Overbeek et al., 2014) for open reading frame prediction and gene annotation.

Phylogenetic analysis

A core genome-based alignment was performed on Parsnp version 1.2 integrated in

the Harvest suite (Treangen et al., 2014), with *Sgg* UCN34 (GenBank accession no.

FN597254) served as the reference genome. The other *S. gallolyticus* complete

closed genomes available on NCBI, *i.e. Sgg* ATCC43143 (GenBank accession no.

NC_017576), ATCC BAA_2069 (GenBank accession no. NC_015215) and DSM

16831 (GenBank accession no. NZ_CP018822); *Sgp* ATCC43144 (GenBank

accession no. NC_015600), NCTC13784 (GenBank accession no. NZ_LS483462)

and WUSP067 (GenBank accession no. NZ_CP039457); and two additional draft

genomes – *Sgm* ACA-DC-206 (unpublished genome) and the *Sgg* TX20005 (NCBI

Assembly ID GCA_000146525) - were also included in the analysis for a clear

distinction of these closely related strains. The resulted alignment was subsequently

visualized on Gingr (integrated in the Harvest suite) and used for phylogenetic

inference on FigTree2 based on approximate-maximum-likelihood algorithm (Price et

al., 2010).

Comparative genomics

A genome-wide comparison among *Sgg* UCN34, three representative *Sgg* clinical

isolates, and three representative *Sgp* was performed using BLAST Ring Image

Generator v0.95 based on BLASTn search (Alikhan et al., 2011). Genomic islands of

the *Sgg* UCN34 were downloaded from IslandViewer 4(Bertelli et al., 2017). NCBI

tBLASTn program was used to identify the protein homologs of the T7SS core

components across all the *Sgg* and *Sgp* clinical isolates. Easyfig v2.2.2 was used to

visualize the region of interest and to perform pairwise comparisons on the protein

sequences using tBLASTx program (Sullivan et al., 2011).

Comparative proteomics on the *Sgg* secretomes

Single colonies of *Sgg* UCN34 WT and the Δ*essC* mutant were inoculated into 20 mL

of M9YEG and grown to OD_{600nm} 3.0 corresponding to the early stationary phase of

growth. The culture supernatant was collected and filtered through 0.22 μm filter units.

15 mL of the filtered culture supernatant was first concentrated using Amicon Ultra 15

mL Centrifugal Filters (cut-off 10 kDa, Merck), and the proteins were precipitated out

of the solution using trichloroacetic acid (TCA) as previously described (Flaugnatti and

Journet, 2017). The samples were further processed and analyzed at the Taplin Mass

Spectrometry Facility. Briefly, precipitated proteins were first treated with trypsin

(Promega) in 50 mM ammonium bicarbonate solution, dried in a speed-vac and were

subsequently reconstituted in HPLC solvent A (2.5% acetonitrile, 0.1% formic acid).

Peptides were loaded using a Famos auto sampler (LC Packings) into a nano-scale

reverse-phase HPLC fused silica capillary column (100 μm inner diameter x ~30 cm

length) packed with 2.6 μm C18 spherical silica beads and eluted with increasing

concentrations of solvent B (97.5% acetonitrile, 0.1% formic acid). Eluted peptides

were subjected to electrospray ionization and entered into an LTQ Orbitrap Velos Pro

ion-trap mass spectrometer (Thermo Fisher Scientific). Peptides were detected,

isolated, and fragmented to produce a tandem mass spectrum of specific fragment

ions for each peptide. Peptide sequences (and hence protein identity) were

determined by comparing the fragmentation pattern acquired by the software program

Sequest (Thermo Fisher Scientific) (Eng et al., 1994) to the protein sequences on

protein databases. All databases include a reversed version of all the sequences and

the data was filtered to between a one and two percent peptide false discovery rate.

Downstream data analysis was performed in R studio using the Differential

Enrichment analysis of Proteomics Data 1.10.0 package (Zhang et al., 2018). Proteins

that were not detected across all the three replicates of either the UCN34 WT or the

Δ*essC* mutant were excluded from the analysis. Next, the total peptide count of the

remaining proteins (n = 162) was scaled and variance stabilized (Huber et al., 2002)

before the missing values (*i.e.* peptide count = 0) were imputed with the assumption

that the missing values are missing not at random (MNAR), using the Deterministic

Minimum Imputation (MinDet) strategy (10^{-4} quantile) (Lazar et al., 2016). The MinDet-

imputed data were subsequently used for differential expression analysis, with a cut

off of false discovery rate of 5%, and a $\log_2$ fold change of 1. The derived data set was

plotted using the EnhancedVolcano package (Blighe, 2018).

Bacterial toxicity experiments

For an intracellular expression of the *Sgg* T7SSb effectors, *E. coli* DH5α, *E. faecalis*

OG1RF, *S. agalactiae* NEM316 and *Sgm* ACA-DC-206 were transformed with

pTCVerm-P$_{tetO}$ plasmid series, whereas *P. aeruginosa* PAO1 was transformed with

pUCP22-P$_{tetO}$ plasmid series (see **Table S1** for the plasmids used in this study).

Overnight cultures of these strains were serially diluted in tenfold increments and 5 µL

of each dilution was spotted onto LB or BHI agar supplemented with an appropriate

amount of antibiotics and inducer (kanamycin, 50 or 500 µg/mL; anhydrotetracycline,

100 or 500 ng/mL). Agar plates incubated overnight at 37°C were documented on the

Gel Doc XR+ Documentation System (Bio-Rad).

For the identification of the TelE cognate immunity protein, the *E. coli* DH5α carrying

pTCVerm-P$_{tetO}$::TelE was transformed with compatible plasmids of either pJN105

(vector control) or pJN105 containing each individual open reading frames found

downstream of *telE*. The expression of these open reading frames was driven by

arabinose-inducible promoter (P$_{BAD}$). Overnight cultures of these strains were first 50x-

diluted into sterile LB broth supplemented with antibiotics and incubated at 37°C with

agitation at 200 rpm for 1.5 hours. These bacterial cultures at the early log-phase of

growth were supplemented with arabinose to a final concentration of 0.5%, and

incubated further for 3 hours, before being subjected to serial dilution and spotted onto

LB agar supplemented with an appropriate amount of antibiotics and inducer

(kanamycin, 50 µg/mL; gentamicin, 20 µg/mL; anhydrotetracycline, 100 ng/mL;

arabinose, 0.5%). Agar plates incubated overnight at 37°C were documented on the

Gel Doc XR+ Documentation System (Bio-Rad).

***In silico* TelE prediction and homologs identification**

The TelE amino acid sequence from the reference strain *Sgg* UCN34 (protein ID

CBI13054.1) was uploaded onto the web-based I-TASSER server (Roy et al., 2010,

Yang et al., 2015, Yang and Zhang, 2015) for protein structural prediction using the

default settings. The same amino acid sequence was also used to identify TelE

homologs among the clinical isolates using the NCBI tBLASTn program. Hit proteins

were considered as TelE homologs if significant similarities (e-value $<10^{-11}$) were

identified at the N terminus, C terminus, or the full-length protein.

Multiple sequence alignment (MSA) and sequence logo generation

MSA was performed on the online MAFFT server version 7

(https://mafft.cbrc.jp/alignment/server/) with E-INS-i iterative refinement method that

assumes all the sequences shared the same conserved motifs (Katoh et al., 2019,

Kuraku et al., 2013). Regional sequence logo (probability-weighted Kullback-Leibler

logo type) was generated on web-based Seq2Logo 2.0 server

(http://www.cbs.dtu.dk/biotools/Seq2Logo/), with the Heuristics clustering method

applied (Thomsen and Nielsen, 2012).

Fluorescence microscopy imaging

For time-lapse microscopy imaging, log-phase *E. coli* (OD_{600nm} ~ 1.0 Abs) cultured in

LB was mixed with propidium iodide (1:1000 diluted from 20 mM stock solution,

Thermo Fisher Scientific) and anhydrotetracycline at a final concentration of 500

ng/mL (Sigma Aldrich), and spotted onto an 1% (w/v) low melting agarose pad

(bioWORLD). Samples were immediately imaged with the microscope Axio Observer

7 fitted with Plan Apochromat 40x/0.95 dry lens for phase contrast imaging, or Plan

Apochromat 100x/1.4 oil lens for differential interference contrast imaging (Carl Zeiss),

with temperature maintained at 37°C, and aided with Definite Focus.2 for autofocus.

When necessary, samples were excited with filter wheel 555/30 to detect red

fluorescence signals.

To image *E. coli* expressing TelE-sfGFP with or without TipE, overnight cultures were diluted 50x into sterile LB broth, incubated at 37°C with 200 rpm for 1.5 hours to grow to early log-phase. Samples were first induced with 0.5% arabinose for 1.5 hours and were subsequently supplemented with anhydrotetracycline to a final concentration of 500 ng/mL and propidium iodide (1:1000 diluted from 20 mM stock solution, Thermo Fisher Scientific), and incubated further for 2 hours at 37°C without agitation. Following incubation, samples were spotted onto an 1% low melting agarose pad (bioWORLD), and imaged with the microscope Axio Observer Z1 fitted with Plan Neofluar 100x/1.3 oil lens, with excitation at 488 nm (GFP) and 535 nm (PI).

Protein expression, localization, purification and pulldown

Protein samples were in general prepared as follow: overnight cultures of *E. coli* BL21(DE3) harboring the respective protein expression pET28b plasmids were 50x diluted into sterile LB broth supplemented with 50 μg/mL of kanamycin and grown to an optical density of ~0.7 Abs measured at 600 nm. Subsequently, cultures were added with isopropyl β-D-1-thiogalactopyranoside (IPTG) to a final concentration of 100 μM and incubated for an additional 3 hours at 37°C, 200 rpm. Following incubation, cells were harvested by centrifugation at 8,000 xg for 10 minutes, and the resulted cell pellets were stored at -80°C (for protein purification) or processed immediately (for protein localization). When needed, cells were resuspended in 100 mM HEPES pH 7.5 and lysed by sonication (10s on-off pulse cycles, at 20 or 40% amplitude, Sonics Vibracell VCX750). Unbroken cells were removed by centrifugation at 8,000 xg for 10 minutes at 4°C, and the resulted supernatant was ultracentrifuged (Beckman Coulter XPN100) at 35,000 rpm for 1 hour at 4°C to separate the soluble (supernatant) and membrane proteins (pellet).

For protein localization study, the membrane pellets were resuspended in 100 mM

HEPES pH 7.5 containing 1% sarkosyl and were ultracentrifuged further to separate

the soluble inner membrane proteins and the insoluble outer membrane proteins. Both

the soluble proteins and inner membrane fractions were subjected to TCA precipitation

prior to SDS-PAGE analysis. Periplasmic proteins were extracted as described

previously(Quan et al., 2013), except that the Tris buffer was replaced with HEPES

buffer.

For protein purification, the membrane pellets were resuspended to a final

concentration of 100 mg/mL in Buffer A (100 mM HEPES pH 7.5, 500 mM NaCl and

1% n-dodecyl-β-D-maltoside (DDM)) and incubated for 1 hour at 4°C aided with top-

down rotation. The solubilized membrane proteins were subsequently supplemented

with imidazole to a final concentration of 50 mM and incubated with nickel-immobilized

HisLink resins (Promega) that had been pre-calibrated in Buffer B (100 mM HEPES

pH 7.5, 500 mM NaCl, 50 mM imidazole and 0.1% DDM), for 30 minutes at 4°C with

top-down rotation. Proteins-bound HisLink resins were washed with 30X resin volume

of ice-cold Buffer B, and finally rinsed with ice-cold buffer containing 100 mM HEPES

pH 7.5, 500 mM NaCl, 500 mM imidazole and 0.1% DDM to elute the proteins.

For protein pulldown, the proteins-bound HisLink resins were incubated with cellular

lysate containing HA-TipE prepared in Buffer B for 20 minutes at 4°C aided with top-

down rotation. The resins were subsequently washed and the bound proteins were

eluted as described above.

Purification of TelE-HA

TelE-HA was expressed in *E. coli* cells BL21(DE3) transformed with the pET28b/TelE-

HA plasmid. At OD_{600} = 0.8, TelE-HA production was induced with 0.5 mM IPTG in

Terrific Broth media (Sigma) supplemented with Kanamycin (50 µg/ml). After overnight

incubation at 18 °C, 160 rpm, cells were collected by centrifugation at 10 000 xg for

20 min at 4°C and the media was discarded. Cell pellet was resuspended in Lysis

buffer containing 50 mM HEPES pH 8, 1 mg/ml lysozyme, 4 tablets of Protease

Inhibitors (Roche), 20X-diluted benzonase. Cells were lysed by sonication (6 bursts of

1 min each at 50%, with 1 min gaps). Unbroken cells were discarded by centrifugation

at 6000 xg for 15 min and membranes were isolated after the cellular lysate was

ultracentrifuged at 100,000 xg for 1 h at 4 °C. Total membrane pellets were

resuspended in solubilization buffer containing 50 mM NaCl, 50 mM Tris pH8, 1 tablet

of Protease Inhibitors, 1 mM TCEP, 5% n-Octyl-β-D-Glucopyranoside (OG, Anatrace),

aided with gentle stirring for 30 min at 6 °C. The resuspended membrane was again

ultracentrifuged at 195 000 xg for 45 min, and the non-solubilized membranes were

removed. Solubilized membrane in the supernatant was loaded overnight onto an

Hitrap SP HP 5ml column (Pharmacia) pre-equilibrated in solubilization buffer. TelE-

HA was eluted using a 100 mM to 2 M NaCl gradient in buffer containing 50 mM Tris

pH8, 0.5 mM TCEP, 0.7% OG. Fractions were analysed by SDS-PAGE. Fractions

containing TelE-HA were pooled and dialyzed overnight against 50 mM Tris pH8, 0.5

mM TCEP, 0.7% OG, 150 mM NaCl. TelE-HA was concentrated 10 times on a

Microcon-10kDa Centrifugal Filter (Millipore). Final protein concentration was

determined to be 1.1 mg/ml by Bradford assay, using BSA in the presence of 0.7 %

OG as a standard. Glycerol was added at 8 % final and the aliquoted protein was flash-

frozen in liquid nitrogen. The protein identity was confirmed using mass-spectrometry.

SDS-PAGE and Western blot

Protein samples were mixed with an equal amount of 2X Laemmli reducing buffer and boiled for 10 minutes before being subjected to SDS-PAGE analysis on 12% polyacrylamide gels. Proteins were stained with InstantBlue solution (Expedeon) for visualization or were transferred onto PVDF membranes using iBlot Dry Blotting System (Thermo Fisher Scientific) at 20V for 7 minutes. Following protein transfer, the PVDF membranes were immediately blocked with 1X casein blocking buffer (Sigma-Aldrich), and were probed with either horseradish peroxidase (HRP)-conjugated primary antibodies against His-tag (Qiagen) or with rabbit primary antibodies against HA-tag (Sigma-Aldrich) followed by anti-rabbit HRP-conjugated secondary antibodies. Membranes were thoroughly rinsed with PBS containing 0.1% Tween 20 between the incubations. Chemiluminescent signals were detected using Amersham ImageQuant 800 System (GE Healthcare) after a brief incubation of the PVDF membranes with Immobilon Forte Western Chemiluminescent HRP Substrates (EMD Millipore). For reblotting with a different antibody after image acquisition, the membranes were incubated with Restore PLUS Western Blot Stripping Buffer (Thermo Fisher Scientific) for 30 minutes at room temperature, rinsed thoroughly before being blocked with casein blocking buffer and probed with antibodies as described above.

Acknowledgement

We are very grateful to Alexandra Doloy and Nicolas Dmytruk for providing us with the collection of S. *gallolyticus* clinical isolates. We thank Ross Tomaino from Taplin Mass Spectrometry Facility for the mass-spectrometry analysis. We thank Tarek Msadek for critical reading of the manuscript. This work was supported by the National Research Foundation and Ministry of Education Singapore under its Research Centre of Excellence Program (SCELSE). S. Dramsi would like to acknowledge the support of the Institut National contre le Cancer (INCA, grant PLBIO16-025) and from the French Government's Investissement d'Avenir program, Laboratoire d'Excellence Integrative Biology of Emerging Infectious Diseases (grant no. ANR-10-LABX-62-IBEID).

Author contributions

Conceptualization, W.K.T.; Methodology, W.K.T. and Y.D.; Investigation, W.K.T. and S. D.; Protein purification, F. G., Resources – clinical isolates, C.P.; Writing – Original Draft, W.K.T.; Writing – Review & Editing, W.K.T., Y.D., C.P., A.F., S.D. and M.G.; Visualization, W.K.T.; Supervision, A.F., S.D. and M.G.; Funding Acquisition, S.D. and M.G.

    Figure Legends

Figure 1 - A T7SSb locus specific to *Sgg*. (A) An overview of the BLASTn-based

comparative genomics analysis. From the innermost circle, the reference genome *Sgg*

UCN34 (black ring); followed by *Sgp* isolates (Gallo1, Gallo4, Gallo6, blue rings); and

Sgg isolates (Gallo37, Gallo43 and Gallo51, red rings). The outermost black blocks

represent the genomic islands in UCN34 genomes predicted by IslandViewer 4. White

spaces within each ring represent a ≤50% similarities of the respective genomic

regions to UCN34. (B) A pairwise nucleotide sequence comparison of the genomic

region of interest. Each of the T7SSb genes (arranged in the order of *esxA, essA,*

esaB, essB, essC, esaA) was represented by colored arrows. Light gray arrows

represent other open reading frames in this genomic region. Gray shading shows the

similarities between the sequences. (C) Summary of BLAST search using NCBI

tBLASTn program, queried against the protein sequences of each T7SSb components

(EsxA, EssA, EsaB, EssB, EssC, EsaA) from the reference genomes UCN34. The

identification of homologs was based on RAST annotation, or significant hits from the

BLAST search. (D) The general gene arrangement of the T7SS gene locus in *Sgg*, *S.*

intermedius and *S. aureus*. Each T7SSb genes was color-coded and labelled

accordingly. The open reading frames flanked by *esxA* and *essA* in the *S. intermedius*

B196 were excluded in this figure for a clear presentation of the T7SSb gene locus.

Figure 2. A *Sgg* T7SSb effector exhibits specific toxicity in Gram-negative

bacteria. (A) A volcano plot of the relative abundance of all proteins (n=162) detected

in the secretomes of the *Sgg* UCN34 wildtype versus the Δ*essC* mutant, ranked by

false discovery rate (adjusted *p*-value). Each dot represents an individual protein. The

vertical dotted line marked the cut-off value of $\log_2$ fold change of 1, and the horizontal

dotted line marked the cut-off value of adjusted *p*-value of 0.05. The proteins that met

the cut-off requirements were represented by red dots and labelled with the protein

identity. (B) The domain architecture of the six proteins labelled in (A). Both

Gallo_1068 and Gallo_1574 are homologous to the lipid II phosphatase TelC and the

conserved aspartate-rich motifs essential for the protein function were indicated. (C)

Viability of the bacterial cells harboring either an empty vector or a vector encoding

the full-length *telE* with the expression inducible by anhydrotetracycline (aTc). Bar

chart on the right showed the logarithm-transformed CFU count of the *telE*-expressing

cells in relative to the control. Error bars represent mean + standard deviation (n = 3).

Figure 3. T7SSb Effector TelE contains a C-terminal glycine zipper motif

essential for its activity. (A) The genomic region encoding the T7SSb core

components, TelE and TipE in *Sgg* UCN34. Black, T7SSb core components; blue,

TelE (LXG domain-containing protein); red, DUF5085 domain-containing protein);

white, hypothetical proteins. (B) Viability of the *E. coli* cells harboring either an empty

vector or a vector encoding various *telE* variants (*telE1* to *telE7*) with the expression

inducible by anhydrotetracycline (aTc). Bar chart on the right showed the logarithm-

transformed CFU count of the *telE*-expressing cells in relative to the control. Error bars

represent mean + standard deviation (n = 3). (C) Alignment of the C-terminus

sequence of the *telE* variants expressed in (B) revealed a conserved region

(underlined) enriched with hydrophobic residues. (D) Sequence logo of the underlined

region in (C) generated from all aligned TelE variants identified a sequence motif

highly similar to glycine zipper motif (GxxxGxxxG). The number at the bottom of the

sequence logo represents the position of the respective amino acid residue in

reference to the TelE sequence from UCN34. Shrunk residues (*i.e.* residues at position

462 to 465) reflected the presence of gaps in the sequence alignment. (E) Viability of

the *E. coli* cells harboring either an empty vector or a vector encoding the mutated *telE*

with the expression inducible by anhydrotetracycline (aTc). Bar chart on the right

showed the logarithm-transformed CFU count of the *telE*-expressing cells in relative

to the control. Error bars represent mean + standard deviation (n = 3).

Figure 4. TelE is a pore-forming toxin. (A) The amino acid sequence of TelE was

color coded to indicate the predicted secondary structure. (B) Differential interference

contrast (DIC) micrograph of *E. coli* cells harboring an empty vector or a vector

encoding *telE* with the expression inducible by anhydrotetracycline (aTc). The

physiological changes of the *E. coli* cells following protein induction were documented

at the indicated time. Cells with membrane lesions were stained red by propidium

iodide. Arrows indicate the region with visible cell shrinkage. Scale bar, 2 μm. (C)

Phase contrast micrograph of *telE*-expressing *E. coli* documented at the indicated time.

Arrows indicate the region with visible cell shrinkage. Scale bar, 2 μm. (D) Total

proteins collected from the respective cellular fractions were resolved by SDS-PAGE

and stained with Coomassie blue (upper), or transferred onto a PVDF membrane for

anti-His immunoblot (lower). The two most intense proteins bands in the outer

membrane fractions are the outer membrane proteins OmpF/C (~37 kDa) and OmpA

(~35 kDa) serving as the control to approximate the purity of the soluble and inner

membrane fractions. T, total cellular lysate; L, cell-free lysate; S, soluble proteins; IM,

sarkosyl-souble inner membrane proteins; OM, sarkosyl-insoluble outer membrane

proteins.

Figure 5. The cognate immunity protein TipE antagonizes TelE toxicity by

preventing TelE membrane insertion. (A) Viability of *E. coli* cells harboring two

vectors for single expression of *telE* inducible by anhydrotetracycline (aTc), or co-

expression of *telE* and the indicated gene candidates with the expression inducible by

arabinose. Bar chart on the right showed the logarithm transformed CFU count of the

cells co-expressing two genes in relative to the control. Error bars represent mean +

standard deviation (n = 3). (B) Total proteins collected from the respective cellular

fractions were resolved by SDS-PAGE and stained with Coomassie blue (upper) or

transferred onto a PVDF membrane for anti-HA immunoblot (lower). The two most

intense protein bands in the outer membrane fractions are the outer membrane

proteins OmpF/C (~37 kDa) and OmpA (~35 kDa) serving as the control to

approximate the purity of the soluble and inner membrane fractions. OmpF/C (~37

kDa) also found in the periplasm was used as the positive control for the periplasmic

extract. T, total cellular lysate; L, cell-free lysate; S, soluble proteins; IM, sarkosyl-

souble inner membrane proteins; OM, sarkosyl-insoluble outer membrane proteins; P,

periplasmic proteins. (C) Immunoblot with anti-His or anti-HA on the proteins eluted

from the TelE-bound nickel-immobilized resins incubated with cellular lysate

containing HA-TipE. The TelE-free resins subjected to the same treatment were used

as the control. (D) Fluorescence micrograph of *E. coli* cells expressing superfolder

GFP (sfGFP), TelE-sfGFP, or co-expressing TipE and TelE-sfGFP. Cells with

membrane lesions were stained red by propidium iodide. Scale bar 10 µm; 2 µm for

inlets. Bar chart (bottom) showed the quantification of cells fluoresced green or red (n

≈ 300).

Figure 6. Purified TelE readily permeabilize *E. coli* pks+ cells when added in the

external milieu. (A) Flow-cytometry analysis of the CFSE-stained *E. coli* cells showing

the fluorescence intensity of the *E. coli* cell population treated with various detergents

or purified TelE protein for 10 minutes at 37°C. Upper panel, *E. coli* cells treated with

n-Octyl-β-D-Glucopyranoside (OG) at the concentration of 0.7% (present in TelE

elution buffer), 2.1% and 8.4%. Middle panel, *E. coli* cells treated with TelE depleted

of OG either using β-cyclodextrin (TelE1) or BioBeads (TelE2 and TelE3). Lower panel,

E. coli cells treated with pore-forming detergent TritonX100 and Tween20 at 0.05% as

positive control. (B) Representation of the flow-cytometry results as median

fluorescence ± CV. (C) Fluorescence images of live CFSE-stained *E. coli* cells treated

with TelE1, TelE2 and TelE3. Images were acquired with Zeiss microscope fitted with

100X objective lens. Scale bar 2.5 µm.

Figure S1. Phylogenetic tree of the clinical isolates sequenced in this study.

Phylogenetic inference of the isolates was performed using approximate-maximum-

likelihood algorithm based on the core-genome alignment using Parsnp version 1.2.

Scale bar represents substitution per variant site within the core genome. The isolates

sequenced in this study were labeled in black font, whereas the reference strains *Sgp*

(NCTC13784, ATCC43144 and WUSP067) and *Sgg* (DSM16831, ATCC43143,

BAA2069 and UCN34) with complete genomes available on NCBI GenBank, and a

draft genome of *Sgm* (strain ACA-DC-206) were labeled in blue font. Genomes used

for comparative genomics analysis as shown in Figure 1A were highlighted with darker

color shades. Bootstrap value for all the major branches ranges from 0.97 to 1.

Genomes labelled with asterisk contain a different T7SSb gene locus arrangement

than the UCN34 reference strain.

Figure S2. Antibacterial activity of *Sgg* T7SSb effectors. (A) Shortlisted proteins

that were significantly more abundant in the secretome of the UCN34 wildtype than

the Δ*essC* mutant. The total peptide count of each protein was expressed as means

± standard deviation from three biological replicates. N.D., not detected; N. A., not

applicable. (B) Viability of *E. coli* cells harboring an empty vector or a vector encoding

Sgg T7SSb effectors with the expression inducible by anhydrotetracycline (aTc). Bar

chart on the right showed the logarithm-transformed CFU count of the cells expressing

the T7SSb effectors in relative to the control. Error bars represent mean + standard

deviation (n = 3). (C) Viability of *Sgm* ACA-DC-206 and *E. faecalis* OG1RF cells

harboring an empty vector or a vector encoding *telC1* and *telC2* with the expression

inducible by anhydrotetracycline (aTc). Bar chart on the right showed the logarithm-

transformed CFU count of the cells expressing *telC1* or *telC2* in relative to the control.

Error bars represent mean + standard deviation (n = 3).

Figure S3. The prevalence of TelE homologs among *Sgg* isolates. The

phylogenetic inference of the *Sgg* isolates was shown on the left. Genomes in black

font represent the clinical isolates sequenced in this study, whereas the genomes in

blue font represent the complete genomes publicly available. Scale bar represents

substitution per variant site within the core genome. Table on the right summarized

the distribution of TelE homologs among *Sgg* isolates. Based on the C-terminal

sequence (shown in Figure 3B), the identified TelE homologs were broadly

categorized into 7 TelE variants and named as TelE1 to TelE7, where TelE1 refers to

the reference TelE from strain UCN34. Bootstrap values of all the major branches are

1.

Figure S4. Multiple sequence alignment of various TelE subtypes. Amino acid

sequences of TelE1 to TelE7 subtypes were aligned using online MAFFT server

version 7 with E-INS-i iterative refinement method that assumes all the sequences

shared the same conserved motifs. The aligned sequences were viewed using MView

(Brown et al., 1998) available on EMBL-EBI.

Figure S5. Impact of epitope tags or protein fusion on TelE activity. (A) Viability

of *E. coli* cells harboring an empty vector or a vector encoding *telE* fused with epitope

tags or superfolder GFP with the expression inducible by anhydrotetracycline (aTc).

Bar chart on the right showed the logarithm-transformed CFU count of the cells

expressing the *telE* variants in relative to the control. Error bars represent mean +

standard deviation (n ≥ 3). (B) Growth kinetics of the *E. coli* BL21(DE3) harboring

either the empty vector or a vector for IPTG-inducible expression of *telE*-His or co-

expression of *telE*-His and HA-*tipE.* Arrow indicates the time point when the gene

expression was induced. (C) Total protein collected from the respective lysates was

resolved by SDS-PAGE and stained with Coomassie blue. I, insoluble lysate; S,

soluble lysate. Arrow indicates the protein band corresponding to TelE and TipE,

respectively.

References

Abdallah, A. M., Gey Van Pittius, N. C., Champion, P. A., Cox, J., Luirink, J., Vandenbroucke-Grauls, C. M., Appelmelk, B. J. & Bitter, W. 2007. Type VII secretion--mycobacteria show the way. *Nat Rev Microbiol,* 5**,** 883-91.

Alikhan, N. F., Petty, N. K., Ben Zakour, N. L. & Beatson, S. A. 2011. BLAST Ring Image Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics,* 12**,** 402.

Anderson, M., Aly, K. A., Chen, Y. H. & Missiakas, D. 2013. Secretion of atypical protein substrates by the ESAT-6 secretion system of Staphylococcus aureus. *Mol Microbiol,* 90**,** 734-43.

Anderson, M., Chen, Y. H., Butler, E. K. & Missiakas, D. M. 2011. EsaD, a secretion factor for the Ess pathway in Staphylococcus aureus. *J Bacteriol,* 193**,** 1583-9.

Anderson, M., Ohr, R. J., Aly, K. A., Nocadello, S., Kim, H. K., Schneewind, C. E., Schneewind, O. & Missiakas, D. 2017. EssE Promotes Staphylococcus aureus ESS-Dependent Protein Secretion To Modify Host Immune Responses during Infection. *J Bacteriol,* 199.

Arthur, J. C., Perez-Chanona, E., Muhlbauer, M., Tomkovich, S., Uronis, J. M., Fan, T. J., Campbell, B. J., Abujamel, T., Dogan, B., Rogers, A. B., Rhodes, J. M., Stintzi, A., Simpson, K. W., Hansen, J. J., Keku, T. O., Fodor, A. A. & Jobin, C. 2012. Intestinal inflammation targets cancer-inducing activity of the microbiota. *Science,* 338**,** 120-3.

Aymeric, L., Donnadieu, F., Mulet, C., Du Merle, L., Nigro, G., Saffarian, A., Berard, M., Poyart, C., Robine, S., Regnault, B., Trieu-Cuot, P., Sansonetti, P. J. & Dramsi, S. 2018. Colorectal cancer specific conditions promote Streptococcus gallolyticus gut colonization. *Proc Natl Acad Sci U S A,* 115**,** E283-E291.

Aziz, R. K., Bartels, D., Best, A. A., Dejongh, M., Disz, T., Edwards, R. A., Formsma, K., Gerdes, S., Glass, E. M., Kubal, M., Meyer, F., Olsen, G. J., Olson, R., Osterman, A. L., Overbeek, R. A., Mcneil, L. K., Paarmann, D., Paczian, T., Parrello, B., Pusch, G. D., Reich, C., Stevens, R., Vassieva, O., Vonstein, V., Wilke, A. & Zagnitko, O. 2008. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics,* 9**,** 75.

Bae, T., Kozlowicz, B. & Dunny, G. M. 2002. Two targets in pCF10 DNA for PrgX binding: their role in production of Qa and prgX mRNA and in regulation of pheromone-inducible conjugation. *J Mol Biol,* 315**,** 995-1007.

Bertelli, C., Laird, M. R., Williams, K. P., Simon Fraser University Research Computing, G., Lau, B. Y., Hoad, G., Winsor, G. L. & Brinkman, F. S. L. 2017. IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets. *Nucleic Acids Res,* 45**,** W30-W35.

Biswas, I., Gruss, A., Ehrlich, S. D. & Maguin, E. 1993. High-efficiency gene inactivation and replacement system for gram-positive bacteria. *J Bacteriol,* 175**,** 3628-35.

Blighe, K., S Rana, And M Lewis 2018. EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling.

Boleij, A., Van Gelder, M. M. H. J., Swinkels, D. W. & Tjalsma, H. 2011. Clinical Importance of Streptococcus gallolyticus Infection Among Colorectal Cancer Patients: Systematic Review and Meta-analysis. *Clinical Infectious Diseases,* 53**,** 870-878.

Brettin, T., Davis, J. J., Disz, T., Edwards, R. A., Gerdes, S., Olsen, G. J., Olson, R., Overbeek, R., Parrello, B., Pusch, G. D., Shukla, M., Thomason, J. A., 3rd, Stevens, R., Vonstein, V., Wattam, A. R. & Xia, F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep,* 5**,** 8365.

Brown, N. P., Leroy, C. & Sander, C. 1998. MView: a web-compatible database search or
multiple alignment viewer. *Bioinformatics,* 14**,** 380-1.
Burts, M. L., Dedent, A. C. & Missiakas, D. M. 2008. EsaC substrate for the ESAT-6 secretion
pathway and its role in persistent infections of Staphylococcus aureus. *Mol Microbiol,*
69**,** 736-46.
Burts, M. L., Williams, W. A., Debord, K. & Missiakas, D. M. 2005. EsxA and EsxB are secreted
by an ESAT-6-like system that is required for the pathogenesis of Staphylococcus
aureus infections. *Proc Natl Acad Sci U S A,* 102**,** 1169-74.
Cao, Z., Casabona, M. G., Kneuper, H., Chalmers, J. D. & Palmer, T. 2016. The type VII secretion
system of Staphylococcus aureus secretes a nuclease toxin that targets competitor
bacteria. *Nat Microbiol,* 2**,** 16183.
Chen, W. C., Lee, P. I., Lin, H. C., Chang, L. Y., Lee, T. F., Chen, J. M. & Hsueh, P. R. 2020.
Clustering of Streptococcus gallolyticus subspecies pasteurianus bacteremia and
meningitis in neonates. *J Microbiol Immunol Infect*.
Corredoira, J., Grau, I., Garcia-Rodriguez, J. F., Alonso-Garcia, P., Garcia-Pais, M. J., Rabunal,
R., Garcia-Garrote, F., Ardanuy, C., Coira, A., Lopez-Alvarez, M. J. & Pallares, R. 2015.
The clinical epidemiology and malignancies associated with Streptococcus bovis
biotypes in 506 cases of bloodstream infections. *J Infect,* 71**,** 317-25.
Corredoira, J. C., Alonso, M. P., Garcia-Pais, M. J., Rabunal, R., Garcia-Garrote, F., Lopez-Roses,
L., Lancho, A., Coira, A., Pita, J., Velasco, D., Lopez-Alvarez, M. J., Tjalsma, H. & Varela,
J. 2014. Is colonoscopy necessary in cases of infection by Streptococcus bovis biotype
II? *Eur J Clin Microbiol Infect Dis,* 33**,** 171-7.
Corredoira-Sanchez, J., Garcia-Garrote, F., Rabunal, R., Lopez-Roses, L., Garcia-Pais, M. J.,
Castro, E., Gonzalez-Soler, R., Coira, A., Pita, J., Lopez-Alvarez, M. J., Alonso, M. P. &
Varela, J. 2012. Association between bacteremia due to Streptococcus gallolyticus
subsp. gallolyticus (Streptococcus bovis I) and colorectal neoplasia: a case-control
study. *Clin Infect Dis,* 55**,** 491-6.
Cuevas-Ramos, G., Petit, C. R., Marcq, I., Boury, M., Oswald, E. & Nougayrede, J. P. 2010.
Escherichia coli induces DNA damage in vivo and triggers genomic instability in
mammalian cells. *Proc Natl Acad Sci U S A,* 107**,** 11537-42.
Dai, Y., Wang, Y., Liu, Q., Gao, Q., Lu, H., Meng, H., Qin, J., Hu, M. & Li, M. 2017. A Novel Esat-
6 Secretion System-Secreted Protein EsxX of Community-Associated Staphylococcus
aureus Lineage ST398 Contributes to Immune Evasion and Virulence. *Front Microbiol,*
8**,** 819.
Danne, C., Guerillot, R., Glaser, P., Trieu-Cuot, P. & Dramsi, S. 2013. Construction of isogenic
mutants in Streptococcus gallolyticus based on the development of new mobilizable
vectors. *Research in Microbiology,* 164**,** 973-978.
Dekker, J. P. & Lau, A. F. 2016. An Update on the Streptococcus bovis Group: Classification,
Identification, and Disease Associations. *J Clin Microbiol,* 54**,** 1694-1699.
Dunny, G. M., Lee, L. N. & Leblanc, D. J. 1991. Improved electroporation and cloning vector
system for gram-positive bacteria. *Appl Environ Microbiol,* 57**,** 1194-201.
Eng, J. K., Mccormack, A. L. & Yates, J. R. 1994. An approach to correlate tandem mass spectral
data of peptides with amino acid sequences in a protein database. *J Am Soc Mass
Spectrom,* 5**,** 976-89.
Flaugnatti, N. & Journet, L. 2017. Identification of Effectors: Precipitation of Supernatant
Material. *Methods Mol Biol,* 1615**,** 459-464.

Franco, A. T., Johnston, E., Krishna, U., Yamaoka, Y., Israel, D. A., Nagy, T. A., Wroblewski, L.
979   E., Piazuelo, M. B., Correa, P. & Peek, R. M., Jr. 2008. Regulation of gastric
980   carcinogenesis by Helicobacter pylori virulence factors. *Cancer Res,* 68**,** 379-87.
981   Fridman, C. M., Keppel, K., Gerlic, M., Bosis, E. & Salomon, D. 2020. A comparative genomics
982   methodology reveals a widespread family of membrane-disrupting T6SS effectors. *Nat*
983   *Commun,* 11**,** 1085.
984   Garcia-Bayona, L., Guo, M. S. & Laub, M. T. 2017. Contact-dependent killing by Caulobacter
985   crescentus via cell surface-associated, glycine zipper proteins. *Elife,* 6.
986   Georgalaki, M. D., Van Den Berghe, E., Kritikos, D., Devreese, B., Van Beeumen, J.,
987   Kalantzopoulos, G., De Vuyst, L. & Tsakalidou, E. 2002. Macedocin, a food-grade
988   lantibiotic produced by Streptococcus macedonicus ACA-DC 198. *Appl Environ*
989   *Microbiol,* 68**,** 5891-903.
990   Green, M. R., Sambrook, J. & Sambrook, J. 2012. *Molecular cloning : a laboratory manual,*
991   Cold Spring Harbor, N.Y., Cold Spring Harbor Laboratory Press.
992   Holberger, L. E., Garza-Sanchez, F., Lamoureux, J., Low, D. A. & Hayes, C. S. 2012. A novel
993   family of toxin/antitoxin proteins in Bacillus species. *FEBS Lett,* 586**,** 132-6.
994   Huber, W., Von Heydebreck, A., Sultmann, H., Poustka, A. & Vingron, M. 2002. Variance
995   stabilization applied to microarray data calibration and to the quantification of
996   differential expression. *Bioinformatics,* 18 Suppl 1**,** S96-104.
997   Inoue, H., Nojima, H. & Okayama, H. 1990. High-Efficiency Transformation of Escherichia-Coli
998   with Plasmids. *Gene,* 96**,** 23-28.
999   Jamet, A. & Nassif, X. 2015. New players in the toxin field: polymorphic toxin systems in
1000   bacteria. *mBio,* 6**,** e00285-15.
1001   Katoh, K., Rozewicki, J. & Yamada, K. D. 2019. MAFFT online service: multiple sequence
1002   alignment, interactive sequence choice and visualization. *Brief Bioinform,* 20**,** 1160-
1003   1166.
1004   Kim, S., Jeon, T. J., Oberai, A., Yang, D., Schmidt, J. J. & Bowie, J. U. 2005. Transmembrane
1005   glycine zippers: physiological and pathological roles in membrane proteins. *Proc Natl*
1006   *Acad Sci U S A,* 102**,** 14278-83.
1007   Kim, S. Y., Joo, S. I., Yi, J. & Kim, E. C. 2010. A case of Streptococcus gallolyticus subsp.
1008   gallolyticus infective endocarditis with colon cancer: identification by 16S ribosomal
1009   DNA sequencing. *Korean J Lab Med,* 30**,** 160-5.
1010   Klein, T. A., Pazos, M., Surette, M. G., Vollmer, W. & Whitney, J. C. 2018. Molecular Basis for
1011   Immunity Protein Recognition of a Type VII Secretion System Exported Antibacterial
1012   Toxin. *J Mol Biol,* 430**,** 4344-4358.
1013   Kuraku, S., Zmasek, C. M., Nishimura, O. & Katoh, K. 2013. aLeaves facilitates on-demand
1014   exploration of metazoan gene family trees on MAFFT sequence alignment server with
1015   enhanced interactivity. *Nucleic Acids Res,* 41**,** W22-8.
1016   Kwong, T. N. Y., Wang, X. S., Nakatsu, G., Chow, T. C., Tipoe, T., Dai, R. Z. W., Tsoi, K. K. K.,
1017   Wong, M. C. S., Tse, G., Chan, M. T. V., Chan, F. K. L., Ng, S. C., Wu, J. C. Y., Wu, W. K.
1018   K., Yu, J., Sung, J. J. Y. & Wong, S. H. 2018. Association Between Bacteremia From
1019   Specific Microbes and Subsequent Diagnosis of Colorectal Cancer. *Gastroenterology,*
1020   155**,** 383-+.
1021   Lacourse, K. D., Peterson, S. B., Kulasekara, H. D., Radey, M. C., Kim, J. & Mougous, J. D. 2018.
1022   Conditional toxicity and synergy drive diversity among antibacterial effectors. *Nat*
1023   *Microbiol,* 3**,** 440-446.

Lazar, C., Gatto, L., Ferro, M., Bruley, C. & Burger, T. 2016. Accounting for the Multiple Natures
of Missing Values in Label-Free Quantitative Proteomics Data Sets to Compare
Imputation Strategies. *J Proteome Res,* 15**,** 1116-25.

Maragkoudakis, P. A., Papadelli, M., Georgalaki, M., Panayotopoulou, E. G., Martinez-
Gonzalez, B., Mentis, A. F., Petraki, K., Sgouras, D. N. & Tsakalidou, E. 2009. In vitro
and in vivo safety evaluation of the bacteriocin producer Streptococcus macedonicus
ACA-DC 198. *Int J Food Microbiol,* 133**,** 141-7.

Mcclain, M. S., Iwamoto, H., Cao, P., Vinion-Dubiel, A. D., Li, Y., Szabo, G., Shao, Z. & Cover, T.
L. 2003. Essential role of a GXXXG motif for membrane channel formation by
Helicobacter pylori vacuolating toxin. *J Biol Chem,* 278**,** 12101-8.

Morrison, D. A., Guedon, E. & Renault, P. 2013. Competence for natural genetic
transformation in the Streptococcus bovis group streptococci S. infantarius and S.
macedonicus. *J Bacteriol,* 195**,** 2612-20.

Ohr, R. J., Anderson, M., Shi, M., Schneewind, O. & Missiakas, D. 2017. EssD, a Nuclease
Effector of the Staphylococcus aureus ESS Pathway. *J Bacteriol,* 199.

Onoyama, S., Ogata, R., Wada, A., Saito, M., Okada, K. & Harada, T. 2009. Neonatal bacterial
meningitis caused by Streptococcus gallolyticus subsp. pasteurianus. *J Med Microbiol,*
58**,** 1252-1254.

Overbeek, R., Olson, R., Pusch, G. D., Olsen, G. J., Davis, J. J., Disz, T., Edwards, R. A., Gerdes,
S., Parrello, B., Shukla, M., Vonstein, V., Wattam, A. R., Xia, F. & Stevens, R. 2014. The
SEED and the Rapid Annotation of microbial genomes using Subsystems Technology
(RAST). *Nucleic Acids Res,* 42**,** D206-14.

Pleguezuelos-Manzano, C., Puschhof, J., Rosendahl Huber, A., Van Hoeck, A., Wood, H. M.,
Nomburg, J., Gurjao, C., Manders, F., Dalmasso, G., Stege, P. B., Paganelli, F. L., Geurts,
M. H., Beumer, J., Mizutani, T., Miao, Y., Van Der Linden, R., Van Der Elst, S., Genomics
England Research, C., Garcia, K. C., Top, J., Willems, R. J. L., Giannakis, M., Bonnet, R.,
Quirke, P., Meyerson, M., Cuppen, E., Van Boxtel, R. & Clevers, H. 2020. Mutational
signature in colorectal cancer caused by genotoxic pks(+) E. coli. *Nature,* 580**,** 269-273.

Price, M. N., Dehal, P. S. & Arkin, A. P. 2010. FastTree 2--approximately maximum-likelihood
trees for large alignments. *PLoS One,* 5**,** e9490.

Quan, S., Hiniker, A., Collet, J. F. & Bardwell, J. C. 2013. Isolation of bacteria envelope proteins.
Methods Mol Biol, 966**,** 359-66.

Roy, A., Kucukural, A. & Zhang, Y. 2010. I-TASSER: a unified platform for automated protein
structure and function prediction. *Nat Protoc,* 5**,** 725-38.

Rusniok, C., Couve, E., Da Cunha, V., El Gana, R., Zidane, N., Bouchier, C., Poyart, C., Leclercq,
R., Trieu-Cuot, P. & Glaser, P. 2010. Genome sequence of Streptococcus gallolyticus:
insights into its adaptation to the bovine rumen and its ability to cause endocarditis. *J*
Bacteriol, 192**,** 2266-76.

Schlegel, L., Grimont, F., Ageron, E., Grimont, P. A. & Bouvet, A. 2003. Reappraisal of the
taxonomy of the Streptococcus bovis/Streptococcus equinus complex and related
species: description of Streptococcus gallolyticus subsp. gallolyticus subsp. nov., S.
gallolyticus subsp. macedonicus subsp. nov. and S. gallolyticus subsp. pasteurianus
subsp. nov. *Int J Syst Evol Microbiol,* 53**,** 631-45.

Shi, H., Westfall, C. S., Kao, J., Odermatt, P. D., Cesar, S., Anderson, S., Sievert, M., Moore, J.,
Gonzalez, C. G., Zhang, L., Elias, J. E., Chang, F., Huang, K. C. & Levin, P. A. 2020.
Starvation induces shrinkage of the bacterial cytoplasm. *bioRxiv,* 2020.12.06.413849.

Smith, A. W. & Iglewski, B. H. 1989. Transformation of Pseudomonas aeruginosa by
electroporation. *Nucleic Acids Res,* 17**,** 10509.
Sullivan, M. J., Petty, N. K. & Beatson, S. A. 2011. Easyfig: a genome comparison visualizer.
Bioinformatics, 27**,** 1009-10.
Sutterlin, H. A., Shi, H., May, K. L., Miguel, A., Khare, S., Huang, K. C. & Silhavy, T. J. 2016.
Disruption of lipid homeostasis in the Gram-negative cell envelope activates a novel
cell death pathway. *Proc Natl Acad Sci U S A,* 113**,** E1565-74.
Taylor, J. C., Gao, X., Xu, J., Holder, M., Petrosino, J., Kumar, R., Liu, W., Hook, M., Mackenzie,
C., Hillhouse, A., Brashear, W., Nunez, M. P. & Xu, Y. 2021. A type VII secretion system
of Streptococcus gallolyticus subsp. gallolyticus contributes to gut colonization and
the development of colon tumors. *PLoS Pathog,* 17**,** e1009182.
Teh, W. K., Dramsi, S., Tolker-Nielsen, T., Yang, L. & Givskov, M. 2019. Increased Intracellular
Cyclic di-AMP Levels Sensitize Streptococcus gallolyticus subsp. gallolyticus to Osmotic
Stress and Reduce Biofilm Formation and Adherence on Intestinal Cells. *J Bacteriol,*
201.
Telford, J. L., Ghiara, P., Dell'orco, M., Comanducci, M., Burroni, D., Bugnoli, M., Tecce, M. F.,
Censini, S., Covacci, A., Xiang, Z. & Et Al. 1994. Gene structure of the Helicobacter
pylori cytotoxin and evidence of its key role in gastric disease. *J Exp Med,* 179**,** 1653-
58.
Thomsen, M. C. & Nielsen, M. 2012. Seq2Logo: a method for construction and visualization
of amino acid binding motifs and sequence profiles including sequence weighting,
pseudo counts and two-sided representation of amino acid enrichment and depletion.
Nucleic Acids Res, 40**,** W281-7.
Treangen, T. J., Ondov, B. D., Koren, S. & Phillippy, A. M. 2014. The Harvest suite for rapid
core-genome alignment and visualization of thousands of intraspecific microbial
genomes. *Genome Biol,* 15**,** 524.
Ulhuq, F. R., Gomes, M. C., Duggan, G. M., Guo, M., Mendonca, C., Buchanan, G., Chalmers, J.
D., Cao, Z., Kneuper, H., Murdoch, S., Thomson, S., Strahl, H., Trost, M., Mostowy, S.
& Palmer, T. 2020. A membrane-depolarizing toxin substrate of the Staphylococcus
aureus type VII secretion system mediates intraspecies competition. *Proc Natl Acad*
Sci U S A, 117**,** 20836-20847.
Unnikrishnan, M., Constantinidou, C., Palmer, T. & Pallen, M. J. 2017. The Enigmatic Esx
Proteins: Looking Beyond Mycobacteria. *Trends Microbiol,* 25**,** 192-204.
Wen, S. & Moss, S. F. 2009. Helicobacter pylori virulence factors in gastric carcinogenesis.
Cancer Lett, 282**,** 1-8.
Wending, G. K., Metzger, P. P., Dozois, E. J., Chua, H. K. & Krishna, M. 2006. Unusual bacterial
infections and colorectal carcinoma - Streptococcus bovis and Clostridium septicum:
Report of three cases. *Diseases of the Colon & Rectum,* 49**,** 1223-1227.
Whitney, J. C., Peterson, S. B., Kim, J., Pazos, M., Verster, A. J., Radey, M. C., Kulasekara, H. D.,
Ching, M. Q., Bullen, N. P., Bryant, D., Goo, Y. A., Surette, M. G., Borenstein, E., Vollmer,
W. & Mougous, J. D. 2017. A broadly distributed toxin family mediates contact-
dependent antagonism between gram-positive bacteria. *Elife,* 6.
Yang, J., Yan, R., Roy, A., Xu, D., Poisson, J. & Zhang, Y. 2015. The I-TASSER Suite: protein
structure and function prediction. *Nat Methods,* 12**,** 7-8.
Yang, J. & Zhang, Y. 2015. I-TASSER server: new development for protein structure and
function predictions. *Nucleic Acids Res,* 43**,** W174-81.

Zhang, X., Smits, A. H., Van Tilburg, G. B., Ovaa, H., Huber, W. & Vermeulen, M. 2018.
Proteome-wide identification of ubiquitin interactions using UbIA-MS. *Nat Protoc,* 13**,**
530-550.
Zhang, Y., Li, S., Zhu, C. G., Guo, Y. L., Wu, N. Q., Xu, R. X., Dong, Q., Liu, G. & Li, J. J. 2015. Risk
Factors, Coronary Severity, Outcome and ABO Blood Group: A Large Chinese Han
Cohort Study. *Medicine (Baltimore),* 94**,** e1708.
1122

Figure 1

Figure 2

A

B

C

Figure 3

Figure 4

A

MKIKMSEVLSQKESLASSISSAKSQLESASRKLKSTADSDALQGNVKEAIDNKITNYQVPLLTNY
VNSLAVISQGYDNLISTFKETVSETSDSAIIDTDVLQQMVDKFDSPLEQLKTCSDEINKAIDEVAD
IVSLTKVTTADAISEFKSAKKVLTNTIKDMGIFNSTVFTSEAGDILEEQNTQLSSLSDLGSSSYTS
KKAKEFYKDSTFKAEVKEVKSVVNGKARRDQLKNALAKNLYDSKYSGYMLETSGLEVKAGILIT
STKKSIGKGKKVVKILDATDKALKQKKFIGGRELMVDSKGRVKVGERFLYKKPSKGKTKTVHLY
SKGTKEYKKTVGEDFSFKKTTLYNGRKLKANGKVGIDWKGVGTSGKEAFKSGLKDNFKSTINP
LNEFKGFKEASNFSKAGKVLGFAGTVMTIGSNIKTDFIDDRRSSTKEKVRNFAVDTTVDVVSSS
SAAAA**GAAVGTLVGGPVGTVVGAAVG**IGFSVAMNHRVKFLGGKSLTEVTKDSLKGVAGKFGL

glycine zipper motif

X coil   X sheet   X helix

B

	t = 5 min	t = 15 min	t = 30 min	t = 60 min
control				
telE				

C

telE

t = 0

t = 30 min

D

Coomassie blue

α-His

Figure 5

Figure 6