Title: Environmental determinants of round goby invasion refuges at a river scale:

implications for conservation of native biodiversity

Authors and affiliation

Olivier Morissette[1*], Cristina Charette[2,3], Matthew J.S. Windle[3], Abraham Francis[4], Annick

Drouin[5], Jesica Goldsmit[5] and Alison M. Derry[2]

1. Chaire de recherche sur les espèces aquatiques exploitées, Université du Québec à

Chicoutimi, Chicoutimi, QC, Canada

2. Département des sciences biologiques, Université du Québec à Montréal, Montréal,

QC, Canada

3. St. Lawrence River Institute of Environmental Sciences, Cornwall, ON, Canada

4. Environment Program, Mohawk Council of Akwesasne, Akwesasne, QC, Canada

5. Direction de l'expertise sur la faune aquatiques, Ministère des Forêts, de la Faune et

des Parcs, Québec, QC, Canada

* Contact and corresponding author

Keyword: round goby, biological invasion, risk management, wetlands, biodiversity

conservation, environmental heterogeneity

Abstract

Introductions of exotic invasive species are a global disturbance for natural habitats. The severity of invasions can greatly vary from local to global scales, as observed in invasion refuges, exhibiting lower-than-expected invasion intensity. In this study, we analyzed the effects of water conductivity and wetland presence on the density of the round goby (*Neogobius melanostomus*) in a large-scale study (> 1300 sites), spanning a 400 km stretch of the St. Lawrence River (Canada). Our results showed that round goby density was null in sites with water conductivity under 100 µS/cm and increased toward a probable biological optimum at 300 µS/cm. The presence of wetlands appeared to also decrease round goby density along the conductivity continuum. Similarly, fish community diversity was maximal outside of the round goby water conductivity optimum. Hence, low water conductivity (<100 µS/cm), in interaction with the presence of wetlands, can provide a refuge for native aquatic species, establishing a simple risk assessment tool for managers. Our results also highlighted the high value of wetland conservation for conservation of native species biodiversity.

Introduction

Biological invasions are devastating, global anthropogenic disturbances that threaten biodiversity, and disrupt ecosystem functions and services (Bacher et al. 2017; IPBES 2019; Lewis et al. 2016; Pejchar and Mooney 2009; Simberloff et al. 2013). Distressingly, the number of non-native species introductions continues to increase around the world (Seebens et al. 2017; Seebens et al. 2018). The magnitude and frequency of invasions, however, can be highly variable (Dawson et al. 2017). From the global to regional scale, some locations are experiencing a disproportionate invasion pressure and impacts while other neighbouring habitats seem to be far less impacted (Dawson et al. 2017; Dyer et al. 2017; Latzka et al. 2016). This phenomenon is crystallized in the identification of invasion hotspots and refuges, exhibiting contrasting invasion prevalence (number and abundance of invasive species) (Boon et al. 2023; Dawson et al. 2017). Disproportionate levels of invasion across an entire landscape can often be explained by differences in propagule pressure or ecosystem characteristics that mediate invasion success (Catford et al. 2009; Lockwood et al. 2009). However, within invaded ecosystems, other factors that determine whether local sites become invasion hotspots or invasion refuges remain to be better described and understood. Importantly, it has been suggested that spatial environmental heterogeneity (or complexity) and social context (i.e. conservation zones or Indigeneous Peoples presence) within invaded ecosystems seems to play an important role—in conjunction with climate—to promote or suppress invasive species density at local and regional scale and their associated impacts on native species (Melbourne et al. 2007; Schuster et al. 2019; Vander Zanden et al. 2017). Identification and delineation of invasion hotspots and refuges are therefore an effective tool for integrated, multidisciplinary planning of invasive species management and conservation of native biodiversity (Byers et al. 2002).

Invasion refuges could arise within invaded regions or habitats where ecological, social or

environmental conditions create the deterrence of invaders, either by decreasing the

propagule pressure or the survival or performance of invaders (Anton et al. 2014; Boon et al.

2023; Melbourne et al. 2007; Vander Zanden et al. 2017). The importance of invasion refuges

can be crucial as even a small to moderate decrease of invasion pressure or invasive species

density can allow coexistence of invasive and native species in neighbouring habitats, hence

promoting native species persistence (Bradley et al. 2019; DeRoy et al. 2020; Sofaer et al.

2018; Strayer 2020; Strayer et al. 2018; Yokomizo et al. 2009). The existence of invasive

refuges is in accordance with the environmental matching hypothesis, stating that the impacts

of an invader decrease with environmental conditions that diverge from its physiological

optimum (Iacarella and Ricciardi 2015; Ricciardi et al. 2013). Hence, a growing body of

literature has explored environmental factors that promote invasion refuges within invaded

freshwater ecosystems (Astorg et al. 2020; Boddy and McIntosh 2021; Boddy et al. 2020;

Grabowski et al. 2009; Iacarella and Ricciardi 2015; Kestrup and Ricciardi 2009; Latzka et

al. 2016; Priddis et al. 2009). However, many of those studies have not identified the broad

role of environmental heterogeneity in the establishment of invasion refuges, mainly because

of their limited spatial scales or the context of laboratory experiments (Astorg et al. 2020;

Iacarella and Ricciardi 2015; Kestrup and Ricciardi 2009), potentially limiting inferences at

the whole ecosystem scale.

The round goby is an invasive littoral benthic fish that was first observed in the St. Lawrence

River in 1997 where it became increasingly abundant in many, but not all, habitats (Astorg et

al. 2022; Kipp et al. 2012; Morissette et al. 2018). Its geographic range distribution in the St.

Lawrence River spans from Lake Ontario to the salt front of the St. Lawrence Estuary,

~30 km downstream of Québec City. When present, the round goby has strong density-

dependant ecological effects, notably by reducing the density of native benthic fishes such as

darters (*Etheostoma* spp.) and sculpins (*Cottus* spp.) that use similar habitats (Janssen and

Jude 2001; Jude et al. 2018; Morissette et al. 2018; Poos et al. 2009). The round goby can

also ravage the diversity and abundance of macroinvertebrate communities (Astorg et al.

2022; Kipp et al. 2012). However, some types of habitats and environmental conditions

appear to reduce round goby abundance. In the Great Lakes, low oxygen concentration and

low abundance of wetlands or riparian vegetation were linked to invasion success of the

round goby in some tributaries (Kornis et al. 2013; Krabbenhoft and Kashian 2022;

McAllister et al. 2022). The concentration of dissolved ions (often measured by water

conductivity or specific conductance, which is conductivity standardized at 25 °C) seem to

also predict the presence and abundance of the exotic round goby in the Great Lakes

(Baldwin et al. 2011) and in the St. Lawrence River (Astorg et al. 2020). Another laboratory

experiment from a limited portion of the St. Lawrence River showed that the impact of

conductivity on round goby performance was influenced by water calcium concentration,

where low dissolved calcium ions (Ca^{2+}) resulted in low predation rates and, thus, reduced

ecological impacts (Iacarella and Ricciardi 2015). Other studies have found evidence to

support that wetlands, through their substrate properties and associated submerged aquatic

vegetation, offer buffering effects from negative round goby impacts on native biodiversity at

invaded sites (Astorg et al. 2020; Cooper et al. 2007). Those results, albeit highly pertinent

for risk assessment and management of the round goby, were based on observations from a

limited portion of the St. Lawrence River, impeding the potential for a generalization of the

results.

The objective of our study was to explore the relative influence of local environmental

conditions on the round goby density in a large-scale ecosystem. We used two extensive fish

surveys characterizing round goby density and littoral fish community diversity along the

entire non-tidal freshwater portion of the St. Lawrence River. Specifically, we aimed to test

the hypothesis that variation of water conductivity and presence of wetlands can influence

round goby density and native littoral fish diversity, establishing invasion refuges and

hotspots. We predicted that non-wetland invaded sites under high water conductivity act as

invasion hotspots by increasing round goby density and causing reduced littoral fish

diversity. Conversely, we predicted that wetland invaded sites under lower water conductivity

would act as invasion refuges from the round goby by reducing round goby density and

maintaining or increasing littoral fish diversity. Our study provides information for delimiting

round goby hotspots and refuges in the St. Lawrence River, helping to refine risk assessment

of Ponto-Caspian invasion in this ecosystem and others under imminent threat of round

gobies, as well as promoting the restoration and conservation of invasion refuge in the region.

Materials and Methods

Study area

We conducted our study over an area spanning approximately 400 km along the St. Lawrence

River, extending from Kingston (Ontario, Canada) at the outflow of Lake Ontario to Batiscan

(Québec, Canada) in the St. Lawrence River (Figure 1). The St. Lawrence River is a large,

fluvial ecosystem draining the Laurentian Great Lakes toward the Atlantic Ocean. Its narrow

fluvial sections successively (upstream to downstream) drain three widening of the St.

Lawrence River, called the fluvial lakes (St. François, St. Louis, and St. Pierre lakes). Where

the St. Lawrence reaches the Archipelago of Montréal, the water from the Ottawa River,

characterized by its low conductivity (~60 to 100 µS/cm) and brown coloration, joins the

Great Lakes water mass, which has a higher water conductivity (200 to 400 µS/cm) and green

coloration. Downstream of Montréal, both water masses flow side by side, with very little

lateral mixing (Frenette et al. 2003; Vis et al. 1998), until the St. Lawrence reaches its tidal

portion downstream of Lake St. Pierre. Hence, the water masses keep their contrasted

chemical properties for hundreds of km, creating a large ecotone where brown waters

(Ottawa River) are mostly restricted to the north shore and green waters (Great Lakes) to the

south shore. The relative position of the mixing zone and the two water masses stay stable,

with seasonal variations depending on the magnitude of spring floods or summer low water.

Along the large-scale chemical gradients, various habitats can be observed, from highly

vegetated wetlands to sandy and rocky bottom sections, creating an additional environmental

variation of interest for fish and invertebrate communities.

Environmental and fish data

Fish community data were acquired using beach seines as part of both the Fish Identification

Nearshore Survey (FINS) conducted by the River Institute and the Mohawk Council of

Akwesasne Environment Program, and the *réseau de suivi ichtyologique* (RSI) operated by

ministère des Forêts, de la Faune et des Parcs fish surveys. The sampling occurred in mid to

late summer over 14 years, between July 2007 to September 2021. The FINS fishing was

conducted using a 9.1 m long and 1.8 m deep purse seine, operated on three 20 m long hauls

for every studied site, corresponding approximately to 182 m^2 of surveyed habitat per effort.

The RSI used a 12.5 m long by 4 m deep beach seine deployed on the site and closed on itself

by operators, corresponding to approximately 120 m^2 of surveyed habitat per site. Both nets

were used in coastal, wadable (depth <1.5 m) habitats. Captured fish were identified to

species and counted by capture sites. As the spatiotemporal effect of round goby presence on

those littoral fish communities have already been assessed (Morissette et al. 2018), we

limited the study to the analysis of goby density and fish diversity. Site-specific round goby

counts were converted to density (no. gobies/m^2) to allow comparisons between both FINS

and RSI datasets.

Environmental data, conductivity (specific conductance), and wetland presence were either

measured and observed on the site during sampling (FINS) or estimated from geographical

data available (RSI). During FINS surveys, conductivity (µS/cm) was measured and recorded

nearshore (maximum 1-meter depth) at each sampling site using a calibrated water quality

multi-parameter sonde (Professional plus model YSI ProDSS). Conductivity estimation was

conducted using an independent 1660 sampling points acquired between 2017 and 2021

(MFFP, *unpublished*) spanning all the entire RSI (Québec) sampling area. Conductivity in the

RSI sampling zone was interpolated using this dataset and ordinary kriging analysis in the

Spatial Analyst toolbox implemented in the software ArcMap 10.7.1 (ESRI). Validation on

the interpolation performance was conducted by regression of interpolated and measured

conductivity values, showing a reliable agreement ($R^2_{adj} = 0.84$, Figure S1). Conductivity

values for RSI fish sampling sites were extracted from the interpolated layer using the

sampling coordinates and the function *extract* coded within the R package *terra* (Hijmans et

al. 2022) implemented in R version 4.1.2 (R Core Team 2021). For both FINS and RSI

surveys, presence of wetlands on sites was extracted using two open-access, high-resolution

cartography of Ontario (Ministry of Natural Resources and Forestry 2021) and Quebec

wetlands (Canards Illimités Canada and ministère de l'Environnement et Lutte contre les

changements climatiques 2021). To simplify and correct for wetland definition discrepancy in

the datasets, only "open water" class was considered as *non-wetlands* and all remaining

classes (e.g., marsh, swamps, wet meadows) were considered as *wetlands*, creating a binary

habitat type factor.

Statistical modelling

All statistical analyses were computed in R version 4.1.2 R (R Core Team 2021) and

RStudio v2022.02.3+492 (RStudio Team 2020). Generalized additive models (GAMs) were

used to assess the effect of conductivity and habitat type on round goby density using the

mgcv R package v1.8-38 (Wood 2011). GAM is a useful regression-based framework to

model non-linear relationships using smooth functions and can incorporate different family

error distributions (Wood 2017). GAMs allow for estimation of partial effects of a predictor

variable, which represent the weighted sum of multiple basic functions of increasing

wiggliness. Three models were built in relation to our hypothesis and predictions. The first

two models included round goby density (individual/m^2) as the response variable. The first

model included 4 predictors that were habitat type as a parametric effect, conductivity by

habitat factor as a smooth interaction, longitude and latitude as smooth interaction, and year

as a random smooth. The second model included all listed predictors, but the conductivity

smooth was considered without the interaction with habitat types and its parametric effect

(Table S1). For both models, latitude and longitude smooth interaction was included to

consider any spatial correlation and to ensure that model residuals are more likely to be

independent. Year was identified as a random predictor to generalize the effect over the

sampling years and to account for any effect of time since invasion (Astorg et al. 2022). The

default thin plate spline was used for the conductivity smooth or for the smooth by factor

interaction, the Duchon spline was applied to the latitude and longitude smooth interaction,

and a random smooth intercept was used for years. Since the response variable was

continuous with left-skewed, non-integer values including zeros, an identity log link function

with a Tweedie error distribution was applied to both round goby models.

Model selection was performed on the first two models using the Akaike's Information

Criterion (AIC) score to identify the best-fitted model with the lowest AIC score (Wood et

al., 2017) (Table S1). The third GAM included littoral native fish diversity as a response

variable. For each site, Shannon diversity index (H) using Hellinger transformed species

abundance, without round goby count, was calculated for all sampling sites. The model

predictors were also habitat type as a parametric effect, conductivity by habitat factor-smooth

interaction, longitude and latitude smooth interaction, and year as a random smooth. The

predictors' splines were identical to preceding models; however, a Gaussian error distribution

was applied since response variable (H) was not skewed and normally distributed. Model

diagnostics and maximum basis functions (k) were verified using the *gam.check* function and

autocorrelation was assessed using the *acf* function of the model residuals from the *stats* R

package. No significant autocorrelation was detected in the GAMs and diagnostic plots

suggested normality and homoscedasticity of the residuals (Figure S2). (Wood et al. 2017).

Model visualization was accomplished using the *mgcViz* package v0.1.9 (Fasiolo et al. 2019)

and *ggplot2* package v3.3.5 (Wickham et al. 2016).

Results

The fish dataset comprised 1310 sampling sites, with an average density of 0.07 goby/m^2

($\pm$0.17 SD), ranging from 0 to 1.8 goby/m^2. Conductivity was variable in the study area,

ranging from 43 to 576 µS/cm, with an average value of 245 µS/cm ($\pm$70 µS/cm SD). The

expected spatial structure in water conductivity structure between north shore and south shore

of the Upper St. Lawrence River downstream of Montréal was observed (Figure 1), where

north shore has a lower conductivity while in the south shore is higher. Measured

conductivity in the Upper St. Lawrence was more homogeneous, where variations were

mainly influenced by small tributary outflows. Wetlands and non-wetlands were equally

represented in the study area, with 662 wetland sites and 648 non-wetland sites in the dataset,

with both habitats present in approximately equivalent amounts on both shores (Figure 1).

According to our predictions, round goby density showed low abundance from sites of water

low conductivity and a sharp increase in median values of water conductivity along this

gradient (Figure 2).

Our results indicate that both habitat type (wetland/non-wetland) and conductivity were

important predictors of round goby density and littoral fish diversity in the St. Lawrence

River. The best-fitted GAM model showed the lowest AIC and included habitat type as a

fixed effect, conductivity by habitat type factor-smooth interaction along with the latitude and

longitude smooth interaction, and year as a random smooth (Table S1 and S2, Figure S2 and

S3). A statistical summary of the selected model (Table 1) showed both the fixed parametric

effect and partial smooth effects were significant (p-value <0.05) and explained 55.8% of the

null deviance with an R^2-adjusted value of 30.5%. Similarly, the fixed parametric effect of

wetlands had a significant (p-value <0.05) on littoral fish diversity, with higher diversity in

wetland sites. The GAM explained 21.4% of the null deviance with an R^2-adjusted value of

18.4%, but the smooth interaction between conductivity and wetlands did not produced a

significant effect, only showing a trend of non-wetlands interaction (p = 0.07).

Habitat type showed a direct effect on round goby density and littoral fish community

diversity. Wetlands generally hosted lower round goby density and preserved a higher fish

diversity compared to non-wetlands (Figure 3). Water conductivity showed a non-linear,

partial effect on the round goby density and diversity. However, the magnitude of the effect

was different between habitat types and was dependent on the range of the observed values.

Overall, from 40 µS/cm to approximately 220 µS/cm, conductivity showed a negative effect

on round goby density (Figure 3). This negative effect was found to be amplified in wetlands

compared to non-wetland habitats (Figure 3). Above this threshold, water conductivity

displayed a positive effect on round goby density in both habitats. At higher levels of

conductivity (> 380 µS/cm), a negative effect on round goby density was only observed in

wetlands. However, the confidence interval is large and includes the null effect. Overall, the

round goby exhibited a maximum density around 295 µS/cm, a potential biological optimum.

Accordingly, fish community diversity increased from 40 µS/cm to 190 µS/cm, but declined

as round goby density increased (220 to 300 µS/cm), to reach its minimum around the round

goby optimum (295 µS/cm). The confidence intervals were also large and included the null

effect.

The best-fitted GAM model demonstrated a significant partial effect of latitude and longitude

on round goby density (Figure 4), showing a spatially coherent weighted sum of models'

predictors on round goby density. A considerable portion of the south shore of the St.

Lawrence River displayed an important positive effect on round goby density while some

locations, notably the Ottawa River and Deux-Montagnes Lake (a widening of the Ottawa

River), had the most negative partial effect on round goby density. We consider that all the

acquired information could guide round goby risk assessment in a simple two-metrics fashion

(Table 2), considering both conductivity and wetland presence. Herein, we propose that non-

wetland habitats with water conductivity under 100 µS/cm have a very low risk of goby

impacts, evaluated as nearly null in wetlands under this conductivity. This risk increases with

increasing water conductivity until higher than 300 µS/cm, reaching the ecological optimum

where impacts are predicted to be maximal. Freshwater habitats over this threshold are

known to host round goby populations with substantial impacts (Baldwin et al. 2011; Kipp et

al. 2012; Kornis et al. 2013). However, under these conductivity conditions, wetlands still

have buffering effects, notably by preserving littoral fish community diversity.

Discussion

Our study aimed to examine the effects of two environmental conditions—water conductivity

and presence of wetlands—on the invasive round goby density and littoral fish community

diversity at a large-scale ecosystem, the St. Lawrence River. This study represented a test of

the hypothesis stating that those condition could establish large-scale and local invasion

refuges and hotspots. Using two unpublished and extensive fish survey datasets, we applied

GAMs to determine thresholds for ecological optimum along the water conductivity gradient

and showed large-scale effects of water conductivity on round goby density, as previously

recognized in its invaded range (Astorg et al. 2020; Baldwin et al. 2011). Our results revealed

that round goby density was positively influenced by increasing conductivity until reaching a

probable biological optimum of ~300 µS/cm (invasion hotspots). Sites of water conductivity

under 200 µS/cm had low density to complete absence of round gobies (<100 µS/cm), even

in regions hosting established populations for more than 10 years (Morissette et al. 2018).

These sites therefore represent possible invasion refuges, a hypothesis supported by the

higher diversity observed in those sites. Additionally, even if the presence of wetlands did not

completely disrupt the pattern of water conductivity effects on round goby density in invaded

habitats, wetlands appeared to decrease round goby density and preserve littoral fish

community diversity along the conductivity continuum, contributing to the invasion refuge

effect. Our study addressed the influence of those two environmental factors along an over

400 km stretch of the Upper St. Lawrence River, compiling more than 1300 capture sites on a

14-year-old-time series. Hence, our results provide an empirical criterion delineating round

goby invasion hotspots and refuges. The sheer size of this study underlines the possibility of

generalizing the phenomenon and emphasizes the importance of habitat diversity (i.e.,

wetland conservation and restoration) to buffer the effects of biological invasion.

The detection of a predictable response of round goby density in the Upper St. Lawrence

River suggests that the risk of potential impact, that we consider density-dependent, could be

identified by two simple metrics, water conductivity and wetlands presence, participating in

identification of sites likely to be more or less impacted by round goby presence. Hence,

wetlands and low conductivity could represents significant contribution to establishment of

ecological refuges from the round goby, even if these habitats characteristics can be limited

in spatiotemporal scale, depending on submerged vegetation abundance, density and

phenology (Janetski and Ruetz 2015; Seilheimer and Chow-Fraser 2007). The simple risk

grid that our study provides can offer a fast and an easy-to-implement tool for managers and

could be disseminated on a large landscape with relatively accessible data. Management of

aquatic invasive species is often confronted with the challenge of rapidly identifying the

present or upcoming threat of a given species which can cause a vast array of impacts (Lodge

et al. 2006). Identification of those impacts can largely vary in the invaded range and depart

from what has been identified in laboratory experiments and small-scale field studies. A

broader understanding of context-dependent potential density and their related impacts is

needed to implement an efficient management in heterogeneous landscapes and populate a

more refined impacts evaluation framework (Vander Zanden et al. 2017). Our finding that a

local-scale phenomenon, observed in 120 m^2 habitat patches, had a uniform response across a

large environment scale underscores the value of observational studies in invasion biology

and its unique potential to offer generalizable risk assessment tools.

The negative effect of low water conductivity on round goby density is concordant with field

observations made by Astorg et al. (2020) and Baldwin et al. (2011), and our findings add to

a known effect of low conductivity on Ponto-Caspian invaders (Kestrup and Ricciardi 2009;

Zhulidov et al. 2004). Reduced Ponto-Caspian invader density in waters with low

concentrations of dissolved ions are suggested to be linked with the evolutionary history of

these euryhaline species, which experienced repeated marine invasions—characterized by

extreme variations of salinity—favouring physiological processes requiring availability of

high-dissolved ion concentrations in ambient waters. Differences in round goby performance

along freshwater conductivity gradients may be linked to the concentration of calcium ions

(Ca^{2+}), which contribute to physiological processes and represent a likely common element

between freshwater and saltwater (Baldwin et al. 2011; Hunn 1985; Iacarella and Ricciardi

2015; Mayer et al. 1994). Calcium represents a critical component of homeostasis in fish, by

intervening on rates and efficiency of gill-water exchanges. Accordingly, field observations

suggest that the fish calcium uptake from the environment seems to represent an

evolutionarily conserved process that does not proportionally respond to large differences in

environmental calcium, even after long-term (multiple generations) residence along this

environmental gradient (Sanderson et al. 2023). Hence, the poor performance of round

gobies—along with other Ponto-Caspian invaders—can be a manifestation of the limited

plasticity of this trait under the unusual environmental conditions imposed by low

conductivity tributaries or water masses. This is an eloquent example of environmental

matching hypotheses and how the intolerance for low conductivity waters translates into

natural refuge to invasion (Ricciardi et al. 2013).

The mitigation of round goby presence and density in the St. Lawrence River by

environmental heterogeneity in water conductivity and local presence of wetlands could

represent an important factor for the resilience of native fishes and macroinvertebrates. The

resistance of wetlands to round goby invasion was previously identified within the St.

Lawrence and Great Lakes Basin (Astorg et al. 2020; Cooper et al. 2007), but the

mechanisms remain to be identified. Cooper et al. (2007) suggested that such a phenomenon

could be linked to time since invasion or lack of access to rocky and hard substrates, a

consideration also considered by Kornis et al. (2013). Our study, however, shows that round

goby invasions are severe at local sites with favourable environmental conditions and remain

absent in unfavourable sites, even if the establishment occurred 2 decades ago (Astorg et al.

2022; Morissette et al. 2018). This refutes the first hypothesis and supports Krabbenhoft &

Kashian's (2022) conclusions that sites showing riparian degradation may be more subject to

invasion and their impacts. Although our models did not consider local substrate (non-

available), rocky and boulder banks are marginal in the studied region (Sergy 2008),

suggesting that substrate may be of minor influence on observed density in and out of the

wetlands. Hence, we suggest that factors promoting invasion resilience of wetlands may be

linked to the impediment of movements caused by submerged vegetation, predation pressure

or different feeding opportunities for the round goby (e.g. dreissenids mussels presence or

macroinvertebrates); these are research avenues that remain to be tested.

Aside from its role buffering invasion impacts identified in our study, wetlands in freshwater

habitats are contributing to a vast array of ecological functions and services and are of great

importance for indigenous peoples living at their proximity (i.e., medicine cabinet and

sources of food). The sum of those ecosystem services and socioeconomics benefits

underlines the critical importance of wetlands in aquatic habitats. Sadly, wetlands have

experienced drastic declines, disappearing three times faster than forested areas (Convention

on wetlands 2021). Such declines, fuelled by anthropogenic modifications of freshwater

habitats, pollution and climate change (Johnson et al. 2010; Patenaude et al. 2015), can still

be contained and even reverted with the application of emergency plans of conservation

(Tickner et al. 2020). Based on our results, we argue that wetlands can also be viewed as a

pragmatic tool in the integrated management of water quality and prevention of biological

invasions (Astorg et al. 2020; Patenaude et al. 2015; Trepel 2010). Promotion of a diverse,

yet complex mosaic of heterogeneous habitats could potentially decrease invasion impacts

and offer refuge for native species, including the imperiled species susceptible to round goby

impacts, notably the eastern sand darter (*Ammocrypta pelludica*), the channel darter (*Percina*

copelandi) and the pugnose shiner (*Notropis anogenus*), thus favouring their coexistence and

persistence (Melbourne et al. 2007; Priddis et al. 2009). Additionnally, conservation of

wetlands along the St. Lawrence River have been important for many indigenous

communities, which may have established critical invasion refuges in their surrounding

(Schuster et al. 2019), however, our actual dataset cannot serve to test this hypothesis.

We recognize that factors for invasion success and ecological impacts of biological invasions

are complex and subject to multiple—and potentially synergistic—biotic and abiotic

influences (Catford et al. 2009; Ricciardi et al. 2021; Spear et al. 2021). Notably, over the

large study region, the influence of climatic variations (i.e., water temperature, amount of

precipitation, winter severity) could also contribute to the observed spatial effect. However,

contrasted round goby density observed in close sampling sites (i.e., in the vicinity of

Montréal and within Lake St. Pierre) suggest that climatic variations were probably of low

importance in the observed patterns that we report here. Round goby invasion success could

be influenced by other factors that were not addressed in our study, such as the importance of

propagule pressure, feeding opportunities (i.e., abundance of dreissenid mussels or

invertebrates), other environmental variables (i.e., sediment type and water flow), or

ecological relationships (i.e., predation). We also recognize that maximum density observed

in the St. Lawrence River (1.8 goby/m^2) remains lower than the maximum observed in the

Great Lakes, reaching at least 3.5 goby/m^2 (Taraborelli et al. 2009), which raises questions of

other underlying factors contributing to round goby density. Overall, however, the striking

effect of water conductivity on round goby density, and therefore ecological impact on native

fish and macroinvertebrate communities, offers an interesting avenue to study mechanisms

under invasion success and magnitude.

In conclusion, our study offered a unique opportunity to combine two ongoing and long-term

systematic fish surveys along a highly structured and complex freshwater ecosystem. We

observed a consistent effect of water chemistry and wetland presence in decreasing the

density of a notable Ponto-Caspian invader. Our results showed that low water conductivity

(<200 µS/cm), in interaction with the presence of wetlands, is linked to a lower round goby

density, buffering the potential ecological impacts of this invader and providing a refuge for

native aquatic species such as macroinvertebrates and fish. The combination of easily

measurable environmental metrics can provide a simple, yet robust risk assessment tool for

managers. Our results also highlight the high value of wetland conservation and restoration—

led by a diverse community of peoples practicing various uses—as a promoter of biodiversity

and an integrated management tool for conservation of native species, especially in invaded

ecosystems subjected to anthropogenic disturbances.

Acknowledgments

The authors are grateful for the field crew and partners who contributed to the FINS and RSI,

notably Chantal Côté, Yves Paradis, Kate Schwartz, and staff at the Mohawk Council of

Akwesasne's Environment Program. All fish collections were approved under the

Institutional Animal Protection Committee either at MELCCFP, UQAM, and the River

Institute and authorized by licences for scientific fish collections from the Ontario Ministry of

Natural Resources and Forestry and the Ministère de l'Environnement, de la Lutte contre les

changements climatiques, de la Faune et des Parcs du Québec.

Funding statement

This research was supported by the Chaire de recherche sur les espèces aquatiques exploitées

(MELCCFP and UQAC) to OM and NSERC Discovery Grants to AMD and OM. Funding to

MW and the FINS Project was provided by Fisheries and Oceans Canada's Habitat

Stewardship Program for Aquatic Species at Risk, the Ontario Ministry for Natural Resources

and Forestry's Species at Risk Stewardship Fund, WWF Canada's Loblaw Water Fund,

Ontario Power Generation, the Trottier Family Foundation, and the River Institute. Work of

CC was supported by scholarships provided by MITACS and Fonds du Québec – nature et

technologie (FRQNT).

Data availability statement

All raw data files and R scripts used in this work are freely available in the following

Borealis repository [Borealis link will be provided upon acceptance].

Authors' contributions

All authors contributed to the ideation of the study, the manuscript writing and the revision.

MW and CC contributed to the FINS survey and OM and AD contributed to the RSI survey.

MW, CC, and OM did the data curation and quality assessment, while CC and OM performed

the statistical analyses and produced figures and tables.

Conflict of interest statement

The authors declare no financial or personal conflict of interest.

References

Anton, A., Simpson, M.S., and Vu, I. 2014. Environmental and biotic correlates to lionfish invasion success in Bahamian coral reefs. PLoS One **9**(9): e106229. doi:10.1371/journal.pone.0106229.

Astorg, L., Charette, C., Windle, M.J.S., and Derry, A.M. 2022. Two decades since first invasion: Revisiting round goby impacts on nearshore aquatic communities in the Upper St. Lawrence River. Journal of Great Lakes Research **48**(2): 581-592. doi:10.1016/j.jglr.2022.01.017.

Astorg, L., Sanderson, S., Côté-Gravel, V., Sorbara, F., Windle, M.J.S., Hendry, A.P., and Derry, A.M. 2020. Different refuge types dampen exotic invasion and enhance diversity at the whole ecosystem scale in a heterogeneous river system. Biological Invasions **23**(2): 443-460. doi:10.1007/s10530-020-02374-7.

Bacher, S., Blackburn, T.M., Essl, F., Genovesi, P., Heikkilä, J., Jeschke, J.M., Jones, G., Keller, R., Kenis, M., Kueffer, C., Martinou, A.F., Nentwig, W., Pergl, J., Pyšek, P., Rabitsch, W., Richardson, D.M., Roy, H.E., Saul, W.C., Scalera, R., Vilà, M., Wilson, J.R.U., Kumschick, S., and Ramula, S. 2017. Socio-economic impact classification of alien taxa (SEICAT) [https://doi.org/10.1111/2041-210X.12844]. Methods in Ecology and Evolution **9**(1): 159-168. doi:10.1111/2041-210x.12844.

Baldwin, B.S., Carpenter, M., Rury, K., and Woodward, E. 2011. Low dissolved ions may limit secondary invasion of inland waters by exotic round gobies and dreissenid mussels in North America. Biological Invasions **14**(6): 1157-1175. doi:10.1007/s10530-011-0146-0.

Boddy, N.C., and McIntosh, A.R. 2021. Could spatial heterogeneity in flow disturbance drive temporal stability of native–invasive species co-occurrence in riverscapes? [https://doi.org/10.1111/fwb.13685]. Freshwater Biology **66**(5): 902-913. doi:10.1111/fwb.13685.

Boddy, N.C., Booker, D.J., and McIntosh, A.R. 2020. Heterogeneity in flow disturbance around river confluences influences spatial patterns in native and non-native species co-occurrence. Biological Invasions **22**(11): 3457-3475. doi:10.1007/s10530-020-02334-1.

Boon, J.S., Keith, S.A., Exton, D.A., and Field, R. 2023. The role of refuges in biological invasions: A systematic review. Global Ecology and Biogeography **n/a**(n/a). doi:https://doi.org/10.1111/geb.13701.

Bradley, B.A., Laginhas, B.B., Whitlock, R., Allen, J.M., Bates, A.E., Bernatchez, G., Diez, J.M., Early, R., Lenoir, J., Vila, M., and Sorte, C.J.B. 2019. Disentangling the abundance-impact relationship for invasive species. Proceedings of the National Academy of Sciences **116**(20): 9919-9924. doi:10.1073/pnas.1818081116.

Byers, J.E., Reichard, S.H., Randall, J.M., Parker, I.M., Smith, C.S., Lonsdale, W.M., Atkinson, I.A.E., Seastedt, T., Williamson, M.H., Chornesky, E.A., and Hayes, D.C.

2002. Directing research to reduce the impacts of nonindigenous species. Conservation Biology **16**(3): 630-640.

Canards Illimités Canada, and ministère de l'Environnement et Lutte contre les changements climatiques. 2021. Cartographie détaillée des milieux humides des secteurs habités du sud du Québec – Données du projet global.

Catford, J.A., Jansson, R., and Nilsson, C. 2009. Reducing redundancy in invasion ecology by integrating hypotheses into a single theoretical framework. Diversity and Distributions **15**(1): 22-40. doi:10.1111/j.1472-4642.2008.00521.x.

Convention on wetlands. 2021. Global wetlands outlook: Special edition 2021. Secretariat on the convention on wetlands, Gland, Switzerland.

Cooper, M.J., Ruetz, C.R., Uzarski, D.G., and Burton, T.M. 2007. Distribution of round gobies in coastal areas of lake Michigan: Are wetlands resistant to invasion? Journal of Great Lakes Research **33**(2): 303-313. Available from https://doi.org/10.3394/0380-1330(2007)33[303:DORGIC]2.0.CO;2 [accessed.

Dawson, W., Moser, D., van Kleunen, M., Kreft, H., Pergl, J., Pyšek, P., Weigelt, P., Winter, M., Lenzner, B., Blackburn, T.M., Dyer, E.E., Cassey, P., Scrivens, S.L., Economo, E.P., Guénard, B., Capinha, C., Seebens, H., García-Díaz, P., Nentwig, W., García-Berthou, E., Casal, C., Mandrak, N.E., Fuller, P., Meyer, C., and Essl, F. 2017. Global hotspots and correlates of alien species richness across taxonomic groups. Nature Ecology & Evolution **1**(7): 0186. doi:10.1038/s41559-017-0186.

DeRoy, E.M., Scott, R., Hussey, N.E., and MacIsaac, H.J. 2020. Density dependence mediates the ecological impact of an invasive fish [https://doi.org/10.1111/ddi.13063]. Diversity and Distributions **26**(7): 867-880. doi:10.1111/ddi.13063.

Dyer, E.E., Cassey, P., Redding, D.W., Collen, B., Franks, V., Gaston, K.J., Jones, K.E., Kark, S., Orme, C.D., and Blackburn, T.M. 2017. The Global Distribution and Drivers of Alien Bird Species Richness. PLoS Biology **15**(1): e2000942. doi:10.1371/journal.pbio.2000942.

Fasiolo, M., Nedellec, R., Goude, Y., and Wood, S.N. 2019. Scalable Visualization Methods for Modern Generalized Additive Models. Journal of Computational and Graphical Statistics **29**(1): 78-86. doi:10.1080/10618600.2019.1629942.

Frenette, J.-J., Arts, M.T., and Morin, J. 2003. Spectral gradients of downwelling light in a fluvial lake (Lake Saint-Pierre, St-Lawrence River). Aquatic Ecology **37**(1): 77-85. doi:10.1023/a:1022133530244.

Grabowski, M., Bacela, K., Konopacka, A., and Jazdzewski, K. 2009. Salinity-related distribution of alien amphipods in rivers provides refugia for native species. Biological Invasions **11**(9): 2107-2117. doi:10.1007/s10530-009-9502-8.

Hijmans, R.J., Bivand, R., Forner, K., Ooms, J., Pebesma, E., and Sumner, M.D. 2022. terra: Spatial Data Analysis R package.

Hunn, J.B. 1985. Role of calcium in gill function in freshwater fishes. Comparative
Biochemistry and Physiology Part A: Physiology **82**(3): 543-547. doi:10.1016/0300-
9629(85)90430-x.

Iacarella, J.C., and Ricciardi, A. 2015. Dissolved ions mediate body mass gain and predatory
response of an invasive fish. Biological Invasions **17**(11): 3237-3246.
doi:10.1007/s10530-015-0949-5.

IPBES. 2019. Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
IPBED Secretariat, Bonn, Germany.

Janetski, D.J., and Ruetz, C.R. 2015. Spatiotemporal patterns of fish community composition
in Great Lakes drowned river mouths. Ecology of Freshwater Fish **24**(4): 493-504.
doi:10.1111/eff.12161.

Janssen, J., and Jude, D.J. 2001. Recruitment failure of mottled sculpin *Cottus bairdi* in
Calumet Harbor, Southern Lake Michigan, induced by the newly introduced round
goby *Neogobius melanostomus*. Journal of Great Lakes Research **27**(3): 319-328.
doi:https://doi.org/10.1016/S0380-1330(01)70647-8.

Johnson, W.C., Werner, B., Guntenspergen, G.R., Voldseth, R.A., Millett, B., Naugle, D.E.,
Tulbure, M., Carroll, R.W.H., Tracy, J., and Olawsky, C. 2010. Prairie Wetland
Complexes as Landscape Functional Units in a Changing Climate. Bioscience **60**(2):
128-140. doi:10.1525/bio.2010.60.2.7.

Jude, D.J., Hensler, S.R., and Murray, M.M. 2018. Round goby and zebra mussel interactions
with darters in a warm-water stream community in southern Michigan, USA. Journal
of Freshwater Ecology **33**(1): 395-412. doi:10.1080/02705060.2018.1483270.

Kestrup, Å.M., and Ricciardi, A. 2009. Environmental heterogeneity limits the local
dominance of an invasive freshwater crustacean. Biological Invasions **11**(9): 2095-
2105. doi:10.1007/s10530-009-9490-8.

Kipp, R., Ricciardi, A., and Ramcharan, C.W. 2012. Impacts of the Eurasian round goby
(*Neogobius melanostomus*) on benthic communities in the upper St. Lawrence River.
Canadian Journal of Fisheries and Aquatic Sciences **69**(3): 469-486.
doi:10.1139/f2011-139.

Kornis, M.S., Sharma, S., Jake Vander Zanden, M., and Ricciardi, A. 2013. Invasion success
and impact of an invasive fish, round goby, in Great Lakes tributaries. Diversity and
Distributions **19**(2): 184-198. doi:10.1111/ddi.12001.

Krabbenhoft, C.A., and Kashian, D.R. 2022. Invasion success of a freshwater fish
corresponds to low dissolved oxygen and diminished riparian integrity. Biological
Invasions **24**(10): 3049-3063. doi:10.1007/s10530-022-02827-1.

Latzka, A.W., Hansen, G.J.A., Kornis, M., Vander Zanden, M.J., and Peters, D.P.C. 2016.
Spatial heterogeneity in invasive species impacts at the landscape scale
[https://doi.org/10.1002/ecs2.1311]. Ecosphere **7**(3): e01311. doi:10.1002/ecs2.1311.

Lewis, M.A., Petrovskii, S.V., and Potts, J.R. 2016. Dynamics of biological invasions. *In* The
Mathematics Behind Biological Invasions. Springer Cham, Switzerland. pp. 19-68.

Lockwood, J.L., Cassey, P., and Blackburn, T.M. 2009. The more you introduce the more
you get: the role of colonization pressure and propagule pressure in invasion ecology
[https://doi.org/10.1111/j.1472-4642.2009.00594.x]. Diversity and Distributions
15(5): 904-910. doi:10.1111/j.1472-4642.2009.00594.x.

Lodge, D.M., Williams, S., MacIsaac, H.J., Hayes, K.R., Leung, B., Reichard, S., Mack,
R.N., Moyle, P.B., Smith, M., Andow, D.A., Carlton, J.T., and McMichael, A. 2006.
Biological invasions: recommendations for U.S. policy and management. Ecological
Applications **16**(6): 2035-2054. doi:10.1890/1051-
0761(2006)016[2035:birfup]2.0.co;2.

Mayer, F.L.J., Marking, L.L., Bills, T.D., and Howe, G.E. 1994. Physicochemical factors
affecting toxicity in freshwater: hardness, pH and temperature. *In* Bioavailability:
physical, chemical and biological interactions. *Edited by* J.L. Hamelink and P.F.
Landrum and H.L. Bergman and W.H. Benson. Lewis Publishers, London. p. 256
pages.

McAllister, K., Drake, D.A.R., and Power, M. 2022. Round Goby (*Neogobius melanostomus*)
impacts on benthic fish communities in two tributaries of the Great Lakes. Biological
Invasions **24**(9): 2885-2903. doi:10.1007/s10530-022-02816-4.

Melbourne, B.A., Cornell, H.V., Davies, K.F., Dugaw, C.J., Elmendorf, S., Freestone, A.L.,
Hall, R.J., Harrison, S., Hastings, A., Holland, M., Holyoak, M., Lambrinos, J.,
Moore, K., and Yokomizo, H. 2007. Invasion in a heterogeneous world: resistance,
coexistence or hostile takeover? Ecology Letters **10**(1): 77-94. doi:10.1111/j.1461-
0248.2006.00987.x.

Ministry of Natural Resources and Forestry. 2021. Ontario land information (GeoHub).
Government of Ontario, Ottawa.

Morissette, O., Paradis, Y., Pouliot, R., and Lecomte, F. 2018. Spatio-temporal changes in
littoral fish community structure along the St. Lawrence River (Québec, Canada)
following round goby (Neogobius melanostomus) invasion. Aquatic Invasions **13**(4):
501-512. doi:10.3391/ai.2018.13.4.08.

Patenaude, T., Smith, A.C., and Fahrig, L. 2015. Disentangling the effects of wetland cover
and urban development on quality of remaining wetlands. Urban Ecosystems **18**(3):
663-684. doi:10.1007/s11252-015-0440-1.

Pejchar, L., and Mooney, H.A. 2009. Invasive species, ecosystem services and human well-
being. Trends in Ecology & Evolution **24**(9): 497-504.
doi:10.1016/j.tree.2009.03.016.

Poos, M., Dextrase, A.J., Schwalb, A.N., and Ackerman, J.D. 2009. Secondary invasion of
the round goby into high diversity Great Lakes tributaries and species at risk hotspots:
potential new concerns for endangered freshwater species [journal article]. Biological
Invasions **12**(5): 1269-1284. doi:10.1007/s10530-009-9545-x.

Priddis, E., Rader, R., Belk, M., Schaalje, B., and Merkley, S. 2009. Can separation along the
temperature niche axis promote coexistence between native and invasive species?
Diversity and Distributions **15**(4): 682-691. doi:10.1111/j.1472-4642.2009.00573.x.

R Core Team. 2021. R: A language and environment for statistical computing. R foundation
for statistical computing, Vienna, Austria.

Ricciardi, A., Hoopes, M.F., Marchetti, M.P., and Lockwood, J.L. 2013. Progress toward
understanding the ecological impacts of nonnative species
[https://doi.org/10.1890/13-0183.1]. Ecological Monographs **83**(3): 263-282.
doi:10.1890/13-0183.1.

Ricciardi, A., Iacarella, J.C., Aldridge, D.C., Blackburn, T.M., Carlton, J.T., Catford, J.A.,
Dick, J.T.A., Hulme, P.E., Jeschke, J.M., Liebhold, A.M., Lockwood, J.L., MacIsaac,
H.J., Meyerson, L.A., Pyšek, P., Richardson, D.M., Ruiz, G.M., Simberloff, D., Vilà,
M., and Wardle, D.A. 2021. Four priority areas to advance invasion science in the
face of rapid environmental change. Environmental Reviews **29**(2): 119-141.
doi:10.1139/er-2020-0088.

RStudio Team. 2020. RStudio: Integrated Development for R. RStudio. PBC, Boston, MA.

Sanderson, S., Astorg, L., Haines, G.E., Beaumont-Courteau, S., Langerhans, R.B., Derry,
A.M., and Hendry, A.P. 2023. Freshwater fishes maintain multi-trait phenotypic
stability across an environmental gradient in aqueous calcium
[https://doi.org/10.1111/jfb.15412]. Journal of Fish Biology **n/a**(n/a).
doi:https://doi.org/10.1111/jfb.15412.

Schuster, R., Germain, R.R., Bennett, J.R., Reo, N.J., and Arcese, P. 2019. Vertebrate
biodiversity on indigenous-managed lands in Australia, Brazil, and Canada equals that
in protected areas. Environmental Science & Policy **101**: 1-6.
doi:https://doi.org/10.1016/j.envsci.2019.07.002.

Seebens, H., Blackburn, T.M., Dyer, E.E., Genovesi, P., Hulme, P.E., Jeschke, J.M., Pagad,
S., Pysek, P., Winter, M., Arianoutsou, M., Bacher, S., Blasius, B., Brundu, G.,
Capinha, C., Celesti-Grapow, L., Dawson, W., Dullinger, S., Fuentes, N., Jager, H.,
Kartesz, J., Kenis, M., Kreft, H., Kuhn, I., Lenzner, B., Liebhold, A., Mosena, A.,
Moser, D., Nishino, M., Pearman, D., Pergl, J., Rabitsch, W., Rojas-Sandoval, J.,
Roques, A., Rorke, S., Rossinelli, S., Roy, H.E., Scalera, R., Schindler, S., Stajerova,
K., Tokarska-Guzik, B., van Kleunen, M., Walker, K., Weigelt, P., Yamanaka, T., and
Essl, F. 2017. No saturation in the accumulation of alien species worldwide. Nature
Communications **8**(1): 14435. doi:10.1038/ncomms14435.

Seebens, H., Blackburn, T.M., Dyer, E.E., Genovesi, P., Hulme, P.E., Jeschke, J.M., Pagad,
S., Pysek, P., van Kleunen, M., Winter, M., Ansong, M., Arianoutsou, M., Bacher, S.,
Blasius, B., Brockerhoff, E.G., Brundu, G., Capinha, C., Causton, C.E., Celesti-
Grapow, L., Dawson, W., Dullinger, S., Economo, E.P., Fuentes, N., Guenard, B.,
Jager, H., Kartesz, J., Kenis, M., Kuhn, I., Lenzner, B., Liebhold, A.M., Mosena, A.,
Moser, D., Nentwig, W., Nishino, M., Pearman, D., Pergl, J., Rabitsch, W., Rojas-
Sandoval, J., Roques, A., Rorke, S., Rossinelli, S., Roy, H.E., Scalera, R., Schindler,
S., Stajerova, K., Tokarska-Guzik, B., Walker, K., Ward, D.F., Yamanaka, T., and
Essl, F. 2018. Global rise in emerging alien species results from increased

accessibility of new source pools. Proceedings of the National Academy of Sciences
115(10): E2264-E2273. doi:10.1073/pnas.1719429115.

Seilheimer, T.S., and Chow-Fraser, P. 2007. Application of the wetland fish index to northern
Great Lakes marshes with emphasis on Georgian Bay coastal wetlands. Journal of
Great Lakes Research **33**(sp3): 154-171. doi:10.3394/0380-
1330(2007)33[154:Aotwfi]2.0.Co;2.

Sergy, G. 2008. The shoreline classification scheme for SCAT and oil spill response in
Canada. *In* Proceedings of the 31st Arctic and Marine Oil Spill Program Technical
Seminar. *Edited by* E. Canada, Ottawa, ON.

Simberloff, D., Martin, J.L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., Courchamp,
F., Galil, B., Garcia-Berthou, E., Pascal, M., Pysek, P., Sousa, R., Tabacchi, E., and
Vila, M. 2013. Impacts of biological invasions: what's what and the way forward.
Trends in Ecology & Evolution **28**(1): 58-66. doi:10.1016/j.tree.2012.07.013.

Sofaer, H.R., Jarnevich, C.S., and Pearse, I.S. 2018. The relationship between invader
abundance and impact [https://doi.org/10.1002/ecs2.2415]. Ecosphere **9**(9): e02415.
doi:10.1002/ecs2.2415.

Spear, M.J., Walsh, J.R., Ricciardi, A., and Zanden, M.J.V. 2021. The invasion ecology of
sleeper populations: prevalence, persistence, and abrupt shifts. Bioscience **71**(4): 357-
369. doi:10.1093/biosci/biaa168.

Strayer, D.L. 2020. Non-native species have multiple abundance-impact curves
[https://doi.org/10.1002/ece3.6364]. Ecology and Evolution **10**(13): 6833-6843.
doi:10.1002/ece3.6364.

Strayer, D.L., Solomon, C.T., Findlay, S.E.G., and Rosi, E.J. 2018. Long-term research
reveals multiple relationships between the abundance and impacts of a non-native
species [https://doi.org/10.1002/lno.11029]. Limnology and Oceanography **64**(S1):
S105-S117. doi:10.1002/lno.11029.

Taraborelli, A.C., Fox, M.G., Schaner, T., and Johnson, T.B. 2009. Density and habitat use
by the round goby (Apollonia melanostoma) in the Bay of Quinte, Lake Ontario.
Journal of Great Lakes Research **35**(2): 266-271.
doi:https://doi.org/10.1016/j.jglr.2008.12.004.

Tickner, D., Opperman, J.J., Abell, R., Acreman, M., Arthington, A.H., Bunn, S.E., Cooke,
S.J., Dalton, J., Darwall, W., Edwards, G., Harrison, I., Hughes, K., Jones, T.,
Leclere, D., Lynch, A.J., Leonard, P., McClain, M.E., Muruven, D., Olden, J.D.,
Ormerod, S.J., Robinson, J., Tharme, R.E., Thieme, M., Tockner, K., Wright, M., and
Young, L. 2020. Bending the curve of global freshwater biodiversity loss: An
emergency recovery plan. Bioscience **70**(4): 330-342. doi:10.1093/biosci/biaa002.

Trepel, M. 2010. Assessing the cost-effectiveness of the water purification function of
wetlands for environmental planning. Ecological Complexity **7**(3): 320-326.
doi:10.1016/j.ecocom.2010.02.006.

Vander Zanden, M.J., Hansen, G.J.A., and Latzka, A.W. 2017. A framework for evaluating
heterogeneity and landscape-level impacts of non-native aquatic species. Ecosystems
20(3): 477-491. doi:10.1007/s10021-016-0102-z.

Vis, C., Cattaneo, A., and Hudon, C. 1998. Periphyton in the clear and colored water masses
of the St. Lawrence River (Quebec, Canada): A 20-year overview. Journal of Great
Lakes Research **24**(1): 105-117. doi:10.1016/s0380-1330(98)70803-2.

Wickham, H., Navarro, D., and Pedersen, T.L. 2016. ggplot2: Elegant graphics for data
analysis. Use R! Springer, Houston, Texas, USA.

Wood, S.N. 2011. Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models
[https://doi.org/10.1111/j.1467-9868.2010.00749.x]. Journal of the Royal Statistical
Society: Series B (Statistical Methodology) **73**(1): 3-36. doi:10.1111/j.1467-
9868.2010.00749.x.

Wood, S.N. 2017. Generalized Additive Models. Chapman and Hall/CRC, New York.

Wood, S.N., Pya, N., and Säfken, B. 2017. Smoothing parameter and model selection for
general smooth models. Journal of the American Statistical Association **111**(516):
1548-1563. doi:10.1080/01621459.2016.1180986.

Yokomizo, H., Possingham, H.P., Thomas, M.B., and Buckley, Y.M. 2009. Managing the
impact of invasive species: the value of knowing the density-impact curve
[https://doi.org/10.1890/08-0442.1]. Ecological Applications **19**(2): 376-386.
doi:10.1890/08-0442.1.

Zhulidov, A.V., Pavlov, D.F., Nalepa, T.F., Scherbina, G.H., Zhulidov, D.A., and Gurtovaya,
T.Y. 2004. Relative distributions of Dreissena bugensis and Dreissena polymorpha in
the Lower Don River system, Russia. International Review of Hydrobiology **89**(3):
326-333. doi:10.1002/iroh.200310727.

Table 1. Statistical summary of the generalized additive model (GAM) selected including

habitat type as a fixed effect, conductivity by habitat type factor-smooth interaction,

longitude and latitude smooth interaction, and year as a random smooth. Edf indicates the

estimated degree of freedom and Ref.df represents the residual degree of freedom. Significant

parametric (A) and partial effects (B) are indicated by p-value <0.05.

A. parametric coefficients	Estimate	Standard Error	t-value	p-value
(Intercept)	-3.8625	0.1487	-25.97	<0.0001
Wetlands	-0.4896	0.1479	-3.31	<0.005
B. smooth terms	edf	Ref.df	F-value	p-value
s (conductivity): non-wetlands	7.198	8.941	4.127	<0.0001
s (conductivity): wetlands	4.15	5.246	6.764	<0.0001
s (longitude, latitude)	55.791	79	17.365	<0.0001
s (year)	9.85	14	3.003	<0.0001
R2-adjusted	0.305			
Deviance explained	55.8 %			

708   **Table 2.** Binary risk of round goby density (and density-dependent impacts) evaluation grid

709   based on specific conductance (conductivity) and local presence of wetlands on a local to

710   large-scale study site.

	Water specific conductance (conductivity)			
Local habitat	<100 µS/cm	100 to 200 µS/cm	200 to 300 µS/cm	> 300 µS/cm
Non-wetlands	very low	low	medium	high
Wetlands	none	very low	medium*	high*

* General agreement to large-scale risk with possibility of local or seasonal buffering effects

of wetlands

Figure 1. Round goby density (proportional red circles, black circles are null abundance) in

relationship to the water conductivity gradient (red to blue colour shade) presented on a

subset of the study for visualization (zone of maximum variation). The complete study area is

presented in the smaller context map with fish sampling sites locations (gray dots).

Figure 2. Round goby density (individual/m^2) by conductivity (μS/cm) for all studied

sampling sites. Only 250 to 349 μS/cm groups show a significant difference (ANOVA,

$F_{10,1297} = 16.73$, $p < 0.001$) in round goby density compared to others.

Figure 3. Generalized additive model (GAM) partial smooth effect of conductivity (μS/cm)

on round goby density (top, individual/m^2) and littoral fish community diversity (bottom,

Shannon H) by habitat types: non-wetlands (orange), wetlands (green) or all types (gray,

showing non-significant interaction). Shaded area indicates the 95% confidence interval.

Figure 4. Generalized additive model (GAM) of partial spatial smooth effect on round goby

density (individual/m^2) along the geographic gradient of the St. Lawrence River study area.

The range of colours represents the effect: yellow shows the most positive effect (higher

density) and dark blue the most negative effect (lower density). Partial residuals are

represented by dots.

Figure 1.

Figure 2.

Figure 3.

Figure 4.