Title: Environmental determinants of invasion hotspots and refugia at a river scale: implications

for conservation of native biodiversity

Authors and affiliation

Morissette, O.[1]*, Charette, C.[2,3], Windle, M.[3], Drouin, A.[4] and Derry, A.[2]

1. Chaire de recherche sur les espèces aquatiques exploitées, Université du Québec à

Chicoutimi, Chicoutimi, QC, Canada

2. Département des sciences biologiques, Université du Québec à Montréal, Montréal, QC,

Canada

3. St. Lawrence River Institute of Environmental Sciences, Cornwall, ON, Canada

4. Direction de l'expertise sur la faune aquatiques, Ministère des Forêts, de la Faune et des

Parcs, Québec, QC, Canada

* Contact and corresponding author

Keyword: round goby, biological invasion, risk management, wetlands, biodiversity

conservation, environmental heterogeneity

Abstract

Introductions of exotic invasive species are a global disturbance for natural habitats, native

species and their related ecological services. The severity of invasions and impacts, however, can

greatly vary from local to global scales. This variation can be observed in both invasion hotspots,

exhibiting disproportionate invasion and impacts, or refugia, exhibiting lower than expected

invasion and impacts. The latter are believed to be habitats where ecological, social or

environmental conditions create deterrence of invaders. Dissolved ions and presence of wetlands

have been suggested as environmental conditions that buffer the impacts of Ponto-Caspian

invaders, notably the round goby (*Neogobius melanostomus*). In this study, we used a

generalized additive model (GAM) to analyze the effects of local water conductivity and wetland

presence on the density of the round goby in a large-scale study (> 1300 sites), spanning the

major part of the non-tidal freshwater portion of the St. Lawrence River (Canada). Our results

showed that, even in long-invaded regions, round goby density was null in sites with water

conductivity under 100 µS/cm and increased toward a probable biological optimum of water

conductivity ~300 µS/cm. Additionally, the presence of wetlands, even if they did not

completely disrupt surpass the effect of conductivity, appeared to decrease the round goby

densities along the conductivity continuum. Our study demonstrated that low water conductivity

(<100 µS/cm), in interaction with the presence of wetlands can provide a refugium for native

aquatic species, establishing a simple, yet robust, risk assessment tool for managers. Our results

also highlighted the high value of wetland conservation and restoration as a promoter of

biodiversity and an integrated management tool for conservation of native species.

Introduction

Biological invasions are devastating global anthropogenic disturbances that threaten biodiversity, and disrupt ecosystem functions and services (Pejchar and Mooney 2009, Simberloff et al. 2013, Lewis et al. 2016, Bacher et al. 2018, IPBES 2019). Distressingly, the number of non-native species introductions continues to increase around the world (Seebens et al. 2017, Seebens et al. 2018). The magnitude and frequency of invasions, however, can be highly variable (Dawson et al. 2017). At the global to the regional scale, some locations are experiencing a disproportionate invasion pressure while other neighbouring habitats seem to have been far less impacted (Latzka et al. 2016, Dawson et al. 2017, Dyer et al. 2017). This phenomenon is crystallized in the identification of invasion hotspots and refugia, exhibiting contrasting invasion prevalence (number and abundance of invasive species) (Dawson et al. 2017). Disproportionate levels of invasion across the landscape can often be explained by differences in propagule pressure or ecosystem characteristics that mediate invasion success (Catford et al. 2009, Lockwood et al. 2009). However, within invaded ecosystems, other factors that determine whether local sites become invasion hotspots or invasion refugia remain to be better described and understood. For example, invaded ecosystems with homogeneous habitats can become invasion hotspots when environmental conditions are suitable and invasive species become abundant, often resulting in extirpation of native species (Melbourne et al. 2007). By contrast, spatial environmental heterogeneity within invaded ecosystems can sometimes play an important role in promoting or suppressing invasive species density at local and regional scales (Melbourne et al. 2007, Vander Zanden et al. 2017). The identification and delineation of invasion hotspots and refugia are therefore an effective tool for planning invasive species management and conservation of native biodiversity (Byers et al. 2002).

Invasion refugia could arise within invaded regions or habitats where ecological, social or

environmental conditions create a deterrence of invaders, either by preventing the propagule

pressure or decreasing survival or performance of invaders (Melbourne et al. 2007, Anton et al.

2014, Vander Zanden et al. 2017). In an invasion refugia, even a small to moderate decrease of

invasion pressure or invasive species density can allow coexistence of invasive and native

species, hence promoting native species persistence (Yokomizo et al. 2009, Sofaer et al. 2018,

Bradley et al. 2019, Strayer et al. 2019, DeRoy et al. 2020, Strayer 2020). This phenomenon

appears to be in accordance with the environmental matching hypothesis, stating that the impacts

of an invader decrease with environmental conditions that diverge from its physiological

optimum (Ricciardi et al. 2013, Iacarella and Ricciardi 2015). Our study joins a growing body of

literature exploring environmental factors that promote invasion refugia within invaded

freshwater ecosystems (Grabowski et al. 2009, Kestrup and Ricciardi 2009, Priddis et al. 2009,

Iacarella and Ricciardi 2015, Latzka et al. 2016, Astorg et al. 2020, Boddy et al. 2020, Boddy

and McIntosh 2021). More specifically, we aimed to address environmental determinants of

invasion hotspots and refugia of an invasive benthic fish, the round goby (*Neogobius*

melanostomus), in a large and complex river system, the St Lawrence River, which drains the

North American Great Lakes. Other previous studies on the role of environmental heterogeneity

in establishment of invasion refugia in the St Lawrence River have been done at limited spatial

scales or in laboratory experiments (Kestrup and Ricciardi 2009, Iacarella and Ricciardi 2015,

Astorg et al. 2020), potentially limiting inferences at the whole ecosystem scale.

The round goby is an invasive benthic littoral fish that was first observed in in the St. Lawrence

River in 1997 where it became increasingly abundant in many, but not all, habitats (Kipp et al.

2012, Morissette et al. 2018, Astorg et al. 2022). Its geographic range distribution in the St.

Lawrence River spans from Lake Ontario to the salt front of the St. Lawrence Estuary, ~30 km

downstream of Québec City. When present, the round goby has strong ecological effects,

notably by reducing the density of native benthic fishes such as darters (*Etheostoma spp.*) and

sculpins (*Cottus spp.*) that use similar habitats (Jude et al. 1992, Poos et al. 2010). The round

goby can also ravage the diversity and abundance of macroinvertebrate communities (Kipp et al.

2012, Astorg et al. 2022). However, certain types of habitats and environmental conditions can

reduce round goby abundance. In the Great Lakes, oxygen concentration and riparian habitat

quality were linked to invasion success of the round goby in some tributaries (Kornis et al. 2013,

Krabbenhoft and Kashian 2022, McAllister et al. 2022). The concentration of dissolved ions

(often measured by water conductivity or specific conductance, which is conductivity

standardized at 25 °C) can also predict the presence and abundance of the exotic round goby in

the Great Lakes (Baldwin et al. 2012) and in the St. Lawrence River (Astorg et al. 2020).

Another laboratory experiment from a limited portion of the St. Lawrence River showed that the

impact of conductivity on round goby performance was influenced by water calcium

concentration, where low dissolved calcium ions (Ca^{2+}) resulted in low predation rates and, thus,

reduced ecological impacts (Iacarella and Ricciardi 2015). Other studies have found evidence to

support that wetlands, through their substrate properties and associated submerged aquatic

vegetation, offer buffering effects from negative round goby impacts on native biodiversity at

invaded sites (Cooper et al. 2007, Astorg et al. 2020). Those results, albeit highly pertinent for

risk assessment and management of the round goby, were based on observations from a limited

portion of the St. Lawrence River, impeding the potential for generalization of the results.

In this study, we synthesize two extensive fish surveys along the St. Lawrence River, the Fish

Identification Nearshore Survey (FINS) and the *réseau de suivi ichtyologique* (RSI) respectively

operated by the River Institute and the *ministère des Forêts, de la Faune et des Parcs*. This

provided us with the opportunity to test the phenomenon of invasion hotspots and refugia for the

round goby on the largest scale of the entire non-tidal freshwater portion of the St. Lawrence

River. The objective of our study was to explore the relative influence of local environmental

conditions on the density of the round goby in the region covered by FINS and RSI. Specifically,

we aimed to test two hypotheses: 1) round goby density increases positively with water

conductivity in invaded waters and 2) the presence of wetlands in invaded waters reduces round

goby density. We predict that invaded sites with high water conductivity and wetland absence act

as invasion hotspots and invaded sites with lower water conductivity and wetland presence act as

invasion refugia from the round goby. Our study provides information for delimiting round goby

hotspots and refugia in the St. Lawrence River, helping to refine risk assessment of Ponto-

Caspian invasion in this ecosystem and others under imminent threat of round goby, as well as

promoting restoration and conservation of invasion refugium in the region.

Materials and Methods

Study area

We conducted our study over an area spanning approximately 400 km along the St. Lawrence

River, extending from Kingston (Ontario, Canada) at the outflow of Lake Ontario to Batiscan

(Québec, Canada) in the St. Lawrence River (Figure 1). The St. Lawrence River is a large fluvial

ecosystem draining the Laurentian Great Lakes toward the Atlantic Ocean. Its narrow fluvial

sections successively (upstream to downstream) drain three widenings of the St. Lawrence River,

called the fluvial lakes (St. François, St. Louis and St. Pierre lakes). Where the St. Lawrence

reaches the Archipelago of Montréal, the water from the Ottawa River, characterized by its low

conductivity (~60 to 100 µS/cm) and brown coloration, joins the Great Lakes water mass, which

has a higher water conductivity (200 to 400 µS/cm) and green coloration. Downstream of

Montréal, both water masses flow side by side, with very little lateral mixing (Vis et al. 1998,

Frenette et al. 2003), until the St. Lawrence reaches its tidal portion downstream of Lake St.

Pierre. Hence, the water masses keep their contrasted chemical properties for hundreds of

kilometres, creating a large ecotone where brown waters (Ottawa River) are mostly restricted to

the North shore and green waters (Great Lakes) to the South shore. The precise position of the

mixing zone and the two water masses stay relatively stable, with seasonal variations depending

on the magnitude of spring flood. Along the large-scale chemical gradients, various habitats can

be observed, from highly vegetated wetlands to sandy and rocky bottom sections, creating an

additional environmental variation of interest for fish and invertebrate communities.

Environmental and fish data

Fish community data were acquired using beach seine as part of both FINS and RSI annual fish

surveys, in mid to late summer over 14 years, between July 2007 to September 2021. The FINS

fishing was conducted using a 9.1 m long and 1.8 m deep purse seine, operated on three 20 m

long hauls for every studied site, corresponding approximately to 182 m^2 of surveyed habitat per

effort. The RSI used a 12.5 long by 4 m deep beach seine deployed on site and closed on itself by

operators, corresponding to approximately 120 m^2 of surveyed habitat per site. Both nets were

used in coastal, wadable (depth <1.5 m) habitats. Captured fish were identified to species and

counted by capture sites. As the spatiotemporal effect of round goby presence on those littoral

fish communities have already been assessed (Morissette et al. 2018), we limited the study to the

analysis of goby density. Site-specific round goby counts were converted to density (goby/m^2) to

allow comparisons between both FINS and RSI datasets.

Environmental data, conductivity (specific conductance) and wetland presence were either

measured and observed on site during sampling (FINS) or extrapolated from geographical data

available (RSI). During FINS surveys, conductivity was measured and recorded nearshore

(maximum one-meter depth) at each sampling site using a calibrated water quality multi-

parameter sonde (Professional plus model YSI ProDSS). Conductivity extrapolation was

conducted using 1660 points dataset acquired between 2017 and 2021, spanning the entire study

sites. Conductivity was interpolated using ordinary kriging analysis in the Spatial Analyst

toolbox implemented in the software ArcMap 10.7.1 (ESRI). Validation was conducted by

comparison of interpolated and measured values, producing a reliable agreement (R^2_{adj} = 0.84,

Figure S1). Conductivity values on fish sampling sites were extracted using the sampling

coordinates and the function *extract* coded within the package terra (Hijmans et al. 2022)

implemented in R version 4.1.2 (R Core Team 2021). For both FINS and RSI surveys, the

presence of wetlands on sites was extracted using a combination of an open-access high

resolution cartography of Ontario (Ministry of Natural Resources and Forestry 2021) and Quebec

wetlands (Canards Illimités Canada and ministère de l'Environnement et Lutte contre les

changements climatiques 2021). To simplify the dataset, only "open water" class was considered

as *non-wetlands* and all remaining classes (e.g., marsh, swamp, wet meadows) were considered

as *wetlands*, creating a binary *habitat type* factor.

Statistical modelling

All statistical analyses were computed in R version 4.1.2 R (R Core Team 2021) and

RStudio v2022.02.3+492 (RStudio Team 2020). Generalized additive models (GAMs) were used

to assess the effect of conductivity and habitat type on round goby density using the *mgcv* R

package v1.8-38 (Wood 2011). GAM is a useful framework to model linear and non-linear

relationships using smooth functions and can incorporate different family error distributions

(Wood 2017). GAM allow for the estimation of the non-linear "partial effect" of a predictor

variable, which represent the weighted sum of multiple basic functions of increasing wiggliness.

Two models were built in relation to our hypotheses. Both models included round goby density

(individual/m^2) as the response variable. The first model predictors were habitat type as a

parametric effect, conductivity by habitat factor-smooth interaction, longitude and latitude

smooth interaction and year as a random smooth while the second model included all listed

predictors, but the conductivity smooth was considered without the interaction with habitat type

and its parametric effect (Table S1). For both models, latitude and longitude smooth interaction

was included to consider any spatial correlation to ensure that model residuals are more likely to

be independent. Year was identified as a random predictor to generalize the effect over the years

and to account for any effect of time since invasion (Astorg et al. 2022). The default thin plate

spline was used for the conductivity smooth or for the smooth by factor interaction, the Duchon

spline was applied to the latitude and longitude smooth interaction and a random smooth

intercept was used for years. Since the response variable was continuous with left-skewed non-

integer values including zeros, an identity log link function with a Tweedie error distribution was

applied to both models. Model diagnostics and maximum basis functions (k) were verified using

the *gam.check* function and autocorrelation was assessed using the *acf* function of the model

residuals from the *stats* R package. No significant autocorrelation was detected in the GAMs and

diagnostic plots suggested normality and homoscedasticity of the residuals (Figure S2). Model

selection was done using the Akaike's Information Criterion (AIC) score (Wood et al. 2016).

The best-fitted model showed the lowest AIC score (Table S1). Model visualization was

accomplished using the *mgcViz* package v0.1.9 (Fasiolo et al. 2020) and *ggplot2* package v3.3.5

(Wickham et al. 2016).

Results

The fish dataset comprised 1310 sampling sites, with an average density of 0.07 goby/m^2 (±0.17

SD), ranging from 0 to 1.78 goby/m^2. Conductivity was variable in the study area, ranging from

43.8 to 576 µS/cm, with an average value of 244.69 µS/cm (±70.27 µS/cm SD). The expected

conductivity structure between the St. Lawrence North shore and South shore downstream of

Montréal was observed (Figure 1). Conductivity in the upper St. Lawrence was more

homogeneous, where variations were mainly influenced by small tributary outflows. Wetlands

and non-wetlands were equally represented in the study area, with 662 wetland sites and 648

non-wetland sites in the dataset, with both habitats present in approximately equivalent amounts

on both shores (Figure 1).

Our results indicate that both habitat type (wetland/non-wetland) and conductivity were

important predictors of round goby density in the St. Lawrence River. The best-fitted GAM

model showed the lowest AIC and included habitat type as a fixed effect, conductivity by habitat

type factor-smooth interaction along with the latitude and longitude smooth interaction and year

as a random smooth (Table S1). A statistical summary of the selected model is shown in Table 1.

Both the parametric fixed effect and partial smooth effects were significant (p-value <0.05) and

explained 55.8% of the null deviance with an R^2-adjusted value of 30.5%.

Habitat type showed a direct effect on the round goby density. Compared to non-wetland

habitats, wetlands had an important negative effect and therefore hosted lower round goby

densities (Figure 2). Water conductivity showed a non-linear partial effect on the round goby

density. However, the magnitude of the effect was different between habitat types and was

dependent on the range of the observed values. Overall, below approximately 220 µS/cm,

conductivity showed a negative effect on the round goby density (Figure 3). This negative effect

was found to be amplified in wetlands (Figure 3B) compared to non-wetland habitats

(Figure 3A). Above this threshold, water conductivity displayed a positive effect on the round

goby density in both habitats. At higher levels of conductivity (> 380 µS/cm), a negative effect

on round goby density was only observed in wetlands. However, the confidence interval is large

and includes the null effect. Overall, the round goby seems to exhibit a maximum density around

295 µS/cm, a potential biological optimum. The best-fitted GAM model demonstrated a

significant partial effect of latitude and longitude on the round goby density (Figure 4), showing

a spatially-coherent weighted sum of models' predictors on the round goby densities. A

considerable portion of the south shore of the St. Lawrence River displayed an important positive

effect on the round goby density while some locations, notably the Ottawa River and Deux-

Montagnes Lake (a widening of the Ottawa River) were locations with the most negative partial

effects on round goby density.

Discussion

Our study aimed to analyze a previously unpublished and extensive dataset provided by two

large-scale fish surveys conducted in the environmentally heterogeneous St. Lawrence River to

examine the effects of two environmental conditions—water conductivity and presence of

wetlands—on round goby presence and density. We used GAMs to determine thresholds for

ecological optima along the water conductivity gradient and showed the large-scale effects of

water conductivity on round goby density, as previously recognized in its invaded range

(Baldwin et al. 2012, Astorg et al. 2020). Our results showed that round goby density was

positively influenced by increasing conductivity until reaching a probable biological optimum of

~300 µS/cm. Sites of water conductivity under 200 µS/cm had low density to complete absence

of round goby (<100 µS/cm), even in regions hosting established populations for more than 10

years (Morissette et al. 2018), therefore representing possible invasion refugia. Additionally,

even if the presence of wetlands did not completely disrupt the pattern of water conductivity

effects on round goby density in invaded habitats, wetlands appeared to decrease round goby

densities along the conductivity continuum, contributing to the invasion refugium effect. Our

study is different from earlier research on the effect of wetlands and low water conductivity (or

dissolved ions) on round goby density as the detected invasion refugium in these other studies

were of either of limited size (Baldwin et al. 2012, Astorg et al. 2020) or laboratory-based

(Iacarella and Ricciardi 2015). By contrast, our study addressed the influence of those two

environmental factors along a over 400 km stretch of the St. Lawrence River, compiling more

than 1300 capture sites on a 14-year-old-time series. Hence, our results provide an empirical

criterion delineating round goby invasion hotspots and refugia. The sheer size of this study

underlines the possibility of generalizing the phenomenon and emphasizes the importance of

habitat diversity (i.e., wetland conservation and restoration) to buffer the effects of biological

invasion.

The detection of a predicable response of round goby density in the St. Lawrence suggests that

the risk of potential impact, that we consider density-dependent, could be identified by two

simple metrics, water conductivity and wetlands presence (Table 2). Herein, we propose that

non-wetland habitat with water conductivity under 100 µS/cm have a very low risk of goby

impacts, evaluated as null in wetlands under this conductivity. This risk increases with increasing

water conductivity until higher than 300 µS/cm, reaching the ecological optimum where impacts

are predicted to be maximal. Freshwater habitats over this threshold are known to host goby

populations with significant impacts (Baldwin et al. 2012, Kipp et al. 2012, Kornis et al. 2013).

However, In these round goby invasion hotspots, wetlands may still have buffering effects, albeit

limited in spatiotemporal scale, depending on submerged vegetation abundance, density and

phenology (Seilheimer and Chow-Fraser 2007, Janetski and Ruetz 2015). The simple risk grid

that our study provides (Table 2) can offer a fast and an easy to implement tool for managers and

could be disseminated on a large landscape with relatively accessible data. Management of

aquatic invasive species is often confronted with the challenge of rapidly identifying the present

or upcoming threat of a given species which can cause a vast array of impacts (Lodge et al.

2006). Identification of those impacts can largely vary in the invaded range and depart from what

has been identified in laboratory experiments and small-scale field studies. A broader

understanding of context-dependent potential density and their related impacts is needed to

implement an efficient management in heterogeneous landscapes and populate a more refined

impacts evaluation framework (Vander Zanden et al. 2017). Our finding that a local-scale

phenomenon, observed in 120 m^2 habitat patches, had a uniform response acrossat a large

environment scale underscores the value of observational studies in invasion biology and its

unique potential to offer generalizable risk assessment tools.

The negative effect of low water conductivity on round goby density is concordant with field

observations made by Astorg et al (2020) and Baldwin et al (2012), and our findings add to a

known effect of low conductivity on Ponto-Caspian invaders (Zhulidov et al. 2004, Kestrup and

Ricciardi 2009). Reduced Ponto-Caspian invader densities in waters with low concentrations of

dissolved ions are suggested to be linked with the evolutionary history of these euryhaline

species, which experienced repeated marine invasions—characterized by extreme variations of

salinity—favouring physiological processes requiring availability of high-dissolved ion

concentrations in ambient waters. Differences in round goby performance along freshwater

conductivity gradients may be linked to the concentration of calcium ions (Ca^{2+}), which

contribute to physiological processes and represent a likely common element between freshwater

and saltwater (Hunn 1985, Mayer et al. 1994, Baldwin et al. 2012, Iacarella and Ricciardi 2015).

Calcium represents a critical component of homeostasis in fish, by intervening on rates and

efficiency of gill-water exchanges. Accordingly, field observations suggest that the fish calcium

uptake from the environment seems to represent an evolutionarily conserved process, that does

not proportionally respond to large differences in environmental calcium, even after long-term

(multiple generations) residence along this environmental gradient (Sanderson et al. 2021).

Hence, the poor performance of round gobies—along with other Ponto-Caspian invaders—can

be a manifestation of the limited plasticity of those species under the unusual environmental

conditions imposed by low conductivity tributaries or water masses. This is an eloquent example

of environmental matching hypothesis and how the intolerance for low conductivity waters

translates in natural refugium to invasion (Ricciardi et al. 2013).

The mitigation of round goby presence and density in the St. Lawrence River by environmental

heterogeneity in water conductivity and local presence of wetlands are important for the

resilience of native fishes and macroinvertebrates populations in this culturally and socio-

economically important river system. Most ecological impacts of the round goby on littoral fish

community and macroinvertebrates appear to be density-dependant, and they are most severe at

local sites with environmental conditions favourable for the round goby establishment where the

fish has occurred since its introduction in the St. Lawrence River two decades ago (Morissette et

al. 2018, Astorg et al. 2022). In the face of this invasion, eradication and even population control

of the round goby is an unattainable goal with the present technology available to managers.

However, favouring coexistence of native species with round goby appears to be a more realistic

objective, both in the St. Lawrence River and in its different tributaries. Some tributaries of the

St. Lawrence Rivert have still not become completely colonized by the round goby, either

because of limiting environmental conditions or because of natural (e.g., waterfalls and rapids) or

anthropogenic (e.g., dams and weirs) fragmentation. Such habitats (e.g., St. Lawrence River

tributaries) host imperiled species predisposed to round goby impacts, notably the eastern sand

darter (*Ammocrypta pelludica*), the channel darter (*Percina copelandi*) and the pugnose shiner

(*Notropis anogenus*). Identification of the imminent risk is critical in ensuring conservation of

those species. We believe that our proposed risk analysis grid (Table 2) could assist management

agencies in either estimating the risk for round goby invasion and the identification of invasion

refugium meeting either the conductivity or wetlands criteria (or both).

Aside from its role buffering invasion impacts identified in our study, wetlands in freshwater

habitats are contributing to a vast array of ecological functions and services. For example,

wetlands have been shown to act as a buffer for wave-induced erosion (Silliman et al. 2019),

filtering runoffs inputs and contaminants (Díaz et al. 2012, Sileshi et al. 2020) and providing

spawning, nursery and feeding habitats for fishes, waterfowls, invertebrates and mammals

(Trebitz et al. 2009, Tanentzap et al. 2014). The sum of those ecosystem services underlines the

critical importance of wetlands in aquatic habitats. Sadly, wetlands have experienced drastic

declines, disappearing three times faster than forested areas (Convention on wetlands 2021).

Such declines, fuelled by anthropogenic modifications of freshwater habitats, pollution and

climate change (Johnson et al. 2010, Patenaude et al. 2015) can still be contained and even

reverted with the application of emergency plans of conservation (Tickner et al. 2020). Hence,

the preservation and restoration of wetlands does not onlydon't represent an aesthetic choice or

an application for the sake of biodiversity preservation—which already represent valuable

objectives—wetlands can also be viewed as a pragmatic tool in the integrated management of

water quality and prevention of biological invasions (Trepel 2010, Patenaude et al. 2015, Astorg

et al. 2020). Promotion of a diverse, yet complex mosaic of heterogeneous habitats could

potentially decrease invasion impacts and offer refugium for native species, thus favouring

coexistence and persistence (Melbourne et al. 2007, Priddis et al. 2009).

We recognize that factors for invasion success and ecological impacts of biological invasions are

complex and subject to multiple—and potentially synergistic—biotic and abiotic influences

(Catford et al. 2009, Ricciardi et al. 2020, Spear et al. 2021). Notably, over the large study

region, the influence of climatic variations (i.e., water temperature, amount of precipitation,

winter severity) could also contribute to the observed spatial effect. However, the contrasted

densities of the round goby observed in close sampling sites (i.e., in the vicinity of Montréal and

within Lake St. Pierre) suggest that climatic variations were probably of low importance in the

observed patterns that we report here. Round goby invasion success could be influenced by other

factors that we have not addressed by our study, such as the importance of propagule pressure,

the feeding opportunities (i.e., abundance of dreissenid mussels or invertebrate), other

environmental variables (i.e., sediment type and water flow) or ecological relationships (i.e.,

predation). However, overall, the striking effect of water conductivity on round goby density,

and therefore ecological impact on native fish and macroinvertebrate communities, offers an

interesting avenue to study mechanisms under invasion success and magnitude.

In conclusion, our study offers a unique opportunity to combine two ongoing and long-term

systematic fish surveys along a highly structured and complex freshwater ecosystem. We

observed a consistent effect of water chemistry and wetland presence in decreasing the density of

a notable Ponto-Caspian invader. Our results show that low water conductivity (<200 µS/cm), in

interaction with the presence of wetlands, are linked to a lower round goby density, buffering the

potential ecological impacts of this invader and providing a refugium for native aquatic species

such as macroinvertebrates and fish. The combination of easily measurable environmental

metrics can provide a simple, yet robust risk assessment tool for managers. Our results also

highlight the high value of wetlands conservation and restoration as a promoter of biodiversity

and an integrated management tool for conservation of native species, especially in invaded

ecosystems subjected to anthropogenic disturbances.

Acknowledgments

This research was supported by the Chaire de recherche sur les espèces aquatiques exploitées

(MFFP and UQAC) to OM and NSERC Discovery Grant to AMD and OM. Funding to MW and

the FINS Project was provided by Fisheries and Oceans Canada's Habitat Stewardship Program

for Aquatic Species at Risk, the Ontario Ministry for Natural Resources and Forestry's Species at

Risk Stewardship Fund, WWF Canada's Loblaw Water Fund, Ontario Power Generation, the

Trottier Family Foundation, and the River Institute. CC was supported by scholarships provided

by MITACS and Fonds du Québec – nature et technologie (FRQNT). The authors are grateful

for the field crew and partners who contributed to the FINS and RSI, notably Chantal Côté, Yves

Paradis, Kate Schwartz, and staff at the Mohawk Council of Akwesasne's Environment

Program.. All fish collections were approved under the Institutional Animal Protection

Committee either at MFFP, UQAM and the River Institute and authorized by licences for

scientific fish collections from the Ontario Ministry of Natural Resources and Forestry and the

ministère des Forêts, de la Faune, et des Parcs du Québec..

Authors contributions

All authors contributed to the ideation of the study, the manuscript writing and the revision. MW

and CC contributed to the FINS survey and OM and AD contributed to the RSI survey. MW, CC

and OM did the data curation and quality assessment and CC and OM realized the statistical

analyses, the production of the figures and the tables.

Conflict of interest statement

The authors declare no financial or personal conflict of interest.

References

Anton, A., M. S. Simpson and I. Vu (2014). "Environmental and biotic correlates to lionfish
invasion success in Bahamian coral reefs." <u>PLOS ONE</u> **9**(9): e106229.

Astorg, L., C. Charette, M. J. S. Windle and A. M. Derry (2022). "Two decades since first
invasion: Revisiting round goby impacts on nearshore aquatic communities in the Upper St.
Lawrence River." <u>Journal of Great Lakes Research</u>.

Astorg, L., S. Sanderson, V. Côté-Gravel, F. Sorbara, M. J. S. Windle, A. P. Hendry and A. M.
Derry (2020). "Different refuge types dampen exotic invasion and enhance diversity at the whole
ecosystem scale in a heterogeneous river system." <u>Biological Invasions</u>.

Bacher, S., T. M. Blackburn, F. Essl, P. Genovesi, J. Heikkilä, J. M. Jeschke, G. Jones, R. Keller,
M. Kenis, C. Kueffer, A. F. Martinou, W. Nentwig, J. Pergl, P. Pyšek, W. Rabitsch, D. M.
Richardson, H. E. Roy, W.-C. Saul, R. Scalera, M. Vilà, J. R. U. Wilson and S. Kumschick
(2018). "Socio-economic impact classification of alien taxa (SEICAT)." <u>Methods in Ecology and
Evolution</u> **9**(1): 159-168.

Baldwin, B. S., M. Carpenter, K. Rury and E. Woodward (2012). "Low dissolved ions may limit
secondary invasion of inland waters by exotic round gobies and dreissenid mussels in North
America." <u>Biological Invasions</u> **14**(6): 1157-1175.

Boddy, N. C., D. J. Booker and A. R. McIntosh (2020). "Heterogeneity in flow disturbance
around river confluences influences spatial patterns in native and non-native species co-
occurrence." <u>Biological Invasions</u> **22**(11): 3457-3475.

Boddy, N. C. and A. R. McIntosh (2021). "Could spatial heterogeneity in flow disturbance drive
temporal stability of native–invasive species co-occurrence in riverscapes?" <u>Freshwater Biology</u>
66(5): 902-913.

Bradley, B. A., B. B. Laginhas, R. Whitlock, J. M. Allen, A. E. Bates, G. Bernatchez, J. M. Diez,
R. Early, J. Lenoir, M. Vilà and C. J. B. Sorte (2019). "Disentangling the abundance-impact
relationship for invasive species." <u>Proceedings of the National Academy of Sciences</u> **116**(20):
9919-9924.

Byers, J. E., S. H. Reichard, J. M. Randall, I. M. Parker, C. S. Smith, W. M. Lonsdale, I. A. E.
Atkinson, T. Seastedt, M. H. Williamson, E. A. Chornesky and D. C. Hayes (2002). "Directing
research to reduce the impacts of nonindigenous species." <u>Conservation Biology</u> **16**.

Canards Illimités Canada and ministère de l'Environnement et Lutte contre les changements
climatiques (2021). "Cartographie détaillée des milieux humides des secteurs habités du sud du
Québec – Données du projet global."

Catford, J. A., R. Jansson and C. Nilsson (2009). "Reducing redundancy in invasion ecology by
integrating hypotheses into a single theoretical framework." <u>Diversity and Distributions</u> **15**(1):
22-40.

Convention on wetlands (2021). Global wetlands outlook: Special edition 2021. Gland,
Switzerland, Secretariat on the convention on wetlands.

Cooper, M. J., C. R. Ruetz, D. G. Uzarski and T. M. Burton (2007). "Distribution of round
gobies in coastal areas of lake Michigan: Are wetlands resistant to invasion?" Journal of Great
Lakes Research **33**(2): 303-313, 311.

Dawson, W., D. Moser, M. van Kleunen, H. Kreft, J. Pergl, P. Pyšek, P. Weigelt, M. Winter, B.
Lenzner, T. M. Blackburn, E. E. Dyer, P. Cassey, S. L. Scrivens, E. P. Economo, B. Guénard, C.
Capinha, H. Seebens, P. García-Díaz, W. Nentwig, E. García-Berthou, C. Casal, N. E. Mandrak,
P. Fuller, C. Meyer and F. Essl (2017). "Global hotspots and correlates of alien species richness
across taxonomic groups." Nature Ecology & Evolution **1**(7): 0186.

DeRoy, E. M., R. Scott, N. E. Hussey and H. J. MacIsaac (2020). "Density dependence mediates
the ecological impact of an invasive fish." Diversity and Distributions **26**(7): 867-880.

Díaz, F. J., A. T. O′Geen and R. A. Dahlgren (2012). "Agricultural pollutant removal by
constructed wetlands: Implications for water management and design." Agricultural Water
Management **104**: 171-183.

Dyer, E. E., P. Cassey, D. W. Redding, B. Collen, V. Franks, K. J. Gaston, K. E. Jones, S. Kark,
C. D. L. Orme and T. M. Blackburn (2017). "The global distribution and drivers of alien bird
species richness." PLOS Biology **15**(1): e2000942.

Fasiolo, M., R. Nedellec, Y. Goude and S. N. Wood (2020). "Scalable visualization methods for
modern generalized additive models." Journal of Computational and Graphical Statistics **29**(1):
78-86.

Frenette, J.-J., M. T. Arts and J. Morin (2003). "Spectral gradients of downwelling light in a
fluvial lake (Lake Saint-Pierre, St-Lawrence River)." Aquatic Ecology **37**(1): 77-85.

Grabowski, M., K. Bacela, A. Konopacka and K. Jazdzewski (2009). "Salinity-related
distribution of alien amphipods in rivers provides refugia for native species." Biological
Invasions **11**(9): 2107.

Hijmans, R. J., R. Bivand, K. Forner, J. Ooms, E. Pebesma and M. D. Sumner (2022). terra:
Spatial Data Analysis.

Hunn, J. B. (1985). "Role of calcium in gill function in freshwater fishes." Comparative
Biochemistry and Physiology Part A: Physiology **82**(3): 543-547.

Iacarella, J. C. and A. Ricciardi (2015). "Dissolved ions mediate body mass gain and predatory
response of an invasive fish." Biological Invasions **17**(11): 3237-3246.

IPBES (2019). Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. E. S.
Brondizio, J. Settele, S. Diaz and H. T. Ngo. Bonn, Germany, IPBED Secretariat.

Janetski, D. J. and C. R. Ruetz (2015). "Spatiotemporal patterns of fish community composition in Great Lakes drowned river mouths." Ecology of Freshwater Fish **24**(4): 493-504.

Johnson, W. C., B. Werner, G. R. Guntenspergen, R. A. Voldseth, B. Millett, D. E. Naugle, M. Tulbure, R. W. H. Carroll, J. Tracy and C. Olawsky (2010). "Prairie wetland complexes as landscape functional units in a changing climate." BioScience **60**(2): 128-140.

Jude, D. J., R. H. Reider and G. R. Smith (1992). "Establishment of gobiidae in the Great Lakes basin." Canadian Journal of Fisheries and Aquatic Sciences **49**(2): 416-421.

Kestrup, Å. M. and A. Ricciardi (2009). "Environmental heterogeneity limits the local dominance of an invasive freshwater crustacean." Biological Invasions **11**(9): 2095-2105.

Kipp, R., A. Ricciardi and C. W. Ramcharan (2012). "Impacts of the Eurasian round goby (*Neogobius melanostomus*) on benthic communities in the upper St. Lawrence River." Canadian Journal of Fisheries and Aquatic Sciences **69**(3): 469-486.

Kornis, M. S., S. Sharma and J. Vander Zanden (2013). "Invasion success and impact of an invasive fish, round goby, in Great Lakes tributaries." Diversity and Distributions **19**(2): 184-198.

Krabbenhoft, C. A. and D. R. Kashian (2022). "Invasion success of a freshwater fish corresponds to low dissolved oxygen and diminished riparian integrity." Biological Invasions.

Latzka, A. W., G. J. A. Hansen, M. Kornis and M. J. Vander Zanden (2016). "Spatial heterogeneity in invasive species impacts at the landscape scale." Ecosphere **7**(3): e01311.

Lewis, M., S. Petrovskii and J. Potts (2016). Dynamics of biological invasions: 19-68.

Lockwood, J. L., P. Cassey and T. M. Blackburn (2009). "The more you introduce the more you get: The role of colonization pressure and propagule pressure in invasion ecology." Diversity and Distributions **15**(5): 904-910.

Lodge, D. M., S. Williams, H. J. MacIsaac, K. R. Hayes, B. Leung, S. Reichard, R. N. Mack, P. B. Moyle, M. Smith, D. A. Andow, J. T. Carlton and A. McMichael (2006). "Biological invasions: Recommendations for U.S. policy and management." Ecol Appl **16**(6): 2035-2054.

Mayer, F. L. J., L. L. Marking, T. D. Bills and G. E. Howe (1994). Physicochemical factors affecting toxicity in freshwater: hardness, pH and temperature. Bioavailability: physical, chemical and biological interactions. J. L. Hamelink, P. F. Landrum, H. L. Bergman and W. H. Benson. London, Lewis Publishers: 256.

McAllister, K., D. A. R. Drake and M. Power (2022). "Round Goby (*Neogobius melanostomus*) impacts on benthic fish communities in two tributaries of the Great Lakes." Biological Invasions.

Melbourne, B. A., H. V. Cornell, K. F. Davies, C. J. Dugaw, S. Elmendorf, A. L. Freestone, R. J. Hall, S. Harrison, A. Hastings, M. Holland, M. Holyoak, J. Lambrinos, K. Moore and H.

Yokomizo (2007). "Invasion in a heterogeneous world: resistance, coexistence or hostile takeover?" Ecology Letters **10**(1): 77-94.

Ministry of Natural Resources and Forestry (2021). Ontario Land Information. Ottawa, Government of Ontario.

Morissette, O., Y. Paradis, R. Pouliot and F. Lecomte (2018). "Spatio-temporal changes in the littoral fish community structure along the St. Lawrence River (Québec, Canada) following the round goby (*Neogobius melanostomus*) invasion." Aquatic Invasions **13**(4): 501-512.

Patenaude, T., A. C. Smith and L. Fahrig (2015). "Disentangling the effects of wetland cover and urban development on quality of remaining wetlands." Urban Ecosystems **18**(3): 663-684.

Pejchar, L. and H. A. Mooney (2009). "Invasive species, ecosystem services and human well-being." Trends in Ecology & Evolution **24**(9): 497-504.

Poos, M., A. J. Dextrase, A. N. Schwalb and J. D. Ackerman (2010). "Secondary invasion of the round goby into high diversity Great Lakes tributaries and species at risk hotspots: potential new concerns for endangered freshwater species." Biological Invasions **12**(5): 1269-1284.

Priddis, E., R. Rader, M. Belk, B. Schaalje and S. Merkley (2009). "Can separation along the temperature niche axis promote coexistence between native and invasive species?" Diversity and Distributions **15**(4): 682-691.

R Core Team (2021). R: A language and environment for statistical computing. Vienna, Austria, R foundation for statistical computing.

Ricciardi, A., M. F. Hoopes, M. P. Marchetti and J. L. Lockwood (2013). "Progress toward understanding the ecological impacts of nonnative species." Ecological Monographs **83**(3): 263-282.

Ricciardi, A., J. C. Iacarella, D. C. Aldridge, T. M. Blackburn, J. T. Carlton, J. A. Catford, J. T. A. Dick, P. E. Hulme, J. M. Jeschke, A. M. Liebhold, J. L. Lockwood, H. J. MacIsaac, L. A. Meyerson, P. Pyšek, D. M. Richardson, G. M. Ruiz, D. Simberloff, M. Vilà and D. A. Wardle (2020). "Four priority areas to advance invasion science in the face of rapid environmental change." Environmental Reviews.

RStudio Team (2020). RStudio: Integrated Development for R. RStudio. Boston, MA, PBC.

Sanderson, S., A. M. Derry and A. P. Hendry (2021). "Phenotypic stability in scalar calcium of freshwater fish across a wide range of aqueous calcium availability in nature." Ecology and Evolution **n/a**(n/a).

Seebens, H., T. M. Blackburn, E. E. Dyer, P. Genovesi, P. E. Hulme, J. M. Jeschke, S. Pagad, P. Pyšek, M. van Kleunen, M. Winter, M. Ansong, M. Arianoutsou, S. Bacher, B. Blasius, E. G. Brockerhoff, G. Brundu, C. Capinha, C. E. Causton, L. Celesti-Grapow, W. Dawson, S. Dullinger, E. P. Economo, N. Fuentes, B. Guénard, H. Jäger, J. Kartesz, M. Kenis, I. Kühn, B. Lenzner, A. M. Liebhold, A. Mosena, D. Moser, W. Nentwig, M. Nishino, D. Pearman, J. Pergl,

W. Rabitsch, J. Rojas-Sandoval, A. Roques, S. Rorke, S. Rossinelli, H. E. Roy, R. Scalera, S. Schindler, K. Štajerová, B. Tokarska-Guzik, K. Walker, D. F. Ward, T. Yamanaka and F. Essl (2018). "Global rise in emerging alien species results from increased accessibility of new source pools." <u>Proceedings of the National Academy of Sciences</u> **115**(10): E2264-E2273.

Seebens, H., T. M. Blackburn, E. E. Dyer, P. Genovesi, P. E. Hulme, J. M. Jeschke, S. Pagad, P. Pyšek, M. Winter, M. Arianoutsou, S. Bacher, B. Blasius, G. Brundu, C. Capinha, L. Celesti-Grapow, W. Dawson, S. Dullinger, N. Fuentes, H. Jäger, J. Kartesz, M. Kenis, H. Kreft, I. Kühn, B. Lenzner, A. Liebhold, A. Mosena, D. Moser, M. Nishino, D. Pearman, J. Pergl, W. Rabitsch, J. Rojas-Sandoval, A. Roques, S. Rorke, S. Rossinelli, H. E. Roy, R. Scalera, S. Schindler, K. Štajerová, B. Tokarska-Guzik, M. van Kleunen, K. Walker, P. Weigelt, T. Yamanaka and F. Essl (2017). "No saturation in the accumulation of alien species worldwide." <u>Nature Communications</u> **8**(1): 14435.

Seilheimer, T. S. and P. Chow-Fraser (2007). "Application of the wetland fish index to northern Great Lakes marshes with emphasis on Georgian Bay coastal wetlands." <u>Journal of Great Lakes Research</u> **33**: 154-171.

Sileshi, A., A. Awoke, A. Beyene, I. Stiers and L. Triest (2020). "Water purifying capacity of natural riverine wetlands in relation to their ecological quality." <u>Frontiers in Environmental Science</u> **8**.

Silliman, B. R., Q. He, C. Angelini, C. S. Smith, M. L. Kirwan, P. Daleo, J. J. Renzi, J. Butler, T. Z. Osborne, J. C. Nifong and J. van de Koppel (2019). "Field experiments and meta-analysis reveal wetland vegetation as a crucial element in the coastal protection paradigm." <u>Current Biology</u> **29**(11): 1800-1806.e1803.

Simberloff, D., J.-L. Martin, P. Genovesi, V. Maris, D. A. Wardle, J. Aronson, F. Courchamp, B. Galil, E. García-Berthou, M. Pascal, P. Pyšek, R. Sousa, E. Tabacchi and M. Vilà (2013). "Impacts of biological invasions: what's what and the way forward." <u>Trends in Ecology & Evolution</u> **28**(1): 58-66.

Sofaer, H. R., C. S. Jarnevich and I. S. Pearse (2018). "The relationship between invader abundance and impact." <u>Ecosphere</u> **9**(9): e02415.

Spear, M. J., J. R. Walsh, A. Ricciardi and M. J. V. Zanden (2021). "The Invasion Ecology of Sleeper Populations: Prevalence, Persistence, and Abrupt Shifts." <u>BioScience</u>.

Strayer, D. L. (2020). "Non-native species have multiple abundance–impact curves." <u>Ecology and Evolution</u> **10**(13): 6833-6843.

Strayer, D. L., C. T. Solomon, S. E. G. Findlay and E. J. Rosi (2019). "Long-term research reveals multiple relationships between the abundance and impacts of a non-native species." <u>Limnology and Oceanography</u> **64**(S1): S105-S117.

Tanentzap, A. J., E. J. Szkokan-Emilson, B. W. Kielstra, M. T. Arts, N. D. Yan and J. M. Gunn (2014). "Forests fuel fish growth in freshwater deltas." <u>Nature Communications</u> **5**(1): 4077.

Tickner, D., J. J. Opperman, R. Abell, M. Acreman, A. H. Arthington, S. E. Bunn, S. J. Cooke, J.
Dalton, W. Darwall, G. Edwards, I. Harrison, K. Hughes, T. Jones, D. Leclère, A. J. Lynch, P.
Leonard, M. E. McClain, D. Muruven, J. D. Olden, S. J. Ormerod, J. Robinson, R. E. Tharme,
M. Thieme, K. Tockner, M. Wright and L. Young (2020). "Bending the curve of global
freshwater biodiversity loss: An emergency recovery plan." BioScience **70**(4): 330-342.

Trebitz, A. S., J. C. Brazner, D. N. P, M. S. Pearson, G. S. Peterson, D. K. Tanner, D. L. Taylor,
C. W. West and T. P. Hollenhorst (2009). "Geographic, anthropogenic, and habitat influences on
Great Lakes coastal wetland fish assemblages." Canadian Journal of Fisheries and Aquatic
Sciences **66**(8): 1328-1342.

Trepel, M. (2010). "Assessing the cost-effectiveness of the water purification function of
wetlands for environmental planning." Ecological Complexity **7**(3): 320-326.

Vander Zanden, M. J., G. J. A. Hansen and A. W. Latzka (2017). "A Framework for Evaluating
Heterogeneity and Landscape-Level Impacts of Non-native Aquatic Species." Ecosystems **20**(3):
477-491.

Vis, C., A. Cattaneo and C. Hudon (1998). "Periphyton in the clear and colored water masses of
the St. Lawrence River (Québec, Canada): A 20-year overview." Journal of Great Lakes
Research **24**(1): 105-117.

Wickham, H., D. Navarro and T. L. Pedersen (2016). ggplot2: Elegant Graphics for Data
Analysis. Houston, Texas, USA, Springer.

Wood, S. N. (2011). "Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models." Journal of the Royal Statistical Society:
Series B (Statistical Methodology) **73**(1): 3-36.

Wood, S. N. (2017). Generalized Additive Models. New York, Chapman and Hall/CRC.

Wood, S. N., N. Pya and B. Säfken (2016). "Smoothing parameter and model selection for
general smooth models." Journal of the American Statistical Association **111**(516): 1548-1563.

Yokomizo, H., H. P. Possingham, M. B. Thomas and Y. M. Buckley (2009). "Managing the
impact of invasive species: the value of knowing the density–impact curve." Ecological
Applications **19**(2): 376-386.

Zhulidov, A. V., D. F. Pavlov, T. F. Nalepa, G. H. Scherbina, D. A. Zhulidov and T. Y.
Gurtovaya (2004). "Relative distributions of Dreissena bugensis and Dreissena polymorpha in
the Lower Don River system, Russia." International Review of Hydrobiology **89**(3): 326-333.
604

607 **Table 1.** Statistical summary of the generalized additive model (GAM) selected including habitat
608 type as a fixed effect, conductivity by habitat type factor-smooth interaction, longitude and latitude
609 smooth interaction and year as a random smooth. Edf indicates the estimated degree of freedom
610 and Ref.df represents the residual degree of freedom. Significant parametric (A) and partial effects
611 (B) are indicated by p-value <0.05.

A. parametric coefficients	Estimate	Standard Error	t-value	p-value
(Intercept)	-3.8625	0.1487	-25.97	<0.0001
Wetlands	-0.4896	0.1479	-3.31	<0.005

B. smooth terms	edf	Ref.df	F-value	p-value
s(conductivity): non-wetlands	7.198	8.941	4.127	<0.0001
s(conductivity): wetlands	4.15	5.246	6.764	<0.0001
s(longitude, latitude)	55.791	79	17.365	<0.0001
s(year)	9.85	14	3.003	<0.0001

R2-adjusted	0.305			
Deviance explained	55.8 %			

613　**Table 2.** Binary risk of round goby density (and density-dependent impacts) evaluation grid

614　based on specific conductance (conductivity) and local presence of wetlands on a local to large-

615　scale study site.

	Water specific conductance (conductivity)			
Local habitat	<100 µS/cm	100 to 200 µS/cm	200 to 300 µS/cm	> 300 µS/cm
Non-wetlands	very low	low	medium	high
Wetlands	null	very low	medium*	high*

* General agreement to large-scale risk with possibility of local or seasonal buffering effects of

wetlands

Figure 1. Round goby density (proportional red circles) in relationship to water conductivity

gradient (blue color shade) and wetlands presence (green polygons) presented on a subset of the

study for visualisation. The complete study area is presented in the smaller context map with fish

sampling sites locations (black dots).

Figure 2. Generalized additive model (GAM) parametric effect of habitat type (non-wetland and

wetland) on round goby density (individual/m^2). Non-wetland habitat represents the intercept in

the GAM, rug plot shows observations of the predictor variable, and the dash lines indicate the

95% confidence intervals.

Figure 3. Generalized additive model (GAM) partial smooth effect of conductivity (µS/cm) on

round goby density (individual/m^2) by habitat type: non-wetlands (A) and wetlands (B). Shaded

area indicates the 95% confidence interval, rug plot shows observations of the predictor variable

and dash lines represent the null effect.

Figure 4. Generalized additive model (GAM) of partial spatial smooth effect on round goby

density (individual/m^2) along the geographic gradient of the St. Lawrence River study area. The

range of colours represents the effect: yellow shows the most positive effect (higher density) and

dark blue the most negative effect (lower density). Partial residuals are represented by dots.

Figure 1.

Figure 2.

Figure 3.

Figure 4.