Successfully treating biofilms of extensively drug resistant *Klebsiella pneumoniae* isolates from hospital patients with N-Acetyl Cysteine

Ankurita Bhowmik[1], Sambuddha Chakraborty[1], Anusha Rohit [2*], Ashwini Chauhan [1*]

1. Department of Microbiology, Tripura University, Suryamaninagar, Agartala West Tripura, India-799022
2. Department of Microbiology, Madras Medical Mission Hospital, Chennai, Tamil Nadu, India-600003

*Equal contributions

Email address: ashwini.chauhan@gmail.com, anusha3rohit@gmail.com

Phone number: +91- 9555128395

Abstract*: Klebsiella pneumoniae* is one of the leading causes of community and nosocomial infections. Reduced treatment options against extensively drug resistant (XDR) - *K. pneumoniae,* is a serious concern in hospital settings, and hence, WHO has categorized it as a "critical priority pathogen". Biofilm forming ability is a common virulence mechanism amongst *K. pneumoniae* that is associated with antibiotic tolerance up to 1000X MIC and hence, are difficult to treat. N-acetyl cysteine (NAC) is an FDA approved mucolytic drug used to treat acetaminophen-associated toxicity and obstructive pulmonary diseases. In this study, we assessed NAC's antibacterial and antibiofilm activity against clinical isolates of XDR *K. pneumoniae,* obtained from Madras Medical Mission Hospital, India. To assess the biofilm eradication ability of NAC, we grew biofilms in 96 well plates and treated the mature biofilms with different concentrations of NAC. We observed that the biofilms of only 3 isolates of XDR *K. pneumoniae* could be eradicated at a concentration as low as 20mg/ml. Although increasing the concentration of NAC to 80mg/mL could significantly reduce the biofilms of all the isolates up to 4-5 Log, NAC at a concentration of 100 mg/mL successfully eradicated the mature biofilms of all the isolates of XDR *K. pneumoniae*. This *in vitro* study demonstrates the potential of NAC as an efficient agent against the biofilms of clinical isolates of XDR-*K. pneumoniae* and thus, provides a promising alternative to antibiotics.

Keywords: *Klebsiella pneumoniae*, Multidrug resistance, Biofilm, N-acetyl cysteine, Nosocomial infections.

1.   Introduction:

Klebsiella pneumoniae is a ubiquitous, Gram negative, non-motile, and encapsulated bacillus that colonizes skin, mouth, respiratory tract, and digestive tract in human. It is a major cause of community and hospital acquired pneumonia, urinary tract infection, sepsis, and meningitis in immunocompetent and immunocompromised patients (Joseph et al., 2021; Liu et al., 2021). Recent studies have demonstrated the association of *K. pneumoniae* with severe infections including pyogenic liver abscesses; endophthalmitis etc. (Russo & Marr, 2019). Diverse virulence factors such as siderophores, capsular polysaccharides, lipopolysaccharides, and fimbrial adhesins contribute to the pathogenicity of *K. pneumoniae* (Remya et al., 2019; Gomes et al., 2021). Moreover, the propensity of *K. pneumoniae* to acquire multidrug resistance has made them more notorious to deal with in the hospital environment. In 2020, the Indian Council of Medical Research (ICMR) categorized *K. pneumoniae* as a Group-I pathogen due to the emergence of resistance toward the last line of antibiotics resulting in increased morbidity and mortality rate. The global priority pathogen (GPP) list published by WHO in 2017 has also enlisted ESBL producing carbapenem-resistant *K.pneumoniae* among the "critical priority pathogens" (WHO 2017). Moreover, *K. pneumoniae* was recently reported to be prevalent in up to 18% of clinical specimens analyzed in India. Furthermore, ICMR has also disclosed an extended resistance in *K. pneumoniae clinical* isolates towards cephalosporins, aminoglycosides, carbapenem, fluoroquinolones, and polymyxin *K. pneumoniae* (Indian Council of Medical Research, 2020; Ashwath et al., 2022). *K. pneumoniae* possesses a natural resistance to the penicillin class of antibiotics due to the production of sulfhydryl variable (SHV-1) penicillinases (Kakoullis et al., 2021). In addition, the presence of plasmid encoded extended spectrum beta-lactamases (ESBL) including cefotaximase-M (CTM-M) in *K. pneumoniae* confers resistance towards broad spectrum aminoglycosides and cephalosporins (Piperaki et al., 2017; Karaiskos & Giamarellou, 2020). Moreover, different carbapenemases such as metallo-beta-lactamases, *Klebsiella pneumoniae* carbapenemases, and oxacillinases confers carbapenem resistance in *K. pneumoniae* (Lopes et al., 2020).

Besides different resistance mechanisms, biofilm forming ability of *K. pneumoniae* confers additional protection from antibiotics and the host immune system (Guilhen et al., 2019; Singh et

al., 2021). The biofilm matrix enables the development of heterogenous bacterial subpopulations
including metabolic inactive persister cells due to gradients of oxygen, nutrients, metabolic
products, and accumulated toxins. The persister cells can survive lethal dosage of antibiotics and
hence, confer increased antibiotic tolerance (Chauhan et al., 2014; Lebeaux et al., 2014). Biofilm
development on indwelling devices and epithelial tissues is clinically significant and is widely
reported (Chauhan, et al., 2012; Lebeaux et al., 2013).

Globally, the rise and spread of XDR-*K. pneumoniae* infections have renewed the research interest
in studying the correlation between antibiotic resistance and biofilm encoding genes in *K.*
pneumoniae (Effah et al., 2020; Ballén et al., 2021; Shadkam et al., 2021). Currently, different
combinations of antibiotics such as meropenem with fosfomycin, and/or gentamicin,
ceftazidime/avibactam with fosfomycin, or gentamicin or tigecycline, etc. are used to treat XDR-*K.*
pneumonia infections (Daikos et al 2014; Jacobs et al 2017; Karaiskos et al 2019; Petrosillo et al
2019; Yu et al 2019; Bassetti et al 2021). Besides, alternatives to antibiotics have also been
explored such as bacteriophage therapy in conjunction with non-active antibiotics such as
sulfamethoxazole-trimethoprim was effective against XDR-*K. pneumoniae* isolate (Bao et al.,
2020). Recent reports demonstrated the potential of antimicrobial peptides such as AA139, SET-
M33, and ΔM2 against colistin resistant *K. pneumoniae* (van der Weide et al., 2019; Rivera-
Sánchez et al., 2020). EDTA in combination with colistin is recently reported to manage colistin
resistant *K. pneumoniae* (Shein et al., 2021). However, to manage the drug resistance challenge
among *K. pneumoniae,* constant exploration of novel non-antibiotic therapeutic interventions is
urgently required.

N-acetyl cysteine (NAC) is a synthetic derivative of endogenous amino acid, L-cysteine, and a
precursor of glutathione. It is an FDA approved mucolytic drug used for decades to treat
acetaminophen overdose and obstructive pulmonary diseases (Tenório et al., 2021). Due to its
mucolytic activity, NAC has attracted microbiologists to explore its potential as an antibacterial and
antibiofilm agent. A number of in vitro studies assessed the activity of NAC against different Gram
positive and Gram-negative bacteria including *Escherichia coli, Acinetobacter baumannii,*
Pseudomonas aeruginosa, Klebsiella pneumoniae, Staphylococcus aureus, and Staphylococcus
epidermidis amongst others (Quah et al.,2012; Choi et al.,2018; Feng et al.,2018; Jun et al.,2019;
Ciacci et al., 2019; Li et al.,2020; Guerini et al.,2020; Kundukad et al., 2020; Manoharan et al.,
2021). Given the broad-spectrum activity of NAC against bacteria, this study aims at repurposing

N-acetyl cysteine (NAC) to treat biofilms of XDR *K. pneumoniae,* isolated from patients in
hospitals.

In our study, a total of 1402 Gram-negative clinical isolates were collected during the period from
2018 to 2020 at Madras Medical Mission Hospital, Chennai India. Out of this, 374 (26.6%) were
identified as *Klebsiella spp*. and intriguingly, 99% of isolates amongst the *Klebsiella spp*. were *K.*
pneumoniae. Based on the prevalence and resistance profile, we selected 12 XDR *K. pneumoniae*
isolates and assessed the *in vitro* antibacterial as well as antibiofilm activity of NAC against these
isolates. We observed that NAC was bactericidal against all the XDR *K. pneumoniae* isolates at a
concentration of 10mg/mL. Furthermore, NAC could successfully eradicate *in vitro* biofilms of all
the XDR *K. pneumonia* at an acceptable concentration of 100mg/mL (Olsson et al., 1988; Hughes et
al., 2018; NIH U.S. National Library of Medicine 2018, Guerini et al., 2022). Safe alternatives to
antibiotics are a compelling need of the hour to treat *K. pneumoniae* infections to improve patient
outcomes. N-acetyl cysteine may be a safe option for treating XDR *K. pneumoniae* leading to
reduced socio-medico-economic burden.

2. Materials and Methods:
2.1. Bacterial strains and Growth media: All the XDR-*Klebsiella pneumoniae* isolates were
grown at 37^{o}C in Luria-Bertani Broth (LB) purchased from Himedia, India (M1245). For viable cell
estimation the serially diluted culture was plated on sterile LB agar plates and incubated overnight
at 37^{o}C.
2.2. Antimicrobial agents: N-acetyl cysteine (NAC) was purchased from Sigma Aldrich
(A7250). Isotonic NAC solution was prepared by dissolving NAC in sterile 0.9% saline.
2.3. Estimation of antibiotic susceptibility: VITEK 2 compact (bioMerieux, France) was used
for biochemical characterization and determining the antibiogram of all the XDR-*K. pneumoniae*
isolates using disposable antibiotic susceptibility Testing (AST) cards. VITEK 2 instrument
continuously monitors the bacterial growth and records the fluorescence, turbidity, and photometric
signals in presence of an increasing concentration of various antibiotics. Wells with no antibiotics
served as the positive control (Ligozzi et al 2002).
2.4. Assessing the Minimum inhibitory Concentration (MIC): The MIC of NAC against the
clinical isolates of XDR-*K. pneumoniae* was determined by broth microdilution assay (Chauhan et
al., 2012; Lebeaux et al., 2014; Lewis et al.,2022). Briefly, NAC was serially diluted in LB in a

microtiter plate to give a 2-fold serial dilution. An equal volume of exponentially growing *K.*

pneumoniae isolate was added to give a final inoculum of 5 X 10^5 CFU/mL in all the wells except

the negative control. Well with bacteria but no NAC, served as an untreated control. The plate was

incubated for 18h at 37°C under static conditions and the first well without visible growth was noted

as the MIC concentration. All the experiments were done at least in triplicate.

2.5.   Effect of NAC on Exponential phase planktonic XDR-*K. pneumoniae*: Growth inhibition

efficacy of NAC against clinical isolates of XDR- *K. pneumoniae* was done in 96 well microtitre

plates using a BioteK Epoch-2 microtiter plate reader (Agilent Technologies). Each treatment was

replicated in at least 3 wells. Log phase culture (OD600 ≈ 0.3-0.5) of *K. pneumoniae* isolates were

treated with sub-MIC (2.5 or 5mg/mL) concentration to 80mg/mL of NAC for 24h at 37°C with

continuous shaking. Untreated wells served as control. The OD600nm were measured by the Biotek

Epoch-2 plate reader for 24 hrs. After 24h, the viable cell count was estimated by plating the serial

dilutions on LB agar media, and plates were incubated at 37°C overnight (Lebeaux et al., 2014).

2.6.   *In vitro* biofilm forming ability of Clinical isolates of XDR *K. pneumoniae*: The biofilm

forming ability of the clinical isolates of XDR *K. pneumoniae* was determined by assessing the

ability of the cells to adhere to the wells of 96-well polystyrene microtiter plates by previously

described crystal violet (CV) staining method (O'Toole et al.,1998; Chauhan et al 2013). An

overnight culture of *K. pneumoniae* grown in LB at 37°C under shaking was diluted 1/100 in fresh

LB (1uL inoculum in 100uL LB media per well) and incubated at 37°C under static conditions for

72h. After incubation, the non-adherent cells were gently removed. The wells were gently washed

thrice with 0.9% saline and 125uL 0.1% CV stain was added to the wells. The plates were incubated

at room temperature for 15 mins and the wells were rinsed with 0.9% saline. The plates were then

air dried for 15mins before solubilizing the CV with acetone-ethanol (1:4), and incubated at room

temperature for 15 minutes. Biofilm biomass was estimated as a function of CV stain by measuring

the absorbance at 595nm. The results are an average of five technical replicate wells from three

independent experiments. The wells containing media without bacteria were used as the negative

control, and strain *K. pneumoniae* KPP1 and KPP2 were used as positive controls as they form

strong biofilms under invitro laboratory conditions (unpublished data). The cut-off OD (ODc) is

defined as three standard deviations (SD) above the mean OD of the negative control, i.e ODc=

average OD of negative control+ (3xSD of negative control). XDR-*K. pneumoniae* isolates were

categorized as negative, weak biofilm formers, moderate or strong biofilm formers as follows:

negative biofilm formers: OD ≤ ODc; weak biofilm producer: $(OD_c < OD \leq 2\times OD_c)$, moderate

biofilm producer $(2\times OD_c < OD \leq 4\times OD_c)$, strong biofilm producer $(OD > 4\times OD_c)$.

2.7.　***In vitro* assessment of the inhibitory potential of NAC against the biofilms formed by**

Clinical isolates of XDR *K. pneumoniae*: The biofilm inhibition assays were done as described by

Lebeaux et al 2014 with slight modifications (Lebeaux et al., 2014). The overnight cultures of *K.*

pneumoniae isolates were inoculated in fresh media at 1:100 dilution in a 96-well microtiter plate

and incubated for 96h to form mature biofilm at 37ºC without shaking. The planktonic bacteria

were removed by gentle pipetting and the wells were rinsed thrice with 0.9% saline. The 96h old

mature biofilms growing in the micro-titre plates were treated with different concentrations of

isotonic solution of NAC (5mg/mL, 10mg/mL, 20mg/mL, 40mg/mL, 80mg/mL and 100mg/mL) by

directly adding 125uL NAC solution into the wells. The treatment was allowed to dwell in the wells

for 24h at 37ºC under static conditions. 0.9% saline treated wells served as untreated controls. After

24h, the NAC solution was removed from the wells and rinsed thrice with 0.9% saline. Wells were

scraped vigorously to remove any residual biofilm and homogenized with 0.9% saline. The viable

cell count was estimated by plating the serial dilutions on LB agar medium and the plates were

incubated at 37ºC. Each treatment was given in replicate of four and each experiment was repeated

at least thrice.

2.8.　**Cytotoxicity assay:** Cytotoxicity of N-acetylcysteine (NAC) was assessed on L929 (mouse

fibroblasts), and J774A.1 (mouse macrophages) cell lines (procured from National Centre for Cell

Sciences, Pune, India). Two concentrations of NAC; 80 mg/mL & 40 mg/mL in 0.9% isotonic

saline solution were subjected to the test. The assay was performed using CyQUANT™ MTT Cell

proliferation assay kit, Invitrogen. Briefly, approximately 4000 cells of both cell types were seeded

per well of 96-well microtiter plates and kept for incubation at 37 °C with 5% CO2. After

attainment of 70-80% confluency, spent media were discarded from all the wells including the

control wells. Two test concentrations of NAC were inoculated in the designated wells. Control

wells were left with the media. The plates were incubated at 37 °C with 5% CO2 for 30 minutes.

Post incubation, NAC was aspirated out from every test well and replaced by fresh media. MTT [3-

(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide] solution was added to all the treated

and control wells, and the plate was incubated at 37 °C for four hrs. in 5% CO2. Post MTT

incubation, all the contents from each well were removed, followed by the addition of DMSO. After

adding DMSO, the plates were incubated for 10 minutes, and the absorbance was measured using

EPOC 2, 96 well plate reader (BioTek) at 570 nm.

2.9.   Statistical analysis: All the results are replicates of at least three independent experiments.

Statistical differences were evaluated using one-way analysis of variance (ANOVA) using

Graphpad Prism version 8.0. *P* values lower than 0.05, were considered statistically significant.

3.   Results:

3.1.   *K. pneumoniae* is most prevalent amongst the Gram negative bacteria isolated from

clinical settings.

A total of 1402 Gram-negative bacterial samples were isolated from patients admitted to Madras

Medical Mission Hospital during 2018-2020. Among these, *Klebsiella spp.,* was found to be the

second predominant bacterial species among all the Gram-negative bacteria. *Klebsiella spp.,*

comprised 26.6% of the isolated Gram-negative bacteria. Intriguingly, 99% of isolates amongst the

Klebsiella spp. were *K. pneumoniae* and only 1% was *K. oxytoca* (Fig.1).

Fig 1: Prevalence of *Klebsiella spp*. and its drug resistance profile in clinical settings of Madras

Medical Mission Hospital, India. A total of 1402 Gram negative bacteria were isolated from

hospital settings and characterized for identification. **A.** *Klebsiella spp.* comprised 26.6%, amongst

the isolated Gram negative bacteria. **B.** Of all the *Klebsiella spp.*, 99% were *K. pneumoniae* that

showed resistance to multiple classes of antibiotics including carbapenems and polymyxin. **C.** *K.*

pneumoniae was more prevalent among samples isolated from patients with respiratory tract

infections (42%). U-Urine, B-Blood, ST-Soft tissues, RT-Respiratory tract, IA-Intraabdominal.

3.2.*K. pneumoniae* isolates showed resistance to the different classes of antibiotics.

The characterization and antibiotic susceptibility profiling of the *K. pneumoniae* isolates was done by an automated Vitek 2 compact system (bioMerieux, France). All the *K. pneumoniae* isolates showed resistance to multiple classes of antibiotics (Fig 1B). We selected 12 XDR-*K. pneumoniae* isolates in our study as they showed resistance to almost all the tested antibiotics (Fig 2). It is worrisome to note that all the isolates showed 100% resistance towards amikacin, amoxicillin-clavulanate, cefazolin, gentamicin, ceftriaxone, cefuroxime, and co-trimoxazole. Furthermore, only less than 20% of isolates were susceptible to imipenem, meropenem, ertapenem, pip-tazobactam, and cefoperazone sulbactam. Most of the isolates (90%) showed resistance towards tigecycline, netilmicin, ciprofloxacin, ofloxacin, and cefepime. Colistin is currently used as a final resort for treating multidrug resistant *K. pneumoniae*. However, genetic mutations in the *phoPQ* and *pmrAB* system result in colistin resistance in *K. pneumoniae* (Nirwan et al., 2021). Recent reports also suggest *mcr* plasmid-mediated colistin resistance in *K. pneumoniae* (Karki et al., 2021; Salloum et al., 2020). In this study, we also observed that 25% *K. pneumoniae* isolates were colistin resistant, which is concerning, and indicative of the probable spread of colistin resistance genes. A recent report in 2021 showed the tendency of carbapenem resistant *K. pneumoniae* to acquire colistin resistance under tigecycline or colistin pressure (Andrew Evans et al., 2021). Similarly, in our study, we observed that all the colistin resistant *K. pneumoniae* isolates were resistant to carbapenems.

Fig 2: Drug resistance profile of 12 XDR-*K. pneumoniae* isolates used in this study. The heatmap represents the antibiogram and extensive antibiotic resistance among the *K. pneumoniae* isolates. Rows represent *K. pneumoniae* isolates and columns represent different antibiotics. The red blocks signify resistance (R), the blue blocks signify intermediate (I), and the green blocks signify susceptibility (S) to antibiotics. AMK-Amikacin, GEN-Gentamicin, CFZ-Cefazolin, SCF-Cefoperazone sulbactam, FEP-Cefepime, CRO-Ceftriaxone, CTX-Cefotaxime, CXM-Cefuroxime, CIP-Ciprofloxacin, OFX-Ofloxacin, CST-Colistin, IPM-Imipenem, MEM-Meropenem, ETP-Ertapenem, NTL-Netilmycin, PIP-TAZ-Pip tazobactam, AMC-Amoxicillin clavulanate, FOF-Fosfomycin, SXT-Co-trimoxazole, TGC-Tigecycline

3.3.   NAC effectively kills clinical isolates of XDR-*K. pneumoniae*.

To assess the anti-bacterial efficacy of NAC, a weak protic acid, we determined its minimum inhibitory concentration (MIC) against the clinical isolates of XDR-*K. pneumoniae* as per CLSI guidelines. Intriguingly, we observed ~ 83% (10 out of 12) of the XDR *K. pneumoniae* isolates that we chose for the study were susceptible to 5mg/mL of NAC. The remaining 2 XDR *K. pneumoniae* isolates were susceptible to a two-fold more NAC concentration i.e., 10mg/mL (Table 1). Furthermore, we checked the effect of NAC on the cell viability of XDR-*K. pneumoniae* isolates by treating exponentially grown cells with NAC from sub-MIC concentration (2.5 or 5mg/mL) to 80mg/mL for 24h and checked the changes in growth kinetics and colony forming units (CFU) (Table 2) (Fig S.1). We observed that NAC was bactericidal at MIC concentration against all the clinical isolates except KPO760. At MIC (5mg/mL), NAC reduced the growth of KPO760 up to 4.8logs, which completely disappeared with a two-fold increase in MIC (10mg/mL).

Table 1: Minimum Inhibitory Concentration of NAC against clinical isolates of XDR *K pneumoniae*

Clinical isolates of XDR *K. pneumoniae (Kp)*	MIC (mg/mL)
ON2352	5
O3875	5
ON4400	5
O3818	5

O3337	5
O760	5
O4444	5
O1125	10
O3283	5
O2040	5
ON2697	5
O3159	10
*MIC results are mean results of at least 3 experiments	

Table 2: Effect of different concentrations of NAC on exponentially growing clinical isolates of XDR-*K. pneumoniae*. * ND: Not done							
	NAC Treatment (mg/mL)						
K. pneumoniae isolates	Untreated	2.5	5	10	20	40	80
	(CFU/mL)						
ON2352	1.1×10^8 (SD:2.94E+07)	7.6×10^3 (SD:2.34E+03)	0 (MIC)	0	0	0	0
O3159	2.7×10^9 (SD:5.27E+08)	ND*	5.3×10^3 (SD:2.42E+03)	0 (MIC)	0	0	0
O3818	8.6×10^9 (SD:2.42E+09)	1.0×10^6 (SD:2.34E+05)	0 (MIC)	0	0	0	0
ON4400	1.2×10^7 (SD:2.31E+06)	4.7×10^6 (SD:6.12E+05)	0 (MIC)	0	0	0	0

ON4569	9.6 x 10^9 (SD:2.94E+09)	1.3 x 10^6 (SD:3.29E+05)	0 (MIC)	0	0	0	0
O4444	1.4 x 10^{10} (SD:1.79E+09)	4.6 x 10^5 (SD:2.07E+05)	0 (MIC)	0	0	0	0
ON2697	2.5 x 10^{10} (SD:4.52E+09)	1.2 x 10^7 (SD:4.27E+06)	0 (MIC)	0	0	0	0
O3337	8.6 x 10^9 (SD:2.42E+09)	9.6 x 10^5 (SD:1.51E+05)	0 (MIC)	0	0	0	0
O3875	1.1 x 10^8 (SD:2.34E+07)	1.0 x 10^5 (SD:4.20E+04)	0 (MIC)	0	0	0	0
O1125	5.6 x 10^9 (SD:5.98E+08)	ND*	6.5 x 10^4 (SD:5.01E+03)	0 (MIC)	0	0	0
O3283	9.6 x 10^7 (SD:6.87E+06)	3.3 x 10^5 (SD:3.88E+04)	0 (MIC)	0	0	0	0
O 760	2.6 x 10^9 (SD:1.97E+06)	1.4 x 10^7 (SD:1.97E+06)	1.2 x 10^5 (MIC; SD:3.15E+04)	0	0	0	0

3.4.   Clinical isolates of XDR *K. pneumoniae* are moderate biofilm formers.

The biofilm forming ability of 12 XDR-*K. pneumoniae* isolates were assessed by the previously described CV staining method (Stepanović et al., 2007; Leoney et al., 2020). The cut off OD (ODc)

obtained was 0.076. We observed that 58% (7/12) of the isolates were moderate biofilm formers,

33% (4/12) were weak biofilm formers and 8% (1/12) were strong biofilm former among the XDR

K. pneumoniae isolates (Fig.3). Interestingly, three of the *K. pneumoniae* isolates, KPON4569,

KPO3875, and KPON4400, that are resistant to all the antibiotics, are weak biofilm formers. All the

moderate and strong biofilm formers showed variable antibiotic resistance patterns. It would be

interesting to study the effect of multidrug resistance on the fitness of *K. pneumoniae* isolates.

Fig 3: Biofilm producers among XDR-*K. pneumoniae* isolates. The biofilm forming ability of all

the isolates was determined by the CV staining method in 96 well plates. The X-axis represents the

K. pneumoniae isolates and the Y-axis represents the optical density at 595nm. The strains were

classified based on their biofilm forming abilities according to previous reports (Stepanović et al.,

2007; Leoney et al., 2020).

3.5. NAC effectively eliminates mature XDR- *K. pneumoniae* biofilms

Managing biofilm associated infections is challenging especially when the infections are due to

antibiotic resistant isolates such as XDR- *K. pneumoniae*. We evaluated the *in vitro* antibiofilm

activity of NAC against the 12 XDR-*K. pneumoniae* isolates. The 96h preformed biofilms were

treated with different concentrations of NAC prepared in an isotonic solution, ranging from

5mg/mL to 100mg/mL for 24h and the cell viability was estimated by CFU/mL plating. Although

we observed a varying level of killing of biofilm-associated XDR-*K. pneumoniae* isolates treated

with *K. pneumoniae* NAC upon 24h exposure, NAC could successfully eradicate the biofilms of all

the XDR isolates (Table 3).

Although, the viable cell count in the mature biofilms of isolates KPON2697 and KPO3875 were

significantly reduced up to 2-log at sub-MIC and MIC concentrations i.e. 5mg/mL respectively, the

complete biofilm eradication could be achieved using 10mg/mL NAC solution. Mature biofilms of

Table 3: Antibiofilm activity NAC against mature biofilm of XDR-*K. pneumoniae* isolates.

Kp isolates	Untreated	NAC concentration(mg/mL)					
		5	10	20	40	80	100
ON2352	2.7×10^{7} (SD: 3.27E+06)	1.9×10^{5} (SD: 4.56E+05)	1.7×10^{5} (SD: 8.14E+04)	4.7×10^{3} (SD: 2.42E+04)	3.3×10^{3} (SD: 3.58E+03)	1.6×10^{3} (SD: 8.16E+02)	0
O3159	1.2×10^{9} (SD: 4.27E+08)	1.1×10^{7} (SD: 2.94E+06)	3.6×10^{6} (SD: 2.42E+06)	1.4×10^{6} (SD: 1.41E+06)	6.3×10^{4} (SD: 7.45E+05)	4.6×10^{3} (SD: 3.44E+05)	0
O3818	1.9×10^{10} (SD: 4.34E+08)	1.3×10^{9} (SD: 2.73E+08)	1.0×10^{8} (SD: 1.74E+08)	9.6×10^{7} (SD: 2.10E+07)	4.3×10^{6} (SD: 2.94E+06)	5.6×10^{4} (SD: 2.66E+04)	0
ON4400	1.0×10^{10} (SD: 3.44E+09)	1.3×10^{7} (SD: 3.01E+06)	1.0×10^{7} (SD: 2.53E+06)	9.0×10^{6} (SD: 2.97E+06)	5.0×10^{6} (SD: 2.10E+06)	4.3×10^{5} (SD: 2.34E+05)	0
ON4569	1.4×10^{9} (SD: 4.34E+08)	7.6×10^{8} (SD: 2.42E+08)	4.6×10^{8} (SD: 2.42E+08)	4.3×10^{7} (SD: 2.42E+07)	4.6×10^{5} (SD: 8.91E+05)	1.1×10^{4} (SD: 1.51E+03)	0
ON2697	9.0×10^{6}	7.0×10^{4}	0	0	0	0	0

	(SD: 2.10E+06)	(SD: 3.03E+04)					
O3875	6.3×10^{7} (SD: 2.8E+07)	3.3×10^{5} (SD: 1.97E+06)	0	0	0	0	0
O3337	1.7×10^{8} (SD: 2.42E+08)	1.0×10^{6} (SD: 2.07E+06)	5.6×10^{5} (SD: 2.97E+04)	0	0	0	0
O1125	6.3×10^{8} (SD: 2.48E+05)	1.7×10^{7} (SD: 1.57E+05)	8.3×10^{6} (SD: 2.06E+04)	1.9×10^{5} (SD: 2.94E+06)	8.6×10^{4} (SD: 3.22E+05)	7.3×10^{3} (SD: 2.12E+04)	0
O760	8.0×10^{9} (SD: 2.57E+05)	3.5×10^{8} (SD: 3.05E+05)	1.9×10^{6} (SD: 1.48E+07)	7.1×10^{5} (SD: 4.52E+0)	1.5×10^{4} (SD: 2.5E+04)	4.0×10^{3} (SD: 1.07E+05)	0
O3283	9.6×10^{8} (SD: 3.03E+04)	5.6×10^{7} (SD: 3.50E+05)	4.6×10^{6} (SD: 2.97E+06)	1.3×10^{5} (SD: 2.32E+0)	8.3×10^{4} (SD: 2.42E+08)	4.6×10^{3} (SD: 1.51E+03)	0
O4444	5.6×10^{8} (SD: 2.34E+08)	2.5×10^{7} (SD: 8.04E+07)	1.5×10^{6} (SD: 4.86E+07)	4.2×10^{5} (SD: 4.73E+0)	4.1×10^{4} (SD: 3.50E+05)	5.6×10^{3} (SD: 1.51E+04)	0

XDR isolate KPO3337, were also markedly reduced by 2-log and 3-log reduction in biofilm cells at
5mg/mL and 10mg/mL NAC, respectively. Increasing the NAC concentration just by 2-fold
(20mg/mL) resulted in the eradication of biofilms formed by isolate KPO3337 (Fig.4). A
concentration dependent anti-biofilm activity of NAC was observed against other XDR-*K.*
pneumoniae isolates. Although 80mg/mL NAC solution reduced the biofilm cell viability up to 6-
log (99.999%), some viable cell population was still recovered. Interestingly, biofilms formed by all
the isolates couldn't tolerate 100mg/mL NAC solution and were completely eradicated after 24hrs
treatment.

| A. KP O3875 | B. KP O3337 |

Fig 4: Representative graphs of *invitro* antibiofilm activity of NAC measured by CFU/ml (effect on cell viability) against XDR-*K. pneumoniae* isolates: A. KP O3875 and B. KP O3337 compared to untreated biofilm.

3.6.   NAC is non-toxic to L929 and J774A.1 cell lines.

NAC is an FDA approved compound and is considered one of the important ingredients to prepare powerful antioxidant rich food (Šalamon et al., 2019). Since NAC is an FDA approved compound and the most efficient antibiofilm concentration of NAC was 100 mg/mL that led to the total eradication of *in vitro* biofilms of XDR- *K. pneumoniae* isolates. We evaluated the cell toxicity by treating the mouse fibroblast cells (L929) and macrophage cells (J774A.1) with 80 or 100mg/mL NAC for 30mins. Incubating cells with 80 mg/mL or 100 mg/mL concentration of NAC did not exert any toxic effect on the cell lines as indicated by the high rate of cell survival of both the cell types used in the study (Fig.5). J774A.1 showed a 100% survival rate at both the tested concentrations. On the same hand, L929 also had a high survival rate of up to 91% survival at both the tested concentrations. A slight decrease in survival rate in L929 is probably due to a shock that appeared from sudden depletion in nutritional media. Higher Survival percentage in J774A.1 cell may be due to the reasons that NAC enhances the activity of macrophages (Vecchiarelli et al.,1994).

Fig 5: Cytotoxicity of NAC against J774A.1 and L929 cell lines after 30 mins of exposure. A) For J774A.1, 100% viability is seen post NAC treatment with both the concentrations, and B) a negligible 9% cytotoxicity was observed against L929 cells.

4. Discussion:

K. pneumoniae are classified as "priority pathogens" by WHO and are one of the most important contributors to nosocomial infections (WHO 2017). The emergence of antibiotic resistance and the biofilm forming ability of pathogens including *K. pneumoniae* pose one of the greatest challenges to clinicians in treating such infections and the health system at large. Adding to the problem is a rapid rise in extensively drug resistant (XDR) bacteria that are extremely difficult to treat sometimes even due to the ineffectiveness of the last line of antibiotics. In this study, we selected XDR- *K. pneumoniae* clinical isolates to assess the efficacy of an FDA approved molecule, N-acetyl cysteine (NAC) against actively growing as well as biofilm bacteria. We report here that, of the total *Klebsiella* isolates from clinical sources, 99% were *K. pneumoniae.* Earlier clinical data also reported the high prevalence of *K. pneumoniae* species in the clinical samples (Heidary et al., 2018; Igrejas et al., 2019; Asri et al., 2021; Odari & Dawadi, 2022). Carbapenems are the first choice of clinicians for treating drug resistant -*K. pneumoniae* (Bassetti et al., 2018). Imipenem is a carbapenem antibiotic that was reported as the most effective antibiotic in treating *K. pneumoniae* associated infections. However, lately, the emergence of carbapenemase producing *K. pneumoniae* has rendered the carbapenem class of antibiotics insensitive (Papadimitriou-Olivgeris et al., 2017; Ashwath et al., 2022). In our study, we observed that imipenem resistance was predominant in up to 83% of clinical isolates. A similar scenario was also demonstrated by Ashwath et al.,2022, where

more than 80% resistance among *K. pneumoniae* isolates was observed against imipenem (Ashwath

et al., 2022). Besides, *K. pneumoniae* isolates also showed the least sensitivity toward other

carbapenems, viz. meropenem and ertapenem (Papadimitriou-Olivgeris et al.,2017; Matovina et al.,

2021., Ashwath et al., 2022)

It is worrisome to note that recent studies have shown the emergence of resistance in clinical

isolates of *K. pneumoniae K. pneumoniae* against third generation cephalosporins and

fluroquinolones (Xie et al.,2018; Geetha et al., 2020; Renggli et al., 2022; Rocha et al., 2022). In the

present study, all the isolates of *K. pneumoniae* were resistant to cefazolin, ceftriaxone, cefotaxime,

and cefuroxime. Moreover, 83% of isolates were resistant to cefoperazone sulbactam and 91% of

isolates were insensitive to cefepime. Among fluoroquinolones including ciprofloxacin and

ofloxacin, only 9% susceptibility was observed.

Furthermore, the rate of colistin resistance has also increased among *K. pneumoniae* (Ah et

al.,2014; Zafer et al.,2019; Azam et al.,2021; Narimisa et al.,2022). In our study also we observed

25% colistin resistant isolates, which is certainly concerning. The increased antibiotic resistance is

attributed to several factors, however, the prevalence of ESBL encoding genes such as *bla* $_{SHV}$, *bla*

$_{CTX-M}$, *bla* $_{TEM}$; and plasmids harboring quinolone resistant *qnrS, qnrA, qnrB,* and colistin resistant

mcr genes, are regarded as the main contributor of resistance among *K. pneumoniae* (Piperaki et al.,

2017; Ashwath et al., 2022).

Another crucial factor responsible for persistent *K. pneumoniae* infections is the development of

biofilm on host tissues and indwelling devices. Moreover, the correlation between biofilm

formation and antibiotic resistance is well established in different studies (Vuotto et al., 2017;

Potempa et al., 2018; Nirwati et al.,2019). Although all the *K. pneumoniae* clinical isolates studied

were capable of forming biofilms *in vitro,* 59% of the isolates were moderate biofilm formers, 33%

were weak biofilm formers and only 8% were strong biofilm formers. Earlier studies also showed

that clinical isolates of *K. pneumoniae* were either moderate or weak biofilm formers (Magesh et

al.,2013; Cusumano et al.,2019; Singh et al., 2019). As reported by earlier studies, we also observed

that all the *K. pneumoniae* isolates (25% of total *K. pneumoniae* isolates) that were resistant to all

the tested antibiotics were weak biofilm formers. Similarly, we noticed that all the moderate biofilm

formers are also antibiotic resistant to several antibiotics, viz. *K. pneumoniae* 100% resistance was

observed against co−trimoxazole, 85% resistance was observed against imipenem, meropenem,

tigecycline, and ofloxacin, and 71% resistance was observed against amoxicillin clavulanate,

amikacin, gentamicin, ciprofloxacin, cefazolin, cefoperazone sulbactam, ceftriaxone, cefepime, cefotaxime, and cefuroxime. Reports suggest that the presence of plasmids carrying resistant genes in *K. pneumoniae,* modulates the expression of diverse biofilm associated genes (Ashwath et al., 2022). Possibly this may be true for the multiple drug resistant *K. pneumoniae* isolated from clinics in our study.

With the rising rates of antibiotic resistance among *K. pneumoniae*, non-antibiotic alternatives are currently being explored. N-acetyl cysteine (NAC) is an FDA approved drug used to treat acetaminophen poisoning and obstructive pulmonary diseases. Recently use of NAC as an antibacterial and antibiofilm agent has gained tremendous interest. Some studies have successfully used NAC to demonstrate its *in vitro* efficacy against planktonically grown as well as biofilms of laboratory strains of bacteria (Li et al., 2020; Aiyer et al., 2021; Chlumsky et al., 2021; Tenório et al., 2021). The antibiofilm activity of NAC is attributed to different aspects such as inhibition of EPS production, bacterial adhesion inhibition, competitive inhibition of cysteine in bacteria, interaction with bacterial proteins via thiol groups, and liquification of biofilm matrix (Zhao et al., 2010; Costa et al.,2017; Manoharan et al.,2020; Guerini et al.,2020; Guerini et al., 2022). Some reports have shown the killing efficacy of NAC is pH-mediated. NAC in its undissociated form can penetrate through the bacterial cell membrane and inside the cell, it dissociates into its conjugate base and H^+ decreasing the intracellular pH that may interfere with DNA replication, RNA transcription, and protein translation (Kundukad et al., 2020).

In this study, we assessed the potential of NAC against the clinical isolates of XDR-*K. pneumoniae*, some of these were resistant to all the tested antibiotics. We observed that the minimum concentration of NAC required for killing the exponentially growing *K. pneumoniae* ranged between 5 to10 mg/mL and a significant number of viable cells were recovered from the cultures treated with NAC concentrations less than 5mg/mL. In 2011, Darouiche and the group also reported that the MIC and MBIC of NAC against the clinical isolates of *K. pneumoniae* were 5 and 10 mg/mL, respectively (Aslam et al., 2011).

Recently, a few groups reported that NAC was bactericidal at a concentration as low as 2mg/mL (Kundukad et al., 2020). This may be attributed to differential strain specific anti-bacterial activity of NAC and probably clinical isolates have comparatively higher tolerance toward NAC. Moreover, our cell viability results indicated that NAC is bactericidal to *K. pneumoniae* isolates contradicting previous studies that reported NAC as bacteriostatic (Manoharan et al., 2020, Aiyer et al., 2021).

However, few other studies demonstrated that NAC is bacteriostatic at low concentrations and
bactericidal at higher concentrations (Perry et al., 1977; Aslam et al., 2011). We then assessed the
antibiofilm activity of NAC at different concentrations against preformed 96 hr old mature biofilms
of *K. pneumoniae*. Although, *in vitro* biofilms of 16.6% of isolates and of 8.3% of isolates were
eradicated at 10mg/mL and 20mg/mL respectively, complete eradication of biofilms of the majority
of the XDR-*K. pneumoniae* isolates could be achieved only at a minimum of 10X MIC of NAC i.e.
100mg/mL.

Biofilms are a source of recurrent infections, which is attributed to the presence of highly antibiotic-
tolerant persister bacteria within biofilms, and is one of the major challenges for the management of
biofilm-associated infections in the clinical setting (Darouiche RO.,2004; Lewis K., 2007; Lebeaux
et al.,2014). NAC proved to be a potential therapeutic option that is able to eradicate even the
persistent bacteria of XDR-*K. pneumoniae* isolates at a concentration of 100mg/mL.

Although the clinical use of acidic compounds may be potentially toxic, our choice of NAC as an
alternative therapeutic to antibiotics is persuasive as several reports suggested the lack of toxicity of
intravenous administration of NAC to patients in various clinical situations (Prescott et al., 1979;
Smilkstein et al., 1991; Perry et al., 1998; Buckley et al., 1999).

Besides, NAC acts as a ROS scavenger and a cysteine prodrug that releases cysteine inside the cells
at a slow rate helping in the restoration of the glutathione levels. (Jiao et al., 2016; Pedre et al.,
2021). NAC showed cytotoxicity preventive effects on U937 monocytes and human kidney
epithelial cells (HK-2 cell line) (Zhang et al., 2011; Ghani et al., 2014). Similarly, we showed that
treating mouse fibroblast cells (L929) and macrophage cells (J774A.1) with NAC up to a
concentration of 100mg/mL had no cytotoxicity even after 30 minutes of exposure.

In conclusion, we summarize that physiologically tolerable concentrations of NAC can completely
eradicate the *in vitro* mature biofilms of all tested XDR-*K. pneumoniae* clinical isolates. Hence, we
believe that the NAC deserves further preclinical and clinical assessments for effective management
of WHO critical priority pathogen XDR- *K. pneumoniae* associated nosocomial infections.

Acknowledgement: AC would like to thank UGC-BSR (F.30-487/2019(BSR)), DST-Nanomission
(DST/NM/NB/2018/203), ICMR (OMI/20/2020-ECD-1) and SERB-CRG (CRG/2021/001974) for
the funding support.

References:

Ah, Y. M., Kim, A. J., & Lee, J. Y. (2014). Colistin resistance in *Klebsiella*
pneumoniae. International journal of antimicrobial agents, 44(1), 8–15.
https://doi.org/10.1016/j.ijantimicag.2014.02.016

Aiyer, A., Visser, S. K., Bye, P., Britton, W. J., Whiteley, G. S., Glasbey, T., Kriel, F. H., Farrell,
J., Das, T., & Manos, J. (2021). Effect of *N*-Acetylcysteine in Combination with Antibiotics
on the Biofilms of Three Cystic Fibrosis Pathogens of Emerging Importance. Antibiotics
(Basel, Switzerland), 10(10), 1176. https://doi.org/10.3390/antibiotics10101176

Andrew Evans, B., Zhou, H., Jabalameli, F., Wang, H., Zhang, Y., Wang, X., Wang, S., Sun, S.,
Li, H., Chen, H., & Wang, Q. (2021). Emergence of Colistin Resistance in Carbapenem-
Resistant Hypervirulent *Klebsiella pneumoniae* Under the Pressure of Tigecycline. *Frontiers*
in microbiology, *12*, 756580.https://doi.org/10.3389/fmicb.2021.756580

Ashwath, P., Deekshit, V. K., Rohit, A., Dhinakaran, I., Karunasagar, I., Karunasagar, I., &
Akhila, D. S. (2022). Biofilm Formation and Associated Gene Expression in Multidrug-
Resistant *Klebsiella pneumoniae* Isolated from Clinical Specimens. Current Microbiology,
79(3). https://doi.org/10.1007/s00284-022-02766-z

Asri, N. A. M., Ahmad, S., Mohamud, R., Hanafi, N. M., Zaidi, N. F. M., Irekeola, A. A., Shueb,
R. H., Yee, L. C., Noor, N. M., Mustafa, F. H., Yean, C. Y., & Yusof, N. Y. (2021). Global
prevalence of nosocomial multidrug-resistant *Klebsiella pneumoniae:*A systematic review and
meta-analysis. Antibiotics (Vol. 10, Issue 12). https://doi.org/10.3390/antibiotics10121508

Azam, M., Gaind, R., Yadav, G., Sharma, A., Upmanyu, K., Jain, M., & Singh, R. (2021). Colistin
Resistance among Multiple Sequence Types of *Klebsiella pneumoniae* Is Associated With
Diverse Resistance Mechanisms: A Report From India. Frontiers in microbiology, 12,
609840. https://doi.org/10.3389/fmicb.2021.609840

Ballén, V., Gabasa, Y., Ratia, C., Ortega, R., Tejero, M., & Soto, S. (2021). Antibiotic Resistance
and Virulence Profiles of *Klebsiella pneumoniae* Strains Isolated From Different Clinical
Sources. Frontiers in Cellular and Infection Microbiology, 11.
https://doi.org/10.3389/fcimb.2021.738223

Bassetti, M., Righi, E., Carnelutti, A., Graziano, E., & Russo, A. (2018). Multidrug-resistant
Klebsiella pneumoniae: Challenges for treatment, prevention and infection control. In Expert

Review of Anti-Infective Therapy (Vol. 16, Issue 10, pp. 749–761).
https://doi.org/10.1080/14787210.2018.1522249

Buckley, N. A., Whyte, I. M., O'Connell, D. L., & Dawson, A. H. (1999). Oral or intravenous N-
acetylcysteine: which is the treatment of choice for acetaminophen (paracetamol)
poisoning? Journal of toxicology. Clinical toxicology, 37(6), 759–767.
https://doi.org/10.1081/clt-100102453

Chauhan, A., Lebeaux, D., Decante, B., Kriegel, I., Escande, M. C., Ghigo, J. M., & Beloin, C.
(2012). A rat model of central venous catheter to study establishment of long-term bacterial
biofilm and related acute and chronic infections. PLoS ONE, 7(5).
https://doi.org/10.1371/journal.pone.0037281

Chauhan, A., Lebeaux, D., Ghigo, J. M., & Beloin, C. (2012). Full and broad-spectrum in vivo
eradication of catheter-associated biofilms using gentamicin-EDTA antibiotic lock therapy.
Antimicrobial Agents and Chemotherapy, 56(12), 6310–6318.
https://doi.org/10.1128/AAC.01606-12

Chauhan, A., Sakamoto, C., Ghigo, J. M., & Beloin, C. (2013). Did I pick the right colony? Pitfalls
in the study of regulation of the phase variable antigen 43 adhesin. PloS one, 8(9), e73568.
https://doi.org/10.1371/journal.pone.0073568

Chauhan, A., Bernardin, A., Mussard, W., Kriegel, I., Estève, M., Ghigo, J.-M., Beloin, C., &
Semetey, V. (2014). Preventing Biofilm Formation and Associated Occlusion by Biomimetic
Glycocalyx like Polymer in Central Venous Catheters. The Journal of Infectious Diseases
(Vol. 210, Issue 9). https://www.jstor.org/stable/43708066

Chlumsky, O., Smith, H. J., Parker, A. E., Brileya, K., Wilking, J. N., Purkrtova, S., Michova, H.,
Ulbrich, P., Viktorova, J., & Demnerova, K. (2021). Evaluation of the antimicrobial efficacy
of n-acetyl-l-cysteine, rhamnolipids, and usnic acid—novel approaches to fight food-borne
pathogens. International Journal of Molecular Sciences, 22(21).
https://doi.org/10.3390/ijms222111307

Choi, Y. S., Kim, C., Moon, J. H., & Lee, J. Y. (2018). Removal and killing of multispecies
endodontic biofilms by N-acetylcysteine. Brazilian journal of microbiology, 49(1), 184–188.
https://doi.org/10.1016/j.bjm.2017.04.003

Ciacci, N., Boncompagni, S., Valzano, F., Cariani, L., Aliberti, S., Blasi, F., Pollini, S., Rossolini,
G. M., & Pallecchi, L. (2019). *In vitro* Synergism of Colistin and N-acetylcysteine

against *Stenotrophomonas maltophilia.* Antibiotics (Basel, Switzerland), 8(3), 101. https://doi.org/10.3390/antibiotics8030101

Darouiche R. O. (2004). Treatment of infections associated with surgical implants. The New England journal of medicine, 350(14), 1422–1429. https://doi.org/10.1056/NEJMra035415

Dludla, P. V., Mazibuko-Mbeje, S. E., Nyambuya, T. M., Mxinwa, V., Tiano, L., Marcheggiani, F., Cirilli, I., Louw, J., & Nkambule, B. B. (2019). The beneficial effects of N-acetyl cysteine (NAC) against obesity associated complications: A systematic review of pre-clinical studies. Pharmacological research, 146, 104332. https://doi.org/10.1016/j.phrs.2019.104332

Effah, C. Y., Sun, T., Liu, S., & Wu, Y. (2020). *Klebsiella pneumoniae:* An increasing threat to public health. Annals of Clinical Microbiology and Antimicrobials (Vol. 19, Issue 1). BioMed Central Ltd. https://doi.org/10.1186/s12941-019-0343-8

Feng, J., Liu, B., Xu, J., Wang, Q., Huang, L., Ou, W., Gu, J., Wu, J., Li, S., Zhuo, C., & Zhou, Y. (2018). *In vitro* effects of N-acetylcysteine alone and combined with tigecycline on planktonic cells and biofilms of *Acinetobacter baumannii.* Journal of thoracic disease, 10(1), 212–218. https://doi.org/10.21037/jtd.2017.11.130

Geetha, P. V., Aishwarya, K. V. L., Mariappan, S., & Sekar, U. (2020). Fluoroquinolone Resistance in Clinical Isolates of *Klebsiella pneumoniae*. Journal of Laboratory Physicians, 12(02), 121–125. https://doi.org/10.1055/s-0040-1716478

Ghani, S., Khan, N., Koriyama, C., Akiba, S., & Yamamoto, M. (2014). N-acetyl-L-cysteine reduces arsenite-induced cytotoxicity through chelation in U937 monocytes and macrophages. Molecular medicine reports, 10(6), 2961–2966. https://doi.org/10.3892/mmr.2014.2612

Gomes, A. É. I., Pacheco, T., Santos, C. da S. dos, Pereira, J. A., Ribeiro, M. L., Darrieux, M., & Ferraz, L. F. C. (2021). Functional Insights from KpfR, a New Transcriptional Regulator of Fimbrial Expression That Is Crucial for *Klebsiella pneumoniae* Pathogenicity. Frontiers in Microbiology, 11. https://doi.org/10.3389/fmicb.2020.601921

Guerini, M., Perugini, P., & Grisoli, P. (2020). Evaluation of the Effectiveness of N-Acetylcysteine (NAC) and N-acetylcysteine-cyclodextrins Multi-Composite in *Pseudomonas aeruginosa* Biofilm Formation. Applied Sciences, 10(10), 3466. http://dx.doi.org/10.3390/app10103466

Guilhen, C., Miquel, S., Charbonnel, N., Joseph, L., Carrier, G., Forestier, C., & Balestrino, D. (2019). Colonization and immune modulation properties of *Klebsiella pneumoniae* biofilm-

dispersed cells. Npj Biofilms and Microbiomes, 5(1). https://doi.org/10.1038/s41522-019-0098-1

Guerini, M., Condrò, G., Friuli, V., Maggi, L., & Perugini, P. (2022). N-acetylcysteine (NAC) and Its Role in Clinical Practice Management of Cystic Fibrosis (CF): A Review. Pharmaceuticals (Basel, Switzerland), 15(2), 217. https://doi.org/10.3390/ph15020217

Hughes, H., Helen K., & Kahl, L. (2018). The Harriet Lane handbook: a manual for pediatric house officers.

Heidary, M., Nasiri, M. J., Dabiri, H., & Tarashi, S. (2018). Prevalence of Drug-resistant *Klebsiella pneumoniae* in Iran: A Review Article. Iranian journal of public health, 47(3), 317–326. (Vol. 47, Issue 3).

Igrejas, G., Zhang, J., da Silva Pranchevicius, M.-C., Ferreira, R. L., M da Silva, B. C., Rezende, G. S., Nakamura-Silva, R., Pitondo-Silva, A., Boni Campanini, E., A Brito, M. C., L da Silva, E. M., César de Melo Freire, C., & da Cunha, A. F. (2019). High Prevalence of Multidrug-Resistant *Klebsiella pneumoniae* Harboring Several Virulence and β-Lactamase Encoding Genes in a Brazilian Intensive Care Unit. Frontiers in microbiology, 9, 3198 https://doi.org/10.3389/fmicb.2018.03198

Indian Council of Medical Research. (2020). Antimicrobial Resistance Research and Surveillance Network January 2020 to December 2020. https://main.icmr.nic.in/sites/default/files/guidelines/AMRSN_annual_report_2020.pdf

Jiao, Y., Ma, S., Wang, Y., Li, J., Shan, L., Liu, Q., Liu, Y., Song, Q., Yu, F., Yu, H., Liu, H., Huang, L., & Chen, J. (2016). N-Acetyl Cysteine Depletes Reactive Oxygen Species and Prevents Dental Monomer-Induced Intrinsic Mitochondrial Apoptosis *Invitro* in Human Dental Pulp Cells. PloS one, 11(1), e0147858. https://doi.org/10.1371/journal.pone.0147858

Joseph, L., Merciecca, T., Forestier, C., Balestrino, D., & Miquel, S. (2021). From *Klebsiella pneumoniae* colonization to dissemination: An overview of studies implementing murine models. Microorganisms, 9(6). https://doi.org/10.3390/microorganisms9061282

Jun, Y., Youn, C. K., Jo, E. R., & Cho, S. I. (2019). *In vitro* inhibitory activity of N-acetylcysteine on tympanostomy tube biofilms from methicillin-resistant *Staphylococcus aureus* and quinolone-resistant *Pseudomonas aeruginosa*. International journal of pediatric otorhinolaryngology, 126, 109622. https://doi.org/10.1016/j.ijporl.2019.109622

Kakoullis, L., Papachristodoulou, E., Chra, P., & Panos, G. (2021). Mechanisms of antibiotic resistance in important Gram-positive and Gram-negative pathogens and novel antibiotic solutions. Antibiotics (Vol. 10, Issue 4). https://doi.org/10.3390/antibiotics10040415

Karki, D., Dhungel, B., Bhandari, S., Kunwar, A., Joshi, P. R., Shrestha, B., Rijal, K. R., Ghimire, P., & Banjara, M. R. (2021). Antibiotic resistance and detection of plasmid mediated colistin resistance mcr-1 gene among *Escherichia coli* and *Klebsiella pneumoniae* isolated from clinical samples. Gut Pathogens, 13(1). https://doi.org/10.1186/s13099-021-00441-5

Kundukad, B., Udayakumar, G., Grela, E., Kaur, D., Rice, S. A., Kjelleberg, S., & Doyle, P. S. (2020). Weak acids as an alternative anti-microbial therapy. Biofilm, 2. https://doi.org/10.1016/j.bioflm.2020.100019

Lebeaux, D., Chauhan, A., Rendueles, O., & Beloin, C. (2013). From in vitro to in vivo Models of Bacterial Biofilm-Related Infections. *Pathogens (Basel, Switzerland)*, *2*(2), 288–356. https://doi.org/10.3390/pathogens2020288

Lebeaux, D., Chauhan, A., Létoffé, S., Fischer, F., de Reuse, H., Beloin, C., & Ghigo, J.-M. (2014). pH-Mediated Potentiation of Aminoglycosides Kills Bacterial Persisters and Eradicates *Invivo* Biofilms. In Source: The Journal of Infectious Diseases (Vol. 210, Issue 9). https://about.jstor.org/terms

Lebeaux, D., Fernández-Hidalgo, N., Chauhan, A., Lee, S., Ghigo, J. M., Almirante, B., & Beloin, C. (2014). Management of infections related to totally implantable venous-access ports: challenges and perspectives. The Lancet. Infectious diseases, 14(2), 146–159. https://doi.org/10.1016/S1473-3099(13)70266-4

Lewis K. (2007). Persister cells, dormancy and infectious disease. Nature reviews. Microbiology, 5(1), 48–56. https://doi.org/10.1038/nrmicro1557

Leoney, A., Karthigeyan, S., Asharaf, A. S., & Felix, A. (2020). Detection and Categorization of Biofilm-forming *Staphylococcus aureus, Viridans streptococcus, Klebsiella pneumoniae,* and *Escherichia coli* Isolated from Complete Denture Patients and Visualization Using Scanning Electron Microscopy. Journal of International Society of Preventive & Community Dentistry, 10(5), 627–633. https://doi.org/10.4103/jispcd.JISPCD_256_20

Ligozzi, M., Bernini, C., Bonora, M. G., De Fatima, M., Zuliani, J., & Fontana, R. (2002). Evaluation of the VITEK 2 system for identification and antimicrobial susceptibility testing of

medically relevant Gram-positive cocci. Journal of clinical microbiology, 40(5), 1681–1686. https://doi.org/10.1128/JCM.40.5.1681-1686.2002

Li, X., Kim, J., Wu, J., Ahamed, A. I., Wang, Y., & Martins-Green, M. (2020). N -Acetyl-cysteine and Mechanisms Involved in Resolution of Chronic Wound Biofilm. Journal of Diabetes Research, 2020, 9589507. https://doi.org/10.1155/2020/9589507

Liu, Y., Liu, Y., Dai, J., Liu, A., Li, Y., Xu, J., Yu, X., Liu, J., & Zhu, H. (2021). *Klebsiella pneumoniae* in patients with rheumatic autoimmune diseases: clinical characteristics, antimicrobial resistance and factors associated with extended-spectrum β-lactamase production. BMC Infectious Diseases, 21(1). https://doi.org/10.1186/s12879-021-06055-1

Lopes, E., Saavedra, M. J., Costa, E., de Lencastre, H., Poirel, L., & Aires-de-Sousa, M. (2020). Epidemiology of carbapenemase-producing *Klebsiella pneumoniae* in northern Portugal: Predominance of KPC-2 and OXA-48. Journal of Global Antimicrobial Resistance, 22, 349–353. https://doi.org/10.1016/j.jgar.2020.04.007

Manoharan, A., Das, T., Whiteley, G. S., Glasbey, T., Kriel, F. H., & Manos, J. (2020). The effect of N-acetylcysteine in a combined antibiofilm treatment against antibiotic-resistant *Staphylococcus aureus*. Journal of Antimicrobial Chemotherapy, 75(7), 1787–1798. https://doi.org/10.1093/jac/dkaa093

Manoharan, A., Ognenovska, S., Paino, D., Whiteley, G., Glasbey, T., Kriel, F. H., Farrell, J., Moore, K. H., Manos, J., & Das, T. (2021). N-Acetylcysteine Protects Bladder Epithelial Cells from Bacterial Invasion and Displays Antibiofilm Activity against Urinary Tract Bacterial Pathogens. Antibiotics,10(8), 900. https://doi.org/10.3390/antibiotics10080900

Narimisa, N., Goodarzi, F., & Bavari, S. (2022). Prevalence of colistin resistance of *Klebsiella pneumoniae* isolates in Iran: a systematic review and meta-analysis. Annals of clinical microbiology and antimicrobials, 21(1), 29. https://doi.org/10.1186/s12941-022-00520-8

NIH U.S. National Library of Medicine. (2018). ACETYLCYSTEINE-acetylcysteine injection, solution Fresenius Kabi USA, LLC. Retrieved on August 5, 2022. Available from: https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=d990ef51-ced2-4359-b99b-092334d2dc9b

Nirwan, P. K., Chatterjee, N., Panwar, R., Dudeja, M., & Jaggi, N. (2021). Mutations in two component system (PhoPQ and PmrAB) in colistin resistant *Klebsiella pneumoniae* from

North Indian tertiary care hospital. Journal of Antibiotics, 74(7), 450–457. https://doi.org/10.1038/s41429-021-00417-2

Nirwati, H., Sinanjung, K., Fahrunissa, F., Wijaya, F., Napitupulu, S., Hati, V. P., Hakim, M. S., Meliala, A., Aman, A. T., & Nuryastuti, T. (2019). Biofilm formation and antibiotic resistance of *Klebsiella pneumoniae* isolated from clinical samples in a tertiary care hospital, Klaten, Indonesia. BMC Proceedings. https://doi.org/10.1186/s12919-019-0176-7

Odari, R., & Dawadi, P. (2022). Prevalence of Multidrug-Resistant *Klebsiella pneumoniae* Clinical Isolates in Nepal. In Journal of Tropical Medicine (Vol. 2022). Hindawi Limited. https://doi.org/10.1155/2022/5309350

Olsson, B., Johansson, M., Gabrielsson, J., & Bolme, P. (1988). Pharmacokinetics and bioavailability of reduced and oxidized N-acetylcysteine. European journal of clinical pharmacology, 34(1), 77–82. https://doi.org/10.1007/BF01061422

O'Toole, G. A., & Kolter, R. (1998). Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signaling pathways: a genetic analysis. Molecular microbiology, 28(3), 449–461. https://doi.org/10.1046/j.1365-2958.1998.00797.x

Papadimitriou-Olivgeris, M., Fligou, & F., Bartzavali, & C., Zotou, & A., Spyropoulou, & A., Koutsileou, & K., Vamvakopoulou, & S., Sioulas, & N., Karamouzos, & v, Anastassiou, E. D., Spiliopoulou, I., Christofidou, & M., & Marangos, & M. (2017). Carbapenemase-producing *Klebsiella pneumoniae* bloodstream infection in critically ill patients: risk factors and predictors of mortality. Eur J Clin Microbiol Infect Dis, 36(7), 1125–1131. https://doi.org/10.1007/s10096-017-2899-6

Perry, H. E., & Shannon, M. W. (1998). Efficacy of oral versus intravenous N-acetylcysteine in acetaminophen overdose: results of an open-label, clinical trial. The Journal of pediatrics, 132(1), 149–152. https://doi.org/10.1016/s0022-3476(98)70501-3

Piperaki, E. T., Syrogiannopoulos, G. A., Tzouvelekis, L. S., & Daikos, G. L. (2017). *Klebsiella pneumoniae*: Virulence, Biofilm and Antimicrobial Resistance. Pediatric Infectious Disease Journal, 36(10), 1002–1005. https://doi.org/10.1097/INF.0000000000001675

Zheng, J. X., Lin, Z. W., Chen, C., Chen, Z., Lin, F. J., Wu, Y., Yang, S. Y., Sun, X., Yao, W. M., Li, D. Y., Yu, Z. J., Jin, J. L., Qu, D., & Deng, Q. W. (2018). Biofilm Formation in *Klebsiella pneumoniae* Bacteremia Strains Was Found to be Associated with CC23 and the Presence

of *wcaG*. *Frontiers in cellular and infection microbiology*, *8*, 21. https://doi.org/10.3389/fcimb.2018.00021

Prescott, L. F., Illingworth, R. N., Critchley, J. A., Stewart, M. J., Adam, R. D., & Proudfoot, A. T. (1979). Intravenous N-acetylcysteine: the treatment of choice for paracetamol poisoning. British medical journal, 2(6198), 1097–1100. https://doi.org/10.1136/bmj.2.6198.1097

Quah, S. Y., Wu, S., Lui, J. N., Sum, C. P., & Tan, K. S. (2012). N-acetylcysteine inhibits growth and eradicates biofilm of *Enterococcus faecalis*. Journal of endodontics, 38(1), 81–85. https://doi.org/10.1016/j.joen.2011.10.004

Remya, P. A., Shanthi, M., & Sekar, U. (2019). Characterisation of virulence genes associated with pathogenicity in *Klebsiella pneumoniae*. Indian Journal of Medical Microbiology, 37(2), 210–218. https://doi.org/10.4103/ijmm.IJMM_19_157

Renggli, L., Gasser, M., Plüss-Suard, C., Harbarth, S., & Kronenberg, A. (2022). Temporal and structural patterns of extended-spectrum cephalosporin-resistant *Klebsiella pneumoniae* incidence in Swiss hospitals. Journal of Hospital Infection, 120, 36–42. https://doi.org/10.1016/j.jhin.2021.11.006

Rivera-Sánchez, S. P., Agudelo-Góngora, H. A., Oñate-Garzón, J., Flórez-Elvira, L. J., Correa, A., Londoño, P. A., Londoño-Mosquera, J. D., Aragón-Muriel, A., Polo-Cerón, D., & Ocampo-Ibáñez, I. D. (2020). Antibacterial Activity of a Cationic Antimicrobial Peptide against Multidrug-Resistant Gram-Negative Clinical Isolates and Their Potential Molecular Targets. Molecules, 25(21), 5035. https://doi.org/10.3390/molecules25215035

Rocha, J., Ferreira, C., Mil-Homens, D., Busquets, A., Fialho, A. M., Henriques, I., Gomila, M., & Manaia, C. M. (2022). Third generation cephalosporin resistant *Klebsiella pneumoniae* thriving in patients and in wastewater: what do they have in common? BMC Genomics, 23(1). https://doi.org/10.1186/s12864-021-08279-6

Russo, T. A., & Marr, C. M. (2019). Hypervirulent *Klebsiella pneumoniae*. Clinical microbiology reviews, 32(3), e00001-19. https://doi.org/10.1128/CMR.00001-19

Šalamon, Š., Kramar, B., Marolt, T. P., Poljšak, B., & Milisav, I. (2019). Medical and Dietary Uses of N-Acetylcysteine. Antioxidants, 8(5),111. https://doi.org/10.3390/antiox8050111

Salloum, T., Panossian, B., Bitar, I., Bitar, I., Hrabak, J., Hrabak, J., Araj, G. F., & Tokajian, S. (2020). First report of plasmid-mediated colistin resistance mcr-8.1 gene from a clinical

Klebsiella pneumoniae isolate from Lebanon. Antimicrobial Resistance and Infection Control,
9(1). https://doi.org/10.1186/s13756-020-00759-w

Schwalfenberg G. K. (2021). N-Acetylcysteine: A Review of Clinical Usefulness (an Old Drug
with New Tricks). Journal of nutrition and metabolism, 2021,9949453.
https://doi.org/10.1155/2021/9949453

Shadkam, S., Goli, H. R., Mirzaei, B., Gholami, M., & Ahanjan, M. (2021). Correlation between
antimicrobial resistance and biofilm formation capability among *Klebsiella pneumoniae*
strains isolated from hospitalized patients in Iran. Annals of Clinical Microbiology and
Antimicrobials, 20(1). https://doi.org/10.1186/s12941-021-00418-x

Sharma, A., Chandraker, S., Patel, V. K., & Ramteke, P. (2009). Antibacterial Activity of
Medicinal Plants Against Pathogens causing Complicated Urinary Tract Infections. Indian
journal of pharmaceutical sciences, 71(2), 136–139. https://doi.org/10.4103/0250-474X.54279

Shein, A., Wannigama, D. L., Higgins, P. G., Hurst, C., Abe, S., Hongsing, P., Chantaravisoot, N.,
Saethang, T., Luk-In, S., Liao, T., Nilgate, S., Rirerm, U., Kueakulpattana, N., Laowansiri,
M., Srisakul, S., Muhummudaree, N., Techawiwattanaboon, T., Gan, L., Xu, C., Kupwiwat,
R., Chatsuwan, T. (2021). Novel colistin-EDTA combination for successful eradication of
colistin-resistant *Klebsiella pneumoniae* catheter-related biofilm infections. Scientific
reports, 11(1), 21676. https://doi.org/10.1038/s41598-021-01052-5

Smilkstein, M. J., Bronstein, A. C., Linden, C., Augenstein, W. L., Kulig, K. W., & Rumack, B. H.
(1991). Acetaminophen overdose: a 48-hour intravenous N-acetylcysteine treatment
protocol. Annals of emergency medicine, 20(10), 1058–1063. https://doi.org/10.1016/s0196-
0644(05)81352-6

Stepanović, S., Vuković, D., Hola, V., Di Bonaventura, G., Djukić, S., Cirković, I., & Ruzicka, F.
(2007). Quantification of biofilm in microtiter plates: overview of testing conditions and
practical recommendations for assessment of biofilm production by *staphylococci*. APMIS:
acta pathologica, microbiologica, et immunologica Scandinavica, 115(8), 891–899.
https://doi.org/10.1111/j.1600-0463.2007.apm_630.x

Tenório, M. C. D. S., Graciliano, N. G., Moura, F. A., de Oliveira, A. C. M., & Goulart, M. O. F.
(2021). N-acetylcysteine (NAC): Impacts on human health. Antioxidants (Vol. 10, Issue 6).
https://doi.org/10.3390/antiox10060967

van der Weide, H., Vermeulen-de Jongh, D., van der Meijden, A., Boers, S. A., Kreft, D., Ten Kate, M. T., Falciani, C., Pini, A., Strandh, M., Bakker-Woudenberg, I., Hays, J. P., & Goessens, W. (2019). Antimicrobial activity of two novel antimicrobial peptides AA139 and SET-M33 against clinically and genotypically diverse *Klebsiella pneumoniae* isolates with differing antibiotic resistance profiles. International journal of antimicrobial agents, 54(2), 159–166. https://doi.org/10.1016/j.ijantimicag.2019.05.019

Vasconcelos, N. G., Vaz, M., Radai, J., Kassuya, C., Formagio, A., Graciani, F. S., Leal, M. L., Oliveira, R. J., da Silva, K. E., Croda, J., & Simionatto, S. (2020). Antimicrobial activity of plant extracts against carbapenem-producing *Klebsiella pneumoniae* and *invivo* toxicological assessment. Journal of toxicology and environmental health. Part A, 83(23-24), 719–729. https://doi.org/10.1080/15287394.2020.1824135

Vecchiarelli, A., Dottorini, M., Pietrella, D., Cociani, C., Eslami, A., Todisco, T., & Bistoni, F. (1994). Macrophage activation by N-acetylcysteine in COPD patients. Chest, 105(3), 806–811. https://doi.org/10.1378/chest.105.3.806

Vuotto, C., Longo, F., Pascolini, C., Donelli, G., Balice, M. P., Libori, M. F., Tiracchia, V., Salvia, A., & Varaldo, P. E. (2017). Biofilm formation and antibiotic resistance in *Klebsiella pneumoniae* urinary strains. Journal of applied microbiology, 123(4), 1003–1018. https://doi.org/10.1111/jam.13533

WHO (2017). WHO publishes list of bacteria for which new antibiotics are urgently needed. 27 February 2017 News Release GENEVA. Retrieved April 19, 2022, Available from https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed

WHO (2017). Media Centre. News Release. WHO publishes list of bacteria for which new antibiotics are urgently needed. Retrived on August 28, 2022. Available from: http://www.who.int/mediacentre/news/releases/2017/bacteria-antibiotics-needed/en/

Xie, Y., Tian, L., Li, G., Qu, H., Sun, J., Liang, W., Li, X., Wang, X., Deng, Z., Liu, J., & Ou, H. Y. (2018). Emergence of the third-generation cephalosporin-resistant hypervirulent *Klebsiella pneumoniae* due to the acquisition of a self-transferable bla$_{DHA-1}$-carrying plasmid by an ST23 strain. *Virulence*, *9*(1), 838–844. https://doi.org/10.1080/21505594.2018.1456229

Zhang, F., Lau, S. S., & Monks, T. J. (2011). The cytoprotective effect of N-acetyl-L-cysteine against ROS-induced cytotoxicity is independent of its ability to enhance glutathione

synthesis. Toxicological sciences: an official journal of the Society of Toxicology, 120(1),

87–97. https://doi.org/10.1093/toxsci/kfq364

Zafer, M. M., El-Mahallawy, H. A., Abdulhak, A., Amin, M. A., Al-Agamy, M. H., & Radwan, H.

H. (2019). Emergence of colistin resistance in multidrug-resistant *Klebsiella pneumoniae* and

Escherichia coli strains isolated from cancer patients. Annals of clinical microbiology and

antimicrobials, 18(1), 40. https://doi.org/10.1186/s12941-019-0339-4