Changes driven by the evolution of higher group polarization in the guppy are consistent across different predation pressures and associated with neuroanatomical changes

Alberto Corral-Lopez*[1,2,3,4], Alexander Kotrschal[2,5], Alexander Szorkovszky[6], Maddi Garate-Olaizola[2,4], James Herbert-Read[8,9], Wouter van der Bijl[1], Maksym Romenskyy[10], Hong-Li Zeng[11], Severine Denise Buechel[2,5], Ada Fontrodona-Eslava[2,12], Kristiaan Pelckmans[7], Judith E. Mank[1], Niclas Kolm[2]

*corresponding author: alberto.corral@ebc.uu.se

Affiliations

1 - Department of Zoology and Biodiversity Research Centre, University of British Columbia, Vancouver, Canada

2 - Department of Zoology/Ethology, Stockholm University, Stockholm, Sweden.

3 - Division of Biosciences, University College London, London, United Kingdom

4 – Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden

5- Behavioural Ecology, Wageningen University & Research, Wageningen, Netherlands.

- RITMO Centre for Interdisciplinary Studies in Rhythm, Time and Motion, University of Oslo, Oslo, Norway

- Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden.

- Department of Zoology, University of Cambridge, Cambridge, UK.

- Aquatic Ecology, Lund University, Lund, Sweden.

- Department of Life Sciences, Imperial College London, London, UK.

- School of Science, Nanjing University of Posts and Telecommunications, Nanjing, China.

- Centre for Biological Diversity, School of Biology, University of St Andrews, St Andrews, UK

Abstract

One of the most spectacular displays of social behavior is the synchronized movements that many animal groups perform to travel, forage and escape from predators. However, elucidating the neural mechanisms underlying the evolution of collective behaviors, as well as their fitness effects, remains challenging. Here, we study anti-predator behavior in guppies experimentally selected for divergence in polarization, an important behavioral aspect of coordinated movement. We find that groups from artificially selected lines remain more polarized than control groups in the presence of a threat. Neuroanatomical measurements of polarization-selected individuals indicated changes in brain regions previously suggested to be important regulators of perception, fear and attention, and motor response. Additional visual acuity and temporal resolution tests performed in polarization-selected and control individuals indicate that observed differences in anti-predator and schooling behavior should not be attributable to changes in visual perception, but rather are more likely the result of the more efficient relay of sensory input in the brain of polarization-selected fish. Our findings highlight that brain morphology may play a fundamental role in the evolution of coordinated movement and anti-predator behavior.

Introduction

Animals regularly gather - for safety, for exploiting resources, or for mating. Group-living often leads to spectacular forms of collective behavior, and individuals in many taxa coordinate their movements in order to increase efficiency in foraging and travelling, or to confuse predators[1]. Collective motion has evolved multiple times in fishes and is widely understood as a behavioral adaptation aimed at reducing the risk of predation[2]. This behavioral adaptation is underpinned by the efficient acquisition of information from external cues through visual and lateral line sensory systems[3–6]. To date, we have a detailed understanding that coordinated movements in animal groups likely emerge from decision rules that individuals use to interact in groups[e.g. 7,8]. Similarly, correlation-based analyses have revealed how predation levels are associated with variation in collective motion in wild populations (see for instance [9]). Yet, the causes of collective motion are still unclear, particularly how evolutionary changes in collective motion contribute to anti-predator specific situations, or what type of sensory information and information processing schooling fishes use to identify and avoid predators as groups.

The brain, as the central organ controlling locomotion, sensory systems and decision-making, can play an important role in the ability to coordinate movements and decisions made in the context of grouping. As such, variation in the anatomy of the brain could be an important mechanism behind the evolution of collective motion. The link between social factors and changes in multiple brain structures across taxa is well established[10–13], and well-studied in fishes where approximately half of marine and freshwater species come together in groups at different life stages[14]. For instance, a comparative study on Lake Tanganyikan cichlids showed that species with more complex social structures have larger telencephali and hypothalami[15]. These regions are part of the fish forebrain, which has an important function in social behavior[16], and has been associated with social competence in cleaner fish and social orienting in zebrafish[12,17]. In nine-spined stickleback, exposure to larger groups during development is correlated with larger size of the optic tectum, the visual center in the fish brain[18].

Despite its potential importance, empirical studies explicitly testing the association between neuroanatomy and collective motion are to date scarce. Brain regions associated with fish social behavior have also been identified in the few studies explicitly testing the link between neuroanatomy and schooling behavior. Lesion studies in goldfish (*Carassius auratus*) showed that individuals with ablated telencephalon exhibited reduced activity and association with

conspecifics[19]. Also, a study on surface and cavefish populations of *Astyanax mexicanus* living

in different light environments showed an underlying positive correlation between optic tectum

size and schooling behavior differences between populations[20]. These few previous studies

highlight the potential role of neuroanatomy in schooling, as well as the need to account for

environmental variation in analyses.

Here we use artificial selection lines of guppies (*Poecilia reticulata*) with divergence in

polarization, the degree of alignment of members of a group when swimming. Polarization-

selected guppies offer a unique opportunity to empirically evaluate the link between evolution

of general schooling behavior, specific anti-predator behavior and neuroanatomy[21]. In relation

to many fish species that associate in large schools, guppies have relatively low schooling

propensity, with high levels of variation across individuals and across populations as a function

of external factors such as predation risk or food availability[22,23]. In our selection lines, intrinsic

schooling propensity was increased in female guppies by over 15% compared to controls in

just three generations by selecting on individuals that exhibited higher polarization, the level

of alignment between individuals moving together in a group[21,24]. Previous assays in the

polarization-selected females showed that differences in polarization were caused by the

combined effect of the likelihood to align with neighbors' movements, the strength of

alignment to larger groups and individual swimming speed[21].

We use these lines to investigate potential changes in the visual system and anti-predator

behavior following directional selection for polarization, as well as to study the association

between increased schooling propensity and changes in brain anatomy. For this, we performed

a series of experiments and measurements to: i) evaluate collective motion patterns and

predator inspection of individuals from polarization-selected and control female guppies when

swimming in groups, ii) quantify brain region sizes with microcomputed tomography

(microCT) of female guppies from polarization-selected and control lines, and iii) perform

comprehensive tests of visual capabilities of these fish, spanning eye size, visual acuity and

temporal resolution measurements of individual fish. Through the study of collective behavior

in an ecologically relevant setting, we identify potential evolutionary pathways for how

selection for higher polarization is associated with brain structure size variation.

Results

Collective motion in response to predation threat in guppies following artificial selection

We investigated whether selection for higher polarization affected cohesiveness and how
individuals from groups reacted in response to neighbor movement in a predation context.
These behavioral decisions should have major fitness consequences in this species[25].
Specifically, we recorded and tracked fish in an experimental arena to obtain positional data
and assessed collective motion of groups of eight guppies when exposed to an imminent threat,
the presence of an artificial replica of a pike cichlid (*Crenicichla frenata*), a natural predator in
wild populations of Trinidadian guppies. Furthermore, we exposed these groups to a non-
predator-shaped object to allow for comparisons when presented to a novel object. These
assays were performed in combination with open field tests (OFT's) on the same fish groups.
In total, we obtained data for 83 polarization-selected groups and 81 control groups of
approximately six months of age and of similar body size (polarization-selected: standard
length mean [95% CIs] = 24.8 mm [24.0, 25.5]; control: standard length mean = 24.5 mm [23.8,
25.3]). Previous analyses of the data for OFT's in these groups provided evidence that selection
for polarization altered individuals' speed, how individuals aligned with, and how individuals
were attracted towards conspecifics during group motion[21].

Our analyses of collective motion of female groups exposed to a predator model and a novel
object showed predictable results in relation to previous findings observed in OFT's. In the
presence of a novel object or a predator model, we observed overall declines of 12% and 17%
in the polarization of the groups, while individuals' speed showed 20-fold and 24-fold
decreases in the presence of a novel object and a predator model respectively (Supplementary
Table 1, Fig. 1A). Previously observed differences between polarization-selected and control
groups in these traits were still present in the presence of these stimuli in the experimental arena
with approximate differences of 7% (Supplementary Table 2, Fig. 1A). Overall, nearest
neighbor distance towards conspecifics in female showed a 10-fold increase in tests with a
novel object and with a predator model in relation to OFT's (Supplementary Table 1, Fig. 1A).
Such strong reduction in median nearest neighbor distance observed between fish in the
presence of these stimuli likely lead to minimum possible values of inter-individual distance in
our experimental setup. Indeed, differences between polarization-selected and control lines in
nearest neighbor distance observed in OFT's[21] were not present in the presence of a predator
model and a novel object (Fig. 1A; Supplementary Table 2). For median speed and group

polarization, the predator model elicited a stronger response than the novel object. Specifically,

collective motion data showed that guppy groups showed a 4-fold reduction in speed and were

4.5% less aligned in the presence of a predator model (LMM$_{speed}$: Estimate$_{novel\ object\ vs\ predator\ model}$

= 3.910 [0.709-7.100], t = 4.71; df = 338; p < 0.001; LMM$_{polarization}$: Estimate$_{novel\ object\ vs\ predator}$

$_{model}$ = 0.045 [0.022-0.067], t = 2.87; df = 338; p = 0.011: Supplementary Table 1, Fig. 1A).

Information processing in response to predation threat

Group polarization spatial patterns

We studied spatial patterns associated with collective motion in our experimental setup by

generating heatmaps with average values of group polarization and group centroid within a 20

x 20 mm grid for OFT's, predator model and novel object tests performed to polarization-

selected and control groups. Visual inspection of heatmaps (Fig. 1B) concords with results

obtained from statistical models using summary statistics for each group, suggesting a decline

in polarization of the group in the presence of these stimuli. Yet, differences in polarization

between control and polarization-selected groups in open field and novel object assays were

consistent across all regions of the arena, while the observed differences in assays with a

predator model were more pronounced in positions further away from the head of the model

(Fig. 1B).

Correlation between group polarization and speed

To further characterize potential differences in how information spreads in polarization-

selected and control fish groups we assessed the effect of group speed in the alignment of fish

when exposed to a predation threat. For this, we generated heatmaps showing the correlation

of group speed and group polarization across treatments for our different selection lines (Fig.

2). Heatmaps indicated that group polarization of polarization-selected fish in OFTs was

constantly close to maximal values of one and differences with control groups were partially

independent of the speed of the group (despite larger values of median speed observed in

relation to those in control groups). Overall group polarization reduction in the presence of a

novel object and a predator model in relation to OFTs were likely associated with a strong

reduction in the speed of the group, including periods of no movement of the group. Periods of

no movement in response to a threat were observed more frequently in control groups

suggesting that they might be an important driver of group polarization differences observed

between polarization-selected and control groups. In addition to differences driven by lack of

movement, correlations of group polarization and speed indicated that differences between
polarization-selected and control groups observed during collective motion resembling motion
in OFT's (mean group speed range 2 – 5 mm/frame) were maintained in the presence of stimuli
in the experimental arena (Fig. 2).

Predator inspection behavior
Group polarization spatial patterns and its relationship to swimming speed suggested
differences between polarization-selected and control fish in their attention to stimuli presented
in the experimental arena. As such, we quantified inspection behavior of individuals in our
groups of eight fish in the presence of a predator model. We scored recorded videos for the
start and end point for each predator inspection performed by one randomly selected fish in
each video. Analyses of predator inspection data showed that females from control lines
presented a higher tendency to inspect the predation threat presented in the experimental arena
than polarization-selected females. Specifically, we observed that polarization-selected
females spent 21% less time inspecting the predator model and the mean duration of predator
inspections were 18% shorter in polarization-selected females (GLMM$_{time\_inspecting}$: Incidence
Rate Ratio (IRR) $_{polarization}$ = 0.79 (0.66-0.95), t = -2.52, p = 0.011; GLMM$_{mean\_inspection}$: IRR
$_{polarization}$ = 0.82 (0.66-0.96), t = -2.49, p = 0.013, Fig. 2d-e, Supplementary Table 3): We
observed a similar trend for total number of predator inspections performed with a 13% reduced
frequency of inspections observed in polarization-selected females (GLMM$_{inspections}$: IRR
$_{polarization}$ = 0.87 (0.75-1.01), t = -1.85, p = 0.064; Fig. 2c, Supplementary Table 3).

Collective motion during predator inspections
Analysis of positional data and median distance to the stimulus presented in our assays
suggested that most inspection behaviors to the predator model were performed during the
initial three minutes of the assays (Supplementary Figure 1). Further, the majority of
inspections were performed at a range <200 mm and in the tail area of the predator model
presented in the experimental arena (Fig. 1A; Fig. 3A). Consequently, we filtered our data to
evaluate collective motion patterns of fish groups in the time and locations where predator
inspections were performed during our assays (see methods). The overall differences in group
polarization found between polarization-selected and control groups were maintained in areas
within 200 mm of the predator model (Fig. 3B; LMM$_{polarization< 200mm}$: Estimate$_{selection}$ = -0.044
[-0.088,-0.000], t = -1.984, df = 272, p = 0.048; Supplementary Table 4a-b). Inspection
behavior is mainly performed from areas with reduced risk of attack from a predator. In line

with such expectation, we found that polarization of all groups was greatly reduced in the area
of the predator model tail (Fig. 3B). However, we found no differences in group polarization
between selected and control females in the head area of the predator model, but a maintenance
of 8% differences in group polarization between polarization-selected and control females in
close proximity to the tail of the predator model (LMM$_{\text{polarization – head area}}$: Estimate$_{\text{selection}}$ = -
0.037 [-0.102, 0.027], t = -1.53, df = 4.58, p = 0.190; LMM$_{\text{polarization – tail area}}$: Estimate$_{\text{selection}}$ =
-0.080 [-0.146, -0.013], t = -1.53, df = -3.48, p = 0.030; Fig. 3B, Supplementary Table 5a-b).
246

247 **Changes in neuroanatomy following artificial selection**
248

We used microcomputed tomography (micro-CT) to reconstruct the brain anatomy of 13
polarization-selected females, and 15 control females and determine overall brain volume and
the volumes of 11 major brain regions that could be safely identified in these brain scans
covering the whole brain volume: olfactory bulbs, ventral telencephalon, dorsal telencephalon,
thalamus, hypothalamus, nucleus glomerulus, torus semicircularis, optic tectum cup, central
optic tectum, cerebellum, and medulla oblongata (see methods). We next used this data set to
evaluate potential associations between collective behavior and neuroanatomy. Polarization-
selected and control fish showed no differences in relative brain size (whole brain volume in
relation to their body size; LMM$_{\text{relativebrain}}$: t = -0.41, df = 23.29, p = 0.682; Fig. 4a). However,
analyses of relative brain region size (volume of each region in relation to the volume of the
rest of the brain) indicated that the thalamus and optic tectum cups are approximately 7% and
4% larger in polarization-selected than control females respectively (LMM$_{\text{thalamus}}$: Odds Ratio
(OR) $_{\text{polarization}}$ = 0.929 (0.866-0.998); t = 2.187, df = 25, p = 0.038; LMM$_{\text{o.tectum}}$: Odds Ratio
(OR) $_{\text{polarization}}$ = 0.959 (0.924-0.994); t = 2.409, df = 23.09, p = 0.024; Fig. 4a), and the medulla
oblongata is an 8% larger in control females (LMM$_{\text{medulla}}$: Odds Ratio (OR) $_{\text{polarization}}$ =1.08
(1.01-1.14); t = -2.65, df = 23.91, p = 0.013; Fig. 4a). All other eight brain regions measured
presented no differences between polarization-selected and control females in relative region
volumes (Fig. 4a-b, Supplementary Table 6).
267

In parallel, we analyzed brain region volume differences using a more conservative approach
and found similar and consistent differences between selection lines. Specifically, we used a
multivariate Bayesian model that included the relative size of the 11 brain regions as dependent
variables. Posterior samples drawn from the multivariate model indicated that confidence
intervals for the difference in relative volume in the medulla oblongata, the optic tectum cups

and the thalamus between polarization-selected and control females did not overlap with zero

(Supplementary Table 7a).

We then used the multivariate Bayesian model to evaluate the correlation in relative brain

region volume between multiple regions measured. We focused on evaluating correlations with

other brain regions for the three regions significantly differentiated between lines following

artificial selection (Supplementary Table 7b). We found no correlation between optic tectum

cup relative volume and volume of any other region measured. However, we found a significant

inverse correlation between thalamus and medulla relative volume (rescorr$_{Medulla-Thalamus}$ [95%

CIs]: -0.40 [-0.65 / -0.12]). This finding suggests that the opposite differences observed in the

volume of these two brain regions between control and polarization-selected female guppies

may be linked to changes in brain development processes associated with artificial selection

for higher coordinated motion.

Information acquisition through the visual system

Efficiently acquiring information through the sensory system, mainly through visual cues, is a

basic principle of collective motion in shoaling fishes[3]. Given observed differences in the size

of the optic tectum cups between polarization-selected and control fish, we investigated

potential differences in visual perception between lines. For this, we compared eye size and

two key characteristics of the visual system to track movement of conspecifics, visual acuity

and temporal resolution.

Eye size

We quantified eye size in females from polarization-selected (n = 57, standard length mean

[95% CIs] = 27.0 mm [26.6, 27.4]) and control lines (n = 55, standard length mean [95% CIs]

= 27.1mm [26.8, 27.5]). Eye size is a common indicator of visual capacities of organisms[26],

and comparative studies across fish species suggest that larger eyes correlate with improved

visual abilities[27]. In our study, we found no difference between polarization-selected and

control lines in either absolute eye size or relative eye size, the proportional size of the eye in

relation to body size (Fig. 5A; LMM$_{eye size}$: Estimate$_{selection}$ = -0.03 [-0.24, 0.19], t = -0.52, df =

2, p = 0.66; LMM$_{relative eye size}$: Estimate$_{selection}$ = -0.00 [-0.01, 0.01], t = -0.13, df = 2, p = 0.90;

Supplementary Table 8). As previously reported in guppies[28] ,we found a strong positive

correlation between eye size and body size (Pearson's correlation test: t = 11.72, df = 110, p <

0.001, cor [95% CIs] = 0.74 [0.65, 0.82]), but this correlation did not differ between

polarization-selected and control lines (LMM$_{eye\ size}$: Estimate$_{selection*body\ size}$ = 0.010 [-0.016,

0.039], t = 0.75, df = 101, p = 0.45).

Visual acuity

We further assessed potential differences in visual perception between selection and control

lines by quantifying visual acuity in the same individuals for which eye size was measured.

Visual acuity allows an individual to resolve spatial detail and can be critical for an organism's

fitness[29]. We measured visual acuity in our fish by quantifying their innate optomotor response

in contrasting rotating gratings. This a widely used method to study visual acuity in multiple

fish species, including guppies[30–32], and we have previously used this approach to evaluate the

visual system of guppies in similar contexts[28,33]. Following the methods in ([28]), we exposed our

fish to a series of six stimuli with rotating and static gratings of different widths at the lower

end of the known guppy visual acuity, where thinner widths are more difficult to perceive. If

polarization-selected and control groups differed in their ability to resolve spatial ability,

differences in their optomotor response towards these rotating stimuli at the lower end of the

guppy visual acuity should be expected. However, we found no difference in their average

optomotor response combining data from all stimuli (LMM$_{acuity}$: Estimate$_{selection}$ = 0.00 [-0.08,

0.009], t = 0.11, df = 12.88, p = 0.913; Supplementary Table S9a), or in analyses independently

evaluating specific optomotor response for any of the six stimuli presented (Fig. 5B;

Supplementary Table 9c).

Visual resolution tracking movement

Although the ability to resolve spatial detail, acuity, is arguably an important visual parameter

for guppies to recognize conspecific positions in shoals, it provides no information on an

individual's ability to track movement[34]. Similar to many social fish species, guppies swim

with a saltatory movement style that features discrete changes in speed and direction[9].

Consequently, we implemented an additional experiment that evaluated potential differences

between polarization-selected and control fish in their temporal assessment of speed and

direction changes. Using the same experimental apparatus used to evaluate visual acuity, we

video recorded female guppies from our selection and control lines and exposed them to a

single-width rotating stimulus that moved in multiple directions and at different speeds (see

methods). We next used automated tracking to obtain orientation and speed of the fish for each

frame and to quantify their direction and speed in relation to the stimuli presented at each time

point. If polarization-selected and control groups differed in their ability to track movement,

differences in the time following the correct direction of the stimuli and the deviation of their
speed in relation to the stimulus speed should be expected.

Overall, fish followed the correct direction of the rotating stimulus for a significant proportion
of the time. This was the case when the stimuli were presented in both a clockwise direction
an in counterclockwise direction (Supplementary Figure 2, Supplementary Table 10). Overall,
swimming speed did not significantly deviate from the stimuli rotating speed at the two lowest
speed levels (Supplementary Figure 2, Supplementary Table 10), but was less than the stimuli
speed at the two highest speed levels (Supplementary Figure 2, Supplementary Table 10). This
was true for both directions in which stimuli were presented, but the mismatch between
swimming and stimuli rotation speed was greater at the higher speed when the stimulus rotated
anticlockwise (Supplementary Figure 2, Supplementary Table 10).

We compared the performance of polarization-selected and control fish in the test to evaluate
their visual temporal resolution while shoaling. We found no differences between polarization-
selected and control lines in their deviation of their swimming speed in relation to the stimuli
rotating speed for their combined scores across speeds and direction of rotation (Fig. 5C;
$LMM_{speed\ deviation}$: $Estimate_{selection}$ = -0.46 [-8.74, 7.83], t = -0.46, df = 2.64, p = 0.863;
Supplementary Table 10a), or for their speed observed at any particular speed at direction of
rotation (Supplementary Figure 2, Supplementary Table 10a). Similarly, polarization-selected
and control females spent similar proportions of time following the stimuli during changes in
stimuli rotating speed (Supplementary Figure 2; Supplementary Table 10a; $LMM_{proportion\ time}$:
$Estimate_{selection}$ = 0.00 [-0.14, 0.15], t = 0.10, df = 2.52, p = 0.928).

Discussion

366
Our work demonstrates that selection for schooling behavior in female guppies has important
implications for how this species behaves in the presence of potential threat. Analyses of
motion patterns in these fish show that polarization-selected groups maintain higher
polarization and speed when exposed to a potential threat. In addition, our analyses indicate
that individuals from polarization-selected groups spend less time inspecting the threat than
individuals from control lines. We further studied visual capacities in these fish and find no
differences between polarization-selected and control individuals in visual acuity, temporal
resolution or eye size, suggesting that effective processing of visual information in the brain is

key for the observed differences in their ability to synchronize their swimming with close

neighbors at multiple contexts.

In parallel, our results suggest that artificial selection for higher polarization has produced

significant changes in the brain anatomy of female guppies. Neuroanatomical measurements

indicate that polarization-selected fish exhibit a larger thalamus and a large optic tectum cup,

but a smaller medulla oblongata, compared to control fish. These rapid changes in brain region

sizes in response to selection for polarization behavior are consistent with previous artificial

selection directly on neuroanatomy, which resulted in rapid shifts in both relative brain size

and relative telencephalon size, in just a few generations in guppies[35,36].

Below, we discuss the implications of these discoveries for our understanding on how the

association between brain morphology and anti-predator behavior might drive the evolution of

collective behavior.

Information processing in a predation threat context

Speed plays a prominent role in structuring fish schools, as individuals adjust their distance

and alignment with neighbours through changes in speed[21,37,38]. Indeed, empirical studies

across several species, as well as modelling approaches, have recently demonstrated a positive

correlation between individual's speed and group polarization[38–43]. In our experiments, the

presence of a novel object and a predator model prompted strong reductions in the median

speed of individuals in the test. Yet, similar differences between polarization-selected and

control females in individual's median speed and polarization were maintained across all

contexts. Swimming performance tests previously performed in these fish demonstrated that

selection did not target fish with better physiological capacities to swim faster, but those

showing higher speed in response to neighbor positions[21]. Taken together, our results from

these selection lines suggest that selection for coordinated movement is largely mediated by

individuals' speed, and that group coordination characteristics such as polarization can emerge

from individuals' tendencies to group with conspecifics and fitness benefits acquired from such

tendencies at specific contexts.

Schooling in fishes is widely understood as a behavioral adaptation to escape the effect of

predation[1]. These synchronized movements have been shown to confer two major benefits to

schools, facilitating escape through transfer of information from closer neighbors[44], or by

confusing the predator in which individual to attack[45]. The use of a static predator model in our

assays does not allow us to infer any potential benefit of higher polarization on the confusion

effect towards predators. However, our results show that directional selection and associated

changes in the brain are correlated to behavioral changes in the presence of a threat and that it

might affect individual ability to acquire private information about a potential predator. The

reduced time spent inspecting the predator model by polarization-selected females, coupled

with the fact that polarization-selected groups remained more polarized closer to the predator

model, especially around the tail of the predator, suggest this a likely possibility. However,

further comparisons within asocial and social contexts should be implemented to disregard the

alternative explanation that directional selection leads to changes in predator inspection

behavior also when fish have no access to social information.

Our study only measured fitness effects indirectly, using a predator model. Yet, previous work

demonstrated that shorter inspection times towards the same predator models are associated

with higher survival in the species[46,47]. Experiments with virtual prey analyses have also shown

that individuals in groups with higher polarization are at lower risk of predation[45]. As such, our

results following directional selection for polarization are consistent with the idea that selection

acting on how individuals interact during motion might confer important fitness benefits for

individuals showing higher coordination with conspecifics and reinforced in specific contexts,

such as high-predation environments. Yet, such benefits might trade-off with a reduced level

of private information obtained by individuals in relation to potential threats or foraging

sources. In sticklebacks less sociable individuals are more likely to discover food first in a

foraging context[37]. This agrees with our results in experiments with a predator model, where

faster individuals' speed and higher tendencies to group with conspecifics seem to drive the

observed lower time inspecting the predator model in the arena and acquire private information

on a potential threat. Overall, these factors are arguably important selective pressures in natural

populations where guppies from high predation habitats swim with higher coordination and in

larger groups[39,48]. Indeed, guppies from higher predation populations have been shown to rely

less on private information for foraging resources, than those from lower predation

populations[49]. Further studies evaluating fitness effects of relying in social versus private

information across different predation pressures is paramount to understand how anti-predator

behavior and collective motion drive evolutionary patterns at the proximate level.

Brain morphology and collective behavior

We found two brain regions that were larger in relation to the rest of the brain in polarization-
selected fish, the optic tectum cup and the thalamus. The optic tectum is the terminus of a vast
majority of optic nerve fibers and axons of retinal cells[50] and as such is the primary vertebrate
visual center. Despite wide variation in optic tectum size across teleost species, this region is
known to control eyes and head movement in fish, as well as forming instant representations
of the immediate surroundings to transform visual cues into behavioral responses[50,51]. This
function is primarily achieved in superficial layers of the tectum[52], which corresponds to the
optic tectum cup region used in our neuroanatomical parcellation of major brain regions in the
guppy[53].

The evolved differences in optic tectum cup size between polarization-selected and control
female guppies we found are concordant with phenotypic plasticity findings in nine-spined
stickleback where it was found that individuals reared in groups developed larger optic tectum
that those reared individually[18]. Differences in the ability to acquire sensory input have
previously been associated with differences in schooling propensity[54]. In our experiment, rapid
evolution of higher polarization did not lead to changes in visual performance or eye size.
Together, these findings suggest that differences found between polarization-selected and
control lines in this section of the optic tectum should have an effect in their ability to control
body orientation during complex social maneuvers such as predator evasion and swimming
with low inter-individual distances, but not in sensory information acquisition. This is
consistent with current considerations of the tectum as more than a visuomotor relay, with
implementation of intrinsic processing of visual information to extract critical features and send
processed outputs to the posterior thalamus and then to further decision-making regions of the
brain[51].

The primary receiving region of processed inputs from the tectum, the thalamus, was also found
to be enlarged in polarization-selected females. To date, the level of homology between the
mammalian and teleost thalamic regions is under strong debate[55,56]. In mammals, the thalamus
plays critical roles in the modification, filtering and distribution of sensory and
mechanosensory information into decision-making regions of the brain[56,57]. Recent findings in
zebrafish suggest that these tasks are performed mainly by the preglomerular complex, whose
developmental origin might be outside the diencephalic area that contains the thalamus in
mammals[58]. The neuroanatomical atlas we created to analyze relative region size does not

allow us to assess whether this particular region is part of the thalamus area which has been
enlarged in polarization-selected females. However, our results of strong gene expression
changes in the telencephalon of polarization-selected and control females in response to
decision-making across social contexts[59], are in agreement with a higher efficiency in the relay
of processed information towards the telencephalon, potentially facilitated by an enlarged
thalamus. Studies across teleosts and mammals do agree on that the thalamus have an important
role in the development of glutamatergic and GABAergic relay nuclei, with crucial roles in
neurotransmission and inhibition. Interestingly, functional characterization of genomic and
transcriptomic analyses performed in these selection lines identified a crucial role of
glutamatergic synaptic function in the evolution of schooling[59]. Taken together, our results
suggest that changes in the development of glutamatergic relay nuclei driven by a larger
thalamus might affect the ability of individuals to synchronize their movements in a shoal. In
parallel, and while not directly addressed in this study, the regulatory role of the thalamus in
aggressiveness is concordant with common expectations of lower aggression levels in group-
living species ([reviewed in 60]). As such, further quantifications of the anatomical characteristics of
the thalamus in relation to aggression levels within and across species might be a promising
avenue for future research.

In contrast to the thalamus and the optic tectum cup, we found that the medulla oblongata was
smaller in polarization-selected lines. The medulla oblongata is an important relay center of
nervous signals between the spinal cord and ascendant brain regions with well-described
functions in teleosts. First, it has an important role in basic motor function through processing
of mechanosensory stimuli from hydrodynamic information[61]. Consistently found across fish
species, the lateral line nerves terminate in the medial and caudal octavo lateralis nucleus in
the medulla[62]. As the lateral line is crucial for schooling through cues that allow fishes to assess
neighbor changes in speed and direction, the reduced relative size of the medulla in
polarization-selected lines could be associated with potential differences in the ability to
integrate and process information received through these nerves. Yet, our previous studies
evaluating motor control capabilities found no difference between polarization-selected and
control female guppies[21,63]. Further studies evaluating information processing of
mechanosensory input in these selection lines is nonetheless paramount, with special focus on
low light and high turbidity conditions.

Finally, studies in fishes indicate that the medulla has an important role for processing
somatosensory signals, with special emphasis in auditory and gustatory signals[64,65].
Specifically, the vagal lobe, part of the medulla, is described to have a prominent role in
processing taste. Consistent with our results, the vagal lobe of larval reef fishes is larger in
solitary as compared to more social species[66]. While not tested in this study, it may be that the
reduction in medulla observed in polarization-selected lines might be associated with important
changes in the auditory system and the ability to perceive different tastes. In line with this
reduction in the size of the medulla oblongata, our results show that three other brain regions
have significant hypoallometric relationships with the medulla (see Supplementary Table 4b):
the cerebellum (motor control center), the thalamus and the hypothalamus (hormonal
regulation center). Gene expression of *angiopoietin*-1, a locus implicated in brain tissue
development, showed contrasting expression levels between the medulla and the thalamic and
hypothalamic regions[67]. Based on this, we hypothesize that selection for more coordinated
motion leads to a trade-off between general sensory capabilities that are not important in
coordinated movements and specific sensory capabilities required to coordinate movement
with neighbors. In the future, it will be interesting to investigate the association between
schooling propensity, brain anatomy and potential trade-offs between sensory and
mechanosensory capacities.
528

Conclusion

Our empirical approach with behavioral assays on artificial selection lines with divergence in
polarization show that collective motion differences are consistent in the presence of a predator
threat and that predator inspection behavior varies between the selection lines and the control
lines. Moreover, we reveal differences in neuroanatomy that could provide a mechanistic
explanation to the observed behavioral differences. Based on our discoveries, we propose that
changes in behavior are intimately intertwined with matching changes in brain morphology
during the evolution of collective behavior.
538

Methods

<u>Artificial selection for higher group polarization</u>

We evaluated the association between brain anatomy and collective motion in female guppies

following artificial selection for higher polarization. Extensive detail on the selection procedure

can be found in ([21]). In short, groups of female guppies were tested in repeated open field tests

and sorted in relation to the mean polarization of the group, the degree to which the individuals

of a group move with higher alignment[21,24,68]. For three generations, females from groups with

higher polarization were bred with males from those cohorts to generate three up-selected

polarization lines. In parallel, random females were exposed to the same experimental

conditions and bred with unselected males to generate three control lines. Third generation

polarization-selected females presented on average a 15% higher polarization, 26% higher

median speed and 10% higher group cohesiveness (i.e. 10% shorter nearest neighbor distances)

in comparison to control females[21]. Further tests in these lines showed that selection for higher

directional coordination was not only driven for selection for faster moving fish. This is

because polarization-selected lines were still 5.7% more polarized after statistically controlling

for speed differences, and the distance to conspecifics was an important factor driving

differences in speed between polarization-selected and control lines[21]. The selection procedure

targeted polarization on female groups and we found a weaker response to selection in males,

and therefore subsequent neuroanatomical, behavioral and physiological studies focused on

females. All fish were removed from their parental tanks after birth, separated by sex at the

first onset of sexual maturation, and afterwards transferred to single-sex groups of eight

individuals in seven liter tanks. We kept all fish used for anti-predator response experiments,

visual capacity tests and brain morphology measurements in these groups throughout their life

span. Fish were not reused for experiments across these three categories. All tanks contained 2

cm of gravel with continuously aerated water, a biological filter, and plants for environmental

enrichment. We allowed for visual contact between the tanks. The laboratory was maintained

at 26°C with a 12-h light:12-h dark schedule. Fish were fed a diet of flake food and freshly

hatched brine shrimp daily.

Anti-predator response in guppies following artificial selection

Collective motion

We evaluated anti-predator behavior in polarization-selected and control female guppies by

conducting assays on 164 groups of eight fish in white arenas with 55 cm diameter and 3 cm

water depth (polarization-selected: n = 83, standard length mean [95% CIs] = 24.8 mm [24.0,

25.5]); control: n = 81, standard length mean [95% CIs] =24.5 mm [23.8, 25.3]). Each group

was initially assessed in an open field assay in the arena for ten minutes, and collective motion

data from these open field assays was previously used to analyze differences in social

interactions[21]. After ten minutes, we sequentially introduced a novel object and a predator

model for 6-minute periods in the centre of the experimental arena. In half the assays, we

introduced the novel object first and the predator model second, with the order reversed in the

other half of the assays. We used a blue coffee mug as a novel object and a fishing lure (18 x

3 cm) custom-painted to resemble the pike cichlid Crenicichla frenata, a natural predator on

the guppy, as the predator model. These objects have been previously used to successfully

reproduce natural behaviors of the guppy in response to a novel object and a predation threat[47].

To facilitate automated data collection, the position and orientation of the predator model was

kept constant by using magnets in the bottom of the fishing lure and in the central position of

the experimental arena. We provided external illumination to avoid shadowed areas in the

circular arena. Disturbances were minimized by performing all assays in an isolated room of

the laboratory. Under these conditions the presence of stimuli should be a major driver of the

spatial locations of fish in the experimental arena. However, we are unable to quantify the

effect of arbitrary spatial preferences in our results. Prior to the start of the assay, the eight-fish

group was confined in the centre of the arena for two minutes in an opaque white 15 cm PVC

cylinder. After this acclimation period, we lifted the cylinder and filmed the arena using a Point

Grey Grasshopper 3 camera (FLIR Systems; resolution, 2048 pixels by 2048 pixels; frame rate,

25 Hz).

We tracked the movement of fish groups in the collected video recordings using idTracker[69]

and used fine-grained tracking data to calculate speed, polarization and nearest neighbor

distance in Matlab 2020 following methods established in ([70]). For speed, we obtained the

median speed across all group members by calculating the first derivatives of the x and y time

series, then smoothed using a third-order Savitzky–Golay filter. For nearest neighbor

distance, we obtained the median distance to the nearest neighbor for every fish across all

frames. For group polarization, we calculated the median global alignment, which indicates

the angular alignment of all fish in the arena. Calculations of median global alignment in each

frame were only calculated if six out of the eight members of the group presented tracks

following the optimization of our tracking protocol in the setup in ([24]). No differences

between completeness of tracks were observed between polarization-selected and control

groups for any of the treatments in our experiment (Supplementary Figure 3). For all

measurements trials with less than 70% complete tracks were disregarded for further

analyses. We additionally used R (v4.1.3), RStudio (v2022.07.0) and the tidyverse package[71–

73] to generate heatmaps with average values of group polarization and group centroid within

20 x 20 mm grid cells for each frame (Fig. 1B). To avoid biases in group centroid

measurements potentially resulting from single individuals or small subgroups at large

distances, we limited the analyses to frames in which at least six individuals formed a

connected group, with an interindividual distance of 10cm counting as a connection. Grid

cells that did not contain values for a minimum of 8 groups per treatment were disregarded.

No differences between the proportion of frames used were observed between polarization-

selected and control groups for any of the treatments (Supplementary Figure 3). To evenly

compare motion patterns when presented with a novel object and a predator model to those

obtained during the open field assays, we limited our analysis of the open field assay data to

the initial six minutes of the recording.

We used LMMs with median speed, polarization and nearest neighbor distance as dependent

variables to test for potential differences between polarization-selected and control lines.

Selection regime, the type of stimulus presented and the interaction between these two were

included as fixed effects, and body size of fish was coded as a covariate, with a random

intercept for each replicated selection line and the order of presentation of stimuli as random

factors. All models were run in R (v4.1.3) and RStudio (v2022.07.0)[71,73] using lme4 and

lmerTest packages[74,75]. Model diagnostics showed that residual distributions were roughly

normal with no evidence of heteroscedasticity. We obtained post-hoc comparisons of the

response between selection line regimes at different levels of other fixed effects in the previous

models using the emmeans package[76] with the tukey-adjustment method for multiple

comparisons.

Predator inspection behavior

Behavioral scoring. Positional data and analyses of median distance to center in our data

indicated that groups of fish swam closer to the stimuli presented in the initial minutes

following the addition of a predator model in the experimental arena, when compared to the

same time periods following the addition of a novel object (Fig. 1A; Fig. 3A; Supplementary

Figure 1). This observation matched previous findings in similar experiments performed on

guppies[47] and likely corresponds to the stereotypical behavioral response of guppies to inspect

and gain information of a potential threat[25]. A predator inspection in guppies is characterized

by an approach to the predator, monitoring predator activity and swimming sideways with an

arched body. Based on this information, we manually visualized the videos during the first

three minutes after addition of the predator model and scored the behavior of one randomly

selected fish in the group using BORIS[77]. While blind to the selection line treatment, the start

and end time of each predator inspection performed by the focal fish was scored for each

video. We used the start and end time of predator inspections to calculate the number of

inspections, average inspection duration and the total time that was spent inspecting per fish.

Next, we fit a statistical model for each variable as a dependent variable using a zero-inflated

beta distribution and a logit link function for the conditional mean in the package glmmTMb[78].

We used the selection line regime as a fixed effect. A random intercept for each replicated

selection line, and the order of presentation of the stimuli in the arena were included as random

factors in the model. We evaluated the adequacy of our fitted model using scaled-residuals

quantile-quantile plots, residual versus predicted values plots and a zero-inflation test in the

DHARMa package[79].

Group collective motion during predator inspections. We analyzed positional data for each

group by binning the observations in a grid, with 20 x 20 mm cells. For each trial, we calculated

a density map, where the value for each grid cell was the fraction of all observations that

occurred within that cell. The resulting density maps are a normalized representation on how

often each grid cell was visited by individuals in our groups of eight fish when exposed to

different stimuli (Fig. 2A). We used information from positional data to calculate summary

statistics in different areas of interest. Predator model assays presented unique spatial patterns

in areas closer to the stimulus presented, with higher densities in the tail area of the predator

model (Fig. 2A). Based on these factors, we calculated two new summary variables for each

group: i) median polarization of the group when the average position of the group was <200

mm to the predator model and novel object assays; and ii) median polarization of the group in

locations closer to the head (y-position > 0) and the tail (y-position < 0) in predator model

assays. We used LMMs with these new calculated variables as dependent variables in the

model to test for potential differences between polarization-selected and control lines.

Selection regime, the type of stimulus presented or location in the tank were included as fixed

effects, with a random intercept for each replicated selection line and the order of presentation

of stimuli as random factors. [74,75]Model diagnostics showed that residual distributions were

roughly normal with no evidence of heteroscedasticity. We obtained post-hoc comparisons of

the response between selection line regimes at different levels of other fixed effects in the

previous models using the emmeans package[76] with the tukey-adjustment method for multiple
comparisons.
680
<u>Brain morphology of female guppies following artificial selection</u>
We assessed neuroanatomical features of polarization-selected and control F3 females,
approximately six months old (polarization-selected: n = 15, standard length mean [95% CIs]
= 238 mm [235, 242]; control: n = 15, standard length mean [95% CIs] = 237 mm [232, 241]).
We euthanized animals with an overdose of benzocaine and fixated the whole fish in 2%
glutaraldehyde and 4% paraformaldehyde in phosphate buffered saline (PBS) for five days.
Following two PBS washes, the brains were dissected out and stained for 48 h in 1%
osmiumtetraoxide. We embedded the stained brains in 3% agar and scanned them using
microcomputed tomography (microCT, Skyscan 1172, Bruker microCT, Kontich,
Belgium). The scanner operated at a voltage of 80 kV, a current of 125 μm, with a 0.5 mm
aluminum filter. Images were acquired using an isotropic pixel size of 2.4 μm. We
reconstructed cross-sections from scanned images following a in NRecon (Bruker microCT)
following a protocol successfully implemented in a previous study evaluating neuroanatomical
differences between guppies up- and down-selected for relative brain size[53]. This protocol
allowed us to obtain measurements of whole brain size volume and brain region volume in 11
major brain regions in the guppy brain: olfactory bulbs, ventral telencephalon, dorsal
telencephalon, thalamus, hypothalamus, nucleus glomerulus, torus semicircularis, optic tectum
cup, central optic tectum, cerebellum, and medulla oblongata (Fig. 4B). We chose all regions
that we could safely identify in the brain scans based on an adult swordtail brain atlas[80] and our
own knowledge in fish neuroanatomy. Extended details on guppy brain region reconstruction
from digital images can be found in ([53]). Two brains from polarization-selected lines were
damaged during the protocol, which reduced the sample size to 28 samples.
703
We tested for overall differences in relative brain size between polarization-selected and
control lines using a linear mixed model (LMM) with brain volume as dependent variable,
body size (standard length) as covariate, selection regime as fixed effect, and replicate as
random effect. For each brain region, we used two different approaches to determine whether
polarization-selected and control lines differ in relative brain region size. First, we used 11
independent LMMs with each region's volume as dependent variable, whole brain volume
(excluding the volume of the region of interest) as covariate, selection regime as fixed effect,
and replicate as random effect. LMMs were implemented in in R (v4.1.3) and RStudio

(v2022.07.0)[71,73] using lme4 and lmerTest packages[74,75]. Second, to take into consideration that

brain region volumes may be interdependent, we used a more conservative approach and

analyzed the data using a Bayesian multilevel model that included 11 brain regions as

dependent variables in a fully multivariate context. The full model included an analogous

structure to those used in the independent LMMs for each brain region. Parameter values were

estimated using the brms interface[81,82] to the probabilistic programming language Stan[83]. We

used default prior distributions with student-t distribution (3, 0, 2.5) for all parameters. The

model estimated residual correlations among all brain region volumes with a Lewandowski-

Kurowicka-Joe (LKJ) prior with $\eta = 1$, which is uniform over the range -1 to 1. Posterior

distributions were obtained using Stan's no-U-turn HMC with six independent Markov chains

of 4000 iterations, discarding the first 2000 iterations per chain as warm-up and resulting in

12000 posterior samples overall. Convergence of the six chains and sufficient sampling of

posterior distributions were confirmed by a scale reduction factor below 1.01, and an effective

size of at least 10% of the number of iterations. For each model, posterior samples were

summarized on the basis of the Bayesian point estimate (median), SE (median absolute

deviation), and posterior uncertainty intervals by HDIs.

Visual information processing in response to predation threat

Visual acuity

We evaluated the ability to perceive detail (visual acuity) in 9-12 months old female guppies

(polarization-selected: n = 59, standard length [95% CIs] = 270 mm [266, 274], control: n =

57, standard length [95% CIs] = 271mm [268, 275]). For this, we assessed their optomotor

response, an innate orient behavior induced by whole-field visual stimulation[84], a widely used

method to assess visual acuity across tax ([reviewed in 85]). Extended methods and the optimization

procedure for the stimuli used here can be found in ([28]). Briefly, we projected a video recording

with rotating vertical black and white bands of six different widths (stimuli) on the walls of a

white ring-shaped arena of 25/50 cm of inner/outer diameter. Previous optimization of the

methods found that the use of these stimuli allowed us to evaluate the optomotor response at

the lower end of the species' acuity[28]. We placed individual fish in between the inner wall of

the arena and a transparent ring of 40 cm diameter. After a two-minute acclimation period, we

recorded their response towards six different rotating stimuli and the static images of these

stimuli using a Sony Cam HDR-DR11E recorder. Each stimulus was presented for one minute

in random order. We manually scored the videos, recording the time that fish spent circling in

the direction of the stimuli (clockwise) at a constant moving pattern using BORIS[77]. Behavioral

scoring was performed blind to the treatment since only running numbers identified recordings.

Likewise, scoring was blind to the rotation and bandwidth of the stripes since only the fish, but

not the rotating stimuli, were visible during scoring. From the scoring, we calculated the

proportion of time that a fish spent swimming in the direction of rotation of the stimuli, out of

the total time that the different vertical black and white bands were presented to them.

Temporal resolution

Two weeks after visual acuity tests were completed in all fish, we measured the ability to track

movement stimuli of different speeds (temporal resolution) in the same females from the

polarization-selected (n = 58) and control lines (n= 55). We did not keep track of fish identity

as fish were kept in groups with conspecifics of the same selection line and replicate between

experiments. To evaluate temporal resolution, we placed fish in a white arena (50 cm diameter,

4 cm water depth) and exposed them to a projection of black and white bands of 3.5 cm width

on the walls rotating clockwise and counterclockwise at four different angular speeds (14.4,

25, 36 and 45 degrees/sec). We chose this bandwidth to focus on the assessment of fish

response towards direction and speed changes, as this stimulus showed a maximal optomotor

response in prior visual acuity tests[28]. The movement of each individual was recorded for a

total of 1380 seconds with a Sony Cam HDR-DR11E; a 300 seconds acclimation period and

1080 seconds of clockwise and counterclockwise rotation of vertical bands at multiple speeds.

Specifically, during each individual test, the 8 stimulus combinations (4 speeds, 2 directions)

were presented separately five independent times for 23 sec. The total time of each individual

test was 920 sec (23 sec per stimulus x 5 times during the test x 8 stimulus combinations). We

randomized the order of presentation of different stimuli a priori, but this order was consistent

for all fish. The stimulus changed speed with smooth transitions of 3 sec, accelerating or

decelerating to the next speed.

To quantify speed and direction changes of fish in our experimental setup, we automated

behavioral scoring and obtained positional data using the Loopy Deep Learning Module

(Loopbio 2020) in MATLAB (v. 2020a). X and y coordinates were transformed into a polar

coordinate system centered on 0 and estimated from positional data. We calculated fish

orientation by taking the difference in the fish's position between frames and defined their

relative orientation (with respect to the arena) with the arcsin ($\sin(\theta - \vartheta)$), where θ was the

orientation of the fish and ϑ is the angle of the arena radius going through the fish position.
Positive values represent a fish swimming clockwise around the arena, while negative values
represent swimming counterclockwise. For each frame, we identified whether the fish was
swimming in the same direction of the stimulus projected and calculated the total proportion
of time swimming in the direction of the stimulus. We also calculated the speed (in degrees per
second) of the fish at each frame by using the dot product of the
positional vector between consecutive frames. Using these values, we calculated for each
individual the average total speed for each of the stimuli presented, and the average speed
deviation between the speed of the stimulus presented and the speed of the fish.
788

Eye size
After the temporal resolution experiments were completed, we measured eye size in the
females from polarization-selected lines (n = 57) and control lines (n =55) that were previously
assessed for visual acuity and temporal resolution. For morphological measurements, we
anesthetized fish with 0.2 mg/l of benzocaine and took pictures of their left side. We measured
eye diameter and body length in these pictures using ImageJ[86]. Relative eye size was calculated
as the ratio of these two variables. Image analyses were performed by a single scorer who was
blind to the selection line treatment in the photographs.
797

Statistics
Analyses were conducted in in R (v4.1.3) and RStudio (v2022.07.0)[71,73]. We analyzed
potential differences in optomotor response, temporal resolution and eye size between
polarization-selected and control females using LMM's. For the visual acuity trials, the
proportion of time rotating was the dependent variable of the model. Fixed effects included
selection line regime and bandwidth of the rotating stimuli. To account for differences in
activity between fish, we used the proportion of time moving when presented with a static
image of the stimuli as a covariate in the model. A random intercept for each replicated
selection line, identity of the fish, and an observation-level variable were included as random
factors in the model. For temporal resolution, we used selection line regime, the speed of
rotation and the direction of rotation as fixed effects. The full model included the interaction
between the selection regime with both speed and direction of rotation. This model included
the identity of the fish, and a random intercept for each replicated selection line as random
factors.
812

For eye size, eye diameter and relative eye size were dependent variables and models included
selection regime as a fixed effect and a random intercept for each replicated selection line as a
random factor. Model diagnostics showed that residual distributions were roughly normal with
no signs of heteroscedasticity in optomotor response and eye size analyses. Model diagnostics
on both models for temporal resolution analyses indicated unequal residual variance across the
range of predicted values and a potential unequal influence of outliers. While estimates in linear
mixed-effects models (LMMs) are argued to be robust to violations of such assumptions[87], we
used the robustlmm package to compare the estimates obtained with LMM's to robust models
with the same predictors that provide reduced weights to outliers in the data[88]. Results were
consistent regarding the modelling approach (Supplementary Table 11). We obtained post-hoc
comparisons of the response between selection line regimes at different levels of other fixed
effects in the previous models using the emmeans package[76] with the tukey-adjustment method
for multiple comparisons.

826
827

Acknowledgements

We thank D. Sumpter for important contributions to the conceptualization of the artificial
selection procedure. We thank two anonymous reviewers for insightful comments in earlier
versions of this manuscript. We thank A. Rennie, E. Trejo, M. Amcoff and A. Boussard for
help with fish husbandry. **Funding**: This work was supported by the Knut and Alice
Wallenberg Foundation (102 2013.0072 to N.K., and K.P.), the Canada 150 Research Chair
Program, and the European Research Council (680951 to J.E.M.), the Swedish Research
Council (2016-03435 to N.K., 2017-04957 to A.K., and 2018-04076 to J.H-R.), and the
Whitten Lectureship in Marine Biology, University of Cambridge (to J.H-R.). **Author**
contributions: N.K., K.P and J.E.M. contributed to conceptualization and funding acquisition
of the project. A.K., A.S., M.R., S.D.B., J.H-R., H.Z., K.P., and N.K. contributed to the design
of the selection procedure. A.C-L., A.K., A.S., A.F-E., M.R., S.D.B. conducted research to
obtain collective behavior data and predator inspection data. A.K. H.Z. and K.P. conducted
research to obtain neuroanatomical data. A.C-L., M.G-O., and J.H-R. conducted research to
obtain visual capacities data. A. C-L. and W.vdB. performed formal analyses and visualization.
A.C-L., J.E.M, and N.K. wrote the original draft. All authors contributed to the final version
of the manuscript. **Competing interests**: The authors declare that they have no competing
interests. **Data and materials availability**: All data and code needed to evaluate the

conclusions in the paper will be deposited in an online repository. Additional data related to this paper may be requested from the authors.

References

1. Krause, J. & Ruxton, G. D. *Living in Groups*. (Oxford University press, 2002).
2. Pitcher, T. J. Functions of Shoaling Behaviour in Teleosts. in *The Behaviour of Teleost Fishes* (ed. Pitcher T. J.) 294–337 (Springer, Boston, MA, 1986).
3. Strandburg-Peshkin, A. *et al.* Visual sensory networks and effective information transfer in animal groups. *Current biology* **23**, R711 (2013).
4. Pitcher, T. J. Shoaling and Schooling in Fishes Evaluations of compliance with the FAO (UN) Code of Conduct for Responsible Fisheries View project Haida Gwaii ecosystem models View project. (1998).
5. Partridge, B. L. & Pitcher, T. J. The sensory basis of fish schools: Relative roles of lateral line and vision. *Journal of Comparative Physiology □ A* **135**, 315–325 (1980).
6. Blaxter, J. H. S., Gray, J. A. B. & Denton, E. J. Sound and Startle Responses in Herring Shoals. *Journal of the Marine Biological Association of the United Kingdom* **61**, 851–869 (1981).
7. Couzin, I. D., Krause, J., Franks, N. R. & Levin, S. A. Effective leadership and decision-making in animal groups on the move. *Nature* **433**, 513–516 (2005).
8. Parrish, J. K. & Edelstein-Keshet, L. Complexity, Pattern, and Evolutionary Trade-Offs in Animal Aggregation. *Science (1979)* **284**, 99–101 (1999).
9. Herbert-Read, J. E. *et al.* How predation shapes the social interaction rules of shoaling fish. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20171126 (2017).
10. Barton, R. A. Neocortex size and behavioural ecology in primates. *Proceedings of the Royal Society B: Biological Sciences* **263**, 173–177 (1996).
11. Burish, M. J., Kueh, H. Y. & Wang, S. S. H. Brain Architecture and Social Complexity in Modern and Ancient Birds. *Brain Behav Evol* **63**, 107–124 (2004).
12. Triki, Z., Levorato, E., McNeely, W., Marshall, J. & Bshary, R. Population densities predict forebrain size variation in the cleaner fish Labroides dimidiatus. *Proceedings of the Royal Society B* **286**, 20192108 (2019).
13. Chee, S. S. A. *et al.* Social status, breeding state, and GnRH soma size in convict cichlids (Cryptoheros nigrofasciatus). *Behavioural Brain Research* **237**, 318–324 (2013).
14. Shaw, E. Schooling Fishes: The school, a truly egalitarian form of organization in which all members of the group are alike in influence, offers substantial benefits to its participants . *Am Sci* **2**, 166–175 (1978).
15. Pollen, A. A. *et al.* Environmental complexity and social organization sculpt the brain in Lake Tanganyikan cichlid fish. *Brain Behav Evol* **70**, 21–39 (2007).
16. O'Connell, L. A. & Hofmann, H. A. Genes, hormones, and circuits: An integrative approach to study the evolution of social behavior. *Front Neuroendocrinol* **32**, 320–335 (2011).
17. Stednitz, S. J. *et al.* Forebrain Control of Behaviorally Driven Social Orienting in Zebrafish. *Current biology* **28**, 2445-2451.e3 (2018).
18. Gonda, A., Herczeg, G. & Merilä, J. Habitat-dependent and -independent plastic responses to social environment in the nine-spined stickleback (Pungitius pungitius) brain. *Proceedings of the Royal Society B: Biological Sciences* **276**, 2085–2092 (2009).

19.   Shinozuka, K. & Watanabe, S. Effects of telencephalic ablation on shoaling behavior
in goldfish. *Physiol Behav* **81**, 141–148 (2004).

20.   Loomis, C. *et al.* An Adult Brain Atlas Reveals Broad Neuroanatomical Changes in
Independently Evolved Populations of Mexican Cavefish. *Front Neuroanat* **13**, 88
(2019).

21.   Kotrschal, A. *et al.* Rapid evolution of coordinated and collective movement in
response to artificial selection. *Sci Adv* **6**, eaba3148 (2020).

22.   Magurran Anne E. *Evolutionary Ecology: The Trinidadian Guppy*. (Oxford University
Press, 2005).

23.   Bierbach, D. *et al.* Using a robotic fish to investigate individual differences in social
responsiveness in the guppy. *R Soc Open Sci* **5**, 181026 (2018).

24.   Szorkovszky, A. *et al.* Assortative interactions revealed by sorting of animal groups.
Anim Behav **142**, 165–179 (2018).

25.   Dugatkin, L. A. & Godin, J. G. J. Predator inspection, shoaling and foraging under
predation hazard in the Trinidadian guppy, Poecilia reticulata. *Environ Biol Fishes* **34**,
265–276 (1992).

26.   Veilleux, C. C. & Kirk, E. C. Visual Acuity in Mammals: Effects of Eye Size and
Ecology. *Brain Behav Evol* **83**, 43–53 (2014).

27.   Caves, E. M., Sutton, T. T. & Johnsen, S. Visual acuity in ray-finned fishes correlates
with eye size and habitat. *Journal of Experimental Biology* **220**, 1586–1596 (2017).

28.   Corral-López, A., Garate-Olaizola, M., Buechel, S. D., Kolm, N. & Kotrschal, A. On
the role of body size, brain size, and eye size in visual acuity. *Behav Ecol Sociobiol* **71**,
1–10 (2017).

29.   Caves, E. M., Brandley, N. C. & Johnsen, S. Visual Acuity and the Evolution of
Signals. *Trends in Ecology and Evolution* vol. 33 358–372 Preprint at
https://doi.org/10.1016/j.tree.2018.03.001 (2018).

30.   Neave, D. A. The development of visual acuity in larval plaice (Pleuronectes platessa
L.) and turbot (Scophthalmusmaximus L.). *J Exp Mar Biol Ecol* **78**, 167–175 (1984).

31.   Anstis, S., Hutahajan, P. & Cavanagh, P. Optomotor test for wavelength sensitivity in
guppyfish (Poecilia reticulata). *Vision Res* **38**, 45–53 (1998).

32.   Haug, M. F., Biehlmaier, O., Mueller, K. P. & Neuhauss, S. C. F. Visual acuity in
larval zebrafish: Behavior and histology. *Front Zool* **7**, 1–7 (2010).

33.   Corral-López, A. *et al.* Female brain size affects the assessment of male attractiveness
during mate choice. *Sci Adv* **3**, e1601990 (2017).

34.   Cronin, T. W., Johnsen, S., Marshall, N. J. & Warrant, E. J. *Visual Ecology. Visual*
Ecology (Princeton University Press, 2014).

35.   Kotrschal, A. *et al.* Artificial Selection on Relative Brain Size in the Guppy Reveals
Costs and Benefits of Evolving a Larger Brain. *Current Biology* **23**, 168–171 (2013).

36.   Fong, S. *et al.* Rapid mosaic brain evolution under artificial selection for relative
telencephalon size in the guppy (Poecilia reticulata). *Sci Adv* **7**, eabj4314 (2021).

37.   Jolles, J. W., Boogert, N. J., Sridhar, V. H., Couzin, I. D. & Manica, A. Consistent
Individual Differences Drive Collective Behavior and Group Functioning of Schooling
Fish. *Current Biology* **27**, (2017).

38.   Kent, M. I. A., Lukeman, R., Lizier, J. T. & Ward, A. J. W. Speed-mediated properties
of schooling. *R Soc Open Sci* **6**, (2019).

39.   Sumpter, D. J. T., Szorkovszky, A., Kotrschal, A., Kolm, N. & Herbert-Read, J. E.
Using activity and sociability to characterize collective motion. *Philosophical*
Transactions of the Royal Society B: Biological Sciences **373**, 20170015 (2018).

40. Mishra, S., Tunstrøm, K., Couzin, I. D. & Huepe, C. Collective dynamics of self-
propelled particles with variable speed. *Phys Rev E Stat Nonlin Soft Matter Phys* **86**,
011901 (2012).
41. Klamser, P. P. *et al.* Impact of Variable Speed on Collective Movement of Animal
Groups. *Front Phys* **9**, 715996 (2021).
42. Pettit, B., Ákos, Z., Vicsek, T. & Biro, D. Speed Determines Leadership and
Leadership Determines Learning during Pigeon Flocking. *Current Biology* **25**, 3132–
3137 (2015).
43. Jolles, J. W. *et al.* Group-level patterns emerge from individual speed as revealed by
an extremely social robotic fish. *Biol Lett* **16**, (2020).
44. Handegard, N. O. *et al.* The dynamics of coordinated group hunting and collective
information transfer among schooling prey. *Current Biology* **22**, 1213–1217 (2012).
45. Ioannou, C. C., Guttal, V. & Couzin, I. D. Predatory fish select for coordinated
collective motion in virtual prey. *Science (1979)* **337**, 1212–1215 (2012).
46. Kotrschal, A. *et al.* Brain size affects female but not male survival under predation
threat. *Ecol Lett* **18**, 646–652 (2015).
47. van der Bijl, W., Thyselius, M., Kotrschal, A. & Kolm, N. Brain size affects the
behavioural response to predators in female guppies (Poecilia reticulata). *Proceedings
of the Royal Society B: Biological Sciences* **282**, 20151132 (2015).
48. Magurran, A. E., Seghers, B. H., Carvalho, G. R. & Shaw, P. W. Behavioural
consequences of an artificial introduction of guppies (Poecilia reticulata) in N.
Trinidad: Evidence for the evolution of anti-predator behaviour in the wild.
Proceedings of the Royal Society B: Biological Sciences **248**, 117–122 (1992).
49. Clément, R. J. G. *et al.* Collective decision making in guppies: A cross-population
comparison study in the wild. *Behavioral Ecology* **28**, 919–924 (2017).
50. Northmore, D. P. M. The Optic Tectum . *Encyclopedia of fish physiology: From
genome to environment* **1**, 131–142 (2011).
51. Isa, T., Marquez-Legorreta, E., Grillner, S. & Scott, E. K. The tectum/superior
colliculus as the vertebrate solution for spatial sensory integration and action. *Current
Biology* **31**, R741–R762 (2021).
52. Jones, M. R., Grillner, S. & Robertson, B. Selective projection patterns from subtypes
of retinal ganglion cells to tectum and pretectum: Distribution and relation to behavior.
Journal of Comparative Neurology **517**, 257–275 (2009).
53. Kotrschal, A. *et al.* Evolution of brain region volumes during artificial selection for
relative brain size. *Evolution (N Y)* **71**, 2942–2951 (2017).
54. Kowalko, J. E. *et al.* Loss of schooling behavior in cavefish through sight-dependent
and sight-independent mechanisms. *Current Biology* **23**, 1874–1883 (2013).
55. Wullimann, M. F. Neural origins of basal diencephalon in teleost fishes: Radial versus
tangential migration. *J Morphol* **281**, 1133–1141 (2020).
56. Mueller, T. What is the Thalamus in Zebrafish? *Front Neurosci* **6**, 64 (2012).
57. Goldberg, J. H., Farries, M. A. & Fee, M. S. Basal ganglia output to the thalamus: Still
a paradox. *Trends Neurosci* **36**, 695–705 (2013).
58. Bloch, S. *et al.* Non-thalamic origin of zebrafish sensory nuclei implies convergent
evolution of visual pathways in amniotes and teleosts. *Elife* **9**, 1–27 (2020).
59. Corral-Lopez, A. *et al.* Functional convergence of genomic and transcriptomic genetic
architecture underlying sociability in a live-bearing fish. *bioRxiv* 2023.02.13.528353
(2023) doi:10.1101/2023.02.13.528353.
60. Kutsukake, N. Complexity, dynamics and diversity of sociality in group-living
mammals. *Ecological Research* **24**, 521–531 (2009).
61. Bleckmann, H. The Lateral Line System of Fish. *Fish Physiology* **25**, 411–453 (2006).

62. Puzdrowski, R. L. Peripheral distribution and central projections of the lateral-line
nerves in goldfish, Carassius auratus. *Brain Behav Evol* **34**, 110–131 (1989).

63. Vega-Trejo, R. *et al.* Artificial selection for schooling behaviour and its effects on
associative learning abilities. *Journal of Experimental Biology* **223**, 235093 (2020).

64. Kozloski, J. & Crawford, J. D. Transformations of an auditory temporal code in the
medulla of a sound- producing fish. *Journal of Neuroscience* **20**, 2400–2408 (2000).

65. Finger, T. E. Evolution of gustatory reflex systems in the brainstems of fishes. *Integr*
Zool **4**, 53–63 (2009).

66. Lecchini, D. *et al.* Variation in brain organization of coral reef fish larvae according to
life history traits. *Brain Behav Evol* **83**, 17–30 (2014).

67. Chen, Y. C. *et al.* Expression change in angiopoietin-1 underlies change in relative
brain size in fish. *Proceedings of the Royal Society B: Biological Sciences* **282**,
20150872 (2015).

68. Szorkovszky, A. *et al.* An efficient method for sorting and quantifying individual
social traits based on group-level behaviour. *Methods Ecol Evol* **8**, 1735–1744 (2017).

69. Pérez-Escudero, A., Vicente-Page, J., Hinz, R. C., Arganda, S. & De Polavieja, G. G.
idTracker: tracking individuals in a group by automatic identification of unmarked
animals. *Nature Methods 2014 11:7* **11**, 743–748 (2014).

70. Kotrschal, A. *et al.* Brain size does not impact shoaling dynamics in unfamiliar groups
of guppies (Poecilia reticulata). *Behavioural Processes* **147**, 13–20 (2018).

71. Posit team. RStudio: Integrated Development Environment for R. Posit Software,
PBC, Boston, MA. URL http://www.posit.co/ (2023).

72. Wickham, H. *et al.* Welcome to the Tidyverse. *J Open Source Softw* **4**, 1686 (2019).

73. R Core Team. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/
(2021).

74. Bates, D., Sarkar, D., Bates, M. D. & Matrix, L. The lme4 Package. *R package version*
2, 74 (2007).

75. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in
Linear Mixed Effects Models. *J Stat Softw* **82**, 1–26 (2017).

76. Lenth, R. Package 'lsmeans'. *Am Stat* **34**, 216–221 (2018).

77. Friard, O. & Gamba, M. BORIS: a free, versatile open-source event-logging software
for video/audio coding and live observations. *Methods Ecol Evol* **7**, 1325–1330 (2016).

78. Brooks, M. E. *et al.* glmmTMB balances speed and flexibility among packages for
zero-inflated generalized linear mixed modeling. *R J* **9**, 378–400 (2017).

79. Hartig, F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed)
regression models. *R package version 0.3* (2018).

80. Anken, R. H. & Rahmann, H. Brain atlas of the adult swordtail fish, Xiphophorus
helleri, and of certain developmental stages. 88 (1994).

81. Bürkner, P. C. Advanced Bayesian multilevel modeling with the R package brms. *R J*
10, 395–411 (2018).

82. Bürkner, P. C. brms: An R package for Bayesian multilevel models using Stan. *J Stat*
Softw **80**, 1–28 (2017).

83. Carpenter, B. *et al.* Stan: A probabilistic programming language. *J Stat Softw* **76**,
(2017).

84. Smith, K. U. & Bojar, S. The nature of optokinetic reactions in mammals and their
significance in the experimental analysis of the neural mechanisms of visual functions.
Psychol Bull **35**, 193–219 (1938).

85. Caves, E. M., Troscianko, J. & Kelley, L. A. A customizable, low-cost optomotor apparatus: A powerful tool for behaviourally measuring visual capability. *Methods Ecol Evol* **11**, 1319–1324 (2020).

86. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods 2012 9:7* **9**, 671–675 (2012).

87. Schielzeth, H. *et al.* Robustness of linear mixed-effects models to violations of distributional assumptions. *Methods Ecol Evol* **11**, 1141–1152 (2020).

88. Koller, M. Robustlmm: An R package for Robust estimation of linear Mixed-Effects models. *J Stat Softw* **75**, 1–24 (2016).

Figures

Fig 1. Collective motion patterns in female guppies artificially selected for higher
polarization. A) Boxplots of median polarization, speed and nearest neighbor distance for
groups of eight individuals of polarization-selected (pink) and control (blue) female guppies
assayed in an open field test (OFT), with a novel object and with a predator model. Horizontal
lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points
within 1.5 times the interquartile range. Asterisks indicate p < 0.05 (see methods;
Supplementary Tables 1-2). **B)** Heatmaps of group polarization across different locations of
the experimental arena when control (top) and polarization-selected (bottom) groups were
exposed to open field, novel object and predator model assays. Grid cells with data for less
than 8 groups were not depicted.

1110
Fig 2. Correlation between group polarization and speed. Heatmaps showing the association
between group polarization and speed in control (top) and polarization-selected (bottom) groups of eight
guppy females exposed to open field, novel object and predator model assays. Group speed depicts the
mean median swimming speed of the individuals in a group in mm per frame. Heatmaps indicate that
the overall decrease in polarization in the presence of a novel object and a predator model was associated
with a reduction in speed (higher relative density of group speed < 2 mm/frame). The consistent
differences in polarization across treatments between polarization-selected and control groups seem to
be driven by: i) lower reduction in mean group speed in polarization-selected groups, including periods
of no movement (area delimited by blue arrows): and ii) consistent differences in motion characteristics
regardless of the presence of stimuli in the assay (lower range of group polarization at group speed >2
mm/frame, area delimited by red arrows).
1122
1123

1124
Fig 3. Predator inspection behavior and collective motion. Boxplots of number of predator
inspections **(A)**, total time inspecting **(B),** and mean inspection duration **(C)** for individuals
when swimming in a group of eight polarization-selected (pink) and control (blue) female
guppies in the presence of a predator model. Horizontal lines indicate medians, boxes indicate
the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range.
Asterisks indicate p < 0.05 (see methods, Supplementary Tables 3-5). **(D)** Density maps based
on positional data of control (top) and polarization-selected (bottom) groups exposed to open
field, novel object and predator model assays. **(E)** Boxplots of median group polarization in
locations closer than 200 mm of the stimulus presented (left) and in the head and tail area of a
predator model (right) in control (blue) and polarization-selected (pink) guppy groups.
1135
1136

1137
1138
1139
Fig 4. The effect of artificial selection for higher polarization in neuroanatomical allometric
relationships. (A) The top left panel shows the allometric relationship between whole brain size volume
and standard length of the fish (SL). Remaining panels show the relationship between each separate
brain region with the rest of the brain ordered rostrally to caudally. Asterisks indicate brain regions with
non-overlapping confidence intervals between polarization-selected females (pink; n = 13) and control
females (blue; n = 15) in two consistent statistical analyses (Supplementary Tables 6-7). **(B)**
Reconstructed brain regions from microCT - scanned guppy brains. A dorsal (left) and lateral (right)
view of a guppy brain with the major brain regions color coded: olfactory bulbs (OB; dark blue), dorsal
telencephalon (DT; red), ventral telencephalon (VT; light green), optic tectum (OT; yellow),
hypothalamus (Hyp; turquoise), thalamus (Th; purple), cerebellum (Cb; brown), medulla oblongata
(MO; dark green). Not visible but measured brain regions include: torus semicircularis, nucleus
glomerulus and optic tectum core.

Fig 5. Eye size and visual capacities of female guppies artificially selected for higher polarization. **(A)** Boxplots of eye morphological measurements. **(B)** Boxplots and density plots of the proportion of time following 6 different rotating stimuli with rotating and static gratings of different widths at the lower end of guppy visual acuity (thinner widths represent a higher degree of difficulty to be perceived). **(C)** Boxplots and density plots of the deviation of fish swimming speed in relation to the speed that a rotating stimulus presented. in polarization-selected. For all morphological measurements and vision assays we measured the same polarization-selected (pink; n = 57-59) and control females (blue; n = 55-57). In all boxplots, horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. Optomotor response average values not sharing any letter are significantly different ($p < 0.05$) in post-hoc contrasts (see Supplementary Table 9b). No significant differences were observed for any comparison between control and polarization-selected fish (see Supplementary Tables 8-10).

1179 <u>**Supplementary Figures for:**</u>
1180
1181 **Changes driven by the evolution of higher group polarization in the guppy are consistent**
1182 **across different predation pressures and associated with neuroanatomical changes**
1183

1184 Alberto Corral-Lopez*, Alexander Kotrschal, Alexander Szorkovszky, Maddi Garate-
1185 Olaizola, James Herbert-Read, Wouter van der Bijl, Maksym Romenskyy, Hong-Li Zeng,
1186 Severine Denise Buechel, Ada Fontrodona-Eslava, Kristian Pelckmans, Judith E. Mank, Niclas
1187 Kolm
1188
1189 *corresponding author: alberto.corral@ebc.uu.se

1190
1191

1192
1193
1194 **Fig S1. Proximity to stimuli in female guppies artificially selected for higher polarization.**
1195 Boxplots of median distance to the stimulus combining data for groups of polarization-selected and
1196 control female guppies (n = 164) in predator model and novel object assays. Horizontal lines indicate
1197 medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the
1198 interquartile range. Asterisks indicate p < 0.05.
1199
1200

1201
1202
Fig S2. Temporal resolution of female guppies artificially selected for higher polarization.
Boxplots and density plots of the deviation of fish swimming speed **(A),** and the proportion of time that
fish followed the direction of the stimulus **(B)** in relation to four different rotating stimuli presented at
different speeds that rotated clockwise and anti-clockwise to polarization-selected (pink; n = 58) and
control females (blue; n = 56). In the boxplots, horizontal lines indicate medians, boxes indicate the
interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. No
significant differences were observed for any comparison between control and polarization-selected
fish (see Tables S9-S10).
1211
1212

1213
1214 **Fig S3. Tracking data reliability between groups.** We found no differences between polarization-
1215 selected groups (pink) and control groups (blue) in **(A)** the total proportion of frames tracked by
1216 idTracker in open field, novel object and predator model assays, or in **(B)** the data used to generate
1217 heatmaps that only considered frames in which at least six individuals formed a connected group, with
1218 an interindividual distance of 10cm counting as a connection.

Supplementary Tables for:

Changes driven by the evolution of higher group polarization in the guppy are consistent across different predation pressures and associated with neuroanatomical changes

Alberto Corral-Lopez*, Alexander Kotrschal, Alexander Szorkovszky, Maddi Garate-Olaizola, James Herbert-Read, Wouter van der Bijl, Maksym Romenskyy, Hong-Li Zeng, Severine Denise Buechel, Ada Fontrodona-Eslava, Kristiaan Pelckmans, Judith E. Mank, Niclas Kolm

*corresponding author: alberto.corral@ebc.su.se

Table S1. Statistical tests for overall comparisons in tests evaluating polarization-selected and control female guppies in their shoaling patterns when exposed to an open field test (OFT), a novel object (cup) and a predator model.

Polarization

Predictors	Estimates	CI	t	p	df
(Intercept)	0.80	0.68 – 0.91	14.54	**< 0.001**	17.59
Selection [P]	0.08	0.02 – 0.13	3.90	**0.017**	3.98
treatment [Cup]	-0.12	-0.15 – -0.10	-9.16	**< 0.001**	338.42
treatment [Predator]	-0.17	-0.20 – -0.14	-12.61	**< 0.001**	340.73
Body size	-0.00	-0.00 – -0.00	-2.35	**0.021**	88.01
Selection [P] * treatment [Cup]	-0.01	-0.05 – 0.03	-0.46	0.644	340.46
Selection [S] * treatment [Predator]	-0.00	-0.04 – 0.03	-0.18	0.860	341.08

Median speed

Predictors	Estimates	CI	t	p	df
(Intercept)	63.25	46.15 – 80.34	7.50	**< 0.001**	35.87
Selection [P]	13.63	8.09 – 19.16	5.96	**< 0.001**	6.29
treatment [Cup]	-20.14	-23.91 – -16.37	-10.51	**< 0.001**	337.87

treatment [Predator]	-24.34	-28.14 – -20.54	-12.60	**< 0.001**	339.82
Body size	-0.06	-0.12 – 0.01	-1.77	0.080	85.64
Selection [P] * treatment [Cup]	-4.55	-9.85 – 0.76	-1.69	0.093	339.52
Selection [S] * treatment [Predator]	-3.95	-9.27 – 1.36	-1.46	0.144	340.05

Nearest neighbor distance

Predictors	Estimates	CI	t	p	df
(Intercept)	39.17	33.89 – 44.45	14.96	**< 0.001**	43.22
Selection [P]	-3.43	-5.34 – -1.53	-4.55	**< 0.001**	5.31
treatment [Cup]	-10.77	-11.94 – -9.60	-18.13	**< 0.001**	337.80
treatment [Predator]	-10.88	-12.06 – -9.70	-18.17	**< 0.001**	339.69
Body size	-0.01	-0.03 – 0.01	-0.63	0.530	105.88
Selection [P] * treatment [Cup]	2.88	1.24 – 4.53	3.45	**< 0.001**	339.38
Selection [S] * treatment [Predator]	2.82	1.17 – 4.47	3.37	**< 0.001**	339.88

Table S2. Independent contrasts for comparisons between polarization-selected and control female guppies in their shoaling patterns when exposed to an open field test (OFT), a novel object (cup) and a predator model

Polarization

contrast	treatment	estimate	SE	df	lower.CL	upper.CL	t.ratio	p.value
C - P	OFT	-0.077	0.020	3.980	-0.133	-0.022	-3.898	**0.018**
C - P	Cup	-0.069	0.020	4.078	-0.124	-0.014	-3.435	**0.026**
C - P	Predator	-0.074	0.020	4.106	-0.129	-0.019	-3.698	**0.020**

Median speed

contrast	treatment	estimate	SE	df	lower.CL	upper.CL	t.ratio	p.value
C - P	OFT	-13.627	2.288	6.292	-19.164	-8.091	-5.956	**0.001**
C - P	Cup	-9.079	2.311	6.560	-14.619	-3.538	-3.928	**0.006**
C - P	Predator	-9.673	2.321	6.618	-15.226	-4.121	-4.168	**0.005**

Nearest neighbor distance

contrast	treatment	estimate	SE	df	lower.CL	upper.CL	t.ratio	p.value
C - P	OFT	3.434	0.754	5.309	1.530	5.339	4.555	**0.005**
C - P	Cup	0.552	0.761	5.510	-1.350	2.455	0.726	0.498
C - P	Predator	0.613	0.764	5.560	-1.292	2.517	0.802	0.455

Table S3. Statistical tests for comparisons in tests evaluating polarization-selected and control female guppies in their anti-predator behavior when exposed to a predator model

Number of inspections

Predictors	Incidence Rate Ratios	CI	t	p
(Intercept)	4.47	3.64 – 5.50	14.19	**< 0.001**
Selection [P]	0.87	0.75 – 1.01	-1.85	0.0645

Total time inspecting

Predictors	Incidence Rate Ratios	CI	t	p
(Intercept)	24.04	18.02 – 32.07	21.64	**< 0.001**
Selection [P]	0.79	0.66 – 0.95	-2.52	**0.011**

Mean inspection duration

Predictors	Incidence Rate Ratios	CI	t	p
(Intercept)	5.62	4.84 – 6.53	22.63	**< 0.001**
Selection [P]	0.82	0.69 – 0.96	-2.49	**0.013**

Table S4a. Statistical tests for overall comparisons of group polarization in polarization-selected and control female guppies when the average position of the group was shorter than 200 mm to the stimulus presented in the arena in tests that exposed these fish to a predator model and a novel object (cup).

Polarization (closer than 200 mm from the predator model)

Predictors	Estimates	CI	t	p	df
(Intercept)	0.68	0.60 – 0.75	25.78	**< 0.001**	3.87
Selection [P]	0.07	0.02 – 0.12	2.96	**0.003**	273.80
treatment [Predator model]	-0.12	-0.16 – -0.08	-5.34	**< 0.001**	146.76
Replicate [Rep2]	0.00	-0.04 – 0.04	0.05	0.958	143.72
replicate [Rep3]	0.01	-0.03 – 0.05	0.32	0.749	142.38
Selection [P] * treatment [Predator model]	-0.03	-0.08 – 0.03	-0.83	0.405	143.99

Table S4b. Statistical tests for independent contrasts of group polarization in polarization-selected and control female guppies when swimming at a distance closer than 200 mm to the stimulus presented in the arena in tests that exposed these fish to a predator model and a novel object (cup).

contrast	treatment	estimate	SE	df	lower.CL	upper.CL	t.ratio	p.value
C - P	Cup	-0.070	0.024	273.79	-0.116	-0.023	-2.956	**0.003**
C - P	Predator	-0.044	0.022	272.27	-0.088	-0.000	-1.984	**0.048**

Table S5a. Statistical tests for overall comparisons of group polarization in polarization-selected and control female guppies when the average position of the group was located closer to the tail area or closer to the head area of the predator model

Polarization

Predictors	Estimates	CI	Statistic	p	df
(Intercept)	0.62	0.54 – 0.71	22.54	**< 0.001**	3.15
Selection [P]	0.04	-0.03 – 0.10	1.53	0.191	4.58
location [tail]	-0.11	-0.14 – -0.07	-5.47	**< 0.001**	156.96
line [Social] * location [tail]	0.04	-0.01 – 0.09	1.64	0.103	150.99

Table S5b. Statistical tests for independent contrasts of group polarization in polarization-selected and control female guppies when the average position of the group was located closer to the tail area or closer to the head area of the predator model

contrast	location	estimate	SE	df	lower.CL	upper.CL	t.ratio	p.value
C - P	Head	-0.037	0.024	4.58	-0.102	0.027	-1.534	0.191
C - P	Tail	-0.080	0.023	3.63	-0.146	-0.013	-3.483	**0.030**

Table S6. Results from independent Linear Mixed Models evaluating differences in relative brain and relative brain region size between polarization-selected and control female guppies.

Region	Coefficent	Estimate	SE	df	t	P-value
Whole brain	Intercept	10.706	0.929	22.992	11.520	**< 0.001**
	Sel. Line (Polarization)	-0.015	0.037	23.296	-0.415	0.682
	Log (SL)	1.581	0.292	22.899	5.401	**< 0.001**
Olfactory bulbs	Intercept	-5.639	4.048	24.311	-1.393	0.176
	Sel. Line (Polarization)	-0.054	0.071	23.443	-0.759	0.455
	Log (rest of brain)	0.973	0.257	24.319	3.782	**< 0.001**
Ventral telencephalon	Intercept	-7.112	4.022	23.970	-1.768	0.0897
	Sel. Line (Polarization)	-0.054	0.070	23.366	-0.762	0.453
	Log (rest of brain)	1.107	0.255	23.968	4.329	**< 0.001**
Dorsal telencephalon	Intercept	-3.002	1.502	25.000	-1.998	0.0567
	Sel. Line (Polarization)	0.029	0.027	25.000	1.096	0.2837
	Log (rest of brain)	0.935	0.095	25.000	9.780	**< 0.001**
Thalamus	Intercept	-1.019	1.805	25.000	-0.565	0.577
	Sel. Line (Polarization)	0.073	0.033	25.000	2.187	**0.038**
	Log (rest of brain)	0.950	0.115	25.000	8.197	**< 0.001**
Hypothalamus	Intercept	-3.858	2.745	25.000	-1.405	0.172
	Sel. Line (Polarization)	0.048	0.049	25.000	0.974	0.339
	Log (rest of brain)	1.098	0.175	25.000	6.259	**< 0.001**
Nucleus glomerulus	Intercept	-2.352	9.716	25.000	-0.242	0.811
	Sel. Line (Polarization)	-0.067	0.176	25.000	-0.381	0.706
	Log (rest of brain)	0.807	0.617	25.000	1.308	0.203
Torus semicircularis	Intercept	-0.665	2.368	24.977	-0.281	0.781
	Sel. Line (Polarization)	-0.0325	0.042	24.106	-0.758	0.456
	Log (rest of brain)	0.829	0.150	24.982	5.496	**< 0.001**
Optic tectum cup	Intercept	1.073	0.966	23.494	1.111	0.2780
	Sel. Line (Polarization)	0.042	0.017	23.095	2.409	**0.024**
	Log (rest of brain)	0.800	0.061	23.474	12.933	**< 0.001**
Optic tectum core	Intercept	-2.883	1.658	25.000	-1.739	0.0944
	Sel. Line (Polarization)	0.006	0.030	25.000	0.211	0.8347
	Log (rest of brain)	1.040	0.106	25.000	9.805	**< 0.001**
Cerebellum	Intercept	-0.403	1.458	24.350	-0.277	0.784
	Sel. Line (Polarization)	-0.007	0.026	23.221	-0.275	0.786
	Log (rest of brain)	0.908	0.093	24.358	9.703	**< 0.001**
Medulla oblongata	Intercept	-0.145	1.519	24.920	-0.096	0.9246
	Sel. Line (Polarization)	-0.074	0.028	23.916	-2.656	**0.013**
	Log (rest of brain)	0.969	0.099	24.935	9.761	**< 0.001**

Table S7a Results from a Bayesian multilevel model evaluating differences in relative brain region size between polarization-selected and control female guppies. Stars indicate estimates that do not include zero in the confidence interval range based on the posterior samples drawn from the model.

Covariate	Estimate	Est.Error	l.95..CI	u.95..CI	
Medulla_Intercept	0.19	0.32	-0.37	0.77	
Cerebellum_Intercept	0.05	0.50	-1.08	1.04	
Nucleus glomerulus_Intercept	0.09	0.48	-0.84	1.04	
Torus semicircularis_Intercept	0.14	0.44	-0.69	1.03	
Thalamus_Intercept	-0.25	0.34	-0.97	0.31	
Optic tectum cups_Intercept	-0.19	0.51	-1.26	0.89	
Hypothalamus_Intercept	-0.06	0.39	-0.84	0.73	
Olfactory bulbs_Intercept	0.13	0.54	-0.94	1.26	
Ventral telencephalon_Intercept	0.05	0.63	-1.23	1.43	
Dorsal telencephalon_Intercept	-0.09	0.32	-0.72	0.56	
Optic tectum core_Intercept	-0.02	0.28	-0.56	0.54	
Medulla oblongata_Selection	-0.42	0.18	**-0.79**	**-0.06**	*
Medulla oblongata_Rest of the brain	0.91	0.09	**0.74**	**1.09**	*
Cerebellum_Selection	-0.07	0.21	-0.48	0.35	
Cerebellum_Rest of the brain	0.96	0.11	**0.75**	**1.18**	*
Nucleus glomerolus_Selection	-0.21	0.40	-0.98	0.58	
Nucleus glomerulus_Rest of the brain	0.34	0.20	-0.06	0.75	
Torus semicircularis_Selection	-0.24	0.32	-0.87	0.38	
Torus semicircularis_Rest of the brain	0.72	0.16	**0.40**	**1.04**	*
Thalamus_Selection	0.49	0.23	**0.04**	**0.94**	*
Thalamus_Rest of the brain	0.92	0.11	**0.70**	**1.16**	*
Optic tectum cups_Selection	0.32	0.16	**0.00**	**0.63**	*
Optic tectum cups_Rest of the brain	0.88	0.08	**0.72**	**1.06**	*
Hypothalamus_Selection	0.12	0.26	-0.39	0.64	

Brain region	Estimate	Est.Error	l.95..CI	u.95..CI	
Hypothalamus_Rest of the brain	0.87	0.13	**0.61**	**1.14**	*
Olfactory bulbs _Selection	-0.27	0.33	-0.93	0.38	
Olfactory bulbs_Rest of the brain	0.62	0.17	**0.29**	**0.95**	*
Ventral telencephalon_Selection	-0.17	0.31	-0.76	0.43	
Ventral telencephalon_Rest of the brain	0.58	0.16	**0.27**	**0.89**	*
Dorsal telencephalon_Selection	0.18	0.21	-0.25	0.60	
Dorsal telencephalon_Rest of the brain	0.91	0.11	**0.69**	**1.13**	*
Optic tectum core_Selection	0.05	0.20	-0.34	0.44	
Optic tectum core_Rest of the brain	0.93	0.10	**0.73**	**1.13**	*

Table S7b. Residual correlations of thalamus, optic tectum cups and medulla oblongata relative volume to other brain regions estimated from a Bayesian multilevel model evaluating differences in relative brain region size between polarization-selected and control female guppies. Stars indicate estimates that do not include zero in the confidence interval range based on the posterior samples drawn from the model.

Medulla oblongata

Brain region	Estimate	Est.Error	l.95..CI	u.95..CI
Cerebellum	-0.32	0.15	**-0.59**	**-0.02** *
Nucleus glomerulus	-0.26	0.15	-0.53	0.05
Torus semicircularis	0.07	0.16	-0.25	0.37
Thalamus	-0.40	0.14	**-0.65**	**-0.12** *
Optic tectum cups	-0.04	0.17	-0.36	0.29
Hypothalamus	-0.50	0.12	**-0.72**	**-0.23** *
Olfactory bulbs	-0.24	0.16	-0.52	0.09
Ventral telencephalon	-0.15	0.16	-0.44	0.17
Dorsal telencephalon	-0.12	0.17	-0.43	0.21
Optic tectum core	-0.22	0.15	-0.51	0.09

Optic tectum cups

Brain region	Estimate	Est.Error	l.95..CI	u.95..CI
Medulla oblongata	-0.04	0.16	-0.36	0.28
Cerebellum	-0.07	0.17	-0.41	0.26
Nucleus glomerulus	-0.07	0.17	-0.40	0.26
Torus semicircularis	0.05	0.17	-0.29	0.38
Thalamus	-0.18	0.16	-0.49	0.15
Hypothalamus	-0.15	0.16	-0.46	0.17

Brain region	Estimate	Est.Error	l.95..CI	u.95..CI
Olfactory bulbs	-0.11	0.17	-0.45	0.23
Ventral telencephalon	0.02	0.17	-0.32	0.36
Dorsal telencephalon	-0.01	0.17	-0.36	0.33
Optic tectum core	-0.06	0.17	-0.39	0.27

Thalamus

Brain region	Estimate	Est.Error	l.95..CI	u.95..CI
Medulla oblongata	-0.40	0.14	**-0.65**	**-0.12** *
Cerebellum	-0.27	0.16	-0.55	0.05
Nucleus glomerulus	0.05	0.16	-0.26	0.36
Torus semicircularis	-0.27	0.16	-0.56	0.06
Optic tectum cups	-0.18	0.17	-0.49	0.15
Hypothalamus	-0.15	0.16	-0.44	0.18
Olfactory bulbs	-0.09	0.16	-0.41	0.24
Ventral telencephalon	0.27	0.16	-0.05	0.56
Dorsal telencephalon	-0.05	0.17	-0.37	0.28
Optic tectum core	0.10	0.16	-0.22	0.41

Table S8. Statistical tests for comparisons in eye morphology between polarization-selected and control female guppies.

Eye diameter					
Predictors	*Estimates*	*CI*	*Statistic*	*p*	*df*
(Intercept)	3.12	2.95 – 3.29	78.98	**< 0.001**	2.00
Selection [P]	-0.03	-0.24 – 0.19	-0.52	0.658	2.00

Eye diameter / body length					
Predictors	*Estimates*	*CI*	*Statistic*	*p*	*df*
(Intercept)	0.12	0.11 – 0.12	69.97	**< 0.001**	2.00
Selection [P]	-0.00	-0.01 – 0.01	-0.13	0.906	2.00

Table S9a. Statistical tests for comparisons in visual acuity between polarization-selected and control female guppies.

Optomotor response

Predictors	Estimates	CI	Statistic	p	df
(Intercept)	0.10	0.03 – 0.17	3.16	**0.008**	12.06
Selection [P]	0.00	-0.08 – 0.09	0.11	0.913	12.88
Band width stimulus [0.28]	0.07	0.00 – 0.14	2.09	**0.036**	570.82
Band width stimulus [0.31]	0.15	0.08 – 0.21	4.20	**< 0.001**	569.22
Band width stimulus [0.35]	0.12	0.05 – 0.19	3.51	**< 0.001**	569.28
Band width stimulus [0.39]	0.18	0.12 – 0.25	5.26	**< 0.001**	574.04
Band width stimulus [0.44]	0.17	0.10 – 0.23	4.75	**< 0.001**	572.07
Optomotor response in static	0.56	0.49 – 0.63	14.92	**< 0.001**	531.69
Selection [P]* Band width stimulus [0.28]	0.01	-0.08 – 0.11	0.26	0.793	569.30
Selection [P]* Band width stimulus [0.31]	0.01	-0.09 – 0.10	0.11	0.915	569.21
Selection [P]* Band width stimulus [0.35]	0.03	-0.06 – 0.13	0.63	0.526	569.18
Selection [P]* Band width stimulus [0.39]	-0.03	-0.13 – 0.06	-0.64	0.520	569.16
Selection [P]* Band width stimulus [0.44]	0.01	-0.09 – 0.10	0.17	0.863	569.22

Table S9b. Independent contrasts for optomotor response observed in fish at rotating stimulus of several band widths

contrast	estimate	SE	df	lower.CL	upper.CL	t.ratio	p.value
bw25 - bw28	-0.079	0.024	571.167	-0.149	-0.009	-3.236	**0.016**

bw25 - bw31	-0.148	0.024	569.618	-0.218	-0.079	-6.094	**< 0.001**
bw25 - bw35	-0.137	0.024	569.230	-0.207	-0.068	-5.637	**< 0.001**
bw25 - bw39	-0.168	0.025	577.744	-0.238	-0.097	-6.818	**< 0.001**
bw25 - bw44	-0.169	0.024	573.616	-0.239	-0.099	-6.918	**< 0.001**
bw28 - bw31	-0.069	0.024	569.653	-0.139	0.000	-2.852	0.051
bw28 - bw35	-0.058	0.024	570.338	-0.128	0.011	-2.390	0.161
bw28 - bw39	-0.089	0.024	571.492	-0.159	-0.019	-3.639	**0.004**
bw28 - bw44	-0.090	0.024	569.612	-0.160	-0.021	-3.712	**0.003**
bw31 - bw35	0.011	0.024	569.262	-0.058	0.081	0.460	0.997
bw31 - bw39	-0.019	0.025	574.211	-0.089	0.051	-0.791	0.969
bw31 - bw44	-0.021	0.024	571.155	-0.091	0.049	-0.858	0.956
bw35 - bw39	-0.031	0.025	575.985	-0.101	0.040	-1.245	0.814
bw35 - bw44	-0.032	0.024	572.350	-0.102	0.038	-1.315	0.777
bw39 - bw44	-0.002	0.024	569.833	-0.071	0.068	-0.064	1.000

Table S9c. Independent contrasts for optomotor response observed in polarization-selected and control female guppies at rotating stimulus of several band widths.

contrast	stimulus	estimate	SE	df	lower.CL	upper.CL	t.ratio	p.value
C - P	bw25	-0.004	0.038	601.455	-0.078	0.070	-0.112	0.911
C - P	bw28	-0.017	0.038	601.441	-0.091	0.057	-0.456	0.649
C - P	bw31	-0.009	0.038	601.422	-0.083	0.064	-0.247	0.805
C - P	bw35	-0.035	0.038	601.457	-0.109	0.039	-0.938	0.349
C - P	bw39	0.027	0.038	601.458	-0.047	0.101	0.722	0.471
C - P	bw44	-0.013	0.038	601.452	-0.086	0.061	-0.340	0.734

Table S10a. Statistical tests for comparisons in visual temporal resolution between polarization-selected and control female guppies.

Speed deviation from stimulus rotation

Predictors	Estimates	CI	Statistic	p	df
(Intercept)	8.38	3.23 – 13.53	5.31	**0.014**	2.87
Selection [P]	-0.46	-8.74 – 7.83	-0.19	0.863	2.64
Speed [24]	-5.93	-7.71 – -4.15	-6.54	**< 0.001**	4398.86
Speed [36]	-23.39	-25.17 – -21.61	-25.78	**< 0.001**	4398.86
Speed [45]	-49.74	-51.52 – -47.96	-54.79	**< 0.001**	4398.84
Rotation[Counterclockwise]	-7.11	-8.36 – -5.85	-11.08	**< 0.001**	4398.89
Selection [P] * Speed [24]	0.31	-2.17 – 2.79	0.25	0.805	4398.91
Selection [P] * Speed [36]	1.26	-1.22 – 3.74	0.99	0.320	4398.91
Selection [P] * Speed [45]	-0.48	-2.96 – 1.99	-0.38	0.701	4398.84
Selection [P] * Rotation[Counterclockwise]	1.29	-0.47 – 3.04	1.44	0.150	4398.98

Proportion of time following the stimulus

Predictors	Estimates	CI	Statistic	p	df
(Intercept)	0.87	0.80 – 0.94	39.43	**< 0.001**	3.12
Selection [P]	0.00	-0.14 – 0.15	0.10	0.928	2.52
Speed [24]	0.03	0.00 – 0.06	2.03	**< 0.001**	4398.95
Speed [36]	0.03	-0.00 – 0.05	1.83	0.067	4398.95
Speed [45]	-0.04	-0.06 – -0.01	-2.55	**0.018**	4398.92
Rotation[Counterclockwise]	-0.07	-0.09 – -0.05	-7.33	**< 0.001**	4398.98
Selection [P] * Speed [24]	0.01	-0.02 – 0.04	0.74	0.457	4399.06
Selection [P] * Speed [36]	-0.00	-0.04 – 0.03	-0.20	0.843	4398.99
Selection [P] * Speed [45]	-0.00	-0.04 – 0.04	-0.15	0.884	4398.99

Selection [P] * Rotation[Counterclockwise]	-0.01	-0.05 – 0.03	-0.62	0.536	4398.92

Table S10b. Independent contrasts for speed deviation to stimulus rotation observed in polarization-selected and control females at multiple rotation speeds in clockwise and counterclockwise directions

Selection	Speed	Rotation	emmean	SE	df	lower.CL	upper.CL	t.ratio	p.value
C	14.4	Clockwise	8.379	1.577	2.874	3.233	13.526	5.313	**0.015**
P	14.4	Clockwise	7.922	1.938	2.485	0.966	14.879	4.089	**0.038**
C	24	Clockwise	2.447	1.577	2.872	-2.701	7.594	1.552	0.222
P	24	Clockwise	2.303	1.938	2.488	-4.650	9.256	1.188	0.336
C	36	Clockwise	-15.012	1.577	2.872	-20.160	-9.865	-9.520	**0.003**
P	36	Clockwise	-14.212	1.938	2.488	-21.164	-7.259	-7.332	**0.010**
C	45	Clockwise	-41.357	1.577	2.874	-46.503	-36.210	-26.223	**< 0.001**
P	45	Clockwise	-42.299	1.938	2.485	-49.255	-35.342	-21.830	**0.001**
C	14.4	Counterclockwise	1.272	1.577	2.873	-3.875	6.420	0.807	0.481
P	14.4	Counterclockwise	2.103	1.938	2.487	-4.850	9.057	1.085	0.372
C	24	Counterclockwise	-4.660	1.577	2.869	-9.809	0.489	-2.956	0.063
P	24	Counterclockwise	-3.517	1.939	2.491	-10.465	3.432	-1.814	0.186
C	36	Counterclockwise	-0.483	-22.119	1.577	2.869	-27.268	-14.030	**0.001**
P	36	Counterclockwise	-20.031	1.939	2.491	-26.979	-13.083	-10.331	**0.004**
C	45	Counterclockwise	-48.464	1.577	2.873	-53.611	-43.317	-30.731	**< 0.001**
P	45	Counterclockwise	-48.118	1.938	2.487	-55.071	-41.164	-24.827	**< 0.001**

Table S11. Statistical results using a robust linear mixed model approach for comparisons in the proportion of time following the correct direction of the stimulus in visual temporal assays between polarization-selected and control female guppies.

Proportion of time following the stimulus

Predictors	Estimates	Std. Error	t-value	p	df
(Intercept)	0.91	0.016	57.45	**< 0.001**	3.12
Selection [P]	-0.001	0.031	-0.04	0.971	2.52
Speed [24]	0.014	0.009	1.61	0.106	4398.95
Speed [36]	0.027	0.009	3.07	**0.001**	4398.95
Speed [45]	-0.010	0.009	-1.17	**0.240**	4398.92
Rotation[Counterclockwise]	-0.041	0.006	-6.51	**< 0.001**	4398.98
Selection [P] * Speed [24]	0.004	0.012	0.33	0.744	4399.06
Selection [P] * Speed [36]	0.004	0.012	0.33	0.741	4398.99
Selection [P] * Speed [45]	-0.009	0.012	-0.74	0.460	4398.99
Selection [P] * Rotation[Counterclockwise]	-0.001	0.009	-0.15	0.882	4398.92

Robustness weights for the residuals:
 3546 weights are ~= 1. The remaining 974 ones are summarized as
  Min. 1st Qu. Median   Mean 3rd Qu.   Max.
 0.199 0.328 0.518 0.557 0.773 0.999

Robustness weights for the random effects:
 101 weights are ~= 1. The remaining 18 ones are summarized as
  Min. 1st Qu. Median   Mean 3rd Qu.   Max.
 0.247 0.489 0.739 0.693 0.904 0.995

Rho functions used for fitting:
 Residuals:
  eff: smoothed Huber (k = 1.345, s = 10), sig: smoothed Huber, Proposal II (k = 1.345, s = 10)
 Random Effects, variance component 1 (trial):
  eff: smoothed Huber (k = 1.345, s = 10), vcp: smoothed Huber, Proposal II (k = 1.345, s = 10)
 Random Effects, variance component 2 (rep):
  eff: smoothed Huber (k = 1.345, s = 10), vcp: smoothed Huber (k = 1.345, s = 10)

R syntax for final models used in statistical analyses

Shoaling patterns (Table S1 + Table S2)

- Polarization_model = lmer(polarization ~ selection line * treatment + body size + (selection line | replicate) + (1 | trial id))
  emmeans(Polarization_model, pairwise ~ line | treatment)

- Activity_model = lmer(median speed ~ selection line * treatment + body size + (selection line | replicate) + (1 | trial id))
  emmeans(Activity_model, pairwise ~ line | treatment)

- Speed_model =lmer(nearest neighbor distance ~ selection line * treatment + body size + (selection line | replicate) + (1 | trial id))
  emmeans(Speed_model, pairwise ~ line | treatment)

Predator inspection behavior (Table S3)

- Number_of_inspections = glmmTMB(inspections ~ selection line + (1 | selection line:replicate) + (1 | Order of stimulus), poisson(link = "log"))

- Total time inspecting = glmmTMB(time inspecting ~ selection line + (1 | selection line:replicate) + (1 | Order of stimulus), genpois(link = "log"))

- Mean inspection duration = glmmTMB(mean inspection ~ selection line + (1 | selection line:replicate) + (1 | Order of stimulus), genpois(link = "log"))

Shoaling patterns during predator inspection trials (Tables S4 + S5)

- Polarization_closer_200mm = lmer(polarization ~ selection line * group position + replicate + (1 | trial id) + (1 | Order of stimulus))
  emmeans(Polarization_closer_200mm, pairwise ~ selection line | group position)

- Polarization_in_tail_head_areas = lmer(polarization ~ selection line * group position + (selection line | replicate) + (1 | trial id) + (1 | Order of stimulus))

emmeans(Polarization_in_tail_head_areas, pairwise ~ selection line | group position

Brain anatomy (Independent Linear Mixed Models for brain regions and whole brain size ; Table S6)

Whole_brain = lmer(log(brain) ~ selection line + log(standard lentgh) + (1 | replicate))

Brain_region = lmer(log(brain region) ~ selection line + log(rest of the brain) + (1 | replicate)

Brain anatomy (Multilevel Model for all brain regions; Table S7)

DT = bf(scale(DT) ~ selection line + scale(rest of the brain) + (1 | replicate))
VT = bf(scale(VT) ~ selection line + scale(rest of the brain) + (1 | replicate))
OB = bf(scale(OB) ~ selection line + scale(rest of the brain) + (1 | replicate))
Hyp = bf(scale(Hyp) ~ selection line + scale(rest of the brain) + (1 | replicate))
OT = bf(scale(OT) ~ selection line + scale(rest of the brain) + (1 | replicate))
OTc = bf(scale(OTc) ~ selection line + scale(rest of the brain) + (1 | replicate))
Th = bf(scale(Th) ~ selection line + scale(rest of the brain) + (1 | replicate))
TS = bf(scale(TS) ~ selection line + scale(rest of the brain) + (1 | replicate))
NG = bf(scale(NG) ~ selection line + scale(rest of the brain) + (1 | replicate))
Cb = bf(scale(Cb) ~ selection line + scale(rest of the brain) + (1 | replicate))
MO = bf(scale(MO) ~ selection line + scale(rest of the brain) + (1 | replicate))

Multilevel_model <- brm(mvbf(DT, VT, OB, Hyp, OT, OTc, Th, TS, NG, Cb, MO) + set_rescor(TRUE), chains = 16, cores = 16, control = list(adapt_delta = 0.995, max_treedepth =15), iter = 3000, warmup = 2000)

Eye morphology (Table S8)

Eye_size = lmer(eye size ~ selection line + (selection line | replicate))

Relative_eye_size = lmer(relative eye size ~ selection line + (selection line | replicate))

Visual acuity (Table S9)

Visual_acuity_model = lmer(optomotor response ~ selection line * stimulus bandwidth + time moving in static + (1| fish id) + (selection line | replicate))

Stimulus_bandwidth_contrasts <- emmeans(Visual_acuity_model ~ stimulus bandwidth, type = "response")
pairs(Stimulus_bandwidth_contrasts, adjust="tukey")

Selection_line_contrasts <- emmeans(Visual_acuity_model, ~selection line | stimulus bandwidth, type = "response")
pairs(Selection_line_contrasts)

Temporal resolution (Tables S10 and S11)

Deviation_speed_model = lmer(speed deviation ~ selection line+ speed of rotation + direction of rotation + selection line : speed of rotation + selection line : direction of rotation + (1 | trial id) + (selection line | replicate))
Selection_line_contrasts <- emmeans(Deviation_speed_model, ~ selection line | speed of rotation * direction of rotation, type = "response", pbkrtest.limit = 4520)
pairs(Selection_line_contrasts)

Proportion_time_following_model = lmer(proportional time following ~ selection line+ speed of rotation + direction of rotation + selection line : speed of rotation + selection line : direction of rotation + (1 | trial id) + (selection line | replicate))
Selection_line_contrasts <- emmeans(Proportion_time_following_model, ~ selection line | speed of rotation * direction of rotation, type = "response", pbkrtest.limit = 4520)
pairs(Selection_line_contrasts)

Proportion_time_following_robust_model <- rlmer(proportional time following ~ selection line+ speed of rotation + direction of rotation + selection line : speed of rotation + selection line : direction of rotation + (1 | trial id) + (selection line | replicate))

Relative density
0.00 0.25 0.50 0.75 1.00

Open field | Novel object | Predator model

Control

Polarization-selected

Polarization

Mean group speed (mm/frame)