**Evolution of schooling propensity in the guppy drives changes in anti-predator behavior that are linked to neuroanatomy**

Alberto Corral-Lopez*[1,2,3], Alexander Kotrschal[2,4], Alexander Szorkovszky[5,6], Maddi Garate-Olaizola[2,7], James Herbert-Read[8,9], Wouter van der Bijl[1], Maksym Romenskyy[6,10], Hong-Li Zeng[11,12], Severine Denise Buechel[2,4], Ada Fontrodona-Eslava[2,13], Kristian Pelckmans[12], Judith E. Mank[1], Niclas Kolm[2]

*corresponding author: corral@zoology.ubc.ca

**Affiliations**

- Department of Zoology, University of British Columbia, Vancouver, Canada
- Department of Zoology/Ethology, Stockholm University, Stockholm, Sweden.
- Division of Biosciences, University College London, London, United Kingdom
- Behavioural Ecology, Wageningen University & Research, Wageningen, Netherlands.
- RITMO Centre for Interdisciplinary Studies in Rhythm, Time and Motion, University of Oslo, Oslo, Norway
- Department of Mathematics, Uppsala University, Uppsala, Sweden.
- Department of Ecology and Genetics, Animal Ecology, Uppsala University, Sweden
- Department of Zoology, University of Cambridge, Cambridge, UK.
- Aquatic Ecology, Lund University, Lund, Sweden.
- Department of Life Sciences, Imperial College London, London, UK.
- School of Science, Nanjing University of Posts and Telecommunications, Nanjing, China.
- Department of Information Technology, Uppsala University, Uppsala, Sweden.
- Centre for Biological Diversity, School of Biology, University of St Andrews, St Andrews, UK

## Abstract

One of the most spectacular displays of social behavior is the synchronized movements that many animal groups perform to travel, forage and escape from predators. However, the mechanistic basis of the evolution of such collective behaviors, as well as their fitness effects, remains empirically untested. Here, we study anti-predator behavior in guppies experimentally selected for divergence in polarization, an important behavioral aspect of coordinated movement. We find that groups from artificially selected lines remain more polarized than control groups in the presence of a threat. Neuroanatomical measurements show these behavioral differences are linked to changes in brain regions previously suggested as important regulators of perception, fear and attention, and motor response. We use further analyses of behavior and visual capabilities to show that differences in anti-predator behavior are not attributable to changes in visual perception, but likely to more efficient transfer of social information in polarization-selected fish. Our findings highlight that brain morphology may play a fundamental role in the evolution of coordinated movement and anti-predator behavior.

## Introduction

Animals regularly gather - for safety, for exploiting resources, or for mating. Group-living often leads to spectacular forms of collective behavior, and individuals in many taxa coordinate their movements in order to increase efficiency in foraging and travelling, or to confuse predators [1]. To date, we have a detailed understanding of the interaction rules that produce highly coordinated movements in animal groups [e.g. 2,3], as well as the ecological factors that produce the broad variation observed across and within species [4]. Collective motion has evolved many times in fish, and is underpinned by the efficient acquisition of information through the sensory system, mainly through visual cues [5]. Fish schooling is widely understood as a behavioral adaptation to reduce the risk of predation [6]. But although correlation-based analyses have revealed how predation levels are associated with variation in collective motion in wild populations (see for instance [7]), the causal aspects are still unclear, particularly how evolutionary changes in collective motion contribute to anti-predator specific situations, or what type of visual information and information processing schooling fish use to identify and avoid predators as groups.

The brain, as the central organ controlling locomotion, sensory systems and decision-making,
should play a major role in the ability to coordinate movements in animal groups. As such,
variation in the anatomy of the brain could be an important mechanism behind the evolution of
collective motion. Despite its potential importance, studies explicitly testing the role of
neuroanatomy in collective motion are scarce. However, the link between social factors and
changes in multiple brain structures across taxa is well established[8–11] (Barton 1996, Burish et al.
2004, Chee et al 2013, Triki et al. 2019).

This association is particularly well studied in fish, where approximately half of marine and
freshwater species come together in groups at different life stages[12]. For instance, Tanganyikan
cichlids species with more complex social structures have larger telencephali and hypothalami[13],
both parts of the fish forebrain, a region with important function in social behavior [14], which has
also been associated with social competence in cleaner fish and social orienting in zebrafish[10,15].
In addition, exposure to larger groups during development in nine-spined stickleback is correlated
with larger size of another brain region, the optic tectum, the visual center in the fish brain [16].

Schooling requires sensory perception of neighbors' movements and positions and motor control
to enact speed and directional changes. Brain regions associated with fish social behavior have
also been implicated in the few studies explicitly testing the link between neuroanatomy and
schooling behavior. First, lesion studies in goldfish (*Carassius auratus*) showed that individuals
with ablated telencephalon exhibited reduced activity and association with conspecifics[17]. Second,
a study on surface and cavefish populations of *Astyanax mexicanus* living in different light
environments showed an underlying positive correlation between optic tectum size and schooling
propensity differences between populations[18]. These limited studies highlight the potential role of
neuroanatomy in schooling, as well as the need to account for environmental variation in analyses.

Here we use artificial selection lines of guppies (*Poecilia reticulata*) with divergence in schooling
propensity, which offer a unique opportunity to empirically evaluate the link between evolution of
general collective behavior, specific anti-predator behavior and neuroanatomy[19]. In relation to
many fish species that associate in large schools, guppies have relatively low schooling propensity,
with high levels of variation across individuals and across populations as a function of external

factors such as predation risk or food availability[20,21]. In our selection lines, intrinsic schooling

propensity was increased in female guppies by over 15% compared to controls in just three

generations by selecting individuals that exhibited higher polarization, the level of alignment

between individuals moving together in a group[19,22]. Previous assays in polarization-selected

females provided a good account of changes in their individual movement patterns in relation to

control females, including higher alignment and attraction to neighbors[19].

We use these lines to investigate potential changes in the visual system and anti-predator behavior

following directional selection for schooling propensity, as well as to study the association

between increased schooling propensity and changes in brain anatomy. For this, we first evaluated

collective motion patterns and predator inspection in groups of polarization-selected and control

female guppies, and found that polarization-selected females inspected threats for shorter times

and formed more cohesive groups when exposed to these threats than control females. Second, we

quantified brain region sizes with microcomputed tomography (microCT) and found that these

behavioral differences were linked to changes in brain regions previously suggested as important

regulators of perception, fear and attention, and motor responses. Third, we performed

comprehensive tests of visual capabilities, spanning morphological, visual acuity and temporal

resolution measurements, which revealed that despite the occurrence of significant changes in

brain regions regulating the visual system, the differences in schooling propensity and anti-

predator behavior between the selection lines do not appear to be driven by differences in the

capacity to acquire visual information. By linking changes in schooling propensity to function in

an ecologically relevant setting and to brain structure size variation we identify potential

evolutionary pathways leading to collective motion.

**Results**

**Collective motion in response to predation threat in guppies following artificial selection**

We investigated whether selection for higher schooling propensity affected cohesiveness and how

individuals from groups react in response to neighbor movement in a predation context. These

behavioral decisions should have major fitness consequences in this species[23]. Specifically, we

recorded and tracked fish in an experimental arena to obtain positional data and assessed collective

motion of groups of eight guppies when exposed to an imminent threat, the presence of an artificial

replica of a pike cichlid (*Crenicichla frenata*), a natural predator in wild populations of Trinidadian

guppies. Furthermore, we exposed these groups to a non-predator-shaped object to allow for

comparisons when presented to a novel object. These assays were performed in combination with

open field tests (OFT's) on the same fish groups. Previous analyses of the data for OFT's in these

groups provided evidence that selection for polarization altered individuals' speed, how

individuals aligned with, and how individuals were attracted towards conspecifics during group

motion[19].

Our analyses of collective motion of female groups exposed to a predator model and a novel object

showed predictable results in relation to previous findings observed in OFT's. In the presence of

a predator model or a novel object, we observed an overall strong decline in the polarization of the

groups, as well as in individuals' speed (Supplementary Table 1, Fig. 1A). Differences between

polarization-selected and control groups in these traits were still present, but to a reduced degree

in the presence of these stimuli in the experimental arena (Supplementary Table 1, Fig. 1).

Similarly, overall attraction towards conspecifics in female groups (using median nearest neighbor

distance as proxy) was stronger when exposed to both stimuli than in OFT's. Yet, differences

between polarization and control lines in nearest neighbor distance previously observed in

OFT's[19], were no longer observed when these groups of fish were exposed to a predator model

and a novel object (Fig. 1A; Supplementary Table 2). We observed no differences between the

response towards the predator model compared to the novel object in how these stimuli altered the

attraction towards conspecifics of these groups (LMM$_{attraction}$: novel object vs predator model: t=

0.331; df = 335; p = 0.941). On the contrary, the predator model elicited a stronger response than

the novel object in the speed and polarization of the group. Specifically, collective motion data

showed that guppy groups were slower and less aligned in the presence of a predator model

(LMM$_{polarization}$: novel object vs predator model: t = 2.87; df = 338; p = 0.011; LMM$_{speed}$: novel

object vs predator model: t = 4.71; df = 338; p < 0.001). Visual inspection of heatmaps

summarizing data of group alignment in combination with positional data concords with these

results obtained from statistical models using summary statistics for each group (Fig. 1B). Yet,

differences in polarization between control and polarization-selected groups in open field and
novel object assays were consistent across all regions of the arena, while the observed statistical
differences in assays with a predator model were more pronounced in positions further away from
the head of the model (Fig. 1B).

**Social information processing in response to predation threat**

178   To characterize potential differences in how efficiently social information spreads in polarization-
179   selected and control fish groups when exposed to a predator threat, we quantified inspection
180   behavior of individuals in our groups of eight fish in the presence of a predator model, and
181   collective motion of the group at times and in locations associated to predator inspection behavior
182   during our assays.

*Predator inspection behavior*

We scored recorded videos for the start and end point for each predator inspection performed by
one randomly selected fish in each video. Analyses of predator inspection data showed that females
from control lines presented a higher tendency to inspect the predation threat presented in the
experimental arena than polarization-selected females. Specifically, we observed that total time
inspecting and the mean duration of predator inspections were significantly shorter in polarization-
selected females (GLMM$_{time\_inspecting}$: selection: Ratio = 0.79 (0.66-0.95), t = -2.52, p = 0.011;
GLMM$_{mean\_inspection}$: selection : Ratio = 0.82 (0.66-0.96), t = -2.49, p = 0.013, Fig. 2d-e,
Supplementary Table 3), while the total number of inspections showed a similar trend
(GLMM$_{inspections}$: selection: Ratio = 0.87 (0.75-1.01), t = -1.85, p = 0.064; Fig. 2c, Supplementary
Table 3).

*Collective motion during predator inspections*

Analysis of positional data and median distance to the stimulus presented in our assays suggested
that most inspection behaviors to the predator model were performed during the initial 3 minutes
of the assays (Supplementary Figure 1). Further, the majority of inspections were performed at a
range closer than 200 mm and in the tail area of the predator model presented in the experimental
arena (Fig. 1A; Fig. 2A). Consequently, we filtered our data to evaluate collective motion patterns

of fish groups in the time and locations where predator inspections were performed during our
assays (see methods). The overall differences in group polarization found between polarization-
selected and control groups were maintained in areas within 200 mm of the predator model (Fig.
2B; $LMM_{polarization < 200mm – predator model}$: t = -1.984, df = 272, p = 0.048; Supplementary Table 4a-b).
Inspection behavior is mainly performed from areas with reduced risk of attack from a predator.
In line with such expectation, we found that polarization of all groups was greatly reduced in the
area of the predator model tail (Fig. 2B). However, we found no differences in group polarization
between selected and control females in the head area of the predator model, but a stronger
maintenance of group polarization of polarization-selected females in close proximity to the tail
of the predator model ($LMM_{polarization – head area}$: t = -1.53, df = 4.58, p = 0.190; $LMM_{polarization – tail}$
$_{area}$: t = -1.53, df = -3.48, p = 0.029; Fig. 2B, Supplementary Table 5a-b).
213
214  **Changes in neuroanatomy following artificial selection**
215
We used microcomputed tomography (micro-CT) to determine whether the volumes of 11 major
brain regions and overall brain volume of female guppies might be associated with selection for
higher schooling propensity and changes in collective behavior in response to predator threats.
Specifically, we used micro-CT scans to reconstruct the brain anatomy of 13 polarization-selected
females, and 15 control females (see methods). Polarization-selected and control fish showed no
differences in whole brain volume in relation to their body size ($LMM_{wholebrain}$: t = -0.41, df =
23.29, p = 0.682; Fig. 3a). However, analyses of the volume of each region in relation to the rest
of the brain indicated that the thalamus and optic tectum cups are larger in polarization-selected
than control females ($LMM_{thalamus}$: selection: t = 2.187, df = 25, p = 0.038; $LMM_{o.tectum}$: selection:
t = 2.409, df = 23.09, p = 0.024; Fig. 3a), and the medulla oblongata is larger in control females
($LMM_{medulla}$: selection: t = -2.65, df = 23.91, p = 0.013; Fig. 3a). All other eight brain regions
measured presented no difference between polarization-selected and control females in relative
volume (Fig. 3a-b, Supplementary Table 6).
229
In parallel, we analyzed brain region volume differences using a more conservative approach and
found similar and consistent differences between selection lines. Specifically, we used a
multivariate Bayesian model that included the relative size of the 11 brain regions as dependent

variables. Posterior samples drawn from the multivariate model indicated that confidence intervals

for the difference in relative volume in the medulla oblongata, the optic tectum cups and the

thalamus between polarization-selected and control females did not overlap with zero

(Supplementary Table 7a).

We then used the multivariate Bayesian model to evaluate the correlation in relative brain region

volume between multiple regions measured. We focused on evaluating correlations with other

brain regions for the three regions significantly differentiated between lines following artificial

selection (Supplementary Table 7b). We found no correlation between optic tectum cup relative

volume and volume of any other region measured. However, we found a significant inverse

correlation between thalamus and medulla relative volume (rescorr$_{Medulla-Thalamus}$: -0.40 ± 0.14;

lower/upper 95% CI: -0.65 / -0.12). This finding suggests that the opposite differences observed

in the volume of these two brain regions between control and polarization-selected female guppies

may be linked to changes in brain development processes associated with artificial selection for

higher coordinated motion.

**Information acquisition through the visual system**

Efficiently acquiring information through the sensory system, mainly through visual cues, is a

basic principle of collective motion in shoaling fish[5]. Given observed differences in the size of the

optic tectum cups between polarization-selected and control fish, we investigated potential

differences in visual perception between lines. For this, we compared eye morphology and two

key characteristics of the visual system to track movement of conspecifics, visual acuity and

temporal resolution.

*Eye morphology*

We quantified eye morphology in a total of 112 individuals from polarization-selected and control

lines. Eye size is a common indicator of visual capacities of organisms[24], and comparative studies

across fish species suggest that larger eyes correlate with improved visual abilities[25]. In our study,

we found no difference between the lines in either absolute eye size or relative eye size, the

proportional size of the eye in relation to body size (Fig. 4A; LMM$_{eye size}$: selection: t = -0.52, df =

2, p = 0.658; LMM $_{relative eye size}$: selection: t = -0.13, df = 2, p = 0.906; Supplementary Table 8).

*Visual acuity*

We further assessed potential differences in visual perception between selection and control lines
by quantifying visual acuity in the same individuals for which eye morphology was measured.
Visual acuity allows an individual to resolve spatial detail and can be critical for an organism's
fitness[26]. We measured visual acuity in our fish by quantifying their innate optomotor response in
contrasting rotating gratings. This a widely used method to study visual acuity in multiple fish
species, including guppies[27–29], and we have previously used this approach to evaluate the visual
system of guppies in similar contexts[30,31]. Following the methods in ([30]), we exposed our fish to a
series of six stimuli with rotating and static gratings of different widths at the lower end of the
known guppy visual acuity, where thinner widths are more difficult to perceive. When comparing
optomotor response between polarization-selected and control fish, we found no difference in their
average optomotor response combining data from all stimuli (LMM$_{acuity}$: selection: t = 0.11, df =
12.88, p = 0.913; Supplementary Table S9a), or in analyses independently evaluating specific
optomotor response for any of the 6 stimuli presented (Fig. 4B; Supplementary Table 9c).

*Visual resolution tracking movement*

Although the ability to resolve spatial detail, acuity, is arguably an important visual parameter for
guppies to recognize conspecific positions in shoals, it provides no information on an individual's
ability to track movement[32]. Similar to many social fish species, guppies swim with a saltatory
movement style that features discrete changes in speed and direction[7]. Consequently, we
implemented an additional experiment that evaluated potential differences between polarization-
selected and control fish in their temporal assessment of speed and direction changes. Using the
same experimental apparatus used to evaluate visual acuity, we video recorded female guppies
from our selection and control lines when they were exposed to a single-width rotating stimulus
(see methods). We next used automated tracking to obtain orientation and speed of the fish for
each frame and to quantify their direction and speed in relation to the stimuli presented at each
time point.

Overall, fish followed the direction of the rotating stimulus for a significant proportion of the time.
This was the case when the stimuli were presented in both a clockwise direction an in

counterclockwise direction (Supplementary Figure 2, Supplementary Table 10). Overall, swimming speed did not significantly deviate from the stimuli rotating speed at the two lowest speed levels (Supplementary Figure 2, Supplementary Table 10), but was less than the stimuli speed at the two highest speed levels (Supplementary Figure 2, Supplementary Table 10). This was true for both directions in which stimuli were presented, but the mismatch between swimming and stimuli rotation speed was greater at the higher speed when the stimulus rotated anticlockwise (Supplementary Figure 2, Supplementary Table 10).

We compared the performance of polarization-selected and control fish in the test to evaluate their visual temporal resolution while shoaling. We found no differences between selection and control lines in their deviation of their swimming speed in relation to the stimuli rotating speed for their combined scores across speeds and direction of rotation (Fig. 4C; $\text{LMM}_{speed\_deviation}$: selection: $t = -0.46$, df $= 2.64$, p $= 0.863$; Supplementary Table 10a), or for their speed observed at any particular speed at direction of rotation (Supplementary Figure 2, Supplementary Table 10a). Similarly, polarization-selected and control females spent similar proportions of time following the stimuli during changes in stimuli rotating speed (Supplementary Figure 2; Supplementary Table 10a; $\text{LMM}_{proportion\_time}$: selection: $t = 0.10$, df $= 2.52$, p $= 0.928$).

**Discussion**

Our work demonstrates that selection for schooling behavior in female guppies has important implications for anti-predator responses in this species. Analyses of motion patterns in these fish shows that polarization-selected groups maintain higher activity and sociability when exposed to a potential predator threat. In addition, our analyses suggest that individuals from polarization-selected groups rely more on neighbor information during a predator threat, as they spent less time inspecting individually. We further studied visual capacities in these fish and found no differences between polarization-selected and control fish, suggesting that the differences in collective motion and predator inspection behavior observed are not driven by their ability to distinguish the threat at longer distances or to visually acquire information on neighbor movements.

In parallel, our results suggest that artificial selection for higher schooling propensity has produced
significant changes in the brain anatomy of female guppies. Neuroanatomical measurements
indicate that polarization-selected fish exhibit a larger thalamus and a large optic tectum cup, but
a smaller medulla oblongata, compared to control fish. These rapid changes in brain region sizes
in response to selection for polarization behavior are consistent with previous artificial selection
directly on neuroanatomy, which resulted in rapid shifts in both relative brain size and relative
telencephalon size, in just a few generations in guppies[33,34].

Below, we discuss the implications of these discoveries for our understanding on how the
association between brain morphology and anti-predator behavior might drive the evolution of
collective behavior.

*Information processing in a predation threat context*
Our behavioral analyses indicate that rapid evolution of schooling propensity affects how groups
of fish behave when encountering a threat. Fish schooling is widely understood as a behavioral
adaptation to escape the effect of predation[1]. These synchronized movements have been shown to
confer two major benefits to fish schools, facilitating escape through transfer of information from
closer neighbors[35], or by confusing the predator in which individual to attack[36]. The use of a static
predator model in our assays does not allow us to infer any potential benefit of higher schooling
propensity on the confusion effect towards predators. However, our results show that directional
selection and associated changes in the brain lead to robust behavioral changes across multiple
contexts and that it might affect individual ability to efficiently process social information in
response to predation. The reduced time spent inspecting the predator model by polarization-
selected females, coupled with the fact that polarization-selected groups remained more aligned
closer to the predator model, especially around the tail of the predator, suggest this a likely
possibility. However, further comparisons within asocial and social contexts should be
implemented to disregard the alternative explanation that directional selection leads to changes in
predator inspection behavior also when fish have no access to social information.

Our study only measured fitness effects indirectly, using a predator model, following directional
selection for polarization. Yet, previous work demonstrated that shorter inspection times towards

the same predator models are associated with higher survival in the species[37,38]. Given our findings,

this is likely an important fitness benefit for individuals showing higher coordination with

conspecifics. This benefit might trade-off with a reduced level of private information from

potential threats obtained by these individuals. These factors are arguably important selective

pressures in natural populations where guppies from high predation habitats swim with higher

coordination and in larger groups[39,40]. Indeed, guppies from higher predation populations have

been shown to rely more on social information for foraging resources, than those from lower

predation populations[41]. This is similar in the three-spined stickleback (*Gasterosteus aculeatus*),

where the transfer of information between conspecifics was more effective in more polarized

groups[42]. Further studies evaluating fitness effects of relying in social versus individual

information across different predation pressures is paramount to understand how anti-predator

behavior and collective motion drive evolutionary patterns at the proximate level.

In addition, across all tests performed we consistently found higher activity in polarization-selected

females. Previous assays in these fish evaluating maximal speed and endurance ruled out the

possibility that differences in collective motion patterns observed between polarization-selected

and control fish were driven by motor capacities[19]. In fact, activity is primarily associated with the

exchange of directional information according to a previous analysis of collective motion patterns

in guppies[39]. The difference in activity detected in our study may suggests that there are differences

in the selection lines in their ability to detect neighbor movements, or even to detect the threat

itself. Indeed, synchronized movements require an accurate detection and representation of the

near and far field of view around an individual to orient the body and maneuver accurately in

reaction to fast neighbor movements[5]. While this could possibly be due to differences we observe

in the primary visual center of the fish brain (the optic tectum), our morphological and

physiological tests performed to evaluate visual capacity indicate the opposite. Moreover, our

previous work using the same methods on brain size selection lines in guppies[30] indicates that

differences observed between polarization-selected and control females are not due to differences

in the acquisition of visual information. Rather, the combination of neuroanatomical, behavioral

and physiological data from our study suggests that in a predation context, effective decision-

making based on social information and effective processing of visual information to synchronize

swimming with close neighbors are central for the observed differences in anti-predator behavior.

_Brain morphology and collective behavior_

The study of brain anatomy in artificially selected fish allows to study brain function in relation to

behaviors that have important implications in the evolution of sociality. In our study, we found

two brain regions that were larger in relation to the rest of the brain in polarization-selected fish,

the optic tectum cup and the thalamus. The optic tectum is the terminus of a vast majority of optic

nerve fibers and axons of retinal cells[43] and as such is the primary vertebrate visual center. Despite

wide variation in optic tectum size across teleost species, this region functions to form instant

representations of the immediate surroundings[43]. This function is primarily achieved in superficial

layers of the tectum[44], which corresponds to the optic tectum cup region used in our

neuroanatomical parcellation of major brain regions in the guppy[45].

The evolved differences in optic tectum cup size between polarization-selected and control female

guppies we found are concordant with phenotypic plasticity findings in nine-spined stickleback

where it was found that individuals reared in groups developed larger optic tectum that those reared

individually[16]. Differences in the ability to acquire sensory input have previously been associated

with differences in schooling propensity[46]. In our experiment, rapid evolution of higher schooling

did not lead to changes in visual perception or eye morphology.  Together, these findings suggest

that differences found between polarization-selected and control lines in this section of the optic

tectum should have an effect in their ability to process visual information in order to control body

orientation during complex social maneuvers, but not in sensory information acquisition. This is

consistent with the role of the optic tectum in information processing in relation to the

telencephalon, a region of the brain commonly associated with decision-making in relation to

social behavior. Specifically, representation of the immediate surrounding in the optic tectum is

self-centered while the representation is allocentric in the telencephalon[43]. This self-centered

representation leads to important visuomotor computations of stimuli and can have important roles

in eye and body orienting as well as in predator evasion[47].

The thalamus was also enlarged in female guppies following artificial selection for higher

coordinated motion. While mostly studied in mammals, it seems that the thalamus plays prominent

roles in regulating attention and alertness and motor control through the modification, filtering and

distribution of sensory information into decision-making regions of the brain[48,49]. Recent findings

suggest similar functionality between homologous region of the mammalian and teleost brain as

defined in zebrafish as the *wider thalamus* region[48], and comparable to the guppy thalamic region

parcellated in our measurements. Specifically, the zebrafish thalamic region acts as the origin of

the main inhibitory neurons in the central nervous system (GABAergic neurons)[50], with an

important role in attenuating aggressiveness and the response to fear[51,52]. Our findings in anti-

predator behavior assays are consistent with functional convergence between the mammalian and

fish thalamic regions. The shorter time spent inspecting a predator threat observed in polarization-

selected fish (with larger thalamus) is likely explained by better ability to regulate alertness and

fear response towards a potential threat. While not directly addressed in this study, the regulatory

role of the thalamus in aggressiveness is concordant with common expectations of lower

aggression levels in group-living species (reviewed by [53]). As such, further quantifications of the

anatomical characteristics of the thalamus in relation to aggression levels within and across species

is a promising avenue for future research.

In contrast to the thalamus and the optic tectum cup, we found that the medulla oblongata was

smaller in polarization-selected lines. Consistent with this, in larval coral reef fishes, the inferior

and vagal lobes, which are subregions of the medulla, are larger in solitary as compared to more

social species[54]. The medulla oblongata is an important relay center of nervous signals between

the spinal cord and ascendant brain regions and has three core functions in teleosts. First, the

medulla oblongata functions in motor control through the presence of efferent motor neurons that

relay signals to the cerebellum[55]. Despite this function, we find no differences in motor control

capabilities between polarization-selected and control female guppies[19,56]. The medulla oblongata

also controls anti-predator responses through neuron firing in two large neurons present in this

region, the Mauthner-cells[57]. Interestingly, a previous study found that grouping reduces the

frequency of startle behavior, commonly observed in combination with predator inspections[58].

These findings are consistent with the reduction of predator inspection behavior we observed in

fish with higher schooling propensity. Finally, the medulla has a central function the processing

of somatosensory signals, with special emphasis in auditory and gustatory signals[59,60]. While not

tested in this study, it may be that the reduction in medulla observed in polarization-selected lines

might be associated with important changes in the auditory system and the ability to perceive

different tastes. In line with this reduction in the size of the medulla oblongata, our results show
that three other brain regions have significant hypoallometric relationships with the medulla (see
Supplementary Table 4b): the cerebellum (motor control center), the thalamus and the
hypothalamus (hormonal regulation center). Gene expression of *angiopoietin*-1, a locus implicated
in brain tissue development, showed contrasting expression levels between the medulla and the
thalamic and hypothalamic regions[61]. Based on this, we hypothesize that selection for more
coordinated motion leads to a trade-off between general sensory capabilities that are not important
in coordinated movements and specific sensory capabilities required to coordinate movement with
neighbors.

Unlike the auditory and taste systems, the mechanosensory system (lateral line) is important for
schooling through cues that allow fish to assess neighbor changes in speed and direction[62],
although this is more critical in low light and high turbidity conditions, which are very different to
our experiments[6]. In the future, it will be interesting to investigate the association between
schooling propensity, brain anatomy and potential trade-offs between sensory and
mechanosensory capacities.

**Conclusion**

Our empirical approach with behavioral assays on artificial selection lines with divergence in
polarization show that collective motion differences are consistent in the presence of a predator
threat and that predator inspection behaviour varies between the selection lines and the control
lines. Moreover, we reveal differences in neuroanatomy that could provide a mechanistic
explanation to the observed behavioural differences. Based on our discoveries, we propose that
changes in behavior are intimately intertwined with matching changes in brain morphology during
the evolution of collective behavior.

**Methods**

**Artificial selection for schooling propensity**

We evaluated the association between brain anatomy and collective motion in female guppies

following artificial selection for higher polarization. Extensive detail on the selection procedure

can be found in ([19]). In short, groups of female guppies were tested in repeated open field tests and

sorted in relation to the mean polarization of the group, the degree to which the individuals of a

group move with higher alignment[19,22,63]. For three generations, females from groups with higher

polarization were bred with males from those cohorts to generate three up-selected polarization

lines. In parallel, random females were exposed to the same experimental conditions and bred with

unselected males to generate three control lines. Third generation polarization-selected females

presented on average a 15% higher polarization and 10% higher group cohesiveness (i.e. 10%

shorter nearest neighbor distances) than control females[19]. The selection procedure targeted

polarization on female groups and we found a weaker response to selection in males, and therefore

subsequent neuroanatomical, behavioral and physiological studies focused on females. All fish

were removed from their parental tanks after birth, separated by sex at the first onset of sexual

maturation, and afterwards kept in single-sex groups of eight individuals in 7 L tanks containing

2 cm of gravel with continuously aerated water, a biological filter, and plants for environmental

enrichment. We allowed for visual contact between the tanks. The laboratory was maintained at

26°C with a 12-h light:12-h dark schedule. Fish were fed a diet of flake food and freshly hatched

brine shrimp daily.

**Anti-predator response in guppies following artificial selection**

*Collective motion*

We evaluated anti-predator behavior in polarization-selected and control female guppies by

conducting assays on 164 groups of eight fish in white arenas with 55 cm diameter and 3 cm water

depth. Each group was initially assessed in an open field assay in the arena for 10 minutes, and

collective motion data from these open field assays was previously used to analyze differences in

social interactions[19]. After 10 minutes, we sequentially introduced a novel object and a predator

model for 6-minute periods in the centre of the experimental arena. In half the assays, we

introduced the novel object first and the predator model second, with the order reversed in the

other half of the assays. We used a blue coffee mug as a novel object and a fishing lure custom-

painted to resemble the pike cichlid *Crenicichla frenata*, a natural predator on the guppy, as the
predator model. These objects have been previously used to successfully reproduce natural
behaviors of the guppy in response to a novel object and a predation threat[38]. Prior to the start of
the assay, the eight-fish group was confined in the centre of the arena for two minutes in an opaque
white 15 cm PVC cylinder. After this acclimation period, we lifted the cylinder and filmed the
arena using a Point Grey Grasshopper 3 camera (FLIR Systems; resolution, 2048 pixels by 2048
pixels; frame rate, 25 Hz).

We tracked the movement of fish groups in the collected video recordings using IDTracker[64] and
used fine-grained tracking data to calculate activity, polarization and attraction in Matlab 2020
following methods established in ([65]). These three variables characterize the two main axes of
collective motion in guppies, activity (speed) and sociability (polarization and attraction)[39]. For
activity, we calculated the median speed across all group members and frames in each assay. For
group polarization, we calculated the median global alignment, which indicates the angular
alignment of all fish in the arena. Calculations of median global alignment only considered frames
in which at least six individuals formed a connected group, with an interindividual distance of less
than 10cm counting as a connection. For attraction, we obtained the median distance to the nearest
neighbor for every fish across all frames. For all variables, we disregarded tracking data that did
not present a minimum of 16 consecutive tracked frames. To estimate the effect of the predator
model and novel object on group collective motion patterns, we additionally calculated group
polarization across all frames that contained reliable data for every group. We then generated a
heatmap with average values across polarization-selected and control groups that occurred within
20 x 20 mm grid cells (Fig. 1B). We used the centroid of the group to estimate group position
within the arena. Grid cells that did not contain values for a minimum of 8 groups per treatment
were disregarded. To evenly compare motion patterns when presented with a novel object and a
predator model to those obtained during the open field assays, we limited our analysis of the open
field assay data to the initial six minutes of the recording.

We used LMMs with median speed, polarization and attraction as dependent variables to test for
potential differences between polarization-selected and control lines. Selection regime, the type
of stimulus presented and the interaction between these two were included as fixed effects, and

body size of fish was coded as a covariate, with a random intercept for each replicated selection

line and the order of presentation of stimuli as random factors. All models were run in R

(v.4.1.3) using lme4 and lmerTest packages[66,67]. Model diagnostics showed that residual

distributions were roughly normal with no evidence of heteroscedasticity. We obtained post-hoc

comparisons of the response between selection line regimes at different levels of other fixed

effects in the previous models using the emmeans package[68] with the tukey-adjustment method

for multiple comparisons.

*Predator inspection behavior*

**Behavioral scoring.** Positional data and analyses of median distance to center in our data indicated

that groups of fish swam closer to the stimuli presented in the initial minutes following the addition

of a predator model in the experimental arena, when compared to the same time periods following

the addition of a novel object (Fig. 1A; Fig. 2A; Supplementary Figure 1). This observation

matched previous findings in similar experiments performed on guppies[38] and likely corresponds

to the stereotypical behavioral response of guppies to inspect and gain information of a potential

threat[23]. A predator inspection in guppies is characterized by an approach to the predator,

monitoring predator activity and swimming sideways with an arched body. Based on this

information, we manually visualized the videos during the first three minutes after addition of the

predator model and scored the behavior of one randomly selected fish in the group using BORIS[69].

While blind to the selection line treatment, the start and end time of each predator inspection

performed by the focal fish was scored for each video. We used the start and end time of predator

inspections to calculate the number of inspections, average inspection duration and the total time

that was spent inspecting per fish. Next, we fit a statistical model for each variable as a dependent

variable using a zero-inflated beta distribution and a logit link function for the conditional mean

in the package glmmTMb[70]. We used the selection line regime as a fixed effect. A random intercept

for each replicated selection line, and the order of presentation of the stimuli in the arena were

included as random factors in the model. We evaluated the adequacy of our fitted model using

scaled-residuals quantile-quantile plots, residual versus predicted values plots and a zero-inflation

test in the DHARMa package[71].

***Group collective motion during predator inspections.*** We analysed positional data for each group
by binning the observations in a grid, with cells being 20 x 20 mm. For each trial, we calculated a
density map, where the value for each grid cell was the fraction of all observations that occurred
within that cell. The resulting density maps are a normalized representation on how often each grid
cell was visited by individuals in our groups of 8 fish when exposed to different stimuli (Fig. 2A).
We used information from positional data to calculate summary statistics in different areas of
interest. Predator model assays presented unique spatial patterns in areas closer to the stimulus
presented, with higher densities in the tail area of the predator model (Fig. 2A). Based on these
factors, we calculated two new summary variables for each group: i) median polarization of the
group when the average position of the group was closer than 200 mm for predator model and
novel object assays; and ii) median polarization of the group in locations closer to the head (y-
position > 0) and the tail (y-position < 0) in predator model assays. We used LMMs with these
new calculated variables as dependent variables in the model to test for potential differences
between polarization-selected and control lines. Selection regime, the type of stimulus presented
or location in the tank were included as fixed effects, with a random intercept for each replicated
selection line and the order of presentation of stimuli as random factors. All models were run in R
(v.4.1.3) using lme4 and lmerTest packages[66,67]. Model diagnostics showed that residual
distributions were roughly normal with no evidence of heteroscedasticity. We obtained post-hoc
comparisons of the response between selection line regimes at different levels of other fixed effects
in the previous models using the emmeans package[68] with the tukey-adjustment method for
multiple comparisons.
594
595

**Brain morphology of female guppies following artificial selection**

We assessed neuroanatomical features of 15 polarization-selected and 15 control F3 fully-grown
females (6 months old), divided equally across polarization-selected and control lines. We used
microcomputed tomography (microCT, Skyscan 1172, Bruker microCT, Kontich, Belgium), and
reconstructed cross-sections from scanned images following a protocol successfully implemented
in a previous study evaluating neuroanatomical differences between guppies artificially up- and
down-selected for relative brain size[45]. This protocol allowed us to obtain measurements of whole

brain size volume and relative brain region volume in 11 major brain regions in the guppy

brain: olfactory bulbs, ventral telencephalon, dorsal telencephalon, thalamus, hypothalamus,

nucleus glomerulus, torus semicircularis, optic tectum cup, central optic tectum, cerebellum, and

medulla oblongata (Fig. 3B). Extended details on guppy brain region reconstruction from digital

images can be found in ([45]). Two brains from polarization-selected lines were damaged during the

protocol, which reduced the sample size to 28 samples. We tested for overall differences in whole

brain size between polarization-selected and control lines using a linear mixed model (LMM) with

brain volume as dependent variable, body size (standard length) as covariate, selection regime as

fixed effect, and replicate as random effect. For the brain regions, we used two different approaches

to determine whether selected and control lines differ in neuroanatomical features. First, we ran

11 independent LMMs with each region's volume as dependent variable, whole brain volume

(excluding volume of the region of interest) as covariate, selection regime as fixed effect, and

replicate as random effect. LMMs were run in R (v 4.1.3) using lme4 and lmerTest packages[66,67].

Second, to take into consideration that brain region volumes may be interdependent, we used a

more conservative approach and analyzed the data using a Bayesian multilevel model that included

11 brain regions as dependent variables in a fully multivariate context. The full model included an

analogous structure to those used in the independent LMMs for each brain region. Parameter

values were estimated using the brms interface[72,73] to the probabilistic programming language

Stan[74]. We used default prior distributions with student-t distribution (3, 0, 2.5) for all parameters.

The model estimated residual correlations among all brain region volumes with a Lewandowski-

Kurowicka-Joe (LKJ) prior with $\eta = 1$, which is uniform over the range $-1$ to $1$. Posterior

distributions were obtained using Stan's no-U-turn HMC with six independent Markov chains of

4000 iterations, discarding the first 2000 iterations per chain as warm-up and resulting in 12000

posterior samples overall. Convergence of the six chains and sufficient sampling of posterior

distributions were confirmed by a scale reduction factor below 1.01, and an effective size of at

least 10% of the number of iterations. For each model, posterior samples were summarized on the

basis of the Bayesian point estimate (median), SE (median absolute deviation), and posterior

uncertainty intervals by HDIs,

**Visual information processing in response to predation threat**

*Visual acuity*

We evaluated the ability to perceive detail (visual acuity) in 9-12 months old female guppies (59

polarization-selected, 57 control individuals) by assessing their optomotor response, an innate

orient behavior induced by whole-field visual stimulation[75]. Briefly, we projected a video

recording with rotating vertical black and white bands of six different widths (stimuli) on the

walls of a white ring-shaped arena of 25/50 cm of inner/outer diameter. Previous optimization of

the methods found that the use of these stimuli allowed us to evaluate the optomotor response at

the lower end of the species' acuity[30]. We placed individual fish in between the inner wall of the

arena and a transparent ring of 40 cm diameter. After a 2-min acclimation period, we recorded

their response towards 6 different rotating stimuli and the static images of these stimuli using a

Sony Cam HDR-DR11E recorder. Each stimulus was presented for 1 minute in random order.

Extended methods and the optimization procedure for the stimuli used here can be found in ([30]).

We manually scored the videos, recording the time that fish spent circling in the direction of the

stimuli (clockwise) at a constant moving pattern using BORIS[69]. Behavioral scoring was

performed blind to the treatment since only running numbers identified recordings. Likewise,

scoring was blind to the rotation and bandwidth of the stripes since only the fish, but not the

rotating stimuli, were visible during scoring. From the scoring, we calculated the proportion of

time that a fish spent swimming in the direction of rotation of the stimuli, out of the total time

that the different vertical black and white bands were presented to them.

*Temporal resolution*

Two weeks after visual acuity tests were completed in all fish, we measured the ability to track

movement stimuli of different speeds (temporal resolution) in the same females from the

polarization-selected (n = 58) and control lines (n= 55). We did not keep track of fish identity as

fish were kept in groups with conspecifics of the same selection line and replicate between

experiments. To evaluate temporal resolution, we placed fish in a white arena (50 cm diameter, 4

cm water depth) and exposed them to a projection of black and white bands of 3.5 cm width

rotating clockwise and counterclockwise at four different angular speeds (14.4, 25, 36 and 45

degrees/sec). The movement of each individual was recorded for a total of 1380 seconds with a

Sony Cam HDR-DR11E; a 300 seconds acclimation period and 1080 seconds of clockwise and

counterclockwise rotations of a projection with black and white bands at multiple speeds.

Specifically, during each individual test, the 8 stimulus combinations (4 speeds, 2 directions) were presented separately five independent times for 23 sec. The total time of each individual test was 920 sec (23 sec per stimulus x 5 times during the test x 8 stimulus combinations). We randomized the order of presentation of different stimuli a priori, but this order was consistent for all fish. The stimulus changed speed with smooth transitions of 3 sec, accelerating or decelerating to the next speed.

To quantify speed and direction changes of fish in our experimental setup, we automated behavioral scoring and obtained positional data using the Loopy Deep Learning Module (Loopbio 2020) in MATLAB (v. 2020a). X and y coordinates were transformed into a polar coordinate system centered on 0 and estimated from positional data. We calculated fish orientation by taking the difference in the fish's position between frames and defined their relative orientation (with respect to the arena) with the arcsin ($\sin(\theta - \vartheta)$), where $\theta$ was the orientation of the fish and $\vartheta$ is the angle of the arena radius going through the fish position. Positive values represent a fish swimming clockwise around the arena, while negative values represent swimming counterclockwise. For each frame, we identified whether the fish was swimming in the same direction of the stimulus projected and calculated the total proportion of time swimming in the direction of the stimulus. We also calculated the speed (in degrees per second) of the fish at each frame by using the dot product of the positional vector between consecutive frames. Using these values, we calculated for each individual the average total speed for each of the stimuli presented, and the average speed deviation between the speed of the stimulus presented and the speed of the fish.

*Eye morphology*

After the temporal resolution experiments were completed, we measured eye morphology in the females from polarization-selected lines (n = 57) and control lines (n =55) that were previously assessed for visual acuity and temporal resolution. For morphological measurements, we anesthetized fish with 0.2 mg/l of benzocaine and took pictures of their left side. We measured eye diameter and body length in these pictures using ImageJ[76]. Relative eye size was calculated as the ratio of these two variables. Image analyses were performed by a single scorer who was blind to the selection line treatment in the photographs.

*Statistics*

We analyzed potential differences in optomotor response, temporal resolution and eye morphology between polarization-selected and control females using LMM's. For the visual acuity trials, the proportion of time rotating was the dependent variable of the model. Fixed effects included selection line regime and bandwidth of the rotating stimuli. To account for differences in activity between fish, we used the proportion of time moving when presented with a static image of the stimuli as a covariate in the model. A random intercept for each replicated selection line, identity of the fish, and an observation-level variable were included as random factors in the model. For temporal resolution, we used selection line regime, the speed of rotation and the direction of rotation as fixed effects. The full model included the interaction between the selection regime with both speed and direction of rotation. This model included the identity of the fish, and a random intercept for each replicated selection line as random factors.

For eye morphology, eye diameter and relative eye size were dependent variables and models included selection regime as a fixed effect and a random intercept for each replicated selection line as a random factor. Model diagnostics showed that residual distributions were roughly normal with no signs of heteroscedasticity in optomotor response and eye morphology analyses. Model diagnostics on both models for temporal resolution analyses indicated unequal residual variance across the range of predicted values and a potential unequal influence of outliers. While estimates in linear mixed-effects models (LMMs) are argued to be robust to violations of such assumptions[77], we used the robustlmm package in R (Koller 2016) to compare the estimates obtained with LMM's to robust models with the same predictors that provide reduced weights to outliers in the data[78]. Results were consistent regarding the modelling approach (Supplementary Table 11). We obtained post-hoc comparisons of the response between selection line regimes at different levels of other fixed effects in the previous models using the emmeans package in R[68] with the tukey-adjustment method for multiple comparisons.

**Acknowledgements**

We thank David Sumpter for important contributions to the conceptualization of the artificial
selection procedure. We thank A. Rennie, E. Trejo, M. Amcoff and A. Boussard for help with fish
husbandry. **Funding**: This work was supported by the Knut and Alice Wallenberg Foundation
(102 2013.0072 to D.S., N.K., and K.P.), the Canada 150 Research Chair Program, and the
European Research Council (680951 to J.E.M), the Swedish Research Council (2016-03435 to
N.K., 2017-04957 to A.K., and 2018-04076 to J.H.-R.), and the Whitten Lectureship in Marine
Biology, University of Cambridge (to J.H.-R.). **Author contributions**: N.K., K.P and J.E.M.
contributed to conceptualization and funding acquisition of the project. A.K., A.S., M.R., S.D.B.,
J.H.-R., H.Z., K.P., and N.K. contributed to the design of the selection procedure. A.C.L, A.K.,
A.S., A.F.E., L.S.A, M.R., S.D.B. conducted research to obtain collective behavior data and
predator inspection data. A.K. H.L.Z. and K.P. conducted research to obtain neuroanatomical data.
A.C-L., M.G-O, and J.H-R. conducted research to obtain visual capacities data. A. C-L and
W.vdB. performed formal analyses and visualization. A.C-L., J.E.M, and N.K. wrote the original
draft. All authors contributed to the final version of the manuscript. **Competing interests**: The
authors declare that they have no competing interests. **Data and materials availability**: All data
and code needed to evaluate the conclusions in the paper will be deposited in an online repository.
Additional data related to this paper may be requested from the authors.

## 745 References

1.  Krause, J. & Ruxton, G. D. *Living in Groups*. (Oxford University press, 2002).
2.  Couzin, I. D., Krause, J., Franks, N. R. & Levin, S. A. Effective leadership and decision-
making in animal groups on the move. *Nature* **433**, 513–516 (2005).
3.  Jolles, J. W., Boogert, N. J., Sridhar, V. H., Couzin, I. D. & Manica, A. Consistent
Individual Differences Drive Collective Behavior and Group Functioning of Schooling
Fish. *Current Biology* **27**, (2017).
4.  Parrish, J. K. & Edelstein-Keshet, L. Complexity, Pattern, and Evolutionary Trade-Offs in
Animal Aggregation. *Science (1979)* **284**, 99–101 (1999).
5.  Strandburg-Peshkin, A. *et al.* Visual sensory networks and effective information transfer
in animal groups. *Current biology* **23**, R711 (2013).
6.  Pitcher, T. J. Functions of Shoaling Behaviour in Teleosts. in *The Behaviour of Teleost*
*Fishes* (ed. Pitcher T. J.) 294–337 (Springer, Boston, MA, 1986).
7.  Herbert-Read, J. E. *et al.* How predation shapes the social interaction rules of shoaling
fish. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20171126 (2017).
8.  Barton, R. A. Neocortex size and behavioural ecology in primates. *Proceedings of the*
*Royal Society B: Biological Sciences* **263**, 173–177 (1996).

9.   Burish, M. J., Kueh, H. Y. & Wang, S. S. H. Brain Architecture and Social Complexity in
Modern and Ancient Birds. *Brain Behav Evol* **63**, 107–124 (2004).
10.  Triki, Z., Levorato, E., McNeely, W., Marshall, J. & Bshary, R. Population densities
predict forebrain size variation in the cleaner fish Labroides dimidiatus. *Proceedings of*
*the Royal Society B* **286**, 20192108 (2019).
11.  Chee, S. S. A. *et al.* Social status, breeding state, and GnRH soma size in convict cichlids
(Cryptoheros nigrofasciatus). *Behavioural Brain Research* **237**, 318–324 (2013).
12.  Shaw, E. Schooling Fishes: The school, a truly egalitarian form of organization in which
all members of the group are alike in influence, offers substantial benefits to its
participants . *Am Sci* **2**, 166–175 (1978).
13.  Pollen, A. A. *et al.* Environmental complexity and social organization sculpt the brain in
Lake Tanganyikan cichlid fish. *Brain Behav Evol* **70**, 21–39 (2007).
14.  O'Connell, L. A. & Hofmann, H. A. Genes, hormones, and circuits: An integrative
approach to study the evolution of social behavior. *Front Neuroendocrinol* **32**, 320–335
(2011).
15.  Stednitz, S. J. *et al.* Forebrain Control of Behaviorally Driven Social Orienting in
Zebrafish. *Current biology* **28**, 2445-2451.e3 (2018).
16.  Gonda, A., Herczeg, G. & Merilä, J. Habitat-dependent and -independent plastic responses
to social environment in the nine-spined stickleback (Pungitius pungitius) brain.
*Proceedings of the Royal Society B: Biological Sciences* **276**, 2085–2092 (2009).
17.  Shinozuka, K. & Watanabe, S. Effects of telencephalic ablation on shoaling behavior in
goldfish. *Physiol Behav* **81**, 141–148 (2004).
18.  Loomis, C. *et al.* An Adult Brain Atlas Reveals Broad Neuroanatomical Changes in
Independently Evolved Populations of Mexican Cavefish. *Front Neuroanat* **13**, 88 (2019).
19.  Kotrschal, A. *et al.* Rapid evolution of coordinated and collective movement in response
to artificial selection. *Sci Adv* **6**, eaba3148 (2020).
20.  Magurran Anne E. *Evolutionary Ecology: The Trinidadian Guppy*. (Oxford University
Press, 2005).
21.  Bierbach, D. *et al.* Using a robotic fish to investigate individual differences in social
responsiveness in the guppy. *R Soc Open Sci* **5**, 181026 (2018).
22.  Szorkovszky, A. *et al.* Assortative interactions revealed by sorting of animal groups. *Anim*
*Behav* **142**, 165–179 (2018).
23.  Dugatkin, L. A. & Godin, J. G. J. Predator inspection, shoaling and foraging under
predation hazard in the Trinidadian guppy, Poecilia reticulata. *Environ Biol Fishes* **34**,
265–276 (1992).
24.  Veilleux, C. C. & Kirk, E. C. Visual Acuity in Mammals: Effects of Eye Size and
Ecology. *Brain Behav Evol* **83**, 43–53 (2014).
25.  Caves, E. M., Sutton, T. T. & Johnsen, S. Visual acuity in ray-finned fishes correlates with
eye size and habitat. *Journal of Experimental Biology* **220**, 1586–1596 (2017).
26.  Caves, E. M., Brandley, N. C. & Johnsen, S. Visual Acuity and the Evolution of Signals.
*Trends in Ecology and Evolution* vol. 33 358–372 Preprint at
https://doi.org/10.1016/j.tree.2018.03.001 (2018).
27.  Neave, D. A. The development of visual acuity in larval plaice (Pleuronectes platessa L.)
and turbot (Scophthalmusmaximus L.). *J Exp Mar Biol Ecol* **78**, 167–175 (1984).
28.  Anstis, S., Hutahajan, P. & Cavanagh, P. Optomotor test for wavelength sensitivity in
guppyfish (Poecilia reticulata). *Vision Res* **38**, 45–53 (1998).

29.   Haug, M. F., Biehlmaier, O., Mueller, K. P. & Neuhauss, S. C. F. Visual acuity in larval
zebrafish: Behavior and histology. *Front Zool* **7**, 1–7 (2010).
30.   Corral-López, A., Garate-Olaizola, M., Buechel, S. D., Kolm, N. & Kotrschal, A. On the
role of body size, brain size, and eye size in visual acuity. *Behav Ecol Sociobiol* **71**, 1–10
(2017).
31.   Corral-López, A. *et al.* Female brain size affects the assessment of male attractiveness
during mate choice. *Sci Adv* **3**, e1601990 (2017).
32.   Cronin, T. W., Johnsen, S., Marshall, N. J. & Warrant, E. J. *Visual Ecology. Visual*
*Ecology* (Princeton University Press, 2014).
33.   Kotrschal, A. *et al.* Artificial Selection on Relative Brain Size in the Guppy Reveals Costs
and Benefits of Evolving a Larger Brain. *Current Biology* **23**, 168–171 (2013).
34.   Fong, S. *et al.* Rapid mosaic brain evolution under artificial selection for relative
telencephalon size in the guppy (Poecilia reticulata). *Sci Adv* **7**, eabj4314 (2021).
35.   Handegard, N. O. *et al.* The dynamics of coordinated group hunting and collective
information transfer among schooling prey. *Current Biology* **22**, 1213–1217 (2012).
36.   Ioannou, C. C., Guttal, V. & Couzin, I. D. Predatory fish select for coordinated collective
motion in virtual prey. *Science (1979)* **337**, 1212–1215 (2012).
37.   Kotrschal, A. *et al.* Brain size affects female but not male survival under predation threat.
*Ecol Lett* **18**, 646–652 (2015).
38.   van der Bijl, W., Thyselius, M., Kotrschal, A. & Kolm, N. Brain size affects the
behavioural response to predators in female guppies (Poecilia reticulata). *Proceedings of*
*the Royal Society B: Biological Sciences* **282**, 20151132 (2015).
39.   Sumpter, D. J. T., Szorkovszky, A., Kotrschal, A., Kolm, N. & Herbert-Read, J. E. Using
activity and sociability to characterize collective motion. *Philosophical Transactions of*
*the Royal Society B: Biological Sciences* **373**, 20170015 (2018).
40.   Magurran, A. E., Seghers, B. H., Carvalho, G. R. & Shaw, P. W. Behavioural
consequences of an artificial introduction of guppies (Poecilia reticulata) in N. Trinidad:
Evidence for the evolution of anti-predator behaviour in the wild. *Proceedings of the*
*Royal Society B: Biological Sciences* **248**, 117–122 (1992).
41.   Clément, R. J. G. *et al.* Collective decision making in guppies: A cross-population
comparison study in the wild. *Behavioral Ecology* **28**, 919–924 (2017).
42.   MacGregor, H., Herbert-Read, J. & Ioannu, C. Information can explain the dynamics of
group order in animal collective behaviour. *Nat Commun* **11**, 1–8 (2020).
43.   Northmore, D. P. M. The Optic Tectum . *Encyclopedia of fish physiology: From genome*
*to environment* **1**, 131–142 (2011).
44.   Jones, M. R., Grillner, S. & Robertson, B. Selective projection patterns from subtypes of
retinal ganglion cells to tectum and pretectum: Distribution and relation to behavior.
*Journal of Comparative Neurology* **517**, 257–275 (2009).
45.   Kotrschal, A. *et al.* Evolution of brain region volumes during artificial selection for
relative brain size. *Evolution (N Y)* **71**, 2942–2951 (2017).
46.   Kowalko, J. E. *et al.* Loss of schooling behavior in cavefish through sight-dependent and
sight-independent mechanisms. *Current Biology* **23**, 1874–1883 (2013).
47.   Suzuki, D. G., Pérez-Fernández, J., Wibble, T., Kardamakis, A. A. & Grillner, S. The role
of the optic tectum for visually evoked orienting and evasive movements. *Proc Natl Acad*
*Sci U S A* **116**, 15272–15281 (2019).
48.   Mueller, T. What is the Thalamus in Zebrafish? *Front Neurosci* **6**, 64 (2012).

49.    Goldberg, J. H., Farries, M. A. & Fee, M. S. Basal ganglia output to the thalamus: Still a
856             paradox. *Trends Neurosci* **36**, 695–705 (2013).
857    50.    Achim, K., Salminen, M. & Partanen, J. Mechanisms regulating GABAergic neuron
858             development. *Cellular and Molecular Life Sciences* **71**, 1395–1415 (2014).
859    51.    Chou, M. Y. *et al.* Social conflict resolution regulated by two dorsal habenular subregions
860             in zebrafish. *Science (1979)* **352**, 87–90 (2016).
861    52.    Duboué, E. R., Hong, E., Eldred, K. C. & Halpern, M. E. Left Habenular Activity
862             Attenuates Fear Responses in Larval Zebrafish. *Current Biology* **27**, 2154–2162 (2017).
863    53.    Kutsukake, N. Complexity, dynamics and diversity of sociality in group-living mammals.
864             *Ecological Research* **24**, 521–531 (2009).
865    54.    Lecchini, D. *et al.* Variation in brain organization of coral reef fish larvae according to life
866             history traits. *Brain Behav Evol* **83**, 17–30 (2014).
867    55.    Morita, Y. & Finger, T. E. Topographic representation of the sensory and motor roots of
868             the vagus nerve in the medulla of goldfish, Carassius auratus. *J Comp Neurol* **264**, 231–
869             249 (1987).
870    56.    Vega-Trejo, R. *et al.* Artificial selection for schooling behaviour and its effects on
871             associative learning abilities. *Journal of Experimental Biology* **223**, 235093 (2020).
872    57.    Medan, V. & Preuss. T. The Mauthner-cell circuit of fish as a model system for startle
873             plasticity. *Journal of Physiology-Paris* **108**, 129–140 (2014).
874    58.    Fischer, E. K., Schwartz, A. J., Hoke, K. L. & Soares, D. Social context modulates
875             predator evasion strategy in guppies. *Ethology* **121**, 364–371 (2015).
876    59.    Kozloski, J. & Crawford, J. D. Transformations of an auditory temporal code in the
877             medulla of a sound- producing fish. *Journal of Neuroscience* **20**, 2400–2408 (2000).
878    60.    Finger, T. E. Evolution of gustatory reflex systems in the brainstems of fishes. *Integr Zool*
879             **4**, 53–63 (2009).
880    61.    Chen, Y. C. *et al.* Expression change in angiopoietin-1 underlies change in relative brain
881             size in fish. *Proceedings of the Royal Society B: Biological Sciences* **282**, 20150872
882             (2015).
883    62.    Partridge, B. L. & Pitcher, T. J. The sensory basis of fish schools: Relative roles of lateral
884             line and vision. *Journal of comparative physiology* **135**, 315–325 (1980).
885    63.    Szorkovszky, A. *et al.* An efficient method for sorting and quantifying individual social
886             traits based on group-level behaviour. *Methods Ecol Evol* **8**, 1735–1744 (2017).
887    64.    Pérez-Escudero, A., Vicente-Page, J., Hinz, R. C., Arganda, S. & de Polavieja, G. G.
888             idTracker: tracking individuals in a group by automatic identification of unmarked
889             animals. *Nature Methods 2014 11:7* **11**, 743–748 (2014).
890    65.    Kotrschal, A. *et al.* Brain size does not impact shoaling dynamics in unfamiliar groups of
891             guppies (Poecilia reticulata). *Behavioural Processes* **147**, 13–20 (2018).
892    66.    Bates, D., Sarkar, D., Bates, M. D. & Matrix, L. The lme4 Package. *R package version* **2**,
893             74 (2007).
894    67.    Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in
895             Linear Mixed Effects Models. *J Stat Softw* **82**, 1–26 (2017).
896    68.    Lenth, R. Package 'lsmeans'. *Am Stat* **34**, 216–221 (2018).
897    69.    Friard, O. & Gamba, M. BORIS: a free, versatile open-source event-logging software for
898             video/audio coding and live observations. *Methods Ecol Evol* **7**, 1325–1330 (2016).
899    70.    Brooks, M. E. *et al.* glmmTMB balances speed and flexibility among packages for zero-
900             inflated generalized linear mixed modeling. *R J* **9**, 378–400 (2017).

901    71.    Hartig, F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression
902           models. *R package version 0.3* (2018).
903    72.    Bürkner, P. C. Advanced Bayesian multilevel modeling with the R package brms. *R J* **10**,
904           395–411 (2018).
905    73.    Bürkner, P. C. brms: An R package for Bayesian multilevel models using Stan. *J Stat*
906           *Softw* **80**, 1–28 (2017).
907    74.    Carpenter, B. *et al.* Stan: A probabilistic programming language. *J Stat Softw* **76**, (2017).
908    75.    Smith, K. U. & Bojar, S. The nature of optokinetic reactions in mammals and their
909           significance in the experimental analysis of the neural mechanisms of visual functions.
910           *Psychol Bull* **35**, 193–219 (1938).
911    76.    Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
912           image analysis. *Nature Methods 2012 9:7* **9**, 671–675 (2012).
913    77.    Schielzeth, H. *et al.* Robustness of linear mixed-effects models to violations of
914           distributional assumptions. *Methods Ecol Evol* **11**, 1141–1152 (2020).
915    78.    Koller, M. Robustlmm: An R package for Robust estimation of linear Mixed-Effects
916           models. *J Stat Softw* **75**, 1–24 (2016).
917

**Figures**

**Fig 1. Collective motion patterns in female guppies artificially selected for higher schooling propensity. A)** Boxplots of median polarization, speed and nearest neighbor distance for groups of eight individuals of polarization-selected (pink) and control (blue) female guppies assayed in an open field test (OFT), with a novel object and with a predator model. Horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. Asterisks indicate $p < 0.05$ (see methods; Supplementary Tables 1-2). **B)** Heatmaps of group polarization across different locations of the experimental arena when control (top) and polarization-selected (bottom) groups were exposed to open field, novel object and predator model assays. Grid cells with data for less than 8 groups were not depicted.

**Fig 2. Social information processing in predator model assays. (A)** Density maps based on positional data of control (top) and polarization-selected (bottom) groups exposed to open field, novel object and predator model assays. **(B)** Boxplots of median group polarization in locations closer than 200 mm of the stimulus presented (left) and in the head and tail area of a predator model (right) in control (blue) and polarization-selected (pink) guppy groups. Boxplots of number of predator inspections **(C)**, total time inspecting **(D),** and mean inspection duration **(E)** for individuals when swimming in a group of eight polarization-selected (pink) and control (blue) female guppies in the presence of a predator model. Horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. Asterisks indicate p < 0.05 (see methods, Supplementary Tables 5-7).

**Fig 3. The effect of artificial selection for higher schooling propensity in neuroanatomical allometric relationships. (A)** The top left panel shows the allometric relationship between whole brain size volume and standard length of the fish (SL). Remaining panels show the relationship between each separate brain region with the rest of the brain ordered rostrally to caudally. Asterisks indicate brain regions with non-overlapping confidence intervals between polarization-selected females (pink; n = 13) and control females (blue; n = 15) in two consistent statistical analyses (Supplementary Tables 3-4). **(B)** Reconstructed brain regions from micro CT - scanned guppy brains. A dorsal (left) and lateral (middle) view of a guppy brain with the major brain regions color coded: olfactory bulbs (OB; dark blue), dorsal telencephalon (DT; red), ventral telencephalon (VT; light green), optic tectum (OT; yellow), hypothalamus (Hyp; turquoise), thalamus (Th; purple), cerebellum (Cb; brown), medulla oblongata (MO; dark green); as some regions are not visible from the outside a partially segmented and slightly tilted image (right) reveals: torus semicircularis (TS; orange), nucleus glomerulus (NG; lilac-blue), optic tectum core (OTc; light turquoise).

**Fig 4. Eye morphology and visual capacities of female guppies artificially selected for higher schooling propensity. (A)** Boxplots of eye morphological measurements. **(B)** Boxplots and density plots of the proportion of time following 6 different rotating stimuli with rotating and static gratings of different widths at the lower end of guppy visual acuity (thinner widths represent a higher degree of difficulty to be perceived). **(C)** Boxplots and density plots of the deviation of fish swimming speed in relation to the speed that a rotating stimulus presented. in polarization-selected. For all morphological measurements and vision assays we measured the same polarization-selected (pink; n = 57-59) and control females (blue; n = 55-57). In all boxplots, horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. Optomotor response average values not sharing any letter are significantly different (p < 0.05) in post-hoc contrasts (see Supplementary Table 9b). No significant differences were observed for any comparison between control and polarization-selected fish (see Supplementary Tables 8-10).

**Supplementary Figures for:**

**Evolution of schooling propensity in the guppy drives changes in anti-predator behavior that are linked to neuroanatomy**

Alberto Corral-Lopez*, Alexander Kotrschal, Alexander Szorkovszky, Maddi Garate-Olaizola, James Herbert-Read, Wouter van der Bijl, Maksym Romenskyy, Hong-Li Zeng, Severine Denise Buechel, Ada Fontrodona Eslava, Kristian Pelckmans, Judith E. Mank, Niclas Kolm

*corresponding author: corral@zoology.ubc.ca

**Fig S1. Proximity to stimuli in female guppies artificially selected for higher schooling propensity.** Boxplots of median distance to the stimulus combining data for groups of polarization-selected and control female guppies (n = 164) in predator model and novel object assays. Horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. Asterisks indicate p < 0.05.

**Fig S2. Temporal resolution of female guppies artificially selected for higher schooling propensity.** Boxplots and density plots of the deviation of fish swimming speed **(A),** and the proportion of time that fish followed the direction of the stimulus **(B)** in relation to four different rotating stimuli presented at different speeds that rotated clockwise and anti-clockwise to polarization-selected (pink; n = 58) and control females (blue; n = 56). In the boxplots, horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. No significant differences were observed for any comparison between control and polarization-selected fish (see Tables S9-S10).

**Supplementary Tables for:**

**Evolution of schooling propensity in the guppy drives changes in anti-predator behavior that are linked to neuroanatomy**

Alberto Corral-Lopez*, Alexander Kotrschal, Alexander Szorkovszky, Maddi Garate-Olaizola, James Herbert-Read, Wouter van der Bijl, Maksym Romenskyy, Hong-Li Zeng, Severine Denise Buechel, Ada Fontrodona Eslava, Kristian Pelckmans, Judith E. Mank, Niclas Kolm

*corresponding author: corral@zoology.ubc.ca

**Table S1**. Statistical tests for overall comparisons in tests evaluating polarization-selected and control female guppies in their shoaling patterns when exposed to an open field test (OFT), a novel object (cup) and a predator model.

### Polarization

| Predictors | Estimates | CI | t | p | df |
|---|---|---|---|---|---|
| (Intercept) | 0.80 | 0.68 – 0.91 | 14.54 | **< 0.001** | 17.59 |
| Selection [P] | 0.08 | 0.02 – 0.13 | 3.90 | **0.017** | 3.98 |
| treatment [Cup] | -0.12 | -0.15 – -0.10 | -9.16 | **< 0.001** | 338.42 |
| treatment [Predator] | -0.17 | -0.20 – -0.14 | -12.61 | **< 0.001** | 340.73 |
| Body size | -0.00 | -0.00 – -0.00 | -2.35 | **0.021** | 88.01 |
| Selection [P] * treatment [Cup] | -0.01 | -0.05 – 0.03 | -0.46 | 0.644 | 340.46 |
| Selection [S] * treatment [Predator] | -0.00 | -0.04 – 0.03 | -0.18 | 0.860 | 341.08 |

### Median speed

| Predictors | Estimates | CI | t | p | df |
|---|---|---|---|---|---|
| (Intercept) | 63.25 | 46.15 – 80.34 | 7.50 | **< 0.001** | 35.87 |
| Selection [P] | 13.63 | 8.09 – 19.16 | 5.96 | **< 0.001** | 6.29 |
| treatment [Cup] | -20.14 | -23.91 – -16.37 | -10.51 | **< 0.001** | 337.87 |

| Predictors | Estimates | CI | t | p | df |
|---|---|---|---|---|---|
| treatment [Predator] | -24.34 | -28.14 – -20.54 | -12.60 | **< 0.001** | 339.82 |
| Body size | -0.06 | -0.12 – 0.01 | -1.77 | 0.080 | 85.64 |
| Selection [P] * treatment [Cup] | -4.55 | -9.85 – 0.76 | -1.69 | 0.093 | 339.52 |
| Selection [S] * treatment [Predator] | -3.95 | -9.27 – 1.36 | -1.46 | 0.144 | 340.05 |

## Nearest neighbor distance

| Predictors | Estimates | CI | t | p | df |
|---|---|---|---|---|---|
| (Intercept) | 39.17 | 33.89 – 44.45 | 14.96 | **< 0.001** | 43.22 |
| Selection [P] | -3.43 | -5.34 – -1.53 | -4.55 | **< 0.001** | 5.31 |
| treatment [Cup] | -10.77 | -11.94 – -9.60 | -18.13 | **< 0.001** | 337.80 |
| treatment [Predator] | -10.88 | -12.06 – -9.70 | -18.17 | **< 0.001** | 339.69 |
| Body size | -0.01 | -0.03 – 0.01 | -0.63 | 0.530 | 105.88 |
| Selection [P] * treatment [Cup] | 2.88 | 1.24 – 4.53 | 3.45 | **< 0.001** | 339.38 |
| Selection [S] * treatment [Predator] | 2.82 | 1.17 – 4.47 | 3.37 | **< 0.001** | 339.88 |

**Table S2.** Independent contrasts for comparisons between polarization-selected and control female guppies in their shoaling patterns when exposed to an open field test (OFT), a novel object (cup) and a predator model

**Polarization**

| contrast | treatment | estimate | SE | df | lower.CL | upper.CL | t.ratio | p.value |
|---|---|---|---|---|---|---|---|---|
| C - P | OFT | -0.077 | 0.020 | 3.980 | -0.133 | -0.022 | -3.898 | **0.018** |
| C - P | Cup | -0.069 | 0.020 | 4.078 | -0.124 | -0.014 | -3.435 | **0.026** |
| C - P | Predator | -0.074 | 0.020 | 4.106 | -0.129 | -0.019 | -3.698 | **0.020** |

**Median speed**

| contrast | treatment | estimate | SE | df | lower.CL | upper.CL | t.ratio | p.value |
|---|---|---|---|---|---|---|---|---|
| C - P | OFT | -13.627 | 2.288 | 6.292 | -19.164 | -8.091 | -5.956 | **0.001** |
| C - P | Cup | -9.079 | 2.311 | 6.560 | -14.619 | -3.538 | -3.928 | **0.006** |
| C - P | Predator | -9.673 | 2.321 | 6.618 | -15.226 | -4.121 | -4.168 | **0.005** |

**Nearest neighbor distance**

| contrast | treatment | estimate | SE | df | lower.CL | upper.CL | t.ratio | p.value |
|---|---|---|---|---|---|---|---|---|
| C - P | OFT | 3.434 | 0.754 | 5.309 | 1.530 | 5.339 | 4.555 | **0.005** |
| C - P | Cup | 0.552 | 0.761 | 5.510 | -1.350 | 2.455 | 0.726 | 0.498 |
| C - P | Predator | 0.613 | 0.764 | 5.560 | -1.292 | 2.517 | 0.802 | 0.455 |

**Table S3.** Statistical tests for comparisons in tests evaluating polarization-selected and control female guppies in their anti-predator behavior when exposed to a predator model

### Number of inspections

| Predictors | Incidence Rate Ratios | CI | t | p |
|---|---|---|---|---|
| (Intercept) | 4.47 | 3.64 – 5.50 | 14.19 | **< 0.001** |
| Selection [P] | 0.87 | 0.75 – 1.01 | -1.85 | 0.0645 |

### Total time inspecting

| Predictors | Incidence Rate Ratios | CI | t | p |
|---|---|---|---|---|
| (Intercept) | 24.04 | 18.02 – 32.07 | 21.64 | **< 0.001** |
| Selection [P] | 0.79 | 0.66 – 0.95 | -2.52 | **0.011** |

### Mean inspection duration

| Predictors | Incidence Rate Ratios | CI | t | p |
|---|---|---|---|---|
| (Intercept) | 5.62 | 4.84 – 6.53 | 22.63 | **< 0.001** |
| Selection [P] | 0.82 | 0.69 – 0.96 | -2.49 | **0.013** |

**Table S4a.** Statistical tests for overall comparisons of group polarization in polarization-selected and control female guppies when the average position of the group was shorter than 200 mm to the stimulus presented in the arena in tests that exposed these fish to a predator model and a novel object (cup).

**Polarization (closer than 200 mm from the predator model)**

| Predictors | Estimates | CI | t | p | df |
|---|---|---|---|---|---|
| (Intercept) | 0.68 | 0.60 – 0.75 | 25.78 | **< 0.001** | 3.87 |
| Selection [P] | 0.07 | 0.02 – 0.12 | 2.96 | **0.003** | 273.80 |
| treatment [Predator model] | -0.12 | -0.16 – -0.08 | -5.34 | **< 0.001** | 146.76 |
| Replicate [Rep2] | 0.00 | -0.04 – 0.04 | 0.05 | 0.958 | 143.72 |
| replicate [Rep3] | 0.01 | -0.03 – 0.05 | 0.32 | 0.749 | 142.38 |
| Selection [P] * treatment [Predator model] | -0.03 | -0.08 – 0.03 | -0.83 | 0.405 | 143.99 |

**Table S4b.** Statistical tests for independent contrasts of group polarization in polarization-selected and control female guppies when swimming at a distance closer than 200 mm to the stimulus presented in the arena in tests that exposed these fish to a predator model and a novel object (cup).

| contrast | treatment | estimate | SE | df | lower.CL | upper.CL | t.ratio | p.value |
|---|---|---|---|---|---|---|---|---|
| C - P | Cup | -0.070 | 0.024 | 273.79 | -0.116 | -0.023 | -2.956 | **0.003** |
| C - P | Predator | -0.044 | 0.022 | 272.27 | -0.088 | -0.000 | -1.984 | **0.048** |

**Table S5a.** Statistical tests for overall comparisons of group polarization in polarization-selected and control female guppies when the average position of the group was located closer to the tail area or closer to the head area of the predator model

**Polarization**

| Predictors | Estimates | CI | Statistic | p | df |
|---|---|---|---|---|---|
| (Intercept) | 0.62 | 0.54 – 0.71 | 22.54 | **< 0.001** | 3.15 |
| Selection [P] | 0.04 | -0.03 – 0.10 | 1.53 | 0.191 | 4.58 |
| location [tail] | -0.11 | -0.14 – -0.07 | -5.47 | **< 0.001** | 156.96 |
| line [Social] * location [tail] | 0.04 | -0.01 – 0.09 | 1.64 | 0.103 | 150.99 |

**Table S5b.** Statistical tests for independent contrasts of group polarization in polarization-selected and control female guppies when the average position of the group was located closer to the tail area or closer to the head area of the predator model

| contrast | location | estimate | SE | df | lower.CL | upper.CL | t.ratio | p.value |
|---|---|---|---|---|---|---|---|---|
| C - P | Head | -0.037 | 0.024 | 4.58 | -0.102 | 0.027 | -1.534 | 0.191 |
| C - P | Tail | -0.080 | 0.023 | 3.63 | -0.146 | -0.013 | -3.483 | **0.030** |

**Table S6.** Results from independent Linear Mixed Models evaluating differences in relative brain and relative brain region size between polarization-selected and control female guppies.

| Region | Coefficent | Estimate | SE | df | t | P-value |
|---|---|---|---|---|---|---|
| Whole brain | Intercept | 10.706 | 0.929 | 22.992 | 11.520 | **< 0.001** |
| | Sel. Line (Polarization) | -0.015 | 0.037 | 23.296 | -0.415 | 0.682 |
| | Log (SL) | 1.581 | 0.292 | 22.899 | 5.401 | **< 0.001** |
| Olfactory bulbs | Intercept | -5.639 | 4.048 | 24.311 | -1.393 | 0.176 |
| | Sel. Line (Polarization) | -0.054 | 0.071 | 23.443 | -0.759 | 0.455 |
| | Log (rest of brain) | 0.973 | 0.257 | 24.319 | 3.782 | **< 0.001** |
| Ventral telencephalon | Intercept | -7.112 | 4.022 | 23.970 | -1.768 | 0.0897 |
| | Sel. Line (Polarization) | -0.054 | 0.070 | 23.366 | -0.762 | 0.453 |
| | Log (rest of brain) | 1.107 | 0.255 | 23.968 | 4.329 | **< 0.001** |
| Dorsal telencephalon | Intercept | -3.002 | 1.502 | 25.000 | -1.998 | 0.0567 |
| | Sel. Line (Polarization) | 0.029 | 0.027 | 25.000 | 1.096 | 0.2837 |
| | Log (rest of brain) | 0.935 | 0.095 | 25.000 | 9.780 | **< 0.001** |
| Thalamus | Intercept | -1.019 | 1.805 | 25.000 | -0.565 | 0.577 |
| | Sel. Line (Polarization) | 0.073 | 0.033 | 25.000 | 2.187 | **0.038** |
| | Log (rest of brain) | 0.950 | 0.115 | 25.000 | 8.197 | **< 0.001** |
| Hypothalamus | Intercept | -3.858 | 2.745 | 25.000 | -1.405 | 0.172 |
| | Sel. Line (Polarization) | 0.048 | 0.049 | 25.000 | 0.974 | 0.339 |
| | Log (rest of brain) | 1.098 | 0.175 | 25.000 | 6.259 | **< 0.001** |
| Nucleus glomerulus | Intercept | -2.352 | 9.716 | 25.000 | -0.242 | 0.811 |
| | Sel. Line (Polarization) | -0.067 | 0.176 | 25.000 | -0.381 | 0.706 |
| | Log (rest of brain) | 0.807 | 0.617 | 25.000 | 1.308 | 0.203 |
| Torus semicircularis | Intercept | -0.665 | 2.368 | 24.977 | -0.281 | 0.781 |
| | Sel. Line (Polarization) | -0.0325 | 0.042 | 24.106 | -0.758 | 0.456 |
| | Log (rest of brain) | 0.829 | 0.150 | 24.982 | 5.496 | **< 0.001** |
| Optic tectum cup | Intercept | 1.073 | 0.966 | 23.494 | 1.111 | 0.2780 |
| | Sel. Line (Polarization) | 0.042 | 0.017 | 23.095 | 2.409 | **0.024** |
| | Log (rest of brain) | 0.800 | 0.061 | 23.474 | 12.933 | **< 0.001** |
| Optic tectum core | Intercept | -2.883 | 1.658 | 25.000 | -1.739 | 0.0944 |
| | Sel. Line (Polarization) | 0.006 | 0.030 | 25.000 | 0.211 | 0.8347 |
| | Log (rest of brain) | 1.040 | 0.106 | 25.000 | 9.805 | **< 0.001** |
| Cerebellum | Intercept | -0.403 | 1.458 | 24.350 | -0.277 | 0.784 |
| | Sel. Line (Polarization) | -0.007 | 0.026 | 23.221 | -0.275 | 0.786 |
| | Log (rest of brain) | 0.908 | 0.093 | 24.358 | 9.703 | **< 0.001** |
| Medulla oblongata | Intercept | -0.145 | 1.519 | 24.920 | -0.096 | 0.9246 |
| | Sel. Line (Polarization) | -0.074 | 0.028 | 23.916 | -2.656 | **0.013** |
| | Log (rest of brain) | 0.969 | 0.099 | 24.935 | 9.761 | **< 0.001** |

**Table S7a** Results from a Bayesian multilevel model evaluating differences in relative brain region size between polarization-selected and control female guppies. Stars indicate estimates that do not include zero in the confidence interval range based on the posterior samples drawn from the model.

| Covariate | Estimate | Est.Error | l.95..CI | u.95..CI | |
|---|---|---|---|---|---|
| Medulla_Intercept | 0.19 | 0.32 | -0.37 | 0.77 | |
| Cerebellum_Intercept | 0.05 | 0.50 | -1.08 | 1.04 | |
| Nucleus glomerulus_Intercept | 0.09 | 0.48 | -0.84 | 1.04 | |
| Torus semicircularis_Intercept | 0.14 | 0.44 | -0.69 | 1.03 | |
| Thalamus_Intercept | -0.25 | 0.34 | -0.97 | 0.31 | |
| Optic tectum cups_Intercept | -0.19 | 0.51 | -1.26 | 0.89 | |
| Hypothalamus_Intercept | -0.06 | 0.39 | -0.84 | 0.73 | |
| Olfactory bulbs_Intercept | 0.13 | 0.54 | -0.94 | 1.26 | |
| Ventral telencephalon_Intercept | 0.05 | 0.63 | -1.23 | 1.43 | |
| Dorsal telencephalon_Intercept | -0.09 | 0.32 | -0.72 | 0.56 | |
| Optic tectum core_Intercept | -0.02 | 0.28 | -0.56 | 0.54 | |
| Medulla oblongata_Selection | -0.42 | 0.18 | **-0.79** | **-0.06** | * |
| Medulla oblongata_Rest of the brain | 0.91 | 0.09 | **0.74** | **1.09** | * |
| Cerebellum_Selection | -0.07 | 0.21 | -0.48 | 0.35 | |
| Cerebellum_Rest of the brain | 0.96 | 0.11 | **0.75** | **1.18** | * |
| Nucleus glomerolus_Selection | -0.21 | 0.40 | -0.98 | 0.58 | |
| Nucleus glomerulus_Rest of the brain | 0.34 | 0.20 | -0.06 | 0.75 | |
| Torus semicircularis_Selection | -0.24 | 0.32 | -0.87 | 0.38 | |
| Torus semicircularis_Rest of the brain | 0.72 | 0.16 | **0.40** | **1.04** | * |
| Thalamus_Selection | 0.49 | 0.23 | **0.04** | **0.94** | * |
| Thalamus_Rest of the brain | 0.92 | 0.11 | **0.70** | **1.16** | * |
| Optic tectum cups_Selection | 0.32 | 0.16 | **0.00** | **0.63** | * |
| Optic tectum cups_Rest of the brain | 0.88 | 0.08 | **0.72** | **1.06** | * |
| Hypothalamus_Selection | 0.12 | 0.26 | -0.39 | 0.64 | |

| | Estimate | Est.Error | l.95..CI | u.95..CI | |
|---|---|---|---|---|---|
| Hypothalamus_Rest of the brain | 0.87 | 0.13 | **0.61** | **1.14** | * |
| Olfactory bulbs _Selection | -0.27 | 0.33 | -0.93 | 0.38 | |
| Olfactory bulbs_Rest of the brain | 0.62 | 0.17 | **0.29** | **0.95** | * |
| Ventral telencephalon_Selection | -0.17 | 0.31 | -0.76 | 0.43 | |
| Ventral telencephalon_Rest of the brain | 0.58 | 0.16 | **0.27** | **0.89** | * |
| Dorsal telencephalon_Selection | 0.18 | 0.21 | -0.25 | 0.60 | |
| Dorsal telencephalon_Rest of the brain | 0.91 | 0.11 | **0.69** | **1.13** | * |
| Optic tectum core_Selection | 0.05 | 0.20 | -0.34 | 0.44 | |
| Optic tectum core_Rest of the brain | 0.93 | 0.10 | **0.73** | **1.13** | * |

**Table S7b.** Residual correlations of thalamus, optic tectum cups and medulla oblongata relative volume to other brain regions estimated from a Bayesian multilevel model evaluating differences in relative brain region size between polarization-selected and control female guppies. Stars indicate estimates that do not include zero in the confidence interval range based on the posterior samples drawn from the model.

## Medulla oblongata

| Brain region | Estimate | Est.Error | l.95..CI | u.95..CI |
|---|---|---|---|---|
| Cerebellum | -0.32 | 0.15 | **-0.59** | **-0.02** * |
| Nucleus glomerulus | -0.26 | 0.15 | -0.53 | 0.05 |
| Torus semicircularis | 0.07 | 0.16 | -0.25 | 0.37 |
| Thalamus | -0.40 | 0.14 | **-0.65** | **-0.12** * |
| Optic tectum cups | -0.04 | 0.17 | -0.36 | 0.29 |
| Hypothalamus | -0.50 | 0.12 | **-0.72** | **-0.23** * |
| Olfactory bulbs | -0.24 | 0.16 | -0.52 | 0.09 |
| Ventral telencephalon | -0.15 | 0.16 | -0.44 | 0.17 |
| Dorsal telencephalon | -0.12 | 0.17 | -0.43 | 0.21 |
| Optic tectum core | -0.22 | 0.15 | -0.51 | 0.09 |

## Optic tectum cups

| Brain region | Estimate | Est.Error | l.95..CI | u.95..CI |
|---|---|---|---|---|
| Medulla oblongata | -0.04 | 0.16 | -0.36 | 0.28 |
| Cerebellum | -0.07 | 0.17 | -0.41 | 0.26 |
| Nucleus glomerulus | -0.07 | 0.17 | -0.40 | 0.26 |
| Torus semicircularis | 0.05 | 0.17 | -0.29 | 0.38 |
| Thalamus | -0.18 | 0.16 | -0.49 | 0.15 |
| Hypothalamus | -0.15 | 0.16 | -0.46 | 0.17 |

| Brain region | Estimate | Est.Error | l.95..CI | u.95..CI |
|---|---|---|---|---|
| Olfactory bulbs | -0.11 | 0.17 | -0.45 | 0.23 |
| Ventral telencephalon | 0.02 | 0.17 | -0.32 | 0.36 |
| Dorsal telencephalon | -0.01 | 0.17 | -0.36 | 0.33 |
| Optic tectum core | -0.06 | 0.17 | -0.39 | 0.27 |

**Thalamus**

| Brain region | Estimate | Est.Error | l.95..CI | u.95..CI |
|---|---|---|---|---|
| Medulla oblongata | -0.40 | 0.14 | **-0.65** | **-0.12** * |
| Cerebellum | -0.27 | 0.16 | -0.55 | 0.05 |
| Nucleus glomerulus | 0.05 | 0.16 | -0.26 | 0.36 |
| Torus semicircularis | -0.27 | 0.16 | -0.56 | 0.06 |
| Optic tectum cups | -0.18 | 0.17 | -0.49 | 0.15 |
| Hypothalamus | -0.15 | 0.16 | -0.44 | 0.18 |
| Olfactory bulbs | -0.09 | 0.16 | -0.41 | 0.24 |
| Ventral telencephalon | 0.27 | 0.16 | -0.05 | 0.56 |
| Dorsal telencephalon | -0.05 | 0.17 | -0.37 | 0.28 |
| Optic tectum core | 0.10 | 0.16 | -0.22 | 0.41 |

**Table S8.** Statistical tests for comparisons in eye morphology between polarization-selected and control female guppies.

**Eye diameter**

| Predictors | Estimates | CI | Statistic | p | df |
|---|---|---|---|---|---|
| (Intercept) | 3.12 | 2.95 – 3.29 | 78.98 | **< 0.001** | 2.00 |
| Selection [P] | -0.03 | -0.24 – 0.19 | -0.52 | 0.658 | 2.00 |

**Eye diameter / body length**

| Predictors | Estimates | CI | Statistic | p | df |
|---|---|---|---|---|---|
| (Intercept) | 0.12 | 0.11 – 0.12 | 69.97 | **< 0.001** | 2.00 |
| Selection [P] | -0.00 | -0.01 – 0.01 | -0.13 | 0.906 | 2.00 |

**Table S9a**. Statistical tests for comparisons in visual acuity between polarization-selected and control female guppies.

**Optomotor response**

| Predictors | Estimates | CI | Statistic | p | df |
|---|---|---|---|---|---|
| (Intercept) | 0.10 | 0.03 – 0.17 | 3.16 | **0.008** | 12.06 |
| Selection [P] | 0.00 | -0.08 – 0.09 | 0.11 | 0.913 | 12.88 |
| Band width stimulus [0.28] | 0.07 | 0.00 – 0.14 | 2.09 | **0.036** | 570.82 |
| Band width stimulus [0.31] | 0.15 | 0.08 – 0.21 | 4.20 | **< 0.001** | 569.22 |
| Band width stimulus [0.35] | 0.12 | 0.05 – 0.19 | 3.51 | **< 0.001** | 569.28 |
| Band width stimulus [0.39] | 0.18 | 0.12 – 0.25 | 5.26 | **< 0.001** | 574.04 |
| Band width stimulus [0.44] | 0.17 | 0.10 – 0.23 | 4.75 | **< 0.001** | 572.07 |
| Optomotor response in static | 0.56 | 0.49 – 0.63 | 14.92 | **< 0.001** | 531.69 |
| Selection [P]* Band width stimulus [0.28] | 0.01 | -0.08 – 0.11 | 0.26 | 0.793 | 569.30 |
| Selection [P]* Band width stimulus [0.31] | 0.01 | -0.09 – 0.10 | 0.11 | 0.915 | 569.21 |
| Selection [P]* Band width stimulus [0.35] | 0.03 | -0.06 – 0.13 | 0.63 | 0.526 | 569.18 |
| Selection [P]* Band width stimulus [0.39] | -0.03 | -0.13 – 0.06 | -0.64 | 0.520 | 569.16 |
| Selection [P]* Band width stimulus [0.44] | 0.01 | -0.09 – 0.10 | 0.17 | 0.863 | 569.22 |

**Table S9b.** Independent contrasts for optomotor response observed in fish at rotating stimulus of several band widths

| contrast | estimate | SE | df | lower.CL | upper.CL | t.ratio | p.value |
|---|---|---|---|---|---|---|---|
| bw25 - bw28 | -0.079 | 0.024 | 571.167 | -0.149 | -0.009 | -3.236 | **0.016** |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| bw25 - bw31 | -0.148 | 0.024 | 569.618 | -0.218 | -0.079 | -6.094 | **< 0.001** |
| bw25 - bw35 | -0.137 | 0.024 | 569.230 | -0.207 | -0.068 | -5.637 | **< 0.001** |
| bw25 - bw39 | -0.168 | 0.025 | 577.744 | -0.238 | -0.097 | -6.818 | **< 0.001** |
| bw25 - bw44 | -0.169 | 0.024 | 573.616 | -0.239 | -0.099 | -6.918 | **< 0.001** |
| bw28 - bw31 | -0.069 | 0.024 | 569.653 | -0.139 | 0.000 | -2.852 | 0.051 |
| bw28 - bw35 | -0.058 | 0.024 | 570.338 | -0.128 | 0.011 | -2.390 | 0.161 |
| bw28 - bw39 | -0.089 | 0.024 | 571.492 | -0.159 | -0.019 | -3.639 | **0.004** |
| bw28 - bw44 | -0.090 | 0.024 | 569.612 | -0.160 | -0.021 | -3.712 | **0.003** |
| bw31 - bw35 | 0.011 | 0.024 | 569.262 | -0.058 | 0.081 | 0.460 | 0.997 |
| bw31 - bw39 | -0.019 | 0.025 | 574.211 | -0.089 | 0.051 | -0.791 | 0.969 |
| bw31 - bw44 | -0.021 | 0.024 | 571.155 | -0.091 | 0.049 | -0.858 | 0.956 |
| bw35 - bw39 | -0.031 | 0.025 | 575.985 | -0.101 | 0.040 | -1.245 | 0.814 |
| bw35 - bw44 | -0.032 | 0.024 | 572.350 | -0.102 | 0.038 | -1.315 | 0.777 |
| bw39 - bw44 | -0.002 | 0.024 | 569.833 | -0.071 | 0.068 | -0.064 | 1.000 |

**Table S9c.** Independent contrasts for optomotor response observed in polarization-selected and control female guppies at rotating stimulus of several band widths.

| contrast | stimulus | estimate | SE | df | lower.CL | upper.CL | t.ratio | p.value |
|---|---|---|---|---|---|---|---|---|
| C - P | bw25 | -0.004 | 0.038 | 601.455 | -0.078 | 0.070 | -0.112 | 0.911 |
| C - P | bw28 | -0.017 | 0.038 | 601.441 | -0.091 | 0.057 | -0.456 | 0.649 |
| C - P | bw31 | -0.009 | 0.038 | 601.422 | -0.083 | 0.064 | -0.247 | 0.805 |
| C - P | bw35 | -0.035 | 0.038 | 601.457 | -0.109 | 0.039 | -0.938 | 0.349 |
| C - P | bw39 | 0.027 | 0.038 | 601.458 | -0.047 | 0.101 | 0.722 | 0.471 |
| C - P | bw44 | -0.013 | 0.038 | 601.452 | -0.086 | 0.061 | -0.340 | 0.734 |

**Table S10a.** Statistical tests for comparisons in visual temporal resolution between polarization-selected and control female guppies.

**Speed deviation from stimulus rotation**

| Predictors | Estimates | CI | Statistic | p | df |
|---|---|---|---|---|---|
| (Intercept) | 8.38 | 3.23 – 13.53 | 5.31 | **0.014** | 2.87 |
| Selection [P] | -0.46 | -8.74 – 7.83 | -0.19 | 0.863 | 2.64 |
| Speed [24] | -5.93 | -7.71 – -4.15 | -6.54 | **< 0.001** | 4398.86 |
| Speed [36] | -23.39 | -25.17 – -21.61 | -25.78 | **< 0.001** | 4398.86 |
| Speed [45] | -49.74 | -51.52 – -47.96 | -54.79 | **< 0.001** | 4398.84 |
| Rotation[Counterclockwise] | -7.11 | -8.36 – -5.85 | -11.08 | **< 0.001** | 4398.89 |
| Selection [P] * Speed [24] | 0.31 | -2.17 – 2.79 | 0.25 | 0.805 | 4398.91 |
| Selection [P] * Speed [36] | 1.26 | -1.22 – 3.74 | 0.99 | 0.320 | 4398.91 |
| Selection [P] * Speed [45] | -0.48 | -2.96 – 1.99 | -0.38 | 0.701 | 4398.84 |
| Selection [P] * Rotation[Counterclockwise] | 1.29 | -0.47 – 3.04 | 1.44 | 0.150 | 4398.98 |

**Proportion of time following the stimulus**

| Predictors | Estimates | CI | Statistic | p | df |
|---|---|---|---|---|---|
| (Intercept) | 0.87 | 0.80 – 0.94 | 39.43 | **< 0.001** | 3.12 |
| Selection [P] | 0.00 | -0.14 – 0.15 | 0.10 | 0.928 | 2.52 |
| Speed [24] | 0.03 | 0.00 – 0.06 | 2.03 | **< 0.001** | 4398.95 |
| Speed [36] | 0.03 | -0.00 – 0.05 | 1.83 | 0.067 | 4398.95 |
| Speed [45] | -0.04 | -0.06 – -0.01 | -2.55 | **0.018** | 4398.92 |
| Rotation[Counterclockwise] | -0.07 | -0.09 – -0.05 | -7.33 | **< 0.001** | 4398.98 |
| Selection [P] * Speed [24] | 0.01 | -0.02 – 0.04 | 0.74 | 0.457 | 4399.06 |
| Selection [P] * Speed [36] | -0.00 | -0.04 – 0.03 | -0.20 | 0.843 | 4398.99 |
| Selection [P] * Speed [45] | -0.00 | -0.04 – 0.04 | -0.15 | 0.884 | 4398.99 |

| Selection | | | | | | | |
|---|---|---|---|---|---|---|---|
| Selection [P] * Rotation[Counterclockwise] | -0.01 | -0.05 – 0.03 | -0.62 | 0.536 | 4398.92 | | |

**Table S10b.** Independent contrasts for speed deviation to stimulus rotation observed in polarization-selected and control females at multiple rotation speeds in clockwise and counterclockwise directions

| Selection | Speed | Rotation | emmean | SE | df | lower.CL | upper.CL | t.ratio | p.value |
|---|---|---|---|---|---|---|---|---|---|
| C | 14.4 | Clockwise | 8.379 | 1.577 | 2.874 | 3.233 | 13.526 | 5.313 | **0.015** |
| P | 14.4 | Clockwise | 7.922 | 1.938 | 2.485 | 0.966 | 14.879 | 4.089 | **0.038** |
| C | 24 | Clockwise | 2.447 | 1.577 | 2.872 | -2.701 | 7.594 | 1.552 | 0.222 |
| P | 24 | Clockwise | 2.303 | 1.938 | 2.488 | -4.650 | 9.256 | 1.188 | 0.336 |
| C | 36 | Clockwise | -15.012 | 1.577 | 2.872 | -20.160 | -9.865 | -9.520 | **0.003** |
| P | 36 | Clockwise | -14.212 | 1.938 | 2.488 | -21.164 | -7.259 | -7.332 | **0.010** |
| C | 45 | Clockwise | -41.357 | 1.577 | 2.874 | -46.503 | -36.210 | -26.223 | **< 0.001** |
| P | 45 | Clockwise | -42.299 | 1.938 | 2.485 | -49.255 | -35.342 | -21.830 | **0.001** |
| C | 14.4 | Counterclockwise | 1.272 | 1.577 | 2.873 | -3.875 | 6.420 | 0.807 | 0.481 |
| P | 14.4 | Counterclockwise | 2.103 | 1.938 | 2.487 | -4.850 | 9.057 | 1.085 | 0.372 |
| C | 24 | Counterclockwise | -4.660 | 1.577 | 2.869 | -9.809 | 0.489 | -2.956 | 0.063 |
| P | 24 | Counterclockwise | -3.517 | 1.939 | 2.491 | -10.465 | 3.432 | -1.814 | 0.186 |
| C | 36 | Counterclockwise | -0.483 | -22.119 | 1.577 | 2.869 | -27.268 | -14.030 | **0.001** |
| P | 36 | Counterclockwise | -20.031 | 1.939 | 2.491 | -26.979 | -13.083 | -10.331 | **0.004** |
| C | 45 | Counterclockwise | -48.464 | 1.577 | 2.873 | -53.611 | -43.317 | -30.731 | **< 0.001** |
| P | 45 | Counterclockwise | -48.118 | 1.938 | 2.487 | -55.071 | -41.164 | -24.827 | **< 0.001** |

**Table S11.** Statistical results using a robust linear mixed model approach for comparisons in the proportion of time following the correct direction of the stimulus in visual temporal assays between polarization-selected and control female guppies.

**Proportion of time following the stimulus**

| Predictors | Estimates | Std. Error | t-value | p | df |
|---|---|---|---|---|---|
| (Intercept) | 0.91 | 0.016 | 57.45 | **< 0.001** | 3.12 |
| Selection [P] | -0.001 | 0.031 | -0.04 | 0.971 | 2.52 |
| Speed [24] | 0.014 | 0.009 | 1.61 | 0.106 | 4398.95 |
| Speed [36] | 0.027 | 0.009 | 3.07 | **0.001** | 4398.95 |
| Speed [45] | -0.010 | 0.009 | -1.17 | **0.240** | 4398.92 |
| Rotation[Counterclockwise] | -0.041 | 0.006 | -6.51 | **< 0.001** | 4398.98 |
| Selection [P] * Speed [24] | 0.004 | 0.012 | 0.33 | 0.744 | 4399.06 |
| Selection [P] * Speed [36] | 0.004 | 0.012 | 0.33 | 0.741 | 4398.99 |
| Selection [P] * Speed [45] | -0.009 | 0.012 | -0.74 | 0.460 | 4398.99 |
| Selection [P] * Rotation[Counterclockwise] | -0.001 | 0.009 | -0.15 | 0.882 | 4398.92 |

Robustness weights for the residuals:
 3546 weights are ~= 1. The remaining 974 ones are summarized as
  Min. 1st Qu. Median   Mean 3rd Qu.   Max.
 0.199  0.328  0.518  0.557  0.773  0.999

Robustness weights for the random effects:
 101 weights are ~= 1. The remaining 18 ones are summarized as
  Min. 1st Qu. Median   Mean 3rd Qu.   Max.
 0.247  0.489  0.739  0.693  0.904  0.995

Rho functions used for fitting:
 *Residuals*:
   eff: smoothed Huber (k = 1.345, s = 10), sig: smoothed Huber, Proposal II (k = 1.345, s = 10)
 *Random Effects, variance component 1 (trial)*:
   eff: smoothed Huber (k = 1.345, s = 10), vcp: smoothed Huber, Proposal II (k = 1.345, s = 10)
 *Random Effects, variance component 2 (rep)*:
   eff: smoothed Huber (k = 1.345, s = 10), vcp: smoothed Huber (k = 1.345, s = 10)