Testing Wright's Intermediate Population Size Hypothesis – When Genetic Drift is a Good Thing

Mitchell B. Cruzan

Portland State University, Department of Biology

Contact: cruzan@pdx.edu

Abstract – In his 1931 monograph, Sewall Wright predicted genetic drift would overwhelm selection in
small populations, and selection would dominate in large ones, but he also concluded drift could
facilitate selection in populations of intermediate size. The idea that drift and selection would act
together in populations of intermediate size has been almost completely ignored even as empirical
evidence of rapid evolution associated with population bottlenecks has continued to accumulate. I used
forward-time simulations with random mating and discrete generations to test the hypothesis that drift
can facilitate selection. I find that selection generates biased distributions of Δq, and this bias is greatest
for population sizes between 20 and 200, resulting in drift facilitation. Drift facilitation reduces the
accumulation of drift load and segregation load for populations of intermediate size. Small populations
accumulated higher levels of drift load and large populations maintained high levels of segregation load.
Fixation of beneficial mutations is facilitated in intermediate populations when dominance is low and
selection is weak. Accumulation of beneficial mutations over time (fixation flux) was higher across small
to intermediate size populations and declined rapidly for large populations. Compared to predictions of
fixation time for codominant beneficial mutations from diffusion equations, drift facilitation accelerates
fixation in populations of intermediate size, but fixation time is slower in large populations when
selection is weak. These results suggest drift facilitation in small and intermediate populations promote
purging of genetic load and fixation of beneficial mutations, and may result in relatively rapid adaptation
compared to large populations.

Impact Summary – After the recognition of Gregor Mendel's contributions to our understanding of the
inheritance of genetically-determined traits around 1900, there was confusion as to whether the type of
variation Mendel studied could account for evolution by natural selection, as described by Charles
Darwin. This controversy was resolved when three theoreticians (Ronald Fisher, Sewell Wright, and
J.B.S. Haldane) published books that integrated Mendelian genetics with evolution. Their contributions
(referred to as the Modern Synthesis), focused on evolutionary processes occurring within and among
populations of a species, and established a mathematical foundation for our understanding of
evolutionary biology. The mathematical models developed by the three architects of the modern
synthesis, and those who followed, predicted that the effects of natural selection would be
overwhelmed by random genetic changes (referred to as Genetic Drift) in small populations, and that
genetic drift would be minimal, while selection would be most effective in large populations. Even
though one of Wright's major conclusions was that genetic drift and selection would work together
(Drift Facilitation) to promote adaptive evolution in intermediate-sized populations, this idea has been
almost completely ignored since it was first introduced in 1931. In this study, I use simulations of
evolution in natural populations to evaluate the potential for drift facilitation to promote evolution in
intermediate-sized populations. My work largely confirms Wright's predictions; the removal of
deleterious mutations and promotion of adaptive evolution are enhanced in population sizes ranging
from about 10 to 200. These results indicate that a new paradigm for our understanding of evolution

within populations is needed – one that emphasizes the importance of drift facilitation in small and

intermediate size populations, and recognizes that periods of reduced population size are opportunities

for enhanced levels of adaptive evolution.

Keywords: Adaptative evolution, conservation genetics, drift facilitation, drift load, inbreeding load,

population bottleneck, purging, segregation load, simulations.

"In a population of intermediate size … there is continual random shifting of gene frequencies … which

leads to a relatively rapid, continuing, irreversible, and largely fortuitous, but not degenerative series of

changes, even under static conditions." Wright 1931, Page 157.

Introduction

The mathematical models for population genetics developed during the Modern Synthesis (Fisher 1930;

Wright 1931; Haldane 1932) and beyond (Crow 1987; Ewens 2004; Charlesworth and Charlesworth

2017) have provided a foundation for a general understanding of evolutionary processes accounting for

divergence among populations, adaptation, and diversification among lineages. A fundamental aspect of

this framework is the dichotomy between the roles of genetic drift and selection; specifically, that drift

governs the accumulation of neutral and degenerative variation while only selection can promote the

purging of deleterious mutations and the fixation of beneficial mutations (Crow and Kimura 1970;

Kimura 1983). The general prediction is genetic drift will dominate the dynamics of allele frequency

change in small populations when selection is weak (i.e. when $s < 1/2N$), resulting in the accumulation of

genetic load and limited potential for adaptive evolution. In contrast, selection is expected to

overwhelm the effects of drift in large populations leading to fixation of beneficial mutations and the

purging of genetic load (Charlesworth 2002; Gossmann et al. 2012; Lanfear et al. 2014; Charlesworth

and Charlesworth 2017). However, in his 1931 monograph, Wright emphasized the idea that the rate of

fixation of beneficial mutations would be much slower in large populations, but there was potential for

genetic drift and selection to act together to promote rapid adaptation in populations of "intermediate"

size. He proposed a range of population sizes where both drift and selection are effective enough to

allow "fortuitous" shifts in allele frequencies to facilitate selection resulting in accelerated accumulation

of adaptive genetic variation. This idea (hereafter referred to as "drift facilitation") was seized upon by

George Gaylord Simpson (1944), who proposed small population size was a likely explanation for the

lack of fossilization during rapid transitions that resulted in the origin of novel forms, including new

genera and families. This idea of "quantum evolution" (i.e. periods of rapid evolution when selection is

facilitated by genetic drift; Simpson 1944) inspired a number of evolutionary biologists, who

subsequently identified a wide range of examples of adaptation associated with reduced population size

(Mayr 1954; Lewis and Raven 1958). Verne Grant (1963, 1981) expanded on the idea of quantum

evolution to include rapid speciation events in small, isolated populations at the periphery of a

widespread species' range (budding or peripatric speciation; Mayr 1954). While interest in the idea of

quantum evolution has waned, evidence of rapid evolution associated with population bottlenecks

continues to accumulate (e.g., Carvalho et al. 1996; Rosenblum et al. 2007; Tepolt et al. 2009; Marsico et

al. 2011; Cruzan 2019; Rego et al. 2019; Cruzan et al. 2021; Mahrt et al. 2021; Yin et al. 2021; van der

Zee et al. 2022). Surprisingly, there have been no attempts to assess the potential for drift facilitation of

selection in populations of intermediate size using analytical or numerical approaches.

In this study, I use forward-time, individual-based simulations of populations to evaluate the potential
for drift facilitation to promote evolution in intermediate-sized populations. I evaluate the effects of
selection (s), dominance (h), population size ($N = N_e$), and the composite variable of population size and
selection (Ns) on the accumulation of deleterious and beneficial mutations. In all cases, generations are
discrete, and populations are assumed to be dioecious with equal sex ratios and random mating.
Changes in allele frequencies and fixation of alleles are examined for a set of unlinked, biallelic loci that
have independent effects on fitness. For genetic load, I examine the proportion of loci that became fixed
for deleterious mutations (drift load), and the proportion of deleterious mutations that remain
segregating after a number of generations (segregation load). In separate simulations, I examine the
proportion of loci fixing for beneficial mutations, the rate of fixation of individual mutations, and fixation
flux (the rate at which beneficial alleles accumulate; Otto and Whitlock 1997). My results indicate that
drift facilitation of selection promotes purging of genetic load and the accumulation of beneficial
mutations across a range of small and intermediate population sizes.

Methods
Forward-time, individual-based simulations were conducted using the SimuPOP module (Peng and
Kimmel 2005) in Python and the COEUS High-Performance Computing Cluster at Portland State
University. A set of N individuals, each homozygous for a single locus, were generated using the
sim.Population simulator and mutations introduced ($a \rightarrow a'$) that were affected by varying levels of
selection (s) and dominance (h). Transmission of alleles between generations was governed by the sim.
MendelianGenoTransmitter. Selection was imposed at the time of mating by defining the fitness of each
diploid genotype using MapSelector in SimuPOP. Depending on the goals of the simulation, mutations
were either introduced once in the first generation, or at a per-generation rate determined by μ. Each
population size of N individuals was replicated k times. For each replication, the proportion of
individuals with fixed mutations (q_{fixed}), and the average allele frequency across individuals ($\bar{q}$) were
extracted after g generations. Two sets of simulations were run; the first set examined levels of drift and
segregation load, and the second set focused on beneficial mutations by evaluating the proportion of
loci fixed for mutations, the rate of fixation of individual mutations, and the fixation flux over g
generations.

Genetic Load – The frequency of fixed deleterious mutations (drift load) across loci and of the
proportion of loci segregating for deleterious mutations after g generations (segregation load) was
assessed for population sizes ranging from $N = 2$ to 5000. For each of $k = 1000$ replications for each
population size, s was randomly generated from a Poisson distribution (range: $0.0 - 0.25$; $\bar{s} = 0.07$), and
h was generated from a uniform distribution between 0.0 and 0.2 (assuming that deleterious mutations
are generally recessive with $\bar{h} = 0.10$; Willis 1999; García-Dorado and Caballero 2000; Agrawal and
Whitlock 2011; Ruzicka et al. 2021). Fitness of the *aa, aa',* and *a'a'* genotypes were set to 1.0, *1-hs*, and
1-s, respectively. The initial allele frequency of *a'* was set at $q = 0$ and mutations were allowed to
accumulate for *g = 1000* generations with $\mu = 1x10^{-3}$ This assumes that each locus consists of thousands
of base pairs (i.e. if the per bp/generation mutation rate is on the order of *$1x10^{-9}$*), but the same
qualitative results are obtained for lower mutation rates over longer timeframes.

The proportion of loci fixed for deleterious mutations (q_{fixed}) and selection (s) for each mutation were
used to estimate the fitness effects of fixed deleterious mutations (drift load) for each replication of

each population size as the reduction in fitness due to the selection coefficient weighted by the
proportion of loci fixed for the mutation;

$$w_{fixed} = 1 - sq_{fixed}$$

Segregation load was assessed as the inbreeding effects on fitness for segregating mutations (inbreeding
depression; Charlesworth and Willis 2009; Willi et al. 2013) for 1000 replicates for each population size
after 1000 generations. Inbreeding depression was estimated based on the mean fitness of hypothetical
progeny that would be the result of self-fertilization compared to progeny from random outcrossing.
Fitness after selfing was estimated as,

$$\overline{w}_s = 2\bar{p}\bar{q}(1 - (0.25 + 0.5hs + 0.25s)^F);$$

where $\bar{p}$ and $\bar{q}$ are the mean frequencies of the normal (a) and mutant (a') alleles across segregating loci
(individuals), and F is the number of segregating sites at the end of 1000 generations. I calculated the
expected fitness of progeny from random outcrossing within each population as,

$$\overline{w}_o = 1 - (\bar{p}^2 + hs2\bar{p}\bar{q} + s\bar{q}^2)^F$$

Segregation load was then estimated for each N as inbreeding depression for selfed progeny compared
to outcrossed progeny (Charlesworth and Willis 2009) as,

$$\delta_{s/o} = 1 - \frac{\overline{w}_s}{\overline{w}_o}$$

These estimates of drift load and segregation load assume independent fitness effects across unlinked
loci.

Beneficial Mutations – I examined the processes of accumulation of beneficial mutations in populations
using simulations to evaluate the probability of fixation of beneficial mutations, the rate of fixation of
individual alleles, and the fixation flux for a range of population sizes (N), selection (s), dominance (h),
and the combination of population size and selection (Ns). For beneficial mutations ($a \rightarrow a'$), the fitness
of the *aa, aa',* and *a'a'* genotypes were set to *1-s*, *1-hs*, and *1.0*, respectively.

The first simulations compared the fixation of beneficial alleles across a range of initial allele frequencies
from *q = 0.01* to *0.50* when *Ns = 2* for population sizes of *N = 100* (*s = 0.02*) and *N = 1000* (*s = 0.002*).
Initial frequency of the *a'* allele was set at *q = 1/2N* (*μ = 0*) and the proportion of loci fixing for the
mutant allele was compared to the diffusion prediction for the probability of fixation (Φ_q) derived by
Kimura (1957; 1962). Alleles were codominant in simulations (*h = 0.5*) to match the assumptions of the
diffusion approximation;

$$\Phi_q = \frac{1 - \exp(-2Nsq)}{1 - \exp(-2Ns)}$$

A second simulations examined the probability of fixation over population sizes ranging from $N = 2$ to
4000 with *s* randomly drawn from a Poisson distribution (*s = 0.0* to *0.25*) and *h* drawn from a uniform
distribution (*h = 0.0* to *0.2*) as described above. Initial frequency of the *a'* allele was set at $q = 1/2N$ ($\mu =$
0) and the proportion of loci fixing for the mutant allele was recorded after 1000 generations. The
results indicated the greatest degree of drift facilitation occurs when $N \approx 100$ (see below). Consequently,
I considered the probability of fixation over a broader range of dominance (*h = 0.0* to *1.0*) and initial
allele frequencies (*q = 0.0* to *0.5*) for a population size of $N = 100$ with $k = 10$ replicates for each
combination of *q* and *h*.
183
184 I examined the rate of fixation for beneficial alleles starting at an initial frequency of $q = 1/2N$ ($\mu = 0$)
185 across a range of population sizes from $N = 2$ to *1000* for *s = 0.02* to *0.20* and with *h = 0.5*. The observed
186 numbers of generations required for fixation ($\bar{t}_{gens}$ across $k = 10$ replicates per population size) from
187 these simulations were compared to the predicted rate of fixation ($\bar{t}$) from the diffusion approximation
188 (Ewens 1979; Otto and Whitlock 2013);
189

190
$$\bar{t} \approx \int_{1/2N}^{0.999} \frac{(e^{4Nsx} - 1)(e^{4Ns(1-x)} - 1)}{sx(1-x)(e^{4Ns} - 1)} dx$$

191
192 This integral is undefined when the upper limit is 1.0, and the approach to fixation is asymptotic, so an
193 arbitrary upper limit of 0.999 was chosen for comparison to fixation frequencies from the simulations.
194
While this approach describes fixation rates for individual alleles, it does not account for the
contribution of the number of mutations entering populations each generation based on population size
for the accumulation of beneficial mutations over time (fixation flux; Otto and Whitlock 1997). In
particular, we might expect that more mutations would occur in larger populations and consequently
flux should increase with population size. I examined fixation flux for population sizes ranging from $N = 4$
to *3000* with $\mu = 1x10^{-4}$, $Ns = 2$ ($s = 2/N$), and *h = 0.5* with 1000 loci and 10 replicates per population size
for time periods of $g = 1000, 2000,$ and *4000* generations. Fixation flux under selection was compared to
fixation with only genetic drift (*s = 0*) over 1000 generations across the same range of population sizes. I
estimated the mean and standard deviation of *Δq* across generations for beneficial mutations that
ultimately fixed in populations across a range of populations sizes ($N = 4$ to *3000*) and selection
coefficients (*s = 0.01* to *0.20* for $N = 100$ and *1000*.
206
Results
Genetic Load – Fixation frequencies for populations subject to mutation were highest for small
populations (< 100), and was effectively zero for population sizes larger than $N = 3000$ (Fig. 1A).
Mutations fixing in populations > 100 had negligible fitness effects so that drift load was near zero for
populations sizes greater than $N = 100$ (Fig. 1B). Larger proportions of loci continued to segregate for
deleterious mutations after 1000 generations in populations of less than $N = 10$ and more than $N = 75$,
while segregation load was lower in intermediate-sized populations (*10 > N < 75*; Fig. 2).
214
Beneficial Mutations – There was a close fit between the diffusion approximation for the probability of
fixation across initial allele frequencies (Φ_q) and the proportion of loci fixed for beneficial mutations
with population sizes of 100 and 1000 with *Ns = 2* and codominance (*h = 0.5*; Fig. 3). When beneficial

alleles were recessive, the frequency of fixation was low for *N < 30* and *N > 1000*, and was greatest when
population size was between 100 and 200 (Fig. 4). Fixation frequencies were generally greater than the
diffusion approximation for *N = 100* and *Ns = 2* when *h < 0.5* across a range of initial allele frequencies,
but was lower for dominant mutations when starting at a low initial allele frequency (Fig. 5).
222
The rate of fixation of individual alleles ($\bar{t}_{gens}$) was similar to the diffusion approximation ($\bar{t}$) when
selection was stronger (*s = 0.1* and *0.2*), but substantially faster for population sizes *N > 10*, but was
slower when *N > 100* when *s = 0.04*, and for *N > 200* when *s = 0.02* (Fig. 6). Accumulation of beneficial
mutations increased over longer time periods (fixation flux was greater after 4000 generations
compared to 2000, and 1000 generations; Fig. 7A). Fixation flux and the loss of beneficial mutations was
higher when *N < 100*, but generally declined with increasing population size (*h = 0.5*; Figs. 7A, B). The
number of mutations that remained segregating was not strongly affected by the length of the time
period, and was consistently low when *N < 100*, and increased rapidly when *N > 100* (Fig. 7C). Fixation
flux was much higher for *Ns = 4* compared to very weak (*Ns = 0.2*) or no selection (Appendix Fig. S1). The
higher rates of fixation flux correspond closely with higher means and standard deviations of *Δq* across
population sizes of *N < 100* (Figs. S2, S3). Mean *Δq* increased with the strength of selection, but the
standard deviation remained fairly constant across a range from *s = 0.01* to *0.20* (Figs. S4, S5). When
selection was held constant (*s = 0.02*), fixation flux increased rapidly and loss of beneficial mutations
decreased rapidly for populations sizes less than *N = 1000*, and then plateaued across larger populations
(Appendix Fig. S6).
238
Discussion
Forward-time simulations of natural populations indicate that segregation load is lower, and rates of
fixation and fixation flux of beneficial mutations are higher for small- to intermediate-sized populations
(*N < ~100*) compared to larger populations. Drift load was relatively high for small populations, but was
negligible for populations larger than *N* = 100. Results of simulations for beneficial mutations were
consistent with diffusion approximations of the probability of fixation when *h = 0.5* (Kimura 1957;
Kimura 1962), but fixation probabilities for recessive mutations were higher than predicted in
intermediate-sized populations (*N = 100*). Observed rates of fixation matched the diffusion
approximation ($\bar{t}$) fairly well when selection was strong (*s ≥ 0.10* and *h = 0.5*), but rates of fixation were
substantially faster when *s < 0.10* for population sizes between *N = 10* and *100*, and slower for
population sizes of *N > 100*. The accumulation of beneficial mutations (fixation flux) was higher across
small populations, and declined rapidly when population size was greater than *N = 100* with constant *Ns*
across population sizes. Fixation flux was greater under selection than for neutral alleles, which indicates
that genetic drift facilitates selection to promote fixation of beneficial alleles even in very small
populations. These results suggest the roles of drift and selection across population sizes is more
complex than predictions from analytical models, and periods of reduced population size may represent
opportunities for enhanced levels of purging, and for rapid acquisition of adaptive genetic variation.
256
The results presented here generally support Wright's (1931) predictions about the potential for
interactions between drift and selection to promote rapid fixation of beneficial alleles, but the
population sizes where drift facilitation of selection is effective are smaller than he predicted. Wright
specifically defined intermediate populations as ranging between *Ns = 2.5* and *20* (page 148). Simpson
(1944) also assumed that the dynamics described by Wright would apply to very large population sizes

(i.e. up to *N = 25,*000) compared to the range of population sizes where drift facilitation was found to be
effecting in the simulations presented. In contrast to the predictions of Wright and Simpson, higher
rates and probabilities of fixation of beneficial mutations were observed in simulations for *Ns = 2* and
less. Moreover, drift facilitation was effective only for populations sizes less than *N = 100*, regardless of
value of *Ns*. This is an important result because *Ns* has been used to represent the compensatory effects
between genetic drift and selection in analytical models including diffusion equation analyses of
population genetic processes (Kimura 1957; Kimura 1962; Ewens 2004). Given the results presented
above, more careful evaluation of the separate effects of selection and population size may be
warranted.
271
While population genetic models have mostly ignored the potential for selection in small populations,
empirical evidence for the effectiveness of selection during population bottlenecks continues to
accumulate. For example, there are cases of rapid adaptation to novel conditions during establishment
when population sizes were small including in artificial introductions of Trinidadian guppies (Carvalho et
al. 1996; van der Zee et al. 2022), Eastern fence lizards on novel substrates in New Mexico (Rosenblum
et al. 2007), European green crabs on the west coast of North America (Tepolt et al. 2009), the cactus-
feeding moth, *Cactoblastis cactorum*, in Florida (Marsico et al. 2011), steelhead trout in Lake Michigan
(Willoughby et al. 2018), the invasive grass, *Brachypodium sylvaticum*, in Oregon (Cruzan 2019; Marchini
et al. 2019), during evolutionary rescue in the seed beetle *Callosobruchus maculatus* (Rego et al. 2019),
and after migration into Lake Champlain by the Atlantic sea lamprey (Yin et al. 2021). In one case, there
was rapid evolution of antibiotic resistance in *Pseudomonas aeruginosa* even when population
bottlenecks were severe and selection was weak (Mahrt et al. 2021). Rapid acquisition of adaptive
genetic variants also seems to have occurred in a case of recent budding speciation (peripatric
speciation; Mayr 1954) in a range-restricted species of buttercup (Cruzan et al. 2021). In all of these
cases, adaptation to novel conditions appears to have occurred quickly during periods of reduced
population size and suggests drift facilitation of selection may be a widespread phenomenon.
288
One of the more surprising results of simulations was the observation of higher rates of fixation flux of
beneficial mutations when *N < 1000*, suggesting drift facilitation can enhance rates of adaptation in
small to intermediate size populations. This result is in stark contrast to diffusion analyses, which
predicts flux estimated as *4NsU* (where *U* is the genome-wide mutation rate) will remain constant across
population sizes if *Ns* is held constant, and will increase with population size if *s* is held constant (Otto
and Whitlock 1997; Ewens 2004). Contrary to expectations from diffusion estimates, fixation flux was
high across smaller populations, but declined in larger populations (*N > 100*) while the strength of
selection relative to population size remained constant (*Ns = 2*). With constant selection (*s = 0.02*), flux
increased rapidly across smaller population sizes, but then appeared to be limited by demographic
constraints (i.e. the time required for allele replacement in a large population; Otto and Whitlock 1997)
when population size was greater than *N ≈ 500*.
300
Consistent with expectations from diffusion analyses, fixation flux was higher when selection was
stronger (i.e. when *Ns = 4*), closely matched neutral flux when selection was very weak (i.e., when *Ns =
0.2*; and *s < 1/2N*), and fixation of deleterious mutation contributing to drift load increased for
population sizes less than *N = 1500*. In contrast, drift facilitation increased fixation flux in small
populations even under the assumptions of diffusion equation estimates of fixation probability (i.e. with

h = 0.5 and *Ns* constant across *N*), and the influence of drift facilitation for the probability of fixation for
beneficial mutations was even greater when dominance was low (*h < 0.5*) and selection was weak (*s <*
0.1).
309
The observation of higher flux in small populations is consistent with some previous analyses that
predicted genetic drift and selection could interact to increase rates of divergence among populations
(Cohan 1984; Lynch 1986), and with selection experiments in maize and simulations where enhanced
rates of fixation of small effect mutations were observed when subjected to high genetic drift (Desbiez-
Piat et al. 2021). While diffusion analyses have led to numerous insights for evolution in populations,
alternative approaches may provide analysis of a broader range of parameters including population size
in combination with different levels of selection and dominance.
317
Previous studies that have found evidence of drift facilitation did not identify its causes (Cohan 1984;
Lynch 1986), but the evaluation of *Δq* across population sizes sheds some light on the mechanisms
responsible for enhancement of fixation rates in small populations. The mean and standard deviation of
Δq increased in smaller populations suggesting they are at least partly responsible for the observed
effects on allele frequency change and fixation. For mutations at intermediate frequencies, it appears
that "fortuitous" fluctuations due to drift are promoted by selection and generate a "ratcheting effect"
that biases and accelerates changes in allele frequencies. This idea is supported by the observation that
the mean of *Δq* is positively associated with selection. The effects of drift facilitation are different when
allele frequencies are close to *q = 0* (for deleterious mutations) or fixation (i.e. *q = 1* for beneficial
mutations). When allele frequencies are close to fixation, higher levels of drift in smaller populations
increase the probability for random fluctuations to result in absorption, while minimal drift in larger
populations results in persistent segregation. Consequently, larger populations appear to accumulate
higher levels of inbreeding load and standing variation for beneficial mutations while having lower drift
load and rates of fixation flux compared to smaller populations (i.e. for *N < 100*).
332
The observation that large populations maintain higher levels of segregation load and standing variation
for beneficial mutations suggests that episodes of reduced population size may be disproportionally
important for rapid evolution in novel or changing environments. Because of higher frequencies of
segregating mutations, large populations entering bottlenecks would have enhanced potential for
purging of segregation load and for the fixation of beneficial mutations. Moreover, reductions in
population size due to bottlenecks and founder events are likely to be associated with increased
environmental stress, and consequently we expect selection to often increase during population
bottlenecks (Fowler and Whitlock 2002; Matute 2013). Higher frequencies of segregating mutations
along with drift facilitation may be responsible for elevated levels of purging following population
bottlenecks (Bouzat 2010), and may help explain the substantial number of cases of rapid adaption
during episodes of reduced population size (e.g., Carvalho et al. 1996; Rosenblum et al. 2007; Tepolt et
al. 2009; Marsico et al. 2011; Cruzan 2019; Rego et al. 2019; Cruzan et al. 2021; Mahrt et al. 2021; Yin et
al. 2021; van der Zee et al. 2022). On the other hand, we expect population bottlenecks and founder
events to result in elevated accumulation of drift load (Lynch et al. 1995; Willi et al. 2013), but the
simulations presented here indicate the potential for purging segregation load and the fixation of
deleterious mutations depends on the severity of the reduction in population size. It is notable that the
standard guideline of a minimal effective population size of *N = 50* (i.e. the 50/500 rule; Jamieson and

Allendorf 2012) is too small to prevent the accumulation of drift load, but is in the right range for drift
facilitation of beneficial mutations. Further analyses may help resolve more specific effects of changes in
population size for adaptation to novel conditions and for the purging of deleterious mutations.
353
Conclusions
The results of individual-based, forward-time simulations are consistent with a wide range of
observations from natural populations, but contrast with many of the predictions from models based on
diffusion equations (Ewens 2004). In general, it appears that Wright's (1931) intuitive predictions about
the potential for genetic drift to facilitate selection were correct, but the population sizes where drift
facilitation is most effective are smaller than he suggested. Furthermore, while Wright saw the potential
for weak deleterious mutations to fix in small populations, he did not anticipate that drift facilitation
could lead to more effective purging of segregation load. One of the most striking results from these
simulations, and one that stands in stark contrast to widely-accepted assumptions about the effects of
population size for adaptive evolution (Charlesworth 2002; Bouzat 2010; Gossmann et al. 2012; Lanfear
et al. 2014), is that fixation flux of beneficial mutations is substantially higher in small populations ($N <$
100) compared to larger populations under the assumption that Ns remains constant. The assumption
selection is stronger in small populations may generally hold true especially under conditions where
population declines are precipitated by increased levels of environmental stress. Moreover, the
probability of fixation and rate of fixation of beneficial mutations were higher when $h < 0.5$ and selection
was weak, so the prediction of rapid evolution due to drift facilitation is not dependent on the
assumption of increased selection in small populations.
371
The potential for rapid evolution in small populations may be further enhanced as large populations
experience reductions in population size. While it is generally assumed population bottlenecks will result
in decreased fitness due to the accumulation of genetic load, and that there will be limited potential for
adaptive evolution, this is not always true (Bouzat 2010). The simulation results presented here indicate
drift facilitation of selection in small to intermediate size populations leads to effective removal of
segregating deleterious mutations, and increases the rate and probability of fixation for beneficial
mutations. While there has been some speculation and evidence for enhanced evolution in small
populations (Simpson 1944; Grant 1981; Cohan 1984; Lynch 1986), no previous study has provided a
detailed characterization of the potential for drift facilitation to promote rapid evolution.
381
Results from these simulations emphasize the importance of the adaptive evolution potential in small
populations. Furthermore, more attention should be paid to the consequences of deleterious mutations
as they depend heavily on population size; a clear distinction needs to be made between the fixation of
deleterious mutations contributing to drift load in small populations, and the persistent of deleterious
mutations in large populations resulting in inbreeding depression due to segregation load. The results of
these simulations highlight genetic drift as a generative, beneficial process that promotes evolution in
small populations. Alternatively, the lack of drift in large populations effectively stalls purging of
segregation load and the fixation of beneficial mutations. While further analyses of drift facilitation are
needed, the results presented here give more weight to Wright's (1931, 1932) suggestion that adaptive
evolution is more likely to occur in small populations (Wade and Goodnight 1998), then to Fisher's
(1930) argument that large populations are more important for fixation of beneficial mutations (Coyne
et al. 1997). Further work on evolution in small populations from modeling and empirical perspectives

will clarify the potential for drift facilitation to promote evolution, and ultimately determine whether
the current paradigm for our understanding of population genetic processes of evolution needs to be
refined.
397
398
399
400

401 Acknowledgements
402 I thank Michael Wade and Michael Whitlock for suggestions on the development of simulations and
403 relevant parameters to focus on. Karla de Lima Berg, Elizabeth Hendrickson, Jaime Schwoch, and Ariana
404 Walczyk provided suggestions to improve the manuscript and presentation of results. This research was
405 supported by NSF-DEB award 2051235.

406
407 Author Contributions
408 MBC is solely responsible for the development of the SimuPOP simulations in Python, data analyses, and
409 writing of the manuscript.
410
411 Data Accessibility
412 All Python scripts and data generated will be deposited in Data Dryad …
413
414

415   Literature Cited
416
Agrawal, A. F. and M. C. Whitlock. 2011. Inferences about the distribution of dominance drawn from
yeast gene knockout data. Genetics 187:553-U274.
(http://doi.org/10.1534/genetics.110.124560).
Bouzat, J. L. 2010. Conservation genetics of population bottlenecks: the role of chance, selection, and
history. Conserv. Genet. 11:463-478. (http://doi.org/10.1007/s10592-010-0049-0).
Carvalho, G. R., P. W. Shaw, L. Hauser, B. H. Seghers, and A. E. Magurran. 1996. Artificial introductions,
evolutionary change and population differentiation in Trinidadian guppies (Poecilia
reticulata:Poeciliidae). Biol. J. Linn. Soc. 57:219-234. (http://doi.org/10.1111/j.1095-
8312.1996.tb00310.x).
Charlesworth, B. 2002. Effective population size. Curr. Biol. 12:R716-R717.
(http://doi.org/10.1016/S0960-9822(02)01244-7).
Charlesworth, B. and D. Charlesworth. 2017. Population genetics from 1966 to 2016. Heredity 118:2-9.
(http://doi.org/10.1038/hdy.2016.55).
Charlesworth, D. and J. H. Willis. 2009. The genetics of inbreeding depression. Nat. Rev. Genet. 10.
(http://doi.org/10.1038/nrg2664).
Cohan, F. M. 1984. Can uniform selection retard random genetic dif=vergence between isolated
conspecific populations? Evolution 38:495-504. (http://doi.org/10.2307/2408699).
Coyne, J. A., N. H. Barton, and M. Turelli. 1997. A critique of Sewall Wright's shifting balance theory of
evolution. Evolution 51:643-671.
Crow, J. F. 1987. Population genetics history: A personal view. Annu. Rev. Genet. 21:1-22.
(http://doi.org/10.1146/annurev.ge.21.120187.000245).
Crow, J. F. and M. Kimura. 1970. An Introduction to Population Genetics Theory. Harper & Row, New
York.
Cruzan, M. B. 2019. How to make a weed - the saga of the slender fasle brome invasion in the North
American west and lessons for the future. Bioscience 69:469-507.
(http://doi.org/https://doi.org/10.1093/biosci/biz051).
Cruzan, M. B., P. G. Thompson, N. A. Diaz, E. C. Hendrickson, K. R. Gerloff, K. A. Kline, H. M. Machiorlete,
and J. M. Persinger. 2021. Weak coupling among barrier loci and waves of neutral and adaptive
introgression across an expanding hybrid zone. Evolution 75:3098-3114.
(http://doi.org/https://doi.org/10.1111/evo.14381).
Desbiez-Piat, A., A. Le Rouzic, M. I. Tenaillon, and C. Dillmann. 2021. Interplay between extreme drift
and selection intensities favors the fixation of beneficial mutations in selfing maize populations.
Genetics 219. (http://doi.org/10.1093/genetics/iyab123).
Ewens, W. J. 1979. Mathematical Population Genetics. Springer-Verlag, Berlin.
Ewens, W. J. 2004. Mathmatical Population Genetics, Volume I: Theoretical Foundations. Springer-
Verlag, Berlin.
Fisher, R. A. 1930. The Genetical Theory of Natural Selection. Clarendon Press, Oxford.
Fowler, K. and M. C. Whitlock. 2002. Environmental stress, inbreeding, and the nature of phenotypic and
genetic variance in Drosophila melanogaster. Proceedings. Biological sciences / The Royal
Society 269:677-683. (http://doi.org/10.1098/rspb.2001.1931).
García-Dorado, A. and A. Caballero. 2000. On the average coefficient of dominance of deleterious
spontaneous mutations. Genetics 155:1991-2001.
(http://doi.org/10.1093/genetics/155.4.1991).
Gossmann, T. I., P. D. Keightley, and A. Eyre-Walker. 2012. The effect of variation in the effective
population size on the rate of adaptive molecular evolution in eukaryotes. Genome Biol Evol
4:658-667. (http://doi.org/10.1093/gbe/evs027).

Grant, V. 1963. The Origin of Adaptations. Columbia University Press, New York.
Grant, V. 1981. Plant Speciation. Columbia University Press, New York.
Haldane, J. B. S. 1932. The Causes of Evolution. Princeton University Press, Princeton.
Jamieson, I. G. and F. W. Allendorf. 2012. How does the 50/500 rule apply to MVPs? Trends Ecol. Evol.
27:578-584. (http://doi.org/https://doi.org/10.1016/j.tree.2012.07.001).
Kimura, M. 1957. Some problems of stochastic processes in genetics. The Annals of Mathematical
Statistics 28:882-901, 820.
Kimura, M. 1962. On the probability of fixation of mutant genes in a population. Genetics 47:713-719.
(http://doi.org/10.1093/genetics/47.6.713).
Kimura, M. 1983. The Neutral Theory of Molecular Evolution. Cambridge University Press, Cambridge.
Lanfear, R., H. Kokko, and A. Eyre-Walker. 2014. Population size and the rate of evolution. Trends Ecol.
Evol. 29:33-41. (http://doi.org/http://dx.doi.org/10.1016/j.tree.2013.09.009).
Lewis, H. and P. H. Raven. 1958. Rapid evolution in *Clarkia*. Evolution 12:319-336.
(http://doi.org/10.2307/2405854).
Lynch, M. 1986. Random drift, uniform selection, and the degree of population differentiation. Evolution
40:640-643. (http://doi.org/10.2307/2408585).
Lynch, M., J. Conery, and R. Burger. 1995. Mutational meltdowns in sexual populations. Evolution
49:1067-1080. (http://doi.org/10.2307/2410432).
Mahrt, N., A. Tietze, S. Kunzel, S. Franzenburg, C. Barbosa, G. Jansen, and H. Schulenburg. 2021.
Bottleneck size and selection level reproducibly impact evolution of antibiotic resistance. Nature
Ecology & Evolution 5. (http://doi.org/10.1038/s41559-021-01511-2).
Marchini, G. L., C. A. Maraist, and M. B. Cruzan. 2019. Functional trait divergence, not plasticity,
determines the success of a newly invasive plant. Ann. Bot. 123:667-679.
(http://doi.org/https://doi.org/10.1093/aob/mcy200).
Marsico, T. D., L. E. Wallace, G. N. Ervin, C. P. Brooks, J. E. McClure, and M. E. Welch. 2011. Geographic
patterns of genetic diversity from the native range of Cactoblastis cactorum (Berg) support the
documented history of invasion and multiple introductions for invasive populations. Biol.
Invasions 13:857-868. (http://doi.org/10.1007/s10530-010-9874-9).
Matute, D. R. 2013. The role of founder effects on the evolution of reproductive isolation. J. Evol. Biol.
26:2299-2311. (http://doi.org/https://doi.org/10.1111/jeb.12246).
Mayr, E. 1954. Change of genetic environment and evolution *in* J. Huxley, ed. Evolution as a Process.
Allen and Unwin, London.
Otto, S. P. and M. C. Whitlock. 1997. The probability of fixation in populations of changing size. Genetics
146:723-733. (http://doi.org/10.1093/genetics/146.2.723).
Otto, S. P. and M. C. Whitlock. 2013. Fixation Probabilities and Times. eLS.
Peng, B. and M. Kimmel. 2005. simuPOP: a forward-time population genetics simulation environment.
Bioinformatics 21:3686-3687. (http://doi.org/10.1093/bioinformatics/bti584).
Rego, A., F. J. Messina, and Z. Gompert. 2019. Dynamics of genomic change during evolutionary rescue
in the seed beetle Callosobruchus maculatus. Mol. Ecol. 28:2136-2154.
(http://doi.org/10.1111/mec.15085).
Rosenblum, E. B., M. J. Hickerson, and C. Moritz. 2007. A multilocus perspective on colonization
accompanied by selection and gene flow. Evolution 61:2971-2985.
(http://doi.org/10.1111/j.1558-5646.2007.00251.x).
Ruzicka, F., T. Connallon, and M. Reuter. 2021. Sex differences in deleterious mutational effects in
Drosophila melanogaster: combining quantitative and population genetic insights. Genetics 219.
(http://doi.org/10.1093/genetics/iyab143).
Simpson, G. G. 1944. Tempo and Mode in Evolution. Columbia University Press, New York.

Tepolt, C. K., J. A. Darling, M. J. Bagley, J. B. Geller, M. J. Blum, and E. D. Grosholz. 2009. European green
crabs (Carcinus maenas) in the northeastern Pacific: genetic evidence for high population
connectivity and current-mediated expansion from a single introduced source population.
Divers. Distrib. 15:997-1009. (http://doi.org/10.1111/j.1472-4642.2009.00605.x).
van der Zee, M. J., J. R. Whiting, J. R. Paris, R. D. Bassar, J. Travis, D. Weigel, D. N. Reznick, and B. A.
Fraser. 2022. Rapid genomic convergent evolution in experimental populations of Trinidadian
guppies (Poecilia reticulata). Evolution Letters 6:149-161. (http://doi.org/10.1002/evl3.272).
Wade, M. J. and C. J. Goodnight. 1998. The theories of Fisher and Wright in the context of
metapopulations: When nature does many small experiments. Evolution 52:1537-1553.
Willi, Y., P. Griffin, and J. Van Buskirk. 2013. Drift load in populations of small size and low density.
Heredity 110:296-302. (http://doi.org/10.1038/hdy.2012.86).
Willis, J. H. 1999. Inbreeding load, average dominance and the mutation rate for mildly deleterious
alleles in *Mimulus guttatus*. Genetics 153:1885-1898.
Willoughby, J. R., A. M. Harder, J. A. Tennessen, K. T. Scribner, and M. R. Christie. 2018. Rapid genetic
adaptation to a novel environment despite a genome-wide reduction in genetic diversity. Mol.
Ecol. 27:4041-4051. (http://doi.org/https://doi.org/10.1111/mec.14726).
Wright, S. 1931. Evolution in Mendelian populations. Genetics 16:97 - 159.
Wright, S. 1932. The roles of mutation, inbreeding, crossbreeding and selection in evolution.
Proceedings of the Sixth International Congress of Genetics 1:356-366.
Yin, X. S., A. S. Martinez, M. S. Sepulveda, and M. R. Christie. 2021. Rapid genetic adaptation to recently
colonized environments is driven by genes underlying life history traits. BMC Genomics 22.
(http://doi.org/10.1186/s12864-021-07553-x).

532
533
534

Fig. 1. The proportion of deleterious mutations fixing (A) and the levels of drift load (B) across population sizes of $N = 2$ to 4000 after 1000 generations ($\mu = 1x10^{-3}$). Selection on mutations ranged from 0.0 to 0.25 and were drawn randomly from a Poisson distribution, and dominance ranged from 0.0 to 0.2 and were drawn randomly from a uniform distribution. The blue lines represent the loess-smoothed means from 1000 replications for each N, and gray shading indicate 95% confidence intervals.

Fig. 2. Levels of inbreeding load due to loci segregating for deleterious mutations across population sizes of $N = 2$ to 4000 after 1000 generations ($\mu = 1x10^{-3}$). Selection on mutations ranged from 0.0 to 0.25 and were drawn randomly from a Poisson distribution, and dominance ranged from 0.0 to 0.2 and were drawn randomly from a uniform distribution. The blue lines represent the loess-smoothed means from 1000 replications for each N, and gray shading indicate 95% confidence intervals.

Fig. 3. The diffusion approximation for the probability of fixation (Φ_q) in response to initial allele frequency with $Ns = 2$ (A), and the proportion of fixed alleles across 100 loci after 1000 generations for population sizes of $N = 100$ and 1000 with $Ns = 2$ (B; the relationship for Φ_q is obscured). Lines represent the loess-smoothed relationship of means for 20 replications per population size.

Fig. 4. Mean fitness effects of beneficial mutations across population sizes of *N = 2* to 4000 after 1000 generations with initial *q = 1/2N* (*μ = 0*). Selection on mutations ranged from 0.0 to 0.25 and were drawn randomly from a Poisson distribution, and dominance ranged from 0.0 to 0.2 and were drawn randomly from a uniform distribution. The blue lines represent the loess-smoothed means from 1000 replications for each *N*, and gray shading indicate 95% confidence intervals.

675   Fig. 5. Deviations between the diffusion approximation for the probability of fixation (Φ_q), and observed
676   frequencies of fixation across initial allele frequencies (*q = 0.01* to *0.5*) and dominance (*h = 0.0* to *1.0*) for
677   a population size of 100 and *s = 0.02* (*Ns = 2*) after 1000 generations with initial *q = 1/2N* (*μ = 0*).

694
695   Fig. 6. Effects of population size on rates of fixation of beneficial alleles from simulations (*h = 0.5*)
696   compared to the diffusion approximation averaged across 10 replicates per population size ($\bar{t} - \bar{t}_{gens}$)
697   for *s = 0.02* (red), *0.04* (black), *0.10* (blue), and *0.20* (green). The lines represent the loess-smoothed
698   means across 100 replications for each *N*, and gray shading indicates 95% confidence intervals.
699
700

740 Fig. 7. Accumulation of beneficial mutations (A; fixation flux), mutations lost (B), and the proportion
741 continuing to segregate (C) after 1000 (blue), 2000, (purple), and 4000 (magenta) generations with *Ns =*
742 *2*, and over 1000 generations with *s = 0* (red). The lines represent the loess-smoothed means across 100
743 replications for each *N*, and gray shading indicates 95% confidence intervals.
744

745   Appendix 2. Supplemental Figures.

Fig. S1. Accumulation of beneficial mutations (A; fixation flux), mutations lost (B), and the proportion continuing to segregate (C) after 1000 generations with $Ns = 4$ (blue), $Ns = 0.02$ (purple), and $Ns = 0$ (red). The lines represent the loess-smoothed means across 100 replications for each N, and gray shading indicates 95% confidence intervals.

789
790
791

807 Fig. S2. Mean *Δq* for beneficial mutations going to fixation from an initial frequency of *1/2N* with *s = 2/N*
808 (*Ns = 2*) across population sizes from *N = 4* to *3000*. The line represents the loess-smoothed means
809 across 100 replications for each *N*, and gray shading indicates the 95% confidence interval.
810
811

827 Fig. S3. Standard deviation of *Δq* for beneficial mutations going to fixation from an initial frequency of
828 *1/2N* with *s = 2/N* (*Ns = 2*) across population sizes from *N = 4* to *3000*. The line represents the loess-
829 smoothed means across 100 replications for each *N*, and gray shading indicates the 95% confidence
830 interval.
831
832

848
849
850   Fig. S4. Mean *Δq* for beneficial mutations going to fixation from an initial frequency of *1/2N* across
851   selection coefficients from *s = 0.01* to 0.20 for *N = 100* (green line) and *N = 1000* (blue line). The line
852   represents the loess-smoothed means across 100 replications for each *N*, and gray shading indicates the
853   95% confidence interval.
854
855

871   Fig. S5. Standard deviation of *Δq* for beneficial mutations going to fixation from an initial frequency of
872   *1/2N* across selection coefficients from *s = 0.01* to 0.20 for *N = 100* (green line) and *N = 1000* (blue line).
873   The line represents the loess-smoothed means across 100 replications for each *N*, and gray shading
874   indicates the 95% confidence interval.
875
876

892   Fig. S6. Accumulation of beneficial mutations (fixation flux; blue) and mutations lost (red) across
893   population sizes after 1000 generations with *s = 0.02*. The lines represent the loess-smoothed means
894   across 100 replications for each *N*, and gray shading indicates 95% confidence intervals.
895