Testing Wright's Intermediate Population Size Hypothesis – When Genetic Drift is a Good Thing

Mitchell B. Cruzan

Portland State University, Department of Biology

Contact: cruzan@pdx.edu

Abstract – In his 1931 monograph, Sewall Wright predicted genetic drift would overwhelm selection in
very small populations, and selection would dominate in large ones, but also concluded drift could
facilitate selection in populations of intermediate size. The idea that drift and selection would act
together in smaller populations has not been evaluated using analytical or numerical approaches even
as empirical evidence of rapid evolution associated with population bottlenecks has continued to
accumulate. I used forward-time simulations with random mating and discrete generations to test the
hypothesis that drift can facilitate selection in small populations. I find evidence of drift facilitation of
selection as increases in levels of Δq in small populations ($N<100$) when selection is weak ($s<0.2$) and
when allele frequencies are low ($q<0.5$). Fixation of beneficial mutations is accelerated by drift
facilitation in small populations for recessive and codominant alleles, and less so for dominant alleles.
Drift facilitation accelerated fixation of beneficial mutations in small populations compared to
predictions from diffusion equations, while fixation time was longer than predicted in large populations.
Drift facilitation increases the probability of fixation of new mutations in small populations.
Accumulation of beneficial mutations (fixation flux) over several thousand generations was high in small
populations and declined rapidly for large populations, which accumulated large amounts of standing
genetic variation. Even though selection is more efficient in large populations, the increased time for
allele replacement and lack of drift facilitation can result in substantially slower rates of adaptive
evolution. Small populations were more susceptible to the accumulation of drift load, while larger
populations maintained higher levels of segregation load. These results indicate that drift facilitation in
small populations promotes purging of genetic load and accelerated fixation of beneficial mutations, and
may account for the large number of observations of rapid adaptation during population bottlenecks.

Impact Summary – After the recognition of Gregor Mendel's contributions to our understanding of the
inheritance of genetically-determined traits around 1900, there was confusion as to whether the type of
variation Mendel studied could account for evolution by natural selection, as described by Charles
Darwin. This controversy was resolved when three theoreticians (Ronald Fisher, Sewell Wright, and
J.B.S. Haldane) published books that integrated Mendelian genetics with evolution. Their contributions
(referred to as the Modern Synthesis), focused on evolutionary processes occurring within and among
populations of a species, and established a mathematical foundation for our understanding of
evolutionary biology. The mathematical models developed by the three architects of the modern
synthesis, and those who followed, predicted that the effects of natural selection would be
overwhelmed by random genetic changes (referred to as Genetic Drift) in small populations, and that
genetic drift would be minimal, while selection would be most effective in large populations. Even
though one of Wright's major conclusions was that genetic drift and selection would work together
(Drift Facilitation) to promote adaptive evolution in intermediate-sized populations, this idea has been
almost completely ignored since it was first introduced in 1931. In this study, I use simulations of
evolution in natural populations to evaluate the potential for drift facilitation to promote evolution in

small populations. My work largely confirms Wright's predictions; the removal of deleterious mutations
and promotion of adaptive evolution are enhanced in population sizes ranging from about 10 to 100.
These results indicate that our paradigm for our understanding of evolution within populations needs
refinement to emphasize the importance of drift facilitation in small populations, and to recognize that
periods of reduced population size are opportunities for enhanced levels of adaptive evolution.
50
51 Keywords: Adaptative evolution, conservation genetics, drift facilitation, drift load, inbreeding load,
52 population bottleneck, purging, segregation load, simulations.
53
54 "In a population of intermediate size ... there is continual random shifting of gene frequencies ... which
55 leads to a relatively rapid, continuing, irreversible, and largely fortuitous, but not degenerative series of
56 changes, even under static conditions." Wright 1931, Page 157.
57
Introduction
The mathematical models for population genetics developed during the Modern Synthesis (Fisher 1930;
Wright 1931; Haldane 1932) and beyond (Kimura 1983; Crow 1987; Ewens 2004; Charlesworth and
Charlesworth 2017; Charlesworth 2020) have provided a foundation for a general understanding of
evolutionary processes accounting for divergence among populations, adaptation, and diversification
among lineages. A fundamental aspect of this framework is the dichotomy between the roles of genetic
drift and selection; specifically, that drift governs the accumulation of neutral and degenerative
variation while only selection can promote the purging of deleterious mutations and the fixation of
beneficial mutations (Crow and Kimura 1970; Kimura 1983). The general prediction is that genetic drift
will dominate the dynamics of allele frequency change in small populations when selection is weak (i.e.
when $s < 1/4N_e$; Charlesworth 2009), resulting in the accumulation of genetic load and limited potential
for adaptive evolution. In contrast, selection is expected to overwhelm the effects of drift in large
populations leading to fixation of beneficial mutations and the purging of genetic load (Charlesworth
2002; Gossmann et al. 2012; Lanfear et al. 2014; Charlesworth and Charlesworth 2017). However, in his
1931 monograph, Wright emphasized the idea that the rate of fixation of beneficial mutations would be
much slower in large populations and that drift could overwhelm selection in very small populations, but
there was potential for genetic drift and selection to act together to promote rapid adaptation in
populations of "intermediate" size. This idea was intrinsic to his Shifting Balance Theory of adaptive
evolution where he predicted that adaptive variants were more likely to become fixed in small
populations, and subsequently spread by gene flow (Wright 1932). These predictions were premised on
the idea that there was a range of population sizes where both drift and selection are effective enough
to allow "fortuitous" shifts in allele frequencies to facilitate selection by generating larger numbers of
high-fitness genotypes resulting in accelerated accumulation of adaptive genetic variation. This idea
(hereafter referred to as "drift facilitation") was seized upon by George Gaylord Simpson (1944), who
proposed small population size was a likely explanation for the lack of fossilization during rapid
transitions that resulted in the origin of novel forms, including new genera and families. This idea of
"quantum evolution" (i.e. periods of rapid evolution when selection is facilitated by genetic drift;
Simpson 1944) inspired a number of evolutionary biologists, who subsequently identified a wide range
of examples of adaptation associated with reduced population size (Mayr 1954; Lewis and Raven 1958).
Verne Grant (1963, 1981) expanded on the idea of quantum evolution to include rapid speciation events
in small, isolated populations at the periphery of a widespread species' range (budding or peripatric

speciation; Mayr 1954). While interest in the idea of quantum evolution has waned, evidence of rapid
evolution associated with population bottlenecks continues to accumulate (e.g., Carvalho et al. 1996;
Rosenblum et al. 2007; Tepolt et al. 2009; Marsico et al. 2011; Cruzan 2019; Rego et al. 2019; Chaturvedi
et al. 2021; Cruzan et al. 2021; Mahrt et al. 2021; Yin et al. 2021; van der Zee et al. 2022; Appendix 3).
Surprisingly, there have been no attempts to assess the potential for drift facilitation of selection in
populations of small or intermediate size using analytical or numerical approaches.
95
In this study, I use forward-time, individual-based simulations of populations to evaluate the potential
for drift facilitation to promote evolution in intermediate-sized (small) populations. I evaluate the effects
of selection (s), dominance (h), and population size (N) on changes in allele frequencies between
generations ($\Delta q = q_{n+1} - q_n$) and its consequences for the loss or accumulation of deleterious mutations,
and the time to fixation and probability of fixation for beneficial mutations. I specifically focus on short
time-frames to account for the numerous observations of rapid evolution within a few hundred
generations during population bottlenecks. In all cases, loci are bi-allelic and segregate independently,
generations are discrete, populations are dioecious with equal sex ratios, and mating is random (i.e. $N =$
N_e). I first characterize the effects of population size, selection, and dominance on Δq for beneficial
alleles going to fixation, and then examine the consequences of selection and population size for the
accumulation of genetic load and beneficial mutations. For genetic load, I examine the proportion of loci
that became fixed for deleterious mutations (drift load), and the proportion of deleterious mutations
that remain segregating after a number of generations (segregation load). In separate simulations, I
examine the probability of fixation for beneficial mutations, the time to fixation, and fixation flux (the
rate at which beneficial alleles accumulate; Otto and Whitlock 1997). My results demonstrate the effects
of drift facilitation as elevated levels of Δq in small populations ($N \leq 100$) when allele frequencies are
low and selection is weak. I find that drift facilitation of selection promotes rapid purging of genetic load
and the accumulation of beneficial mutations in small populations while large populations accumulate
greater levels inbreeding load and standing genetic variation for beneficial mutations.
115
Methods
Forward-time, individual-based simulations were conducted using the SimuPOP module (Peng and
Kimmel 2005) in Python and the COEUS High-Performance Computing Cluster at Portland State
University. A set of N individuals, each homozygous for a single locus, were generated using the
sim.Population simulator and mutations introduced ($a \rightarrow a'$) that were affected by varying levels of
selection (s) and dominance (h). Transmission of alleles between generations was governed by the sim.
MendelianGenoTransmitter. Selection was imposed at the time of mating by defining the fitness of each
diploid genotype using MapSelector in SimuPOP (all Python scripts are available in Appendix 1).
Depending on the goals of the simulation, mutations were either introduced once in the first generation,
or at a per-generation rate determined by μ. Each population size of N individuals was replicated k
times. For each replication, the proportion of individuals with fixed mutations (q_{fixed}), and the average
allele frequency across individuals ($\bar{q}$) were extracted after g generations. Two sets of simulations were
run; the first set examined levels of drift and segregation load, and the second set focused on beneficial
mutations by evaluating the proportion of loci fixed for mutations, the rate of fixation of individual
mutations, and the fixation flux over g generations.
131

Testing for Drift Facilitation of Selection in Small Populations – I tested for the effects of drift facilitation
of selection by comparing Δq ($\Delta q = q_{n+1} - q_n$) in small ($N = 50$) and large ($N = 1000$) populations across
generations as beneficial mutations increased from their initial frequency of $q = 1/2N$ to fixation.
Mutations were repeatedly generated and their allele frequencies tracked until they were fixed or lost
(Python script available in Appendix 1.A). For mutations that fixed, Δq for each of the n generations was
calculated along with its mean and standard deviation across generations. Mean Δq for mutations that
fixed was compared for a range of selection coefficients ($s = 0.01, 0.04, 0.12,$ and 0.20) across
population sizes ranging from $N = 4$ to $N = 3000$ for recessive, additive, and dominant mutations ($h = 0.0,$
0.5, and *1.0*). I evaluated the effects of the effects of the strength of selection ($s = 0.01$ to 0.20) in a small
($N = 100$) and large ($N = 1000$) population on the mean and standard deviation of Δq ($k = 10,000$
replications) for beneficial mutations going to fixation (Appendix 1B).

Consequences of Drift Facilitation for Genetic Load – The frequency of fixed deleterious mutations (drift
load) across loci and of the proportion of loci segregating for deleterious mutations after g generations
(segregation load) was assessed for population sizes ranging from $N = 2$ to 5000 (Python script available
in Appendix 1C). For each of $k = 1000$ replications for each population size, s was randomly generated
from a Poisson distribution (range: $0.0 - 0.25$; $\bar{s} = 0.07$), and h was generated from a uniform
distribution between 0.0 and 0.2 (assuming that deleterious mutations are generally recessive with $\bar{h} =$
0.10; Willis 1999; García-Dorado and Caballero 2000; Agrawal and Whitlock 2011; Ruzicka et al. 2021).
Fitness of the *aa, aa',* and *a'a'* genotypes were set to 1.0, *1-hs,* and *1-s,* respectively. The initial allele
frequency of *a'* was set at $q = 0$ and mutations were allowed to accumulate for $g = 1000$ generations
with $\mu = 1x10^{-3}$ This assumes that each locus consists of thousands of base pairs (i.e. if the per
bp/generation mutation rate is on the order of $1x10^{-9}$), but the same qualitative results are obtained for
lower mutation rates over longer timeframes.

The proportion of loci fixed for deleterious mutations (q_{fixed}) and selection (s) for each mutation were
used to estimate the fitness effects of fixed deleterious mutations (drift load) for each replication of
each population size as the reduction in fitness due to the selection coefficient weighted by the
proportion of loci fixed for the mutation;

$$w_{fixed} = 1 - sq_{fixed}$$

Segregation load was assessed as the inbreeding effects on fitness for segregating mutations (inbreeding
depression; Charlesworth and Willis 2009; Willi et al. 2013) for 1000 replicates for each population size
after 1000 generations. Inbreeding depression was estimated based on the mean fitness of hypothetical
progeny that would be the result of self-fertilization compared to progeny from random outcrossing.
Fitness after selfing was estimated as,

$$\bar{w}_s = 2\bar{p}\bar{q}(1 - (0.25 + 0.5hs + 0.25s)^F);$$

where $\bar{p}$ and $\bar{q}$ are the mean frequencies of the normal (a) and mutant (a') alleles across segregating loci
(individuals), and F is the number of segregating sites at the end of 1000 generations. I calculated the
expected fitness of progeny from random outcrossing within each population as,

$$\bar{w}_o = 1 - (\bar{p}^2 + hs2\bar{p}\bar{q} + s\bar{q}^2)^F$$

177   Segregation load was then estimated for each N as inbreeding depression for selfed progeny compared
178   to outcrossed progeny (Charlesworth and Willis 2009) as,

$$\delta_{s/o} = 1 - \frac{\bar{w}_s}{\bar{w}_o}$$

These estimates of drift load and segregation load assume independent fitness effects across unlinked
loci.

Consequences of Drift Facilitation for Adaptation – I examined the processes of adaptation in
populations using simulations to evaluate the probability of fixation of beneficial mutations, the rate of
fixation of individual alleles, and the fixation flux for a range of population sizes (N), selection (s), and
dominance (h; Appendix 1D). For beneficial mutations ($a \rightarrow a'$), the fitness of the aa, aa', and $a'a'$
genotypes were set to *1-s*, *1-hs*, and *1.0*, respectively.

The first set of simulations compared the fixation of beneficial alleles across a range of initial allele
frequencies from *q = 0.01* to *0.50* when *Ns = 2* for population sizes of *N = 100* (*s = 0.02*) and *N = 1000* (*s =*
0.002). Initial frequency of the *a'* allele was set at *q = 1/2N* (*μ = 0*) and the proportion of loci fixing for
the mutant allele was compared to the diffusion prediction for the probability of fixation (Φ_q) derived by
Kimura (1957; 1962). Alleles were codominant in simulations (*h = 0.5*) to match the assumptions of the
diffusion approximation;

197
$$\Phi_q = \frac{1 - \exp(-2Nsq)}{1 - \exp(-2Ns)}$$

A second set of simulations examined the number of generations required for fixation over population
sizes ranging from *N = 4* to *3000* with *s* randomly drawn from a Poisson distribution (*s = 0.0* to *0.25*) and
h drawn from a uniform distribution (*h = 0.0* to *1.0*). Initial frequency of the *a'* allele was set at *q = 1/2N*
(*μ = 0*) and the proportion of loci fixing for the mutant allele was recorded after 10,000 generations. I
compared the number of generations required for fixation for beneficial alleles ($\bar{t}_{gens}$ across *k = 10*
replicates per population size) to the predicted time to fixation ($\bar{t}$) from the diffusion approximation
(Ewens 1979; Otto and Whitlock 2013);

$$\bar{t} \approx \int_{1/2N}^{0.999} \frac{(e^{4Nsx} - 1)(e^{4Ns(1-x)} - 1)}{sx(1-x)(e^{4Ns} - 1)} dx$$

This integral is undefined when the upper limit is 1.0, and the approach to fixation is asymptotic, so an
arbitrary upper limit of 0.999 was chosen for comparison to fixation frequencies from the simulations.

While this approach describes fixation times for individual alleles, it does not account for the
contribution of the number of mutations entering populations each generation based on population size
for the accumulation of beneficial mutations over time (fixation flux; Otto and Whitlock 1997). We

expect flux to be affected by the probability of fixation for individual mutations and the number of
mutations entering the population each generation. I evaluated the probability of fixation across a range
of population sizes (N = *4* to *3000*) by introducing random mutations (s = *0.0* to *0.2* following a Poisson
distribution, and h = *0.0* to *1.0* following a flat distribution) to 1000 loci in the first generation and
tracking the proportion that fix after 10,000 generations (100 replications per population size).

We expect that more mutations would occur in larger populations and consequently flux should
increase with population size. On the other hand, fixation will take longer in large populations because
of the time required for replacement of the original allele in all individuals. Since the goal of this study is
to understand conditions leading to rapid adaptation to a novel environment, simulations were limited
to a few thousand generations without a prior burn-in period to allow for the accumulation of standing
genetic variation. I examined fixation flux for population sizes ranging from N = *4* to *3000* with μ = *1x10*$^{-}$
^{4}, Ns = *2* (s = *2/N*), and h = *0.5* with 1000 loci and 10 replicates per population size for time periods of g =
1000, 2000, and *4000* generations (Appendix 1E). Fixation flux under selection was compared to fixation
with only genetic drift (s = *0*) over 1000 generations across the same range of population sizes. In
addition, I estimated the mean and standard deviation of Δq across generations for beneficial mutations
that ultimately fixed in populations across a range of populations sizes (N = *4* to *3000*) and selection
coefficients (s = *0.01* to *0.20* for N = *100* and *1000*).

Results
Drift Facilitation of Selection in Small Populations – Drift facilitation was evident as elevated levels of Δq
when population size was small (N = *1000*) compared to large populations (N = *1000*; Fig. 1). The effects of
drift facilitation were strongest when selection was weak (s < *0.2*), mutations were recessive (h = *0.0*),
and allele frequencies were low, but there were strong effects of drift facilitation for codominant
mutations (Fig. 1), and weaker effects when mutations were completely dominant (Fig. S1). There was a
general trend towards the strongest effects of drift facilitation in small populations with weaker
selection and lower dominance. Mean Δq increased with the strength of selection (Fig. S2) and remained
consistently higher in small (N = *100*) compared to large (N = *1000*) populations. The standard deviation
of Δq was consistently higher for small populations and remained fairly constant across a range from s =
0.01 to *0.20* (Fig. S3).

Consequences for Genetic Load – Fixation frequencies of deleterious mutations were highest for small
populations (< 100), and was effectively zero for population sizes larger than N = 3000 (Fig. 2A).
Mutations fixing in populations when N > *100* had negligible fitness effects so that drift load was near
zero when population size was greater than N = *100* (Fig. 2B). Larger proportions of loci continued to
segregate for deleterious mutations after 1000 generations in populations of less than N = 10 and more
than N = 100, while segregation load was lower when N > *10* and N < *100*; Fig. 3).

Consequences for Adaptive Evolution – There was a close fit between the diffusion approximation for the
probability of fixation across initial allele frequencies (Φ_q) and the proportion of loci fixed for beneficial
mutations with population sizes of 100 and 1000 with Ns = 2 and codominance (h = *0.5*; Fig. S4).

The time to fixation of individual alleles ($\bar{t}_{gens}$) was similar to the diffusion approximation ($\bar{t}$) when
selection was stronger (s = *0.1* and *0.2*) and population size was less than N = *100*, but was slower than

predicted when *N > 100*. Time to fixation was substantially faster when *s = 0.04* and *0.02* when
population sizes were greater than *N = 10* and less than *N = 100* (Fig. 4). The probability of fixation for
beneficial mutations was highest for small populations (*N < 100*), and remained lower for large
populations (Fig. 5). Fixation of random beneficial mutations was positively associated with the strength
of selection (*t = 18.79*; *P < 0.0001*) and negatively associated with dominance (*t = -8.87*; *P < 0.0001*) and
population size (*t = -2.71*; *P = 0.0072*). Fixation flux for beneficial mutations with *Ns = 2* was greater after
4000 generations compared to 2000, and 1000 generations (Fig. 6A). Fixation flux and the loss of
beneficial mutations was high when *N < 100*, but generally declined with increasing population size (*h =*
0.5; Figs. 6A, B) due to the limited time-frame (i.e. < 4000 generations) and the number of generations
required for allele replacement in large populations. The number of mutations that remained
segregating in large populations was not strongly affected by the length of the time for the
accumulation of mutations, and was consistently low when *N < 100*, and increased rapidly when *N > 100*
(Fig. 6C).
271
Discussion
Forward-time simulations of natural populations indicate that drift facilitation of selection elevates *Δq* in
small populations (*N $\leq$ 100*) resulting in lower segregation load, accelerated times to fixation, and
higher probability of fixation for beneficial mutations. Drift load was relatively high for small
populations, but was negligible for populations larger than *N = 100*, while segregation load was lowest
for population sizes between 10 and 100. Results of simulations for beneficial mutations were consistent
with diffusion approximations of the probability of fixation when *h = 0.5* (Kimura 1957; Kimura 1962).
Observed times to fixation matched the diffusion approximation ($\bar{t}$) fairly well when selection was
strong (*s $\geq$ 0.10* and *h = 0.5*), but fixation times were substantially faster when *s < 0.10* for population
sizes between *N = 10* and *100*, and slower for population sizes of *N > 100*. Drift facilitation in smaller
populations led to higher probabilities of fixation and high levels of fixation flux of beneficial mutations
across a broad range of small population sizes while flux rapidly declined for populations larger then *N =*
100 after 1000 generations, and *N = 300* after 4000 generations, which was apparently due to the
extended time to fixation in larger populations. While we expect that more beneficial mutations will
occur in large populations, the lack of genetic drift reduces the chances of fixation and increases the
time to fixation compared to small populations. These results suggest the that drift facilitation provides
some explanation for observations of rapid adaptation during periods of reduced population size by
increasing purging of deleterious alleles and promoting the acquisition of adaptive genetic variation.
290
The results presented here generally support Wright's (1931) predictions about the potential for
interactions between drift and selection to promote rapid fixation of beneficial alleles, but the
population sizes where drift facilitation of selection is effective are smaller than he predicted. Wright
specifically defined intermediate populations as ranging between *Ns = 2.5* and *20* (page 148). Simpson
(1944) also assumed that the dynamics described by Wright would apply to very large population sizes
(i.e. up to *N = 25,*000) compared to the range of population sizes where drift facilitation was found to be
effective in the simulations presented here. In contrast to the predictions of Wright and Simpson, higher
rates and probabilities of fixation of beneficial mutations were observed in simulations for *N < 100*. The
results presented here indicate that we cannot assume there is a linear tradeoff between the effects of
population size and the strength of selection when population sizes small (i.e. for *N < 100*). This is an
important result because *Ns* has been used to represent the compensatory effects between genetic drift

and selection in analytical models including diffusion equation analyses of population genetic processes
(Kimura 1957; Kimura 1962; Ewens 2004). Given the results presented above, more careful evaluation of
the separate effects of selection and population size may be warranted when small populations are
being considered.
306
While population genetic models have mostly ignored the potential for selection in small populations,
empirical evidence for the effectiveness of selection during population bottlenecks continues to
accumulate. For example, a search of Web of Science using the keywords "rapid and (evolution or
adaptation) and bottleneck" yielded 153 publications since 1992 on a wide variety of organisms after
removing reviews, editorials, non-biological studies, and studies focused on cancer (details in Appendix
3). A sample of the reported cases of rapid adaptation to novel conditions during establishment and
bottlenecks includes artificial introductions of Trinidadian guppies (Carvalho et al. 1996; van der Zee et
al. 2022), Eastern fence lizards on novel substrates in New Mexico (Rosenblum et al. 2007), European
green crabs on the west coast of North America (Tepolt et al. 2009), the cactus-feeding moth,
Cactoblastis cactorum, in Florida (Marsico et al. 2011), steelhead trout in Lake Michigan (Willoughby et
al. 2018), the invasive grass, *Brachypodium sylvaticum*, in Oregon (Cruzan 2019; Marchini et al. 2019),
during evolutionary rescue in the seed beetle *Callosobruchus maculatus* (Rego et al. 2019), in founding
populations of *Daphnia* (Chaturvedi et al. 2021), and after migration into Lake Champlain by the Atlantic
sea lamprey (Yin et al. 2021). In one case, there was rapid evolution of antibiotic resistance in
Pseudomonas aeruginosa even when population bottlenecks were severe and selection was weak
(Mahrt et al. 2021). Rapid acquisition of adaptive genetic variants also seems to have occurred in a case
of recent budding speciation (peripatric speciation; Mayr 1954) in a range-restricted species of
buttercup (Cruzan et al. 2021). In all of these cases, adaptation to novel conditions appears to have
occurred quickly during periods of reduced population size and suggests drift facilitation of selection
may be a widespread phenomenon.
327
One of the more surprising results of simulations was the observation of higher rates of fixation flux of
beneficial mutations when $N < 300$, suggesting drift facilitation can enhance rates of adaptation in small
populations. This result is in consistent with diffusion analyses, which predicts flux estimated as $4NsU$
(where U is the genome-wide mutation rate) will remain constant across population sizes if Ns is held
constant (Otto and Whitlock 1997; Ewens 2004), but the decline in flux when $N > 300$ is due to the slow
accumulation of standing genetic variation. The simulations conducted here assumed transitions to
novel conditions or colonization events so adaptation from novel mutations would only be for those
occurring after the transition. The rate of flux may have been equally high for large populations if
substantial "burn-in" periods (> 10,000 generations) had been implemented to allow the accumulation
of standing genetic variation. However, the rate of adaptation from standing genetic variation in large
populations will depend on initial allele frequency, and even for alleles that are at mid-frequencies (e.g.,
Lai et al. 2019), the time to fixation would be substantially longer than for small populations.
340
The observations of elevated Δq, reduced time to fixation, and increased probability of fixation of
beneficial mutations in small populations are consistent with some previous analyses that predicted
genetic drift and selection could interact to increase rates of divergence among populations (Cohan
1984; Lynch 1986), and with selection experiments in maize and simulations where enhanced rates of
fixation of small effect mutations were observed when subjected to high genetic drift (Desbiez-Piat et al.

2021). It's notable that predictions from diffusion analyses are often evaluated only for a range of
population sizes that are much larger than what was considered here (e.g., for N > 1000; Charlesworth
2020), but many species have historically experienced severe population bottlenecks. While diffusion
analyses have led to numerous insights for evolution in large populations, alternative approaches may
be required when population sizes less than $N = 300$ are considered.
351
Previous studies that have found evidence of drift facilitation did not identify its causes (e.g., Cohan
1984; Lynch 1986), but the evaluation of Δq across allele frequencies and population sizes sheds some
light on the mechanisms responsible for enhancement of reduced times to fixation in small populations.
The mean and standard deviation of Δq increased in small populations when selection was weak ($s < 0.2$)
and mostly for allele frequencies of $q < 0.5$. For mutations at low frequencies, it appears that
"fortuitous" fluctuations due to drift generate larger numbers of high-fitness genotypes (i.e.
homozygotes when $h = 0$, and heterozygotes as well when $h = 0.5$). Consequently, the effects of drift
facilitation are greatest for recessive and codominant mutations as drift facilitation elevates Δq and
generates a "ratcheting effect" that accelerates changes in allele frequencies. This idea is supported by
the observation that the mean of Δq is positively associated with selection, and is consistently higher in
small compared to large populations. The effects of drift facilitation are apparently different when allele
frequencies are close to $q = 0$ (for deleterious mutations) or fixation (i.e. $q = 1$ for beneficial mutations).
Previous analyses indicate that diffusion equations do not perform well when allele frequencies are
close to fixation when alleles under selection are completely recessive or dominant (Charlesworth
2020). Simulations performed here indicate that higher levels of drift in smaller populations increase the
probability for random fluctuations to result in absorption, while minimal drift in larger populations
results in persistent segregation. Consequently, larger populations appear to accumulate higher levels of
segregation load and standing variation for beneficial mutations, while having lower drift load lower
probabilities of fixation for beneficial mutations compared to smaller populations (i.e. for $N < 100$).
371
The observation that large populations maintain higher levels of segregation load and standing variation
for beneficial mutations suggests that episodes of reduced population size may be disproportionally
important for rapid evolution in novel or changing environments. While it remains true that more
beneficial mutations are likely to occur in large populations, the number of generations required for
fixation may be too great to keep pace with current rates of climate change. On the other hand, large
populations possess higher levels of standing genetic variation so bottlenecks would have enhanced
potential for the fixation of beneficial mutations (e.g., Chaturvedi et al. 2021). Moreover, reductions in
population size due to bottlenecks and founder events are likely to be associated with increased
environmental stress, and consequently, we expect selection to often increase during population
bottlenecks (Fowler and Whitlock 2002; Matute 2013). Higher frequencies of segregating mutations
along with drift facilitation may be responsible for elevated levels of purging following population
bottlenecks (Bouzat 2010), and may help explain the substantial number of cases of rapid adaption
during episodes of reduced population size (e.g., Carvalho et al. 1996; Rosenblum et al. 2007; Tepolt et
al. 2009; Marsico et al. 2011; Cruzan 2019; Rego et al. 2019; Cruzan et al. 2021; Mahrt et al. 2021; Yin et
al. 2021; van der Zee et al. 2022; Appendix 3). On the other hand, we expect population bottlenecks and
founder events to result in elevated accumulation of drift load (Lynch et al. 1995; Willi et al. 2013), but
the simulations presented here indicate the potential for purging of segregation load and the fixation of
deleterious mutations depends on the severity of the reduction in population size. It is notable that the

standard guideline of a minimal effective population size of $N = 50$ (i.e. the 50/500 rule; Jamieson and
Allendorf 2012) for conservation efforts is too small to prevent the accumulation of drift load, but is in
the right range for drift facilitation of beneficial mutations. Further analyses may help resolve more
specific effects of changes in population size for adaptation to novel conditions and for the purging of
deleterious mutations.
395
Conclusions
The results of individual-based, forward-time simulations are consistent with a wide range of
observations from natural populations, but contrast with many of the predictions from models based on
diffusion equations (e.g., Ewens 2004) when population sizes are small. In general, it appears that
Wright's (1931) intuitive predictions about the potential for genetic drift to facilitate selection were
correct, but the population sizes where drift facilitation is most effective are smaller than he suggested.
Furthermore, while Wright saw the potential for weak deleterious mutations to fix in small populations,
he did not anticipate that drift facilitation could lead to more effective purging of segregation load. One
of the most striking results from these simulations is that the probability of fixation and flux of beneficial
mutations are substantially high in small populations ($N < 100$) and is similar to flux in larger populations
under the assumption that Ns remains constant. The assumption that selection is stronger in small
populations may generally hold true; especially under conditions where population declines are
precipitated by increased levels of environmental stress. Moreover, the time to fixation for beneficial
mutations was lower in small populations when selection was weak ($s < 0.2$), so the prediction of rapid
evolution due to drift facilitation is not dependent on the assumption of increased selection in small
populations.
412
The potential for rapid evolution in small populations may be further enhanced as large populations
experience reductions in population size. While it is generally assumed population bottlenecks will result
in decreased fitness due to the accumulation of genetic load, and that there will be limited potential for
adaptive evolution, this is not always true (Bouzat 2010). The simulation results presented here indicate
drift facilitation of selection in small populations leads to effective removal of segregating deleterious
mutations, and increases the rate and probability of fixation for beneficial mutations. While there has
been some speculation and evidence for enhanced evolution in small populations (Simpson 1944; Grant
1981; Cohan 1984; Lynch 1986), no previous study has provided a detailed characterization of the
potential for drift facilitation to promote rapid evolution.
422
Results from these simulations emphasize the importance of the adaptive evolution potential in small
populations. Furthermore, more attention should be paid to the consequences of deleterious mutations
as they depend heavily on population size; a clear distinction needs to be made between the fixation of
deleterious mutations contributing to drift load in small populations, and the persistence of deleterious
mutations in large populations resulting in inbreeding depression due to segregation load. The results of
these simulations highlight genetic drift as a generative, beneficial process that promotes evolution in
small populations. Alternatively, the lack of drift in large populations effectively stalls purging of
segregation load and the fixation of beneficial mutations. While further analyses of drift facilitation are
needed, the results presented here give more weight to Wright's (1931, 1932) suggestion that adaptive
evolution is more likely to occur in small populations (Wade and Goodnight 1998), then to Fisher's
(1930) argument that large populations are more important for fixation of beneficial mutations (Coyne

et al. 1997). Further work on evolution in small populations from modeling and empirical perspectives
will clarify the potential for drift facilitation to promote evolution, and ultimately determine whether
the current paradigm for our understanding of population genetic processes of evolution needs to be
refined.

Acknowledgements

I thank Michael Wade and Michael Whitlock for suggestions on the development of simulations and

relevant parameters to focus on. Karla de Lima Berg, Elizabeth Hendrickson, Jaime Schwoch, and Ariana

Walczyk provided suggestions to improve the manuscript and presentation of results. Thanks to Bo Peng

for assistance with coding in simuPop, and to Jim Stapleton and the PSU OIT for assistance with

configuration of Coeus HPC for simulations in simuPop. This research was supported by NSF-DEB award

2051235.

Author Contributions

MBC is solely responsible for the development of the SimuPOP simulations in Python, data analyses, and

writing of the manuscript.

Data Accessibility

All Python scripts are available in Appendix 1 and will be made available on Github.

Conflict of Interest

The author declares no conflicts of interest.

Literature Cited

Agrawal, A. F. and M. C. Whitlock. 2011. Inferences about the distribution of dominance drawn from
yeast gene knockout data. Genetics 187:553-U274.
(http://doi.org/10.1534/genetics.110.124560).
Bouzat, J. L. 2010. Conservation genetics of population bottlenecks: the role of chance, selection, and
history. Conserv. Genet. 11:463-478. (http://doi.org/10.1007/s10592-010-0049-0).
Carvalho, G. R., P. W. Shaw, L. Hauser, B. H. Seghers, and A. E. Magurran. 1996. Artificial introductions,
evolutionary change and population differentiation in Trinidadian guppies (Poecilia
reticulata:Poeciliidae). Biol. J. Linn. Soc. 57:219-234. ([http://doi.org/10.1111/j.1095-](http://doi.org/10.1111/j.1095-8312.1996.tb00310.x)
[8312.1996.tb00310.x](http://doi.org/10.1111/j.1095-8312.1996.tb00310.x)).
Charlesworth, B. 2002. Effective population size. Curr. Biol. 12:R716-R717.
(http://doi.org/10.1016/S0960-9822(02)01244-7).
Charlesworth, B. 2009. Effective population size and patterns of molecular evolution and variation.
Nature Reviews Genetics 10:195-205. (http://doi.org/10.1038/nrg2526).
Charlesworth, B. 2020. How long does it take to fix a favorable mutation, and why should we care? The
American Naturalist 195:753-771. (http://doi.org/10.1086/708187).
Charlesworth, B. and D. Charlesworth. 2017. Population genetics from 1966 to 2016. Heredity 118:2-9.
(http://doi.org/10.1038/hdy.2016.55).
Charlesworth, D. and J. H. Willis. 2009. The genetics of inbreeding depression. Nat. Rev. Genet. 10.
(http://doi.org/10.1038/nrg2664).
Chaturvedi, A., J. Zhou, J. A. M. Raeymaekers, T. Czypionka, L. Orsini, C. E. Jackson, K. I. Spanier, J. R.
Shaw, J. K. Colbourne, and L. De Meester. 2021. Extensive standing genetic variation from a
small number of founders enables rapid adaptation in Daphnia. Nature Communications
12:4306. (http://doi.org/10.1038/s41467-021-24581-z).
Cohan, F. M. 1984. Can uniform selection retard random genetic dif=vergence between isolated
conspecific populations? Evolution 38:495-504. (http://doi.org/10.2307/2408699).
Coyne, J. A., N. H. Barton, and M. Turelli. 1997. A critique of Sewall Wright's shifting balance theory of
evolution. Evolution 51:643-671.
Crow, J. F. 1987. Population genetics history: A personal view. Annu. Rev. Genet. 21:1-22.
(http://doi.org/10.1146/annurev.ge.21.120187.000245).
Crow, J. F. and M. Kimura. 1970. An Introduction to Population Genetics Theory. Harper & Row, New
York.
Cruzan, M. B. 2019. How to make a weed - the saga of the slender fasle brome invasion in the North
American west and lessons for the future. Bioscience 69:469-507.
(http://doi.org/https://doi.org/10.1093/biosci/biz051).
Cruzan, M. B., P. G. Thompson, N. A. Diaz, E. C. Hendrickson, K. R. Gerloff, K. A. Kline, H. M. Machiorlete,
and J. M. Persinger. 2021. Weak coupling among barrier loci and waves of neutral and adaptive
introgression across an expanding hybrid zone. Evolution 75:3098-3114.
(http://doi.org/https://doi.org/10.1111/evo.14381).
Desbiez-Piat, A., A. Le Rouzic, M. I. Tenaillon, and C. Dillmann. 2021. Interplay between extreme drift
and selection intensities favors the fixation of beneficial mutations in selfing maize populations.
Genetics 219. (http://doi.org/10.1093/genetics/iyab123).
Ewens, W. J. 1979. Mathematical Population Genetics. Springer-Verlag, Berlin.
Ewens, W. J. 2004. Mathmatical Population Genetics, Volume I: Theoretical Foundations. Springer-
Verlag, Berlin.
Fisher, R. A. 1930. The Genetical Theory of Natural Selection. Clarendon Press, Oxford.

Fowler, K. and M. C. Whitlock. 2002. Environmental stress, inbreeding, and the nature of phenotypic and
genetic variance in Drosophila melanogaster. Proceedings. Biological sciences / The Royal
Society 269:677-683. (http://doi.org/10.1098/rspb.2001.1931).
García-Dorado, A. and A. Caballero. 2000. On the average coefficient of dominance of deleterious
spontaneous mutations. Genetics 155:1991-2001.
(http://doi.org/10.1093/genetics/155.4.1991).
Gossmann, T. I., P. D. Keightley, and A. Eyre-Walker. 2012. The effect of variation in the effective
population size on the rate of adaptive molecular evolution in eukaryotes. Genome Biol Evol
4:658-667. (http://doi.org/10.1093/gbe/evs027).
Grant, V. 1963. The Origin of Adaptations. Columbia University Press, New York.
Grant, V. 1981. Plant Speciation. Columbia University Press, New York.
Haldane, J. B. S. 1932. The Causes of Evolution. Princeton University Press, Princeton.
Jamieson, I. G. and F. W. Allendorf. 2012. How does the 50/500 rule apply to MVPs? Trends Ecol. Evol.
27:578-584. (http://doi.org/https://doi.org/10.1016/j.tree.2012.07.001).
Kimura, M. 1957. Some problems of stochastic processes in genetics. The Annals of Mathematical
Statistics 28:882-901, 820.
Kimura, M. 1962. On the probability of fixation of mutant genes in a population. Genetics 47:713-719.
(http://doi.org/10.1093/genetics/47.6.713).
Kimura, M. 1983. The Neutral Theory of Molecular Evolution. Cambridge University Press, Cambridge.
Lai, Y.-T., C. K. L. Yeung, K. E. Omland, E.-L. Pang, Y. Hao, B.-Y. Liao, H.-F. Cao, B.-W. Zhang, C.-F. Yeh, C.-
M. Hung, H.-Y. Hung, M.-Y. Yang, W. Liang, Y.-C. Hsu, C.-T. Yao, L. Dong, K. Lin, and S.-H. Li. 2019.
Standing genetic variation as the predominant source for adaptation of a songbird. Proceedings
of the National Academy of Sciences 116:2152-2157.
(http://doi.org/doi:10.1073/pnas.1813597116).
Lanfear, R., H. Kokko, and A. Eyre-Walker. 2014. Population size and the rate of evolution. Trends Ecol.
Evol. 29:33-41. (http://doi.org/http://dx.doi.org/10.1016/j.tree.2013.09.009).
Lewis, H. and P. H. Raven. 1958. Rapid evolution in *Clarkia*. Evolution 12:319-336.
(http://doi.org/10.2307/2405854).
Lynch, M. 1986. Random drift, uniform selection, and the degree of population differentiation. Evolution
40:640-643. (http://doi.org/10.2307/2408585).
Lynch, M., J. Conery, and R. Burger. 1995. Mutational meltdowns in sexual populations. Evolution
49:1067-1080. (http://doi.org/10.2307/2410432).
Mahrt, N., A. Tietze, S. Kunzel, S. Franzenburg, C. Barbosa, G. Jansen, and H. Schulenburg. 2021.
Bottleneck size and selection level reproducibly impact evolution of antibiotic resistance. Nature
Ecology & Evolution 5. (http://doi.org/10.1038/s41559-021-01511-2).
Marchini, G. L., C. A. Maraist, and M. B. Cruzan. 2019. Functional trait divergence, not plasticity,
determines the success of a newly invasive plant. Ann. Bot. 123:667-679.
(http://doi.org/https://doi.org/10.1093/aob/mcy200).
Marsico, T. D., L. E. Wallace, G. N. Ervin, C. P. Brooks, J. E. McClure, and M. E. Welch. 2011. Geographic
patterns of genetic diversity from the native range of Cactoblastis cactorum (Berg) support the
documented history of invasion and multiple introductions for invasive populations. Biol.
Invasions 13:857-868. (http://doi.org/10.1007/s10530-010-9874-9).
Matute, D. R. 2013. The role of founder effects on the evolution of reproductive isolation. J. Evol. Biol.
26:2299-2311. (http://doi.org/https://doi.org/10.1111/jeb.12246).
Mayr, E. 1954. Change of genetic environment and evolution *in* J. Huxley, ed. Evolution as a Process.
Allen and Unwin, London.
Otto, S. P. and M. C. Whitlock. 1997. The probability of fixation in populations of changing size. Genetics
146:723-733. (http://doi.org/10.1093/genetics/146.2.723).

Otto, S. P. and M. C. Whitlock. 2013. Fixation Probabilities and Times. eLS.

Peng, B. and M. Kimmel. 2005. simuPOP: a forward-time population genetics simulation environment.
Bioinformatics 21:3686-3687. (http://doi.org/10.1093/bioinformatics/bti584).

Rego, A., F. J. Messina, and Z. Gompert. 2019. Dynamics of genomic change during evolutionary rescue
in the seed beetle Callosobruchus maculatus. Mol. Ecol. 28:2136-2154.
(http://doi.org/10.1111/mec.15085).

Rosenblum, E. B., M. J. Hickerson, and C. Moritz. 2007. A multilocus perspective on colonization
accompanied by selection and gene flow. Evolution 61:2971-2985.
(http://doi.org/10.1111/j.1558-5646.2007.00251.x).

Ruzicka, F., T. Connallon, and M. Reuter. 2021. Sex differences in deleterious mutational effects in
Drosophila melanogaster: combining quantitative and population genetic insights. Genetics 219.
(http://doi.org/10.1093/genetics/iyab143).

Simpson, G. G. 1944. Tempo and Mode in Evolution. Columbia University Press, New York.

Tepolt, C. K., J. A. Darling, M. J. Bagley, J. B. Geller, M. J. Blum, and E. D. Grosholz. 2009. European green
crabs (Carcinus maenas) in the northeastern Pacific: genetic evidence for high population
connectivity and current-mediated expansion from a single introduced source population.
Divers. Distrib. 15:997-1009. (http://doi.org/10.1111/j.1472-4642.2009.00605.x).

van der Zee, M. J., J. R. Whiting, J. R. Paris, R. D. Bassar, J. Travis, D. Weigel, D. N. Reznick, and B. A.
Fraser. 2022. Rapid genomic convergent evolution in experimental populations of Trinidadian
guppies (Poecilia reticulata). Evolution Letters 6:149-161. (http://doi.org/10.1002/evl3.272).

Wade, M. J. and C. J. Goodnight. 1998. The theories of Fisher and Wright in the context of
metapopulations: When nature does many small experiments. Evolution 52:1537-1553.

Willi, Y., P. Griffin, and J. Van Buskirk. 2013. Drift load in populations of small size and low density.
Heredity 110:296-302. (http://doi.org/10.1038/hdy.2012.86).

Willis, J. H. 1999. Inbreeding load, average dominance and the mutation rate for mildly deleterious
alleles in Mimulus guttatus. Genetics 153:1885-1898.

Willoughby, J. R., A. M. Harder, J. A. Tennessen, K. T. Scribner, and M. R. Christie. 2018. Rapid genetic
adaptation to a novel environment despite a genome-wide reduction in genetic diversity. Mol.
Ecol. 27:4041-4051. (http://doi.org/https://doi.org/10.1111/mec.14726).

Wright, S. 1931. Evolution in Mendelian populations. Genetics 16:97 - 159.

Wright, S. 1932. The roles of mutation, inbreeding, crossbreeding and selection in evolution.
Proceedings of the Sixth International Congress of Genetics 1:356-366.

Yin, X. S., A. S. Martinez, M. S. Sepulveda, and M. R. Christie. 2021. Rapid genetic adaptation to recently
colonized environments is driven by genes underlying life history traits. BMC Genomics 22.
(http://doi.org/10.1186/s12864-021-07553-x).

599 Fig. 1. Mean *Δq* across allele frequencies from *q = 1/2N* to *q = 1.0* for beneficial alleles in a small (*N = 50;*
600 blue line) compared to a large (*N = 1000*) population for recessive (*h = 0.0;* left column) and codominant
601 (*h = 0.5;* right column) mutations with selection ranging from *s = 0.01* to *s = 0.20.*
602
603

632
633 Fig. 2. The proportion of deleterious mutations fixing (A) and the levels of drift load (B) across
634 population sizes of *N = 2* to 4000 after 1000 generations ($\mu = 1x10^{-3}$). Selection on mutations ranged
635 from 0.0 to 0.25 and were drawn randomly from a Poisson distribution, and dominance ranged from 0.0
636 to 0.2 and were drawn randomly from a uniform distribution. The blue lines represent the loess-
637 smoothed means from 1000 replications for each *N*, and gray shading indicate 95% confidence intervals.
638
639

657 Fig. 3. Levels of inbreeding load due to loci segregating for deleterious mutations across population sizes
658 of $N = 2$ to 4000 after 1000 generations ($\mu = 1x10^{-3}$). Selection on mutations ranged from 0.0 to 0.25 and
659 were drawn randomly from a Poisson distribution, and dominance ranged from 0.0 to 0.2 and were
660 drawn randomly from a uniform distribution. The blue line represents the loess-smoothed means from
661 1000 replications for each N, and gray shading indicate 95% confidence intervals.
662

678 Fig. 4. Effects of population size on number of generations for fixation of beneficial alleles from
679 simulations ($h = 0.5$) compared to the diffusion approximation averaged across 10 replicates per
680 population size (‒) for $s = 0.02$ (red), 0.04 (black), 0.10 (blue), and 0.20 (green). The lines
681 represent the loess-smoothed means across 100 replications for each N, and gray shading indicates 95%
682 confidence intervals.
683

684
685   Fig. 5. The proportion of loci fixing for random beneficial mutations introduced in the first generation
686   with *s = 0.0* to *0.2* (Poisson distribution) and *h = 0.0* to *1.0* (flat distribution) after 10,000 generations for
687   population sizes ranging from *N = 4* to *3000*.

Fig. 6. Accumulation of beneficial mutations (A; fixation flux), mutations lost (B), and the proportion
continuing to segregate (C) after 1000 (blue), 2000, (purple), and 4000 (magenta) generations with *Ns* =
2, and over 1000 generations with *s = 0* (red) with no prior burn-in. The lines represent the loess-
smoothed means across 100 replications for each *N*, and gray shading indicates 95% confidence
intervals.

Appendix 2. Supplemental Figures.

761   Fig. S1. Mean *Δq* across allele frequencies from *q = 1/2N* to *q = 1.0* for beneficial alleles in a small (*N =*
762   *50;* blue line) compared to a large (*N = 1000*; red line) population for dominant mutations (*h = 1.0*) with
763   selection ranging from *s = 0.01* to *s = 0.20.*

764
765

786   Fig. S2. Mean *Δq* for beneficial mutations going to fixation from an initial frequency of *1/2N* across
787   selection coefficients from *s = 0.01* to 0.20 for *N = 100* (green line) and *N = 1000* (blue line). The line
788   represents the loess-smoothed means across 100 replications for each *N*, and gray shading indicates the
789   95% confidence interval.
790
791

807   Fig. S3. Standard deviation of *Δq* for beneficial mutations going to fixation from an initial frequency of
808   *1/2N* across selection coefficients from *s = 0.01* to 0.20 for *N = 100* (green line) and *N = 1000* (blue line).
809   The line represents the loess-smoothed means across 100 replications for each *N*, and gray shading
810   indicates the 95% confidence interval.
811
812
813

846   Fig. S4. The diffusion approximation for the probability of fixation (   ) in response to initial allele

847   frequency with *Ns = 2* (A), and the proportion of fixed alleles across 100 loci after 1000 generations for

848   population sizes of *N = 100* and 1000 with *Ns = 2* (B; the relationship for   is obscured). Lines represent

849   the loess-smoothed relationship of means for 20 replications per population size.