Structure and Function of Gli123 Involved in *Mycoplasma mobile* Gliding

Daiki Matsuike,[a] Yuhei O Tahara,[a,b] Takahiro Nonaka,[a] Tasuku Hamaguchi,[c] Hisashi Kudo,[d,e] Yuuki Hayashi,[d,f] Munehito Arai,[d,g] and Makoto Miyata[a,b]

[a] Graduate School of Science, Osaka Metropolitan University, Sumiyoshi-ku, Osaka, Japan

[b] The OCU Advanced Research Institute for Natural Science and Technology (OCARINA), Osaka Metropolitan University, Sumiyoshi-ku, Osaka, Japan

[c] Institute of Multidisciplinary Research for Advanced Materials (IMRAM), Tohoku University, Aoba-ku, Miyagi, Japan

[d] Department of Life Sciences, The University of Tokyo, Meguro Tokyo, Japan

[e] Graduate School of Science, Technology and Innovation, Kobe University, Nada, Kobe, Japan

[f] Environmental Science Center, The University of Tokyo, Bunkyo, Tokyo, Japan

[g] Department of Physics, The University of Tokyo, Meguro, Tokyo, Japan

Running Title: *Mycoplasma* surface gliding protein

[#] Address correspondence to Makoto Miyata, miyata@omu.ac.jp

ABSTRACT

Mycoplasma mobile is a fish pathogen that glides on solid surfaces by means of a

unique mechanism. The gliding machinery of *M. mobile* is composed of internal and

surface structures. In the present study, we focused on the function and structure of

Gli123, a surface protein that is essential for the localization of other surface proteins.

The amino acid sequence of Gli123, which is 1128 amino acids long, contains

lipoprotein-specific repeats. We isolated the native Gli123 protein from *M. mobile* cells

and a recombinant protein, rGli123, from *Escherichia coli*. The isolated rGli123

complemented a non-binding and non-gliding mutant of *M. mobile* that lacked Gli123.

Circular dichroism and rotary-shadowing electron microscopy (EM) showed that rGli123

has a structure that is not significantly different from that of the native protein.

Rotary-shadowing EM suggested that the molecules changed their shape between

globular and rod-like structures, depending on the ionic strength of the solution.

Negative-staining EM coupled with single-particle analysis revealed that Gli123 forms a

globular structure featuring a small protrusion with dimensions of 20.0, 14.5, and 16.0

nm. Small-angle X-ray scattering analyses indicated a rod-like structure composed of

several tandem globular domains with total dimensions of approximately 34 nm length

and 4 nm width. Both molecular structures were suggested to be dimers based on the

predicted molecular size and structure. Gli123 may have evolved by multiplication of repeating lipoprotein units and acquired clumping role of surface proteins.

IMPORTANCE

Mycoplasmas are pathogenic bacteria that are widespread in animals. They are characterized by small cell and genome sizes but are equipped with unique abilities to escape host immunity, such as surface variation and gliding. Here, we focused on a surface-localizing protein that is essential for *Mycoplasma mobile* gliding. The findings of this study suggested that the protein undergoes drastic conformational changes between its rod-like and globular structures. These changes may be caused by a repetitive structure common in the surface proteins that is responsible for the modulation of the cell surface structure and related to the assembly process for the surface gliding machinery. An evolutionary process for this unique mycoplasma gliding mechanism has also been suggested in the present study.

KEYWORDS

electron microscopy, small-angle X-ray scattering, repeat sequence, single-particle analysis, mollicutes, lipoprotein, conformational change, motility

INTRODUCTION

Class Mollicutes, which includes the genus *Mycoplasma*, consists of organisms fit to

parasitic or commensal life cycles, as represented by their small genome size, lack of

peptidoglycan layer, and antigenic modulation (1-3). Interestingly, more than 20

Mycoplasma species are known to glide on solid surfaces through unique mechanisms

(4). *Mycoplasma* gliding mechanisms can be classified into two types, as represented

by either *Mycoplasma mobile* or *Mycoplasma pneumoniae* (4, 5). Four species are

known to perform mobile-type gliding, including *M. mobile*, *Mycoplasma pulmonis*,

Mycoplasma testudineum, and *Mycoplasma agassizii*, which serve as pathogens for

freshwater fish, mice, turtles, and turtles, respectively (6, 7). The gliding mechanism has

been studied with a focus on *M. mobile*, which glides at speeds of 2.0 to 4.5 µm/s (4, 8).

The gliding machinery, which forms as a protrusion, can be divided into two parts: the

inside and surface structures (4, 9, 10). Its surface structure is composed of four

proteins: Gli123, Gli349, Gli521, and Gli42. Gli123, Gli349, and Gli521 have been

reported as being essential for binding and gliding, based on analyses of mutants

missing each of these proteins (11-13). Gli349 is a flexible rod-shaped, 95-nm-long

protein with a C-terminal globular domain (11, 14, 15). It catches sialylated

oligosaccharides on the surfaces of hosts as a "leg" for gliding, and pulls the cell (16,

17). Gli521 is a rather rigid rod-shaped 120-nm-long protein with N-terminal globular

and C-terminal hook domains (18, 19) that is suggested to transmit the gliding force

generated in an internal motor to Gli349, like a "crank" (5, 18, 19). Gli123 localizes

Gli349 and Gli521 correctly to the gliding machinery, as a "mount" (13). Gli349 and

Gli521 are conserved in all four species of *M. mobile*, *M. pulmonis*, *M. testudineum*, and

M. agassizii, whereas Gli123 and Gli42 are only found in *M. mobile* and *M. pulmonis*

(20). In this study, we isolated Gli123 and found that a mutant lacking this protein could

be complemented by the isolated protein instead. We analyzed Gli123 structures using

electron microscopy (EM), small-angle X-ray scattering (SAXS), and other biochemical

methods, and suggested a plausible structural model and its conversion, which may be

related to the role of the protein in gliding motility.

RESULTS

Gli123 orthologs in the *Mycoplasma* species

Orthologs of *gli123* and *gli42* genes were not found in *M. agassizii* and *M. testudineum*,

unlike the orthologs of nine other genes involved in *M. mobile* gliding (20). The orthologs

of *gli349* and *gli521* from *M. testudineum* and *M. agassizii* have been found to be

positioned closer to *M. pulmonis* than to *M. mobile* in the phylogenetic tree (20).

Therefore, we performed a PSI-BLAST search (21) using the amino acid sequences of

Gli123 and Gli42 of *M. pulmonis*, and found the orthologs of *gli123* and *gli42* in the

genomes of *M. agassizii* and *M. testudineum* (Table S1). The phylogenetic trees

constructed for *gli123*, *gli42*, *gli349*, and *gli521* genes using the maximum likelihood

method shared their topology, as did the phylogenetic tree of 16S ribosomal DNA

sequences (Fig. S1) (20). Next, we examined the localization of these genes in the

genome. In *M. mobile* and *M. pulmonis*, the genes were clustered in one locus (13, 20),

while in *M. agassizii* and *M. testudineum*, they were split into two loci, *gli123-gli42* and

gli349-gli521, although the actual distances are unknown in the unassembled genome

sequences (Fig. S2).

Sequence analysis of Gli123 and the lipoprotein-17 domain

In this study, we found that the amino acid sequences of Gli123 protein and its orthologs

include four to six lipoprotein-17 domains, which are composed of approximately 100

amino acids (Fig. 1A) (22, 23). The lipoprotein-17 domain is found in 233 proteins of 43

bacterial species, mostly in lipoproteins that play a role in antigenic variation or

adhesion. In *Mycoplasma,* the lipoprotein-17 domain was identified in 221 proteins from

32 species (Table S2). This domain was found in 15 Mvsps, Gli123, and Gli349 in *M.*

mobile (23, 24). The Gli123 protein of *M. mobile* had four lipoprotein-17 domains, while

that of *M. pulmonis* and *M. testudineum* had five and that of *M. agassizii* had six (Fig. 1A

and Table S3). Mvsps have a number of lipoprotein-17 domains, up to 16 (Fig. 1A and

Table S3). This may suggest that the surface proteins adjust their lengths by means of

repetitive fusion of the domain (4). The hidden Markov model scores for the

lipoprotein-17 domain were distributed between 30 and 70 for the Mvsps repeats and

between 10 and 30 for the Gli123 and Gli349 repeats (Fig. 1B).

MvspI mutant strains of *M. mobile*

The isolation of native Gli123 (nGli123) from *M. mobile* cell lysates is a long process

because the lipoprotein, MvspI encoded as MMOB3340, behaves in a similar way in the

isolation process. MvspI is characterized by a mass of 220 kDa, with 16 copies of

lipoprotein-17 domains (24). We isolated a spontaneous MvspI-deficient mutant to

simplify the isolation process of Gli123. Colonies that were non-reactive to a monoclonal

antibody against MvspI were screened and isolated through colony blotting and Amido

Black staining (Fig. S3A). Two colonies that were non-reactive to the antibody were

isolated from an approximate of 10000 colonies. Sodium dodecyl

sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) analysis of the isolate revealed

a lack of MvspI protein (Fig. S3B). The band intensities of the surface protein, MvspF

encoded as MMOB3300, was higher in the isolated mutant than in the wild-type (WT)

strain (Fig. S3B). Genome analysis revealed that the *mvspI* gene of the isolate had a

frameshift mutation, resulting in a truncated peptide of 97 amino acids (Table S4). In

addition to mutations in the *mvspI* gene, nine other mutations were identified in *mvsps*

and others (Table S4). This mutant grew constantly through serial cultivation of large

volumes, without recovery of MvspI. The mutant cells glided normally, but the culture

tended to form clumps, suggesting different surface interactions from those of the WT.

This mutant was named the MvspI⁻ strain and used in subsequent experiments.

Isolation of Gli123 from *M. mobile*

We isolated the nGli123 protein from cultured cells of the *M. mobile* MvspI⁻ strain (Fig.

2A). The collected cells were suspended in a buffer containing 1% Triton™ X-100 and

treated on ice, for 1 h. Following that, the lysates were separated by means of

centrifugation into a precipitate and a soluble fraction containing nGli123, Gli349, Gli521,

and MMOB1650 (Fig. 2B). The soluble fraction was then subjected to anion-exchange

chromatography. The nGli123 protein was bound to an anion-exchange column with 20

mM NaCl and eluted using a gradient of NaCl. Gli521 and MMOB1650 were separated

from nGli123 by means of this elution process (Fig. 2B). Next, we subjected the

fractions containing nGli123 and Gli349 to size-exclusion chromatography (SEC). The

proteins, which were traced by means of absorbance, were divided into four peaks,

other than the void-volume fraction (Fig. 2C). SDS-PAGE and mass spectrometry

identified nGli123 in the second peak near the position of the 150 mL elution (Fig. 2B

and 2C). The Stokes radius of nGli123 was calculated as 5.4 nm from the elution

position (Fig. 2C). The final yield of the nGli123 protein was 0.3 mg, isolated from 3 L of

growth medium. The N-terminal amino acid sequence of the nGli123 protein band

collected from the electrophoresis of the whole-cell lysate was analyzed using Edman

degradation and determined as Ala-Ile-Ala-Ile-Gly, thereby showing that the N-terminus

of 18 amino acids including the partial transmembrane segment of 11 amino acids is

truncated in the mature protein (Fig. 2A) (13).

Expression and purification of recombinant Gli123

The yield of nGli123 was not high enough to perform various experiments. Thus, we

expressed recombinant Gli123 (rGli123) with a 6×His-tag at the C-terminus of mature

Gli123 in *E. coli* (Fig. 2A) (13) and purified it using three steps: (i) nickel-nitrilotriacetic

acid (Ni^{2+}-NTA) affinity chromatography, (ii) hydrophobic interaction chromatography,

and (iii) SEC (Fig. 2B). The purified fraction was recovered by means of SEC, as a

single peak that included only the rGli123 protein, which was then confirmed using

SDS-PAGE and visualized using Coomassie Brilliant Blue (CBB) staining and mass

spectrometry (Fig. 2B and 2C). The Stokes radius of rGli123 was estimated as 5.4 nm,

from the elution position of SEC (Fig. 2C), which agreed with that for the nGli123

isolated from *M. mobile* cells. The final yield of the rGli123 protein was approximately 10

mg per 1 L of *E. coli* culture. The circular dichroism spectrum of rGli123 showed that the

secondary structural contents included 14.0% α-helices and 35.1% β-sheets (Fig. 2D)

(25, 26), consistent with the predictions made from the amino acid sequence using

XtalPred, *i.e.*, 14.0% for α-helices and 43.0% for β-sheets. The purified rGli123 protein

was subjected to limited proteolysis with trypsin, to identify the flexible and rigid parts

(Fig. 2A). The SDS-PAGE profile of the cleaved fragment showed that the rGli123

protein finally converged into three products through several fragmentation steps (Fig.

S4A). Mass spectrometry results showed that the three final fragments (f, g, and h)

corresponded at least to the 405–1128 (+8), 565–1128 (+8), and 559–789 amino acid

sequences, respectively. Based on these results, we believe that some components of

the molecule are rigid, as presented in Fig. 2A.

Complementation of the binding and gliding abilities of the *M. mobile* mutant with additional Gli123 proteins

To examine the complementation activity of Gli123 for the Gli123-lacking mutant, m12 (13), we added purified rGli123 protein to the cell suspension and incubated it for 3 h at 25°C. The cells were inserted into the tunnel slides with video recordings of phase-contrast optical microscopy (27). m12 cells were unable to bind to the glass surface (13), whereas cells with rGli123 protein showed binding to the glass surface (Fig. 3A). The average number of bound cells in a 100 μm^2 field on glass was 15.1 ± 3.4 cells (n = 15), which was 69.1% of that of the WT. 14.9 ± 8.5% of the bound cells showed unidirectional gliding, similar to the WT strain, with a speed of 1.5 ± 0.8 μm/s (n = 45), comparable with that of the WT strain, 2.7 ± 0.5 μm/s (n = 45) (Fig. 3B and 3C), suggesting that the rGli123 protein binds and complements the gliding machinery (13). Next, we examined the binding of the rGli123 proteins to the mutant cells. Mutant cells incubated with rGli123 were recovered by means of centrifugation and analyzed using SDS-PAGE (Fig. 3D). This result showed that rGli123 bound to the mutant cells at an amount comparable to that of the WT strain.

Conformational changes in Gli123

We observed the structures of the nGli123 and rGli123 proteins using rotary-shadowing

EM. In rotary-shadowing EM, protein molecules sprayed on a rotating mica surface are

shadowed by platinum particles from a low angle, to observe their molecular shape (19,

24). We observed particles of similar sizes and shapes in the field of view. The nGli123

and rGli123 proteins showed similar rod-like structures, with lengths of 35.6 ± 5.0 and

35.9 ± 4.1 nm, respectively, under conditions including 100 mM ammonium acetate (Fig.

4A and 4B). Next, we observed the nGli123 and rGli123 proteins in the presence of 500

mM ammonium acetate. Interestingly, both proteins showed globular structures with

long axes of 21.0 ± 3.2 and 20.1 ± 4.1 nm (Fig. 4A and 4B). We then examined the

molecular ratios of the two structures in 100, 250, and 500 mM ammonium acetate. The

proportions of rod-like structures at each concentration were 89.7%, 47.0%, and 5.7%,

respectively, and the ratio changed from rod-like to globular structures depending on the

ammonium acetate concentration (Fig. 4C). Furthermore, we measured the light

scattering of rGli123. Light scattering intensity measured at different concentrations

increased with ammonium acetate concentration, in the range of 50–650 mM,

suggesting that the conformation of rGli123 molecules depends on the concentration of

ammonium acetate (Fig. 4D). The 50% effective concentration (EC_{50}) was 261.0 ± 11.3,

238.3 ± 16.1, and 292.9 ± 4.9 mM at protein concentrations of 0.23, 0.75, and 1.23

mg/mL, respectively (Fig. 4D). The changes in the light scattering intensities of rGli123

proteins due to changes in ionic strength were in good agreement with the ratios of

globular and rod-like structures measured using rotary-shadowing EM (Fig. 4C and 4D).

Globular structure of Gli123

Negative-staining EM was performed to visualize the three-dimensional structure of

globular nGli123 with better resolution. The field images of EM showed uniform globular

particles of 15–20 nm (Fig. 5A), and their density changed depending on the protein

concentration used. In addition, many rod-like structures were observed in the

background. We selected globular particles (Fig. 5A) and obtained 2D-averaged images

using the structural analysis software EMAN2.91 (Fig. 5B). The 2D-averaged images

showed that the Gli123 molecules had dimensions of approximately 17 and 14 nm for

the major and minor axes, respectively. Based on the averaged image of the Gli123

protein, a three-dimensional image was reconstructed, without C1 and with C2, the

two-rotational symmetry enforcement. The three-dimensional (3D)-reconstructed

globular structure of Gli123 had a protrusion that was reminiscent of a 'spinning top' (Fig.

5C). Considering the globular molecular shape, the molecular weight of Gli123 was

estimated to be 244 kDa from the elution position of SEC, indicating that the protein

behaves as a dimer (Fig. 2C).

Rod-like structure of Gli123

SAXS was used to obtain the 3D rod-like structure of Gli123. The rGli123 protein under

low ionic strength condition was subjected to SEC-SAXS. The protein showed an

elution curve with a single peak (Fig. 6A). We analyzed the scattering curve at the peak

top of the elution profile with a Guinier plot [ln $I(Q)$ *vs.* Q^2, where $I(Q)$ is the scattering

intensity at scattering vector Q], to estimate the size of the rGli123 protein (Fig. 6B). The

Guinier plot was based on the scattering curve at small angles, and the square of the

radius of gyration (R_g) was obtained from the slope of the approximate line in the

Guinier region ($Q{\times}R_g$ < 1.3) as ln $I(Q)$ = ln $I(0) - R_g^2Q^2/3$, where $I(0)$ is the zero-angle

scattering intensity (28). R_g was calculated as 8.24 nm from the slope of the

approximate line (Fig. 6B). Next, to discuss the shape of the rGli123 protein, the

scattering curve was analyzed using a Kratky plot [$I(Q)Q^2$ *vs. Q*] (Fig. 6C), which shows

the characteristics of the protein shape (29). The plot shows a mountainous curve with a

peak around Q = 0.1 Å^{-1}, suggesting that the rGli123 protein is not in a random coil but

has a folded structure. For an ideal sphere, the Kratky plot shows a hyperbolic curve

with a single peak in the small-angle regions. However, in the present data, the plot

shows a shoulder around $Q = 0.02$ Å^{-1}, marked by an inverted black triangle in the figure

(Fig. 6C), suggesting the existence of another hidden peak. The bimodal character of

the Kratky plot suggests a dumbbell shape for the protein; the innermost shoulder

reflects the globularity of the whole molecule, while the peak reflects that of each

domain (30). According to the pair-distance distribution function, $P(r)$, the maximum

length (D_{max}) of the rGli123 protein was approximately 31.6 nm (Fig. 6D). The 3D

structure of the rGli123 protein was modeled from SAXS data using PRIMUSQT

software (Fig. 6E) (31). The structural model was rod-shaped and featured several

narrow parts, similar to an articulated robot. This structure was consistent with the

images obtained using rotary-shadowing EM with low ionic intensity (Fig. 4B). In the

cross-section plot [ln $I(Q)Q$ vs. Q^2], which tests the validity of the rod-like structure in the

SAXS model (32), the slope of the plot showed high linearity (Fig. S5). The radius of

gyration of cross-section R_c was 1.14 nm, which was obtained by fitting the

cross-section plot to the equation ln $[I(Q)Q] = $ ln $[I(0)Q] - R_c^2 Q^2/2$ (33). The radius of a

cylinder R_p (a short-axis radius) was 1.61 nm, as calculated using the equation $R_p = $

$\sqrt{2}R_c$ (33). The height of the cylinder L (twice the major-axis radius), determined from

the equation $R_g^2 = R_p^2/2 + L^2/12$ (33), was 28.2 nm. The L value obtained from the

cross-section plot was consistent with the length ($D_{max} = 31.6$ nm) estimated from the

$P(r)$ function, indicating the high validity of the rod-like shape of the SAXS model. We also believe that the rod-like structure is likely to be a dimer, because the SAXS model has a molecular volume that corresponds to two molecules of a model predicted from the amino acid sequence using Robetta (Fig. 6F) (34). SAXS measurements were also performed using static cells at three different concentrations of rGli123 (Fig. S6). These results were consistent with those obtained from the SEC-SAXS measurements, and supported the absence of aggregation in the SAXS measurements.

DISCUSSION

Surface gliding proteins

In this study, *gli123* and *gli42* genes were found in the genomes of *M. testudineum* and *M. agassizii* (Table S3), indicating that the two surface proteins of *M. mobile*, Gli123 and Gli42, are also essential for gliding, similar to Gli349 and Gli521 (20). The topologies of phylogenetic trees were common for all these proteins and 16S rDNA (20), suggesting that these mycoplasmas have acquired gliding motility in evolution, not through horizontal transfer (Fig. S1). In addition, the orthologs of *gli349* and *gli521* were positioned adjacent to the genomes of all gliding *mycoplasmas* (Fig. S2) (20), and the localization of Gli349 is dependent on that of Gli521 (13). These gliding proteins may

form a complex immediately after synthesis and establish a gliding machinery in which

binding and pulling are smoothly interlocked.

Structure and conformational changes of Gli123

We isolated the Gli123 protein and found that its molecular structure changed from

rod-like to globular depending on the ionic strength. This conformational change may

occur in the flexible regions identified through limited proteolysis (Fig. S4). A pair of

Gli123 molecules predicted from the amino acid sequence fit into both the rod-like

structure obtained using SAXS and the globular structure obtained by means of

single-particle analysis, suggesting that these structures of Gli123 are dimers (Figs. 5D

and 6F).

The structure of the isolated molecule changes drastically with ionic strength;

however, *M. mobile* can glide under different ionic strength environments, suggesting

that this conformational change is unlikely to occur in the completed machinery (13). We

observed that the purified rGli123 protein complemented the gliding of mutant cells

lacking Gli123. Conformational changes in the Gli123 molecule may be related to this

observation. These three gliding proteins have a localization hierarchy of

Gli123-Gli521-Gli349, suggesting a physical interaction between these proteins at the

cell surface (13). Gli123 might reach the target position on the cell surface as a flexible

rod shape and bundle Gli521 and Gli349 molecules on the cell surface through a drastic

conformational change, which is induced by the electrostatic fields made by the

neighboring proteins, Gli349 and Gli521.

Roles of the lipoprotein-17 domain

We found that a common repeat sequence of the lipoprotein-17 domain was shared by

the Mvsps and gliding proteins. The relationship between the gliding machinery and

Mvsps has been proposed based on the existence of a repeat comprising of

approximately 100 amino acids (22, 24). In this study, we succeeded in determining the

similarity between the repeat sequences. The existence of the lipoprotein-17 domain in

Gli123, Gli349, and their orthologs may suggest that they originated from a common

surface protein (Figs. 1A and 7A). The structure of the lipoprotein-17 domain in Gli123

and Mvsp predicted using AlphaFold2 featured three β-strands, which is consistent with

previously reported structural features determined using nuclear magnetic resonance

(NMR) and crystallography (Fig. S7A and S7B) (24). The predicted Mvsp structures

suggest that an α-helix with a short loop connects the lipoprotein-17 domains to form a

flexible chain (Fig. S7C). Although the detailed structures of Gli123 and Gli349 are still

unknown, their flexibility may be derived from the structures common with the

lipoprotein-17 domain and adjacent regions. Lipoprotein-17 domain may be useful for

adjusting molecular length and flexibility (Fig. 7A). It may provide length variation

adjustment to Mvsps and Gli349, which are involved in antigenic modulation and gliding

leg, respectively. Flexible and long leg proteins may be advantageous for catching the

host cell surface (17). In case of Gli123, it may provide length and flexibility for its role in

clamping other surface gliding proteins on the machinery surface, where as many as

450 gliding units are packed (Fig. 7B). The gliding units fixed by Gli123 can be aligned

in an ordered way for smooth and efficient gliding (13, 27).

MATERIALS AND METHODS

Sequence analyses

The genome sequences of *M. mobile* (163K), *M. pulmonis* (UAB CTIP), *M. agassizii*

(PS6), and *M. testudineum* (ATCC700618) were obtained from the National Center for

Biotechnology Information (NCBI) GenBank database

(https://www.ncbi.nlm.nih.gov/genome/). A homology search against the NCBI

non-redundant protein sequence databank was then performed using PSI-BLAST

(https://blast.ncbi.nlm.nih.gov/Blast.cgi), with an expected threshold of 100.

Phylogenetic trees of the proteins were constructed using neighbor-joining and maximum likelihood methods. The reliability of the topology of each phylogenetic tree was examined with 1000 bootstrap replicates, using the MEGAX package (https://www.megasoftware.net/). The secondary structure of Gli123 was predicted using XtalPred (https://xtalpred.godziklab.org/XtalPred-cgi/xtal.pl). Hidden Markov model scores were obtained from Pfam (http://pfam.xfam.org/family/PF04200) and UniProtKB (https://ftp.uniprot.org/pub/databases/uniprot/previous_releases/2021_03), based on the HMMER3 package (http://hmmer.org/).

Strains and culture conditions

M. mobile 163K (ATCC 43663) was grown at 25°C in Aluotto medium, to an optical density at the wavelength of 600 nm (OD_{600}) of around 0.08, as previously described (35). *The M. mobile* MvspI$^-$ strain was isolated as follows. Mutants that did not react with monoclonal antibodies against MvspI were selected using colony blotting and Amido Black staining (36). The colonies were isolated and cultured in Aluotto liquid medium, following which SDS-PAGE analysis of the isolates was carried out, which confirmed the absence of MvspI protein. *E. coli* DH5α and BL21(DE3) strains were used for DNA recombination and protein expression, respectively.

Isolation of Gli123 from *M. mobile* cells

The nGli123 protein was purified from the *M. mobile* MvspI⁻ strain cells. Cells stored at –

80°C were precultured and grown in 3 L of Aluotto liquid medium, dispensed into nine

300 cm^2 tissue culture flasks, and statically cultured at 25°C until the OD$_{600}$ reached

0.08. The cells were collected by means of centrifugation at 18,800 × *g* and 4°C, for 30

min, and washed twice with phosphate-buffered saline (PBS; 58 mM Na$_2$HPO$_4$, 17 mM

NaH$_2$PO$_4$, and 68 mM NaCl). The precipitate was suspended in 30 mL of buffer A (20

mM Tris-HCl pH 8.0 and 0.1 mM phenylmethylsulfonyl fluoride) and sonicated for 2 min,

at intervals of 5 s, using a sonicator (US-600T, Nippon Seiki Co., Niigata, Japan), on ice.

The cell lysate was combined with 30 mL of buffer A containing 2% Triton™ X-100 (v/v)

and shaken on ice for 1 h, to solubilize the proteins. The cell lysate was separated by

means of centrifugation at 32,300 × *g* and 4°C, for 30 min, following which the obtained

supernatant was applied to an anion-exchange column (HiTrap™ Q HP 5 mL; Cytiva,

Tokyo, Japan) equilibrated with buffer A. Proteins bound to the column were eluted with

a NaCl concentration gradient of 0–500 mM, at 4°C. The eluted fraction containing

Gli123 was concentrated to a volume of 10 mL and subjected to a SEC column

(HiLoad® 26/600 Superdex® 200 prep grade, Cytiva) equilibrated with SEC buffer (20

mM Tris-HCl pH 8.0 and 150 mM NaCl), at 4°C. Thyroglobulin (669 kDa), ferritin (440

kDa), aldolase (158 kDa), and ovalbumin (43 kDa), with Stokes radii of 8.5, 6.1, 4.8, and

3.1 nm, respectively, were used as standards. Protein fractions were analyzed using

SDS-PAGE with a 10% polyacrylamide gel, followed by CBB staining.

Expression and purification of recombinant Gli123

The N-terminal sequence of Gli123 was determined using Edman degradation, as

previously described (11, 12). According to the sequencing results, the DNA sequence

encoding the amino acid region from the 19th to 1128th position was codon-optimized for

expression in *E. coli* and inserted between the *Bam*HI and *Xho*I sites in the multicloning

site of the pET21a plasmid (Novagen, Madison, WI). The recombinant protein, rGli123,

was expressed in *E. coli* BL21(DE3) and induced using 1 mM

isopropylthio-β-galactoside at 15°C, for 18 h. Following that, the cells were collected,

suspended in His-buffer A (20 mM Tris-HCl pH 8.0 and 20 mM NaCl), and washed twice.

The cells were suspended in 60 mL His-buffer A with 60 μL of 100 mM

phenylmethylsulfonyl fluoride and sonicated for 4 min, at intervals of 5 s, on ice, as

described above. The lysate was centrifuged at 32,300 × *g* and 4°C, for 30 min,

following which the supernatant was subjected to Ni^{2+}-NTA affinity chromatography

(HisTrap™ HP 5 mL, Cytiva). Proteins bound to the column were eluted with an imidazole concentration gradient of 0–250 mM, at 4°C. The fraction containing rGli123 was adjusted to 20% saturation of ammonium sulfate, kept on ice for 1 h, following which the obtained supernatant was collected after centrifugation at 32,300 × g and 4°C, for 30 min. The supernatant was subjected to hydrophobic interaction chromatography column (Phenyl Sepharose HP 1 mL, Cytiva) and eluted in a gradient of 20%–0% ammonium sulfate. The rGli123 protein was concentrated to 5 mL by means of centrifugation at 5,000 × g and 4°C, using a 50k Amicon tube (Amicon Ultra, Merck, Darmstadt, Germany), and subjected to SEC. The protein concentration was adjusted using a 50k Amicon tube, if necessary, for analyses.

Optical analyses

The circular dichroism spectra of the proteins were measured in the range of 200–250 nm using a spectropolarimeter (J-805, Jasco International Co., Tokyo, Japan), at 25°C. A quartz cuvette with a 1-mm path length was filled with 1 mg/mL of rGli123 in SEC buffer. The conditions for the measurements have been previously described (37). The measured spectrum was analyzed using BeStSel (26), followed by estimation of the α-helix, β-strand, and turn contents (25). The secondary structure was predicted from

the amino acid sequence using XtalPred (38). For light scattering, the rGli123 protein

dialyzed against 50 mM ammonium acetate was placed into a 10-mm quartz cell, and

the scattered light at the wavelength of 400 nm was measured at 25°C using a

fluorescence meter (FP-6200, Jasco International Co.). The ammonium acetate

concentration was adjusted by means of addition of 3 M ammonium acetate solution.

Optical and electron microscopy

Rotary-shadowing and negative-staining EM were performed as previously described

(10, 19, 24, 39). Protein images were analyzed using the image analysis software

EMAN2.91 (https://blake.bcm.edu/emanwiki/EMAN2). The approximately 10,000

particle images were selected and classified into 50 groups, averaged, and then

reconstructed into 3D structures from 3,202 particle images. For optical microscopy, *M.*

mobile cells were inserted into a tunnel slide (20 µm) precoated with Aluotto medium, for

60 min. After 1 min, the floating cells were removed. The cells on the glass surface were

observed using phase-contrast microscopy and analyzed using ImageJ v1.51w

(http://rsb.info.nih.gov/ij/), as previously described (17, 20, 40).

Assessment of binding of rGli123 to mycoplasma cells

Cultured *M. mobile* WT and m12 cells were washed twice and suspended in PBS. The

suspended cells were kept in PBS containing 0.5 mg/mL of rGli123 protein for 3 h,

following which they were collected by means of centrifugation at 12,000 × g and 25°C,

for 5 min, suspended in PBS, sonicated, and analyzed using SDS-PAGE with CBB

staining. Protein band intensities were analyzed using ImageJ v1.51w.

SAXS measurements

SEC-SAXS measurements were performed at beamline 10C of the Photon Factory of

the High Energy Accelerator Research Organization (KEK), Tsukuba, Japan. The

rGli123 proteins were adjusted to a concentration of 5.33 mg/mL and subjected to a

SEC column (Superdex® 200 Increase 3.2/300, Cytiva) equilibrated with 100 mM

ammonium phosphate, at 4°C. The camera configuration has been previously described

(37). SAXS measurements using a static cell were performed at three different protein

concentrations. The measurements of the two-dimensional scattering images were

circularly averaged and converted to a one-dimensional scattering curve using the

SAngler software (41). The scattering curves immediately before the elution of the

protein were used as those of the blank samples. The average scattering curve of the

blanks was subtracted from the scattering curve of the protein solution to obtain the

scattering profile $I(Q)$ of the protein. R_g and $I(0)$ were obtained based on Guinier

approximation within the Guinier region ($R_gQ < 1.3$). The pairwise distance distribution

function $P(r)$ was calculated using GNOM software (31). The *ab initio* structural model

was simulated using the DAMMIF and DAMMIN softwares (31). Briefly, the outline of the

3D structure was calculated 10 times independently with DAMMIF, using dummy atoms.

The structural models were averaged using DAMAVER, and the averaged structural

models were calculated again using DAMMIF, to obtain the final structural model. The

constructed model was displayed in UCSF Chimera 1.10 (42). In the cross-section plot

(Fig. S5), the radius of gyration of the cross-section (R_c) was obtained by applying a

linear approximation to a plot of the linear region in the range after $Q = 1/R_g$ (33). The

radius of the short axis (Rp) was calculated using the equation $R_p = \sqrt{2}R_c$ (33). The

height of the cylinder (L) was calculated by substituting the values of R_g and R_p in $R_g{}^2 =$

$R_p{}^2/2 + L^2/12$ (33).

Limited proteolysis

The rGli123 protein in SEC-buffer, at the concentration of 1 mg/mL, was treated with

0.001 mg/mL trypsin, at 25°C, for various reaction times, following which the reaction

was stopped by means of heat treatment at 95°C for 5 min, in SDS-sample solution

containing 5% glycerol, 0.025% Bromophenol Blue, 62.5 mM Tris-HCl pH 6.8, 2.5%

SDS, and 5% β-mercaptoethanol. The protein digests were analyzed using SDS-PAGE

with a 10% acrylamide gel and peptide mass fingerprinting, as described previously (19,

39).

ACKNOWLEDGEMENTS

We thank Dr. Nobutaka Shimizu (KEK), Dr. Shinya Saijo (KEK), and the students of Arai

Lab (Department of Life Sciences, The University of Tokyo) for their assistance with the

SAXS measurements. We also thank Toshiaki Arata (Graduate School of Science,

Osaka City University) and Ikuko Fujiwara (Department of Bioengineering, Nagaoka

University of Technology) for their experimental advice in the light scattering analyses,

and Ms. Aya Takamori (Graduate School of Science, Osaka City University) for her

technical assistance with the MALDI-TOF MASS spectrometry. The SAXS

measurements were performed with the approval of the Photon Factory Program

Advisory Committee. This study was supported by Grants-in-Aid for Scientific Research

on the Innovative Area Harmonized Supramolecular Motility Machinery and Its Diversity

(MEXT KAKENHI grant numbers JP24117002, JP25117504, and JP15H01311),

grants-in-aid for scientific research (B) and (A) (MEXT KAKENHI grant numbers

JP24390107, JP17H01544, and JP19H02521), and JST CREST grant number

JPMJCR19S5 to MM.

REFERENCES

1.   Kirchhoff H, Rosengarten R. 1984. Isolation of a motile mycoplasma from fish. J

        Gen Microbiol 130:2439-45.

2.   Razin S, Yogev D, Naot Y. 1998. Molecular biology and pathogenicity of

        mycoplasmas. Microbiol Mol Biol Rev 62:1094-156.

3.   Grosjean H, Breton M, Sirand-Pugnet P, Tardy F, Thiaucourt F, Citti C, Barre A,

        Yoshizawa S, Fourmy D, de Crecy-Lagard V, Blanchard A. 2014. Predicting the

        minimal translation apparatus: lessons from the reductive evolution of *mollicutes*.

        PLoS Genet 10:e1004363.

4.   Miyata M, Hamaguchi T. 2016. Prospects for the gliding mechanism of

        Mycoplasma mobile. Curr Opin Microbiol 29:15-21.

5.   Miyata M, Hamaguchi T. 2016. Integrated information and prospects for gliding

        mechanism of the pathogenic bacterium *Mycoplasma pneumoniae*. Front

        Microbiol 7:960.

6.   Hatchel JM, Jurkovic DA, Balish RS, Balish MF. 2006. Differences and

correlations in gliding motility and cytadherence between species and strains of the *Mycoplasma pneumoniae* group. Abstract book for 16th International Congress of International Organization for Mycoplasmology:p137.

7.   Jurkovic DA, Friedberg AJ, Hatchel JM, Relich RF, Balish MF. 2006. Adherence and gliding motility at low temperatures by tortoise-associated mycoplasmas. Abstract book for 16th International Congress of International Organization for Mycoplasmology:p24.

8.   Miyata M. 2008. Centipede and inchworm models to explain *Mycoplasma* gliding. Trends Microbiol 16:6-12.

9.   Nishikawa MS, Nakane D, Toyonaga T, Kawamoto A, Kato T, Namba K, Miyata M. 2019. Refined mechanism of *Mycoplasma mobile* gliding based on structure, ATPase activity, and sialic acid binding of machinery. *mBio* 10.

10.   Toyonaga T, Kato T, Kawamoto A, Kodera N, Hamaguchi T, Tahara YO, Ando T, Namba K, Miyata M. 2021. Chained structure of dimeric F1-like ATPase in *Mycoplasma mobile* gliding machinery. *mBio* 12:e0141421.

11.   Uenoyama A, Kusumoto A, Miyata M. 2004. Identification of a 349-kilodalton protein (Gli349) responsible for cytadherence and glass binding during gliding of *Mycoplasma mobile*. J Bacteriol 186:1537-45.

12.   Seto S, Uenoyama A, Miyata M. 2005. Identification of a 521-kilodalton protein

      (Gli521) involved in force generation or force transmission for *Mycoplasma*

      mobile gliding. J Bacteriol 187:3502-10.

13.   Uenoyama A, Miyata M. 2005. Identification of a 123-kilodalton protein (Gli123)

      involved in machinery for gliding motility of *Mycoplasma mobile*. J Bacteriol

      187:5578-84.

14.   Adan-Kubo J, Uenoyama A, Arata T, Miyata M. 2006. Morphology of isolated

      Gli349, a leg protein responsible for *Mycoplasma mobile* gliding via glass

      binding, revealed by rotary shadowing electron microscopy. J Bacteriol

      188:2821-8.

15.   Hamaguchi T, Kawakami M, Furukawa H, Miyata M. 2019. Identification of novel

      protein domain for sialyloligosaccharide binding essential to *Mycoplasma mobile*

      gliding. FEMS Microbiol Lett 366.

16.   Mizutani M, Miyata M. 2017. Force measurement on *Mycoplasma mobile* gliding

      using optical tweezers. Bio Protoc 7:e2127.

17.   Mizutani M, Tulum I, Kinosita Y, Nishizaka T, Miyata M. 2018. Detailed analyses

      of stall force generation in *Mycoplasma mobile* gliding. Biophys J

      114:1411-1419.

18. Uenoyama A, Seto S, Nakane D, Miyata M. 2009. Regions on Gli349 and Gli521 protein molecules directly involved in movements of *Mycoplasma mobile* gliding machinery, suggested by use of inhibitory antibodies and mutants. J Bacteriol 191:1982-5.

19. Nonaka T, Adan-Kubo J, Miyata M. 2010. Triskelion structure of the Gli521 protein, involved in the gliding mechanism of *Mycoplasma mobile*. J Bacteriol 192:636-42.

20. Tulum I, Kimura K, Miyata M. 2020. Identification and sequence analyses of the gliding machinery proteins from *Mycoplasma mobile*. Sci Rep 10:3792.

21. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 25:3389-402.

22. Mani R, Vorobiev S, Swapna GV, Neely H, Janjua H, Ciccosanti C, Xiao R, Acton TB, Everett JK, Hunt J, Montelione GT. 2011. Solution NMR and X-ray crystal structures of membrane-associated lipoprotein-17 domain reveal a novel fold. J Struct Funct Genomics 12:27-32.

23. Wu HN, Kawaguchi C, Nakane D, Miyata M. 2012. "Mycoplasmal antigen modulation," a novel surface variation suggested for a lipoprotein specifically

localized on *Mycoplasma mobile*. Curr Microbiol 64:433-40.

24. Adan-Kubo J, Yoshii SH, Kono H, Miyata M. 2012. Molecular structure of

    isolated MvspI, a variable surface protein of the fish pathogen *Mycoplasma*

    mobile. J Bacteriol 194:3050-7.

25. Micsonai A, Wien F, Kernya L, Lee YH, Goto Y, Refregiers M, Kardos J. 2015.

    Accurate secondary structure prediction and fold recognition for circular

    dichroism spectroscopy. Proc Natl Acad Sci U S A 112:E3095-103.

26. Micsonai A, Wien F, Bulyaki E, Kun J, Moussong E, Lee YH, Goto Y, Refregiers

    M, Kardos J. 2018. BeStSel: a web server for accurate protein secondary

    structure prediction and fold recognition from the circular dichroism spectra.

    Nucleic Acids Res 46:W315-W322.

27. Morio H, Kasai T, Miyata M. 2016. Gliding direction of *Mycoplasma mobile*. J

    Bacteriol 198:283-90.

28. Trewhella J, Duff AP, Durand D, Gabel F, Guss JM, Hendrickson WA, Hura GL,

    Jacques DA, Kirby NM, Kwan AH, Perez J, Pollack L, Ryan TM, Sali A,

    Schneidman-Duhovny D, Schwede T, Svergun DI, Sugiyama M, Tainer JA,

    Vachette P, Westbrook J, Whitten AE. 2017. 2017 publication guidelines for

    structural modelling of small-angle scattering data from biomolecules in solution:

an update. Acta Crystallogr D Struct Biol 73:710-728.

29. Arai M, Kondrashkina E, Kayatekin C, Matthews CR, Iwakura M, Bilsel O. 2007.
Microsecond hydrophobic collapse in the folding of Escherichia coli dihydrofolate
reductase, an alpha/beta-type protein. J Mol Biol 368:219-29.

30. Kataoka M, Nishii I, Fujisawa T, Ueki T, Tokunaga F, Goto Y. 1995. Structural
characterization of the molten globule and native states of apomyoglobin by
solution X-ray scattering. J Mol Biol 249:215-28.

31. Manalastas-Cantos K, Konarev PV, Hajizadeh NR, Kikhney AG, Petoukhov MV,
Molodenskiy DS, Panjkovich A, Mertens HDT, Gruzinov A, Borges C, Jeffries
CM, Svergun DI, Franke D. 2021. ATSAS 3.0: expanded functionality and new
tools for small-angle scattering data analysis. J Appl Crystallogr 54:343-355.

32. Arai K, Horikawa Y, Shikata T, Iwase H. 2020. Reconsideration of the
conformation of methyl cellulose and hydroxypropyl methyl cellulose ethers in
aqueous solution. RSC Adv 10:19059-19066.

33. Fukuyama K, Kasahara Y, Kasahara N, Oya A, Nishikawa K. 2001. Small-angle
X-ray scattering study of the pore structure of carbon fibers prepared from a
polymer blend of phenolic resin and polystyrene. Carbon 39:287-290.

34. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,

Tunyasuvunakool K, Bates R, Zidek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. Nature 596:583-589.

35. Aluotto BB, Wittler RG, Williams CO, Faber JE. 1970. Standardized bacteriologic techniques for the characterization of *Mycoplasma* species. Int J Syst Bacteriol 20:35-58.

36. Goldman A, Harper S, Speicher DW. 2016. Detection of Proteins on Blot Membranes. Curr Protoc Protein Sci 86:10 8 1-10 8 11.

37. Kenri T, Kawakita Y, Kudo H, Matsumoto U, Mori S, Furukawa Y, Tahara YO, Shibayama K, Hayashi Y, Arai M, Miyata M. 2019. Production and characterization of recombinant P1 adhesin essential for adhesion, gliding, and antigenic variation in the human pathogenic bacterium, *Mycoplasma pneumoniae*. Biochem Biophys Res Commun 508:1050-1055.

38. Slabinski L, Jaroszewski L, Rychlewski L, Wilson IA, Lesley SA, Godzik A. 2007. XtalPred: a web server for prediction of protein crystallizability. Bioinformatics

23:3403-5.

39.   Kawakita Y, Kinoshita M, Furukawa Y, Tulum I, Tahara YO, Katayama E, Namba

      K, Miyata M. 2016. Structural study of MPN387, an essential protein for gliding

      motility of a human-pathogenic bacterium, *Mycoplasma pneumoniae*. J Bacteriol

      198:2352-9.

40.   Mizutani M, Miyata M. 2019. Behaviors and energy source of *Mycoplasma*

      gallisepticum gliding. J Bacteriol 201.

41.   Shimizu N, Yatabe K, Nagatani Y, Saijyo S, Kosuge T, Igarashi N. 2016.

      Software development for analysis of small-angle x-ray scattering data. AIP

      Conference Proceedings 1741.

42.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC,

      Ferrin TE. 2004. UCSF Chimera--a visualization system for exploratory research

      and analysis. J Comput Chem 25:1605-12.

FIG. 1 Lipoprotein-17 domains in Mvsps, Gli123, and Gli349. (A) Distributions of lipoprotein-17 domains in Mvsps, Gli349, and Gli123 sequences have been shown using white boxes. MYPU_2160, CJF60_02770, and EI74_0698 are Gli123 orthologs of *M. pulmonis*, *M. agassizii*, and *M. testudineum*, respectively. (B) Column distribution of the scores calculated using the hidden Markov model (HMM) for each lipoprotein-17 domain. The scores for each lipoprotein-17 domain of Gli123 and Gli349 and orthologs in *M. mobile, M. pulmonis*, *M. agassizii*, and *M. testudineum* and Mvsps in *M. mobile* are shown.

A

nGli123

MGFSLKKSLIIGPIIAVGAIAIGVPVIVATT
1   1128

rGli123

T7-Tag
MASMTGGQQMGRGSAIAIGVPVIVATT

His-Tag
LEHHHHHH

(+14)19
Transmembrane
1128(+8)

Limited proteolysis

100 200 300 400 500 600 700 800 900 1000 1100 (aa)

B

kDa 1 2 3 4 5
250 —
150 —
100 —
75 —
50 —
35 —

kDa 1 2 3 4 5 6
250 —
150 —
100 —
75 —
50 —
37 —

C

nGli123
rGli123
Marker

A₂₈₀(AU) — A_{280}(AU)
Stokes radius (nm)
Elution volume (ml)

D

α-helix : 14.0%
β-sheet : 35.1%

θ(deg·cm²/decimol), × 10⁴ — θ(deg·cm^2/decimol), × 10^4
Wavelength (nm)

FIG. 2 Amino acid sequence and isolation process of native and recombinant Gli123 proteins (nGli123 and rGli123, respectively). (A) Schematic illustration of the amino acid sequences of nGli123 and rGli123. Transmembrane segment and tagging sequences are marked in red and green, respectively. The blue line and triangle indicate the truncated part of the mature protein. The hatched boxes indicate the lipoprotein-17 domains. The bottom diagram for limited proteolysis shows overlap of degradation products in the digest, suggesting rigid areas, as indicated by the four shades. (B) Left: protein profiles of the fractions obtained in the process of isolation of nGli123 from *M. mobile*. Lane 1, whole-cell lysate; Lane 2, Triton™-insoluble fraction; Lane 3, Triton™-soluble fraction; Lane 4, nGli123 fraction eluted by means of anion-exchange chromatography; Lane 5, nGli123 fraction eluted by means of size-exclusion chromatography (SEC). The protein bands of Gli521, Gli349, MMOB1650, and nGli123 are marked using green, red, purple, and magenta circles, respectively. Right: protein profiles of fractions in the rGli123 purification process. Lane 1, whole-cell lysate; Lanes 2 and 3, insoluble and soluble fractions of the lysate after sonication, respectively; Lane 4, rGli123 fraction after Ni²⁺-NTA affinity chromatography; Lanes 5, rGli123 fraction after hydrophobic interaction chromatography; Lanes 6, rGli123 fraction after SEC. The protein fractions were applied to SDS-PAGE with a 10% acrylamide gel and stained with Coomassie Brilliant Blue. The molecular masses are indicated on the left. (C) SEC of nGli123 and rGli123. Thyroglobulin (669 kDa, 8.5 nm), ferritin (440 kDa, 6.1 nm), aldolase (158 kDa, 4.8 nm), and ovalbumin (43 kDa, 3.1 nm) were used as the markers for molecular mass and Stokes radius. The triangles mark the peaks of nGli123 and rGli123. (D) CD spectrum of rGli123.

FIG. 3 Recovery of gliding and binding by Gli123 proteins in m12 mutant cells. (A) Cell movements analyzed using optical microscopy. The cells were observed using phase-contrast microscopy (left) and a video was recorded. The video frame (at every 0.03 s) was overlaid as a stack for 10 s, changing its color from red to blue (right). (B) Number of cells binding and gliding on glass surfaces. (C) Gliding speeds. (D) Left: Protein profiles of m12 mutant and m12 wild-type cells, after addition of rGli123 protein. The protein bands for Gli521, Gli349, and Gli123 are marked using green, red, and magenta circles, respectively. Right: Molecular ratios of Gli123, Gli349, and Gli521 were calculated from the band intensities normalized to the value of MvspI.

FIG. 4 Ionic strength-induced conformational changes in the nGli123 and rGli123 proteins. Field images of the nGli123 (A) and rGli123 (B) proteins under the conditions of 100 mM (left) and 500 mM (right) ammonium acetate. Scale bar: 50 nm. The representative particle images are magnified in the bottom. Scale bar: 10 nm. The distributions of diameter and length are shown for the globular and rod-like structures on the right. (C) Ratio of globular and rod-like structures of the rGli123 protein under the conditions of 100, 250, and 500 mM ammonium acetate. (D) Light scattering of rGli123 proteins, as a function of ammonium acetate concentration.

FIG. 5 Globular structure of Gli123 protein, as observed by means of negative-staining EM. (A) Field image of Gli123 molecules that were negatively stained with 2% uranyl acetate. Representative particle images from the fields are shown in the bottom. Single-particle analysis was performed using EMAN2.91 software. (B) Representative averaged images of Gli123 were selected from 50 classes which were obtained by averaging approximately 10,000 particles using the command "2D Analysis". (C) Three-dimensional reconstruction of Gli123 molecule by means of *ab initio* reconstruction with the parameters; C1, non-symmetrical; C2, two-rotational symmetry. (D) Superposition of the C2 model structures obtained from the single-particle analysis and those predicted from the amino acid sequence (1–1000 aa) using Robetta.

FIG. 6 Rod-like structure of rGli123 protein, as predicted using SEC-SAXS. (A) SEC of rGli123 was carried out by measuring the UV absorbance at the wavelength of 280 nm, in the presence of 100 mM ammonium acetate. (B) Scattering intensities transformed as a Guinier plot [ln I(Q) versus Q^2]. Data-points in a Guinier region ($R_gQ < 1.3$), which are shown in the figure using filled circles, were then used to determine the R_g value and zero-angle scattering intensity I(0) with Guinier approximation. (C) Scattering intensities transformed as a Kratky-plot [I(Q)Q^2 versus Q]. White and black triangles indicate the peak and hidden peak (shoulder), respectively. (D) Pair-distance distribution function P(r) calculated from the scattering curve. (E) SAXS models of rGli123. (F) Overlay of SAXS models and the structures predicted from the amino acid sequence (1–1000 aa) using Robetta.

FIG. 7 Role of lipoprotein-17 domain repeats and Gli123. (A) The repeat sequence is possibly used to adjust the length and flexibility of surface molecules that are involved in antigenic modulation (Mvsp) and gliding (Gli349 and Gli123). (B) The flexibility and molecular length of Gli123 allow for conformational changes in it, which facilitate its clamping to other gliding proteins at appropriate positions in the machinery.