Expression Ratios of the Anti-Apoptotic BCL2 Family Members Dictate the Selective

Addiction of KSHV-Transformed Primary Effusion Lymphoma Cell Lines to MCL1

Daniel Dunham[a,b,c], Prasanth Viswanathan[a,b,c], Jackson Gill[d], Mark Manzano[a,b,c]#

[a]Department of Microbiology and Immunology, University of Arkansas for Medical Sciences,

Little Rock, Arkansas, USA

[b]Center for Microbial Pathogenesis and Host Inflammatory Responses, University of Arkansas

for Medical Sciences, Little Rock, Arkansas, USA

[c]Winthrop P. Rockefeller Cancer Institute, University of Arkansas for Medical Sciences Little

Rock, Arkansas, USA

[d]Department of Biological Sciences, Henderson State University, Arkadelphia, Arkansas, USA

Running Head: MCL1 Addiction in KSHV-transformed PEL Cells

#Address correspondence to Mark Manzano, mmanzano@uams.edu

Abstract Word Count: 247, Article Word Count: 3546

ABSTRACT

Kaposi's sarcoma-associated herpesvirus (KSHV) causes several malignancies in people living

with HIV, including primary effusion lymphoma (PEL). PEL cell lines exhibit oncogene addictions

to both viral and cellular genes. Using CRISPR screens, we previously identified cellular

oncogene addictions in PEL cell lines, including MCL1. MCL1 is a member of the BCL2 family,

which functions to prevent intrinsic apoptosis and has been implicated in several cancers.

Despite the overlapping functions of the BCL2 family members, PEL cells are only dependent

on MCL1 suggesting that MCL1 may have non-redundant functions. To investigate why PEL

cells exhibit selective addiction to MCL1, we inactivated the intrinsic apoptosis pathway by

engineering BAX/BAK1 double knockout cells. In this context, PEL cells become resistant to

MCL1 knockdown or MCL1 inactivation by the MCL1 inhibitor S63845, indicating that the main

function of MCL1 in PEL cells is to prevent BAX/BAK1-mediated apoptosis. The selective

requirement to MCL1 is due to MCL1 being expressed in excess over the BCL2 family. Ectopic

expression of several BCL2 family proteins ,as well as the KSHV BCL2 homolog, significantly

decreased basal caspase 3/7 activity and buffered against staurosporine-induced apoptosis.

Finally, over-expressed BCL2 family members can functionally substitute for MCL1, when it is

inhibited by S63845. Together our data indicate that the expression levels of the BCL2 family

likely explain why PEL tumor cells are highly addicted to MCL1. Importantly, our results suggest

that caution should be taken when considering MCL1i as a monotherapy regimen for PEL,

because resistance can easily develop.

IMPORTANCE

Primary effusion lymphoma (PEL) is caused by Kaposi's sarcoma-associated herpesvirus. We

previously showed that PEL cell lines require the anti-apoptotic protein MCL1 for survival, but

not the other BCL2 family proteins. This selective dependence to MCL1 is unexpected as the

BCL2 family functions similarly in preventing intrinsic apoptosis. Recently, new roles for MCL1

not shared with the BCL2 family have emerged. Here, we show that non-canonical functions of

MCL1 are unlikely essential. Instead, MCL1 mainly functions to prevent apoptosis. The specific

requirement to MCL1 is due to MCL1 being expressed in excess over the BCL2 family.

Consistent with this model, shifting these expression ratios changes the requirement away from

MCL1 and towards the dominant BCL2 family gene. Together, our results indicate that although

MCL1 is an attractive chemotherapeutic target to treat PEL, careful consideration must be taken

as resistance to MCL1-specific inhibitors easily develops through BCL2 family overexpression.

KEYWORDS: KSHV, primary effusion lymphoma, MCL1, cell death, apoptosis, cell death,

BCL2, vBCL2, S63845

INTRODUCTION

Kaposi's sarcoma-associated herpesvirus (KSHV) establishes lifelong latency in B cells. While

KSHV infections are often asymptomatic, they can develop into lymphoproliferative disorders

such as multicentric Castleman disease and primary effusion lymphoma (PEL) in people living

with HIV (1). PEL is a monoclonal B cell malignancy often found as effusions in body cavities

(2). Patients treated with modified etoposide, vincristine, and doxorubicin with

cyclophosphamide and prednisone (EPOCH) have a median survival time of 22 months and a

three-year survival rate of 47% (3). To date, no cancer-specific treatment is available for PEL.

PEL cell lines exhibit gene expression profiles closest to plasma cells (4) with elevated levels of

key plasma cell-associated transcripts such as the master transcription factor interferon

regulator factor 4 (IRF4) and genes involved in the unfolded protein response and endoplasmic

reticulum stress (ER) pathways (4, 5). The majority of PEL tumors (~80%) are also co-infected

with the related gammaherpesvirus Epstein-Barr virus (EBV). The presence of EBV appears to

improve prognosis in patients (3). Genetic analyses of PEL tumors do not show mutations or

translocations found in other hematological malignancies such as p53, MYC, BCL2, or BCL6 (2,

6-8) although alterations in fragile site tumor suppressors have been reported (9). Instead, the

defining feature of PEL is that all tumor cells are latently infected with KSHV. The latent virus

constitutively expresses only a narrow subset of latency genes and at least 20 mature viral

microRNAs.

It is thought that constitutive expression of the KSHV latency genes drives the transformation of

PEL. Knockdown of the viral cyclin, viral FLICE inhibitory protein, viral interferon regulatory

factors 1-3 (vIRF1 to 3), or viral interleukin 6 all led to a decrease in survival of PEL cells in

culture (10-14). These latency genes regulate cellular networks to constitutively activate

pathways necessary to support the survival and proliferation of these infected cells. This creates

"oncogenic dependencies or addictions" to cellular genes, because tumor cells rely on the

continuous signaling of these pathways. For instance, addictions to several oncogenes have

been reported in PEL cell lines including the constitutive NF-κB signaling, phosphatidylinositol-

3-kinase-mTOR signaling, gp130-STAT3 signaling, high expression of the p53 inhibitor MDM2

and the IRF4 transcription factor, and c-Myc stabilization (7, 15-23).

We recently performed genome-wide CRISPR knockout screens to comprehensively identify

human oncogenic addictions or cellular genes that are required for the survival and/or

proliferation of PEL cell lines (24). Our study confirmed the addictions to well-described

pathways above (Fig. 1A). Importantly, we uncovered new dependencies that are consistently

required in eight different PEL cell lines which included genes that are commonly essential

regardless of cell type and genes that are strongly selective. Of note, we identify a strong

addiction to the myeloid cell leukemia-1 gene (MCL1). MCL1 belongs to the BCL2 gene family,

of which there are six members: MCL1, BCL2, BCL-xL (encoded by *BCL2L1*), BCL-W (encoded

by *BCL2L2*), A1 (encoded by *BCL2A1*), or Diva (encoded by *BCL2L10*). The BCL2 family

functions to inhibit intrinsic apoptosis. In healthy cells, monomers of the apoptotic effector BAX

or BAK1 are sequestered on the mitochondrial outer membrane (MOM) by the BCL2 family

proteins (Fig. 1B). When intrinsic apoptosis is initiated, pro-apoptotic BH3-only proteins (e.g.

PUMA, NOXA) bind to the BCL2 family proteins to release the BAX/BAK1 monomers. The

monomeric effectors oligomerize to form pores in the MOM and trigger an apoptotic cascade

(25). MOM permeabilization is often considered to be the "point-of-no-return" for cell death.

Although the BCL2 family members share the same function in BAX/BAK1 sequestration, PEL

cell lines appear to be only addicted to MCL1 (Fig. 1C). In recent years, new roles beyond

apoptosis have been described for MCL1. MCL1 has been implicated in mitochondrial fusion,

respiration, fatty acid oxidation, and autophagy (26-28). Thus, we hypothesized that the

selective oncogene addiction to MCL1 by PEL cell lines is due to essential functions that are not

shared with other BCL2 family proteins. In this study, we investigate the basis for why KSHV-

transformed PEL cell lines exhibit selective dependence on only MCL1 among the BCL2 gene

family. Contrary to our initial hypothesis, we show that the main essential function of MCL1 in

PEL cells is to block intrinsic apoptosis. PEL cell lines likely depend on MCL1 because it is

expressed at 10-60-fold higher levels than the other BCL2 family members. Altering the

expression ratios by over-expression of the other BCL2-related proteins, including the KSHV

BCL2 homolog, rescues cells from intrinsic apoptosis induced by an inhibitor of MCL1. This

suggests that endogenous expression levels of the BCL2 proteins determine the selective

addiction to MCL1 in KSHV-transformed PEL cell lines.

RESULTS

Primary effusion lymphoma cells are selectively addicted to MCL1

We previously identified host gene addictions of KSHV-transformed PEL cell lines using

genome-wide CRISPR screens (24). Our work showed that these tumor cells require high

expression of the oncogene MCL1 for survival. Although the BCL2 family proteins share similar

roles in preventing intrinsic apoptosis (Fig. 1B), none of the other BCL2 family genes scored in

our CRISPR screens (Fig. 1C). To exclude the possibility that the selective addiction to MCL1 is

due to the absence of the other BCL2 family proteins, we performed Western blotting for

endogenously expressed BCL2, BCL-xL, BCL-W, A1, Diva, or MCL1 proteins in a panel of PEL

cell lines (Fig. 1D). This panel included five cell lines that are co-infected with EBV (BC-1, BC-2,

BC-5, CRO-AP5, and JSC-1) and five that are only infected with KSHV (APK-1, BC-3, BCBL-1,

BCLM, VG-1). Our immunoblots show that PEL cell lines uniformly express BCL-xL, BCL-W,

and MCL1. In contrast, BCL2 expression is detectable only in BC-1 and BC-3, while we do not

detect A1 or Diva in any of the cell lines. Similar expression patterns of MCL1, BCL2, and BCL-

xL were recently reported in additional PEL cell lines (29). Together, our data indicate that PEL

cell lines express the other BCL2 family proteins, but are selectively addicted to MCL1.

BAX/BAK1 double knockout cell lines

We hypothesized that the selective addiction to MCL1 in PEL cell lines reflects an essential

function that is not redundant with the other BCL2 family proteins. Roles beyond inhibiting

apoptosis have emerged for MCL1, including facilitating mitochondrial respiration and fatty acid

oxidation (26-28). We hypothesized that MCL1 may have critical non-canonical functions in PEL

cells that necessitate MCL1 expression, but not the expression of other BCL2 proteins.

Loss-of-function studies on the non-canonical roles of MCL1 pose a technical hurdle because

these are confounded by effects from cell death. To address this issue, we derived clonal BAX

and BAK1 double knockout (DKO) cell lines from BC-1 (KSHV+EBV+) and BC-3 (KSHV+) cells

(Figs. 2, 3A, 3B). This inactivates the intrinsic apoptosis pathway since pore formation by BAX

or BAK is considered to be the necessary step to irreversibly initiate intrinsic apoptosis. We

nucleofected single-vector plasmids encoding Cas9 and guide RNAs (gRNAs) targeting the

coding region of *BAX* or *BAK1* (Fig. 2). To increase the efficiency of editing in all alleles, we co-

transfected two gRNA constructs for each gene and repeated nucleofection for a total of three

times. Single cell clones were isolated and screened for knockouts. Co-transfection of two

gRNAs allows for quick confirmation of editing by detecting ~100-400 bp deletions with

genotyping primers. Clones that did not express full length BAX and BAK1 proteins by Western

blots and had at least one large deletion in the BAX or BAK1 alleles by PCR were chosen for

final validation by Sanger or Illumina sequencing of the region spanning the predicted editing

sites (Figs. 2, 3A, 3B).

Finally, we tested whether the DKO mutations successfully inactivate the intrinsic apoptotic

pathway after treatment with staurosporine. Staurosporine induces intrinsic apoptosis by broadly

inhibiting protein kinases. In parallel, we also tested the sensitivities of the DKO cells to different

antibiotics. Dose response curves for these inhibitors show that DKO cell clones are

substantially resistant to treatment with staurosporine, but remain sensitive to the antibiotics

puromycin, blasticidin, zeocin, and hygromycin (Figs. 3C-G). Our data thus confirm that our

DKO cell clones successfully inactivate the intrinsic apoptotic pathway but not other cell death

mechanisms.

Knockout of BAX and BAK1 eliminates the dependency of PEL cells to MCL1

We used the BC-1 and BC-3 DKO cell clones to investigate their dependency on MCL1. If

inactivation of MCL1 continues to affect the viability of DKO cells, this would suggest that MCL1

has an essential role in PEL that is independent of intrinsic apoptosis. In contrast, if the DKO

cells are unaffected by MCL1 inhibition, then the main essential function of MCL1 is to inhibit

BAX/BAK1-mediated cell death.

We therefore treated BC-1 and BC-3 wild type (WT) and DKO cells with increasing doses of the

MCL1 inhibitor S63845 (MCL1i) (30). Consistent with our previous work (24), WT BC-1 and BC-

3 cells are sensitive to MCL1i with an estimated half maximal inhibitory concentration (IC50) of

50-150 nM (Fig. 4A). However, both DKO cell clones were resistant to MCL1i and viable even at

the maximum dose used (20 μM). Apoptotic cell death induced by the pharmacological inhibition

of MCL1 is thus prevented by ablating BAX and BAK1.

Since MCL1i was specifically designed to bind to the BH3-binding groove of MCL1 (30), it is

possible that MCL1i induces cell death by only inhibiting the anti-apoptotic role of MCL1 without

affecting putative non-canonical functions. We therefore targeted MCL1 for RNAi knockdown in

both BC-3 WT and DKO cells using a doxycycline (DOX)-inducible lentiviral shRNA. Total MCL1

protein was reduced by ~70% by quantitative Western blot in both WT and DKO cells using the

sh2MCL1 construct after Dox addition (Fig. 4B). This is not seen in the absence of DOX nor in a

negative control shRNA construct targeting the *Renilla* luciferase (31). Similar to our previous

work (24), MCL1 knockdown lead to a significant decrease in the total number of live cells in

BC-3 WT cells as early as two days after DOX treatment (Fig. 4C, dotted red line). In contrast,

MCL1 knockdown in BC-3 DKO cells did not substantially affect viability even after 12 days of

DOX treatment (Fig. 4C, dotted purple line). We note the efficiency of MCL1 knockdown is

sustained at this time point as evidenced by reduced MCL1 protein (Fig. 4B, bottom). Taken

together, our results from MCL1i treatment or shRNA knockdown of the DKO cells suggest that

the essential function of MCL1 in PEL cell lines is to prevent intrinsic apoptosis.

Over-expression of BCL2 proteins in PEL cell lines

Our results thus far indicate that the main role of MCL1 in PEL cell lines is to inhibit cell death.

However, this does not address why we observe a selective addiction to MCL1 in our CRISPR

screens despite co-expression of functionally similar BCL2 proteins. Because a minimum

number of pro-survival BCL2 family protein molecules is required to successfully neutralize

BAX/BAK1 monomers and prevent pore formation, we hypothesized that the expression ratios

of the BCL2 family directly influence the addiction to MCL1. Although we detected BCL-xL and

BCL-W proteins in all PEL cell lines and BCL2 in BC-1 and BC-3 (Fig. 1D), Western blotting

does not allow for relative quantification between different proteins even in the same sample.

We therefore reanalyzed publicly available mRNA-seq datasets from different PEL cell lines to

determine the abundance of the BCL2 family transcripts (19, 32, 33). As seen in Figure 5A,

endogenous MCL1 is the most highly expressed BCL2 family gene in all eight PEL cell lines

across the three independent datasets. In contrast, BCL2, BCL-xL, and BCL-W mRNA levels

are only modestly expressed (10- to 60-fold less compared to MCL1) while A1 and Diva mRNAs

are not detected. This expression profile reflects the degree of depletion of gRNAs from the

CRISPR screens (Fig. 1C) consistent with the model that expression ratios of the BCL2 family

may influence the selective addiction to MCL1.

If our hypothesis is true, we expect that changing the ratios of the BCL2 family genes would shift

the dependency to the most abundant BCL2 family protein and away from MCL1. We thus

transduced BC-1 or BC-3 cells with lentiviruses constitutively expressing the different BCL2

genes under the major immediate-early promoter and enhancer of the human cytomegalovirus

at a multiplicity of infection of 5. We chose to only over-express the proteins that are expressed

endogenously at the RNA and protein levels: BCL2, BCL-xL, BCL-W, or MCL1. In addition, we

also expressed the KSHV homolog of BCL2, vBCL2 (34, 35), that has been dually tagged with

FLAG-hemagglutinin (HA) epitopes at the N-terminus (N-vBCL2). Although vBCL2 is a lytic

gene and is not expressed in latently infected PEL cells, we were curious if this can functionally

substitute for MCL1 in PEL. Western blots confirm that each protein is expressed above

endogenous levels in both BC-1 and BC-3 backgrounds (Fig. 5B).

To confirm that the over-expressed proteins are functional in inhibiting intrinsic apoptosis, we

induced cell death in these cell lines using staurosporine and measured cellular viability using

CellTiterGlo 2.0 and apoptosis using Caspase 3/7 Luciferase Assay. All of the ectopically

expressed BCL2 family proteins and vBCL2 reduced basal caspase 3/7 activity in the absence

of staurosporine in both BC-1 and BC-3 cells (Figs. 5C, 5D). Reduction of caspase 3/7 activity

was consistently stronger in the BC-1 cell lines compared to the BC-3 cell lines. In both cases,

over-expression of BCL2, BCL-xL, or BCL-W inhibited basal caspase 3/7 activity the most while

MCL1 or N-vBCL2 had modest to no effects. It is possible that this minor effect of MCL1 or N-

vBCL2 in decreasing basal caspase 3/7 activity may be due to insufficient over-expression of

these proteins.

In the presence of staurosporine, naïve and mCherry-expressing cells had significantly reduced

cellular viability after overnight treatment as expected (Figs. 5E, 5F). Cell death was rescued in

all instances by all the BCL2 family proteins including N-vBCL2 although to a lesser extent

compared to the mCherry control. These effects in cell viability are conversely accompanied by

a reduction in caspase 3/7 activities (Figs. 5G, 5H). These experiments demonstrate that the

ectopically expressed BCL2 family proteins in BC-1 or BC-3 cells are functional in inhibiting

intrinsic apoptosis.

Over-expression of BCL2 family proteins desensitizes PEL cells to MCL1i-induced

apoptosis

We next tested whether over-expression of the BCL2-related proteins in BC-1 or BC-3 can shift

the dependency to a different BCL2 family protein and cause to cells to be less reliant to MCL1

for survival. We therefore inactivated MCL1 with MCL1i. Similar to staurosporine, overnight

treatment with MCL1i reduced the viability of naïve or mCherry-expressing cells (Figs. 6A, 6B).

In contrast, ectopic BCL2, BCL-xL, or BCL-W fully protected BC-1 and BC-3 cells against MCL1i

toxicity. N-vBCL2- and surprisingly, MCL1-expressing cells were sensitive to MCL1i with only a

marginal rescue of cell viability (20-30%) compared to mCherry. These effects in cell viability

are likely due to apoptosis as seen with the opposite pattern of caspase 3/7 activity in these cell

lines (Figs. 6C, 6D). Relative to naïve and mCherry-expressing cells, ectopic BCL2, BCL-xL, or

BCL-W inhibited caspase 3/7 activity induced by MCL1i to the same degree in both BC-1 and

BC-3. Ectopic N-vBCL2 moderately reduced caspase 3/7 activity while MCL1 had no

measurable effect.

Finally, we treated the cells with varying doses of MCL1i to calculate their relative sensitivities to

this inhibitor. Consistent with what we observed in Figs. 6A and 6B, over-expression of BCL2,

BCL-xL or BCL-W desensitized cells from MCL1i and increased the IC50 values by 100- to

1000-fold (Figs. 6E, 6F). The N-vBCL2 construct did not alter the over-all sensitivities of BC-1 or

BC-3 cells to MCL1i. While ectopic MCL1 increased the average MCL1i IC50 by ~2- (BC-3) to

3-fold (BC-1), these effects were not statistically significant. In sum, our data demonstrates that

expression levels of the BCL2 family genes is the major determinant for the dependency to

MCL1 by PEL cell lines.

DISCUSSION

In this report, we explore why KSHV-transformed PEL cell lines exhibit selective addiction to the oncogene MCL1, despite having shared functions with the related BCL2 family proteins. While new roles beyond those shared with the other BCL2 family members have been attributed to MCL1, we show that the essential function of MCL1 in PEL cell lines is to prevent intrinsic apoptosis. This selective addiction to MCL1 is explained by the 10- to 60-fold stoichiometric excess of MCL1 expression relative to other BCL2-related proteins as demonstrated by mRNA-seq and functional rescue from MCL1i-induced cell death by over-expression of any of the BCL2 family proteins.

The failure to rescue MCL1i-induced apoptosis by the MCL1 or N-vBCL2 constructs may be due to the degree of ectopic expression achieved in these cells. It is possible that the total amount of BCL2 family proteins in any of these cases did not reach the minimum required to neutralize all available MCL1i molecules.

It is not clear why BCL2 family expression is skewed towards MCL1 in PEL cell lines. This may be a consequence of the plasma cell-like B cell differentiation stage of PEL tumors (4). Indeed, the survival of long-lived plasma cells in mice is dependent on high levels of Mcl-1 expression (36). Similarly, the Cancer Dependency Map (37) shows that multiple myeloma cell lines, which are of plasma cell lineage, have high MCL1 expression and are selectively dependent on MCL1. The bias of high expression of MCL1 in the plasma cell compartment may be driven by master transcription factors such as IRF4 that occupy lineage-specific super-enhancer regulatory regions. Indeed, we and others have identified a super-enhancer element around the MCL1 locus bound by IRF4 (19, 38, 39).

Upon activation and during maturation, B cells undergo dramatic changes to support their effector functions in immune defense (Reviewed in 40). A large part of the plasma cell

differentiation program is the remodeling and expansion of the endoplasmic reticulum (ER) to

prepare for immunoglobulin production and secretion (41). Nutrient transporters and metabolic

enzymes are upregulated to sustain antibody production and organelle maintenance by the

lineage transcription factor Blimp-1 (42). High expression of MCL1 may thus act as a buffer

from this increased ER stress and metabolic load to ensure the survival of these B cells.

Knockdown of the glucose transporter GLUT4 in multiple myeloma cell lines induced apoptosis

by decreasing MCL1 mRNA expression (43), thereby supporting this link between metabolism

and apoptosis.

In the case of PEL, the infected tumor cells neither express nor secrete immunoglobulin

molecules. However, recent work from the Gottwein laboratory shows that PEL cell lines need

to constitutively counteract death signals originating from the ER and Golgi compartments in the

absence of any exogenous trigger (44). Consistent with this, PEL cell lines have elevated

expression of genes involved in the unfolded protein response, often triggered by ER stress (4).

These suggest that the ER and Golgi bodies in PEL cells are also under stress despite the

absence of antibody production. In line with this, KSHV is known to activate ER stress

responses during lytic reactivation (45, 46). Therefore, KSHV needs to counter these ER, Golgi

and metabolic stresses by possibly reinforcing or amplifying existing survival pathways in

plasma cells such as MCL1. More recently, the latency-associated nuclear antigen (LANA) was

shown to indirectly enhance the steady state levels of the MCL1 protein by binding to the E3

ubiquitin ligase FBW7 and preventing MCL1 degradation (47). As mentioned above, although

we find binding sites for the lineage transcription factor IRF4 on the MCL1 super-enhancer in

PEL cells, we recently reported that the B cell-specific latency gene vIRF3 largely co-occupies

similar sites as IRF4, including the MCL1 super-enhancer and promoter (19). This suggests that

vIRF3 may function to enhance MCL1 expression. We are currently investigating how vIRF3

regulates MCL1 expression at the transcriptional level.

A parallel relationship between lineage-specific expression and dependence to the BCL2 family proteins is also seen in lymphatic endothelial cells (LECs) which are thought to be candidates for the cellular origin of Kaposi's sarcoma spindle cells (48). In this case, LECs express and depend on high levels of BCL-xL instead of MCL1 for survival. This dependency dramatically increases during KSHV infection (Dr. Michael Lagunoff, personal communication) which supports the model where the virus reinforces pro-survival pathways already present in the cell.

The selective dependence of PEL cell lines to MCL1 makes it an attractive target for therapy. Candidate MCL1-selective drugs are now in various stages of clinical trials to treat hematologic malignancies (49) and have the potential to be repurposed to treat PEL. Venetoclax, the first approved BH3 mimetic drug, has over-all response rates of ~80% in relapsed and/or refractory chronic lymphocytic leukemia (50) and ~75% in relapsed and/or refractory mantle cell lymphoma (51) with a durable response even after ~3 years (52). However, other cancers like acute myelogenous leukemia exhibit a lower response rate to venetoclax (19%) where non-responsive patients had an increase in BCL-xL or MCL1 expression (53). Our data similarly shows that over-expression of any of the BCL2 family proteins or loss of BAX or BAK1 leads to resistance to MCL1 inhibitors. Thus, care must be taken when testing MCL1 inhibitors in PEL patients to avoid resistance. One strategy is using high but tolerable doses of MCL1 drugs to minimize selection for resistant tumor cells. Another approach is to combine this with other regimens that do not act on the same pathway. These strategies need to be explored to provide the most effective treatment for KSHV-induced PEL.

MATERIALS AND METHODS

Cells. Cells were cultured as described (19). HEK293T cells were cultured in Dulbecco's

modified Eagle medium with L-glutamine and 4.5 g/L glucose (Corning Life Sciences,

Tewksbury, MA) supplemented with 10% Serum Plus II medium (Millipore-Sigma, Burlington,

MA) and 10 µg/ml gentamicin (Gibco, Waltham, MA). BC-1 and BC-3 cell lines were grown in

RPMI 1640 (Corning) supplemented with 10 µg/ml gentamicin (Life Technologies, Carlsbad,

CA), 0.05 mM β-mercaptoethanol (Sigma-Aldrich, St. Louis, MO), and 10% (BC-1) or 20% (BC-

3) Serum Plus II medium. All PEL cell lines were grown at 37°C with 5% CO_2 and maintained at

log phase between $2x10^5$ and $1x10^6$ cells/mL.

Primers and synthetic genes. All primers were purchased from Integrated DNA Technologies

(Coralville, IO). Synthetic genes for eGFP fragment, shRNA cassettes and MCL1 cDNA were

synthesized by Twist Bioscience (South San Francisco, CA). Sequences and primer

combinations for cloning are listed in Tables 1 and 2.

shRNAs. A self-inactivating version of the pLVX-TetOne-Zeo lentiviral vector (54) (a gift from

Dr. Britt Glaunsinger) was first made via PCR mutagenesis. A deletion in the U3 region of the

3'LTR was introduced using primer pairs 169/171 and 172/170. The PCR products were fused

with primers 169/170 and cloned into pLVX-TetOne-Zeo using the KpnI and NheI restriction

sites (New England Biolabs, Ipswich, MA). The new vector is renamed pLVX-SIN-TetOne-Zeo

(pSTZ) to denote the self-inactivating deletion (SIN).

shRNAs were designed using the SplashRNA algorithm (31) using default parameters

(http://splashrna.mskcc.org). sh2MCL1 was synthesized as a gene fragment (Twist Bioscience)

while shREN was amplified from pZIP-ZsGreen-T2A-Hyg-shRen.713 (44). These shRNA

cassettes were assembled in the 3'UTR of a synthetic eGFP gene (Twist Bioscience) via two-

fragment Gibson assembly into the pSTZ backbone linearized by EcoRI and BamHI (NEB).

BCL2 cDNAs. cDNAs for human BCL2, BCL-xL, and BCL-W were amplified from pCDNA3-HA-

BCL2, pCDNA-HA-BCL-xL (55) (gifts from Dr. Timothy Chambers), and pLX304-BCL2L2 (Clone

HsCD00440394 from Human ORFeome Collection from DNASU (56)). vBCL2 sequence was

amplified from BC-1 genomic DNA. The Flag-HA dual tag cassette was amplified from pLC-

FLAG-HA-vIRF3-P2A-hyg (Manzano and Gottwein, unpublished). Internal restriction sites were

repaired using PCR mutagenesis. Details on primer sequences and primer combinations for

each PCR step are listed in Tables 1 and 2.

Lentiviruses. The day prior to transfections, $4\text{-}5\text{x}10^6$ 293T cells were plated in 10 cm dishes.

Lentiviral transfer vectors (5 μg) were co-transfected with pMD2.G (2.44 μg) and psPAX2 (3.74

μg) packaging vectors and 39.13 μL 1 mg/mL polyethylenimine (Sigma Catalog #408727) in

500 μL OptiMEM (Gibco). After 4-6 hours, media was replaced with complete RPMI medium

supplemented with 10% Serum Plus II medium. Supernatants containing lentiviruses were

harvested after three days by filtering through a 0.45 μm pore membrane.

Lentiviral titers were determined by functional titration. Increasing volumes of supernatant

containing lentiviruses were added to $1\text{x}10^5$ naïve BC-1 or BC-3 cells in 0.5 mL media with 0.5

μL 8 μg/mL polybrene. The day after, appropriate antibiotics were added to select for infected

cells. Number of surviving cells was compared to uninfected cells to determine infection rates

and viral titers. Final lentiviral transductions were performed in $4\text{x}10^5$ BC-1 or BC-3 cells in 2 mL

media with 8 μg/mL polybrene at defined MOIs.

Generation of BAX/BAK1 DKO cells. Two gRNAs targeting the human *BAX* or *BAK1* gene

were chosen from the genome-wide Brunello CRISPR library (57). Based on these gRNAs,

primer pairs were designed to create dsDNA duplexes with compatible BbsI overhangs. These

primer pairs were assembled by heat-denaturation and annealing in Buffer 3.1 (NEB), then

cloned into BbsI-linearized pX458 (58) (a gift from Dr. Feng Zhang, Addgene #48138).

BC-1 and BC-3 cells were adjusted to $2x10^5$ cells/mL the day before transfection. On the day of

transfection, $1x10^6$ BC-1 or BC-3 cells were pelleted in 50 mL tubes at 90g, for 10 min. Each

pellet was carefully resuspended in 20 μL of SF Cell Line Nucleofector Solution with

Supplement 1 (Lonza Cat. #V4XC-2032, Basel, Switzerland) and 1.5 μL of pX458 plasmids (150

ng each of BAX g2, BAX g3, BAK1 g1, BAK1 g3; 600 ng total). Cells transferred to

Nucleocuvette strips and pulsed with program CA-189 using the 4D-Nucleofector X (Lonza).

Transfection reactions were allowed to recover at room temperature for 10 min. Afterwards, 80

μL of pre-warmed complete media was added. Cells in the Nucleocuvette strips were incubated

at 37°C for 10 min. Finally, cells were collected with an additional 120 μL of pre-warmed

complete media and transferred to a round-bottom 96-well plate. Cells were expanded for 5-7

days to at least $2x10^6$ total cells. Two additional rounds of nucleofection were performed as

above to ensure efficient editing in all four alleles.

After a week of recovery from the third nucleofection, cells were individually cultured in round

bottom 96-well plate using dilution cloning. Clones were screened for successful editing as

outlined in Fig. 2. We first triaged the clones based on the absence of full length BAX and BAK1

proteins by Western blotting. Next, we further screened clones using genotyping primers (189

and 190 for *BAX*; 215 and 216 for *BAK1*). We would expect that simultaneous cutting of the two

guides in an allele will lead to ~100-400 bp deletions that can be distinguished from wt or single

cut alleles by PCR. While this assay does not discriminate between unedited from single editing

events, this allows us to limit our final validation to those with at least one allele deleted. Finally,

we confirmed frameshift indels by amplifying the targeted loci for Sanger or Illumina sequencing.

Illumina sequencing and analysis were performed by the Massachusetts General Hospital

Center for Computational and Integrative Biology DNA Core Facility.

Western blots. Cells were washed with phosphate buffered saline and lysed with RIPA lysis

buffer (50 mM Tris, 150 mM NaCl, 1% IGEPAL CA-630, 0.5% sodium deoxycholate, 0.1%

sodium dodecyl sulfate, pH 7.4) with protease inhibitor cocktail III (Calbiochem, EMD Millipore).

Lysates were incubated on ice for 10 minutes and clarified by spinning at 12,000 g for 10

minutes. Total protein was quantified the Pierce BCA Protein Assay Kit (Thermo Fisher).

Equivalent amounts of total proteins were run in Bolt 4%-12% Bis-Tris gradient gels (Thermo

Fisher Scientific) and transferred to 0.2 µm-pore size nitrocellulose membranes in Towbin buffer

(25 mM Tris, 192 mM glycine, 20% methanol, pH 8.3) for 90 min at 400 mA. Membranes were

blocked with a 1:3 dilution of Intercept (TBS) Blocking Buffer (Li-Cor Biosciences, Lincoln, NE)

in TBS for 1 h at room temperature. Membranes were incubated with primary antibodies diluted

1:3000 in blocking buffer–0.1% Tween 20 overnight at 4°C. Primary antibodies were purchased

from Cell Signaling Technology (Danvers, MA): anti-BCL2 clone D55G8, anti-BCL-xL clone

54H6, anti-BCL-W clone 31H4, anti-MCL1 clone D2W9E, anti-HA clone C29F4, anti-A1 clone

D1A1C, anti-Diva (Catalog # 3869), anti-BAK clone D4E4, and anti-BAX clone D2E11; or

Sigma: anti-beta-actin clone AC-15 and anti-alpha-tubulin clone B-5-1-2; or Santa Cruz

Biotechnology (Dallas, TX): anti-GAPDH clone 0411.

Immunoblots were washed five times with TBST for 5 min each then incubated for 1 hour with

IRDye 800 CW-conjugated secondary antibodies (Li-Cor Biosciences) diluted in TBS (1:20,000).

Membranes were finally washed four times with TBST and once in TBS. Western blots were

imaged on the Odyssey Fc or M Imager (Li-Cor Biosciences).

Dose Response Curves. Serial dilutions of the different inhibitors/drugs were added to each

well containing 5 x 10^3 BC-1 or BC-3 cells in a U-bottom 96-well plate. At designated time points

(2 days for puromycin (Cayman Chemicals, Ann Arbor, MI); 3 days for MCL1i

(MedChemExpress, Monmouth Junction, NJ) and staurosporine (APExBIO, Houston, TX); 6

days for blasticidin (Gibco), zeocin (Invivogen, San Diego, CA), and hygromycin (Gibco)), 20 µL

aliquots of the cells were used to measure cell viability using 20 µL CellTiter-Glo 2.0 Cell

Viability Reagent (Promega). Luminescence was read using Synergy H1 microplate reader

(Biotek, Winooski, VT) with an integration time of 2 sec. Raw luminescence readings were

normalized as a percentage of the DMSO control. Normalized values were used to calculate the

IC50 using the "[inhibitor] vs. normalized response (variable slope)" function in Prism 9

(GraphPad, San Diego, CA).

Caspase 3/7 Assay. 2×10^4 BC-1 or BC-3 cells were seeded in each well of a U-bottom 96-well

plate and treated with staurosporine (20 nM for BC-1, 100 nM for BC-3), 100 nM MCL1i, or

DMSO. These concentrations were chosen because they approximate the IC50 values of each

drug for each cell line. After 24 hours, 20 µL of cells were mixed with 20 µL of the CellTiter-Glo

2.0 Cell Viability Reagent or Caspase-Glo 3/7 Assay substrate (Promega). Luminescence were

measured after 2 min incubation for CellTiter-Glo 2.0 or 30 min incubation for Caspase-Glo 3/7

Assay using the Synergy H1 microplate reader as above. Raw luminescence values for caspase

3/7 activities were sequentially normalized to the matched raw CellTiter-Glo 2.0 readings, then

to the DMSO controls and finally, to the mCherry-expressing cell line.

ACKNOWLEDGMENTS

We thank Logan Moorhead for initial help with engineering the DKO cell lines, Dr. Eva Gottwein

for the pZIP-ZsGreen-T2A-Hyg-shRen.713 and pLC-FLAG-HA-vIRF3-P2A-hyg plasmids, Dr.

Britt Glaunsinger for the pLVX-TetOneZeo plasmid, and Dr. Timothy Chambers for the BCL-xL

and BCL2 expression vectors. We thank Dr. Eva Gottwein for helpful discussion and feedback

on the manuscript.

This work is supported by a Transition Career Development Award K22 CA241355 from the

National Cancer Institute of the National Institutes of Health (NIH) to M.M. Additional support

was provided by the Center for Microbial Pathogenesis and Host Inflammatory Responses

program grant P20 GM103625 from the National Institute of General Medical Sciences (NIGMS)

of the NIH, Winthrop P. Rockefeller Cancer Institute at UAMS, and an equipment grant from the

UAMS Office of the Vice Chancellor for Research and Innovation. J.G. is a recipient of the

Undergraduate Summer Research Fellowship from the Arkansas IDeA Network of Biomedical

Research Excellence (INBRE) funded by program grant P20 GM103429 from NIH/NIGMS. The

content is solely the responsibility of the authors and does not necessarily represent the official

views of the funding bodies. The authors declare no conflicts of interests.

FIGURE LEGENDS

Figure 1. *A*. Genome-wide CRISPR screens identify MCL1 as a top gene dependency in PEL cell lines (24). Genes are classified using the DepMap criteria in distinguishing genes that are "Strongly Selective" in a cell line from "Common Essential." ***B*.** The BCL2 family proteins (including MCL1) block intrinsic apoptosis by sequestering activated BAX or BAK1 monomers on the outer membrane of the mitochondria. During intrinsic apoptosis, pro-apoptotic BH3-only proteins inactivate the BCL2 family proteins and release the BAX/BAK1 monomers. Free monomers assemble to create mitochondrial pores and release cytochrome c to initiate the apoptotic cascade in the cytoplasm. ***C*.** Only gRNAs targeting MCL1 among the BCL2 family are significantly depleted in all PEL cell lines that were previously screened (24). ***D*.** Western blots of the different BCL2 family proteins in a panel of PEL cell lines that are infected with KSHV only or co-infected by KSHV and EBV. A1 and Diva proteins are not detected.

Figure 2. Experimental flow chart that summarizes the derivation of the BAX/BAK1 DKO clones. BC-1 or BC-3 cells were nucleofected three times with four plasmids encoding Cas9 and different gRNAs for either BAX or BAK1. Isolated clones were screened for the absence of full length BAX or BAK1 proteins, presence of 100-400 bp deletion by PCR to confirm dual editing, and sequencing of PCR-amplified BAX or BAK1 locus. Two DKO clones were confirmed for both BC-1 and BC-3 backgrounds and are summarized below. Also shown are representative Western blots for BAX and BAK1 as well as agarose gel electrophoresis images for PCR genotyping to confirm dual editing events.

Figure 3. *A*. Western blots for BAX, BAK1 and tubulin proteins of WT or BAX/BAK1 double knockout (DKO) clone 28 (BC-1) and clone 12 (BC-3). ***B*.** Summary of CRISPR edits in BC-1 DKO clone #28 and BC-3 DKO clone #12. ***C-G*.** Dose response curves of BC-1 or BC-3 cells with BAX/BAK1 wildtype or DKO background upon exposure to different inducers of cell death:

C. staurosporine, *D.* puromycin, *E.* blasticidin, *F.* hygromycin, or *G*. zeocin. Statistical

differences were calculated using two-way ANOVA with Tukey's multiple comparison post-hoc

test (n=3). Adjusted *p* values reflect post-hoc comparisons between the responses of the wt cell

line compared to the DKO cell line at the specific time point. Error bars, standard error of mean.

n.s., not significant.

Figure 4. *A*. MCL1i dose response curves of BC-1 or BC-3 cells with BAX/BAK1 wildtype or

DKO background. ***B****.* Western blots for total MCL1 in BC-3 BAX/BAK wt or DKO #12 cells

transduced with lentiviruses expressing a doxycycline (DOX)-inducible shRNA against Renilla

luciferase or MCL1. Shown are samples taken after 2, 3, or 12 days of DOX-treatment (1

µg/mL). ***C****.* Cumulative live cell count of BC-3 cells upon Dox treatment (dotted lines). Cell

counts were terminated for DOX-treated BC-3 wt cells with sh2MCL1 (dotted red line) at day 9

of Dox-treatment due to significant amount of cell death. Statistical differences were calculated

using one- or two-way ANOVA with Tukey's multiple comparison post-hoc test (n=3). One-way

ANOVA for each time point was instead done for *C*. since two-way ANOVA requires matched

data for all time points. Adjusted *p* values reflect post-hoc comparisons between the responses

of the wt cell line compared to the DKO cell line at the specific time point (in *B*.) or Dox-treated

cell line compared to matched untreated cell line (in *C.*) (n=3). Error bars, standard error of

mean. n.s., not significant.

Figure 5. *A*. Normalized expression values of BCL2 family transcripts in PEL cell lines from

published RNA-seq datasets (19, 32, 33). ***B.*** Western blots of the BCL2 family members and

HA-tagged vBCL2 in BC-1 or BC-3 cells transduced with lentiviruses over-expressing these

cDNAs. ***C-D.*** Basal caspase 3/7 activity of BC-1 (n=11) or BC-3 cells (n=7) normalized to total

live cells measured by CellTiterGlo 2.0 assay. ***E-F.*** Cell viability measured by CellTiterGlo 2.0

assay or ***G-H****.* relative caspase 3/7 activity in live BC-1 (n=3) or BC-3 cells (n=4) after overnight

treatment of 20 nM or 100 nM staurosporine. Statistical differences were calculated using one-

or two-way ANOVA with Tukey's multiple comparison post-hoc test. Adjusted *p* values reflect

post-hoc comparisons between the cell lines in response to staurosporine or DMSO unless

indicated otherwise. Error bars, standard error of mean. n.s., not significant.

Figure 6. *A-B.* Cell viability measured by CellTiterGlo 2.0 assay or *C-D.* relative caspase 3/7

activity in live BC-1 or BC-3 cells over-expressing BCL2 family cDNAs after overnight treatment

of 100 nM MCL1i. Statistical differences were calculated using one- or two-way ANOVA with

Tukey's multiple comparison post-hoc test (n=5). Adjusted *p* values reflect post-hoc

comparisons between the cell lines in response to staurosporine or DMSO unless indicated

otherwise. *E-F.* IC50 values of MCL1i in BC-1 or BC-3 cells. For BC-1, one-way ANOVA with

Tukey's multiple comparison post-hoc test was used for statistical tests (n≥3). Since BC-3 cell

lines over-expressing BCL2, BCL-xL or BCL-W were highly resistant to MCL1i even at high

doses, IC50 values derived from these incomplete sigmoidal dose-response curves had low

confidence intervals. Thus, non-parametric Kruskal-Wallis post-test was used. Adjusted *p*

values reflect post-hoc comparisons between the cell lines to the mCherry control. Error bars,

standard error of mean. n.s., not significant.

1   **TABLES**

2   **Table 1. Primer sequences.**

5	CACCGACGGCAGCTCGCCATCATCG
6	AAACCGATGATGGCGAGCTGCCGTC
50	GCTAGCGCTACCGGTGCCGCCACCATGGCGCACGCTGGGAGAACAGGG
51	GGGAGAGGGGCGGATCCTCACTTGTGGCCCAGATAGGC
54	GCTAGCGCTACCGGTGCCGCCACCATGTCTCAGAGCAACCGGGAGCTGG
55	GGGAGAGGGGCGGATCCTCATTTCCGACTGAAGAGTGAGCCC
56	CCACCTAGAGCCTTGGATTCAGGAGAACGGCGGCTGG
57	CCAGCCGCCGTTCTCCTGAATCCAAGGCTCTAGGTGG
62	GCTAGCGCTACCGGTGCCGCCACCATGGCGACCCCAGCCTCGGCCCCAG
63	GGGAGAGGGGCGGATCCTCACTTGCTAGCAAAAAAGGCCCCTAC
64	GCAGCTGGCTGACTGGATTCACAGCAGTGGGGGCTGG
65	CCAGCCCCCACTGCTGTGAATCCAGTCAGCCAGCTGC
66	CCCGATTACGCTAGCGGCGACGAGGACGTTTTGCCT
67	AGGCAAAACGTCCTCGTCGCCGCTAGCGTAATCGGG
68	GAGATTTCACAGCACCACTGGTATAAACCAGCTTGGG
69	CCCAAGCTGGTTTATACCAGTGGTGCTGTGAAATCTC
70	GGGAGAGGGGCGGATCCTTATCTCCTGCTCATCGCGACCGC
97	CACCGTTTCATCCAGGATCGAGCA
98	AAACTGCTCGATCCTGGATGAAAC
99	CACCGGACGAACTGGACAGTAACA
100	AAACTGTTACTGTCCAGTTCGTCC
103F	CACCGCTCACCTGCTAGGTTGCAG
103R	AAACCTGCAACCTAGCAGGTGAGC
161	GACACCGGTGCAGCGGCCGCTGTTTGAATGAGGCTTCAGTAC
162	GTCCAGCTTAGCTCGCAGGGGAGGTGGTCTGGATCCTAAAGTGATTTAATTTATACCATT
169	CTACCAATGCTGATTGTGCCTGGC
170	AACAGCTATGACCATGATTACGCC
171	CTAACCAGAGAGACCCAGTACAGTCCGGATGCAGCTCTCGGGCCAC
172	GTGGCCCGAGAGCTGCATCCGGACTGTACTGGGTCTCTCTGGTTAG
189	GAGTTTGAGGTCTGATTATTGTGG
190	GAAGTTGCCGTCAGAAAACATGTC
215	CTCCCTGACTCCCAGCTTTG
216	CCTGGTGGCAATCTTGGTGA
EHG4087	GAGCTGGTTTAGTGAACCG
eGFP cassette	ACTTTCCGTACCACTTCCTACCCTCGTAAAGAATTCATGGTGAGCAAGGGCGAGGAGCTGTTCACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGCCCACCCTCGTGACCACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACTACAACAGCCACAACGTCTATATCAT

	GGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCAAGATCCGCCACA ACATCGAGGACGGCAGCGTGCAGCTCGCCGACCACTACCAGCAGAACAC CCCCATCGGCGACGGCCCCGTGCTGCTGCCCGACAACCACTACCTGAGC ACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGT CCTGCTGGAGTTCGTGACCGCCGCCGGGATCACTCTCGGCATGGACGAG CTGTACAAGTAGGACACCGGTG CAGCGGCCGCTGTTTGAATGAGGCTTCAGTACTTTACAGA
sh2MCL1 cassette	TGTTTGAATGAGGCTTCAGTACTTTACAGAATCGTTGCCTGCACATCTTGG AAACACTTGCTGGGATTACTTCTTCAGGTTAACCCAACAGAAGGCTAAAGA AGGTATATTGCTGTTGACAGTGAGCGCTCCCAGTATACTTCTTAGAAATAG TGAAGCCACAGATGTATTTCTAAGAAGTATACTGGGAATGCCTACTGCCTC GGACTTCAAGGGGCTACTTTAGGAGCAATTATCTTGTTTACTAAAACTGAAT ACCTTGCTATCTCTTTGATACATTTTTACAAAGCTGAATTAAAATGGTATAAA TTAAATCACTTTAGGATCCAGACCACCTCCCCTGCGAGCTAAGCTGGAC

Table 2. Combinations of primers used for cloning constructs.

Construct	PCR	Primers	Template	Source
pLV-SIN-TetOne-ZeoR (pSTZ)	1	169, 171	pLVX-TetOne-Zeo	(54)
	2	170, 172		
	Overlap	169, 170	PCR 1 and 2	
shREN	N/A	161, 162,	pZIP-ZsGreen-T2A-Hyg-shRen.713	(44)
BCL2	N/A	50, 51	pCDNA3 HA-BCL2	(55)
BCL-xL	1	54, 57	pcDNA3 HA-BCL-xL	
	2	55, 56		
	Overlap	54, 55	PCR 1 and 2	
BCL-W	1	62, 65	pLX304 BCL2L2	DNASU
	2	63, 64		
	Overlap	62, 63	PCR 1 and 2	
vBCL2 N-FLAG-HA	1	EHG4087, 67	pLC-FLAG-HA-vIRF3-P2A-hyg	Manzano & Gottwein, unpublished
	2	66, 69	BC-1 genomic DNA	
	3	68, 70	BC-1 genomic DNA	
	1+2	EHG4087, 69	PCR 1 and 2	
	Final	EHG4087, 70	PCR 1+2 and 3	
BAX g2	Anneal	97, 98		
BAX g3	Anneal	99, 100		
BAK1 g1	Anneal	5, 6		
BAK1 g2	Anneal	103F, 103R		

REFERENCES

1. Cesarman E. 2014. Gammaherpesviruses and lymphoproliferative disorders. Annu Rev Pathol 9:349-72.
2. Nador RG, Cesarman E, Chadburn A, Dawson DB, Ansari MQ, Sald J, Knowles DM. 1996. Primary effusion lymphoma: a distinct clinicopathologic entity associated with the Kaposi's sarcoma-associated herpes virus. Blood 88:645-56.
3. Lurain K, Polizzotto MN, Aleman K, Bhutani M, Wyvill KM, Goncalves PH, Ramaswami R, Marshall VA, Miley W, Steinberg SM, Little RF, Wilson W, Filie AC, Pittaluga S, Jaffe ES, Whitby D, Yarchoan R, Uldrick TS. 2019. Viral, immunologic, and clinical features of primary effusion lymphoma. Blood 133:1753-1761.
4. Jenner RG, Maillard K, Cattini N, Weiss RA, Boshoff C, Wooster R, Kellam P. 2003. Kaposi's sarcoma-associated herpesvirus-infected primary effusion lymphoma has a plasma cell gene expression profile. Proc Natl Acad Sci U S A 100:10399-404.
5. Carbone A, Gloghini A, Cozzi MR, Capello D, Steffan A, Monini P, De Marco L, Gaidano G. 2000. Expression of MUM1/IRF4 selectively clusters with primary effusion lymphoma among lymphomatous effusions: implications for disease histogenesis and pathogenesis. Br J Haematol 111:247-57.
6. Petre CE, Sin SH, Dittmer DP. 2007. Functional p53 signaling in Kaposi's sarcoma-associated herpesvirus lymphomas: implications for therapy. J Virol 81:1912-22.
7. Bubman D, Guasparri I, Cesarman E. 2007. Deregulation of c-Myc in primary effusion lymphoma by Kaposi's sarcoma herpesvirus latency-associated nuclear antigen. Oncogene 26:4979-86.
8. Gaidano G, Capello D, Cilia AM, Gloghini A, Perin T, Quattrone S, Migliazza A, Lo Coco F, Saglio G, Ascoli V, Carbone A. 1999. Genetic characterization of HHV-8/KSHV-positive primary effusion lymphoma reveals frequent mutations of BCL6: implications for disease pathogenesis and histogenesis. Genes Chromosomes Cancer 24:16-23.
9. Roy D, Sin SH, Damania B, Dittmer DP. 2011. Tumor suppressor genes FHIT and WWOX are deleted in primary effusion lymphoma (PEL) cell lines. Blood 118:e32-9.
10. Godfrey A, Anderson J, Papanastasiou A, Takeuchi Y, Boshoff C. 2005. Inhibiting primary effusion lymphoma by lentiviral vectors encoding short hairpin RNA. Blood 105:2510-8.
11. Wies E, Mori Y, Hahn A, Kremmer E, Sturzl M, Fleckenstein B, Neipel F. 2008. The viral interferon-regulatory factor-3 is required for the survival of KSHV-infected primary effusion lymphoma cells. Blood 111:320-7.
12. Xiang Q, Ju H, Nicholas J. 2020. USP7-Dependent Regulation of TRAF Activation and Signaling by a Viral Interferon Regulatory Factor Homologue. J Virol 94.
13. Xiang Q, Ju H, Li Q, Mei SC, Chen D, Choi YB, Nicholas J. 2018. Human Herpesvirus 8 Interferon Regulatory Factors 1 and 3 Mediate Replication and Latency Activities via Interactions with USP7 Deubiquitinase. J Virol 92.
14. Chen D, Sandford G, Nicholas J. 2009. Intracellular signaling mechanisms and activities of human herpesvirus 8 interleukin-6. J Virol 83:722-33.
15. Bhatt AP, Bhende PM, Sin SH, Roy D, Dittmer DP, Damania B. 2010. Dual inhibition of PI3K and mTOR inhibits autocrine and paracrine proliferative loops in PI3K/Akt/mTOR-addicted lymphomas. Blood 115:4455-63.
16. Cousins E, Nicholas J. 2013. Role of human herpesvirus 8 interleukin-6-activated gp130 signal transducer in primary effusion lymphoma cell growth and viability. J Virol 87:10816-27.
17. de Oliveira DE, Ballon G, Cesarman E. 2010. NF-kappaB signaling modulation by EBV and KSHV. Trends Microbiol 18:248-57.

18. Gopalakrishnan R, Matta H, Tolani B, Triche T, Jr., Chaudhary PM. 2016. Immunomodulatory drugs target IKZF1-IRF4-MYC axis in primary effusion lymphoma in a cereblon-dependent manner and display synergistic cytotoxicity with BRD4 inhibitors. Oncogene 35:1797-810.

19. Manzano M, Gunther T, Ju H, Nicholas J, Bartom ET, Grundhoff A, Gottwein E. 2020. Kaposi's Sarcoma-Associated Herpesvirus Drives a Super-Enhancer-Mediated Survival Gene Expression Program in Primary Effusion Lymphoma. mBio 11.

20. Patil A, Manzano M, Gottwein E. 2018. CK1alpha and IRF4 are essential and independent effectors of immunomodulatory drugs in primary effusion lymphoma. Blood 132:577-586.

21. Sarek G, Kurki S, Enback J, Iotzova G, Haas J, Laakkonen P, Laiho M, Ojala PM. 2007. Reactivation of the p53 pathway as a treatment modality for KSHV-induced lymphomas. J Clin Invest 117:1019-28.

22. Sin SH, Roy D, Wang L, Staudt MR, Fakhari FD, Patel DD, Henry D, Harrington WJ, Jr., Damania BA, Dittmer DP. 2007. Rapamycin is efficacious against primary effusion lymphoma (PEL) cell lines in vivo by inhibiting autocrine signaling. Blood 109:2165-73.

23. Keller SA, Schattner EJ, Cesarman E. 2000. Inhibition of NF-kappa B induces apoptosis of KSHV-infected primary effusion lymphoma cells. Blood 96:2537-2542.

24. Manzano M, Patil A, Waldrop A, Dave SS, Behdad A, Gottwein E. 2018. Gene essentiality landscape and druggable oncogenic dependencies in herpesviral primary effusion lymphoma. Nat Commun 9:3263.

25. Westphal D, Kluck RM, Dewson G. 2014. Building blocks of the apoptotic pore: how Bax and Bak are activated and oligomerize during apoptosis. Cell Death Differ 21:196-205.

26. Escudero S, Zaganjor E, Lee S, Mill CP, Morgan AM, Crawford EB, Chen JH, Wales TE, Mourtada R, Luccarelli J, Bird GH, Steidl U, Engen JR, Haigis MC, Opferman JT, Walensky LD. 2018. Dynamic Regulation of Long-Chain Fatty Acid Oxidation by a Noncanonical Interaction between the MCL-1 BH3 Helix and VLCAD. Molecular Cell 69:729-+.

27. Perciavalle RM, Stewart DP, Koss B, Lynch J, Milasta S, Bathina M, Temirov J, Cleland MM, Pelletier S, Schuetz JD, Youle RJ, Green DR, Opferman JT. 2012. Anti-apoptotic MCL-1 localizes to the mitochondrial matrix and couples mitochondrial fusion to respiration. Nat Cell Biol 14:575-83.

28. Germain M, Nguyen AP, Le Grand JN, Arbour N, Vanderluit JL, Park DS, Opferman JT, Slack RS. 2011. MCL-1 is a stress sensor that regulates autophagy in a developmentally regulated manner. Embo Journal 30:395-407.

29. Quentmeier H, Geffers R, Hauer V, Nagel S, Pommerenke C, Uphoff CC, Zaborski M, Drexler HG. 2022. Inhibition of MCL1 induces apoptosis in anaplastic large cell lymphoma and in primary effusion lymphoma. Sci Rep 12:1085.

30. Kotschy A, Szlavik Z, Murray J, Davidson J, Maragno AL, Le Toumelin-Braizat G, Chanrion M, Kelly GL, Gong JN, Moujalled DM, Bruno A, Csekei M, Paczal A, Szabo ZB, Sipos S, Radics G, Proszenyak A, Balint B, Ondi L, Blasko G, Robertson A, Surgenor A, Dokurno P, Chen I, Matassova N, Smith J, Pedder C, Graham C, Studeny A, Lysiak-Auvity G, Girard AM, Grave F, Segal D, Riffkin CD, Pomilio G, Galbraith LC, Aubrey BJ, Brennan MS, Herold MJ, Chang C, Guasconi G, Cauquil N, Melchiore F, Guigal-Stephan N, Lockhart B, Colland F, Hickman JA, Roberts AW, Huang DC, Wei AH, et al. 2016. The MCL1 inhibitor S63845 is tolerable and effective in diverse cancer models. Nature 538:477-482.

31. Pelossof R, Fairchild L, Huang CH, Widmer C, Sreedharan VT, Sinha N, Lai DY, Guan Y, Premsrirut PK, Tschaharganeh DF, Hoffmann T, Thapar V, Xiang Q, Garippa RJ, Ratsch G, Zuber J, Lowe SW, Leslie CS, Fellmann C. 2017. Prediction of potent shRNAs with a sequential classification algorithm. Nat Biotechnol 35:350-353.

32. Zhou F, Shimoda M, Olney L, Lyu Y, Tran K, Jiang G, Nakano K, Davis RR, Tepper CG, Maverakis E, Campbell M, Li Y, Dandekar S, Izumiya Y. 2017. Oncolytic Reactivation of KSHV as a Therapeutic Approach for Primary Effusion Lymphoma. Mol Cancer Ther 16:2627-2638.

33. van der Meer D, Barthorpe S, Yang W, Lightfoot H, Hall C, Gilbert J, Francies HE, Garnett MJ. 2019. Cell Model Passports-a hub for clinical, genetic and functional datasets of preclinical cancer models. Nucleic Acids Res 47:D923-D929.

34. Sarid R, Sato T, Bohenzky RA, Russo JJ, Chang Y. 1997. Kaposi's sarcoma-associated herpesvirus encodes a functional bcl-2 homologue. Nat Med 3:293-8.

35. Cheng EH, Nicholas J, Bellows DS, Hayward GS, Guo HG, Reitz MS, Hardwick JM. 1997. A Bcl-2 homolog encoded by Kaposi sarcoma-associated virus, human herpesvirus 8, inhibits apoptosis but does not heterodimerize with Bax or Bak. Proc Natl Acad Sci U S A 94:690-4.

36. Peperzak V, Vikstrom I, Walker J, Glaser SP, LePage M, Coquery CM, Erickson LD, Fairfax K, Mackay F, Strasser A, Nutt SL, Tarlinton DM. 2013. Mcl-1 is essential for the survival of plasma cells. Nat Immunol 14:290-7.

37. Tsherniak A, Vazquez F, Montgomery PG, Weir BA, Kryukov G, Cowley GS, Gill S, Harrington WF, Pantel S, Krill-Burger JM, Meyers RM, Ali L, Goodale A, Lee Y, Jiang G, Hsiao J, Gerath WFJ, Howell S, Merkel E, Ghandi M, Garraway LA, Root DE, Golub TR, Boehm JS, Hahn WC. 2017. Defining a Cancer Dependency Map. Cell 170:564-576 e16.

38. Wang C, Zhang L, Ke L, Ding W, Jiang S, Li D, Narita Y, Hou I, Liang J, Li S, Xiao H, Gottwein E, Kaye KM, Teng M, Zhao B. 2020. Primary effusion lymphoma enhancer connectome links super-enhancers to dependency factors. Nat Commun 11:6318.

39. Park A, Oh S, Jung KL, Choi UY, Lee HR, Rosenfeld MG, Jung JU. 2020. Global epigenomic analysis of KSHV-infected primary effusion lymphoma identifies functional MYC superenhancers and enhancer RNAs. Proc Natl Acad Sci U S A 117:21618-21627.

40. Lam WY, Bhattacharya D. 2018. Metabolic Links between Plasma Cell Survival, Secretion, and Stress. Trends Immunol 39:19-27.

41. Wiest DL, Burkhardt JK, Hester S, Hortsch M, Meyer DI, Argon Y. 1990. Membrane biogenesis during B cell differentiation: most endoplasmic reticulum proteins are expressed coordinately. J Cell Biol 110:1501-11.

42. Tellier J, Shi W, Minnich M, Liao Y, Crawford S, Smyth GK, Kallies A, Busslinger M, Nutt SL. 2016. Blimp-1 controls plasma cell function through the regulation of immunoglobulin secretion and the unfolded protein response. Nature Immunology 17:323-330.

43. McBrayer SK, Cheng JC, Singhal S, Krett NL, Rosen ST, Shanmugam M. 2012. Multiple myeloma exhibits novel dependence on GLUT4, GLUT8, and GLUT11: implications for glucose transporter-directed therapy. Blood 119:4686-97.

44. Kuehnle N, Manzano M, Gottwein E. 2022. Cellular FLIP Protects KSHV-Transformed Lymphoma Cells from TRAIL-Independent TRAIL Receptor 1-Mediated Cell Death. bioRxiv doi:https://doi.org/10.1101/2022.08.17.504167.

45. De Leo A, Chen HS, Hu CC, Lieberman PM. 2017. Deregulation of KSHV latency conformation by ER-stress and caspase-dependent RAD21-cleavage. PLoS Pathog 13:e1006596.

46. Johnston BP, Pringle ES, McCormick C. 2019. KSHV activates unfolded protein response sensors but suppresses downstream transcriptional responses to support lytic replication. PLoS Pathog 15:e1008185.

47. Kim YJ, Kim Y, Kumar A, Kim CW, Toth Z, Cho NH, Lee HR. 2021. Kaposi's sarcoma-associated herpesvirus latency-associated nuclear antigen dysregulates expression of MCL-1 by targeting FBW7. PLoS Pathog 17:e1009179.

48. Wang HW, Trotter MW, Lagos D, Bourboulia D, Henderson S, Makinen T, Elliman S, Flanagan AM, Alitalo K, Boshoff C. 2004. Kaposi sarcoma herpesvirus-induced cellular reprogramming contributes to the lymphatic endothelial gene expression in Kaposi sarcoma. Nat Genet 36:687-93.

49. Diepstraten ST, Anderson MA, Czabotar PE, Lessene G, Strasser A, Kelly GL. 2022. The manipulation of apoptosis for cancer therapy using BH3-mimetic drugs. Nat Rev Cancer 22:45-64.

50. Roberts AW, Davids MS, Pagel JM, Kahl BS, Puvvada SD, Gerecitano JF, Kipps TJ, Anderson MA, Brown JR, Gressick L, Wong S, Dunbar M, Zhu M, Desai MB, Cerri E, Heitner Enschede S, Humerickhouse RA, Wierda WG, Seymour JF. 2016. Targeting BCL2 with Venetoclax in Relapsed Chronic Lymphocytic Leukemia. N Engl J Med 374:311-22.

51. Davids MS, Roberts AW, Seymour JF, Pagel JM, Kahl BS, Wierda WG, Puvvada S, Kipps TJ, Anderson MA, Salem AH, Dunbar M, Zhu M, Peale F, Ross JA, Gressick L, Desai M, Kim SY, Verdugo M, Humerickhouse RA, Gordon GB, Gerecitano JF. 2017. Phase I First-in-Human Study of Venetoclax in Patients With Relapsed or Refractory Non-Hodgkin Lymphoma. J Clin Oncol 35:826-833.

52. Davids MS, Roberts AW, Kenkre VP, Wierda WG, Kumar A, Kipps TJ, Boyer M, Salem AH, Pesko JC, Arzt JA, Mantas M, Kim SY, Seymour JF. 2021. Long-term Follow-up of Patients with Relapsed or Refractory Non-Hodgkin Lymphoma Treated with Venetoclax in a Phase I, First-in-Human Study. Clin Cancer Res 27:4690-4695.

53. Konopleva M, Pollyea DA, Potluri J, Chyla B, Hogdal L, Busman T, McKeegan E, Salem AH, Zhu M, Ricker JL, Blum W, DiNardo CD, Kadia T, Dunbar M, Kirby R, Falotico N, Leverson J, Humerickhouse R, Mabry M, Stone R, Kantarjian H, Letai A. 2016. Efficacy and Biological Correlates of Response in a Phase II Study of Venetoclax Monotherapy in Patients with Acute Myelogenous Leukemia. Cancer Discov 6:1106-1117.

54. Castaneda AF, Glaunsinger BA. 2019. The Interaction between ORF18 and ORF30 Is Required for Late Gene Expression in Kaposi's Sarcoma-Associated Herpesvirus. J Virol 93.

55. Upreti M, Chu R, Galitovskaya E, Smart SK, Chambers TC. 2008. Key role for Bak activation and Bak-Bax interaction in the apoptotic response to vinblastine. Mol Cancer Ther 7:2224-32.

56. Seiler CY, Park JG, Sharma A, Hunter P, Surapaneni P, Sedillo C, Field J, Algar R, Price A, Steel J, Throop A, Fiacco M, LaBaer J. 2014. DNASU plasmid and PSI:Biology-Materials repositories: resources to accelerate biological research. Nucleic Acids Res 42:D1253-60.

57. Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, Smith I, Tothova Z, Wilen C, Orchard R, Virgin HW, Listgarten J, Root DE. 2016. Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat Biotechnol 34:184-191.

58. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. 2013. Genome engineering using the CRISPR-Cas9 system. Nat Protoc 8:2281-2308.

Figure 1. **A**. Genome-wide CRISPR screens identify MCL1 as a top gene dependency in PEL cell lines (24). Genes are classified using the DepMap criteria in distinguishing genes that are "Strongly Selective" in a cell line from "Common Essential." **B**. The BCL2 family proteins (including MCL1) block intrinsic apoptosis by sequestering activated BAX or BAK1 monomers on the outer membrane of the mitochondria. During intrinsic apoptosis, pro-apoptotic BH3-only proteins inactivate the BCL2 family proteins and release the BAX/BAK1 monomers. Free monomers assemble to create mitochondrial pores and release cytochrome c to initiate the apoptotic cascade in the cytoplasm. **C**. Only gRNAs targeting MCL1 among the BCL2 family are significantly depleted in all PEL cell lines that were previously screened (24). **D**. Western blots of the different BCL2 family proteins in a panel of PEL cell lines that are infected with KSHV only or co-infected by KSHV and EBV. A1 and Diva proteins are not detected.

Fig. 2. Experimental flow chart that summarizes the derivation of the BAX/BAK1 DKO clones. BC-1 or BC-3 cells were nucleofected three times with four plasmids encoding Cas9 and different gRNAs for either BAX or BAK1. Isolated clones were screened for the absence of full length BAX or BAK1 proteins, presence of 100-400 bp deletion by PCR to confirm dual editing, and sequencing of PCR-amplified BAX or BAK1 locus. Two DKO clones were confirmed for both BC-1 and BC-3 backgrounds and are summarized below. Also shown are representative Western blots for BAX and BAK1 as well as agarose gel electrophoresis images for PCR genotyping to confirm dual editing events.

Figure 3. A. Western blots for BAX, BAK1 and tubulin proteins of WT or BAX/BAK1 double knockout (DKO) clone 28 (BC-1) and clone 12 (BC-3). **B.** Summary of CRISPR edits in BC-1 DKO clone #28 and BC-3 DKO clone #12. **C-G**. Dose response curves of BC-1 or BC-3 cells with BAX/BAK1 wildtype or DKO background upon exposure to different inducers of cell death: C. staurosporine, D. puromycin, E. blasticidin, F. hygromycin, or G. zeocin. Statistical differences were calculated using two-way ANOVA with Tukey's multiple comparison post-hoc test (n=3). Adjusted p values reflect post-hoc comparisons between the responses of the wt cell line compared to the DKO cell line at the specific time point. Error bars, standard error of mean. n.s., not significant.

Figure 4. A. MCL1i dose response curves of BC-1 or BC-3 cells with BAX/BAK1 wildtype or DKO background. **B.** Western blots for total MCL1 in BC-3 BAX/BAK wt or DKO #12 cells transduced with lentiviruses expressing a doxycycline (DOX)-inducible shRNA against Renilla luciferase or MCL1. Shown are samples taken after 2, 3, or 12 days of DOX-treatment (1 μ g/mL). **C.** Cumulative live cell count of BC-3 cells upon Dox treatment (dotted lines). Cell counts were terminated for DOX-treated BC-3 wt cells with sh2MCL1 (dotted red line) at day 9 of Dox-treatment due to significant amount of cell death. Statistical differences were calculated using one- or two-way ANOVA with Tukey's multiple comparison post-hoc test (n=3). One-way ANOVA for each time point was instead done for C. since two-way ANOVA requires matched data for all time points. Adjusted p values reflect post-hoc comparisons between the responses of the wt cell line compared to the DKO cell line at the specific time point (in B.) or Dox-treated cell line compared to matched untreated cell line (in C.) (n=3). Error bars, standard error of mean. n.s., not significant.

Figure 5. A. Normalized expression values of BCL2 family transcripts in PEL cell lines from published RNA-seq datasets (19, 32, 33). **B.** Western blots of the BCL2 family members and HA-tagged vBCL2 in BC-1 or BC-3 cells transduced with lentiviruses over-expressing these cDNAs. **C-D.** Basal caspase 3/7 activity of BC-1 (n=11) or BC-3 cells (n=7) normalized to total live cells measured by CellTiterGlo 2.0 assay. **E-F.** Cell viability measured by CellTiterGlo 2.0 assay or **G-H.** relative caspase 3/7 activity in live BC-1 (n=3) or BC-3 cells (n=4) after overnight treatment of 20 nM or 100 nM staurosporine. Statistical differences were calculated using one- or two-way ANOVA with Tukey's multiple comparison post-hoc test. Adjusted p values reflect post-hoc comparisons between the cell lines in response to staurosporine or DMSO unless indicated otherwise. Error bars, standard error of mean. n.s., not significant.

Figure 6. A-B. Cell viability measured by CellTiterGlo 2.0 assay or **C-D.** relative caspase 3/7 activity in live BC-1 or BC-3 cells over-expressing BCL2 family cDNAs after overnight treatment of 100 nM MCL1i. Statistical differences were calculated using one- or two-way ANOVA with Tukey's multiple comparison post-hoc test (n=5). Adjusted p values reflect post-hoc comparisons between the cell lines in response to staurosporine or DMSO unless indicated otherwise. **E-F.** IC50 values of MCL1i in BC-1 or BC-3 cells. For BC-1, one-way ANOVA with Tukey's multiple comparison post-hoc test was used for statistical tests (n≥3). Since BC-3 cell lines over-expressing BCL2, BCL-xL or BCL-W were highly resistant to MCL1i even at high doses, IC50 values derived from these incomplete sigmoidal dose-response curves had low confidence intervals. Thus, non-parametric Kruskal-Wallis post-test was used. Adjusted p values reflect post-hoc comparisons between the cell lines to the mCherry control. Error bars, standard error of mean. n.s., not significant.