A common human *MLKL* polymorphism confers resistance to negative regulation by phosphorylation

Sarah E. Garnish[1,2], Katherine R. Martin[1,2], Maria Kauppi[1,2], Victoria Jackson[1], Rebecca Ambrose[3,4], Vik Ven Eng[3,5], Daniel Frank[1], Emma C. Tovey Crutchfield[1,6], Komal M. Patel[1], Shene Chiou[1,2], Andre L. Samson[1,2], Marcel Doerflinger[1,2], Christopher R. Horne[1,2], Cathrine Hall[1], Samuel N. Young[1], Vicki Athanasopoulos[7], Carola G. Vinuesa[7], Charlotte A. Slade[1,2,8], Jaclyn S. Pearson[3,4,5], Gregor Ebert[9], John Silke[1,2], James M. Murphy[1,2] and Joanne M. Hildebrand[1,2,10]

1. The Walter and Eliza Hall Institute, Parkville, VIC, 3052, Australia
2. University of Melbourne, Department of Medical Biology, Parkville, 3052, Australia
3. Centre for Innate Immunity and Infectious Diseases, Hudson Institute of Medical Research, Clayton, VIC, 3168, Australia
4. Department of Molecular and Translational Science, Monash University, Clayton, VIC, 3168, Australia
5. Department of Microbiology, Monash University, Clayton, 3168, VIC, Australia
6. University of Melbourne, Faculty of Medicine, Dentistry and Health Sciences, Parkville, 3052, Australia
7. Department of Immunology and Infection, John Curtin School of Medical Research, Australian National University, ACT, Australia
8. Department of Clinical Immunology & Allergy, Royal Melbourne Hospital, Parkville, VIC, 3052, Australia
9. Institute of Virology, Technical University of Munich/Helmholtz Munich, Munich, Germany

10. To whom correspondence may be addressed- jhildebrand@wehi.edu.au

ABSTRACT

Across the globe, 2-3% of humans carry the *p.Ser132Pro* single nucleotide polymorphism in

MLKL, the terminal effector protein of the inflammatory form of programmed cell death,

necroptosis. We show that this substitution confers a gain in necroptotic function in human cells,

with more rapid accumulation of activated MLKL^{S132P} in biological membranes and MLKL^{S132P}

overriding pharmacological and endogenous inhibition of MLKL. In mouse cells, the equivalent

Mlkl S131P mutation confers a gene dosage dependent reduction in sensitivity to TNF-induced

necroptosis in both hematopoietic and non-hematopoietic cells, but enhanced sensitivity to IFN-β

induced death in non-hematopoietic cells. *In vivo*, *Mlkl^{S131P}* homozygosity reduces the capacity to

clear *Salmonella* from major organs and retards recovery of hematopoietic stem cells. Thus, by

dysregulating necroptosis, the S131P substitution impairs the return to homeostasis after systemic

challenge. Present day carriers of the *MLKL S132P* polymorphism may be the key to understanding

how MLKL and necroptosis modulate the progression of complex polygenic human disease.

INTRODUCTION

Necroptosis is a caspase independent form of programmed cell death that originated as a defense against pathogens (Kitur et al., 2016; Liu et al., 2021b; Pearson et al., 2017; Petrie et al., 2019; Upton et al., 2010). Highly inflammatory in nature, necroptosis results in the permeabilization of biological membranes and the release of cytokines, nucleic acids, and intracellular proteins into the extracellular space (Weir et al., 2021) Necroptosis is induced by danger- or pathogen-associated molecular patterns that signal via transmembrane receptors or intracellular pattern recognition receptors (He et al., 2011; He et al., 2009; Holler et al., 2000; Kaiser et al., 2013; Thapa et al., 2013). Of the various initiating stimuli, the most well studied necroptotic pathway is downstream of tumor necrosis factor receptor 1 (TNFR1)(Samson et al., 2021). In physiological contexts that favor low cellular inhibitor of apoptosis protein 1 (cIAP1) and caspase-8 activity, TNFR1 signals culminate in the formation of a high molecular weight platform called the necrosome that is nucleated by heterooligomeric RIPK1 and RIPK3 (Li et al., 2012; Meng et al., 2021; Mompean et al., 2018). Here, the terminal executioner protein, MLKL, is phosphorylated and activated by its upstream kinase, RIPK3 (Murphy et al., 2013; Sun et al., 2012; Zhao et al., 2012). Following phosphorylation, MLKL dissociates from RIPK3, oligomerizes and is trafficked to biological membranes where it interacts with Phosphatidylinositol Phosphates, or 'PIPs' (Cai et al., 2014; Chen et al., 2014; Dondelinger et al., 2014; Garnish et al., 2021; Hildebrand et al., 2014; Petrie et al., 2020; Samson et al., 2020; Sethi et al., 2022; Tanzer et al., 2016; Wang et al., 2014b). In human cells, association of activated MLKL oligomers with biological membranes can be inhibited by the synthetic compound necrosulfonamide (Samson et al., 2020; Sun et al., 2012; Wang et al., 2014a) or inhibitory phosphorylation of MLKL at Serine 83 (Zhu et al., 2022).

Pharmacological or mutation driven disruption at any major necroptotic signaling checkpoint

compromises a cell's capacity to execute necroptosis.

In mouse studies, MLKL mediated cell death has been implicated as a driver or suppressor of

diseases spanning almost all physiological systems depending on the pathophysiological context.

The generation of *Mlkl* gene knockout (*Mlkl*[-/-]), knock-in and conditional knock-out mouse models

have enabled the role of necroptosis in infectious and non-infectious challenges to be dissected in

physiological detail (Tovey Crutchfield et al., 2021). Interestingly, genetic deletion of *Mlkl* has

no overt developmental or homeostatic effects in mice (Murphy et al., 2013; Wu et al., 2013). This

is in direct contrast with two mouse models harboring *Mlkl* point mutations, $Mlkl^{D139V}$ and $Mlkl^{S83G}$,

that dysregulate MLKL activation, which results in early neonatal death and severe inflammatory

phenotypes (Hildebrand et al., 2020; Zhu et al., 2022). Altogether, these observations suggest that

while constitutive absence of MLKL-mediated death is benign, imbalanced execution of

necroptotic cell death is deleterious.

Consistent with this notion, more than 20 unique disease-associated human germline gene variants

in the core necroptotic machinery, encompassing *RIPK1, RIPK3, MLKL*, have been identified

(Garnish and Hildebrand, 2022; Zhang et al., 2021). In one family, a haplotype including a rare

MLKL loss-of-function gene variant (*p.Asp369GlufsTer22*, rs561839347) is associated with a

severe and progressive novel neurogenerative spectrum disorder characterized by global brain

atrophy (Faergeman et al., 2020). A more frequent *MLKL* loss-of-function gene variant

(*p.Gln48Ter*, rs763812068) was found to be >20 fold enriched in a cohort of Hong Kong Chinese

patients suffering from Alzheimer's disease (Wang et al., 2018) and common variants that cluster

around the MLKL brace region were shown to be enriched *in trans* in a cohort of chronic recurrent

multifocal osteomyelitis patients (Hildebrand et al., 2020). More recently, a hypomorphic *MLKL*

missense gene variant (*p.G316D*, rs375490660) was reported to be associated with Maturity Onset

Diabetes of the Young (Hildebrand et al., 2021).

Here we present the cellular and physiological characterization of a serine to proline missense

polymorphism at MLKL amino acid 132 (*p.Ser132Pro*; *S132P*). The *S132P* polymorphism is the

third most frequent human *MLKL* missense coding variant in the gnomAD database; a large

repository of whole genome and exome sequence data from humans of diverse ancestry

(Karczewski et al., 2020). To examine the potential human disease-causing effects of this *MLKL*

variant, we exogenously expressed $MLKL^{S132P}$ in human cell lines and introduced the mouse

counterpart variant ($Mlkl^{S131P}$) into a genetically modified mouse model, revealing that this

polymorphism confers MLKL gain-of-function in a cell- and stimulus-dependent manner. This

MLKL gain-of-function manifests in *in vivo* changes to the immune response, impaired bacterial

clearance, and defective emergency hematopoiesis. These observed phenotypes provide important

insights into how this highly frequent human *MLKL S132P* polymorphism may contribute to the

progression of complex disease.

Results

Family members carrying the *p.Ser132Pro* polymorphism exhibit diverse inflammatory

disease profiles.

With a global minor allele frequency (MAF) of 0.0138, the *MLKL S132P* (rs35589326)

polymorphism is predicted to be carried by 2-3% of the human population. It has not been detected

in individuals assigned East Asian ancestry, is rare in individuals of African or Latino/Admixed

American ancestry and is carried by an estimated 6-7% of individuals of Ashkenazi Jewish

ancestry (MAF 0.0315) (www.gnomAD.com, June 2022) **(Figure 1A)**. Notably, Ser132 is

conserved across species, such as mouse, chimpanzee, rat, bovine, and pig **(Figure 1B)**.

We identified two individuals who are heterozygous carriers of the *MLKL S132P* polymorphism

in an Australian registry of patients suffering from immune related disease. Patient 1, a female of

South American heritage, was diagnosed with SAPHO (synovitis, acne, pustulosis, hyperostosis,

osteitis) syndrome (inheritance chart unavailable). Patient 2, a female of European heritage, was

diagnosed with systemic IgG4 disease **(Figure 1C)**. Both patients have one or more immediate

family members carrying the *MLKL S132P* polymorphism who were also diagnosed with

inflammatory diseases in early adulthood. Following whole genome sequencing, Patient 1 was

found to exhibit a region of loss of heterozygosity (5:96031569 – 5:96364063) which covers *CAST,*

ERAP1, ERAP2 and *LNPEP*. These genes have been previously associated with inflammatory

disease (Cortes et al., 2013; Cortes et al., 2015; Evans et al., 2011; Huang et al., 2013; Kirino et

al., 2013; Robinson et al., 2015; Strange et al., 2010; Tsoi et al., 2012; Yin et al., 2015). Patient 2

does not carry any other predicted pathogenic gene variants. Primary peripheral blood

mononuclear cells (PBMCs) isolated from patient 2 showed reduced MLKL protein levels,

accompanied by a nominal increase in pro-inflammatory cytokine production in response to the

Toll-like receptor (TLR) agonists, lipopolysaccharide and poly I:C, relative to age and sex matched

healthy donor PBMCs not carrying the *MLKL S132P* polymorphism (**Supplementary Figure 1A,**

B).

Exogenously expressed *MLKL*^{S132P} confers resistance to chemical and natural regulatory

inhibition.

To examine any changes to MLKL function conferred by this common polymorphism, we stably

transduced parental wild-type (WT) and $MLKL^{-/-}$ versions of the human colonic epithelial cell line

HT29 with doxycycline inducible $MLKL^{WT}$ and $MLKL^{S132P}$ gene expression constructs. We also

included the T357/S358 phosphosite mutant $MLKL^{TSEE}$, previously shown to be inactive, as a

negative control for necroptosis induction (Petrie et al., 2018). Cells expressing $MLKL^{S132P}$ died

with similar kinetics to $MLKL^{WT}$ following necroptotic stimulation (TNF, SMAC mimetic

compound A and pan-caspase inhibitor IDN-665; TSI) and, as expected, $MLKL^{TSEE}$ reconstituted

cells were resistant to necroptotic cell death **(Figure 1D, Supplementary Figure 1C, D**).

Interestingly, in the presence of the MLKL inhibitor necrosulfonamide (NSA), TSI-stimulated

cells expressing $MLKL^{S132P}$ exhibited higher levels of cell death than their $MLKL^{WT}$ counterparts

at later timepoints **(Figure 1D, Supplementary Figure 1C)**. We termed this override of MLKL

inhibition 'breakthrough' cell death. Breakthrough death also occurred when $MLKL^{S132P}$ was co-

expressed with endogenous $MLKL$ and increased when MLKL expression was augmented by

higher doses of doxycycline **(Figure 1D**, **Supplementary Figure 1C, D).** We also observed

breakthrough death in both wild-type and $MLKL^{-/-}$ forms of the human monocytic cell line U937,

indicating this is not a cell type specific phenomenon **(Supplementary Figure 1E, F)**.

Breakthrough death was neither due to differences in MLKL expression, nor changes in MLKL

phosphorylation (T357/S358), because both were equivalent between $MLKL^{WT}$ and $MLKL^{S132P}$

reconstituted cells **(Figure 1E, Supplementary Figure 1G, H)**. Instead, cellular fractionation

experiments suggest that breakthrough death was likely due to enhanced association of MLKL^{S132P}

with biological membranes **(Figure 1F, Supplementary Figure 1I)**.

The small molecule inhibitor, necrosulfonamide (NSA), blocks human MLKL activity through

covalent modification of Cysteine 86, located in the MLKL four helix-bundle, executioner domain

(Figure 1G) (Sun et al., 2012; Wang et al., 2014a)**.** Recently, it was discovered that MLKL can be

endogenously phosphorylated at the proximal residue, Ser83, and this phosphorylation event plays

a species conserved inhibitory role in both human (Ser83) and mouse (Ser82) **(Figure 1G)** (Zhu

et al., 2022). Like NSA, phosphorylation of MLKL Ser83 does not alter the capacity of RIPK3 to

phosphorylate MLKL at Ser357/T358 but prevents necroptosis by blocking the association of

MLKL with cellular membranes (Zhu et al., 2022). To test whether phosphorylation at Ser83 is,

like NSA, less effective at inhibiting MLKL^{S132P} induced death, we created stable HT29 (*MLKL$^{-/-}$*

and wild-type) cell lines that exogenously expressing gene constructs encoding *MLKL^{WT}*,

MLKL^{S132P}, *MLKL^{S83D}* (phosphomimetic) and double mutant *MLKL^{S83D,S132P}*. Consistent with

recent studies, the phosphomimetic mutation of S83, *MLKL^{S83D}*, ablated the capacity of MLKL to

execute necroptotic cell death (Zhu et al., 2022). Also, as previously reported, the inhibitory effect

of *MLKL^{S83D}* overrode the activating effects of endogenous MLKL phosphorylation at

Ser357/Thr358 **(Figure 1H, Supplementary Figure 1J, K)**. Exogenous expression of *MLKL^{S83D}*

also reduced total levels of necroptotic cell death in the presence of endogenous MLKL, indicating

that this S83 phosphomimetic acts in a dominant negative manner **(Supplementary Figure 1L)**.

Strikingly, combining the MLKL^{S132P} and MLKL^{S83D} amino acid replacements to create the

compound mutant MLKL$^{S83D, S132P}$ restored necroptotic cell death responsiveness, albeit with

reduced kinetics and reduced maximal cell death when compared to MLKL^{S132P} alone **(Figure 1H,**

Supplementary Figure 1K). The *MLKL^{S132P}* variant also partially overcomes the dominant

negative effect of Ser83 phosphomimetic mutation in a wild-type background **(Supplementary**

Figure 1L). These data suggest that necroptotic cell death is dysregulated in human carriers of the

common *MLKL S132P* variant. This conclusion, together with the finding of immunoinflammatory

disorders in *MLKL S132P* carriers, raises important questions about how, when, and where, proper

regulation of necroptosis contributes to immune function *in vivo.*

A mouse model of the human *MLKL S132P* polymorphism exhibits differences in steady state

immune cell populations

To address if the *MLKL S132P* polymorphism contributes to immunoinflammatory disease, we

performed detailed histological, immunophenotypic, and experimental analyses of a CRISPR

generated knockin mouse which carried the orthologous mutation, *Mlkl^{S131P}*. *Mlkl^{S131P}*

heterozygotes and homozygotes were born according to the expected Mendelian ratios and had

normal lifespans **(Figure 2A, B)**. The healthy presentation of *Mlkl^{S131P}* mice contrasts with the

lethal phenotypes of mice engineered to encode *Mlkl^{D139V}* and *Mlkl^{S83G}* mutations (Hildebrand et

al., 2020; Zhu et al., 2022). Wild-type, *Mlkl^{S131P}* heterozygote and homozygote mice had

comparable body weight and no gross histological differences up to 9 months of age **(Figure 2C,**

Supplementary Figure 2A). Notably different to *Mlkl^{D139V}* or *Mlkl^{S83G}* homozygotes, no

inflammation was detected in the salivary glands, mediastinum, liver or lungs (**Supplementary**

Figure 2A) (Hildebrand et al., 2020; Zhu et al., 2022). No differences in the number of circulating

platelets, red blood cells, or white blood cells were observed between wild-type, *Mlkl^{S131P}*

heterozygote and homozygote mice across age **(Supplementary Figure 2B-D)**. However, reduced

numbers of classical Ly6C^{hi} 'inflammatory' monocytes in the bone marrow, but not in the

secondary lymphoid organs (spleen and inguinal lymph nodes), were observed in *Mlkl^{S131P}*

homozygotes relative to wild-type controls (**Figure 2D, Supplementary Figure 2E, K, Q**). In the

secondary lymphoid organs, *Mlkl^{S131P}* homozygotes had a small but significant increase splenic

CD4+ T cells and B cells relative to wild-type controls **(Figure 2E, Supplementary Figure 2N-**

P). All other innate and adaptive immune cell populations were comparable between genotypes in

the bone marrow, spleen, and inguinal lymph nodes **(Figure 2D-F, Supplementary Figure 2E-**

V).

We next sought to address whether mouse MLKL^{S131P} exhibited a gain-in-function consistent with

what we described for MLKL^{S132P} in human cells. Primary and immortalized fibroblasts isolated

from the dermis of *Mlkl^{S131P}* homozygote, heterozygote and wild-type mice were examined for

their relative capacity to die in the presence or absence of death stimuli. We did not observe any

differences in TNF-induced apoptosis (TNF and SMAC mimetic; TS) between *Mlkl$^{S131P/S131P}$* and

Mlkl$^{WT/WT}$ immortalized MDFs **(Figure 2G)**. However, *Mlkl$^{S131P/S131P}$* MDFs showed a diminished

capacity to undergo TSI-induced (TNF, SMAC-mimetic compound A, pan-caspase inhibitor IDN-

6556; TSI) necroptotic cell death **(Figure 2G, Supplementary Figure 2W)**. This was driven by a

gene-dosage reduction in MLKL protein level present in MDF cells with the *Mlkl^{S131P}* allele

(Figure 2H, Supplementary Figure 2X). Immortalized MDFs, and to a lesser extent primary

MDFs, did show clear *Mlkl^{S131P}* gene-dose dependent sensitivity to IFN-β, a strong inducer of *Mlkl*

gene expression in mice **(Figure 2G, Supplementary Figure 2W)** (Rodriguez et al., 2016). This

sensitivity was further enhanced by TNF and was refractory to inhibitors of RIPK1 (Nec-1s) or

RIPK3 (GSK-872) **(Figure 2G, Supplementary Figure 2W)**. These results are analogous to cells

expressing the auto active MLKL^{D139V} mutant (Hildebrand et al., 2020) and together support the

notion that MLKL^{S131P} has constitutive RIPK3-independent activity when endogenously expressed

in mouse cells (Hildebrand et al., 2020). These endogenous *Mlkl^{S131P}* observations were not limited

to fibroblasts. In bone marrow derived macrophages (BMDMs), *Mlkl$^{S131P/S131P}$* cells also exhibited

diminished sensitivity to TSI-induced necroptosis and endogenously produced MLKL^{S131P} was

present at reduced levels in comparison to MLKL^{WT} (**Supplementary Figure 2Y, Z**). In contrast

to immortalized dermal fibroblasts, BMDMs derived from *Mlkl^{S131P}* homozygotes did not undergo

cell death in the presence of IFN-β alone, despite clear IFN-β induced upregulation of MLKL

protein relative to untreated cells (**Supplementary Figure 2Y, Z**). Overall, these data show that

across different cell types MLKL^{S131P} is not toxic at steady state, but upon different stimuli, both

gain-of-sensitivity and loss-of-sensitivity to necroptotic cell death is evident. This suggests that

diseases caused by the *MLKL S132P* polymorphism are likely to be diverse, with cell- or context-

specific manifestations.

Emergency hematopoiesis is defective in *Mlkl$^{S131P/S131P}$* mice

Differences in steady state immune cell populations suggest that overt pathological phenotypes

may develop in *Mlkl^{S131P}* homozygotes following experimental challenge. Specifically, reduced

numbers of steady state Ly6C^{hi} monocytes in the bone marrow of *Mlkl^{S131P}* homozygotes could

indicate a defect in hematopoiesis. To test this, we examined recovery from myelosuppressive

irradiation as an assessment of hematopoietic function in mice carrying the *Mlkl^{S131P}* allele.

Following myelosuppressive irradiation, recovery of hematopoietic cell numbers and circulating

peripheral blood cells was significantly delayed in *Mlkl$^{S131P/S131P}$* mice compared with wild-type

controls (**Figure 3A-F**). In *Mlkl$^{S131P/S131P}$* mice, peripheral red blood cell and platelet numbers were

significantly reduced at 14 days post irradiation, with the former also decreased at 21 days (**Figure**

3A, B). Despite equivalent total white blood cell numbers at 21 days post irradiation, a significant

reduction in peripheral monocyte numbers and a non-significant decrease trend in neutrophils was

observed in *Mlkl$^{S131P/S131P}$* mice compared to wild-type controls (**Figure 3C, Supplementary**

Figure 3A, B). These reductions in circulating blood cell numbers were accompanied by

significant decreases in the nucleated viable, progenitor, and LSK cell populations in the bone

marrow of $Mlkl^{S131P/S131P}$ mice **(Figure 3D-F)**. Contrastingly, at 21 days post irradiation, $Mlkl^{S131P}$

heterozygotes displayed a significant increase in their total nucleated viable cells in comparison to

wild-type controls. This was driven predominantly by significant increases in the number of LSK

cells (**Figures 3D, E**).

To investigate whether the $Mlkl^{S131P}$ homozygote defect was intrinsic to the hematopoietic stem

cells, we performed competitive transplantation studies. $Mlkl^{S131P/S131P}$ or $Mlkl^{WT/WT}$ bone marrow

was mixed with GFP$^+$ competitor bone marrow in a 50:50 ratio and injected into Ly5.1 irradiated

hosts. Six weeks post-transplant, $Mlkl^{WT/WT}$ bone marrow had competed with GFP$^+$ bone marrow

effectively, whilst $Mlkl^{S131P/S131P}$ bone marrow performed poorly, contributing to 7%, 5% and 9%

of PBMCs (Ly5$^+$), red blood cells and platelets respectively **(Figure 3G-I)**. Similarly,

$Mlkl^{S131P/S131P}$ bone marrow was outcompeted, again contributing to less than 10 % of the blood

cells, when mixed in excess at 70:30 with GFP$^+$ competitor bone marrow (**Supplementary Figure**

3C-E). Thus, the defect in emergency hematopoiesis was intrinsic to $Mlkl^{S131P/S131P}$ hematopoietic

stem cells, consistent with the intrinsic defect previously reported for the $Mlkl^{D139V}$ autoactivating

mutant (Hildebrand et al., 2020).

Recruitment of Ly6C^{hi} monocytes to sites of sterile inflammation is reduced in $Mlkl^{S131P/S131P}$

mice

To examine if the defects in emergency hematopoiesis displayed by $Mlkl^{S131P/S131P}$ mice could

result in defective immune cell recruitment to sites of inflammation or infection, we employed a

model of localized sterile inflammation induced by zymosan. Intra-peritoneal injection of zymosan

results in a rapid influx of immune cells, predominantly neutrophils, into the peritoneal cavity over

72 hours (Martin et al., 2017). The cellular content of the peritoneal cavity and numbers of

circulating peripheral blood cells was examined at 4- and 24-hours post zymosan injection.

Mlkl$^{S131P/S131P}$ mice displayed equivalent early recruitment of neutrophils and other immune cells

to the peritoneum **(Figure 4A, B, Supplementary Figure 4A-H).** However, at 24 hours post

zymosan injection *Mlkl*$^{S131P/S131P}$ mice had significant reductions in the number of Ly6C^{hi}

monocytes recruited to the peritoneum **(Figure 4C, D, Supplementary Figure 4I-P)**. In the

peripheral blood, monocytes were significantly elevated in *Mlkl*$^{S131P/S131P}$ mice in comparison to

wildtype controls at 4 hours post zymosan **(Figure 4E, Supplementary Figure 4Q-S)**. However,

by 24 hours, the number of peripheral blood monocytes was equivalent between genotypes, with

only a non-significant increase in neutrophils observed in *Mlkl*$^{S131P/S131P}$ mice **(Figure 4F,**

Supplementary Figure 4T-V). Compared to wild-type mice, the recruitment of Ly6C^{hi} monocytes

to the peritoneum of *Mlkl*^{S131P} homozygotes was hampered. These decreases in Ly6C^{hi} monocytes

at the site of sterile inflammation is likely to be driven by the reduced number of this population

in the bone marrow at steady state.

We previously observed that *Mlkl*$^{S131P/S131P}$ cells exhibited reduced response to TSI-induced

necroptosis. To assess whether this defect in necroptotic function was observed in activated

immune cells recruited to sites of inflammation, we isolated neutrophils from the peritoneum at 4

hours post zymosan injection. Consistent with our previous *in vitro* findings, no differences were

observed in the capacity of *Mlkl*$^{S131P/S131P}$ inflammatory neutrophils to undergo spontaneous

apoptosis **(Figure 4G).** However, the percentage of *Mlkl*$^{S131P/S131P}$ inflammatory neutrophils dying

following necroptotic stimulation, TSI, was decreased in comparison to wild-type neutrophils

(Figure 4H).

Mlkl[S131P] homozygotes exhibit reduced capacity to clear *Salmonella* infection

Reductions in the recruitment of $Ly6C^{hi}$ monocytes to sites of inflammation, raise important

questions as to whether challenge with a pathogen may reveal defects in the immune cell response

in *Mlkl*[S131P] homozygotes. *Salmonella enterica* has long been an important pathogenic selective

pressure in humans (Key et al., 2020). For non-typhoidal *Salmonella* serovars such as *S. enterica*

serovar *Typhimurium*, human infection is typically limited to the gastric mucosa (Galan, 2021;

Hohmann, 2001). In mice and immunocompromised humans, *S. Typhimurium* can cause severe

systemic disease after dispersal from the gut by dendritic cells and macrophages to peripheral

organs including the spleen and liver (Wemyss and Pearson, 2019).

Mlkl[S131P] homozygote, heterozygote and wild-type mice were infected with a metabolically

growth-attenuated *Salmonella typhimurium* strain BRD509 (from here referred to as *Salmonella),*

via oral gavage (1 x 10^7 colony forming units (CFU)). Daily monitoring for the 14-day infection

period showed no differences in core body temperature or body mass between genotypes

(Supplementary Figure 5A, B). Consistent with normal weights, there were no obvious

differences in the integrity of the intestinal epithelial barrier or intestinal monocyte and

macrophage populations at infection endpoint **(Figure 5A)**. Despite this, *Mlkl*[S131P/S131P] mice had

increased *Salmonella* burden in both the spleen and liver compared to wild-type controls **(Figure**

5B, Supplementary Figure 5C, D). Bacterial colonization in the feces was increased in female

Mlkl[S131P/S131P] mice relative to wild-type female controls **(Figure 5B, Supplementary Figure 5E)**.

In the peripheral blood, *Mlkl*[S131P/S131P] mice had significantly reduced numbers of circulating

lymphocytes and monocytes, as well as a trend towards decreased numbers of circulating

neutrophils **(Figure 5C, Supplementary Figure 5F-H)**. Infected *Mlkl*^{S131P/S131P} mice also

exhibited significant decreases in the quantity of splenic Ly6C^{hi} monocytes in comparison to

infected wild-type controls **(Figure 5D, E, Supplementary Figure 5I-O)**. No differences in

splenic Ly6C^{hi} monocytes were observed in uninfected mice, therefore the reduction in Ly6C^{hi}

monocytes in infected spleens is likely to be due to the reduced number of this monocyte

population in the bone marrow and defects in emergency monopoiesis. Furthermore, we did not

observe any difference in the kinetics of *Salmonella* induced cell death in ex-vivo BMDM

infection assays, nor inflammasome activation, measured by GSDMD cleavage **(Supplementary**

Figure 5P, Q). Together these data show a hindered pathogen defense in *Mlkl*^{S131P} homozygotes

resulting from widespread immunophenotypic deficiencies. Combined, our investigations of

Mlkl^{S131P} homozygotes under challenge highlight a defect in emergency hematopoiesis that

manifests in a disruption to integral inflammatory and immune responses. This provides important

insights into the potential causes of immunoinflammatory disorders in *MLKL S132P* carriers.

DISCUSSION

Present in 2-3% of the global population, the high frequency *MLKL S132P* polymorphism

confers a gain-of-function to MLKL resulting in hematopoietic dysfunction and immune cell

defects in a genetically modified mouse model. In human cells, MLKL^{S132P} was resistant to

chemical inhibition by necrosulfonamide treatment and endogenous inhibitory phosphorylation at

Ser83. A gain in necroptotic function was also observed when murine equivalent *Mlkl*^{S131P} was

examined *in situ,* with fibroblasts exhibiting RIPK3-independent cell death in the presence of IFN-

β. When subject to endogenous protein regulatory mechanisms, MLKL^{S131P} protein levels are

reduced, manifesting in a reduced capacity for necroptosis in fibroblasts and immune cells

stimulated with TSI. Extensive characterization of the $Mlkl^{S131P}$ mouse unveiled steady state

decreases in the bone marrow pool of inflammatory Ly6C^{hi} monocytes, that manifested in reduced

numbers at the site of sterile inflammation and bacterial infection. Specifically, under infection

challenge, an increased burden of *Salmonella* was present in the spleen and liver of homozygotes,

accompanied by significant reductions in the circulating lymphocytes and monocytes. We

observed significant hematopoietic dysfunction in $Mlkl^{S131P}$ homozygotes, with a catastrophic drop

in emergency hematopoiesis and complete failure of hematopoietic stem cells under transplant. In

sum, both the human $MLKL^{S132P}$ and mouse equivalent $Mlkl^{S131P}$ variants confer a gain-of-

function to MLKL *in vitro*. The resultant dysregulation of necroptotic signaling manifests in

debilitating deviations in baseline immune phenotype and function in our genetically modified

mouse model, $Mlkl^{S131P}$.

Prior to this work two other single point mutant $Mlkl$ mouse models, $Mlkl^{D139V}$ and

$Mlkl^{S83G}$, had been reported, both of which exhibited full or partial homozygous postnatal lethality

characterized by severe inflammation (Hildebrand et al., 2020; Zhu et al., 2022). In stark contrast

to these models, the $Mlkl^{S131P}$ homozygotes were born normal and with no signs of inflammatory

disease at steady state. However, upon challenge, similarities are observed between $Mlkl^{S131P}$ and

$Mlkl^{D139V}$ mice. Hematopoietic dysfunction is observed following myelosuppressive irradiation in

homozygous or heterozygous mice harboring the $Mlkl^{S131P}$ or $Mlkl^{D139V}$ mice respectively

(Hildebrand et al., 2020).

Despite the ostensible basal state phenotypes of $Mlkl^{D139V}$ (Hildebrand et al., 2020) and

recently described $Mlkl^{S83G}$ (Zhu et al., 2022) homozygotes differing from the $Mlkl^{S131P}$

homozygotes, there were similarities observed at the molecular level. As for MLKL^{S131P}, RIPK3-

independent gain-of-function was reported for MLKL^{D139V}, but not MLKL^{S83G} (Hildebrand et al.,

2020; Zhu et al., 2022). This major similarity in molecular function is not unexpected considering

the proximity of the D139 and S131 residues within MLKL. Constitutive activity of both the

S132P and D139V brace helix mutants is restrained by cellular mechanisms that limited protein

level, with decreases in endogenous MLKL observed in cells generated from *Mlkl^{S131P}* and

Mlkl^{D139V} homozygotes. Interestingly, ubiquitin-mediated targeting of MLKL to the lysosome was

identified to be mechanistically involved in the restraint of necroptosis in cells expressing

MLKL^{D139V}, and as a means of clearing endosomal *Listeria monocytogenes* and *Yersinia*

enterocolita (Hildebrand et al., 2020; Liu et al., 2021a; Yoon et al., 2022). Exploring if the

reduction in cellular MLKL levels we observed in the PBMCs of a human carrier of the *S132P*

polymorphism, or cells isolated from the S131P mouse, is likewise mediated by ubiquitination and

lysosomal targeting will be an important next step, particularly as it relates to the clearance of these

intracellular bacteria from human cells. Ablation of the inhibitory phosphorylation site in *Mlkl^{S83G}*

mice similarly subjects MLKL to downregulation in mouse cells (Zhu et al., 2022). In sum, our

analysis of MLKL^{S131P} further highlights the importance of mechanisms that downregulate MLKL

when aberrantly activated.

The finding that MLKL^{S132P} retains necroptotic killing activity despite simulated inhibitory

phosphorylation at Ser83 in human cells is highly notable. The inflammatory phenotypes that

result from homozygosity of phosphomimetic (S83G) mutation provides insight into the potential

disease development that may occur in carriers of the *MLKL S132P* polymorphism under

environmental and cellular scenarios where Ser83 inhibitory phosphorylation is deployed.

Identifying the kinase(s) responsible for phosphorylation at MLKL Ser83 and understanding which

cell types are primed to deploy this kinase(s) is an important next step in determining if this

polymorphism promotes clinically relevant changes to homeostasis or disease outcomes in

humans.

Evidence for positive selection has been found for over 300 immune-related gene loci and

many of these have been found to be associated with the incidence of autoimmune and

autoinflammatory disease in modern humans (Gutierrez-Arcelus et al., 2016; Ramos et al., 2015).

Many of these variants have also been mechanistically linked to defense against a particular

pathogen (Karlsson et al., 2014, Ramos et al., 2015), with pathogenic microbes a major driver of

genetic selection over human history. While a diminished capacity for mice to clear disseminated

Salmonella would argue against the hypothesis that the *MLKL S132P* polymorphism has been

positively selected for in human populations, it is important to note that only 1.4 % of human

carriers are homozygotes (Karczewski et al., 2020). *MLKL S132P* heterozygosity is by far the most

prevalent, and evolutionarily relevant, human scenario. *Mlkl*[S131P] heterozygotes displayed gene

dosage phenotypes consistent with homozygotes, with one exception. Following sublethal

myelosuppressive irradiation, a gain in bone hematopoietic stem cell numbers in heterozygotes

was observed whilst a catastrophic drop occurred in homozygotes. Whether this increase in stem

cell numbers persists over time is currently under investigation. However, increased fitness

conferred to heterozygous mice at this timepoint provides intriguing insights into other selective

pressures that may have promoted accumulation of this polymorphism in humans (Werren et al.,

2021). Sepsis, where emergency hematopoiesis is an essential determinant in survival, is an

interesting and highly evolutionarily significant avenue of exploration for the study of *MLKL*

S132P polymorphism frequency (Boomer et al., 2011; Brinkworth and Valizadegan, 2021). With

nutrition being another important driver of genetic adaptation in humans, exploring the role of

MLKL S132P in metabolic disease is also compelling. An important precedent for this is a recent

report of an association between human RIPK1 promoter polymorphisms and diet-induced obesity

(Karunakaran et al., 2020). The potential for negative selection of this polymorphism over time is

also an important avenue of exploration in light of the recent discovery of a common *TYK2* variant

accounting for and its role in enhancing susceptibility to severe infection by historically important

human pathogen mycobacterium tuberculosis (Boisson-Dupuis et al., 2018; Kerner et al., 2021).

By many, MLKL is viewed as a potential therapeutic target for drug discovery due to its

established involvement in multiple human diseases, especially inflammatory pathologies

(Hildebrand et al., 2020; Pierdomenico et al., 2014; Rickard et al., 2014a; Rickard et al., 2014b).

To date no human clinical trials have been conducted on MLKL-targeted small molecules,

although several inhibitors have been reported in the literature, including first-of-kind human

MLKL inhibitor necrosulfonamide (NSA), which all function by targeting Cys86 (Sun et al., 2012;

Yan et al., 2017). Intriguingly, we show that in the presence of NSA human MLKL^{S132P} protein

displays a gain-of-function that results in non-inhibitable cell death. While NSA has been a

fundamental tool for *in vitro* studies of necroptosis, our findings raise serious questions about the

suitability of Cys86 as a target for MLKL drug discovery programs, because such therapies would

not be suitable for 2-3% of the population that carry the *MLKL S132P* polymorphism.

METHODS

Patient Recruitment, Ethics and Informed consent

Patients and their relatives were recruited from the Department of Clinical Immunology and

Allergy, Royal Melbourne Hospital, Victoria, Australia and the Centre for Personalized

Immunology, Australian National University, Canberra, Australia. Unrelated, age and sex matched

'healthy' controls that did not carry the *MLKL p.Ser132Pro* polymorphism were recruited via the

Volunteer Blood Donor Registry, Parkville. Informed consent was obtained from all participants

for genomic analysis and immunological studies. All procedures were performed and are reported

here with the approval of human ethics review boards of all Institutes that participated in human

genetics studies; Australian National University, The Walter and Eliza Hall Institute of Medical

Research (approved projects 2009.162, 10/02) and with the 1964 Helsinki declaration and its later

amendments or comparable ethical standards.

Genomic analysis

Whole exome sequencing was performed by the Canberra Clinical Genomics service. Libraries

were prepared and enriched using the SureSelect Clinical Research Exome v2 kit (Agilent

Technologies), and targeted regions were sequenced using an Illumina sequencing system with

100bp paired-end reads, with an average depth of >35. Reads were aligned to the human genome

reference sequence (GRCh38) using the Burrows-Wheeler Aligner (BWA-MEM), and variant

calls made using the Genomic Analysis Tool Kit (GATK).

Animal ethics

All mice were housed at the Walter and Eliza Hall Institute of Medical Research (WEHI),

Australia. This facility is a temperature and humidity controlled specific pathogen free facility with

a 12h:12h day night cycle. All experiments were approved by the WEHI Animal Ethics Committee

in accordance with the Prevention of Cruelty to Animals Act (1986) and the Australian National

Health and Medical Research Council Code of Practice for the Care and Use of Animals for

Scientific Purposes (1997).

Mice

Mice were generated on a C57BL/6J background. The *p.Ser131Pro* mutation in the *Mlkl* gene

(*Mlkl^S131P^*) was generated using CRISPR/Cas9 by the Melbourne Advanced Genome Editing

Centre (MAGEC) laboratory at WEHI, following the same methodology as previously described

(Hildebrand et al., 2020; Wang et al., 2013). To introduce a proline-coding mutation in place of

Ser131 within the *Mlkl* gene, an sgRNA of the sequence CTGTCGATCTTCCTGCTGCC was

used to create double stranded breaks within the *Mlkl* locus and initiate homologous

recombination.

An   oligo   donor

(TGTTGCTGCTGCTTCAGGTTTATCATTGGAATACCGTTTCAGATGTCAGCCAGCCAG

CACCATGGCAGCAGGAAGATCGACAGGATGCAGAGGAAGACGGgtgagtctcccaaagactgg

ga) was subsequently used to introduce the S131P mutation. Genotyping of mice was completed

by   Transnetyx   using   custom   probe   Mlkl-8   MUT   (Forward   Primer:

CTGCTTCAGGTTTATCATTGGAATACC,   Reverse   Primer:

TCTGCATCCTGTCGATCTTCCT).

Reagents

Antibodies; Rat anti-mMLKL 8F6 (1:2000), rat anti-mMLKL 5A6 (1:1000), rat anti-hMLKL

10C2 (1:1000), rat anti-hMLKL 7G2 (1:1000), rat anti-hRIPK3 1H2 (1:1000) were produced in-

house. Mouse anti-actin (A-1978; 1:5000) was purchased from Sigma-Aldrich, rabbit anti-

GAPDH (#2118; 1:2000-5000) was purchased from Cell Signalling Technology, rabbit anti-

VDAC (AB10527; 1:10000) was purchased from Millipore, rabbit anti-human pMLKL

(EPR9514; 1:1000-3000) was purchased from Abcam, and rabbit anti-mouse pMLKL (D6E3G;

1:1000) was purchased from CST. Cell treatments were completed with agonists/antagonists at the

following concentrations: 100 ng/ml recombinant hTNF-fc (produced in house as in (Bossen et

al., 2006)), 500 nM SMAC mimetic Compound A (provided by Tetralogic Pharmaceuticals; as in

(Vince et al., 2007), 5 µM Pan-caspase inhibitor IDN-6556 (provided by Tetralogic

Pharmaceuticals), 1 µM necrosulfonamide (NSA; Merck #480073), 5 µM necrostatin (Nec-1s;

Merk #504297), 1 µM GSK-'872 (SynKinase #SYN-5481), 10-20 ng/ml lipopolysaccharide (LPS;

Sigma #L2630), 25 µg/ml polyinosinic:polycytidylic (Poly I:C; Novus), 200 nM MG132 (Merck

#474790), 2 nM PS341 (Sigma #504314), and 30 ng/ml mouse IFNβ (R&D Systems #8234-MB-

010)

Generation of cell lines

Mutations were introduced into a human MLKL DNA template (from DNA2.0, CA) using

oligonucleotide-directed PCR and sub-cloned into the pF TRE3G PGK puro vector(Murphy et al.,

2013). Vector DNA was co-transfected into HEK293T cells with pVSVg and pCMV R8.2 helper

plasmids to generate lentiviral particles. U937 (WT and *MLKL*$^{-/-}$) and HT29 (WT and *MLKL*$^{-/-}$)

were then stably transduced with exogenous human MLKL ligated into pFTRE3G. Successfully

transduced cells were selected using puromycin (2.5µg/mL; StemCell Technologies) using

established procedures (Hildebrand et al., 2014; Murphy et al., 2013; Petrie et al., 2018). The

following oligonucleotides were used for the assembly of constructs:

hMLKL^{S132P} fwd; 5′ GCCAAGGAGCGCCCTGGGCACAG3′

hMLKL^{S132P} rev; 5′ CTGTGCCCAGGGCGCTCCTTGGC 3′

hMLKL^{TSEE} fwd; 5′ GAGGAAAACACAGGAGGAAATGAGTTTGGGAAC 3′

hMLKL^{TSEE} rev; 5′ GTTCCCAAACTCATTTCCTCCTGTGTTTCCTC 3′

hMLKL^{S83A} fwd; 5′ GTTCAGCAATAGAGCCAATATCTGCAG 3′

hMLKL^{S83A} rev; 5′ CCTGCAGATATTGGCTCTATTGCTGAAC 3′

hMLKL^{S83D} fwd; 5′ GAAAAGTTCAGCAATAGAGACAATATCTGCAGGTTTC 3′

hMLKL^{S83D} rev; 5′ GAAACCTGCAGATATTGTCTCTATTGCTGAACTTTTC 3′

Primary mouse dermal fibroblasts (MDFs) were prepared from skin taken from the head and body

of 1-day old mice. These MDFs were immortalised by stable lentiviral transduction with SV40

large T antigen. Bone marrow derived macrophages were generated from the long bones of adult

mice and grown for 7 days in DMEM supplemented with 15% L929 cell supernatant.

Human blood (patient and healthy donor) was collected by collaborators via venipuncture at the

Royal Melbourne Hospital. Collected blood was diluted with PBS and layered on an equal volume

of Histopaque (density 1.077 g/ml) and centrifuged for 30 minutes, 700 x g at room temperature.

The layer containing peripheral blood mononuclear cells (PBMCs) was harvested and washed with

PBS, then frozen in FCS + 10 % DMSO and stored in liquid nitrogen.

Culture of cell lines

Primary MDFs, immortalised MDFs and HT29s were cultured in DMEM + 8% FCS. BMDMs

were cultured in DMEM + 15% FCS + 20% L929. U937 and PBMCs were cultured in RPMI +

8% FCS. All cell lines were grown at 37 °C and 10% (v/v) CO_2.

Western blot

U937 cells were seeded into 48-well plates at 60,000 cells/well and induced for 3 hours with

doxycycline (20 ng/mL, 100 ng/mL or 500 ng/mL) to stimulate MLKL expression. HT29 cells

were seeded into 48-well plates at 45,000 cells/well and following 12-14 hours of cell adherence,

cells were induced overnight with doxycycline (20 ng/mL, 100 ng/mL or 500 ng/mL) for

stimulation of MLKL expression. BMDMs were plated at 400,000 cells/well in a 24-well plate

and MDFs (primary and immortalised) were plated at 30,000 cells/well in a 48-well plate. Cells

were stimulated as indicated at described for 6 hours, except for BMDMs stimulated with LPS for

2 hours before addition of SMAC-mimetic Compound A for a further 4 hours. Human primary

PBMCs were plated at 45,000 cells/well and stimulated for 4 hours. All cells were harvested in 2

x SDS Laemmli's lysis buffer, boiled at 100 °C for 10-15 min, and then resolved by 4-15% Tris-

Glycine SDS-PAGE (Bio-Rad). Proteins were transferred to nitrocellulose or PVDF membrane

and probed with antibodies as indicated.

IncuCyte analysis

Primary and immortalised MDFs were plated at 8,000 cells per well in a 96-well plate. BMDMs

were plated at 150,000 cells per well on day 6 of culture in a 48-well plate. MDFs and BMDMs

were left to settle overnight before stimulation. The next day MDFs and BMDMs were stimulated

in culture media supplemented with propidium iodide. HT29 cells were plated at 45,000 cells per

well in a 48-well plate and left to settle for 6 hours before overnight doxycycline pre-stimulation

(20ng/mL, 100ng/mL or 500ng/mL). HT29 cells were stimulated in Phenol Red-free media

supplemented with 2% FCS, 1mM Na pyruvate, 1mM L-GlutaMAX, SYTOX Green (Invitrogen,

S7020) and either DRAQ5 (Thermofisher, #62251) (MLKL KO HT29) or SPY_620

(Spirochrome, SC401) (WT HT29).

U937 cells were plated at 60,000 cells per well in a 48-well plate and were induced with

doxycycline (20ng/mL, 100ng/mL or 500ng/mL) for 3 hours. Cells were then stimulated in Phenol

Red-free media supplemented with 2% FCS, 1mM Na pyruvate, 1mM L-GlutaMAX, SYTOX

Green and either DRAQ5 (MLKL KO HT29) or SPY_620 (WT HT29).

Neutrophils isolated from the peritoneum at 4 hours post-zymosan injection were counted and

plated at 60,000 cells/well in a 48-well plate. Plating media (RPMI + 8 % FCS) was supplemented

with SYTOX Green and DRAQ5 dyes.

Images were taken every hour using IncuCyte SX5 or S3 imaging and cell death was quantified

by number of dead cells (SYTOX Green or propidium iodide positive). Percentage values were

quantified by number of dead cells out of total live cell number (DRAQ5 positive).

TNF ELISA

100,000 cells were stimulated with LPS (10ng/mL) or Poly I:C (2.5µg/mL) for 3 hours. PBMC

supernatant cytokine content was measured by ELISA (R&D: STA00C) according to the

manufacturer's instructions. The measurements were performed in technical triplicates.

Mouse histopathology

7–9-month-old mice were euthanized by CO_2 and fixed in 10 % buffered formalin. For the full

body, 5-µm sagittal sections were taken at 300-µm intervals of all organs. Histopathologists Aira

Nuguid and Tina Cardamome at the Australian Phenomics Network, Melbourne completed

thorough examination of these sections.

Haematological analysis

Cardiac, submandibular or retro-orbital blood collected from mice at steady state (8-52 weeks old)

or following challenge was placed into EDTA coated tubes. Blood cells were left undiluted or

diluted 2- to 11-fold in DPBS for automated blood cell quantification using an ADVIA 2120i

haematological analyser on the same day as harvest.

Mouse model of *Salmonella* infection

Mice used in this experiment were a mix of littermates and non-littermates aged 6-12 weeks, and

wildtype mice that were littermates behaved equivalently to non-littermates. Mice were infected

with *Salmonella enterica* serovar Typhimurium strain BRD509 (Strugnell et al., 1992) at 10^7

colony forming units (CFU) by oral gavage. Mice were harvested 14 days post-infection. Cardiac

bleeds were taken, and blood populations analysed using an ADVIA hematology analyser. Liver,

spleen, and faeces were harvested for enumeration of viable bacteria on nutrient agar. Organs from

infected mice were weighed and homogenised in 2 mL (spleens), 5 mL (livers) or 1 mL (faeces)

of PBS. Homogenates were serially diluted (in duplicate) in PBS and 10 µl drops plated out in

duplicate onto LB agar (+ streptomycin) and incubated overnight at 37 ℃. CFU/mL was calculated

per organ for each mouse and then standardised to CFU/organ based upon organ weight. A small

portion (1/3) of the spleen was processed for flow cytometry analysis.

In vitro Salmonella **infection**

In vitro infection of BMDMs with *Salmonella enterica* serovar Typhimurium strain SL1344 were

performed as previously reported (Doerflinger et al., 2020). BMDMs on day 6 of differentiation

were plated at 4 x 10^5 cells/well in a 6-well plate for western blot analyses or 5 x 10^4 cells/well in

a 96-well plate for LDH assays. Cells were infected at MOI:25 for western blot or MOI:10 or 50

for LDH assays in antibiotic-free DMEM for denoted incubation times. For all experimental

analyses, following 30-minute incubation, cells were washed and replaced in DMEM media

supplemented with 50 µg/ml gentamycin to ensure growth inhibition of extracellular bacteria.

BMDM cell death levels were measured as a percentage of LDH release using the Promega

CytoTox 96 Non-Radioactive Cytotoxicity Assay (G1780), according to manufacturer's

instructions.

Sublethal irradiation

Mice were irradiated with a 5.5 Gy sub-lethal dose of γ-irradiation and received neomycin (2

mg/mL) in the drinking water for 3 weeks. At 4 days post irradiation, a retro-orbital bleed was

analysed via ADVIA hematology to confirm successful irradiation. Mice had submandibular

bleeds analysed by ADVIA hematology and had long bones harvested for flow cytometry analysis

at either 7-, 14- or 21-days post irradiation to assess stem cell capacity.

Hematopoietic stem cell transplants

Donor bone marrow were injected intravenously into recipient *C57BL/6-CD45$^{ly5.1}$* mice following

11 Gy of γ-irradiation split over two equal doses. Recipient mice received neomycin (2 mg/mL) in

the drinking water for 3 weeks. Long term capacity of stem cells was assessed by flow cytometric

analysis of donor contribution to recipient mouse peripheral blood at 6 weeks.

Zymosan induced peritonitis

1 mg of zymosan A from *Saccharomyces cerevisiae* (Sigma-Aldrich) was intra-peritoneally

injected into mice to induce sterile peritonitis. After 4 or 24 hours, mice were euthanized, cardiac

bled and bone marrow collected. The peritoneal cavity was washed with 1 ml of cold PBS and

cells within the lavage fluid were collected by centrifugation (300 x g, 5 minutes). Bone marrow

and peritoneum immune cells were quantified by flow cytometry.

Flow cytometry

To analyse the innate and adaptive immune cells in peripheral blood, inguinal lymph nodes, spleen

and bone marrow, isolated single cell suspensions were incubated with a combination of the

following antibodies: CD4-BV421, CD8-PECy7, CD19-PerCPCy5.5, CD11b-BV510 or BV786,

GR1-PE, CD45-Alexa700, Ly6G-V450 and Ly6C-APCCy7.

Splenic immune cells were analysed at 14-day post *Salmonella* infection and incubated with a

combination of the following antibodies: CD4-BV421, CD8-PeCy7, CD19-PerCPCy5.5, CD11b-

BV510 or BV786, CD64-PE, CD45-Alexa700, Ly6G-V450 and Ly6C-APCCy7.

To analyse the contribution of donor and competitor cells in transplanted recipients, blood cells

were incubated with either CD41-APC (ThermoFisher) and Ter119-PE or Ly5.1-A700, Ly5.2-PE-

Cy7, CD4-PE, CD8-PE, B220-BV650, Ly6C-APCCy7, Mac1-PerCPCy5.5. To analyse stem- and

progenitor- cell compartment following sub-lethal irradiation, bone marrow cells were incubated

with cKit-PerCPe710 (ThermoFisher), CD48-PECy7 (ThermoFisher), CD150-A647, Sca1-

APCCy7 (ThermoFisher), B220-PE, CD19-PE, CD4-PE, CD8-PE, Gr1-PE. Finally, fluorogold

(AAT Bioquest Cat#17514) was added for dead cell detection where appropriate.

Peritoneal and bone marrow immune cells at 4- or 24- hours post-zymosan injection were

incubated with CD45-Alexa700, CD64-BV650, Ly6G-PE, Ly6C-APCCy7, F4/80-PerCPCy5.5,

CD11b-BV510, CD8-PECy7, CD4-FITC, B220-APC and FC-blocker. Finally, propidium iodide

(2 µg/mL, Sigma-Aldrich) was added for dead cell detection.

All cells were analysed on the Aurora Cytex flow cytometer. With exception of zymosan induced

peritonitis experiments that used the Aurora Cytex automated volume calculator, cells were mixed

with counting beads to quantify absolute cell numbers. Except where denoted, all flow cytometry

antibodies were obtained from BD Biosciences.

Statistical analyses

All data points signify independent experimental repeats or biologically independent data points.

All p values were calculated in Prism using the statistical test identified in figure legends. Asterisks

signify that $p \leq 0.05$ (*), $p \leq 0.01$ (**) or $p \leq 0.001$ (***).

 FIGURE LEGENDS

 Figure 1. MLKL^{S132P} executes cell death in the presence of necrosulfonamide or Ser83

 inhibition.

(A) Minor allele frequency of *MLKL p.Ser132Pro* according to the gnomAD database, stratified

by ancestry. **(B)** Sequence alignment of conserved serine in the MLKL orthologs across different

species. **(C)** Family pedigree of patient 2, identified from Australian registry of patients suffering

from immune related disease. Known diagnoses for family members are indicated. **(D)** MLKL^{WT}

and MLKL^{S132P} reconstitute necroptotic signaling, but as expected control protein harboring

activation loop phosphomimetic substitutions, MLKL^{TSEE}, did not. Human MLKL expression was

induced in *MLKL$^{-/-}$* HT29 cells with 100 ng/ml doxycycline (Dox) and treated with necroptotic

stimulus (TNF, SMAC mimetic, IDN-6556; TSI) in the presence or absence of MLKL inhibitor

Necrosulfonamide (NSA; 1μM). Cell death was measured every hour for 24 hours by percentage

of SYTOX Green positive cells quantified using IncuCyte SX5 live cell imaging. Independent cell

lines were assayed in *n=3-9* experiments, with errors bars indicating the mean ± SEM. **(E)** Western

blot analyses of whole cell lysates taken 3 h post TSI stimulation in the presence or absence of

NSA from doxycycline (100 ng/ml) induced *MLKL$^{-/-}$* HT29 cells expressing *MLKL^{WT}* or

MLKL^{S132P}. **(F)** High molecular weight phosphorylated MLKL^{S132P} is present at enhanced levels

in crude membrane fractions following Blue-Native PAGE under TSI stimulation in the presence

or absence of NSA. **(G)** Ser83 (red), Cys86 (purple) and Ser132 (gold) highlighted as spheres on

cartoon representation of human MLKL. Four-helix bundle domain shown in green, brace helices

shown in beige and pseudokinase domain shown in blue. Homology model is generated from the

N-terminal domain NMR structure (PDB:2MSV) and the C-terminal pseudokinase domain crystal

structure (PDB: 4MWI) of human MLKL, which were aligned using the full-length murine crystal

structure (PDB:4BTF) (Murphy et al., 2013; Murphy et al., 2014; Su et al., 2014). **(H)** *MLKL*

S132P mutation reconstitutes necroptotic signaling in the presence of Ser83 inhibitory

phosphorylation (MLKL^{S83D}). Human MLKL expression was induced in *MLKL$^{-/-}$* HT29 cells with

100 ng/ml doxycycline (Dox) and treated with necroptotic stimulus (TNF, SMAC mimetic, IDN-

6556; TSI) in the presence or absence of MLKL inhibitor Necrosulfonamide (NSA; 1μM). Cell

death was measured every hour for 23 hours by percentage of SYTOX Green positive cells

quantified using IncuCyte SX5 live cell imaging. Independent cell lines were assayed in *n=4*

experiments, with errors bars indicating the mean ± SEM. **(E/F)** Blot images are representative of

at least three independent experiments.

Figure 2. In mice, *Mlkl^{S131P}* homozygosity is tolerated but results in steady state immune cell

population differences.

(A) Macroscopic appearance of *Mlkl$^{WT/WT}$* and *Mlkl$^{S131P/S131P}$* mice at 8-12 weeks of age. **(B)**

Mlkl$^{S131P/S131P}$ mice are born according to Hardy-Weinberg equilibrium as observed in the

distribution of genotypes from *Mlkl$^{WT/S131P}$* heterozygous intercrosses. **(C)** Body weight of

Mlkl$^{WT/WT}$, *Mlkl$^{WT/S131P}$* and *Mlkl$^{S131P/S131P}$* mice at 3-4 or 7-9 months of age. Each dot represents an

individual mouse (*n= 8-17,* mean ± SEM). **(D-F)** Flow cytometry quantification of innate (Ly6C^{hi}

monocytes, Ly6C^{lo} monocytes and neutrophils) and adaptive (CD4$^+$ T cells, CD8$^+$ T cells and B

cells) in the bone marrow **(D)**, spleen **(E)** and inguinal lymph nodes **(F)** of 8–12-week-old

Mlkl$^{WT/WT}$, Mlkl$^{WT/S131P}$, and *Mlkl$^{S131P/S131P}$* mice. Each symbol represents one individual mouse

sampled and error bars represent mean ± SEM for *n=7-11* mice as indicated. **(G, H)** Immortalized

mouse dermal fibroblasts (MDFs) were isolated from *Mlkl$^{WT/WT}$, Mlkl$^{WT/S131P}$,* and *Mlkl$^{S131P/S131P}$*

mice and stimulated as indicated for 6 hours for western blot analysis **(G)** or 24 hours for

quantification of PI-positive cells using IncuCyte S3 live cell imaging **(H)**. Death data represent

mean ± SEM for independently generated cell lines of n=3-5. *p<0.05, **p<0.01 calculated using

an unpaired, two-tailed Students t-test.

Figure 3. $Mlkl^{S131P}$ mice show delayed recovery from myelosuppressive irradiation.

Peripheral red blood cells **(A)**, platelets **(B)** and monocytes **(C)** in $Mlkl^{WT/WT}$, $Mlkl^{WT/S131P}$, and

$Mlkl^{S131P/S131P}$ mice following treatment with 5.5 Gy radiation. Quantified nucleated viable cells

(D), progenitor **(E)** and LSK **(F)** populations in the bone marrow of mice after myelosuppressive

radiation. Mean ± SEM of n=3-10 independent mice from two separate experiments. Bone marrow

from $Mlkl^{WT/WT}$ or $Mlkl^{S131P/S131P}$ mice on CD45$^{Ly5.2}$ background was mixed with wild-type GFP$^+$

competitor bone marrow on a CD45$^{Ly5.2}$ background and transplanted into irradiated CD45$^{Ly5.1}$

recipients. Relative donor contribution to PBMCs **(G)**, red blood cells **(H)** and platelets **(I)** was

assessed at 6 weeks post-transplantation. Mean shown of 2 donors per genotype, and 2-3 recipients

per donor. Host contribution (CD45$^{Ly5.1}$) depicted in pink, GFP competitor in green and test

($Mlkl^{WT/WT}$ or $Mlkl^{S131P/S131P}$) in purple. *p<0.05, **p<0.01, ***p<0.001 calculated using an

unpaired, two-tailed Students t-test **(A-F)**.

Figure 4. $Mlkl^{S131P}$ recruited inflammatory neutrophils are less sensitive to TNF-induced necroptosis.

(A-D) Flow cytometry quantification of peritoneal innate (macrophages, Ly6C^{hi} monocytes,

Ly6C^{lo} monocytes and neutrophils) and adaptive (CD4$^+$ T cells, CD8$^+$ T cells and B cells) immune

cells at 4- **(A, B)** or 24- **(C, D)** hours post zymosan injection. ADVIA hematology quantification

of circulating immune cells (lymphocytes, neutrophils and monocytes) at 4- **(E)** and 24- **(F)** hours

post zymosan injection. Each symbol represents one independent animal, with mice from the 4- or

24-hour timepoint pooled from 3 and 2 independent zymosan experiments respectively. Error bars

represent mean ± SEM for *n=5-14* mice as indicated **(A-H)**. Evaluation of spontaneous apoptotic **(G)** or induced necroptotic signaling **(H)** in neutrophils recruited and isolated from the peritoneum 4 hours post-zymosan injection. Neutrophils were left unstimulated (spontaneous apoptosis) or treated with necroptotic stimulus (TNF, SMAC mimetic, IDN-6556; TSI) for 16 hours and cell death was measured every hour by percentage of SYTOX Green positive cells quantified using IncuCyte SX5 live cell imaging. Data were collected from one independent experiment with male and female data pooled, neutrophils isolated from independent mice with mean ± SEM of *n=6-14* presented. ***p<0.001 calculated using an unpaired, two-tailed Students t-test.

Figure 5. *Mlkl^{S131P}* homozygote mice exhibit bacterial clearance defects following oral *Salmonella* infection.

(A) ZO-1 (yellow) and CD68 (purple) staining in epithelial barrier of intestinal sections taken at 14-days post *Salmonella* infection. **(B)** Increased bacterial burden observed in spleen and liver, but not feces in *Mlkl$^{S131P/S131P}$* mice at infection endpoint. Quantification of circulating white blood cells (lymphocytes, neutrophils and monocytes) using ADVIA hematology **(C)** and splenic adaptive (CD4$^+$ T cells, CD8$^+$ T cells and B cells) and innate (macrophages, Ly6C^{hi} monocytes, Ly6C^{lo} monocytes and neutrophils) immune cells using flow cytometry **(D, E)**. *Salmonella* infection was performed on 3 independent occasions, with each symbol representing an individual mouse sampled. Error bars represent mean ± SEM for *n=9-16* mice as indicated. *p<0.05, **p<0.01, ***p<0.001 calculated using a Mann-Whitney test **(B)** or an unpaired, two-tailed Students t-test **(C-E)**.

Supplementary Figure 1. MLKL^{S132P} is less sensitive to inhibition by necrosulfonamide.

(**A**) Peripheral blood mononuclear cells (PBMCs) isolated from patient 2 and an aged matched

healthy donor control were stimulated as indicated for 4 hours for western blot analysis. (**B**) ELISA

measurement of supernatant TNF in PBMCs stimulated with LPS or Poly I:C for 5 hours. Mean

of technical triplicates presented. (**C-F**) Evaluation of necroptotic signaling by $MLKL^{WT}$,

$MLKL^{S132P}$ and $MLKL^{TSEE}$ in *MLKL$^{-/-}$* (**C**) or WT (**D**) HT29 cells and WT (**E**) or *MLKL$^{-/-}$* (**F**)

U937 cells. Human MLKL expression was induced with doxycycline (Dox) and treated with

necroptotic stimulus (TNF, SMAC mimetic, IDN-6556; TSI) in the presence or absence of MLKL

inhibitor necrosulfonamide (NSA). Cell death was measured every hour for 24 hours by percentage

of SYTOX Green positive cells quantified using IncuCyte SX5 or S3 live cell imaging.

Independent cell lines were assayed in *n=2-9* experiments, with errors bars indicating the mean ±

SEM. (**G, H**) Western blot analyses of whole cell lysates of doxycycline induced WT or *MLKL$^{-/-}$*

HT29 (**G**) or U937 (**H**) cells expressing *MLKL^{WT}*, *MLKL^{S132P}* or *MLKL^{TSEE}*. (**I**) Blue-Native PAGE

crude cytoplasm fractions of *MLKL$^{-/-}$* HT29 cells under TSI stimulation (0-6 hours) in the presence

or absence of NSA. (**J**) Western blot analyses of whole cell lysates taken 5 h post TSI stimulation

in the presence or absence of NSA from doxycycline induced *MLKL$^{-/-}$* HT29 cells expressing

MLKL^{WT}, *MLKL^{S132P}*, *MLKL^{S83D}* or *MLKL^{S83D,S132P}*. (**K, L**) Evaluation of necroptotic signaling in

MLKL$^{-/-}$ (**K**) or WT (**L**) HT29 cells expressing *MLKL^{WT}*, *MLKL^{S132P}*, *MLKL^{S83D}* and

MLKL^{S83D,S132P}. Human MLKL expression was induced with doxycycline (Dox) and cells treated

with necroptotic stimulus (TNF, SMAC mimetic, IDN-6556; TSI) in the presence or absence of

MLKL inhibitor necrosulfonamide (NSA). Cell death was measured every hour for 23 hours by

percentage of SYTOX-green positive cells quantified using IncuCyte SX5 or S3 live cell imaging.

Independent cell lines were assayed in *n=4* experiments, with errors bars indicating the mean ±

SEM. Blots images in **A, G, H, I & J** are representative images of at least two independent repeat

experiments.

Supplementary Figure 2. Cells expressing endogenous MLKL^{S131P} exhibit reduced TSI-

induced necroptotic cell death.

(A) Representative images of H&E staining of salivary glands, mediastinum, liver and lung from

7–9-month-old *Mlkl$^{WT/WT}$* and *Mlkl$^{S131P/S131P}$* mice. Images are representative of *n=2* mice per

genotype. **(B-D)** ADVIA hematology quantification of circulating red blood cells **(B)**, platelets

(C) and white blood cells **(D)** in *Mlkl$^{WT/WT}$*, *Mlkl$^{WT/S131P}$* and *Mlkl$^{S131P/S131P}$* mice across age. Each

symbol represents one independent mouse sampled and error bars represent mean ± SEM for

n=12-38 mice as indicated. **(E-V)** Flow cytometry quantification of innate (Ly6C^{hi}, Ly6C^{lo} and

neutrophils) and adaptative (B cells, CD4$^+$ T cells and CD8$^+$ T cells) immune cells in the bone

marrow **(E-J)**, spleen **(K-P)** and inguinal lymph nodes **(Q-V)** of 8–12-week-old basal state

Mlkl$^{WT/WT}$, *Mlkl$^{WT/S131P}$*, and *Mlkl$^{S131P/S131P}$* mice as indicated. Each symbol represents one

independent mouse sampled and error bars represent mean ± SEM for *n=7-11* mice as indicated.

(W, X) Primary mouse dermal fibroblasts (MDFs) were isolated from *Mlkl$^{WT/WT}$*, *Mlkl$^{WT/S131P}$*, and

Mlkl$^{S131P/S131P}$ mice and stimulated as indicated for 6 hours for western blot analysis **(W)** or 20

hours for quantification of PI-positive cells using IncuCyte S3 live cell imaging **(X)**. **(Y, Z)** Bone

marrow derived macrophages were isolated from *Mlkl$^{WT/WT}$*, *Mlkl$^{WT/S131P}$*, and *Mlkl$^{S131P/S131P}$* mice

and stimulated on day 6 of culture as indicated for 6 hours for western blot analysis **(Y)** or 24 hours

for quantification of PI-positive cells using IncuCyte S3 live cell imaging **(Z)**. *p<0.05 **p<0.01

calculated using an unpaired, two-tailed Students t-test.

Supplementary Figure 3. *Mlkl^S131P/S131P* hematopoietic stem cells transplanted in excess are

outcompeted at 6-weeks post-transplant.

Bone marrow from *Mlkl^S131P/S131P* mice on CD45^Ly5.2 background was mixed with wild-type GFP⁺

competitor bone marrow on a CD45^Ly5.2 background at a 70:30 ratio and transplanted into

irradiated CD45^Ly5.1 recipients. Relative donor contribution to PBMCs **(A)**, red blood cells **(B)** and

platelets **(C)** was assessed at 6 weeks post-transplantation. Mean ± SEM shown of 2 donors per

genotype, and 3 recipients per donor. Host contribution (CD45^Ly5.1) depicted in pink, GFP

competitor in green and test *Mlkl^S131P/S131P* in purple.

Supplementary Figure 4. *Mlkl^S131P* mice have reduced levels of Ly6C^hi monocytes in the

peritoneal lavage at 24-hours post zymosan injection.

(A-P) Flow cytometry quantification of peritoneal innate (macrophages, Ly6C^hi, Ly6C^lo and

neutrophils) and adaptive (CD4⁺ T cells, CD8⁺ T cells and B cells) immune cells at 4- **(A-H)** or

24- **(I-P)** hours post-intraperitoneal injection of zymosan as indicated. **(Q-V)** ADVIA hematology

quantification of peripheral blood cells (lymphocytes, neutrophils, and monocytes) at 4- **(Q-S)** and

24- **(T-V)** post-injection of zymosan. Each symbol represents one independent animal, with mice

from the 4- or 24-hour timepoint pooled from 3 and 2 independent zymosan experiments

respectively. Error bars represent mean ± SEM for *n=2-10* mice as indicated. *p<0.05 **p<0.01

calculated using an unpaired, two-tailed Students t-test.

Supplementary Figure 5. *Mlkl^S131P* does not abrogate death of BMDMs upon *in vitro*

***Salmonella* infection.**

(**A, B**) *Mlkl^{WT/WT}, Mlkl^{WT/S131P},* and *Mlkl^{S131P/S131P}* mice were infected with *Salmonella* via oral gavage and monitored for 14-days by daily body weight (**A**) and temperature (**B**) measurements. (**C-E**) Bacterial burden calculation of *Salmonella* colonization in the spleen (**C**), liver (**D**), and feces (**E**) at experimental endpoint. (**F-H**) ADVIA hematology quantification of peripheral monocytes (**F**), lymphocytes (**G**) and neutrophils (**H**). (**I-O**) Splenic adaptive (CD4$^+$ T cells, CD8$^+$ T cells and B cells) (**I-K**) and innate (macrophages, Ly6C^{hi}, Ly6C^{lo} and neutrophils) (**L-O**) immune cells were quantified by flow cytometry at experimental endpoint. *Salmonella* infection was completed 3 independent times, with each symbol representing an individual mouse sampled. Error bars represent mean ± SEM for *n=3-9* mice as indicated. (**P, Q**) *In vitro* assessment of *Salmonella* SL1344 infection of primary BMDMs generated from *Mlkl^{WT/WT}, Mlkl^{WT/S131P},* and *Mlkl^{S131P/S131P}* mice. (**G**) BMDMs were infected with *Salmonella* (MOI:25) and cleavage associated with Gasdermin-D activation during pyroptosis was analyzed by immunoblotting at the indicated time points. (**H**) LDH release cell death assay of BMDMs after infection with *Salmonella* (MOI:10 or MOI:50) at indicated time points. *In vitro Salmonella* experiment completed once, with mean ± SEM for *n=3* individual mice shown. Blot images in (**G**) are representative of independent duplicates.

ACKNOWLEDGEMENTS

We thank all the following people for their technical assistance; Aira Nuguid and Tina Cardamone (Phenomics Australia Histopathology and Slide Scanning Service- The University of Melbourne). WEHI Cytometry Facility, WEHI Antibody Facility, WEHI Centre for Dynamic Imaging, WEHI Bioservices, Cheree Fitzgibbon (WEHI), Jacinta Hansen (WEHI) and Matthew Cook (ANU). The generation of *Mlkl^{S131P}* mice by CRISPR/Cas9 gene editing was performed by Andrew Kueh and

Marco Herold (WEHI MAGEC laboratory) supported by the Australian Phenomics Network

(APN) and the Australian Government through the National Collaborative Research Infrastructure

Strategy (NCRIS) program. We thank Warren Alexander, Melanie Bahlo and Ian Wicks for the

provision of important resources and expertise. We are grateful to the National Health and Medical

Research Council for fellowship (J.M.H., 1142669; A.L.S., 2002965; J.M.M., 1172929; J.S.,

1107149), grant (J.M.M., 1105023; K.R.M., 1092602; J.S., 1105023; J.M.H., 2011584) and

infrastructure (IRIISS 9000719); Arthritis Australia support to K.R.M. We acknowledge

scholarship support for S.E.G and D.F (Australian Government Research Training Program

Stipend Scholarships), S.E.G (Wendy Dowsett Scholarship), S.C (Walter and Eliza Hall Handman

PhD Scholarship). Victorian State Government Operational Infrastructure Support Scheme.

AUTHOR INFORMATION

The Walter and Eliza Hall Institute of Medical Research, Parkville, VIC, 3052, Australia

Sarah E. Garnish, Katherine R. Martin, Maria Kauppi, Victoria Jackson, Daniel Frank, Emma C.

Tovey Crutchfield, Komal M. Patel, Shene Chiou, Andre L. Samson, Marcel Doerflinger,

Christopher R. Horne, Cathrine Hall, Samuel N. Young, Charlotte Slade, John Silke, James M.

Murphy and Joanne M. Hildebrand

Department of Medical Biology, University of Melbourne, Parkville, VIC, 3052, Australia

Sarah E. Garnish, Katherine R. Martin, Maria Kauppi, Victoria Jackson, Daniel Frank, Shene

Chiou, Andre L. Samson, Charlotte Slade, John Silke, James M. Murphy and Joanne M.

Hildebrand

Centre for Innate Immunity and Infectious Diseases, Hudson Institute of Medical Research, Clayton, VIC, 3168, Australia

Rebecca Ambrose, Vik Ven Eng and Jaclyn S. Pearson

Department of Molecular and Translational Science, Monash University, Clayton, VIC, 3168, Australia

Rebecca Ambrose, Vik Ven Eng and Jaclyn S. Pearson

Department of Microbiology, Monash University, Clayton, VIC, 3168, Australia

Jaclyn S. Pearson

Department of Immunology and Infection, John Curtin School of Medical Research, Australian National University, ACT, Australia

Vicki Athanasopoulos and Carola Vinuesa

Department of Clinical Immunology and Allergy, Royal Melbourne Hospital, Parkville, VIC, 3052, Australia

Charlotte Slade

Institute of Virology, Technical University of Munich/Helmholtz Munich, Munich, Germany

Gregor Ebert

Contributions

Conceptualization: S.E.G., J.S., J.M.M., J.M.H. Methodology: S.E.G., K.R.M., M.K., V.J., R.A., V.E., D.F., S.C., A.L.S., M.D., C.R.H., C.H., S.N.Y., J.S.P., G.E., J.M.H. Resources: K.R.M., M.K., V.A., C.V., C.S., J.S.P., J.S., J.M.M., J.M.H. Supervision: K.M., M.K., J.S., J.M.M., J.M.H. Funding acquisition: J.S., J.M.M., J.M.H. S.E.G and J.M.H co-wrote the paper with input from authors.

ETHICS DECLARATIONS

Competing interests: S.E.G, K.M.P, A.L.S, C.R.H, S.N.Y, J.S, J.M.M and J.M.H contribute, or have contributed, to a project developing necroptosis inhibitors in collaboration with Anaxis Pty Ltd. K.R.M received funding from CSL Pty Ltd. The remaining authors declare no competing interests.

References

Boisson-Dupuis, S., N. Ramirez-Alejo, Z. Li, E. Patin, G. Rao, G. Kerner, C.K. Lim, D.N. Krementsov, N.
    Hernandez, C.S. Ma, Q. Zhang, J. Markle, R. Martinez-Barricarte, K. Payne, R. Fisch, C. Deswarte,
    J. Halpern, M. Bouaziz, J. Mulwa, D. Sivanesan, T. Lazarov, R. Naves, P. Garcia, Y. Itan, B. Boisson,

A. Checchi, F. Jabot-Hanin, A. Cobat, A. Guennoun, C.C. Jackson, S. Pekcan, Z. Caliskaner, J.
Inostroza, B.T. Costa-Carvalho, J.A.T. de Albuquerque, H. Garcia-Ortiz, L. Orozco, T. Ozcelik, A.
Abid, I.A. Rhorfi, H. Souhi, H.N. Amrani, A. Zegmout, F. Geissmann, S.W. Michnick, I. Muller-
Fleckenstein, B. Fleckenstein, A. Puel, M.J. Ciancanelli, N. Marr, H. Abolhassani, M.E. Balcells, A.
Condino-Neto, A. Strickler, K. Abarca, C. Teuscher, H.D. Ochs, I. Reisli, E.H. Sayar, J. El-Baghdadi, J.
Bustamante, L. Hammarstrom, S.G. Tangye, S. Pellegrini, L. Quintana-Murci, L. Abel, and J.L.
Casanova. 2018. Tuberculosis and impaired IL-23-dependent IFN-gamma immunity in humans
homozygous for a common TYK2 missense variant. *Sci Immunol* 3:
Boomer, J.S., K. To, K.C. Chang, O. Takasu, D.F. Osborne, A.H. Walton, T.L. Bricker, S.D. Jarman, 2nd, D.
Kreisel, A.S. Krupnick, A. Srivastava, P.E. Swanson, J.M. Green, and R.S. Hotchkiss. 2011.
Immunosuppression in patients who die of sepsis and multiple organ failure. *JAMA* 306:2594-
2605.
Bossen, C., K. Ingold, A. Tardivel, J.L. Bodmer, O. Gaide, S. Hertig, C. Ambrose, J. Tschopp, and P. Schneider.
2006. Interactions of tumor necrosis factor (TNF) and TNF receptor family members in the mouse
and human. *J Biol Chem* 281:13964-13971.
Brinkworth, J.F., and N. Valizadegan. 2021. Sepsis and the evolution of human increased sensitivity to
lipopolysaccharide. *Evol Anthropol* 30:141-157.
Cai, Z., S. Jitkaew, J. Zhao, H.C. Chiang, S. Choksi, J. Liu, Y. Ward, L.G. Wu, and Z.G. Liu. 2014. Plasma
membrane translocation of trimerized MLKL protein is required for TNF-induced necroptosis. *Nat
Cell Biol* 16:55-65.
Chen, X., W. Li, J. Ren, D. Huang, W.T. He, Y. Song, C. Yang, W. Li, X. Zheng, P. Chen, and J. Han. 2014.
Translocation of mixed lineage kinase domain-like protein to plasma membrane leads to necrotic
cell death. *Cell research* 24:105-121.
Cortes, A., J. Hadler, J.P. Pointon, P.C. Robinson, T. Karaderi, P. Leo, K. Cremin, K. Pryce, J. Harris, S. Lee,
K.B. Joo, S.C. Shim, M. Weisman, M. Ward, X. Zhou, H.J. Garchon, G. Chiocchia, J. Nossent, B.A.
Lie, O. Forre, J. Tuomilehto, K. Laiho, L. Jiang, Y. Liu, X. Wu, L.A. Bradbury, D. Elewaut, R. Burgos-
Vargas, S. Stebbings, L. Appleton, C. Farrah, J. Lau, T.J. Kenna, N. Haroon, M.A. Ferreira, J. Yang, J.
Mulero, J.L. Fernandez-Sueiro, M.A. Gonzalez-Gay, C. Lopez-Larrea, P. Deloukas, P. Donnelly, C.
Australo-Anglo-American Spondyloarthritis, S. Groupe Francaise d'Etude Genetique des, S. Nord-
Trondelag Health, C. Spondyloarthritis Research Consortium of, C. Wellcome Trust Case Control,
P. Bowness, K. Gafney, H. Gaston, D.D. Gladman, P. Rahman, W.P. Maksymowych, H. Xu, J.B.
Crusius, I.E. van der Horst-Bruinsma, C.T. Chou, R. Valle-Onate, C. Romero-Sanchez, I.M. Hansen,
F.M. Pimentel-Santos, R.D. Inman, V. Videm, J. Martin, M. Breban, J.D. Reveille, D.M. Evans, T.H.
Kim, B.P. Wordsworth, and M.A. Brown. 2013. Identification of multiple risk variants for
ankylosing spondylitis through high-density genotyping of immune-related loci. *Nat Genet*
45:730-738.
Cortes, A., S.L. Pulit, P.J. Leo, J.J. Pointon, P.C. Robinson, M.H. Weisman, M. Ward, L.S. Gensler, X. Zhou,
H.J. Garchon, G. Chiocchia, J. Nossent, B.A. Lie, O. Forre, J. Tuomilehto, K. Laiho, L.A. Bradbury, D.
Elewaut, R. Burgos-Vargas, S. Stebbings, L. Appleton, C. Farrah, J. Lau, N. Haroon, J. Mulero, F.J.
Blanco, M.A. Gonzalez-Gay, C. Lopez-Larrea, P. Bowness, K. Gaffney, H. Gaston, D.D. Gladman, P.
Rahman, W.P. Maksymowych, J.B. Crusius, I.E. van der Horst-Bruinsma, R. Valle-Onate, C.
Romero-Sanchez, I.M. Hansen, F.M. Pimentel-Santos, R.D. Inman, J. Martin, M. Breban, B.P.
Wordsworth, J.D. Reveille, D.M. Evans, P.I. de Bakker, and M.A. Brown. 2015. Major
histocompatibility complex associations of ankylosing spondylitis are complex and involve further
epistasis with ERAP1. *Nat Commun* 6:7146.
Dondelinger, Y., W. Declercq, S. Montessuit, R. Roelandt, A. Goncalves, I. Bruggeman, P. Hulpiau, K.
Weber, C.A. Sehon, R.W. Marquis, J. Bertin, P.J. Gough, S. Savvides, J.C. Martinou, M.J. Bertrand,

and P. Vandenabeele. 2014. MLKL compromises plasma membrane integrity by binding to
phosphatidylinositol phosphates. *Cell reports* 7:971-981.
Evans, D.M., C.C. Spencer, J.J. Pointon, Z. Su, D. Harvey, G. Kochan, U. Oppermann, A. Dilthey, M. Pirinen,
M.A. Stone, L. Appleton, L. Moutsianas, S. Leslie, T. Wordsworth, T.J. Kenna, T. Karaderi, G.P.
Thomas, M.M. Ward, M.H. Weisman, C. Farrar, L.A. Bradbury, P. Danoy, R.D. Inman, W.
Maksymowych, D. Gladman, P. Rahman, C. Spondyloarthritis Research Consortium of, A. Morgan,
H. Marzo-Ortega, P. Bowness, K. Gaffney, J.S. Gaston, M. Smith, J. Bruges-Armas, A.R. Couto, R.
Sorrentino, F. Paladini, M.A. Ferreira, H. Xu, Y. Liu, L. Jiang, C. Lopez-Larrea, R. Diaz-Pena, A. Lopez-
Vazquez, T. Zayats, G. Band, C. Bellenguez, H. Blackburn, J.M. Blackwell, E. Bramon, S.J.
Bumpstead, J.P. Casas, A. Corvin, N. Craddock, P. Deloukas, S. Dronov, A. Duncanson, S. Edkins, C.
Freeman, M. Gillman, E. Gray, R. Gwilliam, N. Hammond, S.E. Hunt, J. Jankowski, A. Jayakumar, C.
Langford, J. Liddle, H.S. Markus, C.G. Mathew, O.T. McCann, M.I. McCarthy, C.N. Palmer, L.
Peltonen, R. Plomin, S.C. Potter, A. Rautanen, R. Ravindrarajah, M. Ricketts, N. Samani, S.J.
Sawcer, A. Strange, R.C. Trembath, A.C. Viswanathan, M. Waller, P. Weston, P. Whittaker, S.
Widaa, N.W. Wood, G. McVean, J.D. Reveille, B.P. Wordsworth, M.A. Brown, P. Donnelly, C.
Australo-Anglo-American Spondyloarthritis, and C. Wellcome Trust Case Control. 2011.
Interaction between ERAP1 and HLA-B27 in ankylosing spondylitis implicates peptide handling in
the mechanism for HLA-B27 in disease susceptibility. *Nat Genet* 43:761-767.
Faergeman, S.L., H. Evans, K.E. Attfield, C. Desel, S.B. Kuttikkatte, M. Sommerlund, L.T. Jensen, J. Frokiaer,
M.A. Friese, P.M. Matthews, C. Luchtenborg, B. Brugger, A.B. Oturai, C.A. Dendrou, and L. Fugger.
2020. A novel neurodegenerative spectrum disorder in patients with MLKL deficiency. *Cell death
& disease* 11:303.
Galan, J.E. 2021. Salmonella Typhimurium and inflammation: a pathogen-centric affair. *Nat Rev Microbiol*
19:716-725.
Garnish, S.E., and J.M. Hildebrand. 2022. Rare catastrophes and evolutionary legacies: human germline
gene variants in MLKL and the necroptosis signalling pathway. *Biochem Soc Trans* 50:529-539.
Garnish, S.E., Y. Meng, A. Koide, J.J. Sandow, E. Denbaum, A.V. Jacobsen, W. Yeung, A.L. Samson, C.R.
Horne, C. Fitzgibbon, S.N. Young, P.P.C. Smith, A.I. Webb, E.J. Petrie, J.M. Hildebrand, N. Kannan,
P.E. Czabotar, S. Koide, and J.M. Murphy. 2021. Conformational interconversion of MLKL and
disengagement from RIPK3 precede cell death by necroptosis. *Nat Commun* 12:2211.
Gutierrez-Arcelus, M., S.S. Rich, and S. Raychaudhuri. 2016. Autoimmune diseases - connecting risk alleles
with molecular traits of the immune system. *Nat Rev Genet* 17:160-174.
He, S., Y. Liang, F. Shao, and X. Wang. 2011. Toll-like receptors activate programmed necrosis in
macrophages through a receptor-interacting kinase-3-mediated pathway. *Proc Natl Acad Sci U S
A* 108:20054-20059.
He, S., L. Wang, L. Miao, T. Wang, F. Du, L. Zhao, and X. Wang. 2009. Receptor interacting protein kinase-
3 determines cellular necrotic response to TNF-alpha. *Cell* 137:1100-1111.
Hildebrand, J.M., M. Kauppi, I.J. Majewski, Z. Liu, A.J. Cox, S. Miyake, E.J. Petrie, M.A. Silk, Z. Li, M.C. Tanzer,
G. Brumatti, S.N. Young, C. Hall, S.E. Garnish, J. Corbin, M.D. Stutz, L. Di Rago, P. Gangatirkar, E.C.
Josefsson, K. Rigbye, H. Anderton, J.A. Rickard, A. Tripaydonis, J. Sheridan, T.S. Scerri, V.E. Jackson,
P.E. Czabotar, J.G. Zhang, L. Varghese, C.C. Allison, M. Pellegrini, G.M. Tannahill, E.C. Hatchell, T.A.
Willson, D. Stockwell, C.A. de Graaf, J. Collinge, A. Hilton, N. Silke, S.K. Spall, D. Chau, V.
Athanasopoulos, D. Metcalf, R.M. Laxer, A.G. Bassuk, B.W. Darbro, M.A. Fiatarone Singh, N.
Vlahovich, D. Hughes, M. Kozlovskaia, D.B. Ascher, K. Warnatz, N. Venhoff, J. Thiel, C. Biben, S.
Blum, J. Reveille, M.S. Hildebrand, C.G. Vinuesa, P. McCombe, M.A. Brown, B.T. Kile, C. McLean,
M. Bahlo, S.L. Masters, H. Nakano, P.J. Ferguson, J.M. Murphy, W.S. Alexander, and J. Silke. 2020.
A missense mutation in the MLKL brace region promotes lethal neonatal inflammation and
hematopoietic dysfunction. *Nat Commun* 11:3150.

Hildebrand, J.M., B. Lo, S. Tomei, V. Mattei, S.N. Young, C. Fitzgibbon, J.M. Murphy, and A. Fadda. 2021. A
family harboring an MLKL loss of function variant implicates impaired necroptosis in diabetes. *Cell*
death & disease 12:345.
Hildebrand, J.M., M.C. Tanzer, I.S. Lucet, S.N. Young, S.K. Spall, P. Sharma, C. Pierotti, J.M. Garnier, R.C.
Dobson, A.I. Webb, A. Tripaydonis, J.J. Babon, M.D. Mulcair, M.J. Scanlon, W.S. Alexander, A.F.
Wilks, P.E. Czabotar, G. Lessene, J.M. Murphy, and J. Silke. 2014. Activation of the pseudokinase
MLKL unleashes the four-helix bundle domain to induce membrane localization and necroptotic
cell death. *Proceedings of the National Academy of Sciences of the United States of America*
111:15072-15077.
Hohmann, E.L. 2001. Nontyphoidal salmonellosis. *Clin Infect Dis* 32:263-269.
Holler, N., R. Zaru, O. Micheau, M. Thome, A. Attinger, S. Valitutti, J.L. Bodmer, P. Schneider, B. Seed, and
J. Tschopp. 2000. Fas triggers an alternative, caspase-8-independent cell death pathway using the
kinase RIP as effector molecule. *Nat Immunol* 1:489-495.
Huang, Z., A.H. Rose, F.W. Hoffmann, A.S. Hashimoto, P. Bertino, T. Denk, J. Takano, N. Iwata, T.C. Saido,
and P.R. Hoffmann. 2013. Calpastatin prevents NF-kappaB-mediated hyperactivation of
macrophages and attenuates colitis. *J Immunol* 191:3778-3788.
Kaiser, W.J., H. Sridharan, C. Huang, P. Mandal, J.W. Upton, P.J. Gough, C.A. Sehon, R.W. Marquis, J. Bertin,
and E.S. Mocarski. 2013. Toll-like receptor 3-mediated necrosis via TRIF, RIP3, and MLKL. *J Biol*
Chem 288:31268-31279.
Karczewski, K.J., L.C. Francioli, G. Tiao, B.B. Cummings, J. Alfoldi, Q. Wang, R.L. Collins, K.M. Laricchia, A.
Ganna, D.P. Birnbaum, L.D. Gauthier, H. Brand, M. Solomonson, N.A. Watts, D. Rhodes, M. Singer-
Berk, E.M. England, E.G. Seaby, J.A. Kosmicki, R.K. Walters, K. Tashman, Y. Farjoun, E. Banks, T.
Poterba, A. Wang, C. Seed, N. Whiffin, J.X. Chong, K.E. Samocha, E. Pierce-Hoffman, Z. Zappala,
A.H. O'Donnell-Luria, E.V. Minikel, B. Weisburd, M. Lek, J.S. Ware, C. Vittal, I.M. Armean, L.
Bergelson, K. Cibulskis, K.M. Connolly, M. Covarrubias, S. Donnelly, S. Ferriera, S. Gabriel, J.
Gentry, N. Gupta, T. Jeandet, D. Kaplan, C. Llanwarne, R. Munshi, S. Novod, N. Petrillo, D. Roazen,
V. Ruano-Rubio, A. Saltzman, M. Schleicher, J. Soto, K. Tibbetts, C. Tolonen, G. Wade, M.E.
Talkowski, C. Genome Aggregation Database, B.M. Neale, M.J. Daly, and D.G. MacArthur. 2020.
The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature*
581:434-443.
Karunakaran, D., A.W. Turner, A.C. Duchez, S. Soubeyrand, A. Rasheed, D. Smyth, D.P. Cook, M. Nikpay,
J.W. Kandiah, C. Pan, M. Geoffrion, R. Lee, L. Boytard, H. Wyatt, M.A. Nguyen, P. Lau, M. Laakso,
B. Ramkhelawon, M. Alvarez, K.H. Pietilainen, P. Pajukanta, B.C. Vanderhyden, P. Liu, S.B. Berger,
P.J. Gough, J. Bertin, M.E. Harper, A.J. Lusis, R. McPherson, and K.J. Rayner. 2020. RIPK1 gene
variants associate with obesity in humans and can be therapeutically silenced to reduce obesity
in mice. *Nat Metab* 2:1113-1125.
Kerner, G., G. Laval, E. Patin, S. Boisson-Dupuis, L. Abel, J.L. Casanova, and L. Quintana-Murci. 2021.
Human ancient DNA analyses reveal the high burden of tuberculosis in Europeans over the last
2,000 years. *Am J Hum Genet* 108:517-524.
Key, F.M., C. Posth, L.R. Esquivel-Gomez, R. Hubler, M.A. Spyrou, G.U. Neumann, A. Furtwangler, S. Sabin,
M. Burri, A. Wissgott, A.K. Lankapalli, A.J. Vagene, M. Meyer, S. Nagel, R. Tukhbatova, A. Khokhlov,
A. Chizhevsky, S. Hansen, A.B. Belinsky, A. Kalmykov, A.R. Kantorovich, V.E. Maslov, P.W.
Stockhammer, S. Vai, M. Zavattaro, A. Riga, D. Caramelli, R. Skeates, J. Beckett, M.G. Gradoli, N.
Steuri, A. Hafner, M. Ramstein, I. Siebke, S. Losch, Y.S. Erdal, N.F. Alikhan, Z. Zhou, M. Achtman,
K. Bos, S. Reinhold, W. Haak, D. Kuhnert, A. Herbig, and J. Krause. 2020. Emergence of human-
adapted Salmonella enterica is linked to the Neolithization process. *Nat Ecol Evol* 4:324-333.
Kirino, Y., G. Bertsias, Y. Ishigatsubo, N. Mizuki, I. Tugal-Tutkun, E. Seyahi, Y. Ozyazgan, F.S. Sacli, B. Erer,
H. Inoko, Z. Emrence, A. Cakar, N. Abaci, D. Ustek, C. Satorius, A. Ueda, M. Takeno, Y. Kim, G.M.

Wood, M.J. Ombrello, A. Meguro, A. Gul, E.F. Remmers, and D.L. Kastner. 2013. Genome-wide association analysis identifies new susceptibility loci for Behcet's disease and epistasis between HLA-B*51 and ERAP1. *Nat Genet* 45:202-207.

Kitur, K., S. Wachtel, A. Brown, M. Wickersham, F. Paulino, H.F. Penaloza, G. Soong, S. Bueno, D. Parker, and A. Prince. 2016. Necroptosis Promotes Staphylococcus aureus Clearance by Inhibiting Excessive Inflammatory Signaling. *Cell reports* 16:2219-2230.

Li, J., T. McQuade, A.B. Siemer, J. Napetschnig, K. Moriwaki, Y.S. Hsiao, E. Damko, D. Moquin, T. Walz, A. McDermott, F.K. Chan, and H. Wu. 2012. The RIP1/RIP3 necrosome forms a functional amyloid signaling complex required for programmed necrosis. *Cell* 150:339-350.

Liu, Z., L.F. Dagley, K. Shield-Artin, S.N. Young, A. Bankovacki, X. Wang, M. Tang, J. Howitt, C.A. Stafford, U. Nachbur, C. Fitzgibbon, S.E. Garnish, A.I. Webb, D. Komander, J.M. Murphy, J.M. Hildebrand, and J. Silke. 2021a. Oligomerization-driven MLKL ubiquitylation antagonizes necroptosis. *EMBO J* 40:e103718.

Liu, Z., H. Nailwal, J. Rector, M.M. Rahman, R. Sam, G. McFadden, and F.K. Chan. 2021b. A class of viral inducer of degradation of the necroptosis adaptor RIPK3 regulates virus-induced inflammation. *Immunity* 54:247-258 e247.

Martin, K.R., M. Pederzoli-Ribeil, E. Pacreau, S.S. Burgener, A. Dahdah, C. Candalh, E. Lauret, M. Foretz, L. Mouthon, B. Lucas, N. Thieblemont, C. Benarafa, P. Launay, and V. Witko-Sarsat. 2017. Transgenic Mice Expressing Human Proteinase 3 Exhibit Sustained Neutrophil-Associated Peritonitis. *Journal of immunology* 199:3914-3924.

Meng, Y., K.A. Davies, C. Fitzgibbon, S.N. Young, S.E. Garnish, C.R. Horne, C. Luo, J.M. Garnier, L.Y. Liang, A.D. Cowan, A.L. Samson, G. Lessene, J.J. Sandow, P.E. Czabotar, and J.M. Murphy. 2021. Human RIPK3 maintains MLKL in an inactive conformation prior to cell death by necroptosis. *Nat Commun* 12:6783.

Mompean, M., W. Li, J. Li, S. Laage, A.B. Siemer, G. Bozkurt, H. Wu, and A.E. McDermott. 2018. The Structure of the Necrosome RIPK1-RIPK3 Core, a Human Hetero-Amyloid Signaling Complex. *Cell* 173:1244-1253 e1210.

Murphy, J.M., P.E. Czabotar, J.M. Hildebrand, I.S. Lucet, J.G. Zhang, S. Alvarez-Diaz, R. Lewis, N. Lalaoui, D. Metcalf, A.I. Webb, S.N. Young, L.N. Varghese, G.M. Tannahill, E.C. Hatchell, I.J. Majewski, T. Okamoto, R.C.J. Dobson, D.J. Hilton, J.J. Babon, N.A. Nicola, A. Strasser, J. Silke, and W.S. Alexander. 2013. The Pseudokinase MLKL Mediates Necroptosis via a Molecular Switch Mechanism. *Immunity* 39:443-453.

Murphy, J.M., I.S. Lucet, J.M. Hildebrand, M.C. Tanzer, S.N. Young, P. Sharma, G. Lessene, W.S. Alexander, J.J. Babon, J. Silke, and P.E. Czabotar. 2014. Insights into the evolution of divergent nucleotide-binding mechanisms among pseudokinases revealed by crystal structures of human and mouse MLKL. *The Biochemical journal* 457:369-377.

Pearson, J.S., C. Giogha, S. Muhlen, U. Nachbur, C.L. Pham, Y. Zhang, J.M. Hildebrand, C.V. Oates, T.W. Lung, D. Ingle, L.F. Dagley, A. Bankovacki, E.J. Petrie, G.N. Schroeder, V.F. Crepin, G. Frankel, S.L. Masters, J. Vince, J.M. Murphy, M. Sunde, A.I. Webb, J. Silke, and E.L. Hartland. 2017. EspL is a bacterial cysteine protease effector that cleaves RHIM proteins to block necroptosis and inflammation. *Nat Microbiol* 2:16258.

Petrie, E.J., R.W. Birkinshaw, A. Koide, E. Denbaum, J.M. Hildebrand, S.E. Garnish, K.A. Davies, J.J. Sandow, A.L. Samson, X. Gavin, C. Fitzgibbon, S.N. Young, P.J. Hennessy, P.P.C. Smith, A.I. Webb, P.E. Czabotar, S. Koide, and J.M. Murphy. 2020. Identification of MLKL membrane translocation as a checkpoint in necroptotic cell death using Monobodies. *Proceedings of the National Academy of Sciences of the United States of America* 117:8468-8475.

Petrie, E.J., J.J. Sandow, A.V. Jacobsen, B.J. Smith, M.D.W. Griffin, I.S. Lucet, W. Dai, S.N. Young, M.C. Tanzer, A. Wardak, L.Y. Liang, A.D. Cowan, J.M. Hildebrand, W.J.A. Kersten, G. Lessene, J. Silke,

P.E. Czabotar, A.I. Webb, and J.M. Murphy. 2018. Conformational switching of the pseudokinase
domain promotes human MLKL tetramerization and cell death by necroptosis. *Nat Commun*
9:2422.

Petrie, E.J., J.J. Sandow, W.I.L. Lehmann, L.Y. Liang, D. Coursier, S.N. Young, W.J.A. Kersten, C. Fitzgibbon,
A.L. Samson, A.V. Jacobsen, K.N. Lowes, A.E. Au, H. Jousset Sabroux, N. Lalaoui, A.I. Webb, G.
Lessene, G. Manning, I.S. Lucet, and J.M. Murphy. 2019. Viral MLKL Homologs Subvert Necroptotic
Cell Death by Sequestering Cellular RIPK3. *Cell Rep* 28:3309-3319 e3305.

Pierdomenico, M., A. Negroni, L. Stronati, R. Vitali, E. Prete, J. Bertin, P.J. Gough, M. Aloi, and S. Cucchiara.
2014. Necroptosis is active in children with inflammatory bowel disease and contributes to
heighten intestinal inflammation. *Am J Gastroenterol* 109:279-287.

Ramos, P.S., A.M. Shedlock, and C.D. Langefeld. 2015. Genetics of autoimmune diseases: insights from
population genetics. *J Hum Genet* 60:657-664.

Rickard, J.A., H. Anderton, N. Etemadi, U. Nachbur, M. Darding, N. Peltzer, N. Lalaoui, K.E. Lawlor, H.
Vanyai, C. Hall, A. Bankovacki, L. Gangoda, W.W. Wong, J. Corbin, C. Huang, E.S. Mocarski, J.M.
Murphy, W.S. Alexander, A.K. Voss, D.L. Vaux, W.J. Kaiser, H. Walczak, and J. Silke. 2014a. TNFR1-
dependent cell death drives inflammation in Sharpin-deficient mice. *Elife* 3:

Rickard, J.A., J.A. O'Donnell, J.M. Evans, N. Lalaoui, A.R. Poh, T. Rogers, J.E. Vince, K.E. Lawlor, R.L. Ninnis,
H. Anderton, C. Hall, S.K. Spall, T.J. Phesse, H.E. Abud, L.H. Cengia, J. Corbin, S. Mifsud, L. Di Rago,
D. Metcalf, M. Ernst, G. Dewson, A.W. Roberts, W.S. Alexander, J.M. Murphy, P.G. Ekert, S.L.
Masters, D.L. Vaux, B.A. Croker, M. Gerlic, and J. Silke. 2014b. RIPK1 regulates RIPK3-MLKL-driven
systemic inflammation and emergency hematopoiesis. *Cell* 157:1175-1188.

Robinson, P.C., M.E. Costello, P. Leo, L.A. Bradbury, K. Hollis, A. Cortes, S. Lee, K.B. Joo, S.C. Shim, M.
Weisman, M. Ward, X. Zhou, H.J. Garchon, G. Chiocchia, J. Nossent, B.A. Lie, O. Forre, J.
Tuomilehto, K. Laiho, L. Jiang, Y. Liu, X. Wu, D. Elewaut, R. Burgos-Vargas, L.S. Gensler, S.
Stebbings, N. Haroon, J. Mulero, J.L. Fernandez-Sueiro, M.A. Gonzalez-Gay, C. Lopez-Larrea, P.
Bowness, K. Gafney, J.S. Gaston, D.D. Gladman, P. Rahman, W.P. Maksymowych, H. Xu, I.E. van
der Horst-Bruinsma, C.T. Chou, R. Valle-Onate, M.C. Romero-Sanchez, I.M. Hansen, F.M. Pimentel-
Santos, R.D. Inman, J. Martin, M. Breban, D. Evans, J.D. Reveille, T.H. Kim, B.P. Wordsworth, and
M.A. Brown. 2015. ERAP2 is associated with ankylosing spondylitis in HLA-B27-positive and HLA-
B27-negative patients. *Ann Rheum Dis* 74:1627-1629.

Rodriguez, D.A., R. Weinlich, S. Brown, C. Guy, P. Fitzgerald, C.P. Dillon, A. Oberst, G. Quarato, J. Low, J.G.
Cripps, T. Chen, and D.R. Green. 2016. Characterization of RIPK3-mediated phosphorylation of the
activation loop of MLKL during necroptosis. *Cell Death Differ* 23:76-88.

Samson, A.L., S.E. Garnish, J.M. Hildebrand, and J.M. Murphy. 2021. Location, location, location: A
compartmentalized view of TNF-induced necroptotic signaling. *Science signaling* 14:

Samson, A.L., Y. Zhang, N.D. Geoghegan, X.J. Gavin, K.A. Davies, M.J. Mlodzianoski, L.W. Whitehead, D.
Frank, S.E. Garnish, C. Fitzgibbon, A. Hempel, S.N. Young, A.V. Jacobsen, W. Cawthorne, E.J. Petrie,
M.C. Faux, K. Shield-Artin, N. Lalaoui, J.M. Hildebrand, J. Silke, K.L. Rogers, G. Lessene, E.D.
Hawkins, and J.M. Murphy. 2020. MLKL trafficking and accumulation at the plasma membrane
control the kinetics and threshold for necroptosis. *Nat Commun* 11:3151.

Sethi, A., C.R. Horne, C. Fitzgibbon, K. Wilde, K.A. Davies, S.E. Garnish, A.V. Jacobsen, A.L. Samson, J.M.
Hildebrand, A. Wardak, P.E. Czabotar, E.J. Petrie, P.R. Gooley, and J.M. Murphy. 2022. Membrane
permeabilization is mediated by distinct epitopes in mouse and human orthologs of the
necroptosis effector, MLKL. *Cell Death Differ*

Strange, A., F. Capon, C.C. Spencer, J. Knight, M.E. Weale, M.H. Allen, A. Barton, G. Band, C. Bellenguez,
J.G. Bergboer, J.M. Blackwell, E. Bramon, S.J. Bumpstead, J.P. Casas, M.J. Cork, A. Corvin, P.
Deloukas, A. Dilthey, A. Duncanson, S. Edkins, X. Estivill, O. Fitzgerald, C. Freeman, E. Giardina, E.
Gray, A. Hofer, U. Huffmeier, S.E. Hunt, A.D. Irvine, J. Jankowski, B. Kirby, C. Langford, J. Lascorz,

J. Leman, S. Leslie, L. Mallbris, H.S. Markus, C.G. Mathew, W.H. McLean, R. McManus, R. Mossner, L. Moutsianas, A.T. Naluai, F.O. Nestle, G. Novelli, A. Onoufriadis, C.N. Palmer, C. Perricone, M. Pirinen, R. Plomin, S.C. Potter, R.M. Pujol, A. Rautanen, E. Riveira-Munoz, A.W. Ryan, W. Salmhofer, L. Samuelsson, S.J. Sawcer, J. Schalkwijk, C.H. Smith, M. Stahle, Z. Su, R. Tazi-Ahnini, H. Traupe, A.C. Viswanathan, R.B. Warren, W. Weger, K. Wolk, N. Wood, J. Worthington, H.S. Young, P.L. Zeeuwen, A. Hayday, A.D. Burden, C.E. Griffiths, J. Kere, A. Reis, G. McVean, D.M. Evans, M.A. Brown, J.N. Barker, L. Peltonen, P. Donnelly, and R.C. Trembath. 2010. A genome-wide association study identifies new psoriasis susceptibility loci and an interaction between HLA-C and ERAP1. *Nat Genet* 42:985-990.

Strugnell, R., G. Dougan, S. Chatfield, I. Charles, N. Fairweather, J. Tite, J.L. Li, J. Beesley, and M. Roberts. 1992. Characterization of a Salmonella typhimurium aro vaccine strain expressing the P.69 antigen of Bordetella pertussis. *Infect Immun* 60:3994-4002.

Su, L., B. Quade, H. Wang, L. Sun, X. Wang, and J. Rizo. 2014. A plug release mechanism for membrane permeation by MLKL. *Structure* 22:1489-1500.

Sun, L., H. Wang, Z. Wang, S. He, S. Chen, D. Liao, L. Wang, J. Yan, W. Liu, X. Lei, and X. Wang. 2012. Mixed lineage kinase domain-like protein mediates necrosis signaling downstream of RIP3 kinase. *Cell* 148:213-227.

Tanzer, M.C., I. Matti, J.M. Hildebrand, S.N. Young, A. Wardak, A. Tripaydonis, E.J. Petrie, A.L. Mildenhall, D.L. Vaux, J.E. Vince, P.E. Czabotar, J. Silke, and J.M. Murphy. 2016. Evolutionary divergence of the necroptosis effector MLKL. *Cell Death Differ* 23:1185-1197.

Thapa, R.J., S. Nogusa, P. Chen, J.L. Maki, A. Lerro, M. Andrake, G.F. Rall, A. Degterev, and S. Balachandran. 2013. Interferon-induced RIP1/RIP3-mediated necrosis requires PKR and is licensed by FADD and caspases. *Proceedings of the National Academy of Sciences of the United States of America* 110:E3109-3118.

Tovey Crutchfield, E.C., S.E. Garnish, and J.M. Hildebrand. 2021. The Role of the Key Effector of Necroptotic Cell Death, MLKL, in Mouse Models of Disease. *Biomolecules* 11:

Tsoi, L.C., S.L. Spain, J. Knight, E. Ellinghaus, P.E. Stuart, F. Capon, J. Ding, Y. Li, T. Tejasvi, J.E. Gudjonsson, H.M. Kang, M.H. Allen, R. McManus, G. Novelli, L. Samuelsson, J. Schalkwijk, M. Stahle, A.D. Burden, C.H. Smith, M.J. Cork, X. Estivill, A.M. Bowcock, G.G. Krueger, W. Weger, J. Worthington, R. Tazi-Ahnini, F.O. Nestle, A. Hayday, P. Hoffmann, J. Winkelmann, C. Wijmenga, C. Langford, S. Edkins, R. Andrews, H. Blackburn, A. Strange, G. Band, R.D. Pearson, D. Vukcevic, C.C. Spencer, P. Deloukas, U. Mrowietz, S. Schreiber, S. Weidinger, S. Koks, K. Kingo, T. Esko, A. Metspalu, H.W. Lim, J.J. Voorhees, M. Weichenthal, H.E. Wichmann, V. Chandran, C.F. Rosen, P. Rahman, D.D. Gladman, C.E. Griffiths, A. Reis, J. Kere, P. Collaborative Association Study of, C. Genetic Analysis of Psoriasis, E. Psoriasis Association Genetics, C. Wellcome Trust Case Control, R.P. Nair, A. Franke, J.N. Barker, G.R. Abecasis, J.T. Elder, and R.C. Trembath. 2012. Identification of 15 new psoriasis susceptibility loci highlights the role of innate immunity. *Nat Genet* 44:1341-1348.

Upton, J.W., W.J. Kaiser, and E.S. Mocarski. 2010. Virus inhibition of RIP3-dependent necrosis. *Cell Host Microbe* 7:302-313.

Vince, J.E., W.W. Wong, N. Khan, R. Feltham, D. Chau, A.U. Ahmed, C.A. Benetatos, S.K. Chunduru, S.M. Condon, M. McKinlay, R. Brink, M. Leverkus, V. Tergaonkar, P. Schneider, B.A. Callus, F. Koentgen, D.L. Vaux, and J. Silke. 2007. IAP antagonists target cIAP1 to induce TNFalpha-dependent apoptosis. *Cell* 131:682-693.

Wang, B., S. Bao, Z. Zhang, X. Zhou, J. Wang, Y. Fan, Y. Zhang, Y. Li, L. Chen, Y. Jia, J. Li, M. Li, W. Zheng, N. Mu, L. Wang, Z. Yu, D.S.M. Wong, Y. Zhang, J. Kwan, H. Ka-Fung Mak, A. Ambalavanan, S. Zhou, W. Cai, J. Zheng, S. Huang, G.A. Rouleau, W. Yang, E. Rogaeva, X. Ma, P. St George-Hyslop, L.W. Chu, and Y.Q. Song. 2018. A rare variant in MLKL confers susceptibility to ApoE varepsilon4-

negative Alzheimer's disease in Hong Kong Chinese population. *Neurobiol Aging* 68:160 e161-160
e167.
Wang, H., L. Sun, L. Su, J. Rizo, L. Liu, L.F. Wang, F.S. Wang, and X. Wang. 2014a. Mixed lineage kinase
domain-like protein MLKL causes necrotic membrane disruption upon phosphorylation by RIP3.
Mol Cell 54:133-146.
Wang, H., H. Yang, C.S. Shivalila, M.M. Dawlaty, A.W. Cheng, F. Zhang, and R. Jaenisch. 2013. One-step
generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome
engineering. *Cell* 153:910-918.
Wang, H.Y., L.M. Sun, L.J. Su, J. Rizo, L. Liu, L.F. Wang, F.S. Wang, and X.D. Wang. 2014b. Mixed Lineage
Kinase Domain-like Protein MLKL Causes Necrotic Membrane Disruption upon Phosphorylation
by RIP3. *Molecular Cell* 54:133-146.
Weir, A., S. Hughes, M. Rashidi, J.M. Hildebrand, and J.E. Vince. 2021. Necroptotic movers and shakers:
cell types, inflammatory drivers and diseases. *Curr Opin Immunol* 68:83-97.
Wemyss, M.A., and J.S. Pearson. 2019. Host Cell Death Responses to Non-typhoidal Salmonella Infection.
Front Immunol 10:1758.
Werren, E.A., O. Garcia, and A.W. Bigham. 2021. Identifying adaptive alleles in the human genome: from
selection mapping to functional validation. *Hum Genet* 140:241-276.
Wu, J., Z. Huang, J. Ren, Z. Zhang, P. He, Y. Li, J. Ma, W. Chen, Y. Zhang, X. Zhou, Z. Yang, S.Q. Wu, L. Chen,
and J. Han. 2013. Mlkl knockout mice demonstrate the indispensable role of Mlkl in necroptosis.
Cell research 23:994-1006.
Yan, B., L. Liu, S. Huang, Y. Ren, H. Wang, Z. Yao, L. Li, S. Chen, X. Wang, and Z. Zhang. 2017. Discovery of
a new class of highly potent necroptosis inhibitors targeting the mixed lineage kinase domain-like
protein. *Chem Commun (Camb)* 53:3637-3640.
Yin, X., H.Q. Low, L. Wang, Y. Li, E. Ellinghaus, J. Han, X. Estivill, L. Sun, X. Zuo, C. Shen, C. Zhu, A. Zhang, F.
Sanchez, L. Padyukov, J.J. Catanese, G.G. Krueger, K.C. Duffin, S. Mucha, M. Weichenthal, S.
Weidinger, W. Lieb, J.N. Foo, Y. Li, K. Sim, H. Liany, I. Irwan, Y. Teo, C.T. Theng, R. Gupta, A.
Bowcock, P.L. De Jager, A.A. Qureshi, P.I. de Bakker, M. Seielstad, W. Liao, M. Stahle, A. Franke, X.
Zhang, and J. Liu. 2015. Genome-wide meta-analysis identifies multiple novel associations and
ethnic heterogeneity of psoriasis susceptibility. *Nat Commun* 6:6916.
Yoon, S., K. Bogdanov, and D. Wallach. 2022. Site-specific ubiquitination of MLKL targets it to endosomes
and targets Listeria and Yersinia to the lysosomes. *Cell Death Differ*
Zhang, J., T. Jin, I. Aksentijevich, and Q. Zhou. 2021. RIPK1-Associated Inborn Errors of Innate Immunity.
Front Immunol 12:676946.
Zhao, J., S. Jitkaew, Z. Cai, S. Choksi, Q. Li, J. Luo, and Z.G. Liu. 2012. Mixed lineage kinase domain-like is a
key receptor interacting protein 3 downstream component of TNF-induced necrosis. *Proceedings
of the National Academy of Sciences of the United States of America* 109:5322-5327.
Zhu, X., N. Yang, Y. Yang, F. Yuan, D. Yu, Y. Zhang, Z. Shu, N. Nan, H. Hu, X. Liu, S. Chen, L. Sun, and H.
Wang. 2022. Spontaneous necroptosis and autoinflammation are blocked by an inhibitory
phosphorylation on MLKL during neonatal development. *Cell Res* 32:407-410.
1182
1183

Figure 2