A novel fluorobenzothiazole RBx 10080758 as a dual inhibitor against bacterial GyraseB (GyrB) and Topoisomerase IV (parE) of Gram-positive pathogens causing skin and respiratory infections

Tarani Kanta Barman[1,4], Manoj Kumar[1,4], Tridib Chaira[3,6,7], Smita Singhal[1,4], Tarun Mathur[1,4], Vandana Kalia[1,4], Ramkumar Gangadharan[1,4,7], Madhvi Rao[1,4], Manisha Pandya[1,4], Pragya Bhateja[1,4], Ruchi Sood[1,4], Dilip J. Upadhyay[1,4], Shibu Varughese[2,5], Ajay Yadav[2,5], Lalima Sharma[2,5], Naresh Kumar[2,5], Jitendra Sattigeri[2,5], Pradip K. Bhatnagar[2,5] and V. Samuel Raj[1,4,7]*

[1]Department of Infectious Diseases, [2]Department of Chemistry, [3]Department of Metabolism & Pharmacokinetics, New Drug Discovery Research, Ranbaxy Research Laboratories, R & D III, Sector-18, Gurgaon – 122 015, India
[4]Department of Microbiology, [5]Department of Chemistry, [6]Department of Drug Metabolism & Pharmacokinetics, Daiichi Sankyo India Pharma Private Limited, Village Sarhaul, Sector – 18, Gurgaon – 122 015, India
[7]Centre for Drug Design Discovery and Development (C4D), SRM University, Delhi-NCR, Sonepat – 131 029, Haryana, India

Running title: Efficacy of RB1080758 against Gram positive pathogens

*Address for Correspondence:

Prof. V. Samuel Raj
Centre for Drug Design Discovery and Development (C4D)
SRM University, Delhi NCR,
Rajiv Gandhi Education City,
Sonepat, Haryana – 131029, India
E-mail: directorcd4@srmuniversity.ac.in; deanacademic@srmuniversity.ac.in

Abstract

Development of a novel inhibitor targeting Gyrase B and Topoisomerase IV offers a potential opportunity to combat the drug resistance. In the present study, we extensively investigated the efficacy of RBx 10080758, a novel fluorobenzothiazole, against skin and respiratory infections caused by Staphylococci and Streptococci in *in vitro* and *in vivo* models. RBx 10080758 showed a potent IC_{50} of 0.06 µM against Gyrase and Topoisomerase IV and also exhibited a strong whole cell *in vitro* activity with MIC ranges of 0.015-0.06, 0.015-0.03, 0.008-0.03, 0.008-0.03, 0.008-0.0.06, 0.015-0.06 µg/ml against *Staphylococcus. aureus, Streptococcus pneumoniae,* Coagulase negative Staphylococci, *Streptococcus viridans, Streptococcus pyogenes* and Enterococcus, respectively. As expected from the novel class of molecule, it retains potent even against linezolid and vancomycin resistant strains. Interesting, in mouse systemic infection 10 mg/kg, IV dose protected 100% mice from lethal infections. In rat thigh infection model with MRSA WCUH29 at 45 mg/kg exhibited >3-$\log_{10}$ cfu reduction in thigh muscles. RBx 10080758 displayed potent *in vitro* and *in vivo* activity against a panel of MDR Gram-positive bacteria. As a novel chemical class, the fluorobenzothiazoles have the potential to become clinically viable antibiotics, to address the drug resistance problem by its unique dual targeting mechanism of action.

Keywords: Fluorobenzothiazole, GyrB and Topoisomerase IV, dual inhibitors, Gram positive bacteria, rat thigh infection

Introduction

Gram-positive bacteria, particularly nosocomial methicillin resistant *Staphylococcus aureus* (MRSA), have emerged as a superbug, showing resistance to multiple antibiotics [1, 2]. MRSA infection constitutes the most important cause of health care associated infections, and capable of causing various illness ranging from mild skin infection to life-threatening pneumonia and blood stream infection [3, 4]. The treatment for MRSA is quite complicated and expensive as the infections can recur frequently and can easily progress to life-threatening septicemia or atopic dermatitis [5, 6]. The standard of care (SOC) drug of MRSA such as vancomycin, daptomycin, linezolid and tedizolid are promising antibiotics but not completely devoid of limitations [7, 8]. For example, the extended use of linezolid may cause myelosuppression, while tedizolid shows poor activity in neutropenic patients [7]. Daptomycin has poor activity in pulmonary surfactant and also cause myopathy and neuropathy [9]. Therefore, there is a high unmet medical need for a new class of anti-MRSA antibiotic to overcome the limitations of SOC and fight against emerging resistance.

Development of novel inhibitors against DNA Gyrase B (GyrB) and Topoisomerase IV offers a potential opportunity to meet the challenge of resistance [10, 11]. GyrB and Topoisomerase IV are essential enzymes and play important role in DNA replication and compaction. DNA supercoiling activity is essential in all bacteria but not found in human and hence it is an ideal target [12]. The overall identity between *GyrB* and Topoisomerase IV *parE* is 52%, whereas the identity between *GyrB* and *GyrA* is only 5% in MRSA. In addition, identity of bacterial *GyrB* with human topoisomerase is only 12%, which further reduce the possibility of toxicity [13, 14]. The coumarin class of compounds novobiocin and coumermycin bind to GyrB and parE and both

target ATP binding sites [15]. In addition, Cyclothialidines are potent inhibitors of Gyr B or

GyrB/par E. However, their uses have been limited by poor PK profiles [15, 16].

Therefore, it is of great opportunity to design a dual inhibitor against GyrB and parE, especially

directed at the ATP binding sites. Additionally dual-target inhibitors will pose lower risks of

resistance development than single-target inhibitors. RBx 10080758 is a novel

fluorobenzothiazole inhibits both GyrB and Topoisomerase IV parE of Gram-positive bacteria.

In present study, the *in vitro* and *in vivo* activities of RBx 10080758 with pharmacokinetics

against Gram positive bacteria particularly MRSA were evaluated.

Results

Enzyme inhibitory activity of RBx 10080758 and RBx 10115912.

RBx 10080758 and RBx 10115912 (figure 1) showed a strong inhibitor activity against bacterial

gyrase and topoisomerase IV. The 50% inhibitory concentrations (IC_{50s}) of RBx 10080758 and

RBx 10115912 for *E. coli* gyrase supercoiling and topoisomerase IV decatenation assay are

shown in Table 1. Both compounds showed no inhibitory activity against human topoisomerase

II up to 500 µM (Table 1), suggesting their high selectivity to bacterial topoisomerases.

Antibacterial activity against Gram-positive bacteria

The *in vitro* activities of RBx 10080758 and RBx 10115912 against a panel of Gram-positive

bacteria such as *S. aureus, S. pneumoniae,* coagulase negative Staphylococci, *S. viridians, S.*

pyogenes and Enterococcus are summarized in Table 2. RBx 10080758 displayed stronger

inhibitory activity against recent clinical isolates of *S. aureus* including MSSA, MRSA (n=50)

and multiple drug resistant (MDR) pathogens with MIC_{90} of 0.06 µg/ml and MIC range of 0.015-

0.06 µg/ml (Table 2). Overall, both RBx 10080758 and RBx 10115912 displayed potent activity

against multiple MDR bacteria (Supplementary figure 1a-d).

In vitro **cytotoxicity**

Cytotoxicity assessment of RBx 10080758 and RBx 10115912 was performed using standard

MTT assay. The IC_{50} values were calculated using Graph Pad Prism and both compounds

showed IC_{50} value of ≥ 50 µg/ml.

Frequency of resistance

Though RBx 10080758 and RBx 10115912 showed potent activity against the panel of bacterial

pathogens, RBx 10115912 showed poor solubility hence was not evaluated for further studies.

The frequency of spontaneous resistance (FSR) development was determined at $4 \times$ and $8 \times$ the

MIC of RBx 10080758 against MRSA. The FSR at $4 \times$ MICs of RBx 10080758 was $<3.5 \times 10^{-9}$

against MRSA WCUH-29 and $<6 \times 10^{-9}$ against MRSA 562 (Table 3). RBx 10080758 showed

mutation prevention concentration (MPC) of 0.125 µg/ml and completely prevented the

spontaneous resistant development at 0.125 µg/ml.

Combination studies of RBx 10080758

RBx 10080758 was checked for activity in combination with ten existing drugs against *S. aureus*

strains. The FIC index against the antibiotics are as follows: Linezolid (FIC index 1),

Vancomycin (1.5), Gentamicin (1), Ciprofloxacin (1), Erythromycin (2), Ertapenem (1),

Imipenem (1.2), Augmentin (1.5), Clindamycin (1.5) and Cefoxitin (1.25). The FIC index of

RBx 10080758 against all the antibiotics is indicative of an additive effect (data not shown). RBx

10080758 shows no synergy or antagonism with existing drugs being used for treatment against

S. aureus.

Time- kill kinetic studies

The results of time-kill studies of RBx 10080758 and linezolid against MRSA 562, MRSA

WCUH-29, *S. pneumoniae* MA 80 and *S. pyogenes* are presented in Figure 2 a-d. RBx 10080758

showed rapid concentration dependent killing of MRSA isolates (Figure 2a, 2b). However, RBx

10080758 exhibited time-dependent bactericidal activity against *S. pneumoniae* MA-80 and *S.*

pyogenes even at $1 \times$ MIC (Figure 2c, 2d).

Murine systemic infection model of *MRSA-562*

The *in vivo* efficacy of RBx 100 80758 in murine systemic infection model of MRSA-562 is

presented in Figure 3. All the infected mice treated with placebo, started developing the clinical

symptoms of infection with 8 h of infection and died by following day. RBx 10080758 showed a

dose dependent efficacy against MRSA-562 (Figure 3a) with ED_{50} of 5 mg/kg. The kidney

bacterial load at 8 h or 24 h post-infection, showed significant reduction in bacterial count at 10

and 20 mg/kg dose as compared to placebo control or linezolid (Figure 3b), suggesting the

superior killing potential of RBx 10080758.

Rat thigh and lung infection models

The efficacy of RBx 10080758 (Na salts) was investigated against MRSA WCUH-29 in rat thigh

infection model (Fig. 4a-4c). RBx 10080758 showed a dose dependent efficacy against MRSA

WCUH-29 with a bactericidal potential at 45 mg/kg, IV dose (Figure 4a). When tested against

the highly virulent MRSA PVL-2 strain, RBx 10080758 exhibited similar bactericidal efficacy at

45 mg/kg dose (Figure 4b). Similar activity of RBx 10080758 was also observed against MRSA

WCUH-29 at 45 mg/kg dose in 2 days infection model (Figure 4c).

In the rat lung infection of *S. pneumoniae* MA 80, RBx 10080758 exhibited a bactericidal effect

at 45 mg/kg dose with a significant >3 $\log_{10}$ reduction in bacterial load as compared to placebo

control. Linezolid and telithromycin also exhibited a similar effect against *S. pneumoniae* MA-80

(Figure 4d).

***In vitro* clearance RBx 10080758 in liver microsomes**

The clearance of RBx 10080758 in human liver microsomes contributed by glucuronidation was

assessed. The results revealed a minimal involvement of Phase II glucuronidation in the

clearance of RBx 10080758. The clearance of RBx 10080758 in liver microsomes was stable and

this is not a high clearance molecule.

In CYP inhibition assay, all the compounds from 4-fluorobenzothiazole series (data not shown)

have shown IC_{50} for 5 CYP enzymes (CYP1A2, CYP2C9, CYP2C19, CYP2D6 and CYP3A4)

>10 μM concentration. Intrinsic clearance of 4-fluorobenzthiazole/pyrrole ABH amide series of

compounds are metabolically stable, and they have shown low-moderate clearance in rat and

human liver microsomes (<0.6 mL/min/kg). RBx 10080758 and RBx 10115912 are two

representative compounds of these series.

***In Vivo* pharmacokinetics of RBx 10080758**

In vivo pharmacokinetic data of RBx 10080758 are presented in Table 4. RBx 10080758

displayed similar intravenous pharmacokinetics in rat and dog. In both the species, it showed low

mean plasma clearance (rat 11.34 ml/min/kg and dog 13.67 ml/min/kg), low V_{ss} (< 0. 7) and

short half-life (< 2.0 h). The plasma exposure (AUC_{inf}) of RBx 10080758 was 0.07, 1.5 and 1.27

μg.h/ml in mouse, rat, and dog, respectively by IV route.

Discussion

RBx 10080758 exhibited potent activity against both *S. aureus* DNA GyrB and Topoisomerase

IV enzymes and showed >1000-fold selectivity against human Topoisomerase II. Usually, gyrase

inhibitors are more active against DNA gyrase than topoisomerase, while quinolones show more

inhibitory potency against Topoisomerase IV than that of GyrB [17, 18]. Interestingly, RBx

10080758 inhibited *S. aureus* DNA GyrB and Topoisomerase IV with equal potency, which may

significantly reduce the risk of resistance development. As RBx 10080758 is a novel dual

inhibitor of DNA GyrB and Topoisomerase IV and exhibited potent MIC against various Gram-

positive bacteria with no cross-resistance to fluoroquinolones or other antibiotics, which

highlights its utility against MDR pathogens. RBx 10080758 showed excellent *in vitro* activity

against a panel of skin and soft tissue and respiratory pathogens. RBx 10080758 showed ≥16-

fold potent MIC as compared to ciprofloxacin and ≥60-fold potent MIC as compared to linezolid

against *S. aureus*. Most clinical isolates of MRSA are resistant to fluoroquinolones, which limits

their clinical utilities [17, 19]. In addition, emergence of large number of MDR gram-positive

strains in nosocomial settings even against recently developed linezolid and tedizolid, limiting

the treatment options against these MDR pathogens [20]. Although, in the recent years, several

GyrB and ParE inhibitors demonstrated potent *in vitro* activities against MDR Gram-Positive

pathogens, however none have showed acceptable *in vivo* efficacy or progress for clinical

development.

Interestingly, RBx 10080758 exhibited superior activities against the different MDR strains such

as MDR *S. aureus*, linezolid resistant (LZD[res]) *S. aureus*, vancomycin-intermediate *S. aureus*

(VISA), vancomycin resistant *S. aureus* (VRSA), vancomycin resistant Enterococci (VRE),

LZD[res] *S. pyogenes,* fluoroquinolone resistant (FQ[res]) *S. viridians* and *S. pneumoniae*

(Supplementary figure 1a-d) and showed low frequency of spontaneous resistance, suggest that

RBx 10080758 may effectively cover both susceptible and MDR pathogens and will have a low

propensity for resistance development in clinical settings [21]. RBx 10080758 exhibited low

MPC of 0.125 µg/ml which may be useful in selecting appropriate dosing of drug for efficacy

studies and determining the potential of drug in preventing the selection of resistant mutants.

Pulmonary surfactant reduces the surface tension at the air/liquid interface of alveoli and is

known to significantly impair the antimicrobial activities of moxifloxacin, colistin and

daptomycin [9, 12, 22, 23]. However, the activity of RBx 10080758 was not affected by the

presence of a lung surfactant (Supplementary Figure 2), demonstrating its effectiveness in

pulmonary infections.

Though, RBx 10080758 was found metabolically stable, however, its poor solubility hindered

the further evaluation. Different salt forms of RBx 10080758 such as sodium, lithium and

hydrochloric acid salts were explored to improve its solubility; sodium salt of RBx 10080758

showed significantly better solubility, stability, and exhibited the similar MIC activity against

bacterial pathogens (data not shown). Hence sodium salt of RBx 10080758 was selected for *in*

vivo evaluation. To confirm the broad-spectrum gram-positive pathogen coverage, the efficacy of

RBx 10080758 was assessed, in rat thigh model against *S. aureus* and in lung infection against *S.*

pneumoniae as well as mouse systemic infection models caused by MRSA. These models bear a

resemblance to human skin and soft tissue infection (SSTI), pneumonia and septicemia. RBx

10080758 showed excellent *in vivo* efficacy with ≥ 3 $\log_{10}$ reduction in one day as well as 2 days

rat thigh infection model against *S. aureus* indicating its suitability for SSTI indications. RBx

10080758 also showed potent efficacy against *S. pneumoniae* in rat lung infection model,

indicating the potential for lung infection.

In systemic infection rapid clearance of pathogen is extremely important to control the spread of

infection. RBx 10080758 showed 100 % survival at 10 mg/kg dose in murine systemic infection

model of MRSA-562 and exhibited a significant reduction in the kidney bacterial loads as

compared to linezolid and untreated control. This indicated superior clinical and microbiological

cure by RBx 10080758 compared to linezolid. To calculate the clinical efficacious dose of RBx

10080758 and to confirm safety, further animal studies are warranted [24, 25]. RBx 10080758

with low clearance and low volume of distribution, yielded AUC_{inf} (µg.h/ml) and C_{max} that

correlated well with *in vivo* efficacy in rat. There is likely an improvement in efficacy in dog

(non-rodent species) due to similar pharmacokinetics observed between rat and dog. Taken

together, RBx 10080758 exhibited potent *in vitro* and *in vivo* activities against MRSA and *S.*

pneumoniae. These results indicate that RBx 10080758 can be explored as a new therapeutic

agent against Gram-positive bacterial infections that cause SSTI and pneumonia.

Materials and Methods

Bacterial strains, antibiotics and new chemical entities

Different bacterial isolates, such as, *Staphylococcus aureus* (n=81) including MSSA, MRSA,

vancomycin and linezolid resistant strains, *Staphylococcus epidermidis* (n=35), *Streptococcus*

viridans group (n=20), Enterococci (n=27) including *Enterococcus faecalis* and *Enterococcus*

faecium and *Streptococcus pyogenes* (n=50) were used in this study. Standard strains were

purchased from the American Type Culture Collection (ATCC, Manassas, VA, USA), and

clinical isolates were acquired from Indian hospitals. RBx 10080758 (Figure 1a) and RBx

10115912 (Figure 1b) were synthesized in-house at Daiichi Sankyo India Pharma Pvt. Ltd.,

Gurgaon, India. Standard antibiotics were purchased from commercial sources.

DNA Gyrase supercoiling assay

E.coli DNA gyrase supercoiling assay was performed as we described earlier [18]. Briefly, the

assay was carried out in a reaction volume of 20 µl containing 1 unit of DNA gyrase and 0.5 µg

of relaxed pBR322 DNA in assay buffer in presence of RBx 10080758, RBx 10115912 or

levofloxacin. After incubation at $37^{o}C$ for 60 min with and without inhibitors, the reaction was

stopped. The amount of the supercoiled DNA band was quantified using Bio-Rad's Quantity one

software.

Topoisomerase IV decatenation assay

253 DNA decatenation assay was performed using *E. coli* Topoisomerase IV (subunits A and B) as

254 described earlier [11, 18]. The assay was performed in 20 µl of a reaction mixture containing 1

255 unit of Topoisomerase IV and 0.4 µg of catenated kinetoplast pBR322 DNA in assay buffer in

256 presence or absence of test compounds. After incubation at 37°C for 60 min, the reaction was

257 stopped. The IC_{50} values were determined as described earlier [18].

Activity against human topoisomerase IIα

Human Topoisomerase-IIα was synthesized as a histidine-tagged protein in *E. coli*. Decatenation

assays were performed in 20 µl reaction mixture which include 50 mM Tris-HCl (pH 7.5), 120

mM KCl, 10 mM $MgCl_2$, 0.5 mM ATP, 0.5 mM DTT, 30 µg/ml BSA, 0.4 µg catenated

kinetoplast DNA as a DNA substrate, human topoisomerase IIα enzyme, and test compounds.

After incubation at 37 °C for 60 min, the reaction was terminated and IC_{50}s were determined as

described above. All the above experiment was performed in duplicate and repeated twice.

***In vitro* susceptibility testing**

Minimum inhibitory concentration**s** (MIC) of RBx 10080758, RBx 1011591, and other standard

antibiotics were determined against a panel of Gram-positive bacterial strains using micro-broth

dilution method as recommended by the Clinical and Laboratory Standards Institute (CLSI) [26].

All experiments were performed in triplicate, and highest MIC values were considered as the

final results. To determine the effect of pulmonary surfactant on *in vitro* activity of RBx

10080758, RBx 1011591, the MIC was also determined in presence of pulmonary surfactant as

described earlier [9]

In vitro cytotoxicity assay

Cytotoxicity assay was conducted using MTT assay. HepG2 cells were seeded in a 96-well flat-bottom microtiter plate and allowed to adhere for 24 h at 37°C in a CO_2 incubator. After 24 h of incubation, cells were treated with 200 µl of the test compound. After 48 h of incubation, 12.5 µl of MTT working solution (5 mg/ml) was added and incubated again for 4 h. Finally, the intensity of the DMSO dissolved formazan crystals was quantified at 540 nm.

Combination study of RBx 10080758

Checkerboard assay was performed to evaluate their fractional inhibitory concentration (FIC) index in combinations against standard MRSA strain, and the assay were performed on 96-well plates as described previously [27, 28]. For example, combination A/B was used for interpreting the assay protocol [29]. The FICs were calculated as described earlier. The combining effect of A in combination with B against MRSA was interpreted as described earlier [28-30].

Time-kill study

The time-kill studies of RBx 10080758, linezolid, against MRSA-562, MRSA WCUH-29, *S. pneumoniae* MA-80 and *S. pyogenes* were performed as described earlier [22, 31]. In brief, cation-adjusted Mueller-Hinton broth (CA-MHB) was pre-warmed at 37°C, and the drug was added at concentrations ranging from $0.5 \times$ MIC to 8 µg/ml. Then the exponential-phase cultures were added to the culture flask ($\sim 5 \times 10^6$ cfu/ml). For *S. pyogenes* and *S. pneumoniae*, CA-MHB with 5% lysed horse blood and drug were inoculated with $\sim 5 \times 10^6$ cfu/ml, and the flasks were incubated at 35°C. The viable cell count was assessed at predefined time points. The time-kill kinetics were performed three independent times.

Frequency of resistance

The frequency of resistance was determined as described earlier [18, 32]. Bacterial inoculums of

each *S. aureus* strain were prepared in MHB, and 100 µl of each suspension was spread onto the

surface of MHA containing $4 \times$ and $8 \times$ MIC of RBx 10080758. After incubating for about 48 h

at 37°C, the number of colonies was counted and divided by the number of bacteria inoculated to

determine the frequency of resistance.

***In vitro* ADME studies**

For determination of intrinsic clearance in liver microsomes, the reaction mixture was prepared

by addition of 400 µl of NADPH regenerating solution, 25 µl of liver microsomes and made up

to volume of 995 µl with phosphate buffer (100 mM, pH 7.4). Then 5 µl (100 µM stock) of test

compounds was added to initiate the metabolic reaction at 37°C. Periodically 70 µl of aliquots

were withdrawn every 3 min for liquid chromatography-tandem mass spectrometry (LC-MS/MS)

estimation. The remaining parent compound was estimated at each time point from the samples

and was expressed as percentage of 0 min. The intrinsic clearance was expressed in units of

L/h/kg.

In vitro CYP inhibition assay was performed using a standard commercial recombinant CYPs kit

(BD-Gentest) as recommended by the company. Seven serial half-log dilutions of RBx

10080758 and standard inhibitor starting from 100 µM were prepared. The reaction was initiated

by mixing of 100 µl of test compound/standard inhibitor with 100 µl of enzyme substrate mix.

After incubation, fluorescence was read using a plate reader and IC_{50} was calculated.

Experimental animals and ethical approval

Specific pathogen free (SPF) and Sprague Dawley (SD) rats were used for efficacy studies, while

SPF SD rats and Beagle dogs were used for PK studies. The laboratory environmental conditions

were temperature: 22 ± 2°C; relative humidity: 40 to 60 % and day-light cycle: 12-h light/12-h

dark. All animal protocols (IAEC/2010/38, IAEC/2010/42, DS/IAEC-2012/006 and

IAEC/2010/77/08-E2/10) were approved by the Institutional Animal Ethics Committee.

Murine systemic infection model of *MRSA-562*

The minimum lethal dose of MRSA for systemic infection was determined as described earlier

[33, 34]. The *in vivo* efficacy of RBx 10080758 was determined in murine systemic infection

caused by MRSA 562 strain. The bacterial strain was grown in trypticase soy broth at 37°C. The

culture was mixed with equal volume of 5% hog mucin (Becton, Dickinson and Company) and

0.5 ml of the bacterial suspension (1-5 $\times$ 10^6 cfu/mouse, n=6-10/group) was injected by

intraperitoneal (IP) route into each mouse. RBx 10080758 was administered 2.5, 5, 10, and 20

mg/kg by IP route at +0.5, +4, +7, +10 h post infection (p. i) and linezolid 25 mg/kg was

administered orally at +0.5, +6 h p. i. for 1 day. The survival of animals was recorded daily for 8

days.

Organ load model

331 Additionally, to assess the microbiological cure of RBx 10080758, two satellite groups of mice

332 (n=4/group) in similar experimental conditions as described above, were kept and their kidney

333 bacterial loads were determined as described earlier [35]. Bacterial load was determined at 8 h

334 and 24 h p. i. in treatment group that received 5, 10, and 20 mg/kg of RBx10080758 and 25

335 mg/kg of linezolid.

Efficacy of RBx 10080758 in SD rat thigh infection

SD rats of either sex weighing 90 ± 10 g (n=5-6/group) were used for thigh infection model. The

overnight grown bacterial culture was diluted in MHB and 250 µl containing 1-3 x 10^7 cfu was

injected in the thigh muscles of each rat. Treatment was initiated 2 h p. i and a pre-control group

was sacrificed, and their thigh muscles were evaluated for initial bacterial loads. The efficacy of

RBx 10080758 (sodium salts) was evaluated at 10, 20 and 45 mg/kg (molar equivalent dose of

RBx 10080758) four times a day at +2, +5, +8, and +11 h p. i and linezolid was evaluated at 10

mg/kg by IV route. At the end of 24 h, the animals were sacrificed, and their thigh muscle was

processed for bacterial counts. In addition, the efficacy of RBx 10080758 at 45 mg/kg, q12h for

2 days by IV route was also evaluated against MRSA PVL-2 and WCUH-29 strains in rat thigh

infection model.

Rat lung infection model

To evaluate the efficacy of RBx 10080758 against respiratory tract pathogens, a rat lung

infection model was established using *S. pneumoniae* MA-80. SD rats of either sex weighing 80-

100 g (n=6-14) were used for lung infection. The overnight grown culture was suspended in PBS

and OD_{600nm} was adjusted to 0.3. The bacterial inoculum was further mixed with an equal

volume of 5% mucin. The pre-anesthetized SD rats with isoflurane, were inoculated intra-

tracheally with 1-5 x 10^6 cfu/rat as previously described [36]. The treatment of RBx 10080758

and linezolid was started 2 h p. i. with 45 mg/kg, q12h, IV for 1 day and telithromycin was

administered at 30 mg/kg by oral gavage. Following 12 h of last dose, lungs of each animal were

harvested aseptically and homogenized in 5 ml of PBS and enumerated the viable bacterial

counts.

In vivo **pharmacokinetic study**

Sprague Dawley rats and Beagle dogs were administered single dose of RBx 10080758 at 1

mg/kg by IV route. Serial blood sampling was carried out in rat and dog at different time points

up to 8 h post-dose. Plasma was harvested by centrifugation of blood and kept at -80^{o}C until

analysis. RBx 10080758 concentrations in plasma were determined using LC-MS/MS. The

pharmacokinetic data were analyzed using the NCA module of WinNonlin software (version

4.1).

Statistical analysis

All data were analyzed using GraphPad Prism Software (version 5.03; San Diego, CA). The

statistical significance between the treated and untreated control group was evaluated by

nonparametric Mann-Whitney analysis. The bar chart and scatter line graph presented as Means

$\pm$ SD, were calculated using Microsoft Office Excel. Survival curves were analyzed by log-rank

Mantel-cox test. A P value of <0.05 was considered statistically significant.

Funding: This study was financially supported by Ranbaxy and Daiichi Sankyo India Pharma

Private Limited, Gurgaon, India.

Transparency declarations: None to declare

Acknowledgements

Authors are thankful to Ranbaxy and Daiichi Sankyo India Pharma Private Limited for providing

necessary facility to carry out the studies.

(a) RBx 10080758

(b) RBx 10115912

391   **Figure 1.** Chemical structure of RBx 10080758(a) and RBx 10115912(b)

394
395
396
397
398 **Figure 2:** Time-kill curves of RBx 10080758 against MRSA 562 (a), MRSA WCUH-29 (b), *S.*
399 *pneumoniae* MA-80 (c) and *S. pyogenes* (d). Data represent mean ± SD from three independent
400 experiments.

406 **(a)**

407
408 **(b)**

409
410
Figure 3: Efficacy of RBx 10080758 in murine systemic infection model of MRSA-562. (a)
Kaplan Meier survival curve (n=6-10), (b) Bacterial load in the kidneys mouse (n=4) treated with
RBx 10080758 at 8 h after two doses and at 24 h after 4 four doses. Asterisks indicate a
significant difference compared with the result of the post-control group. Survival data were
analyzed by log-rank Mantel-Cox test. Kidney bacterial loads were compared using one-way
analysis of variance (ANOVA) with Dunnett's multiple-comparison posttest. (*$P < 0.05$, **$P <$
0.01, ***$P < 0.001$, ****$P < 0.0001$).

419
420
421

Figure 4: Efficacy of RBx 10080758 (Na salts) in rat thigh and lung infection models. Each bar represents the mean ± standard deviation (n=5-14). Rat thigh infection was performed with MRSA WCUH-29 with four times a day dosing regimen for 1 day treatment (**a**); MRSA PVL-2 with q12h x 2 days (**b**) and MRSA WCUH-29 with q12h x 2 days treatment (**c**); or intratracheal infection with *S. pneumoniae* MA-80 with q12h x 1 day (**d**). Treatment was initiated 2 h post-infection. Bacterial counts in each treated group were compared with those of pre- and post-control animals. The asterisks indicate a significant difference in bacterial counts after treatment compared to pre-and post-control (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$).

435 **Table 1:** IC$_{50}$ of RBx 10080758 and RBx 10115912 against *E. coli* DNA gyrase, Topoisomerase

436 IV and human Topoisomerase II

| Compound | IC$_{50}$ (µg/ml)[a] | | Human Topoisomerase II |
| | *E. coli* | | |
	DNA Gyrase	Topoisomerase IV	
RBx 10080758	0.06	0.06	>500
RBx 10115912	0.06	0.06	>500
Levofloxacin	54	7.5	>500

439   [a]All reactions were performed in duplicate and repeated twice.

440   The mean values were used to determine the IC$_{50}$.

Table 2: *In vitro* activity of RBx 10080758 and RBx 10115912 in comparison to other
antimicrobial agents against Gram positive clinical isolates

Organisms (no of strains)/new chemical entities/standard drugs	MIC (µg/ml)		
	MIC_{50}	MIC_{90}	MIC Range
Staphylococcus aureus (n=50)[1]			
RBx 10080758	0.03	0.06	0.015-0.06
RBx 10115912	0.25	0.5	0.125 - 1
Linezolid	2	2	1-2
Vancomycin	0.5	1	0.5 - 1
Ciprofloxacin	0.5	>16	0.25->16
Erythromycin	0.5	>16	0.25->16
Mupirocin	0.25	256	<0.125-256
Streptococcus pneumoniae (n=20)			
RBx 10080758	0.015	0.03	0.015-0.03
RBx 10115912	0.125	0.25	0.125–0.5
Linezolid	1	2	0.5- 1
Vancomycin	0.5	1	0.5–1
Ciprofloxacin	0.5	2	0.5-4
Erythromycin	0.06	16	0.06->16
Penicillin	0.03	0.25	0.03->4
Coagulase negative Staphylococci (n=35)[2]			
RBx 10080758	0.015	0.03	0.008-0.03
RBx 10115912	0.25	0.5	0.125 - 1
Linezolid	1	1	0.5-1
Vancomycin	2	2	1-2
Ciprofloxacin	0.5	>16	0.125->16
Erythromycin	0.25	>16	0.125->16
Streptococcus viridans group (n=20)			
RBx 10080758	0.008	0.03	0.008-0.03
RBx 10115912	0.25	0.5	0.125 - 4
Clindamycin	0.03	0.06	0.03-0.06
Amoxicillin	0.06	0.25	0.03-0.25
Erythromycin	0.03	0.06	<0.015-0.06
Streptococcus pyogenes (n=50)[3]			
RBx 10080758	0.03	0.03	0.008-0.0.06
RBx 10115912	0.25	0.5	0.125 - 2
Linezolid	1	2	0.5-2
Erythromycin	0.03	>4	0.015->4
Penicillin	0.008	0.008	<0.004-0.008
Ciprofloxacin	0.5	4	0.5-4
Clindamycin	0.03	4	0.015->4
Enterococci spp (n=27)[4]			
RBx 10080758	0.015	0.06	0.015-0.06

RBx 10115912	0.25	0.5	0.125 - 2
Linezolid	2	4	1-4
Vancomycin	1	>16	1>16
Amoxicillin	0.5	>16	0.25->16
***S. aureus* LZD[R] (n= 10)**			
RBx 10080758	0.03	0.06	0.015-0.06
RBx 10115912	0.25	0.5	0.125 - 1
Linezolid	32	>32	8->32
Vancomycin	1	2	0.5-2
Ciprofloxacin	1	>16	0.25->16
Erythromycin	2	4	0.25-8
***S. aureus* VISA or VRSA strains (n=21)[5]**			
RBx 10080758	0.03	0.03	<0.008-0.03
RBx 10115912	0.25	0.5	0.125 - 1
Linezolid	2	2	1-2
Vancomycin	4	4	4-16
Ciprofloxacin	0.5	16	0.25->16
Erythromycin	>16	>16	0.5 - >16

[1]Methicillin sensitive *S. aureus* (MSSA), methicillin resistant *S. aureus* (MRSA),
fluoroquinolone resistant (FQ[R]) *S. aureus*, Erythromycin[R] *S. aureus*, Mupirocin[R] *S.*
aureus, [2]Methicillin sensitive *S. epidermidis* (MSSE), methicillin resistant *S. epidermidis*
(MRSE), *S. haemolyticus* & other coagulase negative staphylococcus (Cons), 3.
Sensitive, erm & mef strains, 4. *E. faecalis* and *E. faecium,* 5. Vancomycin intermediate
S. aureus (VISA, n=19) and Vancomycin resistant *S. aureus* (VRSA, n=2)

455
456
457

Table 3: Frequency of resistance and MPC RBx 10080758 against *S. aureus*

Bacterial Strains	Frequency of Resistance		MIC (µg/ml)	MPC	MPC/MIC ratio
	$4 \times$ MIC	$8 \times$ MIC			
S. aureus WCU H 29	$<3.5 \times 10^{-9}$	$<3.5 \times 10^{-9}$	0.03	0.125	4
S. aureus MRSA 562	$<6.0 \times 10^{-9}$	$<6.0 \times 10^{-9}$	0.03	0.125	4

Table 4: Pharmacokinetics of RBx 10080758 (0-8 h) in rat and dog by intravenous route

Species	IV Dose (mg/kg)	C_{max} (µg/ml)	$t_{1/2}$ (h)	AUC_{inf} (µg.h/ml)	CL_p (ml/min/kg)	V_{ss} (l/kg)
Rat	1	5.19 ± 0.50	1.63 ± 0.50	1.50 ± 0.52	11.34 ± 0.49	0.17 ± 0.03
Dog	1	4.38 ± 1.63	1.14 ± 0.51	1.27 ± 0.30	13.67 ± 3.39	0.39 ± 0.12

477
References

1. Bozdogan B, Esel D, Whitener C, Browne FA, Appelbaum PC. Antibacterial susceptibility of a
vancomycin-resistant Staphylococcus aureus strain isolated at the Hershey Medical Center. J Antimicrob
Chemother. 2003;52(5):864-8; doi: 10.1093/jac/dkg457.
2. Rice LB. Antimicrobial resistance in gram-positive bacteria. Am J Med. 2006;119(6 Suppl 1):S11-9;
discussion S62-70; doi: 10.1016/j.amjmed.2006.03.012.
3. Kujath P, Kujath C. Complicated skin, skin structure and soft tissue infections - are we threatened by multi-
resistant pathogens? Eur J Med Res. 2010;15(12):544-53; doi: 10.1186/2047-783x-15-12-544.
4. Kumar M, Saadaoui M, Al Khodor S. Infections and Pregnancy: Effects on Maternal and Child Health.
Front Cell Infect Microbiol. 2022;12:873253; doi: 10.3389/fcimb.2022.873253.
5. Hardgrib N, Wang M, Jurik AG, Petersen KK. Life-threatening MRSA sepsis with bilateral pneumonia,
osteomyelitis, and septic arthritis of the knee in a previously healthy 13-year-old boy: a case report. Acta
Radiol Open. 2016;5(10):2058460116677180; doi: 10.1177/2058460116677180.
6. Augustine T, Kumar M, Al Khodor S, van Panhuys N. Microbial Dysbiosis Tunes the Immune Response
Towards Allergic Disease Outcomes. Clinical Reviews in Allergy & Immunology. 2022; doi:
10.1007/s12016-022-08939-9.
7. Kohno S, Yamaguchi K, Aikawa N, Sumiyama Y, Odagiri S, Aoki N, et al. Linezolid versus vancomycin
for the treatment of infections caused by methicillin-resistant Staphylococcus aureus in Japan. J Antimicrob
Chemother. 2007;60(6):1361-9; doi: 10.1093/jac/dkm369.
8. Rybak JM, Roberts K. Tedizolid Phosphate: a Next-Generation Oxazolidinone. Infect Dis Ther. 2015; doi:
10.1007/s40121-015-0060-3.
9. Silverman JA, Mortin LI, Vanpraagh AD, Li T, Alder J. Inhibition of daptomycin by pulmonary surfactant:
in vitro modeling and clinical impact. J Infect Dis. 2005;191(12):2149-52; doi: 10.1086/430352.
10. Azam MA, Thathan J, Jubie S. Dual targeting DNA gyrase B (GyrB) and topoisomerse IV (ParE)
inhibitors: A review. Bioorg Chem. 2015;62:41-63; doi: 10.1016/j.bioorg.2015.07.004.
11. Mathur T, Barman TK, Kumar M, Singh D, Kumar R, Khera MK, et al. In Vitro and In Vivo Activities of
DS-2969b, a Novel GyrB Inhibitor, against Clostridium difficile. Antimicrob Agents Chemother.
2018;62(4); doi: 10.1128/AAC.02157-17.
12. Barman TK, Kumar M, Chaira T, Gangadharan R, Singhal S, Rao M, et al. Potential of the fluoroketolide
RBx 14255 against Streptococcus pneumoniae, Neisseria meningitidis and Haemophilus influenzae in an
experimental murine meningitis model. J Antimicrob Chemother. 2019;74(7):1962-70; doi:
10.1093/jac/dkz119.
13. Durcik M, Tomasic T, Zidar N, Zega A, Kikelj D, Masic LP, et al. ATP-competitive DNA gyrase and
topoisomerase IV inhibitors as antibacterial agents. Expert Opin Ther Pat. 2019;29(3):171-80; doi:
10.1080/13543776.2019.1575362.
14. Lu J, Patel S, Sharma N, Soisson SM, Kishii R, Takei M, et al. Structures of kibdelomycin bound to
Staphylococcus aureus GyrB and ParE showed a novel U-shaped binding mode. ACS Chem Biol.
2014;9(9):2023-31; doi: 10.1021/cb5001197.
15. Stokes NR, Thomaides-Brears HB, Barker S, Bennett JM, Berry J, Collins I, et al. Biological evaluation of
benzothiazole ethyl urea inhibitors of bacterial type II topoisomerases. Antimicrob Agents Chemother.
2013;57(12):5977-86; doi: 10.1128/AAC.00719-13.
16. Gjorgjieva M, Tomasic T, Barancokova M, Katsamakas S, Ilas J, Tammela P, et al. Discovery of
Benzothiazole Scaffold-Based DNA Gyrase B Inhibitors. J Med Chem. 2016;59(19):8941-54; doi:
10.1021/acs.jmedchem.6b00864.
17. Bax BD, Chan PF, Eggleston DS, Fosberry A, Gentry DR, Gorrec F, et al. Type IIA topoisomerase
inhibition by a new class of antibacterial agents. Nature. 2010;466(7309):935-40; doi:
10.1038/nature09197.
18. Barman TK, Kumar M, Mathur T, Namba E, Singh D, Chaira T, et al. In Vitro and In Vivo Activities of
DS-2969b, a Novel GyrB Inhibitor, and Its Water-Soluble Prodrug, DS11960558, against Methicillin-
Resistant Staphylococcus aureus. Antimicrob Agents Chemother. 2018;62(6); doi: 10.1128/AAC.02556-17.
19. Gade ND, Qazi MS. Fluoroquinolone Therapy in Staphylococcus aureus Infections: Where Do We Stand?
J Lab Physicians. 2013;5(2):109-12; doi: 10.4103/0974-2727.119862.

20.   de Kraker ME, Stewardson AJ, Harbarth S. Will 10 Million People Die a Year due to Antimicrobial
Resistance by 2050? PLoS Med. 2016;13(11):e1002184; doi: 10.1371/journal.pmed.1002184.
21.   Jones RN, Fritsche TR, Sader HS, Ross JE. LEADER surveillance program results for 2006: an activity
and spectrum analysis of linezolid using clinical isolates from the United States (50 medical centers). Diagn
Microbiol Infect Dis. 2007;59(3):309-17; doi: 10.1016/j.diagmicrobio.2007.06.004.
22.   Barman TK, Kumar M, Mathur T, Chaira T, Ramkumar G, Kalia V, et al. In Vitro and In Vivo Activities of
a Bi-Aryl Oxazolidinone, RBx 11760, against Gram-Positive Bacteria. Antimicrob Agents Chemother.
2016;60(12):7134-45; doi: 10.1128/AAC.00453-16.
23.   Schwameis R, Erdogan-Yildirim Z, Manafi M, Zeitlinger MA, Strommer S, Sauermann R. Effect of
pulmonary surfactant on antimicrobial activity in vitro. Antimicrob Agents Chemother. 2013;57(10):5151-
4; doi: 10.1128/AAC.00778-13.
24.   Martinez MN, Papich MG, Drusano GL. Dosing regimen matters: the importance of early intervention and
rapid attainment of the pharmacokinetic/pharmacodynamic target. Antimicrob Agents Chemother.
2012;56(6):2795-805; doi: 10.1128/AAC.05360-11.
25.   Kumar M, Mathur T, Joshi V, Upadhyay DJ, Inoue SI, Masuda N. Effect of DS-2969b, a novel GyrB
inhibitor, on rat and monkey intestinal microbiota. Anaerobe. 2018;51:120-3; doi:
10.1016/j.anaerobe.2018.04.017.
26.   Institute. CaLS. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically.
Approved standard M7-A10. Clinical and Laboratory Standards Institute, Wayne, PA. 2015.
27.   Kim SH, Park C, Chun HS, Lee DG, Choi JK, Lee HJ, et al. Pilot Screening to Determine Antimicrobial
Synergies in a Multidrug-Resistant Bacterial Strain Library. Microb Drug Resist. 2016;22(5):372-8; doi:
10.1089/mdr.2015.0251.
28.   Kumar A, Khan IA, Koul S, Koul JL, Taneja SC, Ali I, et al. Novel structural analogues of piperine as
inhibitors of the NorA efflux pump of Staphylococcus aureus. J Antimicrob Chemother. 2008;61(6):1270-
6; doi: 10.1093/jac/dkn088.
29.   Odds FC. Synergy, antagonism, and what the chequerboard puts between them. J Antimicrob Chemother.
2003;52(1):1; doi: 10.1093/jac/dkg301.
30.   Sharma S, Kumar M, Sharma S, Nargotra A, Koul S, Khan IA. Piperine as an inhibitor of Rv1258c, a
putative multidrug efflux pump of Mycobacterium tuberculosis. J Antimicrob Chemother.
2010;65(8):1694-701; doi: 10.1093/jac/dkq186.
31.   Kumar M, Mathur T, Barman TK, Ramkumar G, Bhati A, Shukla G, et al. In vitro and in vivo activities of
the novel Ketolide RBx 14255 against Clostridium difficile. Antimicrob Agents Chemother.
2012;56(11):5986-9; doi: 10.1128/AAC.00015-12.
32.   Kumar M, Rao M, Purnapatre KP, Barman TK, Joshi V, Kashyap A, et al. DS86760016, a Leucyl-tRNA
Synthetase Inhibitor with Activity against Pseudomonas aeruginosa. Antimicrob Agents Chemother.
2019;63(4); doi: 10.1128/aac.02122-18.
33.   Barman TK, Rao M, Bhati A, Kishore K, Shukla G, Kumar M, et al. Non invasive real-time monitoring of
bacterial infection & therapeutic effect of anti-microbials in five mouse models. Indian J Med Res.
2011;134(5):688-95; doi: 10.4103/0971-5916.90997.
34.   Rao MB, T. K.; Kumar, M.; Mathur, T.; Shukla, G.; Bhati, A.; Kishore, K.; Chaira, T.; Pandya, M.;
Upadhyay, D.J. Efficacy of linezolid against Staphylococcus aureus in different rodent skin and soft tissue
infections models. Microbiol Res. 2011;2(1); doi: doi.org/10.4081/mr.2011.e3.
35.   Kumar M, Mathur T, Barman TK, Chaira T, Kumar R, Joshi V, et al. Novel FtsZ inhibitor with potent
activity against Staphylococcus aureus. J Antimicrob Chemother. 2021;76(11):2867-74; doi:
10.1093/jac/dkab270.
36.   Kumar M, Rao M, Mathur T, Barman TK, Joshi V, Chaira T, et al. Azithromycin Exhibits Activity Against
Pseudomonas aeruginosa in Chronic Rat Lung Infection Model. Front Microbiol. 2021;12:603151; doi:
10.3389/fmicb.2021.603151.

578