28 Minutes Later: A behavioral and transcriptomic investigation into the potential role of

aflatrem-like compounds in establishing *Ophiocordyceps* extended phenotypes in zombie

ants

William C. Beckerson,[a,b] Courtney Krider , [a] Umar A. Mohammad, [a] Charissa de Bekker,[a,c]

[a]Department of Biology, College of Sciences, University of Central Florida, Orlando, Florida,

32816

[b]Genomics and Bioinformatics Cluster, University of Central Florida, Orlando, Florida, 32816

[c]Microbiology, Department of Biology, Utrecht University, 3584 CH, Utrecht, The Netherlands

Keywords: secondary metabolites, staggers syndrome, zombie ants, machine learning,

transcriptomics

ABSTRACT

Parasites and their hosts often share intimate coevolutionary relationship that can lead to bizarre

phenotypes over time. *Ophiocordyceps* is one such genus that has evolved to manipulate the

behavior of its various insect and arachnid hosts, the clearest examples of which are found in ants

of the Camponotus tribe, colloquially known as "zombie ants". While the unusual behaviors

induced during infection are well described, the molecular driving force behind these changes are

still unknown. Recent genomic and transcriptomics analyses have identified several candidate

secreted proteins and secondary metabolites highly upregulated during infection. Of particular

interest amongst these compounds is an aflatrem-like ergot alkaloid related to the mycotoxin

aflatrem known to cause neurological issues in other animals, issues collectively called "staggers

syndrome". To test if aflatrem-like compounds play a role in *Camponotus floridanus* manipulation,

healthy ants were injected with purified aflatrem and their behavior was monitored for 30 minutes.

RNA-Seq was also performed on the heads of aflatrem-injected ants to compare any changes at

the transcriptomic level to data available for ants infected by *Ophiocordyceps camponoti-floridani*.

Behavioral assay footage was analyzed using the machine learning tool MARGO, which showed

a negative correlation between activity and speed, and dose of aflatrem. Behaviors were also scored

manually, identifying a positive coloration between aflatrem dose and staggering behavioral

phenotypes. RNA-seq discovered 262 genes significantly up or down regulated compared to sham

injected ants. Of these genes, 108 were also differentially regulated in the heads of late-stage

Ophiocordyceps-infected ants. Together, these results indicate that aflatrem-like compounds may

play an important role during late-stages of infection by reducing activity in ants, keeping them in

elevated microclimates optimal for the production and dispersal of spores.

1. INTRODUCTION

Parasites and their hosts often share an intimate coevolutionary history; associations that over time

can lead to unique phenotypes and subsequent host-specificities. Perhaps one of the more

remarkable symbiotic phenomena known to have convergently evolved in zootic parasites is the

ability to manipulate host behavior. Host manipulations provide an adaptive advantage to the

parasite through induction of specific behaviors that increase its reproduction success (Sánches

and Biron, 2019); behaviors that can be viewed as an example of extended parasitic phenotypes

(Dawkins, 1982). These adaptive phenotypes are different from sickness behaviors, which are

defined as actions exhibited by the infected individuals which are not conducive to the fitness of

the host nor the parasite (e.g., lethargy, in most cases). The difference between behavioral

manipulations and sickness behaviors can be observed in the parasitic flat worm *Leucochloridium*

paradoxum, which manipulates amber snails from the genus *Succinea*. Once inside a snail, the

worm drives the host to ascend nearby structures, a common manipulation behavior known as

"summit disease" (Wesołowska and Wesołowski, 2013). This summiting behavior makes snails

much more susceptible to predatory birds, the primary host of *L. paradoxum*, ultimately increasing

the parasite's likelihood of completing its lifecycle at the expense of its secondary host

(Wesołowska and Wesołowski, 2013). Furthermore, the return of the worm to its bird host allows

it to spread much more effectively through bird droppings, increasing its fitness and providing a

selectable pressure to maintain behavioral manipulation in the snails.

A variety of taxa are implicated in behavioral manipulations across the Tree of Life,

including crustaceans (e.g., *Loxothylacus panopaei* infecting crabs (Blakeslee et al., 2021)),

nematomorphs (e.g., *Spinochordodes tellinii* infecting katydids (Biron et al., 2005)), protozoans

(e.g., *Toxoplasma gondii* infecting mice (Tong et al., 2021)), insects (e.g., *Cotesia congregate*

parasitizing caterpillars (Adamo, 2019)), and fungi (e.g., *Entomophthora muscae* infecting flies

(Elya and Licht, 2021)), and are even found to occur in non-living viruses (e.g., *Rabies lyssavirus*

infecting mammals (Hueffer et al., 2017)). While behavioral manipulation includes a wide array

of tactics, many of these parasites induce summiting behavior to increase their transmission

chances (de Bekker et al., 2021; Araújo & Hughes, 2019; Steinkraus et al., 2017; Latchininsky et

al., 2016; Wesołowski & Wesołowski, 2013). Collectively, behavior-manipulating parasites are

referred to colloquially as "zombie parasites" due to the peculiar actions of infected hosts, many

of which are reminiscent of zombies from pop culture sci-fi media (e.g., movies such as *28 Days*

Later and *The Body Snatchers*).

Amongst these mind-altering parasites, the "zombie fungi" are one of the largest known

groups, infecting various species of insects and arachnids globally (Arruda et al., 2021; de Bekker

et al., 2021; Cooley et al., 2018). This includes wasps (e.g., *O. humberti* (Sobczak et al., 2020)),

beetles (e.g., *O. curculionum* (Shrestha et al., 2016)), scale bugs (e.g., *Ophiocordyceps clavulata*

(Petch, 1933)), caterpillars (e.g., *O. sinensis* (Guo et al., 2021)), flies (e.g., *O. dipterigena* (Chhetri

et al., 2020)), and spiders (e.g., *O. engleriana* (Sung et al., 2007)). While the genus is capable of

infecting a diverse group of hosts, a large portion of the currently described species infect and

manipulate carpenter ants of the Camponotini tribe with extreme host-specificity (Araújo et al.,

2018, Araújo et al., 2020). These *Camponotus*-infecting fungi belong to the species complex

Ophiocordyceps unilateralis sensu lato; one of the more well-studied groups of zombie fungi

known for their clear and tractable behavioral manipulations compared to other *Ophiocordyceps*

species.

Ophiocordyceps unilateralis species begin their lifecycle as a spore, ejected from their

parental fruiting body. These spores sail on the wind to infect new ants, either through immediate

contact or through the production and attachment of secondary capillispores after falling onto the

forest floor (Evans et al., 2018). Once inside the ant, the fungus begins to secrete an array of

biomolecules in a time-specific manner; some of which alter the behavior of their host (Will et al.,

2020; de Bekker et al., 2015) One of the first notable changes occurs as these ordinarily social

hosts begin to show a lessened ability to communicate and participate in foraging tasks (Trinh et

al., 2021). Subsequently, infected ants climb nearby vegetation and latch on with their mandibles

(Andersen et al., 2009). The fungus then completes its lifecycle through the formation of

anastomosis tubes and an endosclerotium that siphons the remaining nutrients from the host and

uses these materials to create a new fruiting body (Fredericksen et al., 2017). Closer observations

and quantification of behaviors in both the lab and the field have led to the classification of other,

more subtle behavioral changes including changes in daily rhythms (Trinh et al., 2021; Hughes et

al., 2011) and the loss of motor coordination and balance in ants that fumble their way around in

a manner described as by Hughes et al. as a "drunkard's walk" (Supplemental Video 1) (Trinh et

al., 2021; Will eat al., 2020; Hughes et al., 2011). These conspicuous behavioral changes caused

by the fungus during infection makes the *O. unilateralis* species complex and their hosts

particularly well-suited for the study of behavior-manipulating parasites in greater detail (de

Bekker et al., 2015).

Recent work establishing techniques for culturing the fungus in a laboratory setting and

manually infecting ants, as well as the genomic sequencing of several *O. unilateralis* species, has

laid the groundwork for research aimed at understanding the molecular framework driving these

behavioral changes (Will et al., 2020; de Bekker et al., 2017; de Bekker et al., 2015).

Transcriptomics analyses of the North American species *Ophiocordyceps kimflemingiae* and

Ophiocordyceps camponoti-floridani have identified secreted proteins and secondary metabolite

clusters highly upregulated during summiting behavior (de Bekker et al., 2015; Will et al., 2020).

Of particular interest is a secondary metabolite cluster with a TRP_DMAT backbone that had a

~5,900 (in *O. kimflemingiae*) and ~12,000-fold (in *O. camponoti-floridani*) increase in expression

during manipulation compared to fungal growth in culture (Will et al., 2020). This cluster was first

identified as an ergot alkaloid producing pathway based on the functional annotation of its

tryptophan dimethyltransferase (Pfam: TRP_DMAT (PF11991)) backbone gene (de Bekker et al.,

2015). More detailed gene-by-gene annotation also found homologs of genes within and

surrounding the cluster to gene that code for aflatrem production proteins in the plant pathogen

Aspergillus flavus (Will et al., 2020). Genomic comparisons with other *Ophiocordyceps* genomes

revealed that this complete secondary metabolite cluster appears to be conserved across species

within *O. unilateralis* but not in other *Ophiocordyceps* species outside the species (e.g.,

Ophiocordycpes australis s.l.)(de Bekker et al., 2017). This begs the question, what evolutionary

purpose may aflatrem-like products, particularly those produced by this cluster, serve for the

fitness of *Ophiocordyceps* species that manipulate the behavior of *Camponotus* hosts.

Aflatrem, a secondary metabolite of the indole-diterpene group called aflatoxins, is a potent

tremorgenic and carcinogenic mycotoxin known to cause neurological disorders in animals

(Nicholson et al., 2009; Duran et al., 2006). Cows exposed to aflatrem through ingestion of *A.*

flavus-contaminated grain feed exhibit neurological problems include muscle tremors and

hyperexcitability, followed shortly by incoordination, ataxia, and seizures (Selala et al., 2008;

Valdes et al., 1985). Collectively, these neurological symptoms linked directly to aflatrem's

positive allosteric modulation of Gamma-aminobutyric acid (GABA) receptors are now referred

to as "staggers syndrome" (Eldefrawi et al, 1990; Yao et al., 1989; Grant et al., 1987). The potential

involvement of GABA receptors in host manipulation of *Camponotus* species could have

important ramifications given their suspected role in caste and behavioral differentiation (Graham,

2018; Gospocic et al., 2017). The likelihood that an aflatrem-like compound may be employed

during the manipulation of carpenter ants is further supported by observations of the "drunkard's

walk" behaviors exhibited during infection, which are reminiscent of the staggers syndrome

detailed in vertebrates (Supplemental Video 1). We, therefore, hypothesize that aflatrem-like

compounds play a role in the behavioral manipulation of carpenter ants during *O. unilateralis*

infection.

In this study, we administered aflatrem to the Florida carpenter ant *Camponotus*

floridanus, the host of *O. camponoti-floridani*, using microinjections to mimic the *in vivo* secretion

of aflatrem-like compounds by fungal cells in the hemolymph during infection (Figure 1).

Figure 1. A project overview of (A) the treatment groups, each consisting of 7 individuals from 3

different colonies of *Camponotus floridanus*, and an example of microinjection, (B) the arena

setup for behavioral assays and parameters for the observational periods, and (C) the behavioral

analyses using machine learning and behavior scoring software.

Given the ease of its procurement and its previous use in other animal models, high purity pre-

synthesized aflatrem (>98%) was used in our experiments (TePaske et al., 1992; Trienens &

Rohlfs, 2011). Through the observation and quantification of various behaviors following

injection, we demonstrate that aflatrem on its own can induce behaviors consistent with both

those exhibited during infection by *O. camponoti-floridani* and those reported for vertebrates

with staggers syndrome. To investigate which host pathways are affected by aflatrem, and

potentially give rise to the exhibited behavioral phenotypes, we also conducted transcriptomics

analyses of aflatrem-injected individuals. Comparison of these results to transcriptomics data

from *Ophiocordyceps*-infected *C. floridanus* ants (Will et al., 2020) yielded several consistent

differentially expressed genes, including genes that encode neuromuscular and chemosensory

proteins.

2. MATERIALS AND METHODS

2.1 Ant Collection

Three separate colonies of *Camponotus floridanus* were collected using a DeWALT Cordless 2

Gallon Wet/Dry ShopVac at the University of Central Florida's arboretum, coordinates: Colony 1

(28.602748, -81.194269), Colony 2 (28.603918, -81.191682), and Colony 3 (28.601430, -

81.191251). Colonies were housed in 9.4 L plastic containers (Rubbermaid) lined with talcum

(Acros Organics), applied using a viscous 20% w/v solution of powder suspended in 100% ethanol

and let dry, to prevent ants from climbing the walls. Throughout the study, ants were provided

with fresh supplies of both double distilled water (ddH2O) and 15% sucrose solutions, *ad libitum*.

The housing was kept in an incubation chamber (Percival) set to the following program: 20°C –

0 lux – 80% relative humidity (RH) with a linear increase in temperature, light, and decrease in

humidity over 4 hours to reach 28°C – 2,100 lux – 65% RH, at which conditions were held for 4

hours followed by a linear decrease in the temperature, light, and increase in humidity, returning

to 20°C – 0 lux – 80% RH over 4 hours; and held for 12 hours.

2.2 Microinjections

To test our hypothesis and identify the effect of aflatrem-like compounds on *C. floridanus*,

we established a behavioral assay using microinjections to introduce aflatrem in a manner

biologically relevant to the secretion of aflatrem-like compounds by *Ophiocordyceps* species in

the hemolymph of their ant hosts (Figure 1). For these assays, we utilized high purity aflatrem,

>98% $C_{32}H_{39}NO_4$, manufactured from *A. flavus* by BioCrick Science Solution Specialists in non-

water-soluble crystal form, which we suspended in 100% acetone (Fisher Chemical) to a

concentration of 10 mg/mL. All injections were performed using borosilicate glass micropipettes

(Fisher) pulled with a glass puller (model: PC-100, Narishige) under the following parameters:

single step, all weights, heated to 62°C. Each ant was held by the dorsal side of the thorax before

piercing the carapace on the ventral side between the first and second pair of legs, followed by

slowly injecting 0.5 μL of solution into the ant while simultaneously releasing grip to alleviate

intracavitary pressure.

Because the aflatrem used in this study is non-water soluble, a preliminary trial was

performed to identify acetone tolerance in ants by injecting subjects with 100%, 50%, 25%, 10%,

and 5% acetone solutions diluted with phosphate buffered saline (PBS) (137 mmol NaCl, 2.7 mmol

KCl, 10 mmol Na_2HPO_4, 1.8 mmol KH_2PO_4, pH adjusted to 7.4), along with a 100% PBS solution

as a control. Prior to injection, containers housing ant colonies were removed from the incubator

and set on the lab bench top to acclimate to lab conditions for 10 minutes. All injections were

performed in triplicate, one ant per colony per treatment. Injected ants were placed into individual

60 mm x 15 mm petri dishes (USA Scientific) and returned to the incubator. Ants were kept under

observation for 48 hours post injection using an infrared GoPro Hero6 set to record time-lapse

footage with 60 second intervals and a CMVision IR30 Illuminator lamp for observation during

the night cycle. Dead and resting *C. floridanus* ants were easily distinguishable from one another

as dying ants coiled their legs and fell to one side upon death while resting ants remained prone

with legs supporting their bodies. Time of death for each ant was estimated by noting the frame in

which each ant exhibited death behaviors, allowing us to estimate time of death within an interval

of 1 minute. Any surviving ants after the 48-hour test period were euthanized by freezing at -20°C

for 24 hours. Results from these preliminary analyses indicated that a PBS/10% acetone solution

was the highest concentration for which there were no statistical differences in average

survivability compared to PBS-injected and non-injected controls.

Using the tolerable PBS/10% acetone standard, a second preliminary trial was performed

to test the survivability of ants post injections and to determine if any observed behavioral

phenotypes may be attributed to "dying behaviors" resulting from aflatrem exposure. To do so, 7

ants were injected with 0.5 μL of a PBS/10% acetone solution containing 500 ng aflatrem. This

was the highest dose possible using the 10 mg/mL aflatrem stock diluted with PBS. Post injection,

ants were recorded for 48 h using the same GoPro setup as the previous preliminary test. Resulting

behaviors were compared to 7 ants injected with a PBS/10% acetone solution without aflatrem and

7 ants injected with a 100% PBS solution. No differences in mortality were observed, with all

individuals surviving for at least 24 h and an average time of death arising at 34 h, indicative of

desiccation from lack of water.

Confident that neither 10% acetone nor 500 ng aflatrem killed the ants or induced death-

related behaviors, an assay to determine the effects of aflatrem on ant behavior was conducted

using PBS/10% acetone solutions containing the following doses of aflatrem; 10 ng, 50 ng, 100

ng, 500 ng. For each dose of aflatrem, 7 ants from each of the 3 colonies were injected (N=21, per

dose) for a total sample size of N=126. Ants were individually lifted from their housing by hand

wearing premium disposable neoprene gloves (NeoTouch) and placed into individual 60 mm x 15

mm petri dish (USA Scientific) arenas on the lab benchtop to acclimate for 10 minutes. These

arenas were fitted with a thread of twine spanning the dish from the bottom of one side to the top

of the other, providing a substrate for biting and more dexterous climbing behaviors (Figure 1-B).

To allow ants to walk around the perimeter of the dish unimpeded, the twine was pressed flat and

taped to the bottom corner of the petri dishes using a small piece of tape (Scotch). To avoid any

behavioral changes that may be induced via chemical ques left by prior occupants, a new arena

was created for each test subject (N=126). After the 10-minute acclimation period, ants were

injected with aflatrem following the microinjection protocols established in the preliminary trials.

The doses of aflatrem used were based on approximate ant equivalent weights, verified by

measuring *C. floridanus* body volume through length and weight following each experiment.

While body lengths were measured to the nearest 0.5 mm for each individual, weight was measured

in groups of seven ants corresponding to their treatment and colony ID. Measurements were

recorded immediately following euthanization at -20°C, preserving the approximate water weight

for each ant. To account for the approximate weight of solution injected into each ant, 3.5 mg was

subtracted from the total weight of each group. Based on an average 8.28 mg weight and 6.68 mm

length calculated from measurements of all 126 subjects used in this study (Supplemental Table

1), the four doses of aflatrem that were used are approximately 6.04e-5, 1.21e-5, 0.60e-5, and

0.12e-5 ant equivalents. These ant equivalent doses are consistent with pervious works testing

insecticidal compounds in ants (Sakamoto & Goka, 2021; Buczkowski & Woosler, 2019; Klotz &

Moss, 1996).

Ant behaviors were recorded for 30 minutes post injections using an iPhone 13 pro. Any

ants exhibiting signs of physical injury from the injection, e.g., immobility in the hind legs or body

curling, were euthanized immediately and a new subject and arena were used in its place. Ants

were recorded three or four arenas at a time in a rolling format to ensure high quality resolution at

60 fps, generating video files of approximately 45 minutes in length. At the end of each recording

period, arenas containing resting ants were tapped several times to induce movement, confirming

the subjects were still alive. All instances of tapping elicited activity.

2.3 Video scoring of behavioral assays post injection

Videos were converted from .MOV to .mp4 files using iMovie for editing in Movavi Video Editor

Plus 2022, which, unlike iMovie, allows for cropping files to show only single arenas. The cropped

files were additionally trimmed to exactly 30 minutes of footage beginning as soon as each arena

was positioned in the field of view. The names for each file were then blinded before manual

scoring by two observers using the behavior tracking software CowLog3 (Pastell, 2016). The

observers were trained using eight test run videos; four of ants injected with PBS/10% acetone/500

ng aflatrem, and four of ants injected with PBS. These videos were made during proof-of-principle

testing and are not included in the data presented in this study. Results generated for these practice

runs were then reviewed together by both observers to form a consensus on scoring methodology.

From these training videos, the observers identified eight scorable behaviors: Climbing Wall,

Climbing Top, Climbing Rope, Walking, Staggering, Grooming, Resting, and Biting

(Supplemental Video 2). While most of these behaviors are self-explanatory, "staggering" was

generally defined as a slow, sluggish walking behavior with repeated lateral movements

(Supplemental Video 3). Behaviors were also marked as "staggering" when trouble balancing,

listing to one side, falling over, walking in tight circles, flipping over, or dramatic wagging of the

gaster were observed (Supplemental Video 3). Each observer also kept notes on unique behaviors

observed during the scoring period, including walking backwards, lunging/bucking, and two

instances of convulsions (Supplemental Video 4).

Blinded videos were assigned to the observers in a randomized block format (Supplemental

Figure 1-A) with each observer scoring three videos at random per treatment group per colony.

The seventh video for each treatment group was scored by both observers and used as a formative

metric to ensure consistent scoring standards throughout the evaluation period. While most

behaviors were readily distinguishable, resulting in very similar scores between observers, walking

and staggering provided the most discrepancy. We therefore used staggering as the primary

measurement to evaluate scoring consistency across observers (Supplemental Figure 1-B).

In addition to the manual observations, videos were also analyzed using the MATLAB-

based machine learning tool Massively Automated Real-time GUI for Object-tracking (MARGO)

(Werkhoven et al., 2019). This tool was used to measure the speed and distance traveled by each

subject. Speed was only measured during activity by calculating the distance between centroid

coordinates in mms traveled per second during movement. Distance traveled was measured as a

total value for distance between centroid coordinates over the entire 30 min duration. To optimize

tracking for the field of view in each video, the following metrics were used: region of interest

(ROI) was set to circular shape and manually overlayed around the arena, background referencing

was set to detect dark objects on light backgrounds, tracking threshold was set between 70-80,

distance of the arena was set to 60 mm using the custom measuring tool, frame rate was set to 30,

and min/max area was set to 5 and 80, respectively.

The results from the PBS/10% acetone controls were first compared to those from the

100% PBS controls to identify any potential behavioral phenotypes induced by exposure to

acetone. Significant differences between the two groups were tested using a two-sample t-test in

RStudio v1.3.959. To test for significant behavioral changes in the aflatrem treatment groups, a

linear regression model using the lme4 package in RStudio was performed to compare the

PBS/10% acetone control (0 ng aflatrem) with PBS/10% acetone containing 10 ng, 50 ng, 100 ng,

and 500 ng aflatrem treatments.

2.4 RNA sequencing

To understand the effects of aflatrem exposure in *C. floridanus* at the genomic level, a total of 10

ants were injected with either PBS/10% acetone (control group, N=5) or PBS/10% acetone/500 ng

aflatrem (treatment group, N=5). Ants were each placed into their own petri dish arena for 4

minutes post injection. This timepoint was selected based on the scaling levels of activity observed

in our behavioral assays (Supplemental Figure 2). Ants were then quickly placed into 2 mL

microcentrifuge tubes and snap frozen in liquid nitrogen. Subsequently, for each ant, the head was

removed using a forceps and placed in a pre-frozen 2.0 mL Self-Standing Impact Resistance

Microtube containing 2 Wheels Manufacturing Grade 25 5/32" metal ball bearings for mechanical

tissue disruption in a SPEX® SamplePrep 1600 MiniG at 1500 rpm for 30 sec. Disrupted samples

were defrosted while dissolving in 500 uL TRIzol (Ambion) to extract RNA followed by sample

clean up using the Qiagen RNeasy® MiniElute® Cleanup Kit (Will et al., 2020). The resulting

RNA was stored at -80C and shipped to Azenta US for Next Generation Sequencing. Libraries

were prepared by Azenta US using PolyA selection and sequenced in duplicate on Illumina HiSeq

with 150 bp paired end reads, resulting in ~350M reads. The quality of resulting paired RNA-seq

reads was verified using FastQC (Andrews 2010). We used Trimmomatic (Bolger et al., 2014) to

trim reads for adapters and quality using the following parameters: Leading: 3, Trailing: 3,

MinLen: 36. Transcripts were then aligned to the *Camponotus floridanus* reference genome (Gene

bank ascension number GCA_003227725.1) using Spliced Transcripts Alignment to a Reference

(STAR) 2.7.10a (Dobin et al., 2013). The resulting .bam files were checked for quality using the

standard metric of uniquely mapped reads > 70%, sorted, and indexed using Samtools (Danecek

et al., 2021). The number of gene hits for each sample were calculated using FeatureCounts

following the suggested counting of strictly uniquely mapped reads within exonic regions (Liao et

al., 2019). The quality of the resulting files was validated using a benchmark of 15-20 million

reads per sample. The hit counts were normalized using DESeq2 (Love et al., 2014) after removing

low level expression (counts less than 100 per gene, i.e., an average less than 10), eliminating

3,227 genes from further analysis. We performed differential expression analysis on the remaining

10,443 genes using DESeq2 with an alpha value of 0.05 and functionally annotated the

differentially expressed genes using Blast2GO (Conesa et al., 2005).

3. RESULTS

3.1 The behavioral effects of aflatrem in *C. floridanus* ants

Before we could analyze the behavioral effects of aflatrem in ants, we screened for any potential

behavioral changes caused by the presence of 10% acetone in the injections. Using an independent

two sample t-test to compare the behavior of ants injected with PBS to those injected with

PBS/10% acetone, we were able to rule out any significant differences in behavior between the

two groups: rest ($p = 0.787$), speed ($p = 0.997$), distance ($p = 0.947$), stagger ($p = 0.205$). Even

though neither group was exposed to aflatrem, we unexpectedly observed a few instances of

staggering in both. To determine if a small amount of staggering behavior is normal in ants, which

would need to be accounted for in our results, we reviewed the footage at timestamps logged for

these staggering events with an increased level of scrutiny. Upon closer review, we discerned that

nearly all events of staggering observed in both groups fell into the "gaster wagging" sub-type of

staggering (Supplemental Video 2), and nearly always occurred in the first 30 seconds of the

recording. This trend was also observed in the aflatrem treatment groups, indicating that this

particular type of staggering behavior is likely a transient physical response to the trauma resulting

from the injection process. This was confirmed by a quick follow-up experiment pricking ants with

empty glass capillaries, which also induced gaster wagging.

Confident that behaviors were not significantly affected by the addition of 10% acetone in

the injections, we compared behaviors across increasing doses of aflatrem. A linear regression

model showed that time spent resting in the arena was positively correlated with increasing doses

of aflatrem ($F_{4,105} = 5.54$, $p = 2.05e\text{-}2$) (Figure 2-A), while the average speed of aflatrem-treated

ants was negatively correlated ($F_{4,105} = 4.91$, $p = 2.89e\text{-}2$) (Figure 2-B). However, the average

distance traveled did not correlate with aflatrem treatment ($F_{4,105} = 3.036$, $p = 8.44e\text{-}2$, ns) (Figure

2-C). Finally, bouts of staggering did positively correlate with increasing doses of aflatrem ($F_{4,105}$

$= 36.65$, $p = 2.32e\text{-}8$) (Figure 2-D).

Figure 2. Behavioral results collected from *C. floridanus* ants injected with increasing doses of

aflatrem. Each boxplot depicts the quartiles of the distribution for each treatment with ants from

colonies 1, 2, and 3 shown as purple, light blue, and dark blue dots, respectively. A) shows the

proportion of time spent resting for each ant, B) shows their average speed during activity in

millimeters per second, C) shows their total distance traveled in meters, and D) shows the

proportion of activity spent staggering for each ant. The linear regression model results are shown

in the center of the graph for each behavior. A & D) both contain pie charts illustrating the relative

duration of each behavior, while B) contains speedometers for each treatment and C) presents the

median tracer plot from MARGO for each group.

3.2 RNA-seq results

With injection of *C. floridanus* with aflatrem resulting in a quantifiable phenotype, we next asked

which genes could be underlying the emergence of staggering phenotypes. As such, we conducted

transcriptomics to compare gene expression in the heads of ants treated with PBS/10% acetone/500

ng aflatrem (treatment group) to that of ants treated with PBS/10% acetone solutions (control

group). We found 262 genes that were significantly differentially expressed (Supplementary File

1). Of these genes, 148 were upregulated and 114 were downregulated in the treatment group

(Figure 3). While a comparable number of genes were up- and down-regulated in our treatment

group, the largest changes in expression were found among the downregulated genes (Figure 3).

We found the largest increase of expression (~22-fold increase) in a gene encoding an

aminopeptidase N-like protein, a common cell surface hydrolase involved in a variety of cellular

processes (Nocek et al., 2008), while to the largest decrease in expression (~512-fold decrease)

was exhibited by a gene coding for a cytochrome P450 6k1, a protein involved in detoxification

of insect cells (Xing et al., 2021) (Figure 3).

Figure 3. Volcano plot for the differential expression of 10,443 *Camponotus floridanus* genes in ants treated with PBS/10% acetone/500 ng aflatrem compared to the PBS/10% acetone control group. Dots in red represent non-significant changes in gene expression, while dots in blue represent significant change using an alpha value of 0.05. Functional annotations are shown for genes with adjusted *P* values less than 5e-5 and an absolute $\log_2$ fold changes lower than -6 or greater than 3.5.

Of all differentially expressed genes, 31 had a greater than 4-fold change in expression in the heads of aflatrem-treated ants. Among these genes were several that have been predicted to

play a role in neuromuscular systems, e.g., a downregulated *anoctamin-4*, which codes for an ion

channel protein involved in Ca^{2+}-dependent conductance (Reichhart et al., 2019), and *membrane*

metallo-endopeptidase (*Neprilysin*), a transmembrane protein found in neurological tissue that

plays a role in tissue perception and preservation (Obulesu, 2019), slightly upregulated in our

aflatrem-treated ants but significantly downregulated in *Ophiocordyceps*-infected ants. This group

also contained many olfactory-related genes, including 4 general odorant-binding protein genes

(OBPs) (i.e., *OBP19a*, *OBP56a*, and 2 *OBP69a*), all of which were downregulated. Such proteins

are located in the antennal sensillary lymph of insects and bind to pheromones that play a role in

insect communication and interactions (Gaubert et al., 2020). Moreover, we found 3 significantly

downregulated *ejaculatory bulb-specific* genes, involved in the release of pheromones in insects

(Pelosi et al., 2018), and a downregulated *sensory neuron membrane protein* gene, which has been

found to play a role in pheromone sensitivity in the moth *Heliothis virescens* (Pregitzer et al.,

2014).

We also found expression signatures of neuromuscular gene dysregulation in the larger

group of 231 differentially expressed genes with a less than 4-fold degrees of change. Amongst

them were 2 upregulated *titins* genes, which code for very large proteins that play an essential role

in sarcomere function (Freiburg et al., 2000; Zhang et al., 2000). We also found an upregulated

ryanodine receptor gene, responsible for releasing Ca^{2+} in the smooth endoplasmic reticulum of

muscle cells during excitation-contraction coupling (Lanner et al., 2010; Shakiryanova et al.,

2007). Furthermore, we detected a downregulated *caveolin-3*, which codes for a scaffold protein

to organize signaling molecules in the membranes surrounding muscle cells (Dewulf et al., 2019;

Galbiati et al., 2000-2013), and 2 downregulated *myosin ID heavy chain-like protein* genes, the

motor proteins of muscles (Wells et al., 1996) (Supplementary File 1). Also of interest, given our

previous exploration of the time-dependent nature of behavioral manipulation and the disruption

of clock-dependent foraging behaviors in *Ophiocordyceps*-infected ants (de Bekker & Das, 2022;

Will et al., 2020; de Bekker et al., 2015), was a gene coding for a circadian clock-controlled protein

with a Haemolymph Juvenile Hormone Binding Protein Pfam domain showing an 18-fold decrease

in expression. These proteins, affected by circadian rhythms in the ant, act as a carrier for juvenile

hormone to its target tissues (Zalewska et. Al., 2009).

Because the behaviors of aflatrem-infected ants are reminiscent of the extended behavioral

phenotype of *Ophiocordyceps*-infected ants, and due to expression data that demonstrates an

aflatrem-like secondary metabolite pathway is highly upregulated during late-stage infection of

carpenter ants by *Ophiocordyceps unilateralis* species (de Bekker et al., 2015; Will et al., 2020),

we also compared our RNA-seq results to transcriptomics data obtained from *C. floridanus*

manipulated by *O. camponoti-floridani* (Will et al., 2020). From this comparison, we found that

108 of the 262 differentially expressed genes observed in aflatrem-treated ants were also found to

be differentially regulated in late-stage *O. camponoti-floridani*-infections (i.e., Alive/Control and

Dead/Control, Supplementary File 1). Of these genes, 34 were differentially regulated in the same

direction, while 74 shared an inverse relationship. The similarly expressed genes contained the

previously mentioned *cytochrome P450*, a *sensory neuron membrane protein* encoding gene,

OBP56a, and *caveolin-3* genes, as well as an upregulated *nuclear hormone receptor E75*, a

circadian clock steroid-producing gene discovered in *Drosophila* linked to the maintenance of

circadian rhythms during stress (Kumar S et al., 2014). Additionally, *neurogenic locus protein*

delta, capable of coding for multiple translational products including neural differentiation factors

and growth hormones (Kopczynski et al., 1988), was upregulated during *Ophiocordyceps* infection

and isolated aflatrem exposure.

Among the genes that were differentially expressed in inverse directions versus the fungal

manipulation dataset, we found the aforementioned *OBP19a-like* gene, *cytochrome P450*, and

ejaculatory bulb-specific protein 3, as well as an *odorant receptor coreceptor*, which is also

expressed in chemosensory organs and forms heteromultimeric complexes with odorant receptors

(Mukunda et al., 2014). The gene set also contained a *short neuropeptide F*, which is implicated

in the regulation of animal behaviors including circadian rhythm and hunger (Lee et al., 2004) and

a *netrin receptor UNC5C*, which aids in the formation of axon connections through axon extension

and cell migration (Kim & Ackerman, 2011). We also found a *receptor-type tyrosine-protein*

phosphatase delta (*PTPRD*), part of the PTP family involved in many known signaling pathways

for which delta is shown to promote neurite growth and regulate axon guidance (Tomita et al.,

2020). Moreover, in both infection and sole aflatrem exposure, *sialin*, a sialic acid transporter-

coding gene that prevents neurodegenerative disorders caused by accumulation of sialic acid

(Tarailo-Graovac et al., 2017), was dysregulated, as well as *obscurin*, which, like *titin*, codes for

another large protein expressed in skeletal muscles that anchors myofibrils to the sarcoplasmic

reticulum for muscle contraction (Feher, 2017). Both datasets also indicate the dysregulation of a

juvenile hormone acid O-methyltransferase-like gene, which codes for an enzyme that converts

precursor molecules into juvenile hormones, regulating many aspects of insect physiology

(Shinoda & Itoyama, 2003). Finally, the two data sets also shared 26 genes with statistically

significant changes in expression coding for proteins of unknown function.

4. DISCUSSION

While sickness behaviors are common in animals infected by parasites, only behaviors that confer

an adaptive advantage and increase the reproduction and/or transmission of parasites are true

behavioral manipulations. It is, therefore, imperative that behaviors exhibited during parasitism be

thoroughly vetted before causal relationships are drawn between parasite genotypes and resulting

host phenotypes. This study aims to test the previously implicated involvement of aflatrem-like

compounds in the establishment of *Ophiocordyceps* extended phenotypes in zombie ants (Will et

al., 2020) by exposing healthy *C. floridanus* ants to high purity aflatrem and quantifying the

behavioral effects, a hypothesis formed from the tremor-like staggers that have been observed in

both *Ophiocordyceps*-infected ants and mammals following ingestion of aflatrem through the

consumption of *A. flavus*-infected feed, (de Bekker et al., 2017; Selala et al., 2008; Valdes et al.,

1985). As such, we observed and quantified staggering, which we found to be positively correlated

to aflatrem dose. This suggests that this behavior is caused by the presence of aflatrem. To rule out

other possible explanations for the staggering behavior, such as death behaviors that may

contradict its status as a true behavioral manipulation, we performed a longevity assay to

demonstrate that aflatrem does not kill *C. floridanus*. This makes it unlikely that the aflatrem-like

compound produced by *Ophiocordyceps* is used as a killing compound, but is rather employed to

induce staggering behaviors prior to death. We were further able to demonstrate that aflatrem has

a temporary yet significant effect on the mobility and coordination of ants and decreased overall

activity. Ants injected with higher doses of aflatrem moved much slower and had more difficulties

participating in climbing behaviors, particularly the more dexterous activities involving the twine

in the arena.

While we do not yet have a direct answer as to how the behaviors observed in this study

confer an adaptive advantage to the fungal parasite during late-stage infection, time-specific

hypotheses regarding the situational use of such a compound provide a presumptive explanation.

Indeed, a reduction in overall activity may seem contradictive to the goal of positioning the host

away from the nest at a higher elevation; however, if we consider the temporal variation and time-

specific expression of fungal genes that have been hypothesized to play a vital role throughout the

infection process, the induction of immobility during late-stage manipulation, during which natural

"drunkard walking" is observed, could have some biological relevance. It is reasonable to suggest

that, prior to the upregulation of aflatrem-like compounds, genes responsible for producing

effectors that induce hyperactivity and climbing behaviors are first produced. This would lead the

host up nearby vegetation and away from aggressive conspecifics of the nest enacted as part of

social immunity before late-stage production of compounds would be required to maintain the ant

in a desirable location (Cremer et al., 2007). In support of this hypothesis, transcriptomics data

from *Ophiocordyceps*-infected ants show that a *protein tyrosine phosphatase* (*ptp*) is upregulated

during manipulation (de Bekker et al., 2015; Will et al., 2020), a gene found to induce hyperactivity

and enhanced locomotion in caterpillars that are infected with a behavior-manipulating baculovirus

(Han et al., 2015). Moreover, enhanced locomotion has been observed in *Ophiocordyceps*-infected

ants during the stages leading up to the final summiting behavior (Trinh et al., 2021). Once away

from the nest, following successful completion of summiting behaviors, the upregulation of

compounds that reduce activity would become vital to maintain the host at optimal microclimates

where conditions suitable for fruiting body production.

In addition to reducing the ant's overall activity, aflatrem-like compounds may also aid in

the positioning of the host for the final and perhaps most iconic behavioral manipulation observed

in zombie ants - the "death grip" - by inhibiting the host's motor capabilities. Under the time-

specific scenario that we propose, the reduction of climbing behavior in concert with reduced

dexterity of the host could aid the fungus in the completion of its lifecycle in a manner that confers

a fitness advantage, as killing the host without proper adherence to the substrate would likely result

in the ant falling to the forest floor, away from the microclimate that promotes fruiting body

formation (Andersen et al., 2012; Andriolli et al., 2019; Will et al., under revision) and negating

the benefits of elevation position for spore dispersal. Field observations of body position for

Ophiocordyceps-infected *Camponotus* species corroborate this hypothesis. In addition to the

"death grip" adherence with their mandibles, infected *Camponotus castaneus* and *C. floridanus*

ants infected with *Ophiocordyceps kimflemingiae*, and *O. camponoti-floridani*, respectively, are

often found wrapping their limbs around the substrate (Figure 4) (Loreto et al., 2018; Will et al.,

2020).   Therefore, the discoordination of the ant host, as indicated by the dysregulation of

neuromuscular and sensory genes in this and previous studies, could play a role in initiating these

hugging behaviors as a prerequisite to biting or ensure adherance in case biting fails during the

final stages of infection (Will et al., 2020).

Figure 4. Examples of "hugging" behavior portrayed by *Ophiocordyceps*-infected *Camponotus*

floridanus ants during the death grip observed in a natural setting.

Even though the exact structure and concentrations of the aflatrem-like compounds

produces by *Ophiocordyceps in vivo* are still unknown, this study was able to replicate the

"drunkard's walk" phenotypes induced in the host during *Ophiocordyceps* infection, as well as

identify significant changes in the expression of 108 previously identified candidate manipulation genes in response to aflatrem exposure. While several of these genes were differentially regulated when compared with previously published transcriptomics data for *Ophiocordyceps*-infected ants (Will et al., 2020), the presence of changes in these genes nevertheless demonstrates their importance as targets for behavioral manipulations. Future work characterizing the 3-dimensional structure of the actual aflatrem-like compound produced by *O. unilaterial* species will shed further light on the role of these alkaloids in behavior manipulation and its precise physiological effects in the host. Even so, this work presents a hypothetical mechanism for time-specific expression of ergot alkaloids for the disruption of host activity in conjunction with summit disease that could play a vital role in facilitating the eventual attachment of the host at an elevated vantage point. By comparing the behaviors and gene expression profiles between ants treated with aflatrem and ants infected with *Ophiocordyceps*, we have glimpsed into the evolutionary purpose that secondary metabolites, upregulated during manipulation, may serve in *Ophiocordyceps* zombie fungi. As such, integrative behavioral studies like this one can bring us one step closer to characterizing the molecular mechanisms that drive zombie-making fungi. Furthermore, the protocols established as part of this research provide a framework for future studies that wish to examine the role of putative effectors produced and secreted by members of the *Ophiocordyceps* species complex, paving the way for further characterization of compounds produced by *Ophiocordyceps* as well as other behavior-manipulating species and their respective insect hosts.

AKNOWLEDGMENTS

We would like to extend our thanks to the King lab for their advice on using, and access to, their ShopVac as a method to collect ants, Biplabendu Das for his recommendations and assistance with

the CowLog3 and MATLAB programs, and Ian G. Will for his knowledgeable insight into the *O. camponoti-floridani* ergot alkaloid gene cluster and advice on statistical analysis of our data. This work is supported by NSF PRFB Award 2109435 to William C. Beckerson and NSF CAREER Award 1941546 to Charissa de Bekker.

Author Contributions:

WCB and CdB designed the experiment, analyzed the data, performed the RNA extractions and wrote the paper. WCB, CK, and UAM collected the ants and performed the injections and video recordings. CK and UAM performed the behavioral analyses using CowLog. WCB ran the MARGO analyses and performed the statistical analyses.

REFERENCES

Adamo SA. (2019). Turning your victim into a collaborator: Exploitation of insect behavioural control systems by parasitic manipulators. *Current Opinion in Insect Science*. 33: 25-29. https://doi.org/10.1016/j.cois.2019.01.004

Adamo SA, Easy RH, Kovalko I, MacDonald J, McKeen A, Swanburg T, Turnbull KF, Reeve C. (2017). Predator exposure-induced immunosuppression: trade-off immune redistribution or immune reconfiguration? *Journal of Experimental Biology* 220, 868-875. https://doi.org/10.1242/jeb.153320

Andersen SB, Gerritsma S, Yusah KM, Mayntz D, Hywel-Jones NL, Billen J, Boomsma JJ, Hughes DP. (2009). The life of a dead ant: the expression of an adaptive extended phenotype. *The American Naturalist* 174(3): 424-33. https://doi.org/10.1086/603640

Andrews S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data

[Online]. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Araújo JPM, Evans HC, Fernandes IO, Ishler MJ, Hughes DP. (2020). Zombie-ant fungi cross

continents: II. *Myrmecophilous hymenostilboid* species and a novel zombie lineage.

Mycologia 112(6): 1138-1170. https://doi.org/10.1080/00275514.2020.1822093

Araújo JPM, Hughes DP. (2019). Zombie-ant fungi emerged from non-manipulating, beetle-

infecting ancestors. *Current Biology* 29(21): 3735-3738.e2.

https://doi.org/10.1016/j.cub.2019.09.004

Araújo JPM, Evans HC, Kepler R, Hughes DP. (2018). Zombie-ant fungi across continents: 15

new species and new combinations within *Ophiocordyceps. L. Myrmecophilous*

hirsutelloids species. *Studies in Mycology* 90: 119-160

https://doi.org/10.1016/j.simyco.2017.12.002

Arruda IDP, Villanueva-Bonilla GA, Faustino ML, Moura-Sobczak JCMS, Sobczak JF. (2021).

Behavioral manipulation of the spider *Macrophyes pacoti* (Araneae: Anyphaenidae) by

the araneopathogenic fungus *Gibellula* sp. (Hypocreales: Cordycipitaceae). *Candian*

Journal of Zoology 99(5): 401-408. https://doi.org/10.1139/cjz-2020-0232

Biron DG, Marché L, Ponton F, Loxdale HD, Galéotti N, Renault L, Joly C, Thomas F. (2005).

Behavioural manipulation in a grasshopper harbouring hairworm: a proteomics approach.

Proceedings of the Royal Society of Biological Sciences 272(1577): 2117-2126.

https://doi.org/10.1098/rspb.2005.3213

Blakeslee AMH, Pochtar DL, Fowler AE, Moore CS, Lee TS, Barnard RB, Swanson KM, Lukas

LC, Ruocchio M, Torchin ME, Miller AW, Ruiz GM, Tepolt CK. (2021). Invasion of the

body snatchers: the role of parasite introduction in host distribution and response to

salinity in invaded estuaries. *Proceedings of the Royal Society of Biological Sciences*

288(1953): 288: 20210703. https://doi.org/10.1098/rspb.2021.0703

Bolger AM, Lohse M, Usadel B. (2014). Trimmomatic: A flexible trimmer for Illumina

Sequence Data. *Bioinformatics*, btu170. https://doi.org/10.1093/bioinformatics/btu170

Buczkowski G, Wossler TC. (2019). Controlling invasive Argentine ants, *Linepithema humile*, in

conservation areas using horizontal insecticide transfer. *Scientific Reports* 9: 19495.

https://doi.org/10.1038/s41598-019-56189-1

Chhetri P.K., Pradhan BK, Chhetri B. (2020) First record of *Ophiocordyceps dipterigena* Berk.

& Broome (Ophiocordycipitaceae) in the Himalayas. *National Academy Science Letters*

43, 367-369. https://link.springer.com/article/10.1007/s40009-019-00857-3

Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. (2005) Blast2GO: a universal

tool for annotation, visualization and analysis in functional genomics research.

Bioinformatics 21: 3674-3676. https://doi.org/10.1093/bioinformatics/bti610

Cooley JR, Marshall DC, Hill KBR. 2018. A specialized fungal parasite (Massospora cicadina)

hijacks the sexual signals of periodical cicadas (Hemiptera: Cicadidae: Magicicada).

Scientific Reports 8:1432. https://doi.org/10.1038/s41598-018-19813-0

Cremer S, Armitage SAO, Schmid-Hempel P. (2007). Social Immunity. *Current Biology* 17(16)

R693-R702. https://doi.org/10.1016/j.cub.2007.06.008

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,

McCarthy SA, Davies RM, Li H. (2021). Twelve years of SAMtools and BCFtools.

GigaScience 10(2): giab008. https://doi.org/10.1093/gigascience/giab008

Das B, de Bekker C. (2022). Time-course RNASeq of *Camponotus floridanus* forager and nurse

ant brains indicate links between plasticity in the biological clock and behavioral division

of labor. *BMC Genomics* 23(57). https://doi.org/10.1186/s12864-021-08282-x

Dawkins R. (1982). *The Extended Phenotype: The Long Reach of the Gene*. Oxford, UK: Oxford

University Press.

De Bekker C, Beckerson WC, Carolyn Elya. (2021). Mechanisms Behind the Madness: How

do zombie-making fungal entomopathogens affect host behavior to increase

transmission? *mBio* 12(5), e01872-21.https://doi.org/10.1128/mBio.01872-21

de Bekker C, Ohm RA, Evans HC, Brachmann A, Hughes DP. (2017). Ant-infecting

Ophiocordyceps genomes reveal a high diversity of potential behavioral manipulation

genes and a possible major role for enterotoxins. Sci Rep 7:12508.

https://doi.org/10.1038/s41598-017-12863-w

de Bekker C, Ohm RA, Loreto RG, Sebastian A, Albert I, Merrow M, Brachmann A, Hughes

DP. (2015). Gene expression during zombie ant biting behavior reflects the complexity

underlying fungal parasitic behavioral manipulation. BMC Genomics 16:620.

https://doi.org/10.1186/s12864-015-1812-x

Dewulf M, Köster DV, Sinha B, Viaris de Lesegno C, Chambon V, Bigot. A, Bensalah M,

Negroni E, Tardif N, Podkalicka. J, Johannes L, Nassov P, Butler-Browne G, Lamaze C,

Blouin CM. (2019). Dystrophy-associated caveolin-3 mutations reveal that caveolae

couple IL6/STAT3 signaling with mechanosensing in human muscle cells. *Nature

Communications* 10: 1974. https://doi.org/10.1038/s41467-019-09405-5

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras

TR. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 29(1):15-21.

https://doi.org/10.1093/bioinformatics/bts635

Duran RM, Cary JW, Calvo AM. (2007). Production of cyclopiazonic acid, aflatrem, and

aflatoxin by *Aspergillus flavus* is regulated by veA, a gene necessary for sclerotial

formation. *Applied Microbiology and Biotechnology* 73(5): 1158-68.

https://doi.org/10.1007/s00253-006-0581-5

Eldefrawi ME, Gant DB, Eldefrawi AT. (1990). The GABA receptor and the action of

tremorgenic mycotoxins. Pohland AE (Ed.). Microbial Toxins in Foods and Feeds:

Cellular and molecular modes of action (291-295). New York, NY, Plenum Press.

https://doi.org/10.1007/978-1-4613-0663-4_27

Elya C, Lich HHdF. (2021). The genus *Entomophthora*: bringing the insect destroyers into the

twenty-first century. *IMA Fungus* 12:34. https://doi.org/10.1186/s43008-021-00084-w

Gant DB, Cole RJ, Valdes JJ, Eldefrawi ME, and Eldefrawi AT. (1987). Action of

tremorgenic mycotoxins on GABAA receptor. *Life Sciences* 41: 2207–2214.

https://doi.org/10.1016/0024-3205(87)90517-0

Evans HC, Araújo JPM, Halfeld VR, Hughes DP. Epitypification and re-description of the

zombie-ant fungus, *Ophiocordyces unilateralis* (*Ophiocordycipitaceae*). *Fungal*

Systematics and Evolution 1(1): 13-22. https://doi.org/10.3114/fuse.2018.01.02

Feher J. (2017). *Contractile Mechanisms in Skeletal Muscle*. in: *Quantitative Human Physiology*

pp. 305-217 :Academic Press. *https://doi.org/10.1016/B978-0-12-800883-6.00028-8*

Fredericksen MA, Zhang Y, Hazen ML, Hughes DP. (2017). Three-dimensional visualization

and a deep-learning model reveal complex fungal parasite networks in behaviorally

manipulated ants. *PNAS 114(47)*: 12590-12595. https://doi.org/10.1073/pnas.1711673114

Freiburg A, Trombitas K, Hell W, Cazorla O, Fougerousse F, Center T, Kolmerer B, Witt C,

Beckmann JS, Gregorio CC, Granzier H, Labeit S. (2000). Series of exon-skipping events

in the elastic spring region of titin as the structural basis for myofibrillar elastic diversity.

Circulation Research 86(11): 1114-1121. https://doi.org/10.1161/01.res.86.11.1114

Galbiati F, Lisanti MP. (2000-2013). *Caveolin-3 and Limb-Girdle Muscular Dystrophy*. In:

Madame Curie Bioscience Database [Internet]. Austin (TX): Landes Bioscience; 2000-

2013. Available from: https://www.ncbi.nlm.nih.gov/books/NBK6073/

Gaubert A, Amigues B, Spinelli S, Cambillau C. (2020). Chapter Seven – Strucutre of odorant

binding proteins and chemosensory proteins determined by X-ray crystallography.

Methods in Enzymology 642: 151-167. https://doi.org/10.1016/bs.mie.2020.04.070

Gospocic J, Shields EJ, Glastad KM, Lin Y, Penick CA, Yan H, Mikheyev AS, Linksvayer TA,

Garcia BA, Berger SL, Liebig J, Reinberg D, Bonasio R. (2017). The neuropeptide

corazonin controls social behavior and caste identity in ants. *Cell* 10; 170(4): 748-

759.e12. https://doi.org/10.1016/j.cell.2017.07.014

Graham RJ. (2018) Epigenetic mechanisms governing behavioral reprogramming in the ant

Camponotus floridanus. *Publicly Accessible Penn Dissertations*. 2883.

https://repository.upenn.edu/edissertations/2883

Guo Y, Zhao Z, Li X. (2021). Moderate warming will expand the suitable habitat of

Ophiocordyceps sinensis and expand the area of *O. sinensis* with high adenosine content.

Science of the Total Environment 787: 147605.

https://doi.org/10.1016/j.scitotenv.2021.147605

Han Y, van Houte S, Drees GF, van Oers MM, Ros VI. (2015). Parasitic Manipulation of Host

Behaviour: Baculovirus SeMNPV EGT Facilitates Tree-Top Disease in Spodoptera

exigua Larvae by Extending the Time to Death. *Insects* 31;6(3): 716-31.

https://doi.org/10.3390/insects6030716

Hueffer K, Khatri S, Rideout S, Harris MB, Papke RL, Stokes C, Schulte MK. (2017). Rabies

virus modifies host behavior through a snake-toxin like region of its glycoprotein that

inhibits neurotransmitter receptors in the CNS. *Scientific Reports* 7(1):12818.

https://doi.org/10.1038/s41598-017-12726-4

Hughes DP, Andersen SB, Hywel-jones NL, Himaman W, Billen J, Boomsma JJ. (2011).

Behavioral mechanisms and morphological symptoms of zombie ants dying from fungal

infection. *BMC Ecology* 11(13) 1-10. https://doi.org/10.1186/1472-6785-11-13

Kim D, Ackerman SL. (2011). The UNC5C netrin receptor regulates dorsal mouse hindbrain

axons. *The Journal of Neuroscience* 31(6): 2167-2179.

https://doi.org/10.1523/JNEUROSCI.5254-10.2011

Klotz JH, Moss JI. (1996). Oral toxicity of a boric acid – sucrose water bait to Florida carpenter

ants (Hymenoptera: Formicidae). *Journal of Entomological Science* 31: 9-12.

Kopczynski CC, Alton AK, Fechtel K, Kooh PJ, Muskavitch MA. (1988). Delta, a *Drosophila*

neurogenic gene, is transcriptionally complex and encodes a protein related to blood

coagulation factors and epidermal growth factor of vertebrates. *Genes and Development*

2(12B): 1723-1735. https://doi.org/10.1101/gad.2.12b.1723

Kumar S, Chen D, Jang C, Nall A, Zheng X, Sehgal A. (2014). An ecdysone-responsive nuclear

receptor regulates circadian rhythms in *Drosophila*. *Nature Communications* 5: 5697.

https://doi.org/10.1038/ncomms6697

Lanner JT, Georgiou DK, Joshi AD, Hamilton SL. (2010). Ryanodine Receptors: Structure,

expression, molecular details, and function in calcium release. *Cold Spring Harbor*

Perspectives in Biology 2(11): 1003996. https://doi.org/10.1101/cshperspect.a003996

Latchininsky AV, Temreshev II, Childebaev MK, Kolov SV. (2016). Host range and recorded

distribution of the fungal pathogen *Entomophaga grylli* (Entomophthoromycota:

Entomophthorales) in Kazakhstan. *BioONE Complete* 25(2): 83-89.

https://doi.org/10.1665/034.025.0207

Lee K-S, You K-H, Choo J-K, Han Y-M, Yu K. (2004). *Drosophila* short neuropeptide F

regulates food intake and body size. *Journal of Biological Chemistry* 279(49): 50781-

50789. https://doi.org/10.1074/jbc.M407842200

Liao Y, Smyth GK and Shi W. (2019). The R package Rsubread is easier, faster, cheaper and

better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Research*

47(8):e47. https://doi.org/10.1093/nar/gkz114

Loreto R, Araújo J, Kepler R, Fleming K, Moreau C, Hughes D. (2018). Evidence for convergent

evolution of host parasitic manipulation in response to environmental conditions.

Evolution 72(10): 2144-2155. https://doi.org/10.1111/evo.13489

Love MI, Huber W, Anders S. (2014). Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology* 15:550.

https://doi.org/10.1186/s13059-014-0550-8

Mukunda L, Lavista-Llanos S, Hansson BS, Wicher D. (2014). Dimerisation of the *Drosophila*

odorant coreceptor Orco. *Frontiers in Cellular Neuroscience* 28.

https://doi.org/10.3389/fncel.2014.00261

Nicholson MJ, Koulman A, Monahan BJ, Pritchard BL, Payne GA, Scott B. (2009).

Identification of two aflatrem biosynthesis gene loci in *Aspergillus flavus* and metabolic

engineering of *Penicillium paxillin* to elucidate their function. *Applied and*

Environmental Microbiology 75(23): 7469-7481. https://doi.org/10.1128/AEM.02146-08

Nocek B, Mulligan R, Bargassa M, Collart F, Joachimiak A. (2008). Crystal structure of

aminopeptidase N from human pathogen Neisseria meningitidis. *Proteins* 70(1): 273-279.

https://doi.org/10.1002/prot.21276

Obulesu M. (2019). *Viral Vector Therapeutics Against Alzheimer's Disease*. In: *Alzheimer's*

Disease Theranostics. Elsevier. https://doi.org/10.1016/C2018-0-00026-2

Pastell M. (2016). CowLog – Cross-platform application for coding behaviours from video.

Journal of Open Research Software 4(1): p.e15. https://doi.org/10.5334/jors.113

Pelosi P, Iovinella I, Zhu J, Wang G, Dani F. (2018). Beyond chemoreception: diverse tasks of

soluble olfactory proteins in insects. *Biological Reviews* 93: 184-200.

https://doi.org/10.1111/brv.12339

Petch, Trans. Br. Mycol. Soc. 18(1): 53 (1933). Ophiocordyceps clavulata (Schwein.)

http://www.indexfungorum.org/Names/NamesRecord.asp?RecordID=251478

Pregitzer P, Greschista M, Breer H, Krieger J. (2014). The sensory neurone membrane protein

SNMP1 contributes sensitivity of a pheromone detection system. *Insect Molecular*

Biology 23(6): 733-742. https://doi.org/10.1111/imb.12119

Reichhart N, Schöberl S, Keckeis S, Alfaar AS, Roubeix C, Cordes M, Crespo-Garcia S,

Haeckel A, Kociok N, Föckler R, Fels G, Mataruga A, Rauh R, Milenkovic VM, Zühlke

K, Klussmann E, Schellenberger E, Strauß O. (2019). Anoctamin-4 is a bona fide Ca^{2+}-

dependent non-selective cation channel. *Scienctific Reports* 9: 2257.

https://doi.org/10.1038/s41598-018-37287-y

Sakamoto H, Goka K. (2021). Acute toxicity of typical ant control agents to the red imported fire

ant, *Solenopsis invicta* (Hymenoptera: Formicidae). *Applied Entomology and Zoology* 56:

217-224. https://doi.org/10.1007/s13355-021-00728-8

Sánchez MI, Biron DG. (2019). Host Manipulation by Parasites. *Frontiers in Ecology and*

Evolution 7:369. https://doi.org/10.3389/fevo.2019.00369

Selala MI, Daelemans F, Schepens PJC. (2008). Fungal Tremorgens: The mechanism of action

of single nitrogen containing toxins – a hypothesis. Drug and Chemical Toxicology 12(3-

4): 237-257. https://doi.org/10.3109/01480548908999156

Shakiryanova D, Klose MK, Zhou Y, Gu T, Deitcher DL, Atwood HL, Hewes RS, Levitan ES.

(2007). Presynaptic ryanodine receptor-activated calmodulin kinase II increases vesicle

mobility and potentiates neuropeptide release. *The Journal of Neuroscience* 27(29):7799-

7806. https://doi.org/10.1523/JNEUROSCI.1879-07.2007

Shinoda T, Itoyama K. (2003). Juvenile hormone acid methyltransferase: A key regulatory

enzyme for insect metamorphosis. *PNAS* 100(21): 11986-11991.

https://doi.org/10.1073/pnas.2134232100

Shrestha B, Tanaka E, Hyun MW, Han J-G, Kim CS, Jo JW, Han S-K, Oh J, Sung G-H. (2016).

Coleopteran and Lepidopteran hosts of the Entomopathogenic genus *Cordyceps sensu*

lato. *Journal of Mycology* 2016: 7648219. https://doi.org/10.1155/2016/7648219

Sobczak JF, Arruda IDP, Fonseca EO, Rabelo PJQ, Ageu F, Nóbrega dS, Pires JC, Somavilla A.

(2020). Manipulation of Wasp (Hymenoptera: Vespidae) Behavior by the

entomopathogenic fungus *Ophiocodyceps humbertii* in the Atlantic forest in Ceará,

Brazil. *Entomological News* 129(1): 98-104. https://doi.org/10.3157/021.129.0115

Steinkraus DC, Hajek AE, Liebherr JK. 2017. Zombie soldier beetles: epizootics in the

goldenrod soldier beetle, Chauliognathus 37ensilvanicus (Coleoptera: Cantharidae)

caused by Eryniopsis lampyridarum (Entomophthoromycotina: Entomophthoraceae).

Journal of Invertebrate Pathology 148: 51–59. https://doi.org/10.1016/j.jip.2017.05.002

Sung G-H, Hywel-Jones NL, Sung J-M, Luangsa-ard JJ, Shrestha B, Spatafora JW. (2007).

Phylogenetic classification of *Cordyceps* and the clavicipitaceous fungi. *Studies in*

Mycology 57: 5-59. https://doi.org/10.3114/sim.2007.57.01

Tarailo-Graovac M, Drögemöller BI, Wasserman WW, Ross CJD, van den Ouweland AMW,

Darin N, Kollberg G, van Kernebeek CDM, Blomqvist M. (2017). Identification of a

large intronic transposal insertion in *SLC17A5* causing sialic acid storage disease.

Orphanet Journal of Rare Diseases 12: 28. https://doi.org/10.1186/s13023-017-0584-6

TePaske MR, Gloer JB, Wicklow DT, Dowd PF. (1991). Aflavarin and β-aflatrem: new anti-

insectan metabolites from the sclerotia of *Aspergillus flavus*. *Journal of Natural Products*

55(8): 1080-1086. https://doi.org/10.1021/np50086a008

Tomita H, Cornejo F, Aranda-Pino B, Woodard CL, Rioseco CC, Neel BG, Alvarez AR, Kaplan

DR, Miller FD, Cancino G. (2020). The protein tyrosin phosphatase receptor delta

regulates developmental neurogenesis. *Cell Reports* 30(1): 215-228.

https://doi.org/10.1016/j.celrep.2019.11.033

Tong EH, Pavey C, O'Handley R, Vyas A. (2021). Behavioral biology of *Toxoplasma gondii*

infection. *Parasites and Vectors* 14(1):77. https://doi.org/10.1186/s13071-020-04528-x

Trienens M, Rohlfs M. (2011). Experimental evolution of defense against a competitive mold

confers reduced sensitivity to fungal toxins but no increased resistance in Drosophila

larvae. *BMC Evolution*ary Biology 11: 206. https://doi.org/10.1186/1471-2148-11-206

Trinh T, Ouellette R, de Bekker C. (2021). Getting lost: the fungal hijacking of ant foraging

behavior in space and time. *Animal Behaviour* 181:165-184.

https://doi.or/10.1016/j.anbehav.2021.09.003

Valdes JJ, Cameron JE, Cole RJ. (1985). Aflatrem: A tremorgenic mycotoxin with acute

neurotoxic effects. Environmental Health Perspectives 62: 459-463.

https://doi.org/10.1289/ehp.8562459

Wells L, Edwards KA, Bernstein SI. (1996). Myosin heavy chain isoforms regulate muscle

function but not myofibril assembly. *The EMBO Journal* 15(17): 4454-4459.

https://doi.org/10.1002/j.1460-2075.1996.tb00822.x

Werkhoven Z, Rohrsen C, Qin C, Brembs B, de Bivort B. (2019). MARGO (Massively

Automated Real-time GUI for object-tracking), a platform for high-throughput ethology.

PLoS one 14(11): e0224243. https://doi.org/10.1371/journal.pone.0224243

Wesołowska W, Wesołowski T. (2013). Do *Leucochloridium* sporocysts manipulate the

behaviour of their snail hosts? *Journal of Zoology* 292(3): 151-155.

https://doi.org/10.1111/jzo.12094

Will I, Das B, Trinh T, Brachmann A, Ohm RA, de Bekker C. 2020. Genetic underpinnings of

host manipulation by *Ophiocordyceps* as revealed by comparative transcriptomics. *G3*

10(7): 2275-2296. https://doi.org/10.1534/g3.120.401290

Xing X, Yan M, Pang H, Wu F, Wang J, Sheng S. (2021). Cytochrome P450s Are Essential for

Insecticide Tolerance in the Endoparasitoid Wasp Meteorus pulchricornis (Hymenoptera:

Braconidae). *Insects* 12(7):651. https://doi.org/10.3390/insects12070651

Yao Y, Peter AB, Baue R, Sigel E. (1989). The tremorigen aflatrem is a positive allosteric

modulator of the gamma-aminobutyric acidA receptor channel expressed in Xenopus

oocytes. *Molecular Pharmacology* 35(3): 319-323.

https://molpharm.aspetjournals.org/content/35/3/319.long

Zalewska M, Kochman A, Estève JP, Lopez F, Chaoui K, Susini C, Ozyhar A, Kochman M.

(2009). Juvenile hormone binding protein traffic - Interaction with ATP synthase and

lipid transfer proteins. *Ciochimica et Biophysica Acta* 1788(9):1695-705.

https://doi.org/10.1016/j.bbamem.2009.04.022

Zhang Y, Featherstone D, Davis W, Rushton E, Broadie K. (2000). *Drosophila* D-Titin is required

for myoblast fusion and skeletal muscle striation. *Journal of Cell Science* 113: 3103-3115.

https://doi.org/10.1242/jcs.113.17.3103

Supplemental Files

Supplemental Table 1: The average length and weight of *Camponotus floridanus* in this study

Colony #	Treatment Group	Average Length (mm)	Average Weight (mg)
Colony 1	C1-PBS	7.07	7.56
	C1-PBS/10% acetone	6.50	6.93
	C1-PBS/10% acetone/500ng aflatrem	6.78	7.94
	C1-PBS/10% acetone/100ng aflatrem	7.00	8.14
	C1-PBS/10% acetone/50ng aflatrem	6.43	8.44
	C1-PBS/10% acetone/10ng aflatrem	7.29	9.01
Colony 2	C2-PBS	7.07	9.49
	C2-PBS/10% acetone	7.07	10.27
	C2-PBS/10% acetone/500ng aflatrem	6.93	11.09
	C2-PBS/10% acetone/100ng aflatrem	6.93	10.48
	C2-PBS/10% acetone/50ng aflatrem	6.86	10.86
	C2-PBS/10% acetone/10ng aflatrem	6.86	10.13
Colony 3	C3-PBS	6.07	5.40
	C3-PBS/10% acetone	6.93	7.93
	C3-PBS/10% acetone/500ng aflatrem	6.79	6.44
	C3-PBS/10% acetone/100ng aflatrem	5.71	5.40
	C3-PBS/10% acetone/50ng aflatrem	5.64	6.23
	C3-PBS/10% acetone/10ng aflatrem	6.29	7.24
	Total	6.68 mm	8.28 mg

 Supplemental Figure 1

Supplemental Figure 1. An overview of the randomized block format used to assign videos to each

observer (A). The videos assigned to observer 1 are shown in green, the videos assigned to observer

2 are shown in blue, and the videos assigned to both for formative analysis are shown in grey. B)

depicts the duration of staggering scores for each grey assignment, with the scores for observer 1

shown as green squares and the scores for observer 2 shown as blue circles.

Supplemental Figure 2

Supplemental Figure 2. Bar graphs depict the number of ants exhibiting staggering behavior

(black) and the number of ants resting (dark grey) across the 30-minute observational period,

broken down into 5 second intervals.

Supplemental Figure 3

Supplemental Figure 3. The path traveled by each ant recorded by the MARGO tracking software

during the 30-minute observational period is shown in dark blue. The three columns on the left

shows the paths traveled by the control group ants injected with PBS/10% acetone, while the three

columns on the right shows the paths traveled by the ants in the PBS/10% acetone/500 ng aflatrem

treatment group. Each group is further divided by colony ID using purple, blue, and dark blue

outline to indicate individuals from colonies 1, 2, and 3, respectively.

Supplemental File 1

RNAseq Data

Supplemental Video 1

O. camponoti-floridani – infected *C. floridanus* staggers syndrome - mp4

Supplemental Video 2

*Observed behaviors – mp4

Supplemental Video 3

*Stagger behaviors – mp4

Supplemental Video 4

*Unique behaviors – mp4

A) Microinjections (0.5 μL)

Per treatment

x7

x7

x7

Aflatrem

Treatments
Phosphate buffered saline (PBS)
PBS/10% acetone
PBS/10% acetone/500 ng aflatrem
PBS/10% acetone/100 ng aflatrem
PBS/10% acetone/50 ng aflatrem
PBS/10% acetone/10 ng aflatrem

B)

30 min

15 mm

60 mm

30 min

60 fps
Video

C)

Output
Average speed
Distance traveled

Behaviors
Resting
Climbing wall
Climbing top
Climbing rope
Walking
Staggering
Grooming
Biting