1   **Short Communication**
2
3   **Cyanotoxins accumulate in Lake St. Clair fish yet their fillets are safe to eat**
4

5   René S. Shahmohamadloo*[a,b], Satyendra P. Bhavsar[c,d], Xavier Ortiz Almirall[c,e], Stephen A. C.
6   Marklevitz[f], Seth M. Rudman[a], and Paul K. Sibley[b]
7
[a] School of Biological Sciences, Washington State University, 14204 NE Salmon Creek Ave,
Vancouver, WA, 98686, United States
[b] School of Environmental Sciences, University of Guelph, 50 Stone Rd E, Guelph, ON, N1G 2W1,
Canada
[c] Ministry of the Environment, Conservation and Parks, 125 Resources Rd, Toronto, ON, M9P 3V6,
Canada
[d] Department of Physical & Environmental Sciences, University of Toronto, 1065 Military Trail,
Toronto, ON, M1C 1A4, Canada
[e] IQS School of Engineering, Universitat Ramon Llull, Via Augusta, 390, 08017, Barcelona
[f] Lake Erie Management Unit, Ministry of Natural Resources and Forestry, 320 Milo Road,
Wheatley, ON, N0P 2P0, Canada
19
20   *Address correspondence to:
21   René S. Shahmohamadloo, PhD
22   School of Biological Sciences | Washington State University
23   14204 NE Salmon Creek Ave | Vancouver | Washington | 98686 | United States
24   Email: rene.shahmohamadloo@wsu.edu
25

26   **Keywords:** *Fish, Harmful Algal Bloom, Human Health, Great Lakes, Food Safety Hazard*
27

Main Text

Abstract

Consuming fish exposed to cyanobacterial harmful algal blooms (HABs) may be a major route of
microcystin toxin exposure to humans. However, it remains unknown whether fish can accumulate
and retain microcystins temporally in waterbodies with recurring seasonal HABs, particularly before
and after a HAB event when fishing is active. We conducted a field study on Largemouth Bass,
Northern Pike, Smallmouth Bass, Rock Bass, Walleye, White Bass, and Yellow Perch to assess
the human health risks to microcystin toxicity via fish consumption. We collected 124 fish in 2016
and 2018 from Lake St. Clair, a large freshwater ecosystem in the North American Great Lakes
that is actively fished pre- and post-HAB periods. Muscles were analyzed using the MMPB
Lemieux Oxidation method for total microcystins, which was used to perform a human health risk
assessment for comparison against fish consumption advisory benchmarks available for Lake St.
Clair. From this collection 35 fish livers were additionally extracted to confirm the presence of
microcystins. Microcystins were detected in all livers at widely varying concentrations (1-1,500 ng
g^{-1} ww), suggesting HABs are an underappreciated and pervasive stressor to fish populations.
Conversely, microcystin levels were consistently low in muscles (0-15 ng g^{-1} ww) and presented
negligible risk, empirically supporting that fillets may be safely consumed before and after HAB
events following fish consumption advisories.

1. Introduction

The global expansion of cyanobacterial harmful algal blooms (HABs) is a pressing challenge for
freshwater ecosystems. Their spread is due, in part, to climatic changes and an expansion in
human activities, particularly intensified agriculture, which render waterbodies increasingly
vulnerable to biodiversity loss and threatened ecosystem services (Hellweger et al., 2022; Ho et
al., 2019; Huisman et al., 2018; Reid et al., 2019; Smith et al., 2019). Freshwater HABs can
produce microcystins, potent hepatotoxins that primarily inhibit protein phosphatases and at
sufficient levels can cause liver necrosis, tumor formation, and organ failure in humans and wildlife
(Chorus and Welker, 2021; Shahmohamadloo et al., 2023b, 2022, 2021).

The World Health Organization's (WHO) guidelines (WHO, 2020) are implemented in many
countries to protect humans from microcystin toxicity in drinking water ($12\,\mu g\,L^{-1}$ short-term, $1\,\mu g\,L^{-1}$
lifetime); however, longstanding concerns remain that fish consumption may be a viable route of
algal toxin exposure (FAO, 2020; IPCC, 2022). These concerns stem from the possibility that
microcystins may be one of several factors involved in fish kills during HAB events (Ibelings et al.,
2005) or that microcystins can accumulate up the food chain (Gene et al., 2019; Shahmohamadloo
et al., 2020a, 2020b). Previous studies measured microcystins in fish tissues at levels exceeding
the WHO's tolerable daily intake (TDI) value (0.04 μg kg^{-1} body weight per day) using the enzyme
linked immunosorbent assay (ELISA) (Dyble et al., 2011; Poste et al., 2011; Wituszynski et al.,
2017). Yet, ELISA is prone to false positives in fish tissues because it only measures free fractions
of microcystins, whereas bound fractions can make up to 90% of the total amount of toxins present
in a sample (Birbeck et al., 2019; Flores et al., 2018; Schmidt et al., 2014). This has raised
skepticism about the accuracy of such data in assessing risks to humans via consumption of fish.
Intertwined in the debate about methodology are projections stating that climate change will
increase the mean number of days of a HAB event from 16-23 days in 2050 to 18-39 days in 2090
in North America (Chapra et al., 2017).

Recent work in Lake Erie addresses some of the methodological and ecological concerns by using
an optimized MMPB Lemieux Oxidation method (Anaraki et al., 2020) to dually assess the human
health risks from fish consumption and fish health risks from microcystin toxin exposure before,
during, and after HAB events (Shahmohamadloo et al., 2023a). Interestingly, the findings suggest
both cause for relief and concern: while Lake Erie fish fillets may be safe to eat, fish livers can
retain high levels of microcystins (mean 460.13 ng g^{-1}ww across species) and may be a pervasive
threat to fish populations (Shahmohamadloo et al., 2023a). Lake St. Clair, a large waterbody that is
important for recreational fishers (OMNRF, 2016), regularly experiences microcystin-HABs (Davis
et al., 2014) and may expose resident fish populations in the future to blooms with extended
residency, severity, and toxicity. Aware that the present duration of a HAB (Chapra et al., 2017) is
a small fraction of the total fish season and the overwhelming body of literature on fish was
collected during HABs (Dyble et al., 2011; Poste et al., 2011; Wituszynski et al., 2017), there is an
invested interest to determine the human health risks from consuming fish before and after HAB
events when recreational fishing is more active.

91
We collected 124 recreationally important sport fish, including some species which have received
little attention previously, namely Largemouth Bass (*Micropterus salmoides*), Northern Pike (*Esox*
lucius), Smallmouth Bass (*Micropterus dolomieu*), Rock Bass (*Ambloplites rupestris*), Walleye
(*Sander vitreus*), White Bass (*Morone chrysops*), and Yellow Perch (*Perca flavescens*) from Lake
St. Clair in 2016 and 2018 throughout pre- and post-HAB periods when recreational fishing is more
active. We measured microcystins in fish muscles using the MMPB Lemieux Oxidation method
(Anaraki et al., 2020). In addition, we dissected 35 livers from these samples to confirm the
presence of microcystins in fish tissues and assess the amount retained pre- and post-HAB
periods. Results from muscles were then compared against fish consumption advisory benchmarks
for microcystins available to Lake St. Clair and applied in a human health risk assessment to
determine whether fish tissues exceeded the WHO's TDI. Here, we hypothesize that microcystin
levels in fish muscles will vary among fish species but, overall, will pose low human health risks in
pre- and post-HAB periods.

105
106 **2. Methods**

107
108 **2.1. Study location and water sampling**

109
Eight locations in Lake St. Clair (42° 25' 20" N, 82° 39' 36" W) were sampled for fish in 2016 and
2018 during pre- (June to July) and post- (October to November) HAB periods. During this time,
Microcystis, *Planktothrix*, and *Dolichospermum* are typically present in Lake St. Clair, with toxigenic
strains of *Microcystis* predominating (Davis et al., 2014). Physical, chemical, and biological water
quality monitoring HABs data was collected by the Ontario Ministry of Natural Resources and
Forestry (MNRF; see Dataset S1). Water data was collected during repeated bi-weekly sampling
trips 'pre', 'during', and 'post' HAB periods to characterize algae blooms and potential microcystin
toxins. This corroborates recent findings demonstrating chlorophyll *a* is a consistent correlate of
microcystins in freshwaters and can better account for the inevitable temporal variation associated
with changing causes of microcystin production (Qian et al., 2021). Notwithstanding, the Ontario
Ministry of the Environment, Conservation and Parks (MECP) additionally collected water samples
during pre- and post-HAB periods from the southern shoreline of Lake St Clair near Belle River
(42° 18' 11" N, 82° 42' 46.7" W) and Puce River (42° 18' 4" N, 82° 42' 19" W) to measure total
phosphorus, chlorophyll *a*, and total microcystins. Total phosphorus (28.7 and 13.4 µg L^{-1}),
chlorophyll *a* (9.7 and 4.6 µg L^{-1}) and total microcystins (0.2 and 1.0 µg L^{-1}) were highest in 2018.
Microcystin concentrations in the present study are similar to previous measurements from Lake
St. Clair (Davis et al., 2014; Taranu et al., 2019).

127
128 **2.2. Fish sampling**
129
A total of 124 samples from Largemouth Bass, Northern Pike, Smallmouth Bass, Rock Bass,
Walleye, White Bass, and Yellow Perch (Table 1, Dataset S1) were collected in 2016 and 2018 for
microcystin screening as a part of the MECP's Fish Contaminant Monitoring Program (Etobicoke,
Ontario, Canada) in partnership with the MNRF (Wheatley, Ontario, Canada). Fish length and
weight were recorded for all samples, after which the dorso-lateral muscle tissues (i.e., skinless,
boneless fillets) were dissected for microcystin analysis. In 2018, a subset of 35 samples of
Walleye, White Bass, and Yellow Perch were additionally dissected for livers to confirm their
exposure to microcystins. All muscles and livers were frozen at -80 °C until homogenization and
microcystin analysis.
139
The total length of fish was measured because most resource users (i.e., fishers) and fish
consumers are unable to determine the age of the fish they consume. For this reason, fish
consumption advisories provided to the public, as well as health and resource managers are given
based on fish length (MECP, 2017). By sampling and presenting the results based on length, we
have provided real-world recommendations using metrics that the general population and decision-
makers can access to help evaluate real-world risks and aid real-world decisions.
146
147 **2.3. Fish tissue analysis**
148
Fish tissues were analyzed for total microcystins (free and bound) using an optimized MMPB
Lemieux Oxidation method that provides the necessary sensitivity and specificity (Anaraki et al.,
2020)). Briefly, frozen tissues were freeze-dried using a Labconco Freezone 2.5 freeze-drier from
Fisher Scientific (Markham, Ontario, Canada) for at least 24 h and then ground to a fine powder
using a pestle and mortar. Next, approximately 100 mg of freeze-dried sample was spiked with the
internal standard (d_3-MMPB), oxidized at pH 8.5 using potassium permanganate (0.3 mM) and
sodium periodate (20 mM) for 2 h. The oxidizing sample was quenched by adjusting the pH to 3
with 10% sulfuric acid and sodium bisulphite. Samples were then centrifuged for 7 min at 5,000 × g
and correspondingly loaded onto Oasis HLB 3 cc (400 mg) LP extraction cartridges (Mississauga,
Ontario, Canada) for clean-up and extraction of MMPB and d_3-MMPB with 0.1% acetic acid and
50% methanol washes. Analytes were eluted with 4 mL of 100% methanol, dried down to 250 μL,
and diluted to 1 mL using 0.1% FA milli-Q water. Samples were finally filtered using Pall GHP
filters (Mississauga, Ontario, Canada) and prepared for instrumental analysis.
162
Samples were quantified using an *in-situ* generated MMPB matrix-matched calibration curve by
isotope dilution with d_3-MMPB by liquid chromatography coupled to time-of-flight mass
spectrometry (LC-QTOF MS) using an Acquity I Class Chromatograph by Waters Corporation
(Milford, Massachusetts, United States). This approach estimates potential matrix effects that may
have impacted the derivatization, sample preparation and instrumental analysis steps. This method
showed 16.7% precision (RSD) and +6.7% accuracy (bias), with a calculated method detection
limit (MDL) of 2.18 ng g^{-1} wet weight (ww).
170
171 **2.4. Human health risk assessment**
172
Total microcystins measured in muscles were compared against Ontario's fish consumption
advisory benchmarks for microcystin-LR developed by the MECP and Ohio's fish consumption
advisory benchmarks for total microcystins developed by the Ohio Environmental Protection
Agency. These benchmarks are based on the World Health Organization's recommended tolerable

daily intake (TDI) of 0.04 μg kg^{-1} bodyweight per day (WHO, 2020). The TDI value is based on the
no-observed-adverse-effect-level (NOAEL) liver pathology observed in a 13-week study on mice
using microcystin-LR (Fawell et al., 1994) and applying an uncertainty factor (UF) of 1000 (Eq. 1).
180

181 $$(1) \quad \text{TDI} = \frac{\text{NOAEL}}{\text{UF}} = \frac{40 \ \mu\text{g kg}^{-1}\text{d}^{-1}}{10 \ \times \ 10 \times 10} = 0.04 \ \mu\text{g kg}^{-1}\text{d}^{-1}$$

183 Ontario's advisory benchmark assumes that 50% of microcystin exposure is through fish
184 consumption. Microcystin-LR is one of the most toxic microcystin congeners (Shimizu et al., 2013).
185 Therefore, application of a microcystin-LR advisory benchmark to total microcystins presents a
186 more conservative and health protective scenario.
187

188 We further calculated the estimated daily intake (EDI) for total microcystins if one meal of fish is
189 consumed every day (Eq. 2) and compared this to the WHO's TDI for microcystin-LR.
190

$$(2) \quad \text{EDI} = \frac{\text{C}_{\text{microcystins}} \times \text{D}_{\text{intake}}}{\text{bw}}$$

Where $\text{C}_{\text{microcystins}}$ is the mean total concentration of microcystins in fish tissue wet weight (Dataset
S1), D_{intake} is the daily fish consumption for Ontarians (one meal is 227 g) following the Ontario's
Guide to Eating Ontario Fish (MECP, 2017), and bw is the body weight of an average healthy adult
(assumed 70 kg). We then calculated the margin of exposure (MOE) for microcystins (Eq. 3),
which is commonly used in human health risk assessment to assess for potentially genotoxic or
carcinogenic compounds.
199

$$(3) \quad \text{MOE} = \frac{\text{NOAEL}}{\text{EDI}}$$

Since the WHO has shown evidence that microcystins are potentially carcinogenic to humans and
animals, an MOE ≥ 10,000 was considered for this study to be protective for genotoxic and
carcinogenic compounds. The value of 10,000 is derived by the multiplication of two UFs; a 100-
fold difference between the calculated reference point and human exposure is applied for species
differences and human variability, and an additional 100-fold difference is applied for inter-
individual variability (EFSA, 2005). Although it is widely documented that MOEs of 10,000 are used
as a protective value in the evaluation of genotoxic and carcinogenic compounds present in food
where a single toxicant is being evaluated (EFSA, 2005), the extent to which microcystins is a risk
remains unclear for MOEs ≤10,000. We therefore adopted the following categories to assist with
MOE classification: 1–1,000 (high risk), 1,000–10,000 (medium risk), 10,000–100,000 (low risk)
(Cunningham et al., 2011).
213

214 We finally calculated the hazard quotient (HQ) for microcystins (Eq. 4) as the ratio of the potential
215 exposure to microcystins and the level at which no adverse effects are expected,
216

217 $$(4) \quad \text{HQ} = \frac{\text{EDI}}{\text{TDI}}$$

219 For adverse effects from microcystin exposure, HQ ≤ 0.1 is considered negligible, HQ ≥ 0.1 ≤ 1 is
220 considered low, HQ ≥ 1 ≤ 10 indicates some hazard, and HQ ≥ 10 indicates a high hazard.
221

222 **2.5. Statistical analysis**

223
224   For statistical analysis any values below the limit of detection of the LC-MS of 2.18 ng g^{-1} ww were
225   treated as zero concentrations. Linear mixed effect models with species treated as a random effect
226   were used to test whether concentrations of microcystins in the muscle and liver were greater than
227   the limit of detection.
228
229   **3. Results and Discussion**
230
Fish muscles contained microcystins ($F_{1,117}$=20.42, p <0.0001), with a mean concentration of 1.29
ng g^{-1} ww across species (Figure 1). Only 24/124 samples had concentrations above the limit of
detection. 2/124 of samples had concentrations exceeding the WHO's TDI and the human health
risk assessment suggests microcystins are a low risk pre- and post-HAB periods (Table 1).
Applying these results against the fish consumption advisory benchmarks in Ontario
(Shahmohamadloo *et al.*, 2022b), 95% of muscles were <6 ng g^{-1} ww microcystins, which would be
classified in the 'unrestricted' fish consumption advisory, and all muscles are in the 'unrestricted'
meal category (<25 ng g^{-1} ww) in Ohio. Considering 98% of muscles had microcystins below the
WHO's TDI, our results indicate that humans can safely consume Lake St. Clair Largemouth Bass,
Northern Pike, Smallmouth Bass, Rock Bass, Walleye, White Bass, and Yellow Perch fillets
ranging from three times a week to every day (12 to 32 meals per month), depending on the
species and its size. A caveat which concerns the validity of our risk assessment result is that
these fish were collected pre- and post-HAB periods. A comprehensive review on wild, freshwater
fish found variable microcystin concentrations in muscles ranging from 0 to 3.27 µg g^{-1} dw during
HABs, yet it was noted that this variability may be due to major limitations in methodology such as
the use of ELISA (Flores et al., 2018). More recent work using the Lemieux Oxidation method on
Nile Tilapia (*Oreochromis niloticus*) collected from fish ponds before, during, and after HABs found
total microcystins (free and bound) could surpass the WHO's EDI, particularly during the warm
months when cyanobacterial blooms are most active (Mohamed et al., 2020). Nile Tilapia are
planktivorous fish, and several studies have shown elevated microcystin concentrations in bottom-
feeders (Adamovský et al., 2007; Flores et al., 2018; Mohamed et al., 2020; Xie et al., 2004; Zhang
et al., 2009). Therefore, caution should be exercised when catching fish before, during, and after
HAB events and can depend on species type.
254
Microcystin concentrations in livers were greater than in muscles with 34/35 samples exceeding
the limits of detection. Fish livers contained a significant amount of microcystins ($F_{1,32}$=21.33, p
<0.0001) with an average value of 224.81 ng g^{-1} ww across Walleye, White Bass, and Yellow
Perch (Figure 2; Table 1). The wide variability and presence of microcystin levels in all livers (1-
1,500 ng g^{-1} ww) supports our previous work in Lake Erie demonstrating that microcystins are
present in fish livers well beyond the cessation of HABs and may be a persistent contaminant in
aquatic ecosystems (Shahmohamadloo et al., 2023a). Microcystin-producing HAB species such as
Microcystis, cyanobacteria common to Lake St. Clair (Davis et al., 2014) and around the globe
(Harke et al., 2016), also possess genetic and ecophysiological diversity that allows them to
effectively overwinter on the sediments and rapidly form blooms under favorable conditions (Dick
et al., 2021). Putative evolutionary adaptations in species such as *Microcystis* (e.g., key genes
involved in biosynthesis of secondary metabolites, carbon concentrating mechanisms,
detoxification of reactive oxygen species, and acquisition of nitrogen and phosphorus (Dick et al.,
2021)) can help to explain the microcystin burden evident in fish livers pre-HAB and offers further
insight into the decades-long postulation that microcystins are one among several stress factors
involved in fish kills during HAB events (Ibelings et al., 2005).
271

To alleviate concerns of microcystin exposure via human consumption, general and sensitive
populations (women of child-bearing age and children) should avoid eating fish organs from Lake
St. Clair and other waterbodies that experience regular HAB events. Our advice is consistent with
current advisories for many other contaminants which could be elevated in fish organs. While it is
convention to gut fish and strictly consume fillets, whole fish consumption is practiced in various
cultures and encouraged for those seeking additional nutrients in their diet (Beveridge et al., 2013;
He, 2009). However, microcystins do not break down during cooking and boiling (Harada et al.,
1996) and may therefore pose some level of risk to consumers eating fish whole. Considering 90%
of anglers consume 8 or less fish meals per month (Awad, 2006) and following the above
recommended practices, our results suggest the risk of microcystin toxicity from fish consumption
may be low for Lake St. Clair before and after HAB events when fishing is active. Our findings
corroborate previous work suggesting microcystin elimination rates from —or degradation rates
in— fish tissues can range from days to weeks, depending on a variety of factors including fish
species, fish tissue type, and environmental signals (Adamovský et al., 2007; Gurbuz et al., 2016;
Shahmohamadloo et al., 2022).
287
Understanding the potential risks of consuming fish exposed to microcystins, and cyanotoxins
more broadly, is critical towards interpreting human and ecological risks. A One Health approach to
safeguarding human and animal health offers an interdisciplinary and collaborative path forward in
monitoring illnesses or deaths associated with HABs (Hilborn and Beasley, 2015). We therefore
exercise prudence in our recommendations and call attention to the need for further work to
investigate the chronic and sublethal impacts of HABs and associated toxins on fish populations.
294

Acknowledgements

This work was supported by an NSERC CREATE Grant (2013-432269), a Banting Postdoctoral
Fellowship, a Canada-Ontario Agreement (2218) through the Ontario Ministry of the Environment,
Conservation and Parks, and a Government of Ontario Grant (GLS 1403). Financial support by the
Government of Ontario does not equal endorsement of this paper. We thank Ngan Diep for
providing microcystin water measurements and the field technicians from the Ontario Ministry of
Natural Resources and Forestry for collecting fish and water samples from Lake St. Clair for this
study.

References

Adamovský, O., Kopp, R., Hilscherová, K., Babica, P., Palíková, M., Pasková, V., Navrátil, S.,
Marsálek, B., Bláha, L., 2007. Microcystin kinetics (bioaccumulation and elimination) and
biochemical responses in common carp (Cyprinus carpio) and silver carp (Hypophthalmichthys
molitrix) exposed to toxic cyanobacterial blooms. Environ. Toxicol. Chem. 26, 2687–2693.

Anaraki, M.T., Shahmohamadloo, R.S., Sibley, P.K., MacPherson, K., Bhavsar, S.P., Simpson,
A.J., Ortiz Almirall, X., 2020. Optimization of an MMPB Lemieux Oxidation method for the
quantitative analysis of microcystins in fish tissue by LC-QTOF MS. Sci. Total Environ. 737,
140209.

Beveridge, M.C.M., Thilsted, S.H., Phillips, M.J., Metian, M., Troell, M., Hall, S.J., 2013. Meeting
the food and nutrition needs of the poor: the role of fish and the opportunities and challenges
emerging from the rise of aquaculture. J. Fish Biol. 83, 1067–1084.
320

321 Birbeck, J.A., Westrick, J.A., O'Neill, G.M., Spies, B., Szlag, D.C., 2019. Comparative Analysis of
322   Microcystin Prevalence in Michigan Lakes by Online Concentration LC/MS/MS and ELISA.
323   Toxins 11.
324
325 Chapra, S.C., Boehlert, B., Fant, C., Bierman, V.J., Jr, Henderson, J., Mills, D., Mas, D.M.L.,
326   Rennels, L., Jantarasami, L., Martinich, J., Strzepek, K.M., Paerl, H.W., 2017. Climate Change
327   Impacts on Harmful Algal Blooms in U.S. Freshwaters: A Screening-Level Assessment.
328   Environ. Sci. Technol. 51, 8933–8943.
329
330 Chorus, I., Welker, M., 2021. Toxic Cyanobacteria in Water:A Guide to Their Public Health
331   Consequences, Monitoring and Management, 2nd Edition. ed. CRC Press.
332
333 Cunningham, F.H., Fiebelkorn, S., Johnson, M., Meredith, C., 2011. A novel application of the
334   Margin of Exposure approach: segregation of tobacco smoke toxicants. Food Chem. Toxicol.
335   49, 2921–2933.
336
337 Davis, T.W., Watson, S.B., Rozmarynowycz, M.J., Ciborowski, J.J.H., McKay, R.M., Bullerjahn,
338   G.S., 2014. Phylogenies of microcystin-producing cyanobacteria in the lower Laurentian Great
339   Lakes suggest extensive genetic connectivity. PLoS One 9, e106093.
340
341 Dick, G.J., Duhaime, M.B., Evans, J.T., Errera, R.M., Godwin, C.M., Kharbush, J.J., Nitschky, H.S.,
342   Powers, M.A., Vanderploeg, H.A., Schmidt, K.C., Smith, D.J., Yancey, C.E., Zwiers, C.C.,
343   Denef, V.J., 2021. The genetic and ecophysiological diversity of Microcystis. Environ.
344   Microbiol. 23, 7278–7313.
345
346 Dyble, J., Gossiaux, D., Landrum, P., Kashian, D.R., Pothoven, S., 2011. A kinetic study of
347   accumulation and elimination of microcystin-LR in yellow perch (Perca flavescens) tissue and
348   implications for human fish consumption. Mar. Drugs 9, 2553–2571.
349
350 European Food Safety Authority (EFSA), 2005. Opinion of the Scientific Committee on a request
351   from EFSA related to A Harmonised Approach for Risk Assessment of Substances Which are
352   both Genotoxic and Carcinogenic. EFSA 3, 282.
353
354 FAO, 2020. Climate change: Unpacking the burden on food safety. Food & Agriculture Org.
355
356 Fawell, J.K., James, C.P., James, H.A., 1994. Toxins from blue-green algae: toxicological
357   assessment of microcystin-LR and a method for its determination in water. Foundation for
358   water research.
359
360 Flores, N.M., Miller, T.R., Stockwell, J.D., 2018. A Global Analysis of the Relationship between
361   Concentrations of Microcystins in Water and Fish. Frontiers in Marine Science 5.
362
363 Gene, S.M., Shahmohamadloo, R.S., Ortiz, X., Prosser, R.S., 2019. Effect of Microcystis
364   aeruginosa-Associated Microcystin-LR on the Survival of 2 Life Stages of Freshwater Mussel
365   (Lampsilis siliquoidea). Environ. Toxicol. Chem. 38, 2137–2144.
366
367 Gurbuz, F., Uzunmehmetoğlu, O.Y., Diler, Ö., Metcalf, J.S., Codd, G.A., 2016. Occurrence of
368   microcystins in water, bloom, sediment and fish from a public water supply. Sci. Total Environ.
369   562, 860–868.
370

371 Harada, K.-I., Tsuji, K., Watanabe, M.F., Kondo, F., 1996. Stability of microcystins from
372   cyanobacteria—III. Effect of pH and temperature. Phycologia 35, 83–88.
373

374 Harke, M.J., Steffen, M.M., Gobler, C.J., Otten, T.G., Wilhelm, S.W., Wood, S.A., Paerl, H.W.,
375   2016. A review of the global ecology, genomics, and biogeography of the toxic
376   cyanobacterium, Microcystis spp. Harmful Algae 54, 4–20.
377

378 He, K., 2009. Fish, long-chain omega-3 polyunsaturated fatty acids and prevention of
379   cardiovascular disease--eat fish or take fish oil supplement? Prog. Cardiovasc. Dis. 52, 95–
380   114.
381

382 Hellweger, F.L., Martin, R.M., Eigemann, F., Smith, D.J., Dick, G.J., Wilhelm, S.W., 2022. Models
383   predict planned phosphorus load reduction will make Lake Erie more toxic. Science 376,
384   1001–1005.
385

386 Hilborn, E.D., Beasley, V.R., 2015. One health and cyanobacteria in freshwater systems: animal
387   illnesses and deaths are sentinel events for human health risks. Toxins 7, 1374–1395.
388 Ho, J.C., Michalak, A.M., Pahlevan, N., 2019. Widespread global increase in intense lake
389   phytoplankton blooms since the 1980s. Nature 574, 667–670.
390

391 Huisman, J., Codd, G.A., Paerl, H.W., Ibelings, B.W., Verspagen, J.M.H., Visser, P.M., 2018.
392   Cyanobacterial blooms. Nat. Rev. Microbiol. 16, 471–483.
393

394 Ibelings, B.W., Bruning, K., de Jonge, J., Wolfstein, K., Pires, L.M.D., Postma, J., Burger, T., 2005.
395   Distribution of microcystins in a lake foodweb: no evidence for biomagnification. Microb. Ecol.
396   49, 487–500.
397

398 IPCC, 2022. Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working
399   Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change
400   [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M.
401   Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University
402   Press. In Press.
403

404 Ministry of the Environment, Conservation and Parks (MECP), 2017. Guide to Eating Ontario Fish.
405   Publications Ontario.
406

407 Mohamed, Z., Ahmed, Z., Bakr, A., Hashem, M., Alamri, S., 2020. Detection of free and bound
408   microcystins in tilapia fish from Egyptian fishpond farms and its related public health risk
409   assessment. J. Consum. Prot. Food Saf. 15, 37–47.
410

411 OMNRF, 2016. Ontario Ministry of Natural Resources and Forestry 2010 Survey of Recreational
412   Fishing in Canada: Selected Results for the Great Lakes Fishery, 2010. Species Conservation
413   Policy Branch, Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario. 29
414   p. + Appendices.
415

416 Poste, A.E., Hecky, R.E., Guildford, S.J., 2011. Evaluating microcystin exposure risk through fish
417   consumption. Environ. Sci. Technol. 45, 5806–5811.
418

419 Qian, S.S., Stow, C.A., Rowland, F.E., Liu, Q., Rowe, M.D., Anderson, E.J., Stumpf, R.P.,
420   Johengen, T.H., 2021. Chlorophyll a as an indicator of microcystin: Short-term forecasting and
421   risk assessment in Lake Erie. Ecol. Indic. 130, 108055.
422
423 Reid, A.J., Carlson, A.K., Creed, I.F., Eliason, E.J., Gell, P.A., Johnson, P.T.J., Kidd, K.A.,
424   MacCormack, T.J., Olden, J.D., Ormerod, S.J., Smol, J.P., Taylor, W.W., Tockner, K.,
425   Vermaire, J.C., Dudgeon, D., Cooke, S.J., 2019. Emerging threats and persistent conservation
426   challenges for freshwater biodiversity. Biol. Rev. Camb. Philos. Soc. 94, 849–873.
427
428 Schmidt, J.R., Wilhelm, S.W., Boyer, G.L., 2014. The fate of microcystins in the environment and
429   challenges for monitoring. Toxins 6, 3354–3387.
430
431 Shahmohamadloo, R.S., Bhavsar, S.P., Ortiz Almirall, X., Marklevitz, S.A.C., Rudman, S.M.,
432   Sibley, P.K., 2023a. Lake Erie fish safe to eat yet afflicted by algal hepatotoxins. Sci. Total
433   Environ. 861, 160474.
434
435 Shahmohamadloo, R.S., Frenken, T., Rudman, S.M., van West, P., Ibelings, B.W., Trainer, V.L.,
436   2023b. Diseases and Disorders in Fish due to Harmful Algal Blooms, in: Woo, P.T.K.,
437   Subasinghe, R.P. (Eds.), Climate Change on Diseases And Disorders Of Finfish In Cage
438   Culture, 3rd Edition. CAB International, pp. 387–429.
439
440 Shahmohamadloo, R.S., Ortiz Almirall, X., Simmons, D.B.D., Lumsden, J.S., Bhavsar, S.P.,
441   Watson-Leung, T., Eyken, A.V., Hankins, G., Hubbs, K., Konopelko, P., Sarnacki, M., Strong,
442   D., Sibley, P.K., 2021. Cyanotoxins within and Outside of Microcystis aeruginosa Cause
443   Adverse Effects in Rainbow Trout (Oncorhynchus mykiss). Environ. Sci. Technol. 55, 10422–
444   10431.
445
446 Shahmohamadloo, R.S., Ortiz Almirall, X., Simmons, D.B.D., Poirier, D.G., Bhavsar, S.P., Sibley,
447   P.K., 2022. Fish tissue accumulation and proteomic response to microcystins is species-
448   dependent. Chemosphere 287, 132028.
449
450 Shahmohamadloo, R.S., Poirier, D.G., Ortiz Almirall, X., Bhavsar, S.P., Sibley, P.K., 2020a.
451   Assessing the toxicity of cell-bound microcystins on freshwater pelagic and benthic
452   invertebrates. Ecotoxicol. Environ. Saf. 188, 109945.
453
454 Shahmohamadloo, R.S., Simmons, D.B.D., Sibley, P.K., 2020b. Shotgun proteomics analysis
455   reveals sub-lethal effects in Daphnia magna exposed to cell-bound microcystins produced by
456   Microcystis aeruginosa. Comp. Biochem. Physiol. Part D Genomics Proteomics 33, 100656.
457
458 Shimizu, K., Sano, T., Kubota, R., Kobayashi, N., Tahara, M., Obama, T., Sugimoto, N., Nishimura,
459   T., Ikarashi, Y., 2013. Effects of the amino acid constituents of microcystin variants on
460   cytotoxicity to primary cultured rat hepatocytes. Toxins 6, 168–179.
461
462 Smith, R.B., Bass, B., Sawyer, D., Depew, D., Watson, S.B., 2019. Estimating the economic costs
463   of algal blooms in the Canadian Lake Erie Basin. Harmful Algae 87, 101624.
464
465 Taranu, Z.E., Pick, F.R., Creed, I.F., Zastepa, A., Watson, S.B., 2019. Meteorological and Nutrient
466   Conditions Influence Microcystin Congeners in Freshwaters. Toxins 11.
467

468 WHO, 2020. Cyanobacterial toxins: microcystins. Background document for development of WHO
469   Guidelines for drinking-water quality and Guidelines for safe recreational water environments.
470   Geneva: World Health Organization; 2020 (WHO/HEP/ECH/WSH/2020.6). Licence: CC BY-
471   NCSA 3.0 IGO.
472
473 Wituszynski, D.M., Hu, C., Zhang, F., Chaffin, J.D., Lee, J., Ludsin, S.A., Martin, J.F., 2017.
474   Microcystin in Lake Erie fish: Risk to human health and relationship to cyanobacterial blooms.
475   J. Great Lakes Res. 43, 1084–1090.
476
477 Xie, L., Xie, P., Ozawa, K., Honma, T., Yokoyama, A., Park, H.-D., 2004. Dynamics of
478   microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity
479   experiment. Environ. Pollut. 127, 431–439.
480
481 Zhang, D., Xie, P., Liu, Y., Qiu, T., 2009. Transfer, distribution and bioaccumulation of microcystins
482   in the aquatic food web in Lake Taihu, China, with potential risks to human health. Sci. Total
483   Environ. 407, 2191–2199.
484

485 **Figures**
486
487 **Figure 1.** Total microcystins (ng g^{-1} ww) in muscles across seven fish species (Largemouth Bass,
488 Northern Pike, Smallmouth Bass, Rock Bass, Walleye, White Bass, Yellow Perch) collected pre-
489 and post-harmful algal bloom in Lake St. Clair. The method detection limit (2.18 ng g^{-1} ww) is
490 denoted by a dashed line. Refer to Table 1 for the number of fish analyzed per species.
491

492
493
494 **Figure 2.** Total microcystins (ng g^{-1} ww) in livers across three fish species (Walleye, White Bass,
495 Yellow Perch) collected pre- and post-harmful algal bloom in Lake St. Clair. The method detection
496 limit (2.18 ng g^{-1} ww) is denoted by a dashed line. Refer to Table 1 for the number of fish analyzed
497 per species.
498

499
500

501 **Table**

503 **Table 1.** Fish collected from Lake St. Clair pre- and post-HAB for microcystin analysis in livers and
504 muscles. Data is reported in mean±SD.

Species	n	Length (cm)	Weight (g)	Microcystins in liver (ng g^{-1} ww)	Microcystins in muscle (ng g^{-1} ww)	EDI (μg kg^{-1} body weight day^{-1}) [a]
Largemouth Bass	3	38±13	1133±833	-	1.9±1.4	0±0.005
Northern Pike	12	74±10	2500±1055	-	0.4±0.9	0±0.003
Rock Bass	16	23±3	250±82	-	1.1±2.1	0±0.007
Smallmouth Bass	25	43±5	1334±559	-	1.0±3.1	0±0.01
Walleye	42	48±10	1059±690	217.2±220.5	1.8±2.5	0±0.008
White Bass	12	36±3	566±161	120.4±17.8	0.6±1.3	0±0.004
Yellow Perch	16	25±2	161±59	233.9±359.8	1.9±2.4	0±0.008

506 [a] Estimated daily intake (EDI) is measured as ng g^{-1} d^{-1}, assuming one fish meal is 227 g and applying the mean tissue concentration as
507 the daily intake value. Values are compared against the World Health Organization's tolerable daily intake value (0.04 μg kg^{-1} body
508 weight day^{-1}).

509