Potentiation of Adipogenesis by Reactive Oxygen Species is a Unifying Mechanism in the Pro-adipogenic Properties of Bisphenol A and its New Structural Analogues.

Radha D. Singh,[1,2,3] Jessica L. Wager,[1,2,3] Taylor B Scheidl,[1,2,3] Liam T. Connors,[1,2,3] Sarah Easson[1,2,3] Mikyla A. Callaghan,[1,3] Samuel Alatorre-Hinojosa,[1] Lucy H. Swift,[1,4] Pina Colarusso,[4] Anshul Jadli,[1,2] Timothy E. Shutt,[3,5,6,7] Vaibhav Patel,[1,2] Jennifer A. Thompson[1,2,3*]

Department of Physiology and Pharmacology,[1] Libin Cardiovascular Institute,[2] Alberta Children's Hospital Research Institute,[3] Snyder Institute for Chronic Diseases,[4] Department of Medical Genetics,[5] Department of Biochemistry and Molecular Biology,[6] Hotchkiss Brain Institute,[7] University of Calgary, Calgary, AB, T2N 4N1

Running title: The role of reactive oxygen species in mediating the effect of bisphenols on adipogenesis

Corresponding Author:
Jennifer Thompson PhD
Department of Physiology and Pharmacology
Cumming School of Medicine
University of Calgary
Calgary, AB
T2N 4N1
Email: jennifer.thompson2@ucalgary.ca
Twitter: @JThompsonlab

Keywords: Stem cells, adipogenesis, bisphenol, environmental toxicants, redox signaling

Singh *et al.*

ABSTRACT

Aims: Structural analogues of bisphenol A (BPA), including BPS and BPF, are emerging

environmental toxicants as their presence in the environment is rising since new regulatory

restrictions were placed on BPA-containing infant products. The adipogenesis-enhancing effect of

bisphenols may explain the link between human exposure and metabolic disease; however,

underlying molecular pathways remain unresolved. Methods: Adipose-derived progenitors were

isolated from mice and exposed to various concentrations of BPS, BPA or BPF before induction

of adipogenesis. RNAseq in BPS-exposed progenitors revealed modulation in redox pathways.

The role of reactive oxygen species (ROS) was assessed by measuring the degree of adipogenesis

in the presence or absence of antioxidants. ROS production and mitochondria function were

determined by fluorescent assays. Fat mass was measured by TD-NMR in adult mice exposed to

BPS during *in utero* establishment of the adipocyte progenitor pool, and in adult mice exposed to

BPS after weaning. Results: Exposure of progenitors to BPS, BPF, BPA or ROS generators

enhanced lipid droplet formation and expression of adipogenic markers after induction of

differentiation. ROS was higher in bisphenol-exposed cells, while co-treatment with antioxidants

attenuated adipogenesis and abolished the effect of BPS. There was a loss of mitochondria

membrane potential in BPS-exposed cells and mitochondria-derived ROS contributed to

potentiation of adipogenesis by BPS and its analogues. Male mice exposed to BPS during gestation

had higher adiposity, while postnatal exposure had no impact on adiposity in either sex. ROS act

as signaling molecules in the regulation of adipocyte differentiation and mediate bisphenol-

induced potentiation of adipogenesis.

INTRODUCTION

The *Lancet* Commission on pollution and health identified chemical intensification of the

environment as one of the most significant causes of premature death (Landrigan et al., 2018). The

plastic industry is a major contributor to the environmental chemical burden, emitting 400 million

tonnes of synthetic chemicals each year (Naidu et al., 2021). Bisphenols, plasticizers used in the

manufacturing of polycarbonate plastics and epoxy resins, are classified as endocrine disrupting

chemicals (EDC), which interfere with developmental and reproductive processes by mimicking

endogenous ligands to receptors and acting as agonists or antagonists. After regulatory bans on the

sale or import of baby products containing bisphenol A (BPA), manufacturers responded by

replacing it with structural analogues, marketing them as 'safe' substitutes. However, evidence

supporting this claim of health benefit is lacking and emerging data suggest that BPA analogues

are another example of 'regrettable substitution', where toxic chemicals are replaced with equally

toxic substitutes (Zimmerman and Anastas, 2015). BPA substitutes, such as bisphenol S (BPS)

and bisphenol F (BPF), are now recognized as emerging toxicants as their production is on the

rise, approaching or exceeding the world average daily intake of BPA in countries that introduced

regulatory restrictions (Wang et al., 2020). Therefore, there is a need to fill the gap in knowledge

with respect to health effects and biological properties of the new BPA substitutes.

A large body of published data show that BPA promotes adipogenesis *in vitro* (Ariemma et al.,

2016, Ohlstein et al., 2014). Fewer studies have investigated the structural analogues of BPA;

however, the available evidence to-date reveal that commonly used substitutes, BPS and BPF,

exhibit similar pro-adipogenic properties as their predecessor, with some studies showing effects

at lower doses (Ramskov Tetzlaff et al., 2020). Investigation into the molecular pathways

mediating the adipogenic effect of bisphenols have almost exclusively focused on endocrine

pathways as it is well known that BPA and its analogues bind to estrogen, androgen and other

hormone receptors.   These studies have yielded inconsistent results with some showing a role for

estrogen receptors and others a role for glucocorticoid receptor (Ahmed and Atlas, 2016, Boucher

et al., 2016). Overall, the cellular mechanisms responsible for the adipogenesis-promoting

properties of bisphenols remain unresolved.

Adipogenesis is a sequential differentiation process by which mesenchymal stem cells commit to

the adipocyte lineage and terminally differentiate into lipid-storing adipocytes. Specification of

stem cells to the adipocyte fate in subcutaneous adipose tissue (SAT), the primary site of energy

storage, is restricted to late fetal life (Jiang et al., 2014, Wang et al., 2013). Establishment of the

progenitor pool is followed by two periods of rapid fat accumulation occurring after birth and

during puberty, with adipocyte numbers stabilizing thereafter (Holtrup et al., 2017). In adult

adipose depots, *de novo* adipogenesis is stimulated to support normal cellular turnover and to

increase lipid storage capacity in response to an obesogenic environment (Jeffery et al., 2015,

Wang et al., 2013, White and Ravussin, 2019). Current evidence suggests that new adipocytes

arising in adult depots derive from a distinct compartment of progenitors that are also specified to

the adipocyte lineage *in utero* (Jiang et al., 2014, Wang et al., 2015). Thus, the early life window

of adipogenesis establishes the setpoint of adiposity and programs lipid buffering capacity in adult

depots.

Herein, we set out to determine the effect of BPS on adipogenesis and elucidate the molecular

pathways involved. Our data reveal a critical role for reactive oxygen species (ROS) in acting as

physiological signaling molecules in the regulation of adipogenesis under normal conditions.

Increased ROS production, contributed by mitochondria dysfunction, was responsible for the

potentiation of adipogenesis in progenitors exposed to BPS as well as other structural analogues.

Thus, bisphenol-induced ROS may be a unifying mechanism that explains the pro-adipogenic

properties of BPA and its substitutes. Second, we exposed mice to an environmentally relevant

dose of BPS during or after the early life window of adipogenesis and determined that only early-

life exposure had an impact on adiposity in adulthood. Therefore, exposure to BPS before birth

may raise the setpoint for adiposity, predisposing to obesity.

MATERIAL AND METHODS

Isolation and differentiation of adipocyte progenitor cells:

All animal procedures were approved by the University of Calgary Animal Care Committee

(AC19-0006) and conducted in accordance with guidelines by the Canadian Council on Animal

Care Ethics. At 6-8 weeks of age, C57BL/J6 male mice were sacrificed for harvesting of inguinal

SAT (iSAT), which was digested in a collagenase I digestion buffer. The pelleted stromal vascular

fraction (SVF) was treated with RBC lysis buffer, resuspended in preadipocyte growth medium

(Cell Applications), and cultured at 37 °C and 5% CO_2. After a single passage, cells were seeded,

and differentiation was induced 48 hours after contact inhibition (Day 0) until day 5 when

differentiation medium was replaced with maintenance medium (Cell Applications). Prior to

contact inhibition (90% confluency) cells were treated with various doses of BPS, BPA, BPF or

ROS generators or vehicle (DMSO), in the presence or absence of antioxidants. Lipid droplets

were stained with an Oil Red O solution (Sigma) or 2 µM BODIPY (Thermo Fisher Scientific),

captured using a Nikon Eclipse Ts2 microscope and processed with NIS-Elements D 5.11.00. After

thorough washing, Oil-Red O-stained cells were incubated with a lysis buffer and the absorbance

of the eluted dye quantified in triplicates at 490nm on a spectrophotometer (Biotek 800TS).

Prenatal and postnatal exposure to BPS

After weaning, female C57BL/J6 mice (Jackson Laboratories) were transferred to cages equipped

with glass water bottles (Lab Products LLC) and absent of plastic enrichment to eliminate

background exposure to plasticizers. For prenatal exposure, 12-week-old dams were randomized

to BPS drinking water (2 µg/kg body weight/day) or vehicle control, started on a phytoestrogen-

low diet (Teklad 2020) and mated with males. Exposure was initiated at gestational day 0 (Gd0)

after confirmation of pregnancy by observation of a copulatory plug. Dams were allowed to deliver

spontaneously, and litters culled to a maximum of 6 to minimize litter effects. Exposure continued

through lactation until postnatal day 21 (Pd21), at which time pups were weaned and maintained

on a plastic-reduced environment. For postnatal exposure, littermates were randomized to BPS or

vehicle drinking water after weaning. At 12-weeks of age, mice were sacrificed and whole-body

fat mass measured with a TD-NMR body composition analyzer (Bruker). Fat mass of litter mates

was averaged for prenatally-exposed offspring to control for litter effects, while litter mates were

compared in postnatally-exposed mice.

RNAseq and Pathway Analysis

RNA samples were assessed with TapeStation and Oubit assays and 1000ng RNA was used in

library preparation using the NEBNext Ultra II directional RNA library prep kit for Illumina, along

with the NEBNext Poly(A) mRNA Magnetic Isolation Module. Final libraries were assessed with

the Kapa library quantification qPCR assay and sequencing performed on the NextSeq 75 cycle

high output run. The sequencing run passed all QC metrics. Samples were quantified using Kallisto

v0.42.4 against the NCBI RefSeq transcriptome (GRCm38) (Bray et al., 2016). A linear regression

model was used to determine differentially expressed transcripts, with terms for treatment as well

as sample admixture proxy markers identified in principal component analysis. Differentially

expressed genes (DEG) were considered as those with a false discover rate (Benjamini-Hochberg

corrected p-value) < 0.05 under the Wald test in Sleuth v0.30.0 (Pimentel et al., 2017). DEGs were

subsequently annotated and analyzed using the Molecular Signatures Database (MSigDB)

hallmark pathway analysis using Enrichr (https://maayanlab.cloud/Enrichr/), a comprehensive

gene set enrichment analysis web server. A heatmap of DEGs involved in adipogenesis and the

Nrf2-mediated oxidative stress response was created using R studio software (2022.02.0+443).

The potential interaction between DEGs involved in these two pathways was presented using

Interaction Network Analysis using the Path Explorer Tool of Qiagen IPA software.

Quantitative real time PCR

Total RNA was isolated using the RNeasy Mini Kit (Qiagen), assessed for integrity using a

TapeStation RNA assay (Agilent) and quantified using a N50 Nanophotometer (Implen Inc.).

Following DNase treatment, cDNA synthesis was performed using a High-Capacity cDNA

Reverse Transcription kit (Applied Biosystems). cDNA products were run in triplicates for

Quantitative real time PCR using a QuantStudio 5 Real-time PCR System (Applied Biosystems)

and Powerup SYBR green master mix (Applied Biosystems). Primers were designed using the

NCBI/Primer Blast tool (Supplementary Table 1). Target mRNA expression was normalized to β-

actin and fold change calculated using $2^{-\Delta\Delta Ct}$ method.

Western blot

Proteins were extracted from cell lysates in buffer containing a protease and phosphatase inhibitor

cocktail (Invitrogen) and resolved in precasted NuPAGE 4%–12% Bis-Tris polyacrylamide gels

(Invitrogen) and then transferred to Amersham Hybond PVDF membranes (GE Healthcare Life

Sciences) at 100V for 2 hr. The membrane was then washed, blocked with 5% blotting grade

blocker (Bio Rad) for 1 hr and probed with primary antibodies. After washing and re-probing with

a horseradish peroxidase-conjugated secondary antibody, bands were visualized with the iBright

CL1500 Imaging system (Applied Biosystems).

ROS detection

For the detection of cellular ROS, cells were prestained with 2′,7′-dichlorodihydrofluorescein

diacetate (DCFDA, Sigma), washed and incubated in BPS at 37 °C for 45 min. CellROX™ deep

red dye (5µM, 640/665 nm, Invitrogen) and MitoSOX™ Red dye (2.5 µM, 510/580 nm,

Invitrogen) were used to detect cellular ROS and mitochondria-derived ROS, respectively, after

24 or 48 hr of BPS exposure. Relative Fluorescent Unit (RFU) was measured with a

spectrophotometer (SpectraMax M2) and normalized to background fluorescence and protein

concentration. Flowcytometric analysis was carried out using an Attune® Acoustic Focusing

Cytometer (ThermoFisher Scientific) after cells were dissociated using 5 mM EDTA in HBSS,

centrifuged at 500g and resuspended in HBSS. Stained cells were imaged with the 60X objective

using a Nikon Ti Eclipse Widefield microscope, employing the DAPI, CY5, and CY4 filters.

Mitochondria function

After 24 hrs of BPS exposure, cells were incubated in 2 µM JC-1 (5, 5', 6, 6'-tetrachloro-1, 1', 3,

3'-tetraethylbenzimidazolylcarbocyanine iodide, Invitrogen) at 37°C for 15 min. After washing,

fluorescence signals were measured at 514/529 (green, monomer) and 514/590 (red, aggregate)

using SpectraMax® M2 and the mitochondria membrane potential expressed as the ratio of green

to red fluorescence. The mitochondria membrane potential was also assessed in live cells exposed

to BPS for 24 hrs using tetramethyl rhodamine ethyl ester (TMRE, Thermo Fisher Scientific)

staining. Live cells were collected in HBSS containing 5 mM EDTA, washed and resuspended in

media and the TMRE mean fluorescent signal quantified on a flow cytometer. To observe

mitochondria morphology, progenitors were stained with MitoTracker Green (Invitrogen).

Statistical Analysis:

Statistical analyses were performed using GraphPad Prism 9. Two-tailed student's t-test or One-

way ANOVA with Dunnett's or Tukey's multiple comparison test were used to evaluate

differences between groups. All data in this study are represented as the mean ± standard error of

mean (SEM) or maximum, minimum, and median. P values <0.05 are considered statistically

significant.

RESULTS

BPS, BPF and BPA potentiate differentiation of adipocyte progenitors *in vitro*

Adipocyte progenitors isolated from inguinal SAT (iSAT) were exposed to various concentrations

of BPS prior to differentiation. Exposure of progenitors to BPS increased lipid droplet formation

on day 2, 4 & 7 post-differentiation, albeit in a non-monotonic manner such that the increase in

differentiation at low doses was attenuated at the highest dose (Figure 2A & B). The MTT assay

revealed a decrease in cell viability at 25 µM, the highest dose of BPS (Supplementary Figure 1),

suggesting that the attenuation of lipid droplet staining at this concentration of BPS is due to

cytotoxicity. On day 7 of differentiation, lipid droplet staining was higher in progenitors exposed

to BPA or BPF (Supplementary Figure 2A &B). The adipogenic response to BPA occurred at

higher doses, while the attenuation of the adipogenic response at 25 µM of BPS was not observed

at the same concentration of BPA or BPF. On day 2 of differentiation in BPS-exposed progenitors,

there was an increase in mRNA expression of peroxisome proliferator-activated receptor gamma

(*Ppar*γ), a nuclear receptor required for adipogenesis, as well as glucose transporter type 4 (*Glut4*),

an adipogenic marker (Figure 2C). On day 2, the mRNA expression of CCAAT/enhancer binding

protein beta (*C/ebpβ*), an early regulator of differentiation, was decreased in BPS-exposed

progenitors (Figure 2C). *Glut4* mRNA expression remained high on day 4 and there was an

increase in stearoyl-CoA desaturase 1 (*Scd1*), a key enzyme involved in lipogenesis (Figure 2C).

Protein expression of adiponectin, SCD1, fatty acid binding protein 4 (FABP4) and C/EBPβ was

increased on day 2 and 4 of differentiation in BPS-exposed progenitors (Figure 2D & E).

BPS exposure modulates gene pathways involved in oxidative stress responses

RNAseq analysis was performed in extracted RNA collected from adipose-derived progenitors on

D0, at which time changes in gene expression patterns and epigenetic signatures stimulated by

contact inhibition prime the cells for differentiation (Guo et al., 2016). There were over 30 DEGs

involved in adipogenesis and over 12 genes involved in Nrf-2-mediated responses to oxidative

stress between vehicle-treated and BPS-exposed cells (Figure 3A). MSigDB Hallmark Pathway

Analysis showed that DEGs were enriched in pathways regulating cellular stress (reactive oxygen

species; hypoxia), inflammatory/immune cascades (TNFα/NF-κB; IL6/JAK/STAT3; Interferon

α/γ), responses to DNA damage (UV light response; G2-M checkpoint), as well as cell growth and

death (p53 pathway; KRAS signaling; apoptosis; mTORC1) (Figure 3B). Interaction Networks

Analysis shows the predicted relationship between DEGs involved in pathways regulating

adipogenesis and Nrf2-mediated oxidative stress responses (Figure 3C). Validation by qPCR of

DEGs identified by RNAseq with fold changes >1 confirmed a downregulation of antioxidant

genes, catalase (*Cat*) and superoxide dismutase 3 (*Sod3*), and upregulation of *c-FOS*, a redox-

sensitive early response gene involved in the regulation of cellular proliferation and differentiation

(Figure 3D).

ROS production is increased in progenitors exposed to bisphenol analogues

In undifferentiated progenitors a surge in ROS generation was observed in response to acute BPS

exposure in a concentration-dependent manner, as assessed by quantification of DCFDA (Figure

4A). Similarly, detection of DCFDA by fluorescence spectroscopy revealed increased ROS

production after acute exposure to BPA (Figure 4B) or BPF (Figure 4C). The number of cells

stained with DCFDA was quantified by flow cytometry, also showing increased ROS production

after acute BPS exposure (Figure 4D & E). After 48 hrs of BPS exposure, staining intensity of

CellROX™ deep red, a fluorescent probe that detects oxidative stress, was higher in BPS-exposed

cells (Figure 4F & G). Similarly, CellROX staining was increased in undifferentiated progenitors

exposed to BPA (Figure 4H) or BPF (Figure 4I) for 48 hrs. These data show that BPA and its

structural analogues stimulate an increase ROS production.

ROS regulate adipogenesis and mediate BPS-induced potentiation of adipogenesis.

Exposure of progenitors to low concentrations of the ROS generator, menadione, augmented lipid

droplet formation on day 4 post-differentiation in a non-monotonic manner similar to BPS, such

that increased lipid droplets at lower concentrations was attenuated with higher concentrations

(Figure 5A). On day 2 post-differentiation in progenitors exposed to menadione, mRNA

expression of *Pparγ* and *Glut4* was increased, while expression of *C/ebpβ* was decreased (Figure

5B). Similarly, exposure of progenitors to another ROS generator, 2,3-dimethoxy-1,4-

napthalenedione (DMNQ), lead to a non-monotonic increase in lipid droplet staining on day 4

(Figure 5C) and an increase in mRNA expression of *Pparγ* and *Glut4* on day 2 (Figure 5D). In the

absence of BPS, suppression of ROS generation via a ROS scavenger (Tempol), led to a

concentration-dependent decrease in lipid droplet staining after differentiation (Figure 5E). Co-

treatment of progenitors with either Tempol or the antioxidant, apocynin, abolished the effect of

BPS on adipogenesis (Figure 5F-H). Together, these data reveal ROS to be critical regulators of

adipogenesis and mediators of BPS-induced potentiation of adipogenesis.

Dysfunctional mitochondria are important sources of ROS in BPS-exposed progenitors

Tightly regulated release of ROS from healthy mitochondria plays a role in modulating signaling

cascades, whereas dysfunctional mitochondria produce excess ROS that contribute to oxidative

stress. Mitochondrial transmembrane polarization in undifferentiated progenitors was measured

by the JC-1 and TMRE assays. A loss of membrane potential in BPS-exposed progenitors was

apparent by a higher ratio of cationic JC-1 monomers relative to the J-aggregates formed when JC-

1 enters the mitochondria (Figure 6A) and a lower accumulation of the cationic dye, TMRE (Figure

6B). Live cell staining with MitoTracker Green revealed elongated mitochondria in vehicle-treated

cells and fragmented mitochondria in BPS-treated progenitors (Figure 6C). Mitochondrial-derived

superoxide (O_2^-) was increased in BPS-exposed progenitors (Figures 6D-F), while co-treatment of

progenitors with the mitochondria-targeted antioxidant, MitoQ, markedly attenuated lipid droplet

formation on day 4 of differentiation and abolished BPS-induced potentiation of adipogenesis

(Figure 6G & I). Similarly, MitoQ attenuated the adipogenesis-enhancing effects of BPA and BPF

(Supplementary Figure 2C &D). Co-treatment with the mitochondria-targeted antioxidant, SS31,

which also protects against membrane depolarization, attenuated lipid droplet formation in

differentiated progenitors exposed to BPS without having an impact on control cells (Figure 6H).

Together, these data highlight mitochondria as sources of ROS that play a role in the regulation of

adipogenesis and in the bisphenol-induced adipogenic response.

Prenatal BPS exposure programs higher adiposity in adulthood

To determine the impact of *in vivo* BPS exposure on adiposity, an environmentally relevant dose

of BPS (2 μg/kg body weight) was administered via glass water bottles to C57BL6 mice housed

in an environment absent of plastic enrichment. Exposure occurred during the early life window

of adipocyte commitment and differentiation (Gd0-Pd21) by exposing pregnant and lactating dams

or in postnatal life after weaning (Figure 7A & B). Exposure to BPS during early life resulted in

higher whole-body fat mass in adult male offspring, but not female offspring (Figure 7C). In

contrast, exposure to BPS during postnatal life did not influence adiposity in adulthood in either

male or female mice (Figure 7D).

DISCUSSION

Studies to-date have shown structural analogues marketed as safer substitutes for BPA to exhibit

similar adipogenic properties as their predecessor (Ahmed and Atlas, 2016, Boucher et al., 2016).

The adipogenesis-enhancing effects of BPA and its analogues have been attributed to their

endocrine disrupting behaviour (Boucher et al., 2016), as they are classified as EDC that bind to

estrogen, androgen, and glucocorticoid receptors, albeit with lower affinity and activity than

endogenous ligands. Studies exploring endocrine pathways as mediators of the pro-adipogenic

effects of bisphenols have yielded inconsistent findings with some reporting a role for estrogen

and others showing a role for glucocorticoids (Ahmed and Atlas, 2016, Boucher et al., 2014).

Results of the present study reveal a novel role for ROS in mediating the adipogenesis-promoting

effect of BPS. Similar to BPS, BPA and BPF increased ROS production and had a potentiating

effect on adipogenesis that was abolished by co-treatment with antioxidants. Therefore, cellular

stress leading to heightened ROS production may be a common mechanism that explains the pro-

adipogenic properties of BPA and its structural analogues. While it is well established that

bisphenols and other EDC like phthalates promote oxidative stress in a variety of cell types

(Špačková et al., 2020, Xie et al., 2020); to our knowledge this is the first study to explore the role

of ROS in mediating the pro-adipogenic properties of EDC.

A handful of published studies demonstrate ROS to act as mediators of adipocyte differentiation

(Schröder et al., 2009, Tormos et al., 2011); however, the role of redox signaling in adipogenesis

has not received wide attention. Our data substantiate these studies as exposure of progenitors to

low concentrations of O_2^- generators enhanced differentiation, while antioxidant treatment

inhibited adipogenesis in the absence of BPS. A non-monotonic response to increasing

concentrations of BPS or ROS generators was observed, such that the increased differentiation at

lower concentrations was attenuated at the highest dose. Cell viability was unaffected at BPS

concentrations that enhanced adipogenesis, but reduced at the highest concentration of BPS,

suggesting that the nonmonotonicity is due to cytotoxicity at higher levels of ROS production.

Thus, cellular ROS promote adipocyte differentiation within a range of physiological

concentrations below the cytotoxic threshold.

When produced at high levels in response to cellular stress, ROS damage macromolecules and

contribute to cellular dysfunction, but at physiological levels, act as second messengers in

signaling cascades that regulate a wide variety of cellular processes. ROS derive from several

sources including mitochondria and enzymes such as the NADPH oxidases and xanthine oxidase.

Silencing of NADPH oxidase 4 (Nox4) by RNA interference in bone marrow-derived

mesenchymal stem cells or by siRNA in 3T3-L1 cells, abrogated lipid droplet formation after

differentiation, implicating Nox4-derived hydrogen peroxide (H_2O_2) in the regulation of

adipogenesis (Kanda et al., 2011, Schröder et al., 2009). Our findings cannot rule out a role for

H_2O_2 as antioxidants and O_2^- generators like DMNQ and menadione will influence levels of both

O_2^- and H_2O_2, as the former quickly dismutates to H_2O_2. Our data show that BPS exposure

increases DCFDA staining of cellular ROS as well as O_2^- levels inside mitochondria detected by

MitoSox. Mitochondria are both generators and targets of ROS and crosstalk with enzymatic

sources of ROS to promote an environment of oxidative stress. Thus, while our findings show

mitochondria to be important contributors to BPS-induced oxidative stress, contributions from

other sources of ROS cannot be ruled out.

Our data support studies showing that mitochondria-derived ROS are required for the induction of

adipogenesis (Carrière et al., 2003, Tormos et al., 2011), as mitochondria-targeted antioxidants

attenuated differentiation even in the absence of BPS. An increase in oxygen consumption during

activation of the adipogenic program is driven by mitochondria biogenesis leading to an increase

in mitochondria content (Li et al., 2017). Distinct subpopulations of mitochondria tethered to lipid

droplets support formation and expansion of lipid droplets (Benador et al., 2019), the final steps

in terminal differentiation. Mitochondria are major sources of ROS as complexes of the electron

transport chain (ETC) leak electrons that combine with oxygen to produce O_2^-. The increase in

oxidative phosphorylation that occurs during adipocyte differentiation is accompanied by an

increase in ROS production (Tormos et al., 2011, Zhang et al., 2013). Decreasing mitochondria

respiration by inhibiting the ETC abrogated differentiation of human mesenchymal stem cells into

adipocytes (Zhang et al., 2013). Interestingly, a study by Tormos *et al*. showed that knockdown of

a complex III subunit required for O_2^- production attenuated differentiation of adipocyte

progenitors, whereas adipogenesis was unaffected when complex III-derived O_2^- was maintained

with knockdown of a complex II subunit (Tormos et al., 2011). These findings demonstrate that

an increase in mitochondria-derived ROS are not only by-products of differentiation but play a

direct role in regulating adipogenesis that is independent of oxidative phosphorylation. Findings

of the current study show that increased mitochondria-derived ROS in BPS exposed

undifferentiated stem cells was accompanied by a loss of mitochondria membrane potential,

indicative of mitochondria injury. Therefore, disruption of mitochondria leading to moderate levels

of ROS may potentiate differentiation, while more extensive mitochondrial damage and oxidative

stress may interfere with the mitochondrial dynamics required for successful differentiation.

Moderate mitochondria membrane depolarization has been shown to serve as a protective

mechanism that preserves energetic capacity while mitigating mitochondria-derived ROS under

conditions of ischemia (Vyssokikh et al., 2020). Therefore, it is possible that the mitochondria

membrane depolarization in BPS-exposed progenitors is a compensatory response to cellular

stress.

Although *in vitro* studies consistently show an adipogenesis-promoting effect of BPA and its

analogues, human and animal studies have yielded inconsistent results with respect to the

relationship between bisphenol exposure and adiposity (Callaghan et al., 2020). Adipogenic

responses *in vitro* do not necessarily reflect adipogenesis *in vivo*, the latter influenced heavily by

the microenvironment of the adipose depot. Further, exposure to bisphenols have pleiotropic

effects with the potential to impact metabolism and adiposity through multiple mechanisms.

Nevertheless, several epidemiological studies have demonstrated relationships between urinary

levels of BPA or its analogues and components of the metabolic syndrome including abdominal

obesity, hypertension, and insulin resistance (Callaghan et al., 2020). Most human studies use

single timepoint measurements of bisphenol levels and are unable to assess low dose affects due

to ubiquitous exposure. Further, humans are not exposed to single toxicants, but rather toxicant

mixtures, and correlates to exposure such as consumption of processed foods are confounding

variables. Animal studies vary widely in the dose, timing, and duration of bisphenol exposure,

some reporting prenatal BPA exposure to result in increased adiposity and some reporting fetal

growth restriction (Rubin et al., 2019). The current study demonstrated that BPS exposure in

pregnant and lactating dams leads to an increase in adiposity in adult male offspring, while

postnatal exposure had no impact on fat mass. Seminal studies by Graff and others show that the

progenitor pool giving rise to adipocytes during postnatal expansion of subcutaneous adipose

depots are specified to the adipocyte fate before birth (Jeffery et al., 2015, Jiang et al., 2014, Wang

et al., 2013, Wang et al., 2015). Thus, our data show that a programmed predisposition to high

adiposity occurs when exposure coincides with this critical window of adipogenesis. However, it

is also possible that higher adiposity in adulthood is a secondary effect of other developmental

perturbations. For instance, some studies report maternal bisphenol exposure to result in fetal

growth restriction, which is well known to cause rapid catch-up growth and program a

predisposition to obesity.   The sexual dimorphism in programming of adiposity by early life BPS

exposure suggests an interaction of BPS with endocrine pathways that warrants further

investigation.

The dose used for *in vivo* exposure was below the tolerable daily intake (TDI) for BPA (4μg/kg

body weight/day) established by the European Food Safety Authority (EFSA). Currently, there are

no established limits for BPA analogues. Recent findings from China show that ingestion of BPS

from contaminated food, the primary route of human exposure, ranged between 5.74 to 56.9 ng/kg

body weight/day, exceeding that of BPA intake (Zhang et al., 2022). In Alberta, Canada, daily 24

hr intake of BPS in pregnant women reached up to 14 nM/kg body weight, approaching the TDI

of BPA (Liu et al., 2018). We acknowledge that prenatally-exposed and postnatally-exposed adult

mice were not exposed to the same level of BPS; however, our approach reflects the real-world

scenario where the fetus and lactating infant is exposed to a fraction of the mother's intake of BPS.

Nevertheless, our findings suggest that BPA analogues are obesogenic and reveal increased ROS

production as a contributing mechanism.

INNOVATION

Pro-adipogenic effects of BPA are well characterized; however, little is known regarding new

structural analogues that are replacing BPA since introduction of new regulatory restrictions and

heightened awareness of its adverse health effects. Molecular pathways underlying the

adipogenesis-promoting properties of bisphenols remain unresolved, and studies to-date have

almost exclusively focused on their endocrine disrupting properties. Herein, we undertook

extensive investigation that substantiates existing data identifying ROS act as signaling molecules

in the regulation of adipocyte differentiation and highlight ROS as a unifying mechanism that

explains the pro-adipogenic properties of BPA and its analogues that is independent of endocrine

disruption.

415 **ACKNOWLEDGEMENTS**

416 We would like to thank Paul Gordan and the Centre for Health Genomics and Informatics at the

417 University of Calgary for support in analyzing the RNAseq data. We also thank Dr. Steven

418 Greenway from the Libin Cardiovascular Institute for providing the mitochondria-targeted

419 antioxidant, SS31.

AUTHOR CONTRIBUTIONS

439 RDS contributed to the generation of hypotheses, experimental design, data collection, analysis,

440 and manuscript writing. JW, MC, TS, LC, SE and SAH contributed to data collection and analysis.

441 AH, VP and TS provided support for mitochondria analyses, while PC and LS provided support

442 for imaging.

AUTHOR DISCLOSURES

463   There are no conflicts of interest to declare.

FUNDING STATEMENT

The salary of RDS was supported by fellowships from the Molly Towell Perinatal Research

Foundation and the Libin Cardiovascular Institute. JW was supported by an O-Brien Summer

Studentship and MC was supported by scholarships from the Alberta Children's Hospital Research

Institute (ACHRI) and NSERC. The salaries of TS and LC were supported by scholarships from

the Libin Cardiovascular Institute. LC also received scholarships from the Cumming School of

Medicine and Faculty of Graduate Studies at the University of Calgary. The research performed

for this study was supported by start-up funds provided to JAT from the Libin Cardiovascular

Institute and an equipment grant from the Canadian Foundation for Innovation (CFI). RDS

received a research grant from ACHRI to support experiments for the current work. JAT was

supported by a National New Investigator Award from the Heart and Stroke Foundation of Canada

(HSFC).

REFERENCES

Ahmed S, Atlas E. Bisphenol S- and bisphenol A-induced adipogenesis of murine preadipocytes
occurs through direct peroxisome proliferator-activated receptor gamma activation. Int J Obes
(Lond) 2016;40(10):1566-1573; doi: 10.1038/ijo.2016.95.
Ariemma F, D'Esposito V, Liguoro D, et al. Low-Dose Bisphenol-A Impairs Adipogenesis and
Generates Dysfunctional 3T3-L1 Adipocytes. PLoS One 2016;11(3):e0150762; doi:
10.1371/journal.pone.0150762.
Benador IY, Veliova M, Liesa M, et al. Mitochondria Bound to Lipid Droplets: Where
Mitochondrial Dynamics Regulate Lipid Storage and Utilization. Cell Metab 2019;29(4):827-
835; doi: 10.1016/j.cmet.2019.02.011.
Boucher JG, Ahmed S, Atlas E. Bisphenol S Induces Adipogenesis in Primary Human
Preadipocytes From Female Donors. Endocrinology 2016;157(4):1397-1407; doi:
10.1210/en.2015-1872.
Boucher JG, Boudreau A, Atlas E. Bisphenol A induces differentiation of human preadipocytes
in the absence of glucocorticoid and is inhibited by an estrogen-receptor antagonist. Nutr
Diabetes 2014;4(1):e102; doi: 10.1038/nutd.2013.43.
Bray NL, Pimentel H, Melsted P, et al. Near-optimal probabilistic RNA-seq quantification. Nat
Biotechnol 2016;34(5):525-527; doi: 10.1038/nbt.3519.
Callaghan MA, Alatorre-Hinojosa S, Connors LT, et al. Plasticizers and Cardiovascular Health:
Role of Adipose Tissue Dysfunction. Front Pharmacol 2020;11:626448; doi:
10.3389/fphar.2020.626448.
Carrière A, Fernandez Y, Rigoulet M, et al. Inhibition of preadipocyte proliferation by
mitochondrial reactive oxygen species. FEBS Lett 2003;550(1-3):163-167; doi: 10.1016/s0014-
5793(03)00862-7.
Guo W, Chen J, Yang Y, et al. Epigenetic programming of Dnmt3a mediated by AP2α is
required for granting preadipocyte the ability to differentiate. Cell Death Dis 2016;7(12):e2496;
doi: 10.1038/cddis.2016.378.
Holtrup B, Church CD, Berry R, et al. Puberty is an important developmental period for the
establishment of adipose tissue mass and metabolic homeostasis. Adipocyte 2017;6(3):224-233;
doi: 10.1080/21623945.2017.1349042.
Jeffery E, Church CD, Holtrup B, et al. Rapid depot-specific activation of adipocyte precursor
cells at the onset of obesity. Nat Cell Biol 2015;17(4):376-385; doi: 10.1038/ncb3122.
Jiang Y, Berry DC, Tang W, et al. Independent stem cell lineages regulate adipose organogenesis
and adipose homeostasis. Cell Rep 2014;9(3):1007-1022; doi: 10.1016/j.celrep.2014.09.049.
Kanda Y, Hinata T, Kang SW, et al. Reactive oxygen species mediate adipocyte differentiation
in mesenchymal stem cells. Life Sci 2011;89(7-8):250-258; doi: 10.1016/j.lfs.2011.06.007.
Landrigan PJ, Fuller R, Acosta NJR, et al. The Lancet Commission on pollution and health.
Lancet 2018;391(10119):462-512; doi: 10.1016/s0140-6736(17)32345-0.
Li Q, Gao Z, Chen Y, et al. The role of mitochondria in osteogenic, adipogenic and
chondrogenic differentiation of mesenchymal stem cells. Protein Cell 2017;8(6):439-445; doi:
10.1007/s13238-017-0385-7.

Liu J, Wattar N, Field CJ, et al. Exposure and dietary sources of bisphenol A (BPA) and BPA-
alternatives among mothers in the APrON cohort study. Environ Int 2018;119:319-326; doi:
10.1016/j.envint.2018.07.001.
Naidu R, Biswas B, Willett IR, et al. Chemical pollution: A growing peril and potential
catastrophic risk to humanity. Environ Int 2021;156:106616; doi: 10.1016/j.envint.2021.106616.
Ohlstein JF, Strong AL, McLachlan JA, et al. Bisphenol A enhances adipogenic differentiation
of human adipose stromal/stem cells. J Mol Endocrinol 2014;53(3):345-353; doi: 10.1530/jme-
14-0052.
Pimentel H, Bray NL, Puente S, et al. Differential analysis of RNA-seq incorporating
quantification uncertainty. Nat Methods 2017;14(7):687-690; doi: 10.1038/nmeth.4324.
Ramskov Tetzlaff CN, Svingen T, Vinggaard AM, et al. Bisphenols B, E, F, and S and 4-
cumylphenol induce lipid accumulation in mouse adipocytes similarly to bisphenol A. Environ
Toxicol 2020;35(5):543-552; doi: 10.1002/tox.22889.
Rubin BS, Schaeberle CM, Soto AM. The Case for BPA as an Obesogen: Contributors to the
Controversy. Front Endocrinol (Lausanne) 2019;10:30; doi: 10.3389/fendo.2019.00030.
Schröder K, Wandzioch K, Helmcke I, et al. Nox4 acts as a switch between differentiation and
proliferation in preadipocytes. Arterioscler Thromb Vasc Biol 2009;29(2):239-245; doi:
10.1161/atvbaha.108.174219.
Špačková J, Oliveira D, Puškár M, et al. Endocrine-Independent Cytotoxicity of Bisphenol A Is
Mediated by Increased Levels of Reactive Oxygen Species and Affects Cell Cycle Progression. J
Agric Food Chem 2020;68(3):869-875; doi: 10.1021/acs.jafc.9b06853.
Tormos KV, Anso E, Hamanaka RB, et al. Mitochondrial complex III ROS regulate adipocyte
differentiation. Cell Metab 2011;14(4):537-544; doi: 10.1016/j.cmet.2011.08.007.
Vyssokikh MY, Holtze S, Averina OA, et al. Mild depolarization of the inner mitochondrial
membrane is a crucial component of an anti-aging program. Proc Natl Acad Sci U S A
2020;117(12):6491-6501; doi: 10.1073/pnas.1916414117.
Wang H, Liu ZH, Zhang J, et al. Human exposure of bisphenol A and its analogues:
understandings from human urinary excretion data and wastewater-based epidemiology. Environ
Sci Pollut Res Int 2020;27(3):3247-3256; doi: 10.1007/s11356-019-07111-9.
Wang QA, Tao C, Gupta RK, et al. Tracking adipogenesis during white adipose tissue
development, expansion and regeneration. Nat Med 2013;19(10):1338-1344; doi:
10.1038/nm.3324.
Wang QA, Tao C, Jiang L, et al. Distinct regulatory mechanisms governing embryonic versus
adult adipocyte maturation. Nat Cell Biol 2015;17(9):1099-1111; doi: 10.1038/ncb3217.
White U, Ravussin E. Dynamics of adipose tissue turnover in human metabolic health and
disease. Diabetologia 2019;62(1):17-23; doi: 10.1007/s00125-018-4732-x.
Xie C, Ge M, Jin J, et al. Mechanism investigation on Bisphenol S-induced oxidative stress and
inflammation in murine RAW264.7 cells: The role of NLRP3 inflammasome, TLR4, Nrf2 and
MAPK. J Hazard Mater 2020;394:122549; doi: 10.1016/j.jhazmat.2020.122549.
Zhang J, Yao K, Yin J, et al. Exposure to Bisphenolic Analogues in the Sixth Total Diet Study -
China, 2016-2019. China CDC Wkly 2022;4(9):180-184; doi: 10.46234/ccdcw2022.044.
Zhang Y, Marsboom G, Toth PT, et al. Mitochondrial respiration regulates adipogenic
differentiation of human mesenchymal stem cells. PLoS One 2013;8(10):e77077; doi:
10.1371/journal.pone.0077077.
Zimmerman JB, Anastas PT. Chemistry. Toward substitution with no regrets. Science
2015;347(6227):1198-1199; doi: 10.1126/science.aaa0812.

Table 1: Primer pairs

Gene		Sequence	Accession
Klf4	Forward	CATTAATGAGGCAGCCACCTG	NM_010637.3
	Reverse	GCTTCATGTGAGAGAGTTCCT	
Fos	Forward	CTTCGACCATGATGTTCTCG	NM_010234.3
	Reverse	CTGGGGAATGGTAGTAGGAA	
Cat	Forward	CTTTGACAGAGAGCGGATTC	NM_009804.2
	Reverse	CTTTGCCTTGGAGTATCTGG	
Sod3	Forward	ACCCAGAAATCTTTTCACGC	NM_011435.3
	Reverse	AGCAGACTCAAAGACTAGGG	
Gpx1	Forward	AGTCCACCGTGTATGCCTTC	NM_008160.6
	Reverse	CAGATCGTTCATCTCGGTGT	
Nr6α1	Forward	ATCATCCAGTAGGTCTGTGG	AF390896.1
	Reverse	GTCACAGCATACCCATCTTC	
β-actin	Forward	GATCAAGATCATTGCTCCTCCT	NM_007393.5
	Reverse	GTAACAGTCCGCCTAGAAGC	
C/ebpβ	Forward	TGATGCAATCCGGATCAAAC	NM_001287738.1
	Reverse	CCGCAGGAACATCTTTAAGT	
Pparγ	Forward	GCTGTTATGGGTGAAACTCT	EF062476.1
	Reverse	TGATGTCAAAGGAATGCGAG	
Glut4	Forward	GGTGGCATGATCTCTTCCTT	NM_001359114.1
	Reverse	CCTGATGTTAGCCCTGAGTA	
Scd1	Forward	CAAGAGTAGCTGAGCTTTGG	NM_009127.4
	Reverse	AATGCATCATTAACACCCCG	
Fasn	Forward	AGGGTACAAAGGTGTTCAGA	NM_007988.3
	Reverse	CTCATGACGAGTGCTGTAAG	

FIGURE LEGENDS

Figure 1: Summary Graphic Illustration. Reactive oxygen species (ROS) act as signaling

molecules that regulate the commitment and differentiation of adipocyte progenitors, a

developmental event that occurs during a discrete window in intrauterine life and establishes the

setpoint of adiposity. Bisphenol S (BPS) potentiates adipogenesis via stimulating an increased

production of ROS in part through inducing mitochondria dysfunction. Therefore, gestational

exposure to BPS and other structural analogues of BPA may predispose to the development of

later-life obesity.

Figure 2: BPS potentiates differentiation of adipocyte progenitors. Representative images of

lipid droplet staining by Oil Red-O or BODIPY/DAPI on day 7 of differentiation in adipose-

derived progenitors exposed to BPS or vehicle control (A). Lipid droplet formation quantified by

measuring the optical density (490 nm) of eluted Oil Red-O dye from stained progenitors on day

2, 4 & 7 of differentiation after BPS exposure (B). Relative mRNA expression of adipogenic

markers on day 2 & 4 of differentiation (C). Representative blots (D) and quantification (E) by

densitometry of Western blots for determination of protein expression of adipogenic markers on

day 2 & 4 of differentiation. Each experiment was conduced in cells isolated from 2-3 animals and

triplicate absorbance readings from each well are averaged. Differences are compared by One-way

ANOVA with Dunnett's multiple comparison test; * $p < 0.05$ BPS vs. vehicle.

Figure 3: Prior to differentiation, pathways involved in adipocyte differentiation and

oxidative stress are modulated in adipose-derived stem cells exposed to BPS. Heat map

showing DEGs involved in adipogenesis and Nrf2-mediated responses to oxidative stress, in

vehicle vs. BPS-exposed undifferentiated progenitors (A). Bubble plot showing number of

upregulated and downregulated genes in the top 10 enriched pathways in MSigDB Hallmark

analysis (B). IPA showing integration of DEGs involved in pathways regulating adipogenesis and

NRF2-mediated responses to oxidative stress (C); Green = downregulated genes; Orange =

upregulated genes. PCR validation of differences in gene expression with > 1fold change between

vehicle and BPS-exposed progenitors (D). RNAseq analysis included n = 5 samples/group. * p <

0.05 BPS vs. vehicle.

Figure 4: BPS and other bisphenol analogues increase production of ROS. Production of O_2^-

was detected by measuring fluorescence of DCFDA staining by the plate reader method in

undifferentiated cells acutely exposed to various concentrations of BPS (A), BPA (B) or BPF (C).

Additionally, DCFDA staining was quantified by flow cytometry in progenitors exposed to BPS

(D) and representative histograms are shown (E). Oxidative stress was detected in cells exposed

to BPS for 48 hrs by CellROX staining. Representative images of CellROX-stained cells with

DAPI nuclear staining are shown (F), and CellROX fluorescence was quantified by spectroscopy

(G). Oxidative stress measured by quantification of CellROX staining was also detected in stem

cells exposed to BPA (H) or BPF (I). Each experiment was conducted in cells isolated from 2-3

animals and triplicate absorbance readings from each well are averaged. Differences are compared

by One-way ANOVA with Dunnett's multiple comparison test. * p < 0.05 BPS vs. vehicle.

Figure 5: ROS regulate adipogenesis and mediate BPS-induced potentiation of adipogenesis.

Lipid droplet formation assessed by measuring optical density (490 nm) of eluted Oil Red-O dye

in progenitors exposed to the O_2^- generator, menadione, on day 4 of differentiation (A). Relative

mRNA expression of adipogenic markers on day 2 of differentiation after exposure to menadione

(B). Quantification of lipid droplet staining by Oil Red-O on day 4 of differentiation of progenitors

treated with the O_{2-} generator, DMNQ (C). Relative mRNA expression of adipogenic markers on

day 2 of differentiation after exposure to DMNQ (D). Quantification of Oil Red-O staining in

progenitors exposed to various doses of the ROS scavenger, Tempol, in the absence of BPS (E).

Absorbance of Oil Red-O staining in vehicle or BPS-exposed progenitors in the presence or

absence of Tempol (F) or the NADPH oxidase inhibitor and antioxidant, apocynin (G).

Representative BODIPY-stained differentiated progenitors exposed to vehicle or BPS in the

presence or absence of Tempol (H). Each experiment was conducted in cells isolated form 2-3

animals and triplicate absorbance readings from each well were averaged. Differences were

assessed by One-way ANOVA with Dunnett's multiple comparison test to compared BPS vs.

vehicle (* $p < 0.05$) or One-way ANOVA with Tukey's post hoc test to compare vehicle or BPS

with inhibitor vs. without inhibitor (ǂ $p < 0.05$ vs. without inhibitor).

Figure 6: Mitochondria are important sources of ROS in BPS-exposed progenitors. Degree

of depolarization calculated as the ratio of JC-1 monomer (green) to aggregate (red) fluorescent

signal in undifferentiated progenitors exposed to BPS for 48 hrs (A). Flow cytometric

quantification of TMRE fluorescence indicating mitochondria membrane polarization in

progenitors exposed to BPS or FCCP, an uncoupler that induces mitochondria membrane

depolarization (B). Representative images of mitochondria stained with MitoTracker Green (C).

Quantification of mitochondria-derived O_2^- by plate reader (D) or flow cytometric (E)

quantification of MitoSOX staining in undifferentiated stem cells exposed to BPS for 24 hrs.

Representative histograms of progenitors stained positive for MitoSOX (F). Quantification of Oil

Red-O staining after differentiation of progenitors exposed to BPS or vehicle in the presence or

absence of mitochondria-targeted antioxidants, MitoQ (G) or SS31 (H). Representative BODIPY

staining of lipid droplets in progenitors differentiated after exposure to BPS in the presence or

absence of MitoQ (I). Each experiment was performed in cells isolated from 2-3 animals and

absorbance of triplicates from each well were averaged. Differences were assessed by One-way

ANOVA with Dunnett's multiple comparison test to compared BPS vs. vehicle (* $p < 0.05$) or

One-way ANOVA with Tukey's post hoc test to compare vehicle or BPS with inhibitor vs. without

inhibitor (⧧ $p < 0.05$ vs. without inhibitor).

Figure 7: Intrauterine exposure to BPS programs an increase in later-life adiposity. In

subcutaneous adipose depots, commitment of mesenchymal stem cells to the adipocyte lineage

occurs prior to birth, followed by rapid expansion of fat accumulation in the early postnatal period

and during a second wave in puberty, with adipocytes numbers stabilizing thereafter (A). In

C57BL6 mice, 2 µg/kg body weight BPS was administered to pregnant dams (prenatal exposure)

starting on gestational day 0 (Gd0) and continuing until the pups weaned on postnatal day 21

(Pd21) or postnatal exposure to BPS was initiated on Pd21 (B). At 12 weeks of age prenatally

exposed mice (C) or postnatally exposed mice (D) were assessed for whole-body fat mass content

using TD-NMR. Fat mass of litter mates in prenatally-exposed mice is averaged for each sample

(n = 7-11 litters). * $p < 0.05$ vs. control.

Singh *et al*.

Supplementary Figure 1: Cell viability is not compromised over the range of BPS

concentrations that promote adipogenesis. Quantification of MTT, as an indicator of cell

viability, in undifferentiated progenitors exposed to various concentrations of BPS for 24 hrs (A)

or 48 hrs (B). Differences compared by One-way ANOVA with Dunnett's post hoc test. * $p < 0.05$

BPS vs. vehicle.

Supplementary Figure 2: Enhanced lipid droplet formation in stem cells exposed to BPA or

BPF is attenuated with a mitochondria-specific antioxidant. Quantification of eluted Oil-Red

O dye in differentiated progenitors after exposure to various concentrations of BPA (A) or BPF

(B). Lipid droplet formation was measured in presence or absence of MitoQ to assess the role of

mitochondria-derived ROS in mediating the effect of BPA (C) or BPF (D) on adipogenesis.

Differences were assessed by One-way ANOVA with Dunnett's multiple comparison test to

compared BPS vs. vehicle (* $p < 0.05$) or One-way ANOVA with Tukey's post hoc test to compare

vehicle or BPS with inhibitor vs. without inhibitor (‡ $p < 0.05$ vs. without inhibitor).

BPS

ROS

Committment
Differentiation

Establishment of the adipocyte progenitor pool

A

B

C

D