Functional characterization of the Co^{2+} transporter AitP in *Sinorhizobium meliloti*: a

new player in Fe^{2+} homeostasis

RUNNING TITLE: AitP Fe-export orchestrates the [Fe-S] salvage pathway

Paula Mihelj[1], Isidro Abreu[2*], Tomás Moreyra[1], Manuel González-Guerrero[2] [3], Daniel

Raimunda[1]

[1]Instituto de Investigación Médica Mercedes y Martín Ferreyra - INIMEC - CONICET, UNC.

Córdoba, Argentina. [2]Centro de Biotecnología y Genómica de Plantas (UPM-INIA/CSIC),

Universidad Politécnica de Madrid, Madrid, Spain. [3]Departamento de Biotecnología-

Biología Vegetal. Escuela Técnica Superior de Ingeniería Agronómica, Alimentaria y de

Biosistemas, Universidad Politécnica de Madrid, Madrid, Spain. *Current address:

Department of Biology, University of Oxford, Oxford, UK.

Abstract

Co^{2+} induces the increase of the labile-Fe pool (LIP) by Fe-S cluster damage, heme synthesis inhibition and "free" iron import, which affects cell viability. The N_2-fixing bacteria, *Sinorhizobium meliloti,* is a suitable model to determine the roles of Co^{2+}-transporting Cation diffusion facilitator exporters (Co-eCDF) in Fe^{2+} homeostasis because it has a putative member of this sub-family, AitP, and two specific Fe^{2+}-export systems. An insertional mutant of AitP showed Co^{2+} sensitivity and accumulation, Fe accumulation and hydrogen peroxide sensitivity, but not Fe^{2+} sensitivity, despite AitP being a *bona fide* low affinity Fe^{2+} exporter as demonstrated by the kinetic analyses of Fe^{2+} uptake into everted membrane vesicles. Suggesting concomitant Fe^{2+}-dependent induced stress, Co^{2+} sensitivity was increased in strains carrying mutations in AitP and Fe^{2+} exporters which did not correlate with the Co^{2+} accumulation. Growth in the presence of sub-lethal Fe^{2+} and Co^{2+} concentrations suggested that free Fe-import might contribute to Co^{2+} toxicity. Supporting this, Co^{2+} induced transcription of Fe-import system and genes associated with Fe homeostasis. Analyses of total protoporphyrin content indicates Fe-S cluster attack as the major source for LIP. AitP-mediated Fe^{2+}-export is likely counterbalanced via a non-futile Fe^{2+}-import pathway. Two lines of evidence support this: i) an increased hemin uptake in presence of Co^{2+} was observed in WT *vs.* AitP mutant, and ii) hemin reversed the Co^{2+} sensitivity in the AitP mutant. Thus, the simultaneous detoxification mediated by AitP aids cells to orchestrate an Fe-S cluster salvage response, avoiding the increase in the LIP caused by the disassembly of Fe-S clusters or free iron uptake.

Importance

Cross-talk between iron and cobalt has been long recognized in biological systems. This is

due to the capacity of cobalt to interfere with proper iron utilization. Cells can detoxify cobalt

by exporting mechanisms involving membrane proteins known as exporters. Highlighting

the cross-talk, the capacity of several cobalt exporters to also export iron is emerging.

Although biologically less important than Fe^{2+}, Co^{2+} induces toxicity by promoting

intracellular Fe release, which ultimately causes additional toxic effects. In this work, we

describe how the N_2-fixating rhizobial cells solve this perturbation by clearing Fe through a

Co^{2+}-exporter, in order to reestablish intracellular Fe-levels by importing non-free Fe, heme.

This piggyback-ride type of transport may aid bacterial cells to survive in free-living

conditions where high anthropogenic Co^{2+} content may be encountered.

Introduction

Cobalt (Co) and iron (Fe) are essential micronutrients in bacteria. While Fe is broadly required to deal with biological reactions in the actual conditions ruled by dioxygen, Co is involved in a handful of metabolic pathways as a catalytic cofactor or as the coenzyme vitamin B_{12} [1-3]. The chemical properties of both Fe and Co mean that they become toxic under certain conditions that promote increased bioavailability, i.e., high concentration in the extracellular environment, or in the case of Fe, an increase in the exchange between intracellular storage and the labile pool (*en route exchange*) [4, 5]. Although Fe toxicity seems to be curbed by Fe-storage systems, recent studies have suggested that Fe accumulation may exceed their buffering capacity and lead to redox stress [6, 7].

The rhizobial α-proteobacteria and alfalfa symbiont, *Sinorhizobium melliloti*, is involved in biological N_2-fixation, a process highly dependent on Fe [8]. Previous studies describing genetic and mechanistic findings on Fe acquisition, export and storage in α-protobacteria, including rhizobia [9], share one of the two transcriptional factors, *irr* [10] and *rirA* [11], which sense cellular Fe status through heme or Fe-S clusters, respectively. RirA regulates iron uptake gene transcription, promoting free (ferric) and non-free (siderophore-iron complexes or heme) iron uptake in low Fe conditions. Irr regulates genes involved in heme synthesis and Fe exporters, derepressing their transcription in high Fe conditions. In *Agrobacterium tumefaciens* and *Bradyrhizobium japonicum*, the membrane-bound ferritin A (MbfA), a member of the CCC1 Fe efflux family, has been described to be involved in Fe homeostasis [12, 13] . In *S. meliloti*, a P-type-IB-ATPase (Nia) has also been described as being specific for Fe transport and upregulated under low O_2 conditions [14]. Cobalt is an essential nutrient for *S. meliloti* and other rhizobia [15], and an uptake system has been identified in this organism [16]. Similarly, the *dmeRF* system, consisting of a member of the FrmR family of transcriptional regulators (DmeR) and a member of the cation diffusion facilitator (DmeF),

has been involved in Co^{2+} homeostasis through export, specifically during free-living stages

previous to establishing symbiosis and nodulation [17-19].

A feature of Co^{2+} dyshomeostasis is related to Fe-S cluster stability. There is evidence that

a high Co^{2+} condition induces Fe-S cluster disassembly, triggering *suf* and *isc* machinery in

response to this injury [20, 21]. The mechanisms proposed leading to Fe-S disruption by

Co^{2+} or other transition metals vary from the non-cognate metal interfering with assembly

process [20], to the Fe-S centers attacked in enzymes [22]. However, the final outcome in

order to supply Fe for cluster reconstitution relies on changes in the dynamic between

intracellular Fe stores and the labile-Fe pool (LIP) [23], and most importantly, the triggering

of free and non-free Fe uptake mechanisms [18, 21, 24].

In rhizobia, the Fe-S cluster synthesis pathway is regulated by the transcription factor RirA,

which represses the *suf* operon, the unique system for Fe-S cluster assembly in *S. meliloti*

[9], under replete Fe conditions [11]. RirA is a functional analog of Fur, the canonical Fe-

responsive transcriptional regulator, but differs from this in that it senses Fe indirectly

through an Fe-S cluster with a labile Fe as a proxy [25, 26]. Thus, under low iron conditions,

the release of RirA from the promoter region (IRO box) activates the transcription of the

genes in its regulon. *S. meliloti* presents a second transcriptional regulator, Irr, which senses

Fe status through heme binding and regulates *rirA* transcription, reinforcing the role of *rirA*

under low iron conditions [4].

Some authors have proposed a link between Co^{2+} and Fe^{2+} homeostasis, providing evidence

of a RirA-mediated response upon high extracellular Co^{2+}, involving upregulation of heme

uptake systems, probably as a result of an increase in the inactive apo form [18]. As a Co^{2+}

Fe-S cluster attack could lead to an increase in Fe^{2+} bioavailability [27-29], an Irr-mediated

response is also expected.

Importantly, previously described Co^{2+} transporters are emerging as Fe-exporters in

bacterial organisms [7, 30, 31]. Therefore, we decided to evaluate the Fe transport capacity

of SMc04167/AitP, a DmeF homolog in *S. meliloti*. Our data, showing that Co^{2+} induces Fe

dyshomeostasis by likely affecting dynamics in the LIP, and that Co^{2+} sensitivity can be

rescued in a non-competitive fashion by heme, together with the observations that AitP

exports Fe^{2+}, indicates a role for the FrmR/DmeR-regulated Co^{2+} exporters of the CDF family

at initial stages of the Co^{2+} induced mediated Fe-S cluster salvage pathway through Fe^{2+}

export.

Results

AitP is a classical CDF Co^{2+} exporter member of a sub-family with non-unique operon

architectures

A survey of more than 17,000 complete bacterial genomes allowed us to identify by Best-

Reciprocal-Hit more than 759 *dmeF/aitP*-like genes (Fig. 1A), showing typical features of

this Co^{2+} CDF subfamily. A multigene blast analysis showed that approximately 62% of

these share the archetypical operon organization of juxtaposed *frmR-aitP/dmeR-dmeF* loci

(Fig. 1A). The remaining percentage appears to have an FrmR-independent regulation with

no archetypical operon configuration and no operator-promoter region for the Co^{2+}

responsive transcriptional regulator FrmR/DmeR (Suppl. Table 1). These findings also

suggest that a significant number of putative Co^{2+} exporter *aitP*-like genes might be part of

Co^{2+}-responsive operons, with effectors other than Co^{2+} generated when this is

accumulated. Moreover, AitP-like members found in Co^{2+}-unresponsive operons could have

a broaden specificity of transport.

The genome of *S. meliloti* encodes a single member of the Co^{2+} efflux CDF subfamily

(SMc04167, AitP), a homolog of the DmeF protein described in *R. leguminosarum* [17], and

also of the AitP transporter of *Pseudomonas aeruginosa* [7]. As in other rhizobia, AitP is

preceded by a transcriptional regulator of the FrmR/DmeR family. Phenotypic analysis of the insertional mutant of the SMc04167 locus in *S. meliloti* Rm1021 (Ω04167 or "AitP mutant") shows increased Co^{2+} sensitivity in this strain (Fig. 1B). The sensitivity was concentration-dependent and it was rescued after gene complementation (Fig. 1C-D). Co^{2+} and Fe^{2+} accumulation occurred in the Ω04167 strain after incubation for 1 hour with 0.2 mM Co^{2+} or 20 minutes with 1.5 mM Fe^{2+},and complementation of Ω04167 reverted the accumulation to WT levels (Fig. 1E). Cell numbers after incubation were the same in all strains, indicating that these treatments (time and Co^{2+} or Fe^{2+} concentrations) are sub-lethal (not shown). Although the Fe^{2+} accumulation in AitP mutant did not induce a defect in cell growth, the double mutant lacking AitP and MbfA transporters was more sensitive to hydrogen peroxide (H_2O_2) than the MbfA single mutant in growth media with no external Fe added (Fig.1F). Free Fe^{2+} catalyzes the formation of reactive oxygen species through a Fenton reaction, which indicates that AitP functions as an Fe^{2+} exporter. As expected, transcription of *aitP* and *frmR*/*dmeR* was upregulated by Co^{2+}(Fig. 1G) but was unresponsive to Fe^{2+} in the case of *aitP* as opposed to *mbfA* (Fig. 1H).

AitP shows a dual specificity of transport

The phenotypical findings (Fig. 1E-F) agree with previous studies on AitP-like members [7], but suggest an alternative role in Fe^{2+} export by AitP. As Fe^{2+} homeostasis plays a significant role for the rhizobia-legume symbiosis, we decided to test whether Fe^{2+} could be transported by AitP. This could be hindered by the apparently redundant *mbfA* and *nia* Fe^{2+} transport systems [12-14]. Therefore, we heterologously overexpressed AitP in the Δ*fieF E. coli* strain (Suppl. Fig. 1A). FieF was previously described as an Fe^{2+} transporter [31, 32], and thus the genotypic background lacking FieF would allow measurement of Fe^{2+} with a low background signal. After protein induction, everted membrane vesicles were obtained and energized by pre-incubation with the electron donor NADH (or lactate), generating a H^+-electrochemical

gradient (Suppl. Fig. 1B). Only in this condition was $^{59}Fe^{2+}$-importing activity to the vesicle

detected, indicating that Fe^{2+} is a substrate transported by AitP, and that the driving force is

the electrochemical gradient of H^+ (Fig. 2A). After initial rate conditions were determined, we

characterized the kinetics of the transport (Fig 2B). The estimated $K_{1/2}$ value at the low

millimolar range (1.35±0.21) may indicate a low affinity for transport, in line with the iron-

saving strategies observed in almost all living organisms toward intracellular Fe^{2+}. The H^+-

dependent transport at sub-saturating concentrations (Fe^{2+} = 2mM) was inhibited only by

Co^{2+}, indicating that competition at the transport site is possible (Fig. 2C). However, Co^{2+}

accumulation observed in the AitP mutant (Fig. 1E) suggests a dual (Fe/Co) transport

specificity, with both ions as cytosolic substrates, likely exported by a sequential mechanism

involving binding to the same transport site. Considering the absence of Fe^{2+} regulation in

aitP transcription, these results suggest a role for AitP in Fe^{2+} transport in conditions in which

concomitant intracellular Co^{2+} accumulation occurs.

AitP participates in Fe-homeostasis

Previous phenotypical characterizations of *aitP*-like genes in other rhizobial organisms point

to a similar role of AitP in Co^{2+} export and detoxification [17-19], and we have previously

shown that the AitP homolog in *P. aeruginosa* is involved in Fe^{2+} homeostasis [7]. However,

a P-IB-type ATPase, *nia*, has been described as participating in Fe^{2+} homeostasis in this

organism [14], and the identification of an *mbfA* ortholog in Rm1021 genome led us to

hypothesize that AitP in *S. meliloti* may play a non-classical role in Fe^{2+} homeostasis. In this

line, Fe^{2+} sensitivity was not observed in the Ω04167 strain, pointing to *nia* and *mbfA* as

major exit routes for Fe^{2+} in rhizobia. As Co^{2+} transport has been described through Fe^{2+}

exporters, we conducted phenotypical analyses to test whether double mutants showed

increased sensitivity towards Co^{2+}. We observed an increased sensitivity in the double

mutants, Ω3004 and Ω03808, beyond the threshold shown by the Ω04167 strain (Fig. 3A)

but, interestingly, no significant increases of Co^{2+} accumulation were observed *vs.* the

Ω04167 strain (Fig. 3B). Supporting the exclusive participation of AitP in Co^{2+} efflux, Co^{2+}-

dependent *nia* transcription is not expected [14], and *mbfA* was not induced by Co^{2+} (Fig.

1G). Thus, we hypothesized a possible Fe^{2+}-induced toxic effect being added to those

promoted by Co^{2+} accumulation. To test this, we measured the Co^{2+} sensitivity of the

Ω04167 *vs.* Rm1021 in the presence of Fe. The increased sensitivity shown by Ω04167 in

this condition (Fig. 3C) indicates that Co^{2+} accumulation induces a shift in iron distribution,

with increased LIP or higher levels of Fe^{2+} bioavailability that result toxic to the cell. As the

LIP can be fed with intracellular and extracellular Fe sources, we compared the transcription

of *fbpA,* the substrate binding protein of the ferric cation ABC transporter, in WT and AitP

mutant exposed to sub-lethal Co^{2+} concentrations, as a proxy of extracellular Fe

contribution. We detected a significant increase of *fbpA* in both strains (Fig. 3D), with higher

levels in the AitP mutant, indicating that Fe import and probably intracellular Fe contribute

to Co^{2+} toxicity. Considering that the Co^{2+} sensitivity of double mutants was observed in

media with no added Fe (ca. 10 µM total Fe), we tested the effect of Co^{2+} on pathways

whose inhibition could lead to an increase in the bioavailability of iron. As Co^{2+} inhibits heme

synthesis and induces Fe-S disassembly, thus promoting the increase of LIP, we quantified

the total protoporphyrin content in both strains, exposed or not to sub-lethal Co^{2+} (Fig. 3E)

and measured the transcription of *sufA* in WT, a gene upregulated in response to Fe-S

cluster attack. Although Co^{2+} treatment decreases the level of protoporphyrin, this is similar

in both strains, suggesting that the Co^{2+}-mediated Fe-S cluster attack is the main

contributing source to LIP. Regarding this, we observed *sufA* up-regulation in WT in the

presence of sub-lethal Co^{2+} concentrations (Fig. 3F). We also analyzed the Co^{2+}-

dependency of transcriptional levels of bacterioferritin (*bfr*) and bacterioferrodoxin (*bfd*) [33],

as an estimate of the Fe storage status and the RirA-mediated response. Only *bfd*

transcription was increased by Co^{2+} (Fig. 3F). Finally, Fe-S cluster or heme synthesis

disruption would lead to increased Fe^{2+} bioavailability, and a suitable proxy to test this are

the *mbfA* and *bfr* transcriptional levels, but increases were not statistically significant (Fig.

1G and 3F).

AitP Fe^{2+} transport likely orchestrates the initial stages of the Co^{2+}-induced Fe-S cluster

salvage pathway

As Co^{2+}-dependent Fe^{2+} toxic effects seem mainly related to Fe-S clusters, the AitP function

may be related to driving a concerted RirA-dependent Fe-S cluster salvage/synthesis

pathway. If Co^{2+} accumulation led to Fe-S disassembly, we hypothesized that genes

regulated by RirA might be dysregulated, similarly to that observed for *fbpA* and *sufA* (Fig

3D and F). In particular, SMc02726 or *shmR*, a hemin-binding outer membrane receptor

[34], is induced in low Fe conditions and in a Δ*rirA* mutant [35]. We detected increased

transcripts of the outer membrane porin, ShmR, and the periplasmic component, HmuT, of

the ABC-heme importer HmuTUV [36](Fig. 4A). This upregulation of the heme uptake

system in the Ω04167 mutant suggested an exacerbated response to Co^{2+} accumulation,

likely due to an induced shift in the equilibrium of RirA towards the apo-form. We then tested

if adding hemin in the media affected the Co^{2+} sensitive phenotype detected in the Ω04167

strain. As shown in Fig. 4B, hemin addition to buffered TY medium (pH = 6,5) with 0.2 mM

Co^{2+} rescued the Co^{2+}-sensitive phenotype at concentrations between 50-200 µM. Chelation

of Co^{2+} by Fe-protoporphyrin IX (hemin) or metal exchange was discarded, as empty

protoporphyrin IX (PPIX) at 200 µM did not decrease Co^{2+} sensitivity (Fig. 4C). The detection

of increased transcripts of hemin transport systems in Co^{2+} treated cells led us to test the

hemin uptake rate under the same treatment. Surprisingly, the AitP mutant did accumulate

hemin at a significantly lower rate *vs.* WT (Fig. 4D and E). Although counterintuitive, this

result indicates that an Fe/Co-dependent post-transcriptional regulation is possible. Overall,

the data point to a role for AitP in Fe export under Co-rich conditions in order to concert a

RirA-dependent heme uptake process likely used for Fe-S cluster assembly.

Discussion

Cobalt is an essential element for life, and organisms like *S. meliloti* have a special

requirement for Co for methionine synthesis [15, 16]. However, in high concentrations it

proves toxic to cells. Co^{2+} efflux transporters of the CDF family have been previously

characterized in rhizobia, and their transcriptional regulation is mainly governed by a

FrmR/DmeR transcription factor [17]. Our bioinformatics analyses show that a significant

number of members of this family may be regulated by different factors, thus responding to

other effectors. Interestingly, several of these members appear under the regulation of the

Multiple Antibiotic Resistance Regulator (MarR) superfamily of transcription factors.

Although analysis of the functional roles of non-FrmR-regulated Co^{2+}-CDF is beyond the

scope of this work, the heterogeneity in *aitP*-like homolog regulation points to their capacity

to transport other divalent cations. Recent works have expanded the substrate of the Co^{2+}

efflux CDF transporters [7], and in this work, we have unequivocally confirmed the Fe^{2+}

transporting capacity of AitP by $^{59}Fe^{2+}$ transport assays in energized everted vesicles. Due

to the sensitivity of the mutant to high Co^{2+} but not to Fe^{2+}, from a physiological perspective

two questions arise: why does AitP export iron? And why is it regulated by Co^{2+}?

The main toxic effect of Co^{2+} in bacteria seems related to Fe-S cluster assembly [20]. In *E.*

coli, the responses to the intoxication involve the induction of the Fe-S cluster salvage

pathway in order to counterbalance this effect, but also, downregulation of free-Fe import

systems, such as Feo [23]. In contrast, *S. meliloti*, and likely all α-proteobacteria of the

Rhizobiaceae clade bearing RirA homologs, handle Co^{2+} toxicity by inducing iron uptake

mechanisms, i.e., triggering an Fe-starvation-like response. In *Agrobacterium tumefaciens*,

RirA represses genes related to free and non-free Fe uptake, and the repression is lost in

the presence of Co^{2+} [18]. Our results also point to an induction of the expression of these

iron uptake systems by Co^{2+} (*fbpA*, *shmR* and *hmuT*; Fig 3D and 4A). Similarly, this is

prompted by a derepression involving Co^{2+} attack on holo-RirA.

It has been proposed that the Fe-S cluster attack might lead to an increase of the labile-Fe

pool (LIP). In the presence of Co^{2+} intoxication and accumulation, the contribution of Fe from

the Fe-S cluster to the LIP may be detrimental by oxidative damage. We detected *suf* up-

regulation induced by Co^{2+}, which is in line with the increase. This could be an important Fe

source in low iron conditions, likely contributing to the increase in sensitivity observed in

double mutants lacking the main Co^{2+} efflux system and one of the Fe^{2+} exporting systems

(Fig 3A). As AitP-like members are also found in δ-proteobacteria, which mostly regulate Fe

homeostasis via direct sensing by Fur, Fe^{2+} export activity may be required to promote apo-

Fur and the Fe-starvation response through non-free Fe import.

Thus, AitP activity counterbalances the increase of LIP coming both from the release of Fe

from Fe-S clusters disassembled by Co^{2+} and from the Fbp-mediated entry. The latter,

although it represents an apparently futile Fe import-export cycle, is promoted only by Co^{2+}

accumulation. In agreement with our hypothesis, the addition of free iron increased Co^{2+}

toxicity in the AitP mutant (Fig. 3C), while the non-free iron from hemin had a protective role

(Fig. 4B,C), likely allowing replenishment of the intracellular pool without incurring in stress

for the bacteria.

Two more plausible, but still speculative mechanisms, involve RirA and the regulatory

protein, *hmuP*/*hemp* [37]. Direct attack on RirA by Co^{2+} explains the transcriptional

derepression observed of Fe-S cluster salvage (Fig. 3F) and hemin uptake genes (Fig 4A),

but still leaves the export role of Fe for AitP dispensable. A role in the fine tuning of the Co^{2+}

toxic response involving Fe-export by AitP might be explained considering the post-

transcriptional regulation of *shmR* in rhizobia [38]. A unique putative binding site for the

HmuP protein (a regulatory protein of *shmR*) was found in 5´UTR of *shmR* transcripts.

Importantly, heme binding to HmuP has been shown *in vitro* [39]. We speculate that *in vivo*

HmuP sensing capacity, likely via free-Fe or PPIX binding, may be involved in the post-

transcriptional regulation of *shmR*. Thus, a LIP increase in the AitP mutant would

downregulate *shmR* expression. Our data showing lower hemin uptake capacity of AitP

mutant *vs*. WT, even when transcripts are elevated, support this. The recovery of Co^{2+}

resistance in the presence of high hemin concentrations (Fig. 4B) points to an unspecific

uptake, or passive diffusion, mechanism in the AitP mutant. These hypotheses are currently

being tested.

Based on the above, we propose a model during Co^{2+} dyshomeostasis in which AitP would

detoxify both Co^{2+} and Fe^{2+}. This dual export allows cells to reduce the primary stressor

(Co^{2+}), while maintaining the labile-Fe pool in range despite the release of free Fe from the

Fe-S clusters or the entry of free Fe via ABC transporter. The lack of heme import activity,

even when transcripts of the importing systems are high, suggests that AitP activity is

involved in the orchestration of the transcription-translation process of ShmR and Hmu.

Future studies will focus on the role of AitP-like members in the regulation of heme uptake

systems and the need for other partners to achieve this regulation.

Materials and methods

Bioinformatic analyses

To analyze the relative abundance of the canonical "FrmR-AitP" operon architecture, *S.*

meliloti AitP (SMc04167/ SM_RS10235) homologs were searched by the best reciprocal hit

method [40]. Briefly, a database was created locally with all the protein sequences from

bacteria with a complete genome (17,972) and chromosome (2,570) assembly level

obtained from ftp://ftp.ncbi.nih.gov/genomes/refseq/bacteria/assembly_summary.txt

(August 2021). A BLAST search in this database, with WP_010969625.1 (SMc04167/

SM_RS10235) as query and an Expect-value of 1.10^{-5}, resulted in 13,570 subject/hits.

Removal of duplicated subject/hits by accession number led to 4,431 SMc04167 homolog

candidates. These were blasted against *S. meliloti* 1021

(GCF_002197445.1_ASM219744v1_protein.faa) individually. In all cases BLAST retrieved

WP_010969625.1 as the subject, confirming best-reciprocal-hits association. After visual

inspection of the subject annotations, CDFs lacking the canonical poly-histidine stretch

sequence (TMS4 and TMS5) were excluded in order to minimize the number of arbitrarily

defined "true candidates" (Expect-value<$2.19.10^{-35}$ and percentage of positive-scoring

matches>54%). There were 1,651 assemblies of bacterial genomes having AitP-like

members with unique accession numbers. This number of organisms reflects several strain

assemblies of the same organism, and we maintained it in order to promote the sensitivity

of the final assessment. These were used later to compare and create a list of 759 non-

redundant bacterial organisms having, or not having, the FrmR-AitP architecture. The FrmR-

AitP architecture search was performed using Multigenblast [41]. For these analyses, the

genomic database was built with approximately 4,200 genebank-formatted (.gbbf)

assemblies of chromosome and complete genome with identical gender and species names

to those drawn from the 1,651 organisms within the AitP list. The SMc04168-SMc04167

sequence was used to run the analyses. Results were parsed to obtain the final list of

organism names with the operon architecture (~500) and compared to the list of the AitP-

possessing organisms (~800).

The remaining organisms with non-FrmR-AitP-like architecture (~300) were further

analyzed. Three search strategies were followed to confirm the lack of FrmR(RcnR/DmeR)-

dependent regulation in those organisms with AitP-like members lacking the FrmR-AitP

architecture. In the first, the 500bp up-stream CDS region was downloaded and used in a

tblastn search, using the protein sequence of SMc04168/*dmeR* as query. In the second,

protein accession numbers of AitP-like members of this group were used to retrieve the up-

and downstream loci annotations, using the E-fetch "identical protein group" tool. A third

strategy consisted in performing a blastn search in the 500bp upstream regions for the

consensus FrmR consensus binding motif, using the *dmeR* binding sequence up-stream

SMc04167 (AAATAGGGTACCCCCCTATGCT) as query. These searches allowed us to

identify 27 non-FrmR-AitP-like previous candidates as false negatives, and to confirm that

284 AitP-like loci lack the FrmR-AitP architecture. After removal of duplicated organism

names, a group diagram was prepared with these numbers.

Culture media, strains and mutant constructions

S. meliloti strains (Table 1) were cultured in TY media (0.5% tryptone, 0.3% yeast extract,

0.09% $CaCl_2$) supplemented with streptomycin 600 μg/ml. The *S. meliloti* strain

Rm1021(StrR) and *E. coli* S17-1 [42] carrying the plasmid pK19mob2ΩHMB for the mutation

of SMc04167 were obtained from University of Freiburg, Germany (Table 1). The deletion

mutant strain of gene SMc04167 (Ω04167) was generated by plasmid insertion following

described protocols [43]. Briefly, freshly grown *S. meliloti* Rm1021 was mixed with *E. coli* S-

17 cells bearing the plasmid pK19mobΩHMB, containing the internal sequence of the gene

SMc04167, named S17-1.PI.G1PELR20F8 (www.rhizogate.de (GenDB)), and then

incubated overnight at 30°C on a nitrocellulose filter disk placed on LB-agar.

Transconjugants were recovered from the filter and selected by plating on TY-agar

supplemented with streptomycin 600 μg/ml and kanamycin 50–200 μg/ml. Complementation

was achieved by conjugation of Ω04167 with the non-integrative plasmid pJB3Tc20-P$_{770}$-

aitP containing gene SMc04167 plus 770 bp upstream. Transconjugants were selected on

TY media with kanamycin (200 μg/ml) and tetracycline (10 μg/ml). Single and double

mutants of *nia* (SMa1163) and *mbfA* (SMc00359) were generated by conjugation and single

crossover of plasmid pJQ200SK, containing a 600 bp internal region of each loci in Rm1021

(single mutant) or Ω04167 (double mutants). Transconjugants were selected in TY agar

supplemented with streptomycin 600 µg/ml and gentamycin (50 µg/ml). Mutation of *fief* was

performed as already described[44]. Briefly, 50 bp up- and downstream were joined to the

Kan^R cassette from plasmid pKD4. After purification, the fragment was introduced by

electroporation (1.8 kV, 600 Ω and 10 µF, 1mm gap cuvette) into BL21(DE3) cells previously

transformed with pKD46. In order to induce the Rec system and to increase recombination,

the electrocompetent cells were prepared by growing cells in LB supplemented with 1mM L-

(+)-arabinose at 30°C. After electroporation, cells were recovered in 5 ml of SOC media.

Transformants were selected in LB-agar plates with 50 µg/ml kanamycin incubated at 37°C

for 16 hours. Curation of the Red system was confirmed in isolated colonies that showed

negative growth after 16 h incubation at 37°C in LB-agar plates supplemented with 100

µg/ml ampicillin and 50 µg/ml kanamycin, but positive growth in plates supplemented with

50 µg/ml kanamycin. All strains were checked by PCR using purified genomic DNA as a

template and internal and external primers (Table 1).

Sensitivity tests

Serial dilutions of *S. meliloti* Rm1021, Ω04167, Ω00359, Ω1163*,* Ω03808 and Ω3004 cells,

initially at OD$_{600nm}$=0.4, were spotted (10 µl) on TY-agar supplemented with the indicated

transition metals (as chloride salts, except Fe^{2+} as SO$_4^{2-}$)$^-$ or H$_2$O$_2$ concentrations for 48 h at

30°C. When hemin or PPIX (dissolved in 50 mM NaOH) were added to the media, the pH

was buffered with MES-KOH 15 mM pH 6.5, and the control conditions (0 mM hemin, 0 mM

PPIX) were supplemented with the same amounts of NaOH.

Cobalt and iron accumulation assays

Cells were incubated with 0.1 mM CoCl$_2$ for 1 hour or with 1.5 mM FeSO$_4$ in TY medium

during early exponential phase growth. Cells were pelleted and washed with 0.9% NaCl and

mineralized with HNO$_3$. Co or Fe were quantified by ICP-MS or Atomic absorption

spectroscopy.

Transcriptional analyses by qRT-PCR

Fold change expression levels ($2^{-\Delta\Delta Ct}$) of iron exporters and iron-related genes were assayed

under 20 µM, 100 µM or 200 µM $CoCl_2$, and low (0.75 mM 2,2'-bipyridyl (Bpy)), or high (0.75

mM $FeSO_4$) Fe^{2+} conditions *in vitro vs.* 16S reference gene. Cells were grown till the mid-

log phase at 30ºC in TY. After this, the culture volume was split, supplemented or not, and

incubated for 45 min at 30ºC before RNA extraction. Primers used are described in Table 1.

Iron transport assay

Fe^{2+} uptake was measured in energized everted membrane vesicles (EMV) obtained from

E. coli ΔfieF expressing AitP. Heterologous expression was detected by electroblotting SDS-

PAGE Tris-glycine gels onto nitrocellulose membranes and immunostaining with primary

rabbit anti-His-tag antibody and secondary goat anti-rabbit IgG phosphatase alkaline

conjugated antibody (GenScript, Piscataway, NJ, USA) (Suppl. Fig. 1). Vesicles were

obtained as previously described [45, 46] Before transport measurement, 10 µl vesicles (or

the equivalent to 100 µg total protein of EMV) were energized in 100 µl of transport buffer

(100 mM HEPES-Na, 250 mM sucrose, 50 mM KCl, 2 mM $MgCl_2$, 5 mM TCEP, pH 6.8) by

2-minute pre-incubation with 2.5 mM NADH at 37ºC and the uptake was initiated by addition

of a 10-fold concentrated $^{59}Fe^{2+}$ solution and mixing. Reaction was stopped after 5-minute

incubation at the same temperature by filtering through 0.22 µm nitrocellulose. Filters were

immediately washed with a standard solution containing 2 mM EDTA and dissolved in

scintillation cocktail for counting.

Hemin uptake assay

Hemin uptake was estimated following a previously described protocol [47]. Briefly,

stationary phase cells cultured in 50 ml of TY medium were incubated for 2 hours with 0.2mM

$CoCl_2$. Cells were centrifuged, washed once with ice-cold phosphate-buffered saline (PBS)

pH 7.45 and resuspended in 50 ml PBS 0.05 mM $CoCl_2$ supplemented with 30 ug/ml of a

hemin solution. At 2.5, 5, 15, 30, 60 and 120 minutes, one milliliter of culture was centrifuged

and the supernatant collected to measure the absorbance at 400 nm. No significant

difference was observed in the cell numbers of WT and AitP mutant strains after treatment,

indicating that the incubation time with Co^{2+} used does not affect cell viability.

Porphyrin quantification

Porphyrins were quantified following the described protocol [48]. Briefly, mid-log phase cells

cultured in 50 ml of TY medium were incubated for 2 hours with or without 0.2mM $CoCl_2$.

Cells were centrifuged, washed twice with ice-cold 50 mM Tris-HCl (pH 8.0), resuspended

in ethyl acetate/glacial acetic acid (3:1, v/v) to 1/100 the original culture volume, and lysed

by sonication on ice. Cell debris was removed by centrifugation at room temperature, and

the non-aqueous (top) phase was washed twice with 1 ml ddH_2O to remove residual water-

soluble contaminants. Extraction of porphyrins from the solution was achieved by the

addition of 0.5 ml 3 M HCl, and the absorbance of the bottom-aqueous phase was assessed

at 408 nm. Porphyrin levels were normalized to optical cell density (OD_{600nm}). No significant

difference was observed in the cell numbers of WT and AitP mutant strains after treatment,

indicating that the incubation time with Co^{2+} used does not affect cell viability.

Conflict of interest statement

All authors declare that research was conducted in the absence of any commercial or

financial relationships.

Author contributions

Conceived and designed the experiments: D.R., M.G.-G, I.A., P.M., T. M.

Analyzed the data: D.R., M.G.-G, I.A., P.M., T. M.

Contributed reagents/samples/analysis/tools: D.R., M.G.-G, I.A., P.M., T. M.

Wrote or edited the manuscript: D.R., M.G.-G, I.A., P.M., T. M.

All authors read and approved the final manuscript.

Funding

This work was supported by the Agencia Nacional de Promoción Científica y Tecnológica

(ANPCyT)(PICT2015-2897 and PICT2018-03139 to D.R.) and the Consejo Nacional de

Investigaciones   Científicas   y   Tecnológicas   (CONICET)   PUE-CONICET

22920160100135CO. D.R. is a CONICET investigator. P.M. is recipient of a CONICET

doctoral fellowship. T.M. is recipient of a doctoral fellowship from ANPCyT (PICT2018-

03139). I.A. was recipient of a Juan de la Cierva-Formación postodoctoral fellowship from

Ministerio de Ciencia, Innovación y Universidades (FJCI-2017-33222).

Abbreviations

TM, transition metal; CDF, cation diffusion facilitator; TMS, transmembrane segment;

LIP, labile-Fe pool; ICP-MS, inductively coupled plasma mass spectrometry; Bpy, 2,2'-

bipyridyl.

Acknowledgments

We thank Dr. J.M Argüello for help and support on the setting of $^{59}Fe^{2+}$ transport assays.

References

1.   Fraústo da Silva, J.J.R. and R.J.P. Williams, *The biological chemistry of the elements : the*
inorganic chemistry of life. 2nd ed. 2001, Oxford: Oxford University Press. xvii, 575 p.
2.   Kobayashi, M. and S. Shimizu, *Cobalt proteins.* Eur J Biochem, 1999. **261**(1): p. 1-9.
3.   Okamoto, S. and L.D. Eltis, *The biological occurrence and trafficking of cobalt.* Metallomics,
2011. **3**(10): p. 963-70.
4.   Bradley, J.M., et al., *Bacterial iron detoxification at the molecular level.* J Biol Chem, 2020.
295(51): p. 17602-17623.
5.   Pi, H. and J.D. Helmann, *Ferrous iron efflux systems in bacteria.* Metallomics, 2017. **9**(7): p.
840-851.
6.   Guan, G., et al., *PfeT, a P1B4 -type ATPase, effluxes ferrous iron and protects Bacillus*
subtilis against iron intoxication. Mol Microbiol, 2015. **98**(4): p. 787-803.
7.   Salusso, A. and D. Raimunda, *Defining the Roles of the Cation Diffusion Facilitators in*
Fe(2+)/Zn(2+) Homeostasis and Establishment of Their Participation in Virulence in
Pseudomonas aeruginosa. Front Cell Infect Microbiol, 2017. **7**: p. 84.
8.   Abreu, I., P. Mihelj, and D. Raimunda, *Transition metal transporters in rhizobia: tuning the*
inorganic micronutrient requirements to different living styles. Metallomics, 2019. **11**(4): p.
735-755.
9.   Rodionov, D.A., et al., *Computational reconstruction of iron- and manganese-responsive*
transcriptional networks in alpha-proteobacteria. PLoS Comput Biol, 2006. **2**(12): p. e163.
10. Hamza, I., et al., *The bacterial irr protein is required for coordination of heme biosynthesis*
with iron availability. J Biol Chem, 1998. **273**(34): p. 21669-74.
11. Todd, J.D., et al., *RirA, an iron-responsive regulator in the symbiotic bacterium Rhizobium*
leguminosarum. Microbiology (Reading), 2002. **148**(Pt 12): p. 4059-4071.
12. Bhubhanil, S., et al., *Roles of Agrobacterium tumefaciens membrane-bound ferritin (MbfA)*
in iron transport and resistance to iron under acidic conditions. Microbiology (Reading),
2014. **160**(Pt 5): p. 863-871.
13. Sankari, S. and M.R. O'Brian, *Synthetic Lethality of the bfr and mbfA Genes Reveals a*
Functional Relationship between Iron Storage and Iron Export in Managing Stress
Responses in Bradyrhizobium japonicum. PLoS One, 2016. **11**(6): p. e0157250.
14. Zielazinski, E.L., et al., *Sinorhizobium meliloti Nia is a P(1B-5)-ATPase expressed in the*
nodule during plant symbiosis and is involved in Ni and Fe transport. Metallomics, 2013.
5(12): p. 1614-1623.
15. Watson, R.J., et al., *Sinorhizobium meliloti cells require biotin and either cobalt or*
methionine for growth. Appl Environ Microbiol, 2001. **67**(8): p. 3767-70.
16. Cheng, J., et al., *An ABC-type cobalt transport system is essential for growth of*
Sinorhizobium meliloti at trace metal concentrations. J Bacteriol, 2011. **193**(17): p. 4405-
16.
17. Rubio-Sanz, L., et al., *Functional and expression analysis of the metal-inducible dmeRF*
system from Rhizobium leguminosarum bv. viciae. Appl Environ Microbiol, 2013. **79**(20): p.
6414-22.
18. Dokpikul, T., et al., *Regulation of the Cobalt/Nickel Efflux Operon dmeRF in Agrobacterium*
tumefaciens and a Link between the Iron-Sensing Regulator RirA and Cobalt/Nickel
Resistance. Appl Environ Microbiol, 2016. **82**(15): p. 4732-4742.
19. Li, Z., et al., *Nickel and cobalt resistance properties of Sinorhizobium meliloti isolated from*
Medicago lupulina growing in gold mine tailing. PeerJ, 2018. **6**: p. e5202.

20.   Ranquet, C., et al., *Cobalt stress in Escherichia coli. The effect on the iron-sulfur proteins.* J
479   Biol Chem, 2007. **282**(42): p. 30442-51.
480   21.   Thorgersen, M.P. and D.M. Downs, *Cobalt targets multiple metabolic processes in*
481   *Salmonella enterica.* J Bacteriol, 2007. **189**(21): p. 7774-81.
482   22.   Macomber, L. and J.A. Imlay, *The iron-sulfur clusters of dehydratases are primary*
483   *intracellular targets of copper toxicity.* Proc Natl Acad Sci U S A, 2009. **106**(20): p. 8344-9.
484   23.   Fantino, J.R., et al., *A genetic analysis of the response of Escherichia coli to cobalt stress.*
485   Environ Microbiol, 2010. **12**(10): p. 2846-57.
486   24.   Chillappagari, S., et al., *Copper stress affects iron homeostasis by destabilizing iron-sulfur*
487   *cluster formation in Bacillus subtilis.* J Bacteriol, 2010. **192**(10): p. 2512-24.
488   25.   Pellicer Martinez, M.T., et al., *Sensing iron availability via the fragile [4Fe-4S] cluster of the*
489   *bacterial transcriptional repressor RirA.* Chem Sci, 2017. **8**(12): p. 8451-8463.
490   26.   Martinez, M.T.P., et al., *Mechanisms of iron-and O2-sensing by the [4Fe-4S] cluster of the*
491   *global iron regulator RirA.* Elife, 2019. **8**: p. e47804.
492   27.   Imlay, J.A., *Pathways of oxidative damage.* Annu Rev Microbiol, 2003. **57**: p. 395-418.
493   28.   Latifi, A., M. Ruiz, and C.C. Zhang, *Oxidative stress in cyanobacteria.* FEMS Microbiol Rev,
494   2009. **33**(2): p. 258-78.
495   29.   Gao, F., *Iron-Sulfur Cluster Biogenesis and Iron Homeostasis in Cyanobacteria.* Front
496   Microbiol, 2020. **11**: p. 165.
497   30.   Patel, S.J., et al., *Fine-tuning of Substrate Affinity Leads to Alternative Roles of*
498   *Mycobacterium tuberculosis Fe2+-ATPases.* J Biol Chem, 2016. **291**(22): p. 11529-39.
499   31.   Brown, J.B., M.A. Lee, and A.T. Smith, *Ins and Outs: Recent Advancements in Membrane*
500   *Protein-Mediated Prokaryotic Ferrous Iron Transport.* Biochemistry, 2021. **60**(44): p. 3277-
501   3291.
502   32.   Grass, G., et al., *FieF (YiiP) from Escherichia coli mediates decreased cellular accumulation*
503   *of iron and relieves iron stress.* Arch Microbiol, 2005. **183**(1): p. 9-18.
504   33.   Costa, D., et al., *The Irr and RirA Proteins Participate in a Complex Regulatory Circuit and*
505   *Act in Concert To Modulate Bacterioferritin Expression in Ensifer meliloti 1021.* Appl
506   Environ Microbiol, 2017. **83**(16).
507   34.   Amarelle, V., M.R. O'Brian, and E. Fabiano, *ShmR is essential for utilization of heme as a*
508   *nutritional iron source in Sinorhizobium meliloti.* Appl Environ Microbiol, 2008. **74**(20): p.
509   6473-5.
510   35.   Chao, T.C., et al., *Role of the regulatory gene rirA in the transcriptional response of*
511   *Sinorhizobium meliloti to iron limitation.* Appl Environ Microbiol, 2005. **71**(10): p. 5969-82.
512   36.   Cuiv, P.O., et al., *The hmuUV genes of Sinorhizobium meliloti 2011 encode the permease*
513   *and ATPase components of an ABC transport system for the utilization of both haem and*
514   *the hydroxamate siderophores, ferrichrome and ferrioxamine B.* Mol Microbiol, 2008.
515   **70**(5): p. 1261-73.
516   37.   Amarelle, V., et al., *A new small regulatory protein, HmuP, modulates haemin acquisition*
517   *in Sinorhizobium meliloti.* Microbiology (Reading), 2010. **156**(Pt 6): p. 1873-1882.
518   38.   Amarelle, V., U. Koziol, and E. Fabiano, *Highly conserved nucleotide motifs present in the*
519   *5'UTR of the heme-receptor gene shmR are required for HmuP-dependent expression of*
520   *shmR in Ensifer meliloti.* Biometals, 2019. **32**(2): p. 273-291.
521   39.   Sato, T., et al., *The Small Protein HemP Is a Transcriptional Activator for the Hemin Uptake*
522   *Operon in Burkholderia multivorans ATCC 17616.* Appl Environ Microbiol, 2017. **83**(16).
523   40.   Hernandez-Montes, G., J.M. Arguello, and B. Valderrama, *Evolution and diversity of*
524   *periplasmic proteins involved in copper homeostasis in gamma proteobacteria.* BMC
525   Microbiol, 2012. **12**: p. 249.

41.   Medema, M.H., E. Takano, and R. Breitling, *Detecting sequence homology at the gene*
cluster level with MultiGeneBlast. Mol Biol Evol, 2013. **30**(5): p. 1218-23.
42.   Simon, R., U. Priefer, and A. Puhler, *A Broad Host Range Mobilization System for In Vivo*
Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. Nat. Biotech.,
1983. **1**(9): p. 784-791.
43.   Luo, L., et al., *Two new Sinorhizobium meliloti LysR-type transcriptional regulators required*
for nodulation. J Bacteriol, 2005. **187**(13): p. 4562-72.
44.   Datsenko, K.A. and B.L. Wanner, *One-step inactivation of chromosomal genes in*
Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A, 2000. **97**(12): p. 6640-5.
45.   Gonzalez-Guerrero, M., et al., *Distinct functional roles of homologous Cu+ efflux ATPases in*
Pseudomonas aeruginosa. Mol Microbiol, 2010. **78**(5): p. 1246-58.
46.   Raimunda, D., et al., *The transport mechanism of bacterial Cu+-ATPases: distinct efflux*
rates adapted to different function. Biometals, 2011. **24**(3): p. 467-75.
47.   Genco, C.A., B.M. Odusanya, and G. Brown, *Binding and accumulation of hemin in*
Porphyromonas gingivalis are induced by hemin. Infect Immun, 1994. **62**(7): p. 2885-92.
48.   Martin, J.E., et al., *The Escherichia coli small protein MntS and exporter MntP optimize the*
intracellular concentration of manganese. PLoS Genet, 2015. **11**(3): p. e1004977.
49.   Simon, R., U. Priefer, and A. Pühler, *A broad host range mobilization system for in vivo*
genetic engineering: transposon mutagenesis in gram negative bacteria. Bio/technology,
1983. **1**(9): p. 784-791.
50.   Quandt, J. and M.F. Hynes, *Versatile suicide vectors which allow direct selection for gene*
replacement in gram-negative bacteria. Gene, 1993. **127**(1): p. 15-21.
51.   Blatny, J.M., et al., *Construction and use of a versatile set of broad-host-range cloning and*
expression vectors based on the RK2 replicon. Appl Environ Microbiol, 1997. **63**(2): p. 370-
9.

Figures

Figure 1.eps

Figure 2.eps

Figure 3.eps

Figure 4.eps

Table/s

552   Table 1. List of strains, plasmids and primers used in this study.

Strain/plasmid/primer

Strains	Relevant characteristics	Reference
Sinorhizobium meliloti		
Rm1021	Wild-type (Str^R)	[43]
Ω04167	Rm1021 SMc04167::pK19mob2ΩHMB- *aitP* (Str^R) (Kan^R)	This work
Ω00359	Rm 1021 SMc00359::pJQ200SK- *mbfA* (Str^R) (Gm^R)	This work
Ω1163	Rm 1021 Sma1163SMa1163::pJQ200SK- *mbfA* (Str^R) (Gm^R)	This work
Ω03808	Rm 1021 SMc04167::pK19mob2ΩHMB- *aitP*; SMc00359::pJQ200SK- *mbfA*(Str^R) (Kan^R)(Gm^R)	This work
Ω3004	Rm 1021 SMc04167::pK19mob2ΩHMB- *aitP*; Sma1163SMa1163::pJQ200SK- *mbfA*(Str^R) (Kan^R)(Gm^R)	This work
C-Ω04167	Rm 1021 SMc04167::pK19mob2ΩHMB- *aitP* / pJB3Tc20-P$_{770}$-*aitP* (Str^R) (Kan^R) (Tc^R)	This work
Escherichia coli		
S17-1	*E. coli 294, thi RP4-2 – Tc: :Mu-Km: :Tn7* chromosomally integrated	[49]
DH5α	F- *Φ80lacZΔM15 Δ(lacZYA-argF) U169 recA1 endA1 hsdR17 (rK–, mK+) phoA supE44 λ– thi-1 gyrA96 relA1*	Invitrogen, Carlsbad, CA, USA
BL21(DE3)pLysS	F- *ompT hsdS*B (rB-mB-) *gal dcm* (DE3) *pLysS* (Cam^R)	Invitrogen, Carlsbad, CA, USA
BL21(DE3)pKD46	F- *ompT hsdS*B (rB-mB-) *gal dcm* (DE3) *pKD46* (Amp^R)	[44]
ΔfieF	BL21(DE3) *fieF::Km* (Kan^R)	This work

Plasmids		
pK19mob2ΩHMB	RP4 mobilizable plasmid, Kan^R	[43]
pK19mob2ΩHMB-*aitP*	SMc04167 insertion plasmid containing a 334 bp *aitP* fragment (AL591688.1:2155814-2156147)	www.rhizogate.de
pJQ200SK	RP4 mobilizable plasmid, Gm^R	[50]
pJQ200SK- *mbfA*	SMc00359 insertion plasmid containing a 568 bp *mbfA* fragment (AL591688.1:314759-315326)	This work
pJB3Tc20	Broad host-range RK2 cloning vector, Tc^R	[51]
pJB3Tc20-P$_{770}$-*aitP*	SMc04167 (*aitP*) containing a 770 bp promoter fragment cloned in pJB3Tc20	This work

Primers	Sequence (5'->3')	Use
qSMc02726-ShmR-F	AGCCCAACCCTACCGATTTC	
qSMc02726-ShmR-R	CCTGTTCCGCTCTGAGATCA	
qSMc01512-hmuT-F	GTCTTCGGCGTTGGGAAATG	
qSMc01512-hmuT-R	GGTAAATGCTGGTGGAATCGCG	
qSMc1021-16S-F	GATAAGCCGAGAGGAAGGTG	
qSMc1021-16S-R	GTGTAGCCCAGCCCGTAAG	

qSMc04167-aitP-F	CGCGCTTTACCTTCGGTACG	qRT-PCR
qSMc04167-aitP-R	AATGGCCTGCGGAAAACTGA	
qSMc04168-dmeR-F	GAAAAGTTGCTCGCCCGTGT	
qSMc04168-dmeR-R	GGAGGCGACGAGGTTGAGAA	
qSMc03787-bfd-F	GGACGCCGATATATTTGATTTC	
qSMc03787-bfd-R	CGTCGCCTTTCAAGATCC	
qSMc03788-bfr-F	CGATTACGTCTCGATGAAGC	
qSMc03788-bfr-R	CTGGCCGTATTTTTCCTCAC	
qSMc00359-mbfA-F	CCTCGGAAGAGGACGATGGA	
qSMc00359-mbfA-R	GCTCTCCTCCTCCGCCATTT	
qSMc00784-fbpA-F	AACTGGTTCGGTCTGACGAC	
qSMc00784-fbpA-R	AGGCCGACATTGTAGGAGTG	
qSMc00301-sufA-F	AATATGCCGTCGATCTCGTG	
qSMc00301-sufA-R	GTCTCGAAATCCATCTGCGT	
DM-F-Sma1163-BamHI	ACTGGGATCCGACAGCACCTATGCGGCA	Cloning of 600 bp and yield of double and single mutants in *S. meliloti*
DM-R-Sma1163-XhoI	CAGTCTCGAGGTCCGTATGTCTCGCGAG	
DM-F-Smc00359-BamHI	ACTGGGATCCCTGCATGTCGCCCGTTTCG	
DM-R-Smc00359-XhoI	CAGTCTCGAGCGAGTGCCGTCATGATGC	
BamHI_AitP_-770_FW	GGGGATCCCCTCCCAGGCTAAAGCACGT	Complementation in *trans*
PstI_aitP_Stop_RV	GGCTGCAGCTCAGGCCGCCAGCGGCTCG	
Delta-fieF-F	ATGAATCAATCTTATGGACGGCTGGTCAGTCGGGCGGCGATTGCTGCGACGTGTAGGCTGGAGCTGCTTC	Generating *E. coli* ΔfieF
Delta-fieF-R	TTATGAAAGCATAGACCGTTTACCCTCCCTGGGTACGACGGAACAGGGGTCATATGAATATCCTCCTTAG	
Smc04167-F	AGCGAAACCCATTCACC	Transconjugant confirmation
Smc04167-R	GCCGCCAGCGGCTCGACCTC	
Smc00359-F	ATGCTCTCCCGACTTTTCTCC	
Smc00359-R	AGCATTCCCTATGACGATGCC	
Sma1163-F	TTGCGGGATATTGCGCAG	
Sma1163-R	CTCGTGCTCGAGCGATC	
up-fieF-F	CCGAACATCAGCGTCAGCA	*E. coli* ΔfieF confirmation
K1-R	CAGTCATAGCCGAATAGCCT	

Figure legends

Figure 1. AitP is involved in Co^{2+}/Fe^{2+} homeostasis and the genetic context of bacterial

AitP-like transporters points to a broader transport specificity and regulation. A)

Operon architecture analysis of AitP-like transporters in bacterial sequenced organisms

indicates the existence of a subgroup with different transcriptional regulation. B) Metal

toxicity was evaluated by inhibition halo assay using *S. meliloti* Rm1021 and Ω04167. Bars

represent the inhibition halo diameter around 7-mm filter paper discs containing 10 (black),

5 (dark gray), 2 (gray) or 1 (light gray) μmole of the indicated metal (as Cl^- except for $FeSO_4$).

Notice the Co^{2+} sensitivity in Ω04167. * = p<0.01 *vs.* Rm1021. C) Serial dilutions (10 μl) of

cells, initially at OD_{600nm}=0.4, were spotted on TY-agar supplemented with the indicated Co^{2+}

(as Cl^-) concentration for 48 h at 30°C. D) Complementation was achieved by transformation

of Ω04167 with pJB3Tc20-P_{770}-*aitP* (C-Ω04167 strain). Notice that Co^{2+} sensitivity was

restored to WT levels in the this strain. E) Cobalt (dark gray) or iron (light gray) content in

cells after incubation with 0.2 mM $CoCl_2$ for 1 h or 1.5 mM $FeSO_4$ for 30 min at 30°C in TY

medium during early exponential phase growth. Data are the mean ± SE of three

independent experiments. * = *p*< 0.05 and # = *p*<0.01 from a two tail Student's t-test *vs.*

Rm1021. F) Serial dilutions (10 μl) of cells initially at OD_{600nm}=0.4 were spotted on TY-agar

supplemented with the indicated hydrogen peroxide concentrations for 48 h at 30°C. G) Fold

change expression levels ($2^{-\Delta\Delta Ct}$ with 16S as reference gene), of *dmeR*, *aitP* and *mbfA* with

Co^{2+} or H) Fe^{2+} treatment with low (0.75 mM 2,2'-bipyridyl (Bpy)) or high Fe^{2+} (0.75 mM

$FeSO_4$). Cells were grown till mid-log phase at 30°C in TY. At this point media were

supplemented accordingly and incubated for 45 min at 30°C before RNA extraction. Data

are the mean ± SE of three independent experiments. * = *p* < 0.05 and # = *p*<0.01 from a

two tail Student's t-test of averaged ΔC_t (treated *vs.* non-treated/Bpy).

Figure 2. AitP transports Fe with low affinity and high specificity. Fe^{2+} uptake was

measured in energized everted membrane vesicles (EMV) obtained from *E. coli* Δ*fieF*

expressing AitP (see materials and methods). A) Time course and NADH dependency of

Fe^{2+} uptake into EMV. Vesicles obtained from Δ*fieF* transformed with empty plasmid (empty

and filled squares) or containing *aitP* (empty and filled circles) were assayed for NADH-

dependent Fe^{2+} uptake activity. Notice that only AitP containing EMV (filled circles) present

NADH-dependent activity. Initial rate conditions were assumed until 5 minutes after Fe^{2+}

addition. Fe^{2+} concentrations for each trace are indicated (for control conditions only 5 mM

traces are shown for clarity as the other concentrations did not show significant differences).

B) Fe^{2+} transport kinetics. Data were fitted to $v= V_{max} [Fe^{2+}]/ K_{1/2} + [Fe^{2+}]$ equation by using

$K_{1/2}$ = 1.35±0.21 mM and V_{max} = 16.4±1.0 nmol.mg^{-1}.min^{-1}. C) Apparent effects on Fe^{2+}

transport rate by AitP of unlabeled divalent transition metals (TM^{2+}) as real substrates or

competitive inhibitors. Dashed line depicts the expected transport rate value obtained using

$v= V_{max} [Fe^{2+}]/ (K_{1/2} (1+ ([TM^{2+}]/K_i))+[Fe^{2+}])$, with $[Fe^{2+}]=[TM^{2+}]$= 2 mM and K_i assumed to be

equal to $K_{1/2}$. Note that only Co^{2+} affects the Fe^{2+} transport rate negatively, probably as real

substrate. In all cases, Fe^{2+} uptake by vesicles from empty vector-transformed strain *E. coli*

Δ*fieF* was subtracted prior to data analysis. In all cases, data are the mean ± SE of three

independent experiments.

Figure 3. AitP is involved in Fe^{2+} homeostasis. A) Serial dilutions (10 µl) of *S. meliloti*

Rm1021, Ω04167, Ω1163*,* Ω00359, Ω3004, and Ω03808 cells, initially at OD$_{600nm}$=0.4, were

spotted on TY-agar supplemented with the indicated Co^{2+} (as Cl$^-$) concentration for 48 h at

30°C. B) Cobalt content in Rm1021, Ω04167, Ω3004, and Ω03808 after incubation with 0.2

or 0.5 mM Co^{2+} for 1 hour at 30°C in TY medium during early exponential phase growth. C)

Serial dilutions (10 µl) of *S. meliloti* Rm1021, Ω04167*,* and C-Ω04167 cells, initially at

OD$_{600nm}$=0.4, were spotted on TY-agar supplemented with the indicated Co^{2+} (as Cl$^-$) and

Fe^{2+} (as SO$_4^{2-}$) concentration for 48 h at 30°C. D) Fold change expression levels ($2^{-\Delta\Delta Ct}$ with

16S as reference gene) of *fbpA*. Cells were grown till mid-log phase at 30°C in TY. At this

point media were supplemented with 200 µM CoCl$_2$ and incubated for 45 min at 30°C before

RNA extraction E) Total heme was estimated after cell incubation with 0.2 mM $CoCl_2$ for 2

hours at 30°C during the exponential growth phase. Data points and the mean of three

independent experiments are shown. * = $p<0.05$. F) Fold change expression levels ($2^{-\Delta\Delta Ct}$

with 16S as reference gene) of *sufA*. Cells were grown till mid-log phase at 30°C in TY. At

this point media were supplemented accordingly and incubated for 45 min at 30°C before

RNA extraction. For D) and F) Data are the mean ± SE of three independent experiments. *

= $p<0.05$ and # = $p<0.01$ from a two tail Student's t-test of averaged ΔC_t (treated *vs.* non-

treated).

Figure 4. AitP activity induces heme uptake transport. A) Fold change expression levels

($2^{-\Delta\Delta Ct}$ with 16S as reference gene) of heme uptake components in WT (light gray) and

Ω04167 (dark gray) *in vitro vs.* 16S reference gene. Cells were grown till mid-log phase at

30°C in TY. After this, half of the culture volume was supplemented with $CoCl_2$ or not, and

incubated for 2 hours at 30°C before RNA extraction. Data are the mean ± SE of four

independent experiments. * = $p<0.05$ and # = $p<0.01$ from a two-tail Student's t-test of

averaged ΔC_t (treated *vs.* non-treated). B) Serial dilution of *S. meliloti* Rm1021, Ω04167*,*

and C-Ω04167 cells, initially at $OD_{600nm}=0.4$, were spotted (10 μl) on TY-agar supplemented

with the indicated Co^{2+} (as Cl^-) and hemin concentration for 48 h at 30°C. C) Serial dilutions

of Rm1021, Ω04167, and C-Ω04167 cells, initially at $OD_{600nm}=0.4$, were spotted (10 μl) on

TY-agar supplemented with the indicated Co^{2+} (as Cl^-) and 200 μM PPIX or 200 μM hemin

concentration for 48 h at 30°C. D) Hemin uptake assay of WT (light gray) and Ω04167 (dark

gray) cells. Different letters designate significantly different means as informed by a

Bonferroni post hoc ($p<0.05$) ANOVA test. E) Illustrative photograph of cell pellets showing

Co^{2+} induced hemin uptake. Cells were grown in TY at 30°C till mid-exponential phase and

then incubated with Co^{2+} and hemin concentrations as indicated for 2 hours (n=2). After

several washes, pellets were imaged. Notice the darker green color of Rm1021 cells (upper-

right) after the Co^{2+}+hemin incubation *vs. ΔaitP,* indicating increased hemin accumulation.

Picture shows representative results of at least three independent experiments. F)

Quantification of total hemin in cell pellets. *=p<0.01 vs Rm1021.

Figure 1

Figure 2

Figure 3

Figure 4