Sedimentary DNA can reveal past pelagic copepod population dynamics

Kai Nakane[1], Xin Liu[2], Hideyuki Doi[3], Gaël Dur[4], Michinobu Kuwae[1], Syuhei Ban[2],

Narumi Tsugeki[5]

[1] Center for Marine Environmental Studies, Ehime University, Matsuyama 790-8577,

Ehime, Japan

[2] Department of Ecosystem Studies, School of Environmental Science, The University of

Shiga Prefecture, Hikone 522-8533, Shiga, Japan

[3] Graduate School of Simulation Studies, University of Hyogo, Kobe 650-0047, Hyogo,

Japan,

[4] Creative Science Unit (Geoscience), Faculty of Science, Shizuoka University, Suruga-

ku 422-8529, Shizuoka, Japan

[5] Faculty of Law, Matsuyama University, Matsuyama 790-8578, Ehime, Japan

*Corresponding author: Narumi Tsugeki

Email: narutsugeki@gmail.com

Keywords: sedDNA, *Eodiaptomus japonicus*, copepod dynamics, Lake Biwa, quantitative

PCR

Abstract

1．Copepods play a key trophic role as secondary producers in transferring primary production to higher trophic levels such as fish. Copepod production significantly contributes to successful fish recruitment. Despite their importance, knowledge of their dynamics over several decades remains limited due to the difficulty to obtain long-term data series with exhaustive sampling and analysis. However, an understanding long-term copepod dynamic is urgently required to head toward better management for sustainable aquatic ecosystems and fish recruitment.

2．Sedimentary DNA (sedDNA) has been developing a useful tool for reconstructing past plankton dynamics. This study evaluates whether sedDNA targeting the pelagic copepod, *Eodiaptomus japonicus*, in Lake Biwa (Japan) can be an effective tool for elucidating its past population dynamics.

3. It was applied a quantitative polymerase chain reaction method targeting the mitochondrial cytochrome c oxidase subunit I gene on sediment cores and compared the detected sedDNA concentrations with the unique long-term dataset of demographic traits, biomass, specific growth rate, production, subitaneous eggs, and resting eggs of *E. japonicus*.

36 4．The sedDNA concentration of *E. japonicus* recovered from sediment layers did not

37 correlate significantly with its resting eggs and its specific growth rate but with *in situ*

38 production, biomass, and subitaneous egg production. The results provided the evidence

39 for the sedDNA as a tracking tool for assessing copepod past production dynamics.

Introduction

Copepods are the most abundant multicellular organisms on Earth (Schminke, 2007) and are important secondary producers that transfer primary production to higher trophic levels such as fish in aquatic ecosystems (Mauchline, 1998). Copepod production as a food resource is essential for sustainable fish management (Mauchline, 1998). Successful fish recruitment is highly dependent on copepod production because it promotes fish larvae survival (Mauchline, 1998). Copepod population growth is influenced by temperature and food availability (Liu *et al*., 2015; Ban, 1994; Koski & Kuosa, 1999; Hart, 1990). Previous studies have shown that global warming decreases copepod abundance (Garzke *et al*., 2015). Also, large-scale climate events showing oscillation cycles over decades, such as the Pacific Decadal Oscillation and Arctic Oscillation, regulate ocean temperature and consequently affect copepod biomass and production (Peterson & Schwing, 2003; Chiba *et al*., 2006; Liu *et al*., 2021). Information about long-term copepod population dynamics is urgently required to achieve better fish management and to understand how large-scale climatic events influence aquatic ecosystems. Nevertheless, copepod dynamics across decades are rarely investigated due to the cost and time required to obtain long-term data series with exhaustive field sampling and sample analysis.

Sedimentary archives are valuable for gaining information on zooplankton long-

term dynamics (Jankowski & Straile, 2003; Brede *et al*., 2009). For specific species, such

as the cladoceran zooplankton *Daphnia*, whose remains and resting eggs are well

preserved in sediment (Jankowski & Straile, 2003; Tsugeki *et al*., 2009), morphological

investigation and genetic analysis are powerful tools to clarify biological responses to

anthropogenic perturbations (Jankowski & Straile, 2003; Brede *et al*., 2009; Monchamp

et al., 2017). However, copepods have been excluded from palaeoecological studies

because their remains are not well preserved in sediments (Selden *et al*., 2010). Although

copepod resting eggs are preserved in sediments (Hairston *et al*., 1995), they are seldom

used in palaeoecological studies because of their poor taxonomic features (Briski *et al*.,

2011). Consequently, traditional investigations of sediment archives are insufficient for

establishing copepod past dynamics. Alternatively, to overcome such difficulties in

reconstructing species lacking remains, sedimentary DNA (sedDNA) could be a useful

complementary method (Capo *et al*., 2021). Using the sedDNA approach, scientists could

reconstruct plankton community (Monchamp *et al*., 2016; Huang *et al*., 2020; Epp *et al*.,

2010) and fish dynamics (Kuwae *et al*., 2020). Combined studies of sedDNA with

morphological techniques have been conducted to track the community composition of

plankton communities (Monchamp *et al*., 2016; Huang *et al*., 2020). However, most

quantitative studies on sedDNA have focused on phytoplankton; thus, the utility of the

sedDNA approach to reconstruct successive zooplankton dynamics remains limited. A

previous study on zooplankton sedDNA targeting for *Daphnia galeata* and *Daphnia*

pulicaria in Lake Biwa (Japan) using quantitative polymerase chain reaction (qPCR)

established successful detection of both *Daphnia* sedDNA concentrations (Tsugeki *et al*.,

2022). The temporal variation in *Daphnia* sedDNA concentrations were consistent with

their resting egg production, supporting that zooplankton sedDNA could be used as a tool

for monitoring spawning activity dating back 100 years (Tsugeki *et al*., 2022). But,

continuous and quantitative reconstruction of copepod population dynamics using

sedDNA has not been reported (Bissett *et al*., 2005; Xu *et al*., 2011). Therefore, there is

an urgent need for additional research to ensure the utility and reliability of sedDNA for

reconstructing important zooplankton taxon, including copepods (Mauchline, 1998).

To ensure the reliability of the reconstruction using sedDNA for the past population

dynamics, accumulating knowledge about the source of sedDNA is imperative. Without

such information, one cannot interpret what is reconstructed. The comparison of historical

variation between sedDNA and regular survey data in the water column could confirm

the reliability of the information provided by sedDNA. In Lake Biwa (Japan), one of the

world's ancient lakes (Hampton *et al*., 2018), regular zooplankton surveys have been

intensively conducted since 1965 (Hsieh *et al*., 2011; Liu *et al*., 2020b). Moreover, the

long-term demographic traits of the pelagic dominant copepod *Eodiaptomus japonicus*,

such as biomass, population growth rate, and production, have been determined from

1971 to 2010 with experimental and microscopic observations (Liu *et al*., 2015; Liu *et al*.,

2021; Liu *et al*., 2020b). *E. japonicus* produces both subitaneous and resting eggs; the

historical variation of resting egg concentration was recovered from sediment archives up

to 60 years ago (Liu *et al*., 2020a). Lake Biwa sediment has had a continuous record with

a high time resolution, such as two or three years per 1-cm layer sample (Tsugeki et al.

2003; Tsugeki *et al*., 2021). Therefore, *E. japonicus* from Lake Biwa is suitable for

comparing historical variation between its sedDNA and demographic traits over the past

several decades.

We first investigated whether the pelagic copepod *E. japonicus* sedDNA were

successively detected from sediment core samples collected in the deep and large northern

basin of Lake Biwa via qPCR analysis. We then tested the efficiency of sedDNA as a tool

to reconstruct its population dynamics (Fig. 1b) by comparing the detected *E. japonicus*

sedDNA and its previously determined demographic traits (Fig. 1g, h). Furthermore, to

investigate the potential sources of *E. japonicus* sedDNA, we conducted a sieving

experiment to evaluate the inclusion effects of resting eggs (Fig. 1e) and the substances

in pore water to sedDNA concentrations (Fig. 1f). We also evaluated the effects of

enzymatic inhibitors in the qPCR process (Fig. 1b) by co-sinking with phytoplankton

debris and their aggregates (Fig. 1d). Based on the findings, we discuss what was

reconstructed for *E. japonicus* from sedDNA and its reliability as a tool for reconstructing

in situ *E. japonicus* population dynamics.

Methods

Sampling site, sediment core, and zooplankton sample collection

We performed a paleogenetic analysis using Lake Biwa sediment core samples to

reconstruct temporal variations of *E. japonicus* eDNA concentrations (Fig. 1b). In this

study, four 30-cm long gravity core samples (2017LB2, 2017LB5, 2017LB7, and

2019IM1; Fig. 1a-f) were collected on August 17, 2017 (2017LB core) and August 27,

2019 (2019IM1 core) in the northern basin of Lake Biwa (Supplementary Fig. S1).

Detailed information of coring sites is shown in Supplementary Fig. S1, and the sampling

methods have been previously reported (Tsugeki *et al*., 2022). Briefly, 2017LB2 and

2017LB5 cores were used to establish the variation of *E. japonicus* sedDNA

concentrations (Fig. 1b) and to explore potential sources of its sedDNA (Fig. 1e, f).

Additionally, to develop the primer-probe for *E. japonicus,* we collected a total of

95 individuals of *E. japonicus* (including 14 egg-bearing individuals) and 91 individuals

of the sympatric copepod *Mesocyclops* spp. (including 17 egg-bearing individuals) from

water samples on May 14, 2021, in Lake Biwa. Among these, six individuals of *E.*

japonicus (including three egg-bearing individuals) and four individuals of *Mesocyclops*

spp. (including two egg-bearing individuals) were randomly selected, stored at –80 °C

until DNA extraction, and then stored at –20 °C until *in vitro* validation.

Chronology of sediment cores

Sediment chronology was established for the 2017LB7 core based on the constant

rate of supply (CRS) model of ^{210}Pb dating (Appleby & Oldfield, 1978) and verified using

the ^{137}Cs peak traced in the period 1962 to 1963 (Appleby, 2001). Details of the

chronological method have been reported elsewhere (Tsugeki *et al*., 2021). The ages and

age errors of the 2017LB2 and 2017LB5 cores were indirectly estimated using

stratigraphic correlations between the cores based on chronological controls in

chlorophyll pigments and magnetic susceptibilities of the chronological 2017LB7 core

(Tsugeki *et al*., 2021). To compare these proxies, the marked peak or trough layers were

used as reference layers (Supplementary Fig. S4). In this study, sedimentary records of

resting eggs for *E. japonicus* (Liu *et al*., 2020a) were newly denoted with improved

chronology estimated using CRS-model (Supplementary Fig. S5a, b).

DNA extraction and primer-prove development for *E. japonicus*

DNA extraction of sediment samples followed the methods of previous studies

(Kuwae *et al*., 2020; Tsugeki *et al*., 2022; Sakata *et al*., 2020) using the 'Experienced User

Protocol 3 to 22' of a fecal-soil DNA extraction kit (Power Soil DNA Isolation Kit, Qiagen,

Germany). Finally, 200 µL of the DNA solution was obtained and stored at –20 °C until

qPCR analysis. Detailed information on the DNA extraction procedure from 9 g sediment

sample has been already reported (Tsugeki *et al*., 2022).

The primer-probe for *E. japonicus* in qPCR analysis was not purchased by a

company, thus we exploited it for this study (Supplementary Table S1a). We downloaded

the mitochondrial cytochrome c oxidase subunit I gene (COI) from the National Center

for Biotechnology Information (NCBI, http://www.ncbi.nlm.nih.gov/). We then

compared these COI genes among the target species *E. japonicus*, and non-target species

of copepod *Mesocyclops thermocyclopoides*, *Cyclops vicinus,* and *Daphnia* in Lake Biwa

which is main zooplankton species (Liu *et al*., 2020b). From the preliminary investigation,

we identified the variability of sequences among *E. japonicus* and other species, obtained

COI sequences of *E. japonicus* and other zooplankton species, and then constructed the

primer-probe using Primer3plus (https://www.bioinformatics.nl/cgi-

bin/primer3plus/primer3plus.cgi).

The reference sequences for the targeted gene regions were queried on possible

amplicons between 50–150 bp using NCBI primer blast. The specificities of the primers

and probe were then assessed *in silico* with homologous sequences from other copepods

and *Daphnia* species in Lake Biwa using NCBI, by targeting 116 bp of the mitochondrial

COI gene. Once PCR amplicons were identified for suitable detection, the respective

primers and probe were tested against template DNA originating from the *E. japonicus*

species to confirm amplification. During the *in silico* screening for specificity, we

performed Primer-BLAST (http://www.ncbi.nlm.nih.gov/tools/primer-blast/). We

checked all species from Japan for the order species. Using the *E. japonicus* primer-set,

we did not detect any species.

qPCR

DNA samples were quantified via real-time TaqMan quantitative PCR using the

QuantStudio™ 1 Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA,

USA) in Center for Marine Environmental Studies, Ehime University. The qPCR method

was followed according to a previous study (Tsugeki *et al*., 2022) on the developed

primer-probe set for *E. japonicus* (Supplementary Table S1a). Briefly, the TaqMan

reaction contained 900 nM of each forward and reverse primer, 125 nM TaqMan-Probe,

10 µL qPCR master mix (TaqPath; Thermo Fisher Scientific), and 4.0 µL template DNA

solution. The final volume of the PCR was 20 µL after adding UltraPure™ DNase/RNase-

Free Distilled Water (Thermo Fisher Scientific). The dilution gradient pattern of 20,000,

2,000, 200, and 20 copies per PCR reaction ($n = 4$) was proposed for the standard curve

using the target DNA cloned into a plasmid. The R^2 values of the regression analysis for

the standard curves ranged from 0.983 to 0.993 (PCR efficiency = 81.2 to 98.4%). The

qPCR data of the DNA copies (copies g^{-1} dry sed.) were shown as mean ± SD of four

replicates including zero (i.e., no detection). We also used four replicates for each sample

and an NTC ($n = 4$). No positives were detected from the NTC and the negative control

of DNA extraction, confirming that there was no cross-contamination in any of the DNA

measurements.

To determine primer-probe specificity, *in vitro* validation was performed using

extracted DNA of *E. japonicus* and other copepod *Mesocyclops* spp. collected from a

water sample in Lake Biwa (respective 5 pg per PCR reaction, $n = 4$). Additionally, *in*

vivo validation was performed using DNA extracted from the 2017LB5 core samples of

9.5 cm, 19.5 cm, and 29.5 cm depth layers ($n = 4$ per layer). Positive samples were

randomly collected among the qPCR amplicons of 2017LB2 ($n = 12$) and were then

directly sequenced following ExoSAP-IT (USB Corporation, Cleveland, OH, USA)

treatment. Sequences were determined using a commercial sequencing service (Eurofins

Genomics, Tokyo, Japan).

Inhibitor test

To test the PCR inhibition effect of substances and minerals included in sediments,

spike tests were performed for the 2017LB2 and 2017LB5 core samples (Fig. 1b).

Detailed information on the spike test was provided in a previous study (Tsugeki *et al.*,

2022) and $\Delta Ct \geq 3$ cycle appeared to be evidence of inhibition (Hartman *et al.*, 2005). We

used the primer-probe sets for IPC as follows:

IPC1-5′ : CCGAGCTTACAAGGCAGGTT

IPC1-3′: TGGCTCGTACACCAGCATACTAG

IPC1-Taq: [FAM] TAGCTTCAAGCATCTGGCTGTCGGC [TAMRA]

Demographic traits and experimental test to explore potential sources of sedDNA

To determine the source of *E. japonicus* sedDNA, we evaluated the correlation

between the sedDNA concentration and demographic traits of this copepod during the

past years, including biomass, specific growth rate, population production, and the

number of subitaneous and resting eggs (Fig. 1g, h). Long-term monthly data sets of

biomass, specific growth rate, and population production in previous studies were

referred to (Liu *et al*., 2021; Liu *et al*., 2020b). The calculation methodology is

summarized in Supplementary Table S4; detailed descriptions are provided in previous

studies (Liu *et al*., 2015; Liu *et al*., 2021; Liu *et al*., 2020b). Briefly, biomass (g dry weight

m^{-2}) was calculated as the product of the population density (individuals m^{-2}) with the

mean body dry-weight (g individual^{-1}). Specific growth rate (g day $^{-1}$) was estimated using

a multiple regression equation with two parameters: mean water temperature (°C) during

the copepod development and the food index as a proxy for availability of food required

for *E. japonicus* growth (Liu *et al*., 2015). The food index was calculated from the ratio

of the prosome length of the females collected from Lake Biwa to the expected prosome

length of the females obtained from laboratory experiments under adequate food

conditions. Population production (gC m^{-2} month^{-1}) was calculated as the product of

biomass and specific growth rate. The concentration of subitaneous eggs (eggs m^{-2}) was

estimated as the product of proportion of ovigerous female (%), ovigerous female density

(individuals m^{-2}), and clutch size (eggs clutch^{-1}). Here, all parameters used in the

calculation of subitaneous eggs were based on counts of individuals collected by field

survey (Dur *et al*., in press). The concentration of resting eggs (eggs g^{-1} dry sediment)

was calculated as the ratio of the number of resting eggs counted by microscopy from

each layer of the 2013LB core (eggs sample^{-1}) to the dry weight for each layer of sediment

(g dry sediment) (Liu *et al*., 2020a).

To determine whether *E. japonicus* sedDNA concentrations were regulated by DNA

derived directly from resting eggs included in the analytical sediment, we divided the

sediment sample into two fractions using a sieve to exclude the eggs (Fig. 1e). Since the

minimum size of the resting egg of this copepod is approximately 80 µm (Liu *et al*.,

2020a), the analytical sediments for DNA extraction were divided into two size classes

using a 38-µm mesh-sieve on three randomly selected-layer samples (specifically,

2017LB5-2.5 cm, 2017LB5-10.5 cm, and 2017LB5-19.5 cm). Furthermore, to test the

possibility of vertical movement of *E. japonicus* sedDNA through pore waters, we

examined the sedDNA concentration in pore water and its residual sediment by separating

them (Fig. 1f). The DNA extractions of the aforementioned samples were performed

according to previous studies (Tsugeki *et al*., 2022).

Possible factors regulating *E. japonicus* sedDNA

To explore potential factors regulating temporal variation in sedDNA concentrations,

we analyzed chlorophyll-a pigments and algal remains. Sedimentary pigments of

chlorophyll-a were investigated for the 2017LB2 core, and algal remains were

investigated for the 2017LB7 core (Fig. 1d). Details of the method used for chlorophyll-

a and algal remains are described in previous studies (Tsugeki *et al*., 2022; Tsugeki *et al*.,

2021).

Data analysis

All statistical analyses were performed using R ver. 4.1.2 (R Core Team 2021) with

the package "smatr" ver. 3.4.8. Type II regression models along with the standardized

major axis method were used to determine the relationship between sedDNA

concentrations of *E. japonicus* and demographic properties of biomass, specific growth

rate, production, and abundance of *E. japonicus* subitaneous, resting eggs, and algal

proxies. Since sedDNA and resting egg analysis for *E. japonicus* (Liu *et al*., 2020a) were

conducted on different sediment cores (2017LB2, 2017LB5, and 2013LB; Fig. 1b, h), the

chronological age of each analytical sample was slightly different. Therefore, prior to

statistical analysis, each sedDNA and resting egg concentration in the respective core was

converted to the value at 0.1-year intervals by linear interpolation using the AnalySeries

2.0.3 software package (Paillard *et al*., 1996). We then recalculated the average values

within the year corresponding to the sedimentation period in each sample of the

chronology 2017LB7 core. The monthly data of demographic traits in *E. japonicus*

including biomass, specific growth rate, production, and abundance of subitaneous eggs

were annually averaged and converted to the value at 0.1-year intervals by linear

interpolation. Thereafter, we calculated the average values in the respective sedimentation

period of the 2017LB7 core samples, as mentioned above. These demographic data were

limited to the year 2010. However, we estimated that the value until 2011 corresponded

to the estimated age at 4.5 cm depth of 2017LB7 core by linear extrapolation using

AnalySeries 2.0.3 software package (Paillard *et al*., 1996). Prior to Type II regression

analysis, distribution was tested by the Shapiro–Wilk test (significant level $p < 0.05$).

Since some data such as sedDNA concentration, subitaneous eggs were identified as non-

Gaussian distribution, they were $\log_{10}$-transformed. We then used Type II regression

models to determine the relationship between the $\log_{10}$-transformed values of sedDNA

concentration and their demographic traits and algal proxies.

Results

sedDNA concentration of *E. japonicus*

The results of qPCR analysis in both cores (2017LB2 and 2017LB5; Fig. 1b) showed

that *E. japonicus* sedDNA copies were detected continuously throughout the study period

(Fig. 2a, b), except for three missing layers for the 2017LB5 core (5–6 cm, 6–7 cm, 7–8

cm layers; hereafter expressed in middle depth, i.e., 5.5 cm, 6.5 cm, and 7.5 cm,

respectively, Supplementary Table S2). The mean DNA copy number of *E. japonicus*

varied depending on the core depth and ranged from 6 ± 11 (mean $\pm$ SD) copies g^{-1} dry

sediment (hereafter copies g^{-1}) in the depth of the 2017LB2-28.5 cm layer (estimated age:

1924.5 ± 39.0) to $290,330 \pm 30,848$ copies g^{-1} in the 2017LB2-4.5 cm layer (estimated

age: 2007.3 ± 0.4). Supplementary Table S2). They were almost abundantly detected in

three to four replicates in the upper 8.5 cm layers (estimated age: 1993.8 ± 1.0 and 1999.0

± 0.7 for 2017LB2 and 2017LB5 cores, respectively; Fig. 2a, Supplementary Table S2).

The DNA copy numbers below 8.5 cm layers were relatively low but were detected from

all layers in one to four replicates (Fig. 2b, Supplementary Table S2).

No DNA of *E. japonicus* was detected in any of the negative controls for sub-

sampling, DNA extraction, or PCR analysis, or the no-template control (NTC), indicating

that sample contamination did not occur during these processes. We also confirmed that

only *E. japonicus* DNA was amplified through direct sequencing of PCR amplicons, using

qPCR analysis of *E. japonicus* DNA.

A spike test was performed on both sediment core samples (2017LB2 and 2017LB5)

for evaluating the effect of qPCR inhibition (Fig. 1b). All ΔCt values were below 1 (–1.0

to 0.87: 2017LB2, –0.49 to 0.21: 2017LB5; Supplementary Table S3), indicating no

inhibition.

318 Long-term variation in demographic traits of *E. japonicus*

All the mean values of demographic traits such as biomass, specific growth rate, and

production for *E. japonicus* were enumerated depending on the time resolution of

deposition period on each sample (see details in data analysis). The average biomass of

E. japonicus till approximately the year 2000 was relatively low with a range of 0.7 to

1.2 g dry weight m^{-2}, except for the values measured around 1970 (1.7 g dry weight m^{-2}),

but the average biomass rapidly increased approximately two-fold after 2010 compared

to that before 2000 (Supplementary Fig. S2a). Likewise, the average mean of their

production was stable before approximately 2000, at less than 1.5 g C m^{-2}, and then

continuously increased up to 2010 (Supplementary Fig. S2c). Contrarily, the average

mean of specific growth rate (*g*) remained relatively constant after 1970 at approximately

0.07 day^{-1}. The lowest value of specific growth rate was observed around 1970 and the

highest value was observed around 2010 (Supplementary Fig. S2b). The average mean of

subitaneous eggs remained constant, being less than 1.5×10^5 eggs m^{-2} up to approximately

2000, excluding the marked peak in the late 1990s, and continuously increased 2000

onward (Supplementary Fig. S2d). The average mean of resting egg concentration

recovered from sediment observation increased markedly from the mid-1980s, peaked

around the early 1990s, and then decreased and stabilized (Supplementary Fig. S2e).

Correlations of the *E. japonicus* sedDNA and demographic traits with the algal

remain concentrations

A highly significant correlation was detected between the *E. japonicus* sedDNA

concentration and *E. japonicus* production from 1971 to 2010 (Type II regression model:

$R^2 = 0.60$, $p = 0.001$, $n = 14$; Fig. 3c), in addition to its biomass concentration ($R^2 = 0.51$,

$p = 0.004$, $n = 14$; Fig. 3a). Significant correlation was also detected between the sedDNA

concentration and number of subitaneous *E. japonicus* eggs ($R^2 = 0.52$, $p = 0.005$, $n = 13$;

Fig. 3d). In contrast, no significant relationship was observed between the sedDNA

concentration and the specific growth rate ($R^2 = 0.02$, $p = 0.66$, $n = 14$; Fig. 3b), the

number of resting eggs ($R^2 = 0.25$, $p = 0.066$, $n = 14$; Fig. 3e). *E. japonicus* sedDNA

concentrations were positively corelated with diatom valves ($R^2 = 0.65$, $p < 0.001$, $n = 27$;

Supplementary Fig. S4a), chlorophyll-a ($R^2 = 0.71$, $p < 0.001$, $n = 29$; Supplementary Fig.

S4b) and green algal cells ($R^2 = 0.37$, $p = 0.001$, $n = 25$; Supplementary Fig. S4c),

although these long-term trends differed from the trend of variation in *E. japonicus*

sedDNA concentrations (Supplementary Fig. S4d, e, f).

Potential sources of *E. japonicus* sedDNA

E. japonicus environmental DNA (eDNA) was not detected from the pore water of

all three samples from the 2017LB5 core by qPCR (Fig. 1f, Table 1). In contrast, *E.*

japonicus eDNA was detected in the residual sediment samples. *E. japonicus* sedDNA

was detected in both sieved and non-sieved samples, except for the sieved sample of

2017LB5-20 (Table 2). High concentrations of sedDNA were detected in the non-sieved

samples for two of the three samples, namely 2017LB5-3 (11,630 ± 5,921 copies g^{-1})

2017LB5-20 (1,449 ± 1,977 copies g^{-1}), compared to those in the sieved samples.

However, for 2017LB5-11, the sieved sample (2,630 ± 4,555 copies g^{-1}) was more

abundant than the non-sieved sample (1,838 ± 3,184 copies g^{-1}), although the difference

between both values was minimal.

Discussion

Several studies have reported the long-term preservation of copepod eDNA in

aquatic sediments (Bissett *et al*., 2005; Xu *et al*., 2011; Coolen *et al*., 2013), such as a

report by Bissett *et al*. (2005), which detected copepod eDNA from sediment layers as

old as 9,950 years in some Antarctic lakes. However, continuous, and quantitative

reconstruction of copepod sedDNA is not yet to be reported (Domaizon *et al*., 2017). This

study is the first to successively detect copepod eDNA for *E. japonicus* from a sediment

core in Lake Biwa over the past 100 years with high-resolution dating.

We found that the temporal variation of *E. japonicus* sedDNA concentration

significantly correlated with its production and biomass, obtained via microscopic

observations of regular field samples. In support of this, the number of reads for copepod

obtained by metabarcoding genetic analysis correlated with their biomass in water

samples (Yang *et al*., 2017) although a meaningful relationship between eDNA abundance

and biomass for some copepod species was not well observed (Djurhuus *et al*., 2018). We

also established the temporal increase in *E. japonicus* sedDNA concentration in the 1970s,

which is more than 50-fold compared to those before the 1970s when eutrophication

occurred in Lake Biwa (Hyodo *et al*., 2008). Concordant with this finding, regular survey

reports in Lake Biwa described the density of *E. japonicus* as being temporally and greatly

increased in 1975 with densities more than 10-times higher than those in 1974 (Shiga

Prefecture, 1964-1979). This coincidence of increased sedDNA concentration and the

marked expansion of *E. japonicus* in the water column provides evidence that copepod

sedDNA concentration can be a powerful tool to estimate their population biomass over

several decades.

The temporal variations in the concentration of *E. japonicus* sedDNA were suspected

to operate by diagenetic degradation. Leaching from sediments had been mainly observed

in coarse-textured sediments that enable fluid flow across layers (Andersen *et al*., 2012;

Haile *et al*., 2007). Thus, leaching is generally assumed as a minor process in water-

saturated sediments (Giguet-Covex *et al*., 2014) like the fresh sediment investigated in

the current study. Since sedimentary compaction formed over an extended period, it is

likely that upward fluid flow into the surface layer arises permanently, inducing DNA

leaching. However, we did not detect *E. japonicus* eDNA in any of the pore water samples,

as previously shown in fish (Kuwae *et al*., 2020) and *Daphnia* (Tsugeki *et al*., 2022)

sedDNA analysis in sediment samples. Therefore, it is reasonable to conclude that leached

E. japonicus eDNA only negligibly influences sedDNA. Moreover, a recent experimental

study showed almost no reduction for sedDNA concentration after initial decline within

one month (Ogata *et al*., 2021) probably induced by decomposition. The concentrations

of *E. japonicus* sedDNA did not always show increasing trends toward the surface layers.

Rather, it decreased toward the surface after the peak observed before 2010, except for

the top surface layer. This indicates that the temporal variation in sedDNA of *E. japonicus*

cannot be simply explained by time-dependent diagenetic degradation. The sedDNA

concentration of *E. japonicus* denoted some correlations with the other proxies, such as

pigment concentration and algal diatom remains. However, the long-term trends were

different from those of *E. japonicus* sedDNA, indicating that *E. japonicus* sedDNA

concentration is not regulated by phytoplankton sinking and/or their debris aggregations.

We detected that the sedDNA concentrations of *E. japonicus* correlated with the

concentration of its subitaneous eggs. Moreover, we observed high concentrations of

sedDNA from the non-sieved samples likely including resting eggs in two of the three

samples compared to those in the sieved samples excluding eggs. However, we did not

detect significant correlation with the number of resting eggs, which is likely because in

Lake Biwa, subitaneous eggs, rather than the resting eggs, mainly regulate the spawning

activity of *E. japonicus* (Liu *et al*., 2020a). The contribution of resting eggs to total *E.*

japonicus eggs in the water column was expected to be < 0.3% (Liu *et al*., 2020a). A

recent study demonstrated that eDNA were released during spawning periods (Takeuchi

et al., 2019; Tillotson *et al*., 2018). In addition, in a report of zooplankton *Daphnia*

sedDNA, substances released during spawning activities were assumed to be a significant

source of DNA archived in sediments (Tsugeki *et al*., 2022). Therefore, these findings

emphasize the importance of sedDNA-sourced material being released during the main

spawning period. As shown in the present study, the sedDNA concentrations of *E.*

japonicus markedly increased more than 3-fold during the 2000s compared to those

before the 1990s, which significantly correlated with their production and biomass. The

production of *E. japonicus* greatly increased more than approximately 2-fold in the 2000s,

mainly due to the decline of predation effect (Liu *et al*., 2021), which occurred in parallel

with their temporal increase of specific growth rate and subitaneous eggs. These facts

support that copepod sedDNA can be applied as a powerful tracking tool for

reconstruction of their production and biomass related to the intensity of spawning

activity.

In conclusion, we successively and qualitatively detected *E. japonicus* sedDNA over

a 100-year period, demonstrating the potential utility of sedDNA in evaluating the long-

term qualitative dynamics of copepod. The absence of *E. japonicus* sedDNA in the pore

water samples suggested that the temporal variation of its sedDNA concentration was not

interrupted by leaching reaction after sedimentation. The historical variation in sedDNA

concentration was similar to its production, biomass, and subitaneous eggs found in

regular *in situ* survey samples. Therefore, the *E. japonicus* sedDNA concentration is a

possible indicator of the magnitude of productive activities in the water. Overall, we

confirmed that sedDNA can be used as a proxy for monitoring the production and

spawning activity of copepods, which can track 100 years of sedimentary records.

References

Andersen, K., Bird, K. L., Rasmussen, M., Haile, J., Breuning-Madsen, H., Kjaer, K. H., Orlando, L., M. THOMAS P. Gilbert, M. T. P., & Willerslev, E. (2012). Meta-barcoding of 'dirt' DNA from soil reflects vertebrate biodiversity. *Molecular Ecology*, 21, 1966–1979. https://doi.org/10.1111/j.1365-294X.2011.05261.x

Appleby, P. G. (2001). Chronostratigraphic Techniques in Recent Sediments in *Track. Environ. Chang. Using Lake Sediments: Basin Analysis, Coring, and Chronological Techniques* (eds. Last, W. M., Smol, J. P.) 171–203 (Kluwer Academic, Dordrecht, The Netherlands).

Appleby, P. G. & Oldfield, F. (1978). The calculation of lead-210 dates assuming a constant rate of supply of unsupported ^{210}Pb to the sediment. *Catena*, 5, 1–8. https://doi.org/10.1016/S0341-8162(78)80002-2

Ban, S. (1994). Effect of temperature and food concentration on post-embryonic development, egg production and adult body size of calanoid copepod *Eurytemora affinis*. *Journal of Plankton Research*, 16, 721–735. https://doi.org/10.1093/plankt/16.6.721

Bissett, A., Gibson, J. A. E., Jarman, S. N., Swadling, K. M. & Cromer, L. (2005).

461   Isolation, amplification, and identification of ancient copepod DNA from lake

462   sediments. *Limnology and Oceanography: Methods*, 3, 533–542.

463   https://doi.org/10.4319/lom.2005.3.533

464 Brede, N., Sandrock, C., Straile, D., Spaak, P., Jankowski, T., Streit, B. & Schwenk, K.

465   (2009). The impact of human-made ecological changes on the genetic

466   architecture of *Daphnia* species. *Proceedings of the National Academy of*

467   *Sciences of the United States of America*, 106, 4758-4763.

468   https://doi.org/10.1073/pnas.0807187106

469 Briski, E., Cristescu, M. E., Bailey, S. A. & MacIsaac, H. J. (2011). Use of DNA

470   barcoding to detect invertebrate invasive species from diapausing eggs.

471   *Biological Invasions*, 13, 1325–1340. https://doi.org/10.1007/s10530-010-9892-

472   7

473 Capo, E., Giguet-Covex, C., Rouillard, A., Nota, K., Heintzman, P. D., Vuillemin, A.,

474   Ariztegui, D., Arnaud, F., Belle, S., Bertilsson, S., Bigler, C., Bindler, R., Brown,

475   A. G., Clarke, C. L., Crump, S. E., Debroas, D., Englund, G., Ficetola, G. F.,

476   Garner, R. E., Gauthier, J., Gregory-Eaves, I., Heinecke, L., Herzschuh, U.,

477   Ibrahim, A., Kisand, V., Kjær, K. H., Lammers, Y., Littlefair, J., Messager, E.,

478   Monchamp, M.-E., Olajos, F., Orsi, W., Pedersen, M. W., Rijal, D. P., Rydberg,

J., Spanbauer, T., Stoof-Leichsenring, K. R., Taberlet, P., Talas, L., Thomas, C.,

Walsh, D. A., Wang, Y., Willerslev, E., van Woerkom, A., Zimmermann, H. H.,

Coolen, M. J. L., Epp, L. S., Domaizon, I., Alsos, I. G., Parducci, L. & Parducci,

L. (2021). Lake sedimentary dna research on past terrestrial and aquatic

biodiversity: Overview and recommendations. *Quaternary*, 4, 6.

https://doi.org/10.3390/quat4010006

Chiba, S., Tadokoro, K., Sugisaki, H. & Saino, T. (2006). Effects of decadal climate

change on zooplankton over the last 50 years in the western subarctic North

Pacific. *Global Change Biology*, 12, 907–920. https://doi.org/10.1111/j.1365-

2486.2006.01136.x

Coolen, M. J. L., Orsi, W. D., Balkema, C., Quince, C., Harris, K., Sylva, S. P.,

Filipova-Marinova M. & Giosan, L. (2013). Evolution of the plankton paleome in

the Black Sea from the Deglacial to Anthropocene. *Proceedings of the National*

Academy of Sciences of the United States of America, 110, 8609–8614.

https://doi.org/10.1073/pnas.1219283110

Djurhuus, A., Pitz, K., Sawaya, N. A., Rojas-Márquez, J., Michaud, B., Montes, E.,

Muller-Karger, F. & Breitbart, M. (2018). Evaluation of marine zooplankton

community structure through environmental DNA metabarcoding. *Limnology*

and Oceanography: Methods, 16, 209–221. https://doi.org/10.1002/lom3.10237

Domaizon, I., Winegardner, A., Capo, E., Gauthier, J. & Gregory-Eaves, I. (2017).

DNA-based methods in paleolimnology: new opportunities for investigating

long-term dynamics of lacustrine biodiversity. *Journal of Paleolimnology*, 58, 1–

21. https://doi.org/10.1007/s10933-017-9958-y

Dur, G., Liu, X., Sakai, Y., Hsieh, C. H., Ban, S., Souissi, S. (in press). Disrupted

seasonal cycle of the warm-adapted and main zooplankter of Lake Biwa, Japan.

Journal of Great Lakes Research.

Epp, L. S., Stoof, K. R., Trauth, M. H. & Tiedemann, R. (2010). Historical genetics on a

sediment core from a Kenyan lake: Intraspecific genotype turnover in a tropical

rotifer is related to past environmental changes. *Journal of Paleolimnology*, 43,

939–954. https://doi.org/10.1007/s10933-009-9379-7

Garzke, J., Ismar, S. M. H. & Sommer, U. (2015). Climate change affects low trophic

level marine consumers: warming decreases copepod size and abundance.

Oecologia, 177, 849–860. https://doi.org/10.1007/s00442-014-3130-4

Giguet-Covex, C., Pansu, J., Arnaud, F., Rey, P. J., Griggo, C., Gielly, L., Domaizon, I.,

Coissac, E., David, F., Choler, P., Poulenard J. & Taberlet, P. (2014). Long

livestock farming history and human landscape shaping revealed by lake

sediment DNA. *Nature Communications*, 5, 1–7.

https://doi.org/10.1038/ncomms4211

Haile, J., Holdaway, R., Oliver, K., Bunce, M., Gilbert, M. T. P., Nielsen, R., Munch,

K., Ho, S. Y. W., Shapiro, B. & Willerslev, E. (2007). Ancient DNA chronology

within sediment deposits: Are paleobiological reconstructions possible and is

DNA leaching a factor? *Molecular Biology and Evolution*, 24, 982–989.

https://doi.org/10.1093/molbev/msm016

Hairston, N. G., Van Brunt, R. A., Kearns, C. M. & Engstrom, D. R. (1995). Age and

survivorship of diapausing eggs in a sediment egg bank. *Ecology*, 76, 1706–1711.

https://doi.org/10.2307/1940704

Hampton, S. E., McGowan, S., Ozersky, T., Virdis, S. G. P., Vu, T. T., Spanbauer, T.

L., Kraemer, B. M., Swann, G., Mackay, A. W., Powers, S. M., Meyer, M. F.,

Labou, S. G., O'Reilly, C. M., DiCarlo, M., Galloway, A. W. E. & Fritz, S. C.

(2018). Recent ecological change in ancient lakes. *Limnology and*

Oceanography, 63, 2277–2304. https://doi.org/10.1002/lno.10938

Hart, R. C. (1990). Copepod post-embryonic durations: pattern, conformity, and

predictability. The realities of isochronal and equiproportional development, and

trends in the opepodid-naupliar duration ratio. *Hydrobiologia*, 206, 175–206.

https://doi.org/10.1007/BF00014085

Hartman, L. J., Coyne, S. R. & Norwood, D. A. (2005). Development of a novel

internal positive control for Taqman® based assays. *Molecular and Cellular*

Probes, 19, 51–59. https://doi.org/10.1016/j.mcp.2004.07.006

Hsieh, C. H., Sakai, Y., Ban, S., Ishikawa, K., Ishikawa, T., Ichise, S., Yamamura, N.

& Kumagai, M. (2011). Eutrophication and warming effects on long-term

variation of zooplankton in Lake Biwa. *Biogeosciences*, 8, 593–629.

https://doi.org/10.5194/bg-8-1383-2011

Huang, S., Herzschuh, U., Pestryakova, L. A., Zimmermann, H. H., Davydova, P.,

Biskaborn, B. K., Shevtsova, I. & Stoof-Leichsenring, K. R. (2020). Genetic and

morphologic determination of diatom community composition in surface

sediments from glacial and thermokarst lakes in the Siberian Arctic. *Journal of*

Paleolimnology, 64, 225–242. https://doi.org/10.1007/s10933-020-00133-1

Hyodo, F., Tsugeki, N., Azuma, J., Urabe, J., Nakanishi, M. & Wada, E. (2008).

Changes in stable isotopes, lignin-derived phenols, and fossil pigments in

sediments of Lake Biwa, Japan : Implications for anthropogenic effects over the

last 100 years. *Science of The Total Environment*, 403, 139-147.

https://doi.org/10.1016/j.scitotenv.2008.05.010

Jankowski, T. & Straile, D. (2003). A comparison of egg-bank and long-term plankton

dynamics of two *Daphnia* species, *D. hyalina* and *D. galeata*: Potentials and

limits of reconstruction. *Limnology and Oceanography*, 48, 1948–1955.

https://doi.org/10.4319/lo.2003.48.5.1948

Koski, M. & Kuosa, H. (1999). The effect of temperature, food concentration and

female size on the egg production of the planktonic copepod *Acartia bifilosa*.

Journal of Plankton Research, 21, 1779–1789. https://doi.org/10.1007/s00227-

005-0132-0

Kuwae, M., Tamai, H., Doi, H., Sakata, M. K., Minamoto, T. & Suzuki, Y. (2020).

Sedimentary DNA tracks decadal-centennial changes in fish abundance.

Communications Biology, 3, 558. https://doi.org/10.1038/s42003-020-01282-9

Liu, X., Ban, S., Beyrend, D., Dur, G., Kuwae, M., Makino, W. & Urabe, J. (2020a).

Resting eggs of the perennial copepod *Eodiaptomus japonicus* in Lake Biwa

(Japan). *Inland Waters*, 10, 89–100.

https://doi.org/10.1080/20442041.2019.1671766

Liu, X., Beyrend, D., Dur, G. & Ban, S. (2015). Combined effects of temperature and

food concentration on growth and reproduction of *Eodiaptomus japonicus*

(Copepoda: Calanoida) from Lake Biwa (Japan). *Freshwater Biology*, 60, 2003–

2018. https://doi.org/10.1111/fwb.12626

Liu, X., Dur, G., Ban, S., Sakai, Y., Ohmae, S. & Morita, T. (2021). Quasi-decadal

periodicities in growth and production of the copepod Eodiaptomus japonicus in

Lake Biwa, Japan, related to the Arctic Oscillation. *Limnology and*

Oceanography, 66, 3783–3795. https://doi.org/10.1002/lno.11918

Liu, X., Dur, G., Ban, S., Sakai, Y., Ohmae, S. & Morita, T. (2020b). Planktivorous fish

predation masks anthropogenic disturbances on decadal trends in zooplankton

biomass and body size structure in Lake Biwa, Japan. *Limnology and*

Oceanography, 65, 667–682. https://doi.org/10.1002/lno.11336

Mauchline, J. (1998). *The biology of calanoid copepods: Advances in marine biology*.

33. (Academic Press: London. UK).

Monchamp, M. E., Enache, I., Turko, P., Pomati, F., Rîşnoveanu, G. & Spaak, P.

(2017). Sedimentary and egg-bank DNA from 3 European lakes reveal

concurrent changes in the composition and diversity of cyanobacterial and

Daphnia communities. *Hydrobiologia*, 800, 155–172.

https://doi.org/10.1007/s10750-017-3247-7

Monchamp, M. E., Walser, J. C., Pomati, F. & Spaak, P. (2016). Sedimentary DNA

reveals cyanobacterial community diversity over 200 years in two perialpine

lakes. *Applied and Environmental Microbiology*, 82, 6472–6482.

https://doi.org/10.1128/AEM.02174-16

Ogata, M., Masuda, R., Harino, H., Sakata, M. K., Hatakeyama, M., Yokoyama, K.,

Yamashita, Y. & Minamoto, T. (2021). Environmental DNA preserved in marine

sediment for detecting jellyfish blooms after a tsunami. *Scientific Reports*, 11, 1–

14. https://doi.org/10.1038/s41598-021-94286-2

Paillard, D., Labeyrie, L. & Yiou, P. (1996). Analyseries 1.0: a Macintosh software for

the analysis of geographical time-series. *Eos*, 77, 379.

Peterson, W. T. & Schwing, F. B. (2003). A new climate regime in northeast pacific

ecosystems. *Geophysical Research Letters*, 30, 1–4.

https://doi.org/10.1029/2003GL017528

Sakata, M. K., Yamamoto, S., Gotoh, R. O., Miya, M., Yamanaka, H. & Minamoito, T.

(2020). Sedimentary eDNA provides different information on timescale and fish

species composition compared with aqueous eDNA. *Environmental DNA*, 2,

505–518. https://doi.org/10.1002/edn3.75

Schminke, H. K. (2007). Entomology for the copepodologist. *Journal of Plankton*

Research, 29, i149–i162. https://doi.org/10.1093/plankt/fbl073

Selden, P. A., Huys, R., Stephenson, M. H., Heward, A. P. & Taylor, P. N. (2010).

Crustaceans from bitumen clast in Carboniferous glacial diamictite extend fossil

record of copepods. *Nature Communications*, 1, 2–7.

https://doi.org/10.1038/ncomms1049

Shiga Prefecture. (1964-1979). Annual report of the regular observation in Lake Biwa.

Shiga Prefectural Fisheries Experiment Station. (in Japanese).

Takeuchi, A., Iijima, T., Kakuzen, W., Watanabe, S., Yamada, Y., Okamura, A., …

Tsukamoto, K. (2019). Release of eDNA by different life history stages and

during spawning activities of laboratory-reared Japanese eels for interpretation of

oceanic survey data. *Scientific Reports*, 9, 1–9. https://doi.org/10.1038/s41598-

019-42641-9

Tillotson, M. D., Kelly, R. P., Duda, J. J., Hoy, M., Kralj, J. & Quinn, T. P. (2018).

Concentrations of environmental DNA (eDNA) reflect spawning salmon

abundance at fine spatial and temporal scales. *Biological Conservation*, 220, 1–

11. https://doi.org/10.1016/j.biocon.2018.01.030

Tsugeki, N. K., Honjo, M. N. & Kuwae, M. (2021). Interspecific variation in ephippial

size between *Daphnia galeata* and *D. pulicaria* in Lake Biwa, Japan. *Limnology*,

22, 197–207. https://doi.org/10.1007/s10201-020-00646-8

Tsugeki, N. K., Ishida, S. & Urabe, J. (2009). Sedimentary records of reduction in

resting egg production of *Daphnia galeata* in Lake Biwa during the 20th century:

A possible effect of winter warming. *Journal of Paleolimnology*, 42, 155–165.

https://doi.org/10.1007/s10933-008-9268-5

Tsugeki, N., Nakane, K., Doi, H., Ochi, N. & Kuwae, M. (2022). Reconstruction of

100-year dynamics in *Daphnia* spawning activity revealed by sedimentary DNA.

Scientific Reports, 12, 1–11. https://doi.org/10.1038/s41598-021-03899-0

Tsugeki, N., Oda , H., Urabe, J. (2003). Fluctuation of the zooplankton community in

Lake Biwa during the 20th century : a paleolimnological analysis. *Limnology*, *4*,

101–107. https://doi.org/10.1007/s10201-003-0097-y

Xu, Z. H., Jiang, X. D., Wang, G. Z., He, J. F., Cai, M. H., Wu, L. S., … Chen, X. L.

(2011). DNA extraction, amplification and analysis of the 28S rRNA portion in

sediment-buried copepod DNA in the Great Wall Bay and Xihu Lake, Antarctica.

Journal of Plankton Research, 33, 917–925.

https://doi.org/10.1093/plankt/fbq165

Yang, J., Zhang, X., Xie, Y., Song, C., Zhang, Y., Yu, H. & Burton, G. A. (2017).

Zooplankton community profiling in a eutrophic freshwater ecosystem-lake tai

basin by DNA metabarcoding. *Scientific Reports*, 7, 1–11.

https://doi.org/10.1038/s41598-017-01808-y

Acknowledgements

We are grateful to M. Honjo, M. Sakata, S. Goda, T. Akatsuka, and J. Kurata for their assistance with laboratory analysis and field sampling. We also thank Shiga Prefectural Fisheries Experiment Station for providing preserved zooplankton sample collection. This study was supported by Grants-in-Aid for Scientific Research (17K20045, 21K12273, 21H01170, and 18H03961) from the Japan Society for the Promotion of Science (JSPS) and partly supported by a special research grant from Matsuyama University; the Academic Research Organization Joint Usage/Research Grants from Leading Academia in Marine and Environment Pollution Research (LaMer), Ehime University; the Center for Ecological Research (2017–2021 jurc-cer), Kyoto University; the cooperative research program (17A065) of the Center for Advanced Marine Core Research, Kochi University, and the Environment Research and Technology Development Fund (JPMEERF20204004) of the Environmental Restoration and Conservation Agency, Japan; and grants from the Ministry of Agriculture, Forestry and Fisheries, Japan for a research project titled *Development of technologies for mitigation and adaptation to climate change in Agriculture, Forestry and Fisheries*.

Author contributions

N.T. and K.N. conceived the idea and designed the study; K.N., N.T., and M.K.

conducted sampling and experiments; X.L., G.D., and S.B. provided data on the target

species and contributed to the data analysis; H.D. performed some of the experiments and

contributed to primer and probe design; K.N. and N.T. wrote the first draft of the

manuscript; All authors revised and commented on the manuscript.

Competing interests

The authors declare no competing interests.

Data Availability Statement

Supplementary information will be available after publication.

 Figure legends

Figure 1. Research and experimental design of the study.

(a) Sediment geochronology; (b) Reconstruction of temporal variation in *Eodiaptomus*

japonicus sedimentary DNA (sedDNA) concentrations and spike test for inhibition; (c)

Checking for contamination; (d) Reconstruction of temporal variation of chlorophyll-a

pigment concentration and algal remains abundance; (e) Evaluating potential sources of

E. japonicus sedDNA between samples with and without sieve; (f) Regulating factors of

E. japonicus sedDNA; (g) Demographic traits of biomass, specific growth rate,

production, and subitaneous eggs of *E. japonicus*; (h) Sedimentary records of resting eggs.

The named LB cores were collected in November 2013 (2013LB) and August 2017

(2017LB2, 2017LB5, and 2017LB7), and IM cores (namely 2019IM1) were collected in

August 2019 (details are described in the section on sampling site and sediment collection

of methods).

685
686

Figure 2. Temporal changes in *Eodiaptomus japonicus* sedimentary DNA concentration

(gene copies of COI, g^{-1} of dry sediments): (a) observed throughout the study period,

and (b) limited to the period before 1992. For both panels, the open triangles and

squares indicate raw data of the 2017LB2 and 2017LB5 cores, respectively. The gray

triangles and squares indicate the average mean (*n* = 4) of 2017LB2 and 2017LB5 core

samples, respectively. For the panel (a) and (b), black circles and thick solid lines

indicate the mean (*n* = 2) of 2017LB2 and 2017LB5 core samples within the year

corresponding to the sedimentation period in the chronology 2017LB7 core.

Figure 3. Relationships among $\log_{10}$-transformed sedimentary DNA concentration of

Eodiaptomus japonicus (gene copies of COI, g^{-1} of dry sediments) and its biomass (g dry

weight m^{-2}; a), specific growth rate (day^{-1}; b), production (gC m^{-2} $month^{-1}$; c), $\log_{10}$-

transformed subitaneous eggs ($\times 10^3$ egg numbers m^{-2}; d), and resting eggs (egg numbers

g^{-1} dry sed.; e). The blue line denotes a Type II regression model with a 95% confidence

interval indicated by the gray zone.

703
704

Tables

Table 1. The sedimentary DNA concentration of *Eodiaptomus japonicus* (gene copies of COI, g^{-1} of dry sediments) for pore water in sediment and its residual pore water-free sediment samples.

Sample ID (estimated age in sample depth and its age error; yr)	Core depth		Pore water in sediment sample	SD	Detection / total replicates	Pore water - free sediment sample	SD	Detection / total replicates
	Top (cm)	Bottom (cm)	(mean copies g^{-1} dry)			(mean copies g^{-1} dry)		
2017LB5-3 (2014 ± 0.1)	2	3	ND	-	0/4	725,030	98,992	4/4
2017LB5-11 (1994 ± 1.3)	10	11	ND	-	0/4	2,665	608	4/4
2017LB5-20 (1964 ± 9.3)	19	20	ND	-	0/4	12	21	1/4

Table 2. The sedimentary DNA concentration of *Eodiaptomus japonicus* (gene copies of COI, g^{-1} of dry sediments) in sediment samples with and without sieve.

Sample ID (estimated age in sample depth and its age error; yr)	Core depth		Sample with sieve	SD	Detection / total replicates	Sample non - sieve	SD	Detection / total replicates
	Top (cm)	Bottom (cm)	(mean copies g^{-1} dry)			(mean copies g^{-1} dry)		
2017LB5-3 (2014 ± 0.1)	2	3	5,246	6,531	2/4	11,630	5,921	4/4
2017LB5-11 (1994 ± 1.3)	10	11	2,630	4,555	1/4	1,838	3,184	1/4
2017LB5-20 (1964 ± 9.3)	19	20	ND	-	0/4	1,449	1,977	2/4