Molecular mechanism of substrate recognition by

folate transporter SLC19A1

Authors:

Yu Dang[1], Dong Zhou[2], Xiaojuan Du[3]†, Hongtu Zhao[4], Chia-Hsueh Lee[4], Jing Yang[3], Yijie

Wang[3], Changdong Qin[5], Zhenxi Guo[5], and Zhe Zhang[1,2,3]*

Affiliations:

[1]State Key Laboratory of Membrane Biology, Peking University-Tsinghua University-National

Institute of Biological Sciences Joint Graduate Program, Academy for Advanced

Interdisciplinary Studies, Peking University, Beijing 100871, China

[2]Center for Life Sciences, Academy for Advanced Interdisciplinary Studies, Peking University,

Beijing 100871, China

[3]School of Life Sciences, Peking University, Beijing 100871, China

[4]Department of Structural Biology, St. Jude Children's Research Hospital, Memphis, TN 38105,

United States

[5]Cryo-EM Platform, School of Life Sciences, Peking University, Beijing 100871, China

†Present address: Peking University First Hospital, Peking University Health Science Center,

Beijing 100191, China

*Correspondence should be addressed to zzhang01@pku.edu.cn.

Abstract

Folate (vitamin B$_9$) is the coenzyme involved in one-carbon transfer biochemical reactions

essential for cell survival and proliferation, with its inadequacy causing developmental defects or

severe diseases. Notably, mammalian cells lack the ability to *de novo* synthesize folate but instead

rely on its intake from extracellular sources via specific transporters or receptors, among which

SLC19A1 is the ubiquitously expressed one in tissues. However, the mechanism of substrate

recognition by SLC19A1 has been unclear. Here we report the cryo-EM structures of human

SLC19A1 and its complex with 5-methyltetrahydrofolate at 3.5-3.6 Å resolution and elucidate the

critical residues for substrate recognition. In particular, we reveal that two variant residues among

SLC19 subfamily members would designate the specificity for folate. Moreover, we identify

intracellular thiamine pyrophosphate as the favorite coupled substrate for folate transport by

SLC19A1. Together, this work has established the molecular basis of substrate recognition by this

central folate transporter.

Introduction

Folate (vitamin B9) is the coenzyme serving as the single-carbon donor in many biochemical reactions, e.g., the synthesis of purine and thymidylate, the metabolism of serine and methionine, and the methylation of nucleic acids and proteins[1]. Given such essential roles in cell growth, proliferation, and differentiation, folate inadequacy would lead to severe developmental defects or neurological disorders in humans[2,3].

Mammalian cells lack the ability to *de novo* synthesize folate and must obtain it from extracellular sources such as foods. Three different systems are known for the transmembrane uptake of folate in mammals, i.e., the proton-coupled folate transporter SLC46A1[4], the folate receptors (FRs)[5-7], and the reduced folate carrier SLC19A1[8,9]. SLC46A1 is predominantly expressed in the gastrointestinal tract and is responsible for dietary folate absorption. Accordingly, SLC46A1 exhibits an optimal activity at acidic pH and couples folate transport to proton influx[10,11]. On the other hand, FRs take folate into cells via receptor-mediated endocytosis, primarily for folate delivery to the brain or folate retention in the kidney[7,12,13]. Notably, both SLC46A1 and FRs exert tissue-specific roles in folate transport. In contrast, SLC19A1 is ubiquitously expressed in the body and represents the major system of folate transport in diverse cell types[13]. For instance, though all three systems could facilitate the cellular uptake of antifolate drugs for cancer chemotherapy, SLC19A1 is the predominant route in many cancer cells[11,14,15]. Indeed, decreased expression or loss-of-function mutations of SLC19A1 in cancers would result in resistance to antifolate treatments[16]. Additionally, while SLC46A1 and FRs have equal affinities to folate and its reduced derivatives (e.g., 5-methyltetrahydrofolate, 5-MTHF), SLC19A1 shows a strong preference for the reduced derivatives[2].

The structural mechanisms of folate transport by SLC46A1 and FRs have been

elucidated[17-19]. However, despite its central role in folate uptake among different tissues, the

molecular basis of substrate recognition by SLC19A1 has remained unclear. Here we report the

cryo-EM structures of human SLC19A1 and its complex with 5-MTHF at 3.5-3.6 Å resolution and

demonstrate the critical residues for substrate binding. In particular, we reveal two variant residues

among SLC19 subfamily members, i.e., Arg133 and Gln377 in SLC19A1 *vs.* Glu138 and Met401

in SLC19A2, or Glu120 and Met384 in SLC19A3, being sufficient to designate the specificity for

folate. Moreover, we identify intracellular thiamine pyrophosphate (TPP) as the favorite coupled

substrate for folate transport by SLC19A1. These results have established the key mechanism of

substrate recognition by SLC19A1.

Results

Structures of human SLC19A1 and its complex with 5-MTHF

Human SLC19A1 has 591 residues with a molecular weight of 65 kDa and is mainly

composed of 12 transmembrane helices (TMs). As a result, the majority of the protein would be

embedded in detergent micelles without obvious features when extracted from the cell membrane,

limiting the structural determination by cryo-EM. To overcome this issue, we exploited the

BRIL/Fab/Nb module[20-22], which helped provide the apparent shape for particle alignment. The

N-terminal 23 residues preceding the TM1 of SLC19A1 were replaced by the BRIL domain (Fig.

1a). Importantly, wild-type SLC19A1 or BRIL-SLC19A1 overexpressed in HEK293F cells

exhibited comparable transport activity for a standard substrate [³H]-radiolabeled methotrexate

([³H]-MTX)[23], indicating that the BRIL-tag would not affect the normal function of SLC19A1

(Fig. 1b and Supplementary Fig. S1a).

We purified the BRIL-SLC19A1 protein and then added anti-BRIL Fab and anti-Fab

nanobody (Nb) to assemble the BRIL-SLC19A1/Fab/Nb ternary complex (Fig. 1c). The cryo-EM

map of the ternary complex was collected and successfully reconstructed to 3.6-Å resolution (Fig.

1d, and Supplementary Fig. S2 and Table S1). In the structure of BRIL-SLC19A1, the last helix

of BRIL rotated around its joint region with the TM1 of SLC19A1, and as a result, the four-helical

bundle of BRIL resided parallel to the cell membrane. In addition, BRIL leaned on the intracellular

loop between TM4 and TM5 of SLC19A1 via the electrostatic interactions between three acidic

residues of BRIL (Glu4, Asp5, and Glu8) and two basic residues (Arg145 and Arg148) of

SLC19A1 (Fig. 1e), stabilizing the current conformation of BRIL-SLC19A1. SLC19A1 adopted

the classical major facilitator superfamily (MFS) fold[24,25] with two discrete TM bundles (TM1-6

and TM7-12) (Fig. 1f, g), and all the TM regions were clearly resolved in the cryo-EM structure.

SLC19A1 was present in the inward-facing conformation, i.e., the intracellular gate between TM4-

5 and TM10-11 was open while the extracellular gate was closed by the regions of TM1, 2, and 7

(Fig. 1g). A notable feature was that a segment of TM1 (Ile41-Phe47) was unwound in the

extracellular leaflet of the membrane (Fig. 1g). It has been documented that the discontinuity of

transmembrane helices could play pivotal roles in transporters and ion channels by creating

substrate-binding sites or providing flexible gating hinges[26-30]. On the other hand, the EM densities

of the intracellular loop between TM6 and TM7 (residues 214-249) and the C-terminal cytoplasmic

region (residues 452-591) of SLC19A1 were invisible (Fig. 1f, g), implicating their high motility

and in line with their dispensable role for the transporter function[11].

In parallel to the strategy of the BRIL module, we also identified one nanobody against

SLC19A1 from a synthetic yeast-display library[31]. Using the nanobody-based legobody strategy[32],

we determined the cryo-EM structure of the SLC19A1/legobody complex to a medium resolution

(~5 Å). The nanobody bound the TM6-7 linker region of SLC19A1, and the overall structure of

the nanobody-bound SLC19A1 was almost identical to that observed in BRIL-SLC19A1

(Supplementary Fig. S3a). Given its higher resolution, the structure of BRIL-SLC19A1 (hereafter

referred to as SLC19A1) was pursued in our further studies.

We next determined the cryo-EM structure of SLC19A1 in complex with its preferred

substrate 5-MTHF at 3.5-Å resolution (Fig. 2a, and Supplementary Fig. S4 and Table S1). Similar

to the apo-structure of SLC19A1, the SLC19A1/5-MTHF complex was in the inward-facing

conformation (Fig. 2b). The 5-MTHF binding did not induce a significant conformational change

of SLC19A1, as the root-mean-square deviation (RMSD) between the apo- and 5-MTHF-bound

structures was 1.2 Å (Fig. 2c). Though the EM densities of the glutamate moiety of 5-MTHF were

unresolved in the complex structure, the assignment of the substrate was unambiguously achieved.

5-MTHF resided inside the central cavity of SLC19A1 in the perpendicular position to the cell

membrane, with the pterin ring pointing to the extracellular gate (Fig. 2b).

Mechanism of substrate recognition by human SLC19A1

In the structure of the SLC19A1/5-MTHF complex, the substrate-binding site was

predominantly formed by TM1, 4, 7, and 10 of SLC19A1 (Fig. 2b). Notably, the electrostatic

distribution of this binding site matched the charge characteristics of 5-MTHF (Fig. 2d), i.e., the

polar pterin ring of 5-MTHF was wrapped in a negatively-charged pocket, and the glutamate

moiety of 5-MTHF was located in a positively-charged environment.

We examined the critical residues involved in the substrate recognition of SLC19A1. On

the extracellular side, the pterin ring of 5-MTHF formed hydrogen bonds with the side chains of

Glu123 and Thr49 (Fig. 2e, f). At the same time, the pterin and *p*-aminobenzoyl groups of 5-MTHF

were in close contact with an array of residues through van der Waals and hydrophobic interactions,

including Glu45, Ile48, Ile68, Thr69, Leu72, Tyr126, Met130, Val285, Tyr286, and His289. It is

worth noting that the extra methyl group on N5 nitrogen atom of 5-MTHF could enhance the

hydrophobic interactions with SLC19A1 (Fig. 2a, e, and f), thus making 5-MTHF (K_t of 1-7 μM)

a better substrate compared to folate (K_t of ~200 μM)[10]. On the intracellular side, the negatively-

charged glutamate moiety of 5-MTHF was accommodated by two arginine residues (Arg133 and

Arg373). Additionally, Tyr281 and Gln377 also participated in polar interactions with the

glutamate moiety (Fig. 2e, f).

To verify the functional relevance of those residues involved in the substrate binding of

SLC19A1, we mutated them individually and tested their effects on the transport activity in

HEK293F cells (Fig. 2g and Supplementary Fig. S1b). Substitutions of Glu123, Arg133, Tyr281,

and Arg373 with alanine abolished the transport activity of SLC19A1, consistent with their

participation in the polar or electrostatic interactions with 5-MTHF. In comparison, the Q377A

mutation only reduced the transport activity by ~40%, indicating that its polar interaction with 5-

MTHF was less critical than that of the four residues above. In addition, T49A, I68A, T69A, L72A,

Y126A, V285A, Y286A, or H289A mutation attenuated the transport activity by 20~50%,

suggesting that their hydrophobic stacking with the pterin or *p*-aminobenzoyl ring would also

contribute to substrate binding. Of importance, these key residues of SLC19A1 involved in the

substrate recognition are mostly conserved among different species (Supplementary Fig. S5). In

contrast, Ile48 and Met130 had minor roles as their mutations barely affect the function of

SLC19A1. Intriguingly, mutation of Glu45 to alanine enhanced the transport activity by ~70%.

This E45A mutation might better stabilize the unique loop structure of TM1 around the substrate-

binding site (Supplementary Fig. S3b). In support of this notion, mutations of the residues adjacent

to Glu45, e.g., G44R and S46I, have been identified in antifolate-drug resistant leukemia cells[33,34].

SLC19 subfamily contains three members, i.e., SLC19A1, SLC19A2, and SLC19A3.

Although sharing over 40% sequence identity, they engage different substrates, i.e., SLC19A1

transports folate, whereas SLC19A2 and SLC19A3 transport thiamine (vitamin B_1)[10]. We

investigated the mechanism designating the folate specificity of SLC19A1. By primary and ternary

structural alignments, five out of the twelve residues of SLC19A1 comprising the binding site

around the pterin and *p*-aminobenzoyl groups of 5-MTHF are not conserved in SLC19A2 and

SLC19A3, i.e., Thr69, Leu72, Met130, Tyr286, and His289 (Fig. 3a-c). However, mutations of

these residues to their cognates in SLC19A2 or SLC19A3 only slightly affected the transport

activity of SLC19A1 (Fig. 3d and Supplementary Fig. S1c). We thus focused on the residues of

SLC19A1 accommodating the negatively-charged glutamate moiety of 5-MTHF, i.e., Arg133,

Tyr281, Arg373, and Gln377. While Tyr281 and Arg373 are conserved in SLC19A2 and

SLC19A3, Arg133 and Gln377 become glutamate and methionine residues in SLC19A2 (Glu138

and Met401) and SLC19A3 (Glu120 and Met384), respectively (Fig. 3a-c). Replacing either of

these two residues with their cognates in SLC19A2 and SLC19A3 (R133E or Q377M) completely

abolished the function of SLC19A1 (Fig. 3d and Supplementary Fig. S1c). In addition, substituting

the alanine residue adjacent to Arg133 to proline, i.e., A132P, caused the malfunction of

SLC19A1[35,36], elucidating the geometry restriction at Arg133. These results have suggested that

these two variant residues among SLC19 subfamily members would be sufficient to determine the

substrate specificity, with the negatively-charged glutamate and hydrophobic methionine residues

of SLC19A2 and SLC19A3 precluding folate via the electrostatic or nonpolar repulsion on its

glutamate moiety.

TPP is the favorite coupled substrate of SLC19A1

Although SLC19A2 and SLC19A3 both transport thiamine but not folate, most of the

residues that comprise the substrate-binding pocket of SLC19A1, particularly those surrounding

the pterin and *p*-aminobenzoyl groups of 5-MTHF, are highly conserved in SLC19A2 and

SLC19A3 (Fig. 3a). Also, the non-conserved residues appeared to have a minor role in the

transport function of SLC19A1 (Fig. 3d). These observations raised a tempting possibility that

SLC19A1 might recognize specific types of thiamine derivatives as its substrate. Notably,

SLC19A1 functions as an antiporter, i.e., coupling folate intake with the transport of another

substrate in the opposite direction. In fact, a variety of organic phosphate anions were reported to

be such coupled substrates of SLC19A1, including TPP, ATP (adenosine triphosphate), ADP

(adenosine diphosphate), AMP (adenosine monophosphate), G6P (glucose 6-phosphate), and

NAD^+ (nicotinamide adenine dinucleotide)[23,37,38]. It thus came to our attention that the majority of

thiamine would be metabolized in cells to its active form, the organic-phosphate derivative TPP

(Fig. 4a), which is the coenzyme involved in biochemical reactions of decarboxylation[39].

We then pursued the cryo-EM structure of the SLC19A1/TPP complex to 3.7-Å resolution

(Supplementary Fig. S6 and Table S1). Strikingly, TPP could be clearly detected in the same

substrate-binding site as in the SLC19A1/5-MTHF structure (Fig. 4b, c). There was no significant

conformational difference between the TPP-bound and 5-MTHF-bound SLC19A1, as the RMSD

of the two structures was 1.3 Å (Fig. 4d). TPP interacted with SLC19A1 in a manner highly

reminiscent of that in 5-MTHF (Fig. 4e, f). On the extracellular side, the pyrimidine ring of TPP

formed hydrogen bonds with the side chain of Glu123 and the main-chain carboxyl of Tyr286. In

addition, the pyrimidine and thiazolium rings of TPP were stabilized through van der Waals and

hydrophobic interactions by the similar collection of residues that interacted with the pterin and *p*-

aminobenzoyl groups of 5-MTHF as described above. On the intracellular side, the negatively-

charged pyrophosphate moiety of TPP was clamped by the same positively-charged Arg133 and

Arg373, as well as the polar interaction with the hydroxyl of Tyr281. Notably, the side chain of

Arg133 assumed the unique conformation for a better adaption of TPP (Fig. 4g and Supplementary

Fig. S3c).

To validate TPP as an authentic substrate of SLC19A1, we tested its ability to compete

with the uptake of [³H]-MTX. As the positive control, 5-MTHF, folate, and MTX all effectively

blocked the SLC19A1-mediated intake of [³H]-MTX at 200 μM concentration. Importantly, TPP

exhibited a comparable inhibitory effect with IC_{50} of 19 μM (Fig. 4h, i). In contrast, ATP, ADP,

AMP, G6P, and NAD^+ showed much weaker or no effect on the [³H]-MTX transport (Fig. 4h). In

parallel, we measured the thermostability of SLC19A1 in the presence of different compounds

(Fig. 4j). As expected, the well-documented substrates 5-MTHF and folate significantly enhanced

the thermostability of SLC19A1. Among the examined organic phosphate compounds, only TPP

could elevate the thermostability of SLC19A1. Furthermore, microscale thermophoresis (MST)

analysis confirmed the binding of TPP to SLC19A1 (dissociation constant $K_d = 327 \pm 76$ μM) (Fig.

4k). All these results together supported that TPP is the favorite coupled substrate of SLC19A1.

Discussion

SLC19A1 is the first identified folate transporter ubiquitously expressed in tissues and

responsible for folate uptake in most types of mammalian cells[9]. Its action is coupled with the

counter-transport of organic phosphate anions[40]. Although multiple cellular metabolites such as

ATP, ADP, AMP, and NAD^+ have been documented as the coupled substrates of SLC19A1, we

showed with the structural and functional analyses that TPP would be the favorite compared to

those commonly recognized ones. According to the alternative access mechanism, SLC19A1

would cycle between the inward-facing and outward-facing conformations to carry its substrates

across the cell membrane[41]. In our current structures, the extracellular (5-MTHF) and intracellular

(TPP) substrates are bound to the identical site in SLC19A1, similar to that observed in some other

antiporters[42,43]. It is plausible that cytosolic TPP could liberate 5-MTHF from the inward-facing

SLC19A1 through competition under physiological conditions, and SLC19A would then adopt the

outward-facing conformation for releasing TPP and binding extracellular 5-MTHF again (Fig. 5).

The complete documentation of such a transport mechanism awaits the future structure of outward-

facing SLC19A1. Folate and TPP are the coenzymes generally characterized for anabolism and

catabolism, respectively. Therefore, the coordinated exchange of these two molecules by

SLC19A1 might represent a novel, intrinsic part of cell metabolic regulation.

Thiamine monophosphate (TMP) might bind to SLC19A1 as a coupled substrate. Indeed,

TMP effectively competed with [^{3}H]-MTX in the transporter assay and enhanced the

thermostability of SLC19A1 (Fig. 4h, j). However, given that TMP is an intermediate of thiamine

metabolism and its intracellular concentration is approximately one to two orders of magnitude

lower than TPP[44,45], it would likely have a limited role in facilitating folate transport. It is also

known that additional organic-phosphate anionic molecules like ZMP (or AICAR, 5-

aminoimidazole 4-carboxamide ribonucleoside) and cGAMP (2'3'-cyclic-GMP-AMP) were

potential substrates of SLC19A1[46-49]. After the submission of our manuscript, two separate studies

reported the structures of SLC19A1 in complexes with some other substrates, including N-

hydroxysuccinimide-conjugated MTX (NHS-MTX), pemetrexed (PMX), and different cyclic

dinucleotides (CDNs)[50,51]. SLC19A1 exhibited the same inward-facing conformation in all of

these structures (Supplementary Fig. S7a). These studies together elucidated the recognition

mechanisms for different substrates by SLC19A1. Specifically, folate (5-MTHF) and antifolate

drugs (MTX and PMX) bound SLC19A1 as a monomer in a deep cavity very close to the

extracellular side, on the contrary, CDNs localized to the broader intracellular entrance of

SLC19A1 in the form of a dimer (Supplementary Fig. S7b). Moreover, the interaction mode

between SLC19A1 and 5-MTHF was very similar in our and Zhang et al.'s studies, even though

the conformations of the pterin ring and glutamate moieties of 5-MTHF showed some differences[51],

indicating these regions might possess some degree of flexibility upon binding to SLC19A1

(Supplementary Fig. S7b, c). However, the antifolate drug MTX was flipped by about 180 degrees

along the axis perpendicular to the cell membrane in Wright et al.'s structure[50], which might be

caused by their NHS-mediated crosslinking (Supplementary Fig. S7d).

SLC46A1 and FRs are often specifically expressed or upregulated in cancer cells[52,53]. Of

importance, the folate-binding sites of these two proteins are distinct from that of SLC19A1

reported here (Supplementary Fig. S8). Such structural divergence would enable the development

of new antifolate drugs that distinguish the different transport systems, thus minimizing potential

adverse effects of cancer chemotherapy on normal non-malignant cells. This work has thus bridged

a gap in the knowledge of the molecular mechanism of folate transport and could have broad

implications for basic biology and translational research.

Materials and methods

Cell cultures

HEK293S GnTI⁻ cells were cultured in Yocon HEK293 medium (Yocon Biotechnology)

supplemented with 1% FBS and 1% Antibiotic-Antimycotic (Gibco). HEK293F cells were

cultured in SMM 293-TI medium (Sino Biological Inc). All the mammalian cells were cultured at

37 °C with 5% CO_2. Sf9 cells were cultured in Sf-900 II SFM medium (Gibco) at 27 °C.

Expression and purification of BRIL-SLC19A1

The codon-optimized complementary DNA sequence of BRIL-SLC19A1 was cloned into

a pEG BacMam expression vector with 10×His-tag and green fluorescent protein (GFP) attached

to the N-terminus[54]. The plasmid was transformed into DH10Bac *Escherichia coli* cells for bacmid

generation, and then the bacmid was transfected into sf9 cells using Cellfectin II reagents (Life

Technologies) to produce recombinant baculoviruses. For protein expression, 10% of passage 3

(P3) baculoviruses were added to HEK293S GnTI⁻ cells at a density of $3×10^6$ cells/ml. After

culturing at 37 °C for 12 hours, 10 mM sodium butyrate was added to boost the protein expression

and the cells were transfered to 30 °C. Cells were harvested 48 hours later by centrifugation at

6,000 rpm for 20 min.

For protein purification, the cells were first dispersed by a hand-held homogenizer in lysis

buffer (50 mM HEPES pH 7.4, 300 mM NaCl, and 15% glycerol) supplemented with protease

inhibitors (1 µg/ml aprotinin, 1 µg/ml leupeptin, 1 µg/ml pepstatin, 20 µg/ml trypsin inhibitor, 1

mM benzamidine, and 1 mM phenylmethylsulfonyl fluoride) and DNase (2 µg/ml), and then lysed

by addition of 1% n-dodecyl β-D-maltoside (DDM) and 0.1% Cholesteryl hemisuccinate (CHS)

at 4 °C for 2 hours. Cell debris was removed by centrifugation at 18,000 rpm for 40 min. The

soluble fraction was mixed with pre-equilibrated anti-GFP nanobody (GFPnb)-coupled cyanogen

bromide-activated Sepharose beads (GE Healthcare) at 4 °C for 2 hours. The beads were

subsequently washed with 25 column volumes of Buffer A (25 mM HEPES pH 7.5, 150 mM NaCl,

and 0.02% DDM-0.002% CHS) and then incubated with PreScission protease (5:1 w/w ratio) at

4 °C overnight to release the target protein. The PreScission protease was removed by incubation

with Glutathione Sepharose beads (GE Healthcare) at 4 °C for 1 hour. The protein was further

purified by size exclusion chromatography (SEC) using a Superose 6 Increase 10/300 GL column

(GE Healthcare) equilibrated with Buffer A. The protein samples of the peak fractions were

collected and concentrated to 6 mg/ml using a 100 kDa molecular weight cut-off concentrator

(Millipore).

Expression and purification of anti-BRIL Fab

DNA sequences of the heavy and light chains of anti-BRIL Fab[20] were cloned into pEG

BacMam expression vectors separately. A GFP-tag was attached to the C-terminus of the heavy

chain. The bacmids and baculoviruses were prepared in the same way as that of BRIL-SLC19A1.

HEK293F cells at a density of 3×10^6 cells/ml were infected with 10% P3 baculoviruses of heavy

and light chains (5% of each). 12 hours after infection, the cells were induced with 10 mM sodium

butyrate and transferred to 30°C for protein expression.

For Fab purification, the cell culture media was centrifuged at 6,000 rpm for 30 min, and

then the supernatant was concentrated and exchanged into Buffer B (25 mM HEPES pH 7.5 and

150 mM NaCl) using a Hydrosart Ultrafilter system (Sartorius). The following purification steps

for anti-BRIL Fab were the same as that of BRIL-SLC19A1. Briefly, after the anti-GFP affinity

chromatography, SEC was applied for further purification.

Expression and purification of anti-Fab Nanobody

The DNA sequence of anti-Fab Nanobody was cloned into pET24 (+) vector which bears an N-terminal PelB signal sequence[55] and a C-terminal 6×His-tag. The nanobody would be expressed and translocated to the periplasm of *Escherichia coli* strain BL21. The bacteria were cultured in LB medium at 37 °C until OD600 reached about 0.8. Then the protein expression was induced by addition of 1 mM isopropyl-β-D-1-thiogalactopyranoside (IPTG) for 6 hours.

For nanobody purification, the cell pellets were lysed by sonication in lysis buffer. Cell debris was removed by centrifuged at 18,000 rpm for 1 hour. Next, the supernatant was mixed with pre-equilibrated Ni-NTA beads (Smart-Lifesciences) and incubated at 4 °C for 1 hour. The beads were sequentially washed with 20 column volumes of Buffer B containing 25 mM and 50 mM imidazole, and then the target protein was eluted with 3 column volumes of Buffer C (25 mM HEPES pH 7.5, 150 mM NaCl, and 300 mM imidazole). The protein was further purified by SEC using a Superdex 200 Increase 10/300 column (GE Healthcare) equilibrated with Buffer B. The protein samples of the peak fractions were collected and concentrated to 5 mg/ml.

The BRIL-SLC19A1/Fab/Nb complex assembly

For complex assembly, the purified BRIL-SLC19A1, anti-BRIL Fab, and anti-BRIL Fab nanobody (Nb) were incubated on ice for 1 hour at the ratio of 1:1.4:2. Then the samples were applied to a Superose 6 Increase 10/300 GL column equilibrated with Buffer A to remove the excess Fab and Nb components. The peak fractions corresponding to the ternary complex were collected and concentrated to 6 mg/ml.

Cryo-EM sample preparation and data collection

Quantifoil R1.2/1.3 Au400 holey carbon grids were glow-discharged for 1 min. 3 μl protein samples were deposited on the grids and blotted for 3 s with filter paper at 10 °C and 100% humidity using a Vitrobot Mark IV (FEI) equipment. The grids were flash frozen in liquid ethane and stored in liquid nitrogen until further use. To prepare the substrate-bound SLC19A1 samples, 5 mM 5-MTHF or TPP was incubated with the BRIL-SLC19A1/Fab/Nb complex on ice for 1 hour prior to grid freezing.

The grids were loaded into a 300 kV Titan Krios electron microscope (FEI) with K3 Summit direct electron detector (Gatan). Data were collected in super-resolution mode at a magnification of ×81,000 using the EPU software (Thermo Fisher Scientific). The physical pixel size was 1.07 Å, and the defocus range was 0.7 to 1.5 μm. The exposure time for each micrograph (40 frames) was 3.2 s, yielding a total dose of 60 e$^-$/Å^2. The numbers of micrographs collected for apo, 5-MTHF-, and TPP-bound SLC19A1 samples were 10,485, 7,382, and 14,262, respectively.

Cryo-EM data processing

For all the three datasets, the micrographs were motion corrected using MotionCor2[56] and binned to a pixel size of 2.675 Å. The images with ice contamination were manually removed. Contrast transfer function (CTF) was estimated using Gctf[57]. Particle picking was carried out using Gautomatch (http://www.mrc-lmb.cam.ac.uk/kzhang). Totally 3,104,540 (apo), 1,802,524 (5-MTHF-bound), and 4,144,030 (TPP-bound) particles were automatically picked for the three datasets. After two rounds of 2D classification, the good subclasses were chosen and subjected to the following 3D classification in RELION 3.1[58]. For all the three datasets, one subclass of particles with the highest resolution was re-extracted at the pixel size of 1.07 Å and applied to 3D

refinement. The particles and maps of the apo and 5-MTHF-bound datasets were further imported

into cryoSPARC for Non-uniform (NU) refinement[59]. The particle numbers used for the final

refinement of these three datasets were 268,975 (apo), 206,075 (5-MTHF-bound), and 229,158

(TPP-bound), respectively. The resolutions of the final maps were 3.6 Å for apo SLC19A1, 3.5 Å

for 5-MTHF-bound SLC19A1, and 3.7 Å for TPP-bound SLC19A1. All the resolutions reported

here were calculated using the 0.143 cutoff criterion. The post-processed maps were generated

using DeepEMhancer[60].

Model building and refinement

The AlphaFold-predicted model of SLC19A1[61], and the reported structures of anti-BRIL

Fab and anti-BRIL Fab nanobody[20] (PDB code: 6WW2) were used for our model building. All

these models were roughly fitted into the SLC19A1 maps using ChimeraX[62], and then manually

adjusted using Coot[63]. Refinement of the final structures in real space was done by PHENIX[64].

Geometries of the structure models were validated using MolProbity[65]. The Fourier shell

correlation (FSC) curves were calculated between the refined models and full maps using PHENIX.

Local resolutions were estimated in cryoSPARC. All the figures were prepared using ChimeraX.

[³H]-methotrexate uptake assay

5 µg plasmid of wild-type or mutant SLC19A1 with a C-terminal GFP tag was transiently

transfected into 5 ml HEK293F cells at a density of 2-3×10^6 cells/ml using PEI (BIOHUB). After

culturing at 37 °C for 12 hours, 10 mM sodium butyrate was added and the cells were transferred

to 30 °C for protein expression. Cells were harvested after 24-30 hours by centrifugation at 4,000

rpm for 5 min. The cells were washed with PBS and resuspended in MHS buffer (20 mM HEPES,

and 225 mM sucrose, pH adjusted to 7.4 with MgO) at a concentration of 1.5×10^7 cells/ml. 10 µl

cells were taken out and used for protein expression detection. Next, 100 µl cells were mixed with

equal volume of MHS buffer containing 17 nM nonradioactive methotrexate (MTX) and 1.25 pM

[^{3}H]-methotrexate (0.05 µCi, American Radiolabeled Chemicals, Inc) at 37 °C for 5 min. To

analyze the inhibitory effect of TPP, the corresponding molecules were added simultaneously at

the indicated concentrations. 800 µl ice cold PBS was added to terminate the MTX transport, and

the cells were further washed three times with PBS to remove the extracellular radioactive

substrate. Then the cells were lysed with 500 µl 1% Triton X-100 at room temperature for 30 min.

400 µl cell lysate was mixed with 250 µl scintillation fluid, and the [^{3}H] radioactive signal was

measured using a scintillation counter (PerkinElmer).

To check the protein expression level of SLC19A1, the cells were lysed with RAPI lysis

buffer (25 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% NP-40, and 0.1% SDS) at room

temperature for 60 min and mixed with 4×SDS-PAGE loading buffer. The samples were separated

using 10 % SDS-PAGE gels. The GFP fluorescence signal of SLC19A1 was directly detected

using a gel image system (Tanon).

Fluorescence-detection size-exclusion chromatography (FSEC)-based thermostability assay

GFP-tagged SLC19A1 expressing cells were lysed with 1% DDM and 0.1% CHS in

exactly the same way as that of the BRIL-SLC19A1 purification. After removing the cell debris

by centrifugation, the supernatant was incubated with different small molecules on ice for 1 hour.

The samples were then treated at 60 °C for 10 min, and subsequently centrifuged at 15,000 rpm

for 1 hour to remove the protein aggregates. FSEC was performed using a Shimadzu HPLC system.

SLC19A1 protein was separated by a Superose 6 Increase 10/300 GL column during which process

the GFP fluorescence signal was monitored. The integral areas of the corresponding fluorescence

peak were calculated to represent the amount of SLC19A1 protein. Samples without heat treatment

were used as control to evaluate the protein thermostability.

MST analysis

The affinity of SLC19A1 for TPP was determined using a NanoTemper Monolith NT.115

instrument (NanoTemper Technologies). The full-length SLC19A1 protein was fluorescently

labelled with a Red-NHS 2nd Generation kit (NanoTemper Technologies) in Buffer D (25 mM

HEPES pH 7.5, 150 mM NaCl, and 0.002% LMNG). 50 nM SLC19A1 was mixed with TPP

prepared in 16 different serial concentrations (0.25 μM to 8 mM) in Buffer D. Then the mixture

was incubated overnight at 16 °C followed by centrifugation at 15,000 g for 15 min before loaded

into the Premium Capillaries (NanoTemper Technologies). The thermophoresis analysis,

including six to ten replicate measurements, was performed with 20% excitation power and 60%

MST power. The K_d value was calculated using the MO.Affinity Analysis software (NanoTemper

Technologies).

Data availability

Cryo-EM density maps of the apo, 5-MTHF-bound, and TPP-bound SLC19A1 have been

deposited in the Electron Microscopy Data Bank under the accession codes EMD-34817, EMD-

34818, and EMD-34819. Their atomic coordinates have been deposited in the Protein Data Bank

under accession codes 8HII, 8HIJ, and 8HIK.

Acknowledgements

We thank the Cryo-EM platform and the School of Life Sciences of PKU for cryo-electron

microscopy data collection. We are grateful to Dr. Guopeng Wang, Bo Shao, Xia Pei, and Dr. Ning

Gao for their help in cryo-EM experiments. We thank Shitang Huang at the National Center for

Protein Sciences of Peking University for assistance with radioactive experiments. We thank

Jinkun Xu in Long Li's lab for assistance with the nanobody screening. We thank Qing Chang and

Wenqi Li at Beijing Advanced Innovation Center for Structural Biology of Tsinghua University

for assistance with our pilot experiments to verify the protein-small molecule interactions. This

research was funded by the National Key Research and Development Program of China (to Z.Z.,

NO. 2021YFA1302300) and the National Natural Science Foundation of China (to Z.Z., NO.

32171201). The study was also supported by Center for Life Sciences (CLS), School of Life

Sciences (SLS) of Peking University, the SLS-Qidong innovation fund, Li Ge-Zhao Ning Life

Science Youth Research Foundation, and the State Key Laboratory of Membrane Biology of

China.

Conflict of interests: The authors declare no competing interests.

Author contributions: Y.D., X.D., and H.Z. did the cloning and protein purification. X.D.

screened nanobodies against SLC19A1. Y.D., C.Q., and Z.G. collected the cryo-EM data. Y.D.

and Z.Z. processed the EM data. Z.Z. built and refined the structural models. Y.D. and D.Z.

performed the MTX uptake experiments. Y.D. carried out the thermostability assay. Y.D. and

Y.W. did the MST analysis. C.-H.L. assisted in model building and structural analysis. Z.Z. wrote

the manuscript with input and support from all co-authors. J.Y. revised the manuscript.

References

1. Bailey, L.B. et al. Biomarkers of Nutrition for Development-Folate Review. *J. Nutr.* **145**,
1636S-1680S (2015).
2. Alam, C., Kondo, M., O'Connor, D.L. & Bendayan, R. Clinical Implications of Folate
Transport in the Central Nervous System. *Trends Pharmacol. Sci.* **41**, 349-361 (2020).
3. Coppede, F. The genetics of folate metabolism and maternal risk of birth of a child with
Down syndrome and associated congenital heart defects. *Front. Genet.* **6**, 223 (2015).
4. Qiu, A. et al. Identification of an intestinal folate transporter and the molecular basis for
hereditary folate malabsorption. *Cell* **127**, 917-28 (2006).
5. Henderson, G.B. Folate-binding proteins. *Annu. Rev. Nutr.* **10**, 319-35 (1990).
6. Kelemen, L.E. The role of folate receptor alpha in cancer development, progression and
treatment: cause, consequence or innocent bystander? *Int. J. Cancer* **119**, 243-50 (2006).
7. Kamen, B.A., Wang, M.T., Streckfuss, A.J., Peryea, X. & Anderson, R.G. Delivery of
folates to the cytoplasm of MA104 cells is mediated by a surface membrane receptor that
recycles. *J. Biol. Chem.* **263**, 13602-9 (1988).
8. Dixon, K.H., Lanpher, B.C., Chiu, J., Kelley, K. & Cowan, K.H. A novel cDNA restores
reduced folate carrier activity and methotrexate sensitivity to transport deficient cells. *J.*
Biol. Chem. **269**, 17-20 (1994).
9. Goldman, I.D., Lichtenstein, N.S. & Oliverio, V.T. Carrier-mediated transport of the folic
acid analogue, methotrexate, in the L1210 leukemia cell. *J. Biol. Chem.* **243**, 5007-17
(1968).
10. Zhao, R. & Goldman, I.D. Folate and thiamine transporters mediated by facilitative carriers
(SLC19A1-3 and SLC46A1) and folate receptors. *Mol. Aspects Med.* **34**, 373-85 (2013).
11. Matherly, L.H., Wilson, M.R. & Hou, Z. The major facilitative folate transporters solute
carrier 19A1 and solute carrier 46A1: biology and role in antifolate chemotherapy of
cancer. *Drug Metab. Dispos.* **42**, 632-49 (2014).
12. Rothberg, K.G., Ying, Y.S., Kolhouse, J.F., Kamen, B.A. & Anderson, R.G. The
glycophospholipid-linked folate receptor internalizes folate without entering the clathrin-
coated pit endocytic pathway. *J. Cell Biol.* **110**, 637-49 (1990).
13. Zhao, R., Matherly, L.H. & Goldman, I.D. Membrane transporters and folate homeostasis:
intestinal absorption and transport into systemic compartments and tissues. *Expert Rev.*
Mol. Med. **11**, e4 (2009).
14. Matherly, L.H., Hou, Z. & Deng, Y. Human reduced folate carrier: translation of basic
biology to cancer etiology and therapy. *Cancer Metastasis Rev.* **26**, 111-28 (2007).
15. Gonen, N. & Assaraf, Y.G. Antifolates in cancer therapy: structure, activity and
mechanisms of drug resistance. *Drug Resist. Updat.* **15**, 183-210 (2012).
16. Zhao, R. & Goldman, I.D. Resistance to antifolates. *Oncogene* **22**, 7431-57 (2003).
17. Parker, J.L. et al. Structural basis of antifolate recognition and transport by PCFT. *Nature*
595, 130-134 (2021).
18. Chen, C. et al. Structural basis for molecular recognition of folic acid by folate receptors.
Nature **500**, 486-9 (2013).
19. Wibowo, A.S. et al. Structures of human folate receptors reveal biological trafficking states
and diversity in folate and antifolate recognition. *Proc. Natl. Acad. Sci. USA* **110**, 15180-8
(2013).

20.   Tsutsumi, N. et al. Structure of human Frizzled5 by fiducial-assisted cryo-EM supports a
485   heterodimeric mechanism of canonical Wnt signaling. *Elife* **9**(2020).

486   21.   Mukherjee, S. et al. Synthetic antibodies against BRIL as universal fiducial marks for
487   single-particle cryoEM structure determination of membrane proteins. *Nat. Commun.* **11**,
488   1598 (2020).

489   22.   Ereno-Orbea, J. et al. Structural Basis of Enhanced Crystallizability Induced by a
490   Molecular Chaperone for Antibody Antigen-Binding Fragments. *J. Mol. Biol.* **430**, 322-
491   336 (2018).

492   23.   Goldman, I.D. The characteristics of the membrane transport of amethopterin and the
493   naturally occurring folates. *Ann. N. Y. Acad. Sci.* **186**, 400-22 (1971).

494   24.   Yan, N. Structural Biology of the Major Facilitator Superfamily Transporters. *Annu. Rev.*
495   *Biophys.* **44**, 257-83 (2015).

496   25.   Bai, X., Moraes, T.F. & Reithmeier, R.A.F. Structural biology of solute carrier (SLC)
497   membrane transport proteins. *Mol. Membr. Biol.* **34**, 1-32 (2017).

498   26.   Doyle, D.A. et al. The structure of the potassium channel: molecular basis of K+
499   conduction and selectivity. *Science* **280**, 69-77 (1998).

500   27.   Dutzler, R., Campbell, E.B., Cadene, M., Chait, B.T. & MacKinnon, R. X-ray structure of
501   a ClC chloride channel at 3.0 A reveals the molecular basis of anion selectivity. *Nature*
502   **415**, 287-94 (2002).

503   28.   Feng, L., Campbell, E.B., Hsiung, Y. & MacKinnon, R. Structure of a eukaryotic CLC
504   transporter defines an intermediate state in the transport cycle. *Science* **330**, 635-41 (2010).

505   29.   Zhang, Z. & Chen, J. Atomic Structure of the Cystic Fibrosis Transmembrane Conductance
506   Regulator. *Cell* **167**, 1586-1597 e9 (2016).

507   30.   Screpanti, E. & Hunte, C. Discontinuous membrane helices in transport proteins and their
508   correlation with function. *J. Struct. Biol.* **159**, 261-7 (2007).

509   31.   McMahon, C. et al. Yeast surface display platform for rapid discovery of conformationally
510   selective nanobodies. *Nat. Struct. Mol. Biol.* **25**, 289-296 (2018).

511   32.   Wu, X. & Rapoport, T.A. Cryo-EM structure determination of small proteins by nanobody-
512   binding scaffolds (Legobodies). *Proc. Natl. Acad. Sci. USA* **118**(2021).

513   33.   Wong, S.C. et al. Impaired membrane transport in methotrexate-resistant CCRF-CEM cells
514   involves early translation termination and increased turnover of a mutant reduced folate
515   carrier. *J. Biol. Chem.* **274**, 10388-94 (1999).

516   34.   Drori, S., Jansen, G., Mauritz, R., Peters, G.J. & Assaraf, Y.G. Clustering of mutations in
517   the first transmembrane domain of the human reduced folate carrier in GW1843U89-
518   resistant leukemia cells with impaired antifolate transport and augmented folate uptake. *J.*
519   *Biol. Chem.* **275**, 30855-63 (2000).

520   35.   Schuetz, J.D. et al. Membrane protein changes in an L1210 leukemia cell line with a
521   translocation defect in the methotrexate-tetrahydrofolate cofactor transport carrier. *J. Biol.*
522   *Chem.* **264**, 16261-7 (1989).

523   36.   Brigle, K.E., Spinella, M.J., Sierra, E.E. & Goldman, I.D. Characterization of a mutation
524   in the reduced folate carrier in a transport defective L1210 murine leukemia cell line. *J.*
525   *Biol. Chem.* **270**, 22974-9 (1995).

526   37.   Henderson, G.B. & Zevely, E.M. Structural requirements for anion substrates of the
527   methotrexate transport system in L1210 cells. *Arch. Biochem. Biophys.* **221**, 438-46 (1983).

528   38.   Zhao, R. et al. Impact of the reduced folate carrier on the accumulation of active thiamin
529   metabolites in murine leukemia cells. *J. Biol. Chem.* **276**, 1114-8 (2001).

530   39.   Lonsdale, D. A review of the biochemistry, metabolism and clinical benefits of thiamin(e)
531   and its derivatives. *Evid. Based Complement Alternat. Med.* **3**, 49-59 (2006).
532   40.   Hou, Z. & Matherly, L.H. Biology of the major facilitative folate transporters SLC19A1
533   and SLC46A1. *Curr. Top. Membr.* **73**, 175-204 (2014).
534   41.   Jardetzky, O. Simple allosteric model for membrane pumps. *Nature* **211**, 969-70 (1966).
535   42.   Huang, Y., Lemieux, M.J., Song, J., Auer, M. & Wang, D.N. Structure and mechanism of
536   the glycerol-3-phosphate transporter from Escherichia coli. *Science* **301**, 616-20 (2003).
537   43.   Fluman, N., Ryan, C.M., Whitelegge, J.P. & Bibi, E. Dissection of mechanistic principles
538   of a secondary multidrug efflux protein. *Mol. Cell* **47**, 777-87 (2012).
539   44.   Tallaksen, C.M., Bohmer, T., Bell, H. & Karlsen, J. Concomitant determination of thiamin
540   and its phosphate esters in human blood and serum by high-performance liquid
541   chromatography. *J. Chromatogr.* **564**, 127-36 (1991).
542   45.   Tallaksen, C.M., Bohmer, T., Karlsen, J. & Bell, H. Determination of thiamin and its
543   phosphate esters in human blood, plasma, and urine. *Methods Enzymol.* **279**, 67-74 (1997).
544   46.   Zhao, R., Gao, F. & Goldman, I.D. Reduced folate carrier transports thiamine
545   monophosphate: an alternative route for thiamine delivery into mammalian cells. *Am. J.*
546   *Physiol. Cell Physiol.* **282**, C1512-7 (2002).
547   47.   Luteijn, R.D. et al. SLC19A1 transports immunoreactive cyclic dinucleotides. *Nature* **573**,
548   434-438 (2019).
549   48.   Ritchie, C., Cordova, A.F., Hess, G.T., Bassik, M.C. & Li, L. SLC19A1 Is an Importer of
550   the Immunotransmitter cGAMP. *Mol. Cell* **75**, 372-381 e5 (2019).
551   49.   Visentin, M., Zhao, R. & Goldman, I.D. Augmentation of reduced folate carrier-mediated
552   folate/antifolate transport through an antiport mechanism with 5-aminoimidazole-4-
553   carboxamide riboside monophosphate. *Mol. Pharmacol.* **82**, 209-16 (2012).
554   50.   Wright, N.J. et al. Methotrexate recognition by the human reduced folate carrier SLC19A1.
555   *Nature* **609**, 1056-1062 (2022).
556   51.   Zhang, Q. et al. Recognition of cyclic dinucleotides and folates by human SLC19A1.
557   *Nature* (2022).
558   52.   Matherly, L.H., Hou, Z. & Gangjee, A. The promise and challenges of exploiting the
559   proton-coupled folate transporter for selective therapeutic targeting of cancer. *Cancer*
560   *Chemother. Pharmacol.* **81**, 1-15 (2018).
561   53.   Assaraf, Y.G., Leamon, C.P. & Reddy, J.A. The folate receptor as a rational therapeutic
562   target for personalized cancer treatment. *Drug Resist. Updat.* **17**, 89-95 (2014).
563   54.   Goehring, A. et al. Screening and large-scale expression of membrane proteins in
564   mammalian cells for structural studies. *Nat. Protoc.* **9**, 2574-85 (2014).
565   55.   Steiner, D., Forrer, P., Stumpp, M.T. & Pluckthun, A. Signal sequences directing
566   cotranslational translocation expand the range of proteins amenable to phage display. *Nat.*
567   *Biotechnol.* **24**, 823-31 (2006).
568   56.   Zheng, S.Q. et al. MotionCor2: anisotropic correction of beam-induced motion for
569   improved cryo-electron microscopy. *Nat. Methods* **14**, 331-332 (2017).
570   57.   Zhang, K. Gctf: Real-time CTF determination and correction. *J. Struct. Biol.* **193**, 1-12
571   (2016).
572   58.   Fernandez-Leiro, R. & Scheres, S.H.W. A pipeline approach to single-particle processing
573   in RELION. *Acta Crystallogr. D Struct. Biol.* **73**, 496-502 (2017).
574   59.   Punjani, A., Rubinstein, J.L., Fleet, D.J. & Brubaker, M.A. cryoSPARC: algorithms for
575   rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290-296 (2017).

60. Sanchez-Garcia, R. et al. DeepEMhancer: a deep learning solution for cryo-EM volume
post-processing. *Commun. Biol.* **4**, 874 (2021).
61. Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021).
62. Pettersen, E.F. et al. UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. *Protein Sci.* **30**, 70-82 (2021).
63. Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr. D Biol. Crystallogr. **66**, 486-501 (2010).
64. Adams, P.D. et al. PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 213-21 (2010).
65. Davis, I.W. et al. MolProbity: all-atom contacts and structure validation for proteins and
nucleic acids. *Nucleic Acids Res.* **35**, W375-83 (2007).
588

 Figure Legends

Fig. 1 Cryo-EM structure of the BRIL-SLC19A1/Fab/Nb complex.

a Schematic diagrams of BRIL-SLC19A1, anti-BRIL Fab, and anti-Fab Nanobody. **b** [^{3}H]-MTX

uptake assay to verify the function of BRIL-SLC19A1. The results are normalized to the activity

of wild-type SLC19A1. All experiments were done in triplicates. (n = 3, mean ± s.d.). ns, non-

significant; **** *P* < 0.0001 (T-test). **c** Profile of size exclusion chromatography (SEC) for the

complex purification and the SDS-PAGE results to show the protein purity. **d** Cryo-EM map of

the BRIL-SLC19A1/Fab/Nb complex. **e** Overall structure of the BRIL-SLC19A1/Fab/Nb complex.

α-helices are shown in cylinders and β-strands are in ribbon. The residues of BRIL and SLC19A1

that are involved in electrostatic interactions are shown with side chains. **f** Cartoon diagram for the

transmembrane (TM) domain of SLC19A1. The TM numbers are labeled, and the plasma

membrane is indicated with dotted lines. **g** Ribbon presentation of the SLC19A1 structure in two

views. Two half TM bundles are colored in blue (TM1-6) and yellow (TM7-12), respectively. A

close-up view of the unwound region of TM1 is shown in an inset.

Fig. 2 Cryo-EM structure of the SLC19A1/5-MTHF complex.

a Chemical structures of several folate analogs, including 5-MTHF (5-methytetrahydrofolate),

folate, and MTX (methotrexate). The constituent moieties of 5-MTHF are indicated. **b** Ribbon

diagram of SLC19A1 in complex with 5-MTHF. The four TMs (TM1, 4, 7, and 10) interacting

with 5-MTHF are colored in cyan and the other ones are colored in blue. 5-MTHF is shown in

sticks and its cryo-EM densities are shown in black meshes. **c** Superposition of the apo and 5-

MTHF-bound SLC19A1 structures. **d** The electrostatic potential (in units of k_BT/e, where k_B is the

Boltzmann constant, *T* is the absolute temperature and *e* is the elementary charge) of the substrate-

binding pocket in SLC19A1, as calculated at pH 7.0 and 0.15 M concentrations of monovalent

cations and anions. 5-MTHF is shown in sticks. **e** Ribbon presentation of the substrate-binding site

in SLC19A1. The residues participating in 5-MTHF binding are indicated with side chains (< 4.0

Å). Hydrogen bonds and salt bridges are depicted as dashed lines. **f** A schematic summary of the

interactions between SLC19A1 and 5-MTHF. The distances of the hydrogen bonds are indicated

in angstroms. **g** The [3H]-MTX uptake activities of SLC19A1 mutants. The results are normalized

to the activity of wild-type SLC19A1. All experiments were done in triplicates (n = 3, mean ± s.d.).

ns, non-significant; * $P < 0.01$; ** $P < 0.005$; *** $P < 0.001$ (T-test).

Fig. 3 Analyses of the substrate discrimination mechanism of SLC19 subfamily members.

a Sequence alignment of the three SLC19 family members. The partially conserved residues are

indicated with blue boxes and the strictly conserved ones are further filled with orange color. The

residues of SLC19A1 that involved in 5-MTHF binding are indicated. The conserved and non-

conserved ones are denoted by black and red arrowheads, respectively. **b-c** Ribbon presentation of

the 5-MTHF binding site in SLC19A2 and SLC19A3. The structures of SLC19A2 and SLC19A3

are predicted by AlphaFold. 5-MTHF is modeled into these structures based on the superposition

with our SLC19A1/5-MTHF structure. The cognates of SLC19A2 and SLC19A3 corresponding

to the 5-MTHF interaction residues of SLC19A1 are indicated. **d** Functional verification of the

non-conserved residues for SLC19A1 using the [3H]-MTX uptake assay. The results are

normalized to the activity of wild-type SLC19A1. All experiments were done in triplicates (n = 3,

mean ± s.d.). ns, non-significant; * $P < 0.01$; ** $P < 0.005$; *** $P < 0.001$ (T-test).

Fig. 4 Verification of TPP as the favorite coupled substrate of SLC19A1.

a The chemical structure of TPP. The constituent moieties are indicated. **b** Ribbon diagram of

SLC19A1 in complex with TPP, TPP is shown in sticks and its cryo-EM densities are shown in

black meshes. **c** The electrostatic potential of the TPP binding site. **d** Structural comparison of the

TPP- and 5-MTHF-bound SLC19A1. **e** Ribbon presentation of the TPP-binding site in SLC19A1.

The residues involved in the interaction with TPP are shown with side chains. **f** A schematic

summary of the interactions between SLC19A1 and TPP. The distances of the hydrogen bonds are

indicated in angstroms. **g** Conformational change of the side chain of Arg133 between apo and

TPP-bound SLC19A1 structures. **h** Inhibitory effect of different compounds on the $[^3H]$-MTX

uptake activity of SLC19A1. All the molecules were tested at the concentration of 200 μM. The

results are normalized to the activity of the control experiment in which no inhibitors are added.

All experiments were done in triplicates (n = 3, mean ± s.d.). ns, non-significant; * P < 0.01; ** P

< 0.005 (T-test). **i** Quantitative measurement of the potency of TPP in inhibiting the $[^3H]$-MTX

delivery by SLC19A1. The results are normalized to the MTX transport activity of SLC19A1 in

the absence of TPP. All experiments were done in triplicates (n = 3, mean ± s.d.). IC_{50} was

calculated by fitting to a nonlinear regression model. **j** Quantification of the fluorescence-detection

size-exclusion chromatography (FSEC)-based thermostability assay. The concentrations of

molecules are indicated. The thermostability is calculated relative to the control experiment in

which SLC19A1 was not incubated with any compounds. All experiments were done in triplicates

(n = 3, mean ± s.d.). ns, non-significant; * *P* < 0.01 (T-test). **k** MST analysis to examine the affinity

of SLC19A1 for TPP. All experiments were done in multiple replicates (n = 6-10, mean ± s.d.).

Fig. 5 Model of the substrate transport cycle of SLC19A1.

SLC19A1 utilizes the alternating access mechanism to reverse transport two substrates. Under

physiological conditions, 5-MTHF and TPP are likely the favorite extracellular and intracellular

substrates of SLC19A1, respectively. They compete for the same binding site within the central

cavity of SLC19A1. The four key residues (Glu123, Arg133, Tyr281, and Arg373) for substrate

recognition are indicated.

a

BRIL | SLC19A1
24   591

Fab   Anti-BRIL Fab

Nb   Anti-Fab Nanobody

b

Relative [³H]-MTX uptake

ns

****

SLC19A1 BRIL-SLC19A1 Control

c

mAU

100 kDa
70 kDa
50 kDa   BRIL-SLC19A1

25 kDa   Fab
20 kDa   Nb

BRIL-SLC19A1/Fab/Nb

Fab/Nb

0   5   10   15   20

Volume (ml)

d

Out

SLC19A1

N   C
In

BRIL

Fab

Nb

e

Out

N   C
In

E8 R148
R145
E4 D5

f

Out

In

1b
1a   2   3   4   5   6   7   8   9   10   11   12

N   C

g

Out

6   1b
3   4
5   8
2
10
11 7
N   9
In   C
1a

6
1b

5
1a

90°

2
3   11
4   7
6
1b   12
1a   10
5   8   9