Development of a new cold hardiness prediction model for grapevine using phased integration of

acclimation and deacclimation responses

Al P. Kovaleski[1,*], Michael G. North[1], Timothy E. Martinson[2], Jason P. Londo[2,*]

[1]Department of Horticulture, University of Wisconsin–Madison, Madison, WI 53706, USA

[2]School of Integrative Plant Science, Cornell AgriTech, Cornell University, Geneva, NY 14456, USA

[*]Authors for correspondence:

AP Kovaleski: al.kovaleski@wisc.edu

JP Londo: jpl275@cornell.edu

Abstract

Cold injury in plants limits distribution of perennial agricultural crops, though replacement of plants and other management practices allow for some damage to be tolerated. Grapes (Vitis spp.) represent a crop that is grown in many marginal areas due to its profitability. However, monitoring of cold hardiness or damage for deployment of cold damage mitigation and management practices is a laborious process. To that effect, a model was published in 2014 to predict cold hardiness using data from Washington state in the US. Although that model works well regionally, it has been found to not predict adequately in other regions. In part, it is known that its description of some dormancy aspects as they relate to cold hardiness are not accurate. Therefore, the objective of this work was to develop a new model that incorporate previous research on cold hardiness dynamics, such as to increase realism. Cold hardiness data from *V. labruscana* 'Concord', and *V. vinifera* 'Cabernet Sauvignon' and 'Riesling' from Geneva, NY, USA were used. Data were separated in calibration (~2/3) and validation (~1/3) datasets. The proposed model uses 3 functions to describe acclimation, and 2 functions to describe deacclimation, with a total of 9 optimized parameters. A shared response to chill between acclimation and deacclimation provides a phased integration where acclimation responses decrease over the course of winter and are overcome by deacclimation. The new model outperforms the currently available model, reducing RMSE by up to 37% depending on cultivar. The new model also tends to slightly underpredict cold hardiness, as opposed to the overprediction from the current model. The underprediction can be preferred as damage mitigation methods may be deployed prior to damage occurring with greater safety. Some optimized parameters were shared between cultivars, suggesting conserved physiological aspects of the process were captured by the new model.

Keywords

Grapevine, *Vitis*, cold hardiness, model, differential thermal analysis

1. Introduction

Perennial agricultural plants are often introduced beyond their native cold range limits and therefore have higher risk of cold injury during winter (Hänninen 2013). This is the case for grapevines (*Vitis* spp.), especially domesticated European grapevines (*V. vinifera*), in northern regions within the United States and in Canada, where low temperatures during the dormant season in winter are a limiting factor. Single freezing events can effectively eliminate plant populations, thus limiting the cold range edge of native plants (Körner, 2021). In agricultural systems, however, injured or dead plants can be replaced, although this increases production costs and can create gaps or delays in production. Therefore, understanding and predicting cold hardiness of perennial plants can be helpful in determining management practices and site selection.

Domesticated European grapevines are native to the Mediterranean region yet are cultivated on all continents except Antarctica. Within the US, New York state is the third largest grape producer in the US (USDA, 2019), growing grapevines in significantly colder conditions than that of the larger two producing states (California and Washington), and colder than some regions with which it shares latitudes within Europe, such as northern Spain, southern France, and central Italy. Viticulture in New York relies on the proximity to large bodies of water (e.g. Lake Erie, Lake Ontario, the Finger Lakes, Atlantic Ocean) to help buffer low temperatures during winter, thus allowing cultivation of *V. vinifera* grapevines, which are generally less cold hardy than American species (e.g., *V. labrusca*, *V. riparia*) and hybrid varieties (Londo and Kovaleski, 2017). Winter survival of mixed buds in grapevines is required for production of grapes in the following growing season. These buds are formed during summer, and must remain dormant and develop cold hardiness throughout fall to survive low temperatures in winter before resuming growth in spring. Predicting bud cold hardiness can thus be used to evaluate winter bud injury, particularly in currently cold growing regions, but can also be used to evaluate current suitability for expansion into other areas, or for identifying regions which will become more/less suitable as climate changes. While climate predictions suggest winter temperatures will rise, more erratic climate is also expected with large swings between warm temperatures and acute low temperatures (IPCC, 2022). As cold damage mitigation tools are developed, accurate predictions of cold hardiness are also needed to assess necessity and timing of deployment of these methods (Li and Dami, 2016; Kovaleski and Londo, 2019). Therefore, accurate cold hardiness models can help evaluate the future of viticulture in cold climates from a winter survival standpoint.

Dormancy and cold hardiness are separate but concomitant processes within a plant (Arora et al.,

2003; Kovaleski, 2022). Both processes are required for survival of low temperatures during the cold

season in subtropical and temperate environments but appear to respond to slightly different

environmental cues. Dormancy establishment in grapevines typically occurs in response to shortening

photoperiod in late summer and fall (Fennel and Hoover, 1991) but other environmental factors can also

play a role [e.g., drought (Shellie et al., 2018)]. Dormancy progression is then modulated by exposure to

low temperatures (chilling), transitioning from a warm temperature non-responsive (endodormancy) to a

responsive phase (ecodormancy), from which plants are released from upon return of warm temperatures

in spring (Lang et al., 1987).

Cold hardiness usually presents a U-shape pattern during the progression from fall to winter to

spring with three distinct phases. In the fall, acclimation leads to gains in cold hardiness of grapevine buds

through the establishment and enhancement of supercooling. In mid-winter, buds reach maximum cold

hardiness and maintain high levels of cold hardiness depending on environmental cues. Finally, in late

winter and early spring as warm temperatures return, deacclimation leads to loss of cold hardiness –

though reacclimation is possible upon return of low temperatures. Photoperiod alone is unable to elicit

significant gains in cold hardiness (Fennel and Hoover, 1991). While some authors have indicated that

only the synergistic effect of short photoperiod and low temperatures can induce deep cold hardiness

(Schnabel and Wample 1987), temperature effects appear to be the largest contributors to cold hardiness

levels of grapevine buds, and models solely based on temperature have shown to produce good regional

predictions (Ferguson et al., 2011, 2014).

Dormancy directly affects cold hardiness dynamics. During the endodormant period, in the fall

and early winter, buds respond to low temperatures to gain cold hardiness. The ability to gain cold

hardiness appears to decrease as chilling accumulates and buds transition to ecodormancy (Ferguson et

al., 2011, 2014; North et al., 2021). The inverse seems to happen with regards to warm temperatures: a

bud's deacclimation responsiveness increases as chilling accumulates (Ferguson et al. 2011, 2014;

Kovaleski et al., 2018; North et al., 2021; Kovaleski, 2022). Therefore, acknowledging dormancy

dynamics is necessary for predicting cold hardiness.

Currently, the most widely used model for predicting grapevine bud cold hardiness was developed

using cold hardiness data from Washington state based on *V. vinifera* 'Cabernet Sauvignon',

'Chardonnay', and *V. labruscana* 'Concord' (Ferguson et al., 2011), and was later enhanced and further

expanded to add twenty other cultivars (Ferguson et al., 2014) – from here on referred to as "Washington

(WA) model". This model has also been recently adapted for cold climate interspecific hybrids grown in a

different environment (North et al., 2021). However, as indicated by the original authors, local adaptation

of the model may be required given the specific climate in which it was developed. Indeed, the model

does not produce reliable results in viticultural regions in the Eastern United states, including the state of

New York. This is likely a consequence of trade-offs among three dimensions in models: accuracy,

realism, and generality (Levins, 1966, 1968). Accuracy represents the fit between predictions and

observations, realism refers to causally appropriate internal model processes, and generality represents

robust applicability across space and/or time (Kramer et al., 2002). A model will ideally maximize all

accuracy, realism, and generality simultaneously; however, in practice, model developers must identify an

optimal point among the three dimensions that matches the overall aim of a model (Hänninen, 2016). In

the case of the WA model, accuracy was prioritized over realism and generality. Recent studies of

dormancy related to cold hardiness have shown that some of the parameters used in the model, such as the

linear response to temperature and the discrete change in responses by the vines based on dormancy status

do not accurately describe the nature of the process (Kovaleski et al., 2018; North et al., 2022). In fact,

linear responses to temperature have been related to errors in predictions of spring phenology as well

(Wolkovich et al., 2021).

The objective of this study was to develop a new model for cold hardiness prediction of grapevine

buds with greater accuracy for the Eastern United States and other viticultural regions more broadly. To

do so, we use a medium-term dataset comprised of three cultivars: *V. labruscana* 'Concord', and *V.*

vinifera 'Cabernet Sauvignon' and 'Riesling'. The choice of cultivars was made based on availability of

data (partially due to regional relevance), and variability in responses. The new model expands from

previously empirically determined concepts of deacclimation in response to chilling and forcing

temperatures (Kovaleski et al., 2018; Kovaleski, 2022; North, 2022), thus with increased realism, and

incorporates theoretical derivation of acclimation functions based on preliminary experimental data to

predict cold hardiness with field-based temperature data.

2. Materials and Methods

2.1. Cold hardiness data. Bud cold hardiness data from three cultivars were used: *V. labruscana* 'Concord', *V. vinifera* 'Cabernet Sauvignon', and *V. vinifera* 'Riesling' (from here on referred to by cultivar name only). Buds were collected from vineyards located in Geneva, NY, throughout multiple dormant seasons in varying numbers depending on cultivar (for details, see **Supplementary Table S1**). Cold hardiness was measured using differential thermal analysis (DTA) (Wolf and Pool, 1987; Mills et al., 2006; Ferguson et al., 2011, 2014; Londo and Kovaleski, 2017, North et al., 2021). Briefly, for DTA, buds are placed on thermoelectric modules connected to a datalogger and placed into a programmable freezer which can gradually lower temperature over time. Once intracellular water freezes, heat release is recorded as a low temperature exotherm (LTE) and signifies the temperature of death for any given bud (see Mills et al., 2006 for additional details). The number of buds used to measure cold hardiness for each cultivar on each sampling date varied between 5 and 30 buds (although number of recovered peaks may have been lower). For model development, average LTE values for each sampling date were used.

Eight seasons of data are available for 'Concord', five for 'Cabernet Sauvignon', and nine for 'Riesling'. The number of observations in each season varied based on cultivar, length of the season, and research program. For some seasons, more than one program collected data independently, but data were combined for modeling without using lab as a factor.

2.2. Weather data. Weather data were obtained from the Network for Environment and Weather Applications (NEWA; www.newa.cornell.edu). Hourly data were downloaded from which daily maximums and minimums were extracted for model development.

2.3. Data analyses. All data were analyzed with R (ver. 4.1.2; R Core Team, 2021) within R Studio (RStudio Team, 2021). Chill accumulation was calculated from hourly temperature data for each winter based on the dynamic model (chill portions; Fishman et al., 1987a, 1987b) using the chillR package (Luedeling, 2020). Three inputs were then used for model development, daily maximum (T_{max}) and minimum temperatures (T_{min}), and chill accumulation (*chill*, which is a function of temperature).

2.4. Model development. The model presented here is based on a generalized function for cold hardiness

(*CH*) from Kovaleski (2022) where the *CH* at any time t_i during the dormant season is a result of the

integration of separate acclimation and deacclimation processes occurring starting from bud set to t_i:

$$CH_{t_i} = CH_{bud\ set} + \int_{bud\ set}^{t_i} Acclimation\ dt + \int_{bud\ set}^{t_i} Deacclimation\ dt \quad [Eq.1]$$

where $CH_{bud\ set}$ is the cold hardiness of buds as they set and mature in late summer and fall, which is a

characteristic of the tissues, and *Acclimation* and *Deacclimation* are comprised of functions described

below. Here, *Acclimation* and *Deacclimation* are integrated over daily time steps for *t*, but both are

functions of temperature, either directly or indirectly through *chill*. However, field temperatures are

themselves a function of time.

2.4.1. For **Acclimation**, three functions were used and are based on expectations of the responses (visual

representations of these are presented in **Figs. 1A-D**). First, a function is used to describe what is the

maximum temperature below which any acclimation occurs as a response to chilling accumulation [the

acclimation temperature threshold ($T_{th_{acc}}$)]:

$$T_{th_{acc}}(chill) = T_{th_{low}} + \left.\frac{(T_{th_{high}} - T_{th_{low}})}{1 + e^{[b\ (\ln(chill) - \ln(c)]}}\right. \quad [Eq.2]$$

where the temperature threshold for acclimation decreases from a high threshold ($T_{th_{high}}$) at low chill

accumulations to a low threshold ($T_{th_{low}}$) at high chill accumulations; *b* is the slope associated with how

quickly the decrease occurs, and *c* is the inflection point of the log logistic curve (**Fig. 1A**). This decrease

in temperature threshold represents physiological changes during chill accumulation and dormancy

progression affecting the directionality of the cold hardiness dynamics process: at high chill

accumulations, the temperature experienced by buds must be lower to promote any acclimation. Both

parameters *b* and *c* are shared with deacclimation potential, which is explained further below (*2.4.2.*

Deacclimation).

Figure 1. Graphic representation of functions used to model cold hardiness of grapevine buds for 'Concord', 'Cabernet Sauvignon', and 'Riesling'. Functions are separated between acclimation (**A-D**) and deacclimation (**E, F**). Solid colored lines represent fits (**A, B, E, F**), or example fits for each cultivar (**C, D**); blue = 'Concord', magenta = 'Cabernet Sauvignon', green = "Riesling". The semi-transparent grey lines represent iterations of combinations for 'Concord' parameters that were not selected. **(A)** Acclimation step 1; see also **Eq. 2**. The temperature below which acclimation occurs is the acclimation temperature threshold ($T_{th_{acc}}$), which changes according to chill accumulation. As chill accumulates, $T_{th_{acc}}$ changes from high acclimation temperature threshold ($T_{th_{high}}$) to low acclimation temperature threshold ($T_{th_{low}}$). The driver of acclimation, DT, is calculated from the difference between the daily low temperature and $T_{th_{acc}}$. **(B)** Acclimation step 2; see also **Eq. 3**. Upper limit of curves, or maximum cold hardiness (CH_{max}) at DT_{acc} = 0°C, represents cold hardiness at bud set ($CH_{bud\ set}$), which was fixed at –6 °C for all cultivars. Lower limit of curves, or CH_{max} at DT_{acc} = 30°C, represents the maximum cold hardiness observed during any season ($CH_{abs\ max}$), which we set equal to the average of the 15 lowest cold hardiness measurements rounded down to the nearest integer. The shape of these curves between the upper and lower limits are determined by fitting the slope (*d*) and intercept (*f*) parameters in **Eq. 3**. **(C)** Acclimation step 3; see also **Eq. 4**. Example curves for artificial cold hardiness (CH^*) for a maximized DT in response to artificial time. These curves represent the time buds would take when exposed to a given DT_{acc} to reach the CH_{max} elicited. **(D)** Final acclimation rate based on partial derivative of step 3; see also **Eq. 5**. The acclimation rate k_{acc} is obtained based on the artificial time calculated from Eq. 4 solved for time using the the cold hardiness of the previous day as CH^*. **(E)** Deacclimation step 1, deacclimation rate; see also **Eq. 6**. The deacclimation rate (k_{deacc}) varies across temperature from 0 °C to 40°C. For all cultivars, k_{deacc} reaches maximum at 25 °C (T_{optim}) and is zero at 40 °C (T_{max}). Two aspects that were fit separately for each cultivar are the maximum deacclimation rate ($k_{deacc_{max}}$) and the coefficient *a*, which determines the slope on either side of $k_{deacc_{max}}$. **(F)** Deacclimation step 2, deacclimation potential; see also **Eq. 7**. Deacclimation potential (Y_{deacc}) modulates the deacclimation rate to an effective deacclimation rate (k^*_{deacc}) based on chill accumulation from low to high (i.e., Y_{deacc} increases as chill accumulates). The inflection point for all cultivars was fixed at 66 chill portions, while the slope parameter of Y_{deacc} curves (*b*) was fit separately for each cultivar and is shared with $T_{th_{acc}}$.

The second function describes the maximum cold hardiness possible (CH_{max}) elicited by the difference between the T_{min} of any given day and the $T_{th_{acc}}$ ($\Delta T_{acc} = T_{th_{acc}} - T_{min}$). Here, we used a log-logistic function to describe the response:

$$CH_{max}(\Delta T_{acc}) = CH_{bud\,set} + \frac{(CH_{abs\,max} - CH_{bud\,set})}{1 + e^{[d\,(\ln(\Delta T_{acc}) - \ln(f))]}} \quad [Eq.3]$$

where $CH_{bud\,set}$ is the minimum cold hardiness, which was fixed at –6 °C for all cultivars (about the limit of detection of our equipment); $CH_{abs\,max}$ is the maximum cold hardiness observed during any season, and here was used as the average of the 15 lowest cold hardiness measurements rounded down to the nearest integer; d is the slope associated with the logistic curve; and f is the inflection point of the logistic curve (**Fig. 1B**).

The third function describes the rate of acclimation based on how close to the CH_{max} the cold hardiness of the previous day ($CH_{t_{i-1}}$) is. For simplicity, here we present this in two steps. First, the shift from $CH_{bud\,set}$ to the CH_{max} elicited by any given ΔT_{acc} is not instant. Should buds be exposed to a constant ΔT_{acc}, acclimation would occur over time, giving rise to rate of acclimation. Because these are based on assumptions related to acclimation rather than experiments, we have dubbed this cold hardiness as an artificial cold hardiness (CH^*). The artificial cold hardiness (CH^*) thus changes from $CH_{bud\,set}$ to CH_{max} as a function of time based on a log-logistic function:

$$CH^* = CH_{bud\,set} + \frac{(CH_{max} - CH_{bud\,set})}{1 + e^{\{g\,[\ln(t) - \ln(h)]\}}} \quad [Eq.4]$$

where g and h are the slope and intercept associated with a log-logistic curve (**Fig. 1C**). CH^* is also indirectly a function of the temperature and chill accumulation since both contribute to defining CH_{max}. An artificial time t_a is found by solving **Eq. 4** for t using $CH^* = CH_{t_{i-1}}$ (the cold hardiness of the previous day) and CH_{max} for T_{min} at t_i (using **Eq. 3**). Using t_a, we can obtain the rate of acclimation (k_{acc}) in the field at any time t_i through the partial derivative of the log logistic function with respect to time:

264
$$k_{acc} = (CH_{max} - CH_{bud\ set}) \times \left\{ 0.001 + \frac{\left(e^{[g\ (\ln(t_a) - \ln(h))]} \right) \times (-g/t_a)}{(1 + e^{[g\ (\ln(t_a) - \ln(h))]})^2} \right\} \quad [Eq.5]$$

Where g and h are the slope and intercept associated with a log-logistic curve, and t_a is the artificial time

based on $CH_{t_{i-1}}$ (**Fig. 1D**). Here, k_{acc} is a function of artificial time (since field temperature changes

continuously), and the field temperature and accumulated chill that define CH_{max}. In addition to the rate

obtained through the partial derivative, a 0.001 (or 0.1%) rate was added to acclimation at all times. This

was incorporated to acknowledge anecdotal observations that buds seem to continue acclimating when

temperatures are very low (Kovaleski personal observations). Therefore, buds can surpass the CH_{max}

elicited by any ΔT_{acc}, and even a cultivar's $CH_{abs\ max}$, if temperatures are continuously low. While the

actual effect of the factor is very small in this model, the factor could be enhanced or reduced to

accommodate new perspectives based on future research. A sample workflow of an *Acclimation* step is

shown in **Fig. 2**.

Figure 2. Example workflow to describe progression for acclimation step and highlight relative importance of changes in conditions. **(A)** The $T_{th_{acc}}$ in this example varies from 12 °C ($T_{th_{low}}$) to 0 °C ($T_{th_{high}}$) across accumulating chill. DT is the difference between T_{min} (points) and $T_{th_{acc}}$. Two DT_{acc} scenarios are included in this example (DT_I and DT_{II}) to highlight importance of conditional changes throughout acclimation calculations. Here we observe that different temperatures can lead to the same DT ($DT_I = DT^*_I$): as chill accumulates and $T_{th_{acc}}$ decreases, a lower T_{min} would elicit the same response. We can also observe that the same temperature can lead to different DTs ($DT_{II} \neq DT^*_I$) if at two different chilling accumulations. **(B)** Based on a given DT_{acc}, a certain CH_{max} is obtained, which is greater (in magnitude) for greater DT_{acc}. **(C)** With the different CH_{max} for each of the scenarios, DT_I and DT_{II}, we can observe the different curves for gains in cold hardiness if buds were constantly exposed to those temperatures. Solving ***Eq. 4*** for time using the CH of the previous day (CH_{i-1}), here −15 °C, as CH^* we obtain the time at which this would occur along that artificial temperature exposure: t_{a_I} and $t_{a_{II}}$. **(D)** Using the first derivative of those acclimation curves, we obtain the curves for rate of acclimation, k_{acc}. The rate of acclimation for a given day, t_i, is calculated by solving ***Eq. 5*** using t_{a_I} and $t_{a_{II}}$.

2.4.2. For ***Deacclimation***, two functions were used (**Figs. 1E, F**) which are based on descriptions from

previous empirical experiments on grapevines and other species (Kovaleski et al., 2018; Kovaleski, 2022;

North et al., 2022). One is for the rate of deacclimation (k_{deacc}) that any given temperature elicits. Other

functions have been used to describe k_{deacc}, such as a logistic (Chuine, 2000; North et al., 2022),

combination of exponential and logarithmic (Kovaleski et al. 2018) or a third-degree polynomial

(Kovaleski, 2022). Here, the temperature response curve used was that based on a temperature-

photosynthesis curve (O'Neil et al., 1972):

$$k_{deacc}(T) = k_{deacc_{max}} \left(\frac{T_{max} - T}{T_{max} - T_{opt}} \right)^a \times e^{\left[a \left\{ 1 - \left(\frac{T_{max} - T}{T_{max} - T_{opt}} \right) \right\} \right]} \quad [Eq.\,6]$$

where k_{deacc} (T) is the rate of deacclimation (°C day^{-1}) at temperature T (°C); a is a coefficient related to

the slope of the curve; $k_{deacc_{max}}$ is the maximum rate of deacclimation at the optimal temperature (T_{optim},

°C), and T_{max} is the maximum temperature (°C) from which point the deacclimation rate becomes zero.

T_{optim} and T_{max} were fixed at 25 °C and 40 °C, respectively. While an absolute decrease in rates with

temperature has not observed in empirical studies (Kovaleski et al., 2018; North et al., 2022), some

considerations can be made: a deviation from exponential increase in rates was observed (Kovaleski et al.,

2018; Kovaleski, 2022); temperatures much higher than 25 °C do not occur frequently during the

deacclimation period in the spring; and the suggestion of deacclimation as an enzymatic process is

acknowledged by this curve (Kalberer, 2006; Kovaleski et al., 2018). The T_{optim} of 25 °C was used based

on previously reported deacclimation rates (Kovaleski et al., 2018; North et al., 2022). Because of the

non-linearity of the response, the amount of deacclimation observed in a day was the average between

k_{deacc} for the daily T_{max} and T_{min}, rather than using the average of the temperature.

The second function defining deacclimation is the deacclimation potential (Ψ_{deacc}), which modulates the

level of effective deacclimation occurring based on a log logistic function of chill accumulation during the

winter:

$$\Psi_{deacc}(chill) = {1}/{1 + e^{[-b\,(\ln(chill) - \ln(c)]}} \quad [Eq.\,7]$$

where b is the slope associated with the log-logistic curve, and c is the inflection point of the log-logistic

curve, a chilling requirement-analogous measurement where 50% of the maximum Ψ_{deacc} is achieved.

Because b and c were shared with $T_{th_{acc}}$, such that the dormancy-related responses to *chill* are symmetric

for *Acclimation* and *Deacclimation*, a negative sign is used for b here (negative values indicate increases

in Ψ_{deacc} as chill accumulation increases). For c, a fixed value of 66 chill portions was used based on

previous work with many species (Kovaleski, 2022). The phased integration of *Acclimation* and

Deacclimation provided by the shared b and c terms mean that as chilling accumulates, *Acclimation*

becomes more difficult, while *Deacclimation* becomes more pronounced.

The effective deacclimation rate at any time t_i, $k^*_{deacc_{t_i}}$, is ultimately a result of the product of the average

daily deacclimation rate and the deacclimation potential:

347
$$k^*_{deacc_{t_i}} = {[k_{deacc}(T_{min}) + k_{deacc}(T_{max})]}/_2 \times \Psi_{deacc} \quad [Eq.8]$$

2.5. Model optimization. The cold hardiness data available was separated in calibration and validation

datasets by season-year. The ratio was approximately 2/3 calibration and 1/3 validation. The years were

randomly assigned to each dataset within each cultivar using set.seed() based on when code preparation

for each cultivar started [17 August for 'Concord' set.seed(1708); 7 June for 'Cabernet Sauvignon'

set.seed(0706); 7 July for 'Riesling' set.seed(0708)]. The dynamic model based on **Eq. 1** predicts cold

hardiness with 1-day time steps. Nine parameters were optimized here: $T_{th_{low}}, T_{th_{high}}, b, d, f, g, h,$

$k_{deacc_{max}}$, and a (see **Table 1** and **Fig. 1**). The parameters were optimized using a stepwise iterative

method based on the minimization of root mean square error (RMSE) between predicted and observed

CH for a calibration dataset (RMSE$_{cal}$). The parameter combination with lowest RMSE$_{cal}$ was then used to

predict CH using a naïve validation dataset, and RMSE of a prediction was then calculated (RMSE$_{val}$).

The RMSE$_{val}$ of the present model was compared to that of the predictions based on the latest update of

the currently available grapevine bud cold hardiness developed in Washington state (RMSE$_{WA}$; Ferguson

et al., 2014, see also Ferguson et al., 2011 for details on equations). Other fitness statistics were also

calculated for both models [*Bias* = $CH_{measured} - CH_{predicted}$; Pearson correlation ($r$)].

Table 1. Parameters optimized and their range

Cultivar	$T_{th_{low}}$	$T_{th_{high}}$	b	c	$CH_{abs\ max}$			$CH_{bud\ set}$	d	f	g	h	k_{deacc}			a
					Con	CS	Rie						Con	CS	Rie	
Start	–6	6	6	66	31	23	2	6	2	2	–2.3	12	1.9	1.3	1.7	3.5
Stop	0	14	11	66	31	23	7	6	7	5	–1.7	24	2.4	1.8	2.2	5.0
Step	2	2	1	0	0			0	1	1	0.1	2	0.1			0.5
n	4	5	6	1	1			1	6	4	7	7	6			4
Equation	**Eq. 2**		**Eqs. 2, 7**		**Eq. 3**			**Eqs. 1, 3, 4, 5**	**Eq. 3**		**Eqs. 4, 5**		**Eq. 6**			

3. Results

The model proposed here using optimized parameters for each cultivar based on calibration dataset (**Table 2**, **Fig. 1**) produced good predictions of cold hardiness for the validation dataset of the three cultivars studied (RMSE$_{val}$ ranging from 2.12 ℃ to 2.72 ℃, bias from –1.46 ℃ to 1.28 ℃; **Table 2**, **Figs. 3 and 4A**).

Figure 3. Observed cold hardiness, based on average low temperature exotherms (points), plotted with predicted bud cold hardiness according to *New York model* (solid lines) and Washington model (dashed lines) with daily maximum and minimum temperature range (grey ribbons) in 2018-2019 for three grapevine cultivars.

The values for RMSE$_{cal}$ (2.09 ℃, 1.47 ℃, 1.61 ℃ for 'Concord', 'Cabernet Sauvignon', and 'Riesling', respectively) and RMSE$_{val}$ (**Fig. 4A**) were similar within each cultivar. This demonstrates the model and parameters chosen are robust, thus predicting similar variability in both datasets, dealing well with the

386 year-to-year weather variations. In absolute values, predictions produced were slightly better for the two

387 *V. vinifera* cultivars compared to 'Concord'. This may be a result of the higher responsiveness of

388 'Concord' compared to either *V. vinifera* (greater changes in *CH* over time), as well as its overall greater

389 cold hardiness.

Figure 4. Comparison between predicted and observed bud cold hardiness, reported as average low temperature exotherm, for three grapevine cultivars using both the (**A**) *New York model* and (**B**) Washington model. Statistics are listed in each panel, including root mean square error (RMSE), bias, correlation coefficient (r), and sample size (n). The prediction and observation correlations are illustrated (solid black line) along with a one-to-one reference (dashed black line).

Compared to the existing Washington model, the newly developed model produced better predictions for

'Concord' and 'Cabernet Sauvignon', and slightly better or comparable predictions for 'Riesling' for data

from Geneva, NY (**Figs. 3,4**). $RMSE_{WA}$ was only computed within the validation datasets, and had values

of 4.14 °C, 3.63 °C, and 2.24 °C for 'Concord', 'Cabernet Sauvignon', and 'Riesling', respectively.

The Washington model consistently overpredicted cold hardiness ($Bias_{WA}$ of 3.12 °C, 3.03 °C, and 1.54

°C for 'Concord', 'Cabernet Sauvignon', and 'Riesling', respectively; **Fig. 4B**). This is likely due to the

development of the model using data from a region with relatively warmer and drier winters. The bias may be enhanced by the model's design, where any temperature below the temperature threshold that separates acclimation and deacclimation can lead to full acclimation to the maximum cold hardiness. In the present model, low temperatures have both a different maximum cold hardiness possible (**Eqs. 2 and 3**) and different acclimation rates (due to the effect of maximum cold hardiness on **Eqs. 4 and 5**). Our model, although selected based on lowest RMSE$_{cal}$, also had low bias for the three cultivars.

In the same season, 'Concord' gains more cold hardiness, and does so faster, than the two *V. vinifera* despite the same starting cold hardiness. This effect is observed in the generally lower CH_{max} (**Fig. 1B**) and higher rates of acclimation in response to ΔT_{acc} (**Fig. 1D**). 'Concord' also loses its cold hardiness faster than the *V. vinifera*, while 'Riesling' is also faster than 'Cab. Sauvignon'. Within the optimized parameters, 'Riesling' has a higher $k_{deacc_{max}}$ than 'Concord'. However, because of the higher a parameter for **Eq. 6** (**Table 2**), 'Riesling' has lower deacclimation rates at temperatures between 0 ºC and 17.5 ºC (**Fig. 1E**), which are temperatures that largely predominate during the spring, when the majority of deacclimation occurs.

While all combinations of parameters were tested for optimization within each cultivar, some parameters were found to be very similar between the cultivars. This could suggest that the model is realistic in describing a physiological process that is conserved among the different cultivars within a species ('Cabernet Sauvignon' and 'Riesling'), and among grapevine species (here *V. vinifera* and *V. labruscana*). 'Cabernet Sauvignon' had a lower $T_{th_{high}}$ than 'Concord' and 'Riesling', but the rate of change in $T_{th_{acc}}$ in response to chilling (b) was very similar. The parameter b, which defines the slope of the log-logistic associated here with both $T_{th_{acc}}$ and Ψ_{deacc}, has previously been suggested to be the same for all species for deacclimation (Kovaleski, 2022), leaving regulation of magnitudes of deacclimation (effective deacclimation) to the effect of temperature alone on the rate of deacclimation, k_{deacc}.

Cultivar	$T_{th_{low}}$	$T_{th_{high}}$	b	c	$CH_{abs\,max}$	$CH_{bud\,set}$	d	f	g	h	k_{deacc}	a	Calibration	Validation
'Concord'	0	12	8	66	–31	–6	3	5	–2.1	24	1.9	3.5	2012-2013, 2013-2014, 2016-2017, 2017-2018, 2019-2020	2014-2015, 2018-2019, 2020-2021
'Cabernet Sauvignon'	0	8	7	66	–23	–6	6	3	–2.2	22	1.8	5.0	2016-2017, 2017-2018, 2018-2019, 2020-2021	2015-2016, 2019-2020
'Riesling'	0	12	7	66	–26	–6	3	4	–1.9	24	2.2	5.0	2012-2013, 2014-2015, 2017-2018, 2018-2019, 2019-2020, 2020-2021	2013-2014, 2015-2016, 2016-2017

4. Discussion

Accurate modeling of cold hardiness is an important step in understanding adaptation of species and cultivars in future climates (Wolkovich et al., 2018; Körner, 2021). However, modeling of this trait is also helpful for growers in guiding management practices and in cultivar choice for different regions in current climates. Here we used known deacclimation responses (Kovaleski et al., 2018; Kovaleski, 2022; North et al., 2022) and posited acclimation responses, along with a medium-term dataset with three cultivars that have relatively different cold hardiness behaviors to create a new cold hardiness prediction model for grapevine buds.

Our new model ("NY model") outperforms the currently available model for grapevine cold hardiness (Ferguson et al., 2014) using New York state data. The WA model consistently overestimates bud cold hardiness in New York, which consequently can underestimate freeze injury risk and occurrence. An overestimated cold hardiness could mislead adaptation of production systems (e.g. site selection or cultivar evaluation) and management decisions (e.g. spring pruning decisions). This is especially consequential as extreme and erratic weather events during winter become more frequent under changing climates (Casson et al., 2019; IPCC, 2022).

Parameters here are optimized using field data, as in other modeling studies (Ferguson et al., 2014;

North et al., 2021). This was done despite some data available on deacclimation rates for the two *V.*

vinifera cultivars (Kovaleski et al., 2018) and a few hybrids (North et al., 2022), though not 'Concord'.

However, the estimated deacclimation rates based on **Eq. 6** for either *V. vinifera* cultivars were

comparable to values previously reported (Kovaleski et al., 2018). This indicates field data, associated

with models that prioritize the realism model dimension, can estimate parameters equivalent to controlled

experiments.

The modulation of acclimation rate based on a logistic regression in response to time has a similar

effect here as that of the Theta (θ) term in the Washington model, effectively attenuating the approach to

any given cold hardiness. However, because each temperature elicits a different cold hardiness in our

model (**Eqs. 2 and 3**), this effect is dynamic and can occur at any cold hardiness and any temperature,

rather than just the asymptotical approach to absolute maximum and minimum cold hardiness as in

Ferguson et al. (2014). This may improve the generalizability of our model in applications to regions

other than New York. Deployment of the model to various cold hardiness monitoring programs (e.g., MI,

PA, OH, CO, and Canada) for validation of fit would enable validation of these assumptions.

The optimized $T_{th_{low}}$ was 0 ºC for all cultivars, which is the upper limit established. For a best fit

of the model, it would be ideal to increase the range for iterations such that RMSE may be further

minimized, ideally finding a global minimum. However, a higher $T_{th_{low}}$ would mean that the small

deacclimations that happen at low but positive temperatures might have been counterbalanced by the

effect of acclimation. By limiting $T_{th_{low}}$ to 0 ºC we prevented this effect, which would otherwise be

particularly consequential during the spring when plants are fully chilled. Should data be gathered in

warmer regions with lower seasonal chill accumulation, this limit should be evaluated and possibly be re-

parameterized to remove these effects.

The calibration and validation datasets were not the same for all cultivars (**Table 2**;

Supplementary Figs. 1-3). This was due to data not having been collected the same number of years for

all cultivars. However, enough variation in the years available for the three cultivars appear to allow for

good parameterization of the models. For example, within the calibration dataset for 'Concord', 2012-

2013 is a much milder winter both in minimum temperature and the length of exposure to very low

temperatures as compared to 2013-2014 (**Supplementary Fig. 1**).

Here, the transition of endodormancy to ecodormancy occurs quantitatively through the

accumulation of chill that modulates the rates of deacclimation (k_{deacc}) through the deacclimation potential

(Ψ_{deacc}). This is very different from the WA model, where a qualitative transition occurs. It is likely that a

qualitative transition works well as an assumption within a single region, given some regularity in the

level of chill accumulation across years. However, for a single model to accomplish the estimation of

deacclimation rates in regions of lower chill accumulation and higher chill accumulation, a quantitative

transition is required based on experimental results (Kovaleski et al., 2018; Kovaleski, 2022, North et al.,

2022). Field data from different regions, particularly those where low chilling accumulation occur will be

required to test this. It is also important to note that some management practices that "supplement

chilling" and break dormancy (e.g., application of hydrogen cyanamide) have not been studied from a

cold hardiness context, and therefore we do not expect the model to excel if applied to regions where such

treatments occur until effects of dormancy breaking treatments has been studied.

The dynamic model was used here to estimate chilling accumulation, due to its reported higher

transferability to warmer environments (Luedeling and Brown, 2011). Within high chill accumulation

regions, it has been reported to be highly correlated to other chilling models, such as Utah and NC [e.g.,

in Boston, MA, USA (Kovaleski, 2022), and Madison, WI, USA (North et al., 2022)]. However, it is still

possible that all chill models are incorrectly describing chilling accumulation, and therefore this may be a

source of error for predictions, especially in regions with low chill accumulation. Deacclimation

experiments, such as in other studies by the present authors (Kovaleski et al., 2018; Kovaleski, 2022;

North et al., 2022), that use material collected from warmer climates should help elucidate the dynamics

of chilling. Should the chilling accumulation models change, parameter changes within the model

presented here may be required.

In our calibration dataset, the rounded average of the 15 most cold hardy points for 'Concord' is –

31 ℃, and the lowest observed cold hardiness was –31.4 ℃. This is below the maximum possible using

the Washington model for that cultivar (–29.5 ℃). This suggests the maximum cold hardiness for this

genotype may not be achieved in Washington's conditions, which are relatively warmer compared to

upstate New York. The higher value of cold hardiness being the maximum possible in their model results

in a hard limit seen in the lower end of the observed vs. predicted plots for the Washington model (**Fig.**

4B). The opposite is observed in 'Cabernet Sauvignon', however, where the present model uses a

maximum cold hardiness of –23 ℃ (although this value would be –26 ℃ if estimated from the validation

dataset), whereas the Washington model uses a value of –25.1 ℃. In the currently proposed model,

besides having a lower maximum, this is not a hard limit given the possibility of a low acclimation rate at

any temperature.

The values estimated for the slope associated with the log logistic for chill responses were similar

to those previously reported (Kovaleski, 2022). Here, both acclimation and deacclimation were guided by

the same slope, providing the phased integration of both, which may have led to slight differences from

measured effects. Future studies in acclimation could help understand whether the response should be

maintained as varying alongside deacclimation or not.

The similarity in parameter values across cultivars for acclimation and deacclimation, particularly

the portions that *do not describe magnitude* (i.e., slope for $T_{th_{acc}}$, slope and intercept for CH^*, and slope

for Ψ_{deacc}), suggests a conserved physiology if the model is correctly approximating responses. This has

been suggested for the dormancy and deacclimation across the land plant phylogeny (Kovaleski, 2022),

though without using multiple genotypes within a species. Deacclimation appears to be a simpler process,

where no temperature sensing is required, only temperature perception (Penfield, 2008). It remains to be

tested whether acclimation has conserved mechanisms that can easily translate to mathematical models of

the process, though acclimation appears to be more difficult to describe and elicit in controlled

environment experiments.

Variability in model performance across regions is at least partially a consequence of model

development with datasets from a single region. However, there is a lack of freely available datasets of

cold hardiness from different climates that restricts improvements in model development, unlike with

other types of data such as spring phenology (e.g., Pan European Phenology Project and USA's National

Phenology Network). Therefore, we are making our dataset public so other efforts may use this medium-

term dataset (https://github. com/apkovaleski/GrapeCHmodel; Kovaleski et al., 2022). It is also noticeable

that most cold hardiness datasets are produced in higher latitudes and temperate climates where

considerably low temperatures occur during winter [e.g., Washington (Ferguson et al., 2014), Wisconsin

(North et al., 2021, 2022) and New York states within the US; Chile (Rubio and Pérez, 2020)]. There is a

need for new data to be collected from warmer locations in lower latitudes such that parameters can be

optimized based on warmer subtropical (or even tropical) climates. It is possible that the model proposed

here, the existing WA model (Ferguson et al., 2014), or regional adaptations (North et al., 2021) perform

variably across different regions, and therefore models should continue to be used in conjunction in new

regions to validate and tune predictions. As models continue to be improved, convergence of responses

should result in high confidence of predictions.

Our model uses only air temperature-related parameters to describe cold hardiness. Some variation in cold

hardiness can also be explained by temperature mediated aspects of microclimate and daylength, such as

through radiative warming (Grace, 2006; Quiñones et al., 2019; Vitasse et al., 2021), wind, water

availability, and daily amplitude in temperature (Gonzalez Antivilo et al., 2017; Matsubara 2018). In

addition to direct effects on cold hardiness, such factors may indirectly affect cold hardiness through

effects on dormancy. However, integrating these relatively minor factors influencing cold hardiness

changes into models, while reducing prediction error somewhat, also increase a model's computational

complexity and potentially make the results of parameter computing less generalizable.

5. Conclusions

The model produced here demonstrates a robust manner of predicting cold hardiness for buds of three

different grapevine cultivars. This model integrates novel deacclimation concepts discussed in recent

literature (Kovaleski et al., 2018; Kovaleski, 2022; North et al., 2022). It is possible that future research

regarding acclimation could lead to a better understanding of the processes leading to gains in cold

hardiness and subsequently improved modeling approaches. However, acclimation seems to be an elusive

process to date and coincides with other significant physiological changes that are only partially

understood (e.g., dormancy induction). Therefore, further research in acclimation and cold hardiness

modeling will need to continually assess the balance between model realism and model accuracy. The

proposed model provides accurate cold hardiness predictions in the meantime and a framework on which

to build. Better predictions were produced using the proposed model when compared to a currently

available model (Ferguson et al., 2011, 2014). Thus, the proposed model may be used in lieu of, or in

addition to, existing models to aid in decision making steps. Despite having been developed using

regional data, the selection of similar parameters across cultivars could indicate a representative

description of physiological processes that may also promote transferability into other regions with some

caveats, such as errors in chilling accumulation models.

Acknowledgements

We would like to thank Hanna Martens, Bill Wilsey, Lex Pike, and Kathleen Deys for help collecting and

processing grape bud tissue for field cold hardiness measurements. We would also like to thank Ravines

Vineyard for access to 'Cabernet Sauvignon' and 'Riesling' vines. This work was partially supported by

the Office of the Vice Chancellor for Research and Graduate Education at the University of Wisconsin–

Madison with funding from the Wisconsin Alumni Research Foundation, and through the USDA ARS

appropriated project 1910-21220-006-00D.

Competing interests

The authors have no competing interests to declare.

References

Arora R, Rowland LJ, Tanino K. 2003. Induction and Release of Bud Dormancy in Woody Perennials: A Science Comes of Age. *HortScience.* **38:** 911-921.

Casson NJ, Contosta AR, Burakowski EA, Campbell JL, Crandall MS, Creed IF, Eimers MC, Garlick S, Lutz DA, Morison MQ, Morzillo AT, Nelson SJ. 2019. Winter weather whiplash: Impacts of meteorological events misaligned with natural and human Systems in Seasonally Snow-Covered Regions. *Earth's Future.* **7:** 1434–1450.

Chuine I. 2000. A unified model for budburst of trees. *Journal of Theoretical Biology.* **207:** 337–347.

Fennel A, Hoover E. 1991. Photoperiod influences growth, bud dormancy, and cold acclimation in *Vitis labruscana* and *V. riparia.* *Journal of the American Society for Horticultural Sciences.* **116:** 270-273.

Ferguson JC, Tarara JM, Mills LJ, Grove GG, Keller M. 2011. Dynamic thermal time model of cold hardiness for dormant grapevine buds. *Annals of Botany* **107**: 389–396.

Ferguson JC, Moyer MM, Mills LJ, Hoogenboom G, Keller M. 2014. Modeling dormant bud cold hardiness and budbreak in twenty-three *Vitis* genotypes reveals variation by region of origin. *American Journal of Enology and Viticulture.* **65**: 59–71.

Fishman S, Erez A, Couvillon GA. 1987a. The temperature dependence of dormancy breaking in plants: computer simulation of processes studied under controlled temperatures. *Journal of Theoretical Biology.* **126:** 309–321.

Fishman S, Erez A, Couvillon GA. 1987b. The temperature dependence of dormancy breaking in plants: mathematical analysis of a two-step model involving a cooperative transition. *Journal of Theoretical Biology.* 124: 473–483.

Gonzalez Antivilo F, Paz RC, Keller M, Borgo R, Tognetti J, Juñent FR. 2017. Macro- and microclimate conditions may alter grapevine deacclimation: variation in thermal amplitude in two contrasting wine regions from North and South America. *International Journal of Biometeorology.* **61:** 2033-2045.

Grace J. 2006. The temperature of buds may be higher than you thought. *New Phytologist.* **170:** 1-3.

Hänninen H. 2013. Modeling bud dormancy release in trees from cool and temperate regions. *Acta Forestalia Fennica.* **213:** 47.

Hänninen H. 2016. Boreal and temperate trees in a changing climate: modelling the ecophysiology of seasonality. *Biometeorology.* Springer. https://doi.org/10.1007/978-94-017-7549-6.

IPCC. 2022. Climate Change 2022: impacts, adaptation, and vulnerability. contribution of working group
II to the sixth assessment report of the intergovernmental panel on climate change [H.-O. Pörtner, D.C.
Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V.
Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press. In Press.

Kalberer SR, Wisniewski M, Arora R. 2006. Deacclimation and reacclimation of cold-hardy plants:
current understanding and emerging concepts. *Plant Science.* **171:** 3–16.

Körner C. 2021. The cold range limit of trees. *Trends in Ecology & Evolution.* **36:** 979-989.

Kovaleski AP. 2022. Woody species do not differ in dormancy progression: differences in time to
budbreak due to forcing and cold hardiness. *Proceedings of the National Academy of Sciences of the*
United States of America **119**: e2112250119.

Kovaleski AP, North MG, Martinson TE, Londo JP. 2022. Data from "Development of a new cold
hardiness prediction model for grapevine using phased integration of acclimation and deacclimation
responses" GitHub. https://github.com/apkovaleski/GrapeCHmodel Deposited ## September 2022.

Kovaleski AP, Londo JP. 2019. Tempo of gene regulation in wild and cultivated Vitis species shows
coordination between cold deacclimation and budbreak. *Plant Science.* **287:** 110178.

Kovaleski AP, Reisch BI, Londo JP. 2018. Deacclimation kinetics as a quantitative phenotype for
delineating the dormancy transition and thermal efficiency for budbreak in *Vitis* species. *AoB Plants* **10**:
ply066.

Kramer K, Leinonen I, Bartelink HH, Berbigier P, Borghetti M, Bernhofer C, Cienciala E, Dolman
AJ, Froer O, Gracia CA, Granier A, Grünwald T, Hari P, Jans W, Kellomäki S, Loustau D,
Magnani F, Markkanen T, Matteucci G, Mohren GMJ, Moors E, Nissinen A, Peltola H, Sabaté S,
Sanchez A, Sontag M, Valentini R, Vesala T. 2002. Evaluation of six process-based forest growth
models using eddy-covariance measurements of CO_2 and H_2O fluxes at six forest sites in Europe. *Global*
Change Biology. **8:** 213-230.

Lang GA, Early JD, Martin GC, Darnell RL. 1987. Endo-, para-, and ecodormancy: physiological
terminology and classification for dormancy research. *HortScience.* **22:** 371–377.

Levins R. 1966. The strategy of model building in population biology. *American Scientist.* **54:** 421-431.

Levins R. 1968. Evolution in changing environments: some theoretical explorations, Princeton University
Press, 2020. *Project MUSE* https://muse.jhu.edu/book/73534.

Li S, Dami IE. 2016. Responses of *Vitis vinifera* 'Pinot gris' grapevines to exogenous abscisic acid (ABA): I. yield, fruit quality, dormancy, and freezing tolerance. *Journal of Plant Growth Regulation* **35:** 245–255.

Londo JP, Kovaleski AP. 2017. Characterization of wild North American grapevine cold hardiness using differential thermal analysis. *American Journal of Enology and Viticulture* **68**: 203–212.

Luedeling E. 2020. chillR: Statistical Methods for Phenology Analysis in Temperate Fruit Trees. R package version 0.70.24. https://CRAN.R-project.org/package=chillR .

Luedeling E, Brown PH. 2011. A global analysis of the comparability of winter chill models for fruit and nut trees. *International Journal of Biometeorology* **55**: 411–421.

Matsubara S. 2018. Growing plants in fluctuating environments: why bother? *Journal of Experimental Botany*. **69:** 4651-4654.

Mills LJ, Ferguson JC, Keller M. 2006. Cold-hardiness evaluation of grapevine buds and cane tissues. *American Journal of Enology and Viticulture*. **57:** 194–200.

North M, Workmaster BA, Atucha A. 2021. Cold hardiness of cold climate interspecific hybrid grapevines grown in a cold climate region. *American Journal of Enology and Viticulture* **72**: 318–327.

North M, Workmaster BA, Atucha A. 2022. Effects of chill unit accumulation and temperature on woody plant deacclimation kinetics. *Physiologia Plantarum* **174**: e13717.

Penfield S. 2008. Temperature perception and signal transduction. *New Phytologist*. **179:** 615-628.

Quiñones AJP, Keller M, Gutierrez MRS, Khot L, Hoogenboom G. 2019. Comparison between grapevine tissue temperature and air temperature. *Scientia Horticulturae* **247**: 407–420.

R Core Team. 2021. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

RStudio Team. 2021. RStudio: integrated development environment for R. RStudio, PBC, Boston, MA.

Rubio S, Pérez FJ. 2020. Testing the Ferguson model for the cold-hardiness of dormant grapevine buds in a temperate and subtropical valley of Chile. *International Journal of Biometeorology* **64:** 1401–1408.

Schnabel BJ, Wample RL. 1987. Dormancy and cold hardiness in *Vitis vinifera* L. cv. White Riesling as influenced by photoperiod and temperature. *American Journal of Enology and Viticulture*. **38:** 265-272

Shellie K, Kovaleski AP, Londo JP. 2018. Water deficit severity during berry development alters timing of dormancy transitions in wine grape cultivar Malbec. *Scientia Horticulturae*. **232:** 226-230.

743 **USDA. 2019.** 2017 Census of agriculture. United States. Summary and State Data. Vol. 1. Geographic
744 Area Series.
745
746 **Vitasse Y, Baumgarten F, Zohner CM, Kaewthongrach R, Fu YH, Walde MG, Moser B. 2021.**
747 Impact of microclimatic conditions and resource availability on spring and autumn phenology of
748 temperate tree seedlings. *New Phytologist* **232:** 537–550.
749
750 **Wolf TK, Pool RM. 1987.** Factors affecting exotherm detection in the differential thermal analysis of
751 grapevine dormant buds. *Journal of American Society for Horticultural Science.* **112:** 520-525.
752
753 **Wolkovich EM, de Cortázar-Atauri IG, Morales-Castilla I, Nicholas KA, Lacombe T. 2018.** From
754 Pinot to Xinomavro in the world's future wine-growing regions. *Nature Climate Change.* **8:** 29-37.
755
756 **Wolkovich EM, Auerbach J, Chamberlain CJ, Buonaiuto DM, Ettinger AK, Morales-Castilla I,**
757 **Gelman A. 2021.** A simple explanation for declining temperature sensitivity with warming. *Global*
758 *Change Biology* **27:** 4947–4949.
759
760

Supplementary Figure 1. *Vitis labruscana* 'Concord' validation, calibration, and N/A datasets. Observed cold
hardiness, based on average low temperature exotherms (blue points) plotted with predicted bud cold hardiness
according to *New York model* (solid blue lines) and Washington model (dashed blue lines) with daily maximum and
minimum temperature range (grey ribbons).

Supplementary Figure 2. *Vitis vinifera* 'Cabernet Sauvignon' validation, calibration, and N/A datasets. Observed
cold hardiness, based on average low temperature exotherms (magenta points) plotted with predicted bud cold
hardiness according to *New York model* (solid magenta lines) and Washington model (dashed magenta lines) with
daily maximum and minimum temperature range (grey ribbons).

Supplementary Figure 3. *Vitis vinifera* 'Riesling' validation, calibration, and N/A datasets. Observed cold
hardiness, based on average low temperature exotherms (green points) plotted with predicted bud cold hardiness
according to *New York model* (solid green lines) and Washington model (dashed green lines) with daily maximum
and minimum temperature range (grey ribbons).