Body mass change over winter is consistently sex-specific across roe deer (*Capreolus capreolus*) populations

Hewison, A.J.M [a]*, Bonnot, N.C.[a], Gaillard, J.M.[b], Kjellander, P.[c], Lemaitre, J.F.[b], Morellet, N.[a] & Pellerin, M.[d]

[a] Université de Toulouse, INRAE, CEFS Castanet-Tolosan, France; *and* LTSER ZA PYRénées GARonne, 31320 Auzeville Tolosane, France

[b] Univ Lyon, Université Lyon 1; CNRS, Laboratoire de Biométrie et Biologie Evolutive UMR5558, 69622 Villeurbanne, France

[c] Grimsö Wildlife Research Station, Department of Ecology, Swedish University of Agricultural Sciences, Riddarhyttan, Sweden

[d] Direction de la Recherche et de l'Appui Scientifique, Office Français de la Biodiversité, Unité Ongulés Sauvages, 38610 Gières, France

* Corresponding author: A.J. Mark Hewison; mark.hewison@inrae.fr

co-authors are listed by alphabetical order

Abstract:

In most polygynous vertebrates, males must allocate energy to growing secondary sexual characteristics, such as ornaments or weapons, that they require to attract and defend potential mates, impacting body condition and potentially entailing fitness costs.

We investigated sex differences in over winter body mass change across five intensively monitored populations of roe deer (*Capreolus capreolus*) with markedly contrasting environmental conditions. At winter onset, males weighed, on average, 8.4% (from 4.7% in the most northerly population to 11.6% in the most southerly one) more than females. However, across all populations, males fared worse over the winter than females, losing more (Sweden) or gaining less (France) mass, so that sexual mass dimorphism was virtually absent prior to the onset of spring.

Our findings reveal that the direction of over-winter change in mass of roe deer depends on winter severity, but that males are consistently more sensitive to this environmental constraint than females. As a result of this sex-specific change in body mass, sexual mass dimorphism is lowest at the onset of the territorial season. We suggest that allocation to antler growth and territory establishment drives this pattern, providing a likely explanation to account for the lower rates of male adult survival that are consistently reported in this weakly dimorphic species.

Introduction

Although the difference in body mass of males and females is often considered as a fixed quantity for a given species, in reality, body mass of large herbivores varies seasonally in relation to resource abundance (Douhard et al. 2018) and the sex-specific schedule of allocation to reproduction (Apollonio et al. 2020). First, because of the greater energy requirements to sustain their larger body size, males are generally more susceptible to lose mass during periods of resource restriction, notably over winter in temperate areas (Clutton-Brock et al. 1982). Second, body condition is expected to fluctuate asynchronously between the sexes in relation to their different schedules of reproductive allocation (Stephens et al. 2009). In species that lack male parental care, females pay the costs of rearing offspring, notably during late gestation and early lactation, which generally coincide with the period of peak resource availability during late spring-early summer. Mothers can therefore offset this marked increase in energy expenditure by either increasing foraging activity (income breeder, sensu Jönsson 1997) or by mobilizing previously accumulated body reserves (capital breeder, sensu Jönsson 1997). In contrast, males must engage in contest competition to ensure access to mates, for example, through tending receptive females (Hogg 1984) or by defending a mating territory (Vanpé et al. 2009), and may lose substantial body condition as a result (Apollonio et al. 2020). In addition, to maximise their competitive ability, males must allocate substantial energy to developing secondary sexual traits including weapons such as antlers, which are

regrown annually as a prerequisite to successful reproduction. Because allocation to these elements of male-

male competition occurs earlier, typically prior to or during winter, when resources are less abundant in

temperate areas, males are expected to adopt a capital breeder tactic (sensu Jönsson 1997), accumulating fat

reserves during the season of highest resource abundance to offset the subsequent costs of reproduction

(Williams et al. 2017).

The roe deer (*Capreolus capreolus*) is a weakly polygynous species (Vanpé et al. 2008) with a low level of

sexual size dimorphism (Hewison et al. 2011) and relatively short antlers (Lemaitre et al. 2018), but where

males are strongly territorial from late winter until the end of the summer (Vanpé et al. 2009). Unusually, roe

deer males cast their antlers in late autumn which then regrow immediately over the following two to three

months, so that the costs of allocation to antler growth are levied during the heart of winter. In contrast, roe deer

females are not territorial, but are considered income breeders (Andersen et al. 2000), with very few fat reserves

(Hewison et al. 1996), that breed every year irrespective of previous reproductive status (Andersen et al. 2000,

Hewison and Gaillard 2001) and offset the annual costs of gestation and lactation during spring and summer

through concurrent intake.

While we previously showed that immature juvenile roe deer of both sexes continued to gain mass at a similar

rate over winter (Hewison et al. 2002 for two populations at 46-48°N latitude), no study has yet analyzed how

sexual mass dimorphism of mature adults is impacted by winter harshness at a broad spatial scale. We

addressed this knowledge gap by investigating how this unusual schedule of allocation to secondary sexual

traits in males shapes sex differences in body mass change over the winter and, hence, the degree of sexual size

dimorphism. We used body mass data derived from the intensive (> 7000 individuals), long-term (> 20 years)

capture-mark-recapture monitoring of five roe deer populations living under markedly different ecological

conditions to investigate the following predictions. First, because males must allocate to antler growth during

the winter months, loss of body mass should be greater (or mass gain should be lower) in males than females so

that sexual mass dimorphism is lowest at the onset of spring. Second, roe deer in the two Swedish populations

should lose more body mass than those in the three French populations because of the much harsher winter in

the north of the species' range, although this may vary among years in relation to specific annual conditions.

Materials & Methods

Study sites

We focused on five intensively monitored populations of roe deer, three in France and two in Sweden, living on

study sites with markedly different environmental conditions (Table 1). The two Swedish study sites are

situated towards the northern limit of the species' range, with harsh winter conditions, whereas the French study

sites are within the southern part of the roe deer core range and have relatively mild winters. Otherwise, the

study sites differ in terms of available habitat types, landscape structure and population density (Table 1).

Table 1: Study site characteristics of the roe deer populations. Sample size indicates the number of body mass
measurements and the number of unique individual roe deer (i.e. the ratio indicates the mean number of
measures per individual, see Table S10 for sample sizes per year). Julian date indicates when body mass was
measured where $1 =$ Jan 1^{st} (see Bonnot et al. 2024 for data and code).

	Bogesund (Sweden)	Grimsö (Sweden)	Aurignac (France)	Chizé (France)	Trois-Fontaines (France)
Latitude, Longitude	59°38′N, 18°28′E	59°73′N, 15°47′E	43°13′N, 0°52′E	46°11'N, 0°34'W	48°43'N, 4°55'E
Surface area (ha)	2 600	8 000	7 500	2 614	1 360
Habitat type	Mixed agricultural	Boreal coniferous forest	Mixed agricultural	Deciduous forest	Deciduous forest
Snow cover (days)	80	130	5	<15	<15
January temperature (°C)	3.7	-1.3	4.9	5.6	3.1
Years monitored	1989-2016	1974-2017	2001-2023	1978-2015	1976-2015
Sample size: observations (individuals)	2432 (493)	1516 (540)	503 (361)	5571 (3297)	3887 (2564)
Julian date: (start, end)	2-92	1-99	5-74	4-84	4-73

Body mass data

We collected data for all animals caught during routine capture-mark-recapture operations that took place each

winter (January to March) over two to four decades depending on the study site (see Table 1). Animals were

caught either in baited box traps (Sweden, see Kjellander et al. 2006 for more details) or drive nets (France, see

Lemaître et al. 2018 and Hewison et al. 2009 for more details). They were subsequently manipulated by

experienced handlers who recorded each individual's sex, body mass (to the nearest 0.1 kg) and age (as either

juveniles in their first winter i.e. around 8 months old, or adult i.e. older than 1.5 years old when both sexes

have attained >90% of their asymptotic body mass, Hewison et al. 2011). Juveniles can be easily distinguished

from older animals on the basis of the presence of a milk tooth at the third pre-molar (Ratcliffe & Mayle 1992).

Animals were marked with individually numbered ear tags and, in some cases, collars, before being released on

site.

 Ethical statement

 All capture and marking procedures were done in accordance with local and European animal welfare laws. For

 Aurignac-VCG: prefectural order from the Toulouse Administrative Authority to capture and monitor wild roe

 deer and agreement no. A31113001 approved by the Departmental Authority of Population Protection. For

 Bogesund and Grimsö: the marking and handling of roe deer were approved by the Ethical Committee on

 Animal Experiments, Uppsala, Sweden (Current approval Dnr: C302/2012). For Chizé and Trois-Fontaines, the

 capture protocol for roe deer under the authority of the Office Français de la Biodiversité (OFB) was approved

 by the Director of Food, Agriculture and Forest (Prefectoral order 2009-14 from Paris). All procedures were

 approved by the Ethical Committee of Lyon 1 University (project DR2014-09, June 5, 2014).

 Data analysis

 We analysed individual body mass of adult animals only in relation to sex and capture date defined as the

 number of days after 1st January (hereafter, Julian date 1, see Bonnot et al. 2024). Although captures

 occasionally took place during October, November or December, we excluded these few data so as to consider a

 common winter start date across all five populations. However, because a given Julian date cannot be

 considered strictly equivalent between France and Sweden from a phenological point of view (e.g. different

 dates for spring vegetation green-up), we performed the analysis separately for each population. Hence, while

 the analysed range for Julian date started from 1 (i.e. January 1st), the end date differed somewhat among

 populations (see Table 1). Note that, as a consequence of this choice, it was not possible to formally test our

 second hypothesis with just five independent data points (populations).

 Preliminary analysis indicated that body mass change over winter was adequately modelled as a linear function

 of date in all populations (little or no improvement in model fit when looking for non-linearity using quadratic,

 cubic or smoothing functions, see Table S6 in Appendix), and that including exact age did not influence the

 outcome (analyses restricted to known aged individuals, results not presented). Therefore, to evaluate sex-

 specific body mass trajectories over winter, we built linear mixed models with the lme4 (Bates et al. 2015)

 package in R where the full model contained sex, Julian date and their two-way interaction. We first scaled

 Julian date for each population by centering (i.e. subtracting each value from the mean Julian date) and then

 dividing it by its standard deviation. For the Aurignac-VCG population only, we also included the spatial sector

 of capture as a two-modality fixed factor (mixed vs. open habitat) to control for body mass differences in

 relation to landscape structure at this study site (i.e. roe deer heaviest in open areas, Hewison et al. 2009); note,

 we did not include animals caught in the strict forest sector because of systematic differences in capture date

 among sectors). Finally, we initially included individual identity (to control for repeated measures) and year (to

 control for annual variation in conditions) as random effects on both the intercept and the slope. While these

models successfully converged in two out of five cases, the low number of repeated measures of individuals

(Table 1) precluded convergence for the three French populations. Therefore, to investigate whether issues of

pseudo-replication might affect model selection for these populations, we re-ran the analysis on a reduced data

set that included a single observation per individual (with year as a random effect on both the intercept and

slope). As we obtained equivalent results with this approach (same model selected, essentially identical

parameter estimates), below we present the analysis based on the full data set in the main text, with the

equivalent analysis on the reduced data set provided in the Appendix (Tables S7-9, Fig. S1). We performed

model selection in relation to AIC values and weights for the candidate model set. For each population, we

retained the model with the lowest AIC value as long as it differed by at least 2 points from any simpler

competing model (see Arnold 2010).

Results

In all five populations, the best supported model describing over-winter variation in body mass consistently

included the sex by date interaction (for all five populations, ∆AIC > 3.5 compared to the second-best model),

showing that average change in body mass over winter differed between males and females (see Tables S1-S5

for scaled parameter estimates). More specifically, in the two Swedish populations, body mass (mean ± sd) of

males decreased by -21.1 g (± 3.1, Bogesund, and -21.5 g (± 3.7, Grimsö) per day between 1st January and the

end of the winter, while this decrease was much less marked for females (-4.5 ± 2.8 g and -12.0 ± 3.7 g /day,

respectively). In contrast, in the three French populations, female body mass increased by between 14.2 g (±

3.1, Trois-Fontaines) and 25.8 g (± 6.1, Aurignac-VCG) per day over winter, while that of males remained more

or less constant (from -2.3 ± 3.0 g/day at Chizé to 3.9 ± 4.3 g/day at Trois-Fontaines). As a result, while males

were clearly heavier, on average, than females at the onset of winter in all five populations, albeit more

pronouncedly in France (mean ± se: Chizé: 23.0 ± 0.2 kg for males vs. 20.7 ± 0.2 kg for females; Trois-

Fontaines: 25.0 ± 0.2 kg for males vs. 22.8 ± 0.2 kg for females; Aurignac-VCG: 23.9 ± 0.3 kg for males vs.

21.4 ± 0.2 kg for females, i.e. a sexual mass dimorphism of about 10%) than in Sweden (Bogesund: 24.8 ± 0.2

kg for males vs. 23.5 ± 0.2 kg for females; Grimsö: 26.3 ± 0.2 kg for males vs. 25.1 ± 0.2 kg for females, i.e. a

sexual mass dimorphism of about 5%), by mid-March (Julian date = 74), males did not weigh substantially

more than females in all populations (Fig. 1). Finally, at Aurignac-VCG only, the best supported model

included an additive effect of sector, indicating that deer weighed, on average, 0.81 kg (± 0.2) more in the open

sector than those in the partially wooded sector.

Fig. 1: Body mass (kg) of male (orange) and female (purple) adults (>1.5 years old) in five intensively monitored populations of roe deer situated in Sweden (Bogesund, Grimsö) and France (Chizé, Trois-Fontaines, Aurignac-VCG) in relation to date over winter. Sexual mass dimorphism (SMD) calculated as the ratio of predicted male mass to female mass at the start (1st January) and end (15th March) of the winter period is indicated for each population.

Lastly, there was some evidence that over-winter body mass change varied among years to a greater extent in

the Swedish populations compared to the French populations: the proportion of the variance attributable to the

random effect of year on the slope of the body-mass-date relationship was 2.5-9.0 times higher in Bogesund and

Grimsö than in the three French populations (Table 2).

Table 2: Variance attributed to the residual, fixed and random components of the model describing sex-specific

over-winter variation in body mass across five populations of roe deer. To facilitate reliable comparison, we

derived these variance components using an identical model structure for each population, with individual

identity as a random effect on the intercept, and year as a random effect on both the intercept and the slope. The

proportion of the total variance explained by the random effect of year on the slope is provided as a percentage.

	Residual	Fixed effects	Random effects			
			ID intercept	*Year intercept*	*Year slope*	*% variance slope(Year)*
Bogesund (SW)	1.67	0.18	5.97	0.63	0.05	0.53%
Grimsö (SW)	1.83	0.30	4.92	0.60	0.06	0.72%
Trois-Fontaines (FR)	2.52	0.88	4.20	1.05	0.01	0.09%
Chizé (FR)	1.87	0.64	4.00	0.88	0.01	0.08%
Aurignac-VCG (FR)	0.94	1.16	3.49	0.06	0.01	0.21%

183
184
185

Discussion

From the analysis of the body mass of over 7000 individual roe deer living along a gradient of ecological

conditions in terms of winter harshness, from near the northern-most extreme to the southern part of their core

geographical range, we found strong support for both our predictions, i/ that over-winter body mass change was

sex-specific whereby males lost more (or gained less) than females, so that sexual dimorphism in mass was

virtually absent by the end of the winter; ii/ but that this pattern was strongly modulated by winter severity such

that while roe deer in Sweden lost mass, on average, those in France gained mass. There was also some

indication that over-winter body mass change was more variable among years in the Swedish populations

compared to the French populations, providing additional support for the latter hypothesis. The costs of

allocation to sex-specific reproductive schedules likely drive seasonal variations in the degree of sexual mass

dimorphism in this weakly polygynous ungulate.

On the energetic cost of allocation to secondary sexual traits during winter

In polygynous mammals, reproductive effort during the mating season can lead to considerable loss of body

mass in males (Apollonio et al. 2020; e.g. in red deer (*Cervus elaphus*): Yoccoz et al. 2002; in elephant seals

(*Mirounga angustirostris*): Deutsch et al. 1990; in moose (*Alces alces*): Mysterud et al. 2005a). Indeed,

reproductive males often abstain from feeding while they court and defend females or a mating territory

(Mysterud et al. 2008). Similarly, although information on the costs of allocation to secondary sexual traits is

sparse, antlers are smaller during less favourable years (Mysterud et al. 2005b), suggesting that growing these

secondary sexual traits is costly. Here, we showed that male roe deer were consistently more constrained by

winter resource restriction than females, losing around two to four times more mass in Sweden, while gaining

up to six times less mass in France. As a result, by the onset of territorial season at the end of March (Hewison

et al. 1998), sexual dimorphism in mass was virtually absent, with the average male only weighing about half a

kilogram more than the average female across all five populations. While gestation in roe deer females begins in

late December or early January following approximately 4.5 months of diapause (Aitken 1974), substantial

allocation to foetal growth is concentrated in the latter third (April-May) so that fetuses weigh no more than a

few grams during the winter period studied here (Beyes et al. 2017). We suggest that this over-winter decrease

in sexual dimorphism of body mass is likely due to sex differences in the schedule of reproductive effort, in

particular, the energetic costs to males of growing weapons and establishing a mating territory during the most

resource-limited season (Williams et al. 2017).

On the impact of winter severity for body mass change

While roe deer are consistently heavier in Sweden than France at winter onset (Fig. 1), the severity of

conditions during the Scandinavian winter caused an average body mass loss of between 0.4 kg (females at

Bogesund) and 2.1 kg (males at Grimsö). Note that these figures are likely conservative, as resource scarcity

during winter may begin well before the New Year in northern environments depending on the annual timing of

first snowfall. For an animal of around 20-25 kg this loss is clearly considerable and indicates that the capacity

to store fat reserves and, therefore, seasonal fluctuations in body mass, are much greater in the north of its range

than previously documented for this medium-sized income breeder (Kjellander et al. 2006). Indeed, there was

also some indication in our data that over-winter body mass change varied from year to year somewhat more

markedly in Sweden compared to France (see Table 2), presumably in response to the harshness of winter

conditions in a given year. This is likely an adaptation to buffer against severe winters, as further south, in the

heart of its range, over-winter body mass is generally stable and may even increase slightly (Hewison et al.

1996, 2002). Larger body size (Linstedt & Boyce 1985) and the capacity to store fat (Trondrud et al. 2021,

Denryter et al. 2022) have been widely interpreted as adaptations which increase fasting endurance in response

to the dramatic fluctuations of resource availability in highly seasonal environments. Although differences in

the operational sex ratio across populations could theoretically influence relative priority of allocation to sexual

secondary characters, such as antlers, in polygynous systems driven by variation in the intensity of male-male competition (Weir et al. 2011), this is highly unlikely in our specific case. Indeed, the roe deer is only weakly polygynous (Vanpé et al. 2008) and the adult sex ratio (number of males/total number of males and females) is ostensibly the same across the five populations (Chizé: 0.44; Trois-Fontaines: 0.47; Aurignac-VCG: 0.41; Bogesund: 0.43; Grimsö: 0.41). Larger body size has often been reported at higher latitudes within species of mammals (Ashton et al. 2000), in line with Bergmann's rule, and is thought to reflect natural selection for greater thermoregulatory buffering in endotherms (He et al. 2023). Our data are also in line with this general pattern, but indicate that sexual selection is the ultimate driver of between-sex differences in over-winter body mass change, suggesting similar priority of energy allocation to this secondary sexual trait across hugely contrasted environments.

On the life history implications of annual body mass loss during winter

The repeated annual cycles of fat accumulation and depletion that underpin a capital breeding tactic are predicted to carry life history costs (Houston et al. 2006). While there is clear evidence to indicate that roe deer females adopt an income breeder tactic relative to other large herbivores (Andersen et al. 2000), our results imply that males must accumulate body condition prior to winter to offset the energetic costs of antler growth and subsequent territory establishment and, in this sense, can be considered capital breeders relative to females (Apollonio et al. 2020). In polygynous mammals, allocation to traits that confer an advantage in contest competition for females are predicted to impose costs in terms of survival (Clinton & Leboeuf 1993). Previous work has established that, despite the low level of polygyny in roe deer (Vanpé et al. 2008), the sex difference in annual survival of adults is equivalent to that of more polygynous and size dimorphic ungulates (Gaillard et al. 1993). We suggest that the repeated energetic cost of allocating to secondary sexual traits every winter is a proximal driver that, together with the direct costs of territorial defense and male-male competition for mates, contributes to the survival deficit for males in this weakly dimorphic ungulate. Most deer species cast and re-grow antlers during spring, when resources are plentiful (Mysterud et al. 2005b). However, because of their unusual schedule of allocation to reproduction, roe deer males are repeatedly faced with a trade-off between maintaining accumulated mass to offset the costs of establishing and defending a mating territory in spring, a full four months prior to the rut, and growing antlers during the winter season of food scarcity. The relative importance of antler size, body mass and territory quality for determining male reproductive success has yet to be established. Despite the huge among individual variation in body mass at the onset of winter within all five populations (Fig. 1, Table 2), there was little evidence for individual variation in over-winter body mass change, although we could not formally evaluate this due to the very low number of repeat measures. Future investigations of inter-individual variation in over-winter body mass change in relation to environmental severity would be highly informative for understanding individual tactics of energy allocation to secondary sexual traits and their life history consequences.

Acknowledgements:

We thank the local hunting associations, the Fédération Départementale des Chasseurs de la Haute Garonne, and numerous co-workers and volunteers for their assistance. We also thank Denis Réale, Patrick Bergeron, Phil McLoughlin and Achaz von Hardenberg for constructive comments on previous versions of this manuscript. Author contributions: A.J.M.H., N.C.B., J.M.G. & P.K. conceived the ideas, designed the study and the methodology; all authors collected the data and discussed the analytical approach; N.C.B. analysed the data.; A.J.M.H. wrote the first draft of the manuscript and all authors contributed critically to revision.

Funding:

This work was supported by an August Larsson Visiting Fellowship to A.J.M.H. and the "DivinT" and "EVORA" projects funded by the Agence Nationale de la Recherche grants ANR-22-CE02-0020-03 and ANR-22-CE02-0021-03 to J.M.G, J.F.L., M.P., N.M. and A.J.M.H.

Conflict of interest disclosure:

The authors declare they have no conflict of interest relating to the content of this article. A.J.M.H. and N.C.B. are recommenders for PCIEcology.

Data, scripts, code and supplementary information availability:

Data and code can be found at Bonnot et al. (2024), while supplementary information is given in an Appendix at the end of this article.

References

Aitken, R.J. (1974) Delayed implantation in roe deer (*Capreolus capreolus*). *Journal of Reproduction and Fertility,* 39, 225-233

Andersen, R., Gaillard, J.M., Linnell, J.D.C., & Duncan, P. (2000) Factors affecting maternal care in an income breeder, the European roe deer. *Journal of Animal Ecology,* 69, 672-682.

Apollonio. M., Merli, E., Chirichella, R., Pokorny, B., Alagic, A., Flajšman, K. & Stephens, P.A. (2020) Capital-income breeding in male ungulates: causes and consequences of strategy differences among species. *Frontiers in Ecology and Evolution*, 8, 521767.

Arnold, T.W. (2010) Uninformative parameters and model selection using Akaike's information criterion. *Journal of Wildlife Management,* 74, 1175-1178.

Ashton, K.G., Tracy, M.C. & de Queiroz, A. (2000) Is Bergmann's rule valid for mammals? *The American Naturalist*, 156, 390-415.

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software,* 67, 1-48.

Beyes, M., Nause, N., Bleyer, M., Kaup, F.J. & Neumann, S. (2017) Description of post-implantation embryonic stages in roe deer (*Capreolus capreolus*) after embryonic diapause. *Anatomia Histologia Embryologia*, 46, 582-591.

Bonnot, N.C., Hewison, A.J.M. & Gaillard, J.M. Kjellander, P., Lemaitre, J.F., Morellet, N. & Pellerin, M. (2024) Data set and code for the paper: "Body mass change over winter is consistently sex-specific across roe deer (Capreolus capreolus) populations", Hewison et al. [Data set]. Zenodo. https://doi.org/10.5281/zenodo.10997219.

Clinton, W.L. & Leboeuf, B.J. (1993) Sexual selection effects on male life-history and the pattern of male mortality. *Ecology*, 74, 1884-1892.

Denryter, K., Conner, M. M., Stephenson, T. R., German, D. W. & Monteith, K. L. (2022) Survival of the fattest: How body fat and migration influence survival in highly seasonal environments. *Functional Ecology*, 36, 2569–2579.

Deutsch, C.J., Haley, M.P. & Leboeuf, B.J. (1990) Reproductive effort of male northern elephant seals: estimates from mass loss. *Canadian Journal of Zoology*, 68, 2580-2593.

Douhard, M., Guillemette, S., Festa-Bianchet, M. & Pelletier, F. (2018) Drivers and demographic consequences of seasonal mass changes in an alpine ungulate. *Ecology,* 99, 724-734.

Gaillard, J.M., Delorme, D., Boutin, J.M., Van Laere, G., Boisaubert, B., & Pradel, R. (1993) Roe deer survival patterns: a comparative analysis of contrasting populations. *Journal of Animal Ecology*, *62*, 778-791.

He, J., Tu, J., Yu, J. & Jiang, H. (2023) A global assessment of Bergmann's rule in mammals and birds. *Global Change Biology*, 29, 5199-5210.

Hewison, A.J.M. & Gaillard, J.M. (2001) Phenotypic quality and senescence affect different components of reproductive output in roe deer. *Journal of Animal Ecology*, 70, 600-608.

Hewison, A.J.M., Angibault, J.M., Boutin, J.M., Bideau, E., Vincent, J.P. & Sempéré, A. (1996) Annual variation in body composition of roe deer (*Capreolus capreolus*) in moderate environmental conditions. *Canadian Journal of Zoology*, 74, 245-253.

Hewison, A.J.M., Vincent, J.P., & Reby, D. (1998) Social organisation of European roe deer. In R. Andersen, P. Duncan, & J. D. C. Linnell (Eds.), *The European roe deer: The biology of success* (pp. 189–219). Scandinavian University Press, Oslo.

Hewison, A.J.M., Gaillard, J.M., Angibault, J.M., Van Laere, G., & Vincent, J.P. (2002) The influence of density on post-weaning growth in roe deer (*Capreolus capreolus*) fawns. *Journal of Zoology, 257*, 303-309.

Hewison, A. J. M., Morellet, N., Verheyden, H., Daufresne, T., Angibault, J.-M., Cargnelutti, B., Merlet, J., Picot, D., Rames, J.-L., Joachim, J., Lourtet, B., Serrano, E., Bideau, E., & Cebe, N. (2009) Landscape fragmentation influences winter body mass of roe deer. *Ecography*, 32, 1062–1070.

Hewison, A.J.M., Gaillard, J.M., Delorme, D., Van Laere, G., Amblard, T., & Klein, F. (2011) Reproductive constraints, not environmental conditions, shape the ontogeny of sex-specific mass-size allometry in roe deer. *Oikos*, 120, 1217-1226.

Hogg, J.T. (1984) Mating in bighorn sheep – multiple creative male strategies. *Science*, 225, 526-529.

Houston, A.I., Stephens, P.A., Boyd, I.L., Harding, K.C. & McNamara, J.M. (2007) Capital or income breeding? A theoretical model of female reproductive strategies. *Behavioral Ecology,* 18, 241–250.

Jonsson, K.I. (1997). Capital and income breeding as alternative tactics of resource use in reproduction. *Oikos,* 78, 57-66.

Kjellander, P., Gaillard, J.M. & Hewison, A.J.M. (2006) Density-dependent responses of fawn cohort body mass in two contrasting roe deer populations. *Oecologia*, 146, 521-530.

Lemaître, J-F., Cheynel, L., Douhard, F., Bourgoin, G., Débias, F., Ferté, H., Gilot-Fromont, E., Pardonnet, S., Pellerin, M., Rey, B., Vanpé, C., Hewison, A.J.M. & Gaillard, J.M. (2018) The influence of early-life allocation to antlers on male performance during adulthood: Evidence from contrasted populations of a large herbivore. *Journal of Animal Ecology* 87, 921-932.

Linstedt, S. & Boyce, M. (1985) Seasonality, fasting endurance, and body size in mammals. *American Naturalist,* 125, 873-878.

Mysterud, A., Solberg, E.J. & Yoccoz, N.G. (2005a) Aging and reproductive effort in male moose under variable levels of intersexual competition. *Journal of Animal Ecology,* 74, 742-754.

Mysterud, A., Meisingset, E., Langvatn, R., Yoccoz, N.G. & Stenseth, N.C. (2005b). Climate-dependent allocation of resources to secondary sexual traits in red deer. *Oikos,* 111, 245-252.

Mysterud, A., Bonenfant, C., Loe, L.E., Langvatn, R., Yoccoz, N.G. & Stenseth, N.C. (2008). Age-specific feeding cessation in male red deer during rut. *Journal of Zoology,* 275, 407-412.

Nilsen, E.B., Gaillard, J.M., Andersen, R., Odden, J., Delorme, D., van Laere, G. & Linnell, J.D.C. (2009) A slow life in hell or a fast life in heaven: demographic analyses of contrasting roe deer populations *Journal of Animal Ecology,* 78, 585-594.

Stephens, P. A., Boyd, I. L., McNamara, J. M., & Houston, A. I. (2009) Capital breeding and income breeding: Their meaning, measurement, and worth. *Ecology*, 90, 2057–2067.

Ratcliffe, P.R. & Mayle, B. (1992) Roe deer biology and management. *Forestry Commission Bulletin,* 105, HMSO, London.

Trondrud, L.M., Pigeon, G., Krol, E., Albon, S., Evans, A.L., Arnold, W., Hambly, C., Irvine, R.J., Ropstad, E., Stien, A., Veiberg, V., Speakman, J.R. & Loe, L.E. (2021) Fat storage influences fasting endurance more than body size in an ungulate. *Functional Ecology,* 35, 1470–1480.

Vanpé, C., Kjellander, P., Galan, M., Cosson, J.F., Aulagnier, S., Liberg, O., & Hewison, A.J.M. (2008) Mating system, sexual dimorphism and the opportunity for sexual selection in a territorial ungulate. *Behavioural Ecology*, 19, 309-316.

Vanpé, C., Morellet, N., Kjellander, P., Goulard, M., Liberg, O., & Hewison, A.J.M. (2009a). Access to mates in a territorial ungulate is determined by the size of a male's territory, but not by its habitat quality. *Journal of Animal Ecology*, 78, 42–51.

Weir, L.K., Grant, J.W.A. & Hutchings, J.A. (2011) The influence of operational sex ratio on the intensity of competition for mates. *American Naturalist,* 177, 167-176.

Williams, C.T., Klaassen, M., Barnes, B.M., Buck, C.L., Arnold, W., Giroud, S., Vetter, S.G. & Ruf, T. (2017) Seasonal reproductive tactics: annual timing and the capital-to-income breeder continuum. *Philosophical Transactions of the Royal Society B*, 372, 20160250.

Yoccoz, N.G., Mysterud, A., Langvatn, R. & Stenseth, N.C. (2002) Age- and density-dependent reproductive effort in male red deer. *Proceedings of the Royal Society of London B*, 269, 1523-1528.

<p>384</p>

Appendix

<p>385</p>

<p>386</p> *1.1 Model selection:*

<p>387</p>

<p>388</p> Table S1a: Model fit and selection (fixed and random effects, difference in AIC score compared to the best
<p>389</p> model, AIC weight) describing sex-specific over-winter variation in body mass in the **Bogesund** population.
<p>390</p> The selected model is shaded grey.

<p>391</p>

Fixed effects	Random intercept	Random slope	ΔAIC	weight
Sex + Julian date + Sex:Julian date	ID & Year	ID & Year	0.0	1.00
Sex + Julian date	ID & Year	ID & Year	26.2	0
Julian date	ID & Year	ID & Year	27.2	0
Sex	ID & Year	ID & Year	37.7	0
null	ID & Year	ID & Year	38.4	0

<p>392</p>
<p>393</p>
<p>394</p>

<p>395</p> Table S1b: Model summary for the selected model: parameter coefficients for the fixed effects (mean and
<p>396</p> standard error, s.e.) and variance components for the fixed, random and residual effects, describing sex-specific
<p>397</p> over-winter variation in body mass in the **Bogesund** population. Note that Julian date was centered and scaled
<p>398</p> (see text). The reference category is female.

<p>399</p>

Parameter	Coefficient	s.e.	Variance
Fixed effects			**0.17**
Intercept	23.38	0.22	
Sex (male)	0.55	0.24	
Julian date	-0.10	0.06	
Sex (male):Julian date	-0.37	0.07	
Random effects			**6.65**
Individual identity (intercept)			5.92
Individual identity (slope)			0.05
Year (intercept)			0.63
Year (slope)			0.04
Residual			**1.63**

<p>400</p>

Table S2a: Model fit and selection (fixed and random effects, difference in AIC score compared to the best model, AIC weight) describing sex-specific over-winter variation in body mass in the **Grimsö** population. The selected model is shaded grey.

Fixed effects	Random intercept	Random slope	ΔAIC	weight
Sex + Julian date + Sex:Julian date	ID & Year	ID & Year	0.0	0.90
Sex + Julian date	ID & Year	ID & Year	4.4	0.01
Julian date	ID & Year	ID & Year	15.1	0.00
Sex	ID & Year	ID & Year	27.4	0.00
null	ID & Year	ID & Year	37.3	0.00

Table S2b: Model summary for the selected model: parameter coefficients for the fixed effects (mean and standard error, s.e.) and variance components for the fixed, random and residual effects, describing sex-specific over-winter variation in body mass in the **Grimsö** population. Note that Julian date was centered and scaled (see text). The reference category is female.

Parameter	Coefficient	s.e.	Variance
Fixed effects			**0.30**
Intercept	24.56	0.19	
Sex (male)	0.74	0.22	
Julian date	-0.30	0.08	
Sex (male):Julian date	-0.24	0.09	
Random effects			**5.65**
Individual identity (intercept)			4.92
Individual identity (slope)			0.06
Year (intercept)			0.60
Year (slope)			0.06
Residual			**1.76**

Table S3a: Model fit and selection (fixed and random effects, difference in AIC score compared to the best model, AIC weight,) describing sex-specific over-winter variation in body mass in the **Aurignac-VCG** population. The selected model is shaded grey.

Fixed effects	Random intercept	Random slope	ΔAIC	weight
Sex + Julian date + Sex:Julian date + Sector	ID & Year	Year	0.0	0.87
Sex + Julian date + Sector	ID & Year	Year	4.0	0.12
Sex + Sector	ID & Year	Year	9.9	0.01
Sex + Julian date + Sex:Julian date	ID & Year	Year	12.0	0.00
Sex + Julian date	ID & Year	Year	15.3	0.00
Sex	ID & Year	Year	22.4	0.00
Julian date + Sector	ID & Year	Year	67.1	0.00
Sector	ID & Year	Year	73.8	0.00
Julian date	ID & Year	Year	82.2	0.00
null	ID & Year	Year	88.7	0.00

Table S3b: Model summary for the selected model: parameter coefficients for the fixed effects (mean and standard error, s.e.) and variance components for the fixed, random and residual effects, describing sex-specific over-winter variation in body mass in the **Aurignac-VCG** population. Note that Julian date was centered and scaled (see text). The reference category is female and the mixed sector.

Parameter	Coefficient	s.e.	Variance
Fixed effects			**1.16**
Intercept	21.30	0.20	
Sex (male)	1.87	0.22	
Julian date	0.43	0.10	
Sector (open)	0.81	0.21	
Sex (male):Julian date	-0.37	0.15	
Random effects			**3.56**
Individual identity (intercept)			3.49
Year (intercept)			0.06
Year (slope)			0.01
Residual			**0.94**

426 Table S4a: Model fit and selection (fixed and random effects, difference in AIC score compared to the best
427 model, AIC weight,) describing sex-specific over-winter variation in body mass in the **Chizé** population. The
428 selected model is shaded grey.

Fixed effects	Random intercept	Random slope	ΔAIC	weight
Sex + Julian date + Sex:Julian date	ID & Year	Year	0.0	1.0
Sex + Julian date	ID & Year	Year	47.2	0.0
Sex	ID & Year	Year	66.2	0.0
Julian date	ID & Year	Year	375.9	0.0
null	ID & Year	Year	396.3	0.0

433 Table S4b: Model summary for the selected model: parameter coefficients for the fixed effects (mean and
434 standard error, s.e.) and variance components for the fixed, random and residual effects, describing sex-specific
435 over-winter variation in body mass in the **Chizé** population. Note that Julian date was centered and scaled (see
436 text). The reference category is female.

Parameter	Coefficient	s.e.	Variance
Fixed effects			**0.64**
Intercept	21.44	0.16	
Sex (male)	1.52	0.08	
Julian date	0.33	0.04	
Sex (male):Julian date	-0.37	0.05	
Random effects			**4.88**
Individual identity (intercept)			4.00
Year (intercept)			0.88
Year (slope)			0.01
Residual			**1.87**

Table S5a: Model fit and selection (fixed and random effects, difference in AIC score compared to the best model, AIC weight,) describing sex-specific over-winter variation in body mass in the **Trois-Fontaines** population. The selected model is shaded grey.

Fixed effects	Random intercept	Random slope	ΔAIC	weight
Sex + Julian date + Sex:Julian date	ID & Year	Year	0.0	0.86
Sex + Julian date	ID & Year	Year	3.7	0.14
Sex	ID & Year	Year	14.1	0.00
Julian date	ID & Year	Year	322.1	0.00
null	ID & Year	Year	333.0	0.00

Table S5b: Model summary for the selected model: parameter coefficients for the fixed effects (mean and standard error, s.e.) and variance components for the fixed, random and residual effects, describing sex-specific over-winter variation in body mass in the **Trois-Fontaines** population. Note that Julian date was centered and scaled (see text). The reference category is female.

Parameter	Coefficient	s.e.	Variance
Fixed effects			**0.88**
Intercept	23.28	0.18	
Sex (male)	1.83	0.10	
Julian date	0.23	0.05	
Sex (male):Julian date	-0.17	0.07	
Random effects			**5.25**
Individual identity (intercept)			4.20
Year (intercept)			1.05
Year (slope)			0.01
Residual			**2.52**

1.2 Test for non-linear variation in over-winter body mass

In preliminary analysis, in addition to a linear function, we modelled body mass change over winter with quadratic, cubic or smoothing functions of date using the 'lme4' and 'gamm4' packages (Bates et al. 2015; Wood & Scheipl 2020) in R. All models included sex, Julian date and their two-way interaction, as well as individual identity (to control for repeated measures on individuals) and year (to control for annual variation in conditions) as random factors. In three out of five cases, the linear function provided the best fit (Table S6), whereas the cubic model provided a somewhat better fit in the Grimsö population, while the quadratic model performed slightly better in the Aurignac-VCG population. Because this improvement was marginal from a biological point of view, and to facilitate comparison among populations, we present results from linear models in the main text, i.e. assuming that the rate of change in body mass during winter is constant over the entire study window.

Wood, S. & Scheipl, F. (2020) gamm4: Generalized Additive Mixed Models using 'mgcv' and 'lme4'. R package version 0.2-6.

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software,* 67, 1-48.

Table S6: Comparison of model fit (AIC values) for the best supported model describing sex-specific over-winter variation in body mass in five roe deer populations (i.e. mass ~ sex * Julian date, with an additive effect of sector for the Aurignac-VCG population only, see main text) when the relationship between body mass and date was modelled as either a linear effect, a quadratic effect, a cubic effect, or as a smoothing spline in a general additive mixed model framework). The selected model is indicated in bold.

Population	linear	quadratic	cubic	GAMM
Bogesund	**9526.9**	9553.2	9530.4	9543.0
Grimsö	6348.3	6348.6	**6342.7**	6364.8
Aurignac-VCG	2057.3	**2056.3**	2058.6	2130.5
Chizé	**24095.6**	24146.1	24102.4	24439.6
Trois-Fontaines	**17777.2**	17782.8	17783.7	18109.9

 2.1 Model selection on a reduced data set, featuring a single observation per individual (with year as a random
effect on both the intercept and slope) for the three populations where the full model (which also included a
random effect of individual identity on both the intercept and slope) did not converge

Table S7: Model fit and selection (fixed and random effects, difference in AIC score compared to the best
model, AIC weight) on the reduced data set, describing sex-specific over-winter variation in body mass in the
Aurignac-VCG population. The selected model is shaded grey.

Fixed effects	Random intercept & slope	ΔAIC	weight
Sex + Julian date + Sex:Julian date + Sector	Year	0.0	0.63
Sex + Julian date + Sector	Year	2.0	0.23
Sex + Sector	Year	3.1	0.13
Sex + Julian date + Sex:Julian date	Year	9.6	0.01
Sex + Julian date	Year	11.1	0.00
Sex	Year	14.7	0.00
Julian date + Sector	Year	64.8	0.00
Sector	Year	65.5	0.00
Julian date	Year	79.0	0.00
null	Year	82.3	0.00

Table S8: Model fit and selection (fixed and random effects, difference in AIC score compared to the best
model, AIC weight) on the reduced data set, describing sex-specific over-winter variation in body mass in the
Chizé population. The selected model is shaded grey.

Fixed effects	Random intercept & slope	ΔAIC	weight
Sex + Julian date + Sex:Julian date	Year	0.0	1.0
Sex + Julian date	Year	26.7	0.0
Sex	Year	31.2	0.0
Julian date	Year	329.9	0.0
null	Year	335.9	0.0

Table S9: Model fit and selection (fixed and random effects, difference in AIC score compared to the best
model, AIC weight) on the reduced data set, describing sex-specific over-winter variation in body mass in the
Trois-Fontaines population. The selected model is shaded grey.

Fixed effects	Random intercept & slope	ΔAIC	weight
Sex + Julian date + Sex:Julian date	Year	0.0	0.50
Sex + Julian date	Year	0.3	0.42
Sex	Year	3.7	0.08
Julian date	Year	291.9	0.00
null	Year	298.3	0.00

2.2 Estimated over-winter changes in body mass in the three French populations (Chizé, Trois-Fontaines, Aurignac-VCG) based on a reduced data set, featuring a single observation per individual

Fig. S1: Body mass (kg) of male (purple) and female (orange) adults (>1.5 years old) in three intensively monitored populations of roe deer situated in France (Chizé, Trois-Fontaines, Aurignac-VCG) in relation to date over winter (based on the model selection presented in Tables S7-S9).

Table S10: Sample size per year and sex for the five populations of roe deer situated in Sweden (Bogesund, Grimsö) and France (Aurignac-VCG , Chizé, Trois-Fontaines), with the number of unique individuals measured that year in brackets.

Year	Grimsö		Bogesund		Aurignac-VCG		Chizé		Trois-Fontaines	
	F	M	F	M	F	M	F	M	F	M
1974	3 (2)	3 (3)	/	/	/	/	/	/	/	/
1975	4 (4)	1 (1)	/	/	/	/	/	/	/	/
1976	2 (2)	3 (3)	/	/	/	/	/	/	23 (23)	23 (23)
1977	11 (11)	6 (6)	/	/	/	/	/	/	53 (53)	55 (55)
1978	2 (2)	4 (4)	/	/	/	/	41 (41)	24 (24)	39 (39)	41 (41)
1979	6 (6)	1 (1)	/	/	/	/	53 (53)	32 (32)	57 (57)	59 (59)
1980	2 (2)	3 (3)	/	/	/	/	56 (56)	49 (49)	60 (60)	44 (44)
1981	/	/	/	/	/	/	66 (66)	56 (56)	39 (39)	45 (45)
1982	/	/	/	/	/	/	68 (68)	55 (55)	66 (66)	41 (40)
1983	/	/	/	/	/	/	87 (87)	68 (67)	75 (75)	56 (56)
1984	/	1 (1)	/	/	/	/	117 (117)	76 (76)	72 (72)	58 (58)
1985	8 (8)	/	/	/	/	/	79 (79)	65 (65)	80 (78)	62 (62)
1986	3 (3)	1 (1)	/	/	/	/	127 (127)	70 (70)	79 (79)	63 (63)
1987	41 (20)	16 (9)	/	/	/	/	97 (97)	68 (68)	54 (54)	46 (46)
1988	16 (10)	10 (8)	/	/	/	/	73 (73)	54 (54)	37 (37)	42 (42)
1989	12 (9)	3 (3)	6 (4)	1 (1)	/	/	74 (74)	58 (58)	38 (38)	33 (33)
1990	28 (19)	28 (11)	18 (11)	21 (11)	/	/	66 (66)	52 (52)	53 (53)	44 (44)
1991	36 (24)	24 (16)	18 (13)	12 (11)	/	/	64 (64)	54 (54)	58 (58)	43 (43)
1992	33 (16)	30 (15)	43 (36)	26 (20)	/	/	72 (72)	55 (55)	55 (55)	38 (38)
1993	36 (19)	32 (19)	21 (13)	12 (9)	/	/	58 (58)	39 (39)	42 (42)	39 (39)
1994	58 (31)	43 (20)	8 (8)	12 (12)	/	/	49 (49)	40 (40)	52 (52)	45 (45)
1995	43 (25)	30 (13)	32 (25)	19 (12)	/	/	55 (55)	39 (39)	45 (45)	49 (49)
1996	37 (22)	21 (12)	49 (33)	26 (15)	/	/	55 (55)	41 (41)	39 (39)	41 (41)
1997	40 (17)	10 (6)	87 (41)	34 (19)	/	/	67 (67)	63 (63)	37 (37)	40 (40)
1998	17 (17)	6 (6)	7 (7)	16 (10)	/	/	81 (81)	89 (89)	34 (34)	29 (29)
1999	25 (20)	7 (6)	36 (24)	31 (16)	/	/	62 (62)	74 (74)	40 (40)	26 (26)
2000	9 (9)	5 (4)	34 (23)	19 (11)	/	/	48 (48)	46 (46)	8 (8)	10 (10)
2001	26 (21)	13 (10)	61 (36)	59 (34)	/	2 (2)	85 (84)	77 (77)	47 (47)	37 (37)
2002	8 (8)	8 (7)	46 (29)	57 (34)	5 (5)	5 (5)	83 (83)	65 (65)	44 (44)	59 (59)
2003	14 (14)	3 (3)	37 (26)	44 (29)	8 (8)	5 (5)	91 (90)	77 (77)	63 (63)	69 (69)
2004	23 (17)	8 (6)	62 (42)	66 (37)	/	/	136 (136)	99 (99)	64 (64)	75 (75)
2005	15 (14)	12 (10)	95 (31)	92 (34)	12 (12)	8 (8)	126 (126)	93 (93)	85 (85)	75 (75)
2006	36 (29)	29 (19)	62 (28)	64 (19)	14 (14)	12 (12)	127 (127)	111 (111)	57 (57)	51 (51)
2007	31 (28)	19 (14)	22 (12)	19 (9)	13 (13)	5 (5)	79 (79)	69 (69)	33 (33)	38 (38)
2008	16 (14)	8 (7)	16 (11)	11 (7)	15 (15)	5 (5)	85 (85)	51 (51)	36 (36)	25 (25)
2009	40 (29)	23 (17)	30 (14)	15 (12)	12 (12)	8 (8)	79 (79)	73 (73)	37 (37)	30 (30)
2010	58 (29)	37 (22)	97 (43)	73 (21)	15 (14)	10 (10)	108 (108)	73 (73)	45 (45)	36 (36)
2011	39 (22)	28 (14)	93 (36)	51 (26)	16 (16)	11 (11)	111 (110)	91 (91)	62 (62)	41 (41)
2012	16 (13)	14 (13)	64 (27)	44 (18)	19 (19)	8 (8)	90 (90)	61 (61)	42 (42)	37 (37)
2013	21 (16)	24 (19)	110 (37)	75 (25)	13 (13)	10 (10)	110 (110)	82 (82)	59 (59)	55 (55)
2014	21 (14)	20 (14)	46 (20)	48 (12)	17 (17)	8 (8)	94 (94)	77 (77)	58 (58)	52 (52)
2015	19 (15)	26 (17)	80 (22)	43 (12)	18 (17)	11 (11)	108 (108)	78 (78)	42 (42)	38 (38)
2016	18 (10)	35 (24)	96 (18)	66 (12)	15 (15)	12 (12)	/	/	49 (49)	39 (39)
2017	23 (16)	25 (17)	/	/	19 (19)	7 (7)	/	/	/	/
2018	/	/	/	/	14 (14)	15 (14)	/	/	/	/
2019	/	/	/	/	12 (12)	12 (12)	/	/	/	/
2020	/	/	/	/	16 (16)	15 (15)	/	/	/	/
2021	/	/	/	/	13 (13)	9 (9)	/	/	/	/
2022	/	/	/	/	16 (16)	13 (13)	/	/	/	/
2023	/	/	/	/	16 (16)	14 (14)	/	/	/	/

514