Tethered agonist activated ADGRF1 structure reveals molecular preference for $G\alpha_q$

signalling

Daniel T. D. Jones[1], Andrew N. Dates[1], Shaun D. Rawson[1], Maggie M. Burruss[1], Colin H. Lipper[1]

and Stephen C. Blacklow[1,2]*

[1]Department of Biological Chemistry and Molecular Pharmacology, Blavatnik Institute, Harvard

Medical School, Boston, MA 02115, USA

[2]Department of Cancer Biology, Dana Farber Cancer Institute, Boston, MA 02215, USA

*Lead contact. Correspondence: stephen_blacklow@hms.harvard.edu and

daniel_jones@hms.harvard.edu.

Abstract

Adhesion G-Protein Coupled Receptors (aGPCRs) have evolved an activation mechanism to translate extracellular force into liberation of a tethered agonist (TA) to modulate cell signalling. We report here that ADGRF1 is the first class B GPCR shown to signal through all major G-protein classes and identify the structural basis for its $G\alpha_q$ preference by cryo-EM. Our structure shows that $G\alpha_q$ over $G\alpha_s$ preference in ADGRF1 derives from tighter packing at the conserved F569 of the TA, altering contacts between TM helix I and VII, with a concurrent rearrangement of TM helices VII and VIII at the site of $G\alpha$ recruitment. $G\alpha_s$ signalling is also more sensitive to mutation of TA or binding site residues than $G\alpha_q$. Our work advances the understanding of aGPCR TA activation in molecular detail, identifying structural features that potentially explain preferential signal modulation.

Main

Adhesion G protein coupled receptors (aGPCRs) are the second largest class of GPCRs, constituting 33 members across 9 subfamilies [1]. aGPCRs control a multitude of cellular processes involved in organ development and tissue homeostasis, responding to external stimuli that sense extracellular physical forces, such as binding to protein ligands presented on cells, components of extracellular matrix and shear flow[2]. Mutations in aGPCRs are genetically responsible for human developmental disorders, such as ADGRV1 in Usher syndrome[3], and EMR2 in vibratory urticaria[4].

aGPCRs have a unique modular architecture, with an N-terminal ectodomain containing extracellular adhesive modules, followed by a G-protein coupled receptor autoinducing (GAIN) domain linked to a seven-transmembrane (7TM) bundle that relays extracellular events to the cell. A distinguishing hallmark of aGPCRs is that the GAIN domain undergoes autoproteolysis at a short hairpin turn between the final two beta-strands, separating the aGPCR into non-covalently attached N-terminal and C-terminal fragments, named NTF and CTF respectively[5,6]. Currently, models for aGPCR activation suggest that mechanical force applied to the adhesive modules in aGPCRs induces dissociation of the NTF from the CTF[7,8], enabling the newly liberated N-terminal end of the CTF to act as an intramolecular agonist to the 7TM domain. Intramolecular ligation of this tethered agonist (TA) to the 7TM domain relays the initial extracellular cue to the cell through G-protein coupling or beta-arrestin activity.

The first structure of an aGPCR 7TM domain was solved for ADGRG3 (GPR97) bound to glucocorticoids in the orthosteric pocket, revealing key residues involved in ligand recognition, conformational switches important for receptor activation, and residues for coupling to G-protein[9]. Cryo-EM structures were then reported for seven different aGPCR family members activated by their TAs, including representative GPCR-G-protein complexes for all major families of $G\alpha$, except for $G\alpha_q$ [10–13]. These studies showed that the TA acts an intramolecular ligand for the 7TM domain and identified a canonical binding pose for a conserved hydrophobic interaction motif (HIM) of the TA in the orthosteric site.

ADGRF1 (GPR110) is an aGPCR belonging to aGPCR subfamily VI [1,14]. ADGRF1 was identified as an oncogene overexpressed in lung and prostate cancer[15], and has been implicated in synaptamide dependent neural outgrowth and repair[16,17], though this activity remains

controversial[18]. ADGRF1 was used as a prototypical aGPCR to determine that membranes

expressing ADGRF1 treated with urea at high concentration induced shedding of the NTF,

liberating the TA to activate the 7TM and recruit $G\alpha_q$ preferentially over other $G\alpha$ proteins[8].

Further work has also showed that mouse ADGRF1 can be activated by synthetic TA peptides when

added in trans, signalling through both $G\alpha_s$ and $G\alpha_q$[19].

Recent structures of the TA-engaged state of the ADGRF1 CTF coupled to either $miniG\alpha_s$

and $miniG\alpha_i$ also identified a lysophosphatidylcholine (LPC) lipid bound at the intracellular

convergence of TM helices II, III and IV[13]. Interestingly, fragments of ADGRF1 expressed with the

GAIN domain included also yielded TA-engaged structures of ADGRF1 coupled to miniG but lacked

density for the GAIN domain, suggesting that the CTF may have dissociated from the NTF portion

of the protein during purification.

To understand the molecular events driving TA activation of ADGRF1 and how its structure

determines $G\alpha$ class preference, we determined a cryo-EM structure of TA-activated ADGRF1

coupled to $miniG\alpha_{s/q}$-$\beta_1\gamma_2$, stabilised by Nb35. Combining structural insights with extensive

functional studies, we identify key residues of the orthosteric binding site that are essential for

$G\alpha_s$ signalling but dispensable for $G\alpha_q$ signalling. Overall, our work complements other recent

aGPCR structures and extends them by visualizing ADGRF1 in complex with its most relevant G-

protein partner, elucidating a functional and structural landscape by which ADGRF1 is activated

and uncovering how its structure determines $G\alpha$ class preference. Furthermore, the mechanistic

insights determined here regarding TA engagement to strong and weakly associated G-proteins

offers valuable insight into how TA mediated activation of aGPCRs may be functionally modulated.

Results

ADGRF1 signals through all G-protein classes

We recently developed a platform for analysis of the signalling activity of aGPCRs, in which the CTF

of interest is expressed as a protein fusion with an N-terminal IL2 signal sequence followed by

maltose binding protein (MBP) and a tobacco etch virus cleavage site (denoted MBP-CTF) [20]

(Extended Data Fig. 1). Using this approach to measure signalling activity in reporter assays, we

observed transcriptional responses with both CRE and NFAT reporters, indicative respectively of

$G\alpha_s$ and $G\alpha_q$ signalling, consistent with previous studies[13,21]. We also observed transcriptional

responses with the SRE and SRF-RE reporters, which suggests that ADGRF1 can also signal through

$G\alpha_{i/o}$ and $G\alpha_{12/13}$ (Fig. 1a).

To further evaluate the potential for signalling through $G\alpha_{12/13}$ (a structure of $G\alpha_i$ has been

reported[13]) we used HEK293T cells with deletion of six $G\alpha$ subunits ($G\alpha_s$ short, $G\alpha_s$ long, $G\alpha_q$,

$G\alpha_{11}$, $G\alpha_{16}$, $G\alpha_{12}$ and $G\alpha_{13}$, gift from Associate Prof. A. Inoue, Tohoku university, Japan)[22], and

deletion complementation by transient transfection to determine which $G\alpha$ subunits support

ADGRF1 signalling in the SRF-RE reporter assay. ADGRF1 MBP-CTF signals strongly through $G\alpha_q$

and $G\alpha_{11}$, and also exhibits a smaller significant signal for $G\alpha_{12}$ and $G\alpha_{13}$ (Fig. 1b). These data show

that ADGRF1 has the functional capacity to signal through all $G\alpha$ classes, making ADGRF1 the first

of the class B GPCRs shown to have this capability[23].

Protein engineering of ADGRF1 and Cryo-EM structure determination

To obtain an active state ADGRF1 G-protein protein complex amenable for structural and

functional investigation, we probed complex assembly with a suite of miniGα proteins[24,25], using

a NanoBiT recruitment assay[26, 27]. We found that LgBiT-miniGα_s, miniG$\alpha_{s/q}$ and miniGα_o were all

recruited to C-terminal SmBiT-tagged ADGRF1-CTF as judged by increased luminescence upon

SmBiT-LgBiT complementation, whereas LgBiT-miniGα_{12}, and matched CΔ5 controls were not

(Extended Data Fig. 2a). To assess tractibility for structure determination, we co-expressed these

constructs in Expi293T suspension cells, and also observed a similar pattern of $G\alpha$ recruitment to

ADGRF1 in these cells (Extended Data Fig. 2b, left). To further test the stability of the complex, we

solubilised cells with mild detergent supplemented with cholesterol in the presence of apyrase.

The only complex that retained luminescence under these lysis conditions was the combination of

SmBiT-tagged ADGRF1-CTF with LgBiT-miniG$\alpha_{s/q}$ (Extended Data Fig. 2b, right). In addition, only

miniG$\alpha_{s/q}$ specifically purified with ADGRF1-CTF using anti-FLAG M2 magnetic beads (Extended

Data Fig. 2c), indicating that ADGRF1-Gα complex stability is greater for miniG$\alpha_{s/q}$ compared to

miniGα_s and miniGα_o. Furthermore, miniG$\alpha_{s/q}$ co-expression with ADGRF1-CTF appears to

increase the amount of receptor purified, suggesting that the miniG$\alpha_{s/q}$ may improve receptor

expression by facilitating the folding of the receptor, increasing its stability, or by blocking toxic

signalling through endogenous signalling pathways.

Structure of the ADGRF1-miniG$\alpha_{s/q}$ heterotrimeric G protein complex with the Nb35 nanobody

To elucidate the basis for selective recruitment of G$\alpha_{s/q}$, we complexed ADGRF1 CTF with

a miniG$\alpha_{s/q}$ heterotrimeric G protein complex and the Nb35 nanobody, purified the complex by

size-exclusion chromatography (Fig. 2a), and determined its structure by cryo-EM to 3.4 Å

resolution (Extended Data Fig. 3-4). The cryo-EM map was of sufficient resolution to permit the

building of an atomic model for the receptor, G-protein complex, and nanobody (Fig. 2b, Extended

Data Fig. 5). The cryo-EM map in the orthosteric pocket on the extracellular face of the receptor

permitted unambiguous modelling of the TA (Extended Data Fig. 5), which adopts a short alpha-

helical motif with a loop capping the TA *en route* to TM1, akin to other resolved TA structures of

aGPCRs[10–13]. The polypeptide backbone of the 7TM domain of ADGRF1 bound to miniG$\alpha_{s/q}$ has an

RMSD of 0.6 Å when aligned to the 7TM domain of ADGRF1 bound to either miniGα_s and miniGα_i,

highlighting that the overall organization of these structures is highly similar despite their different

G-protein partners. An LPC lipid is present at the intracellular interface of TM helices II, III and IV,

as seen in ADGRF1 structures bound to miniGα_s or miniGα_i[13]. The acyl chain is similarly positioned

in our structure, whereas the headgroup has the phosphate embedded deeper in the detergent

micelle (Fig. 2b, Extended Data Fig. 5).

Structural differences between the ADGRF1 miniG$\alpha_{s/q}$ and ADGRF1 miniGα_s complexes

MiniG$\alpha_{s/q}$ was engineered as a chimera with six key Gα_q residues on the C-terminal α5 helix

that distinguish Gα_q coupling over Gα_s, even though the bulk of the chimera is derived from

miniGα$_s$[24]. We also observe a preference for miniGα$_{s/q}$ recruitment to the CTF of ADGRF1,

consistent with the previously observed preference for Gα$_q$ in GTPγS activity assays[8], arguing that

the preference for Gα$_q$ is driven largely by the differences in its C-terminal α5 helix. At the ADGRF1

G-protein interface we observe a canonical pose of the C-terminal α5 helix at the open cavity at

the intracellular opening of TA activated ADGRF1 (Fig. 2b and Fig. 3a). The N387 amide side chain

nitrogen of miniGα$_{s/q}$ approaches within hydrogen bonding distance of the backbone carbonyl of

ADGRF1 R685$^{3.57}$, and the side chain nitrogen of N392 of miniGα$_{s/q}$ approaches within hydrogen

bonding distance of the receptor side chain carboxylate of D842$^{8.47}$. The analogous residues of

miniGα$_s$ are H387 and E392, which do not form the same hydrogen bonding arrangement in the

Gα$_s$ complex. In the Gα$_s$ complex, H387 does not make any polar interactions, and instead E392

approaches within hydrogen bonding distance of the backbone amine of S843$^{8.48}$ (Extended Data

Fig. 6a). Given the close proximity of E392 to D842$^{8.47}$, It is likely that charge repulsion destabilises

the miniGα$_s$-bound state by comparison to the Gα$_q$-bound state. These interactions are associated

with a shift in the positioning of the segment linking TM helix VII to helix VIII relative to the Gα

subunit, with the Cα of D392 in the α5 helix of miniGα$_s$ situated 1.9 Å closer to the Cα of ADGRF1

S843$^{8.48}$ (Fig. 3c). The arrangement seen in the miniGα$_s$ complex is also present in the miniGα$_i$

structure. Superposition of helices I and VII of miniGα$_{s/q}$ on miniGα$_s$ shows a 3.2 Å displacement

of the L384 Cα miniGα$_{s/q}$ relative to that of N384 of miniGα$_s$ (Extended Data Fig. 6b).

At the orthosteric site, we also observe a 1.2 Å displacement of L593$^{1.47}$ towards F569$^{P3'}$ in

the ADGRF1 complex with miniGα$_{s/q}$, compared with the miniGα$_s$ and miniGα$_i$ complexes (Fig. 3d).

This inward movement leads to a rearrangement of the TM I alpha helix, allowing the conserved

S596$^{1.50}$ hydroxyl to approach within hydrogen bond distance of the backbone carbonyl of A828$^{7.48}$

at the kink in the TM VII helix. This interaction of S569$^{1.50}$ with TM helix VII is not observed in either

the miniGα$_s$ or miniGα$_i$ structures.

Distinguishing preferential Gα$_q$ over Gα$_s$ tethered agonism at the orthosteric site

Because ADGRF1 shows preference for Gα$_q$ over other Gα partners[8], we investigated

whether Gα$_q$ and Gα$_s$ signalling exhibited differential sensitivity to alanine mutants of the receptor

stalk residues (Fig. 4a), using NFAT and CRE reporter gene assays for Gα$_q$ and Gα$_s$ coupling,

respectively (Fig. 4b). Positions 574-581, corresponding to P8'-P15', were largely unaffected by

mutation to alanine in NFAT and CRE assays, suggesting this region is not directly important for TA

dependent agonism, and therefore functioning mainly as a linker connecting the TA to TM1 in

ADGRF1.

Alanine substitutions at $F569^{P3'}$, $L572^{P6'}$ and $M573^{P7'}$ of the HIM affected TA-dependent

signalling. The $L572^{P6'}A$ and $M573^{P7'}A$ mutations both reduced the response of the CRE reporter,

whereas these substitutions either did not affect ($L572^{P6'}A$) or modestly increased ($M573^{P7'}A$) the

response of the NFAT reporter, arguing that $G\alpha_s$, but not $G\alpha_q$, selectively requires bulky

hydrophobic residues at positions P6' and P7' for TA-dependent signalling in ADGRF1.

Strikingly, we found that the lack of signalling activity for $F569^{P3'}A$ in both NFAT and CRE is

accompanied by greatly decreased receptor abundance, as judged by the lack of a band on

Western blot when probing for the N-terminal 3X-FLAG tag with an anti-Flag antibody (Fig. 4c). An

antibody raised to a C-terminal portion (residues 831-880) of ADGRF1 showed a similar reduction

in Western blot signal for the $F569^{P3'}A$ mutant (Extended Data Fig. 7), confirming that the loss of

anti-FLAG reactivity was not due to proteolysis of the N-terminal tag. We propose that failure of

the mutated TA to engage the 7TM leads to the exposure of the TA hydrophobic sequence,

creating a neoepitope that stimulates heat-shock protein binding in the secretory pathway and

subsequent degradation (*i.e.*, the inactivity of the TA is the cause of the reduced protein amount,

and not a consequence of it).

We explored the tolerance of $F569^{P3'}$ more extensively by testing a larger range of amino

acid substitutions, including a conservative leucine substitution, a methionine substitution, and a

suite of charged amino acids (Fig. 4d, e). Among these substitutions, only the $F569^{P3'}L$ mutation

produced a detectable signal (Fig. 4d), and only this substitution resulted in similar amounts of

expressed protein as the wild-type CTF as judged by Western blot (Fig. 4e).

Because of the variation in protein abundance for most of the $F569^{P3'}$ mutants, we also

assessed the role of the P3', P6', and P7' hydrophobic buried residues in activation using *trans*

addition of TA peptide sequences. For these assays, we used a form of the 7TM domain lacking

the TA, which we refer to as stalk-less, or Δstalk. We incubated cells expressing the Δstalk form of

ADGRF1 with a 12mer TA mimicking peptide (P12) at 100 μM or 10 μM, and with analogous

peptides containing alanine substitutions at P3', P6' or P7', and measured signalling activity using

the NFAT reporter. We found that the Δstalk construct signaled robustly through NFAT with the

P12 peptide at 100 μM, but not 10 μM (Fig. 4f, g). Together, our data show that a large bulky

hydrophobic residue is required at P3' for ADGRF1 tethered agonism, that $G\alpha_q$ signalling in TA

mode is more tolerant of substitutions at the positions at P6' or P7' in the HIM motif than $G\alpha_s$

signalling, and that signal activation by peptides added in *trans* is intolerant of alanine substitutions

at any of the HIM positions (P3', P6', or P7') even for $G\alpha_q$ signalling.

To further investigate tethered agonism of ADGRF1 through $G\alpha_s$ and $G\alpha_q$, we performed

extensive alanine mutagenesis of the TA binding site in the 7TM domain and measured the

signalling response of the NFAT and CRE reporters. The signalling activity of the mutated proteins

clustered into three distinct categories: mutations that did not affect the CRE or NFAT response,

those that were detrimental to both CRE and NFAT, and those that were detrimental to CRE only

(Fig. 5a, Extended Data Fig. 8).

Residues where alanine substitutions reduced signalling in both the NFAT and CRE

reporters include conserved residues W734$^{45.51}$ and W737 of ECL2, and several residues that

surround F569$^{P3'}$ in the 7TM core, including conserved 'switch' residue W804$^{6.53}$, neighboring

residues F823$^{7.41}$ and N827$^{7.45}$, and T589A$^{1.43}$, L593A$^{1.47}$, A828L$^{7.46}$ and D638A$^{2.61}$ (Fig. 5b-d). These

sets of mutations outline the importance of ECL2, the binding site arrangement around the

'switch' residue, and the conserved F569$^{P3'}$ position for general ADGRF1 tethered agonism. We

also found that positions V812$^{6.61}$ and S814$^{6.63}$ lowered signalling in both NFAT and CRE assays,

but this was likely due to a profound decrease in protein expression (Extended Data Fig. 8)

Strikingly, mutations that decreased CRE signalling but not NFAT signalling were in two

separate clusters that interact with positions other than F569$^{P3'}$ of the TA sequence. The first

cluster contains residues within van der Waals contact of L572$^{P6'}$ and M573$^{P7'}$ at positions Y668$^{3.40}$,

F672$^{3.44}$ and F747$^{5.39}$ on TM helices III and V, with the F672$^{3.44}$A substitution actually enhancing the

NFAT response (Fig 5b-d). The second cluster consists of T810$^{7.28}$ and H820$^{7.38}$; T810$^{7.28}$ interacts

with M573$^{P7'}$ and S574$^{P8'}$, and H820$^{7.38}$ approaches within H-bonding distance of S570^{P4}. These

results show that CRE signalling is more sensitive that NFAT signalling upon disruption of the

pocket proximal to P6' and P7', as well as upon disruption of interactions with S570$^{P4'}$ and S574$^{P8'}$.

Discussion

For a typical G-protein in which the agonist is a soluble ligand, G-protein coupling in a

biological system is guided by intrinsic G-protein binding preferences, agonist selectivity for G-

protein, and the local reservoir of G-proteins in that cellular environment. aGPCRs are unique in

that the 7TM bundle and its TA are encoded in the same polypeptide chain. As a result, aGPCR G-

protein signalling is simplified to a combination of two factors – the G-protein preference of the

TA-engaged CTF and the G-protein context of the cellular environment.

Here, we combine cryo-EM structural studies with mutational analysis and cell-based

signalling assays to elucidate structure-function relationships in ADGRF1 and deduce its G-protein

coupling profile preferences. Our work establishes that ADGRF1 prefers coupling to Gα_q, as

reported previously[8], provides a structure-based rationale for this coupling preference, and, most

strikingly, identifies a cluster of residues in the TA and in the binding pocket that selectively

suppress Gα_s coupling upon mutation while retaining Gα_q coupling (Fig. 6).

While the structure of the ADGRF1-miniG$\alpha_{s/q}$ complex shows that the overall architecture

of the complex and the pose of the TA in the orthosteric site are similar to recently observed

structures of ADGRF1 when complexed to miniGα_s and miniGα_i[13], there are subtle differences in

the complexes that explain why Gα_q coupling is more robust than Gα_s coupling. We also establish

that ADGRF1 can signal through Gα_{12} or Gα_{13}, which to our knowledge makes ADGRF1 the first

class B GPCR to show the functional capacity to signal through all major G-protein classes[28].

Among the structural differences, tighter packing of L593[1.47] with the highly conserved

phenylalanine (P3') of the HIM appears to be critical for Gα_q preference, because the positional

shift of L593[1.47] propagates its effects down helix I, introducing a hydrogen bond between S596[1.50]

and A828[7.48] to stabilize the kink in TM helix VII. Our functional data suggests that these

interactions contribute towards an increased stability of the ADGRF1-Gα_q complex relative to the

Gα_s and Gα_i complexes. Both the TA and the binding pocket of the 7TM domain show increased

mutational lability for Gα_s signalling at several residues, whereas Gα_q signalling remains

unaffected by those same mutations, showing that Gα_q signalling is more tolerant to mutations

than Gα_s. Our data also establish the critical importance of the F569[P3'] residue for ADGRF1 G-

protein signalling, because a bulky hydrophobic residue at the P3' position is essential for

expression and TA-dependent activation of the ADGRF1-CTF and mutations of residues that
surround the F569$^{P3'}$ pocket are detrimental in both NFAT and CRE assays.

Unlike TA signaling of ADGRF1, which is tolerant of substitutions at positions other than
F569$^{P3'}$, we found that *trans* addition of 12mer peptides to a stalkless form of ADGRF1 mimics the
TA only when the wild-type sequence is used at the relatively high concentration of 100 µM.
Mutations at the P6' and P7' positions of the peptides, which are tolerated in TA signaling assays,
failed to activate the stalkless ADGRF1 protein. This observation indicates that the development
of peptidomimetic agonists for ADGRF1 will require extensive optimization if high potency is to be
achieved in a soluble ligand.

Finally, our data elucidating a molecular basis for the preference of ADGRF1 for coupling
to Gα_q over Gα_s identifies key sites in the 7TM that influence G protein coupling preferences.
These findings should also inform the study of G protein coupling preferences and signaling biology
for other aGPCRs.

Methods

Cloning and construct design

MBP-CTF and signalling assay plasmids

DNA sequences encoding full length (20-910) and CTF (567-910) forms of ADGRF1 (Uniprot
Q5T601) were subcloned into the pFuse-hIgG1-Fc2 vector behind an N-terminal IL2-signal
sequence followed by a HA epitope tag, maltose binding protein and TEV protease site[29]. The CTF
sequence was also subcloned to create the M-CTF protein with the N-terminal tags and mannose-
binding protein omitted. Firefly reporter plasmids (CRE, NFAT, SRE and SRF-RE) were assembled
as described[29]. The pRL-TK plasmid (Promega) was used to express *Renilla* luciferase.

Receptor and G-protein plasmids

For recruitment assays, sequences encoding full-length (20-910) and CTF (567-910) ADGRF1 were
subcloned into pcDNA3.1(+) with an N-terminal haemagglutinin signal sequence followed by a
3XFLAG purification tag and a TEV protease cleavage site. For the CTF construct the TEV protease
site P1' residue encoded a threonine so that the correct residue would occupy T567 in the CTF

(e.g., ENLYFQ/TSFSILM…). The C-terminus has a "GGGGSGGGGSSG" linker followed by a SmBiT tag

"VTGYRLFEEIL"[30]. For proteins probed by Western blot the C-terminal SmBiT tag was replaced with

a HA epitope tag. Mutagenesis was performed with a Q5 site directed mutagenesis kit (New

England Biolabs) or using a modified QuikChange™ method[31].

MiniG protein sequences (miniGα$_s$, miniGα$_o$, miniGα$_{s/q}$ and miniGα$_{12}$) with capacity to

complex with Gβ$_1$γ$_2$ (e.g. 399 variants[24]) were subcloned into pcDNA3.1(+) with an N-terminal

LgBiT fusion followed by a "GGGGSGGGGSSGEF" linker that includes a translated EcoRI site. For

structure determination and mutagenesis studies of ADGRF1, the constructs described above for

recruitment assays were used.

Protein expression, purification and complexation

ADGRF1 and miniGα$_{s/q}$ were co-expressed in Expi293F cells. Cells were seeded at a density of 3.3

x 10^6 cells/mL in 900 mL of Expi293 expression media. To transfect cells, equal amounts of receptor

(250 ug) and miniG (250 ug) DNA were resuspended in 50 mL opti-MEM, FectoPro (Polyplus)

transfection reagent (0.5 mL) was resuspended in 50 mL opti-MEM, the DNA and FectoPro

solutions were mixed to give a final 1:1 DNA/FectoPro ratio, and the mixture incubated at room

temperature for 10 minutes before adding to cells. Approximately 24 hours later, filter-sterilised

Valproic acid sodium salt (Sigma-Aldrich) was added to 5 mM, along with 10 mL of 45 % D-(+)-

Glucose solution (Sigma-Aldrich). Cells were cultured for a further 24 hours before they were

harvested by centrifugation at 4000 g for 15 minutes. The pellet was flash frozen and then stored

at -80 °C until purification.

Cells were thawed and resuspended in 250 mL of ice-cold hypotonic lysis buffer (10 mM

HEPES, pH 7.5, containing Roche protease inhibitor tablets and 0.1 mU/mL apyrase to prevent

separation of the GPCR-miniG complex). The pellet was stirred on ice for 30 minutes and lysed by

dounce homogenization. The broken cells were pelleted by centrifugation at 50,000 g and

resuspended in 225 mL solubilisation buffer (10 mM HEPES, pH 7.5 containing 1 mM MgCl$_2$, 2 mM

CaCl$_2$, and 100 mM NaCl) with 0.1 mU/mL apyrase and 1:100,000 Benzonase (v/v). The

resuspended material was dounce homogenized, and DDM-CHS (10:1 pre-mix, Anatrace) was then

added to a final concentration of 1 % (w/v). This solution was stirred for 75 min minutes at 4 °C,

clarified by centrifugation at 50,000 g for 1h, and passed through a glass microfibre filter. The

filtrate was then loaded onto M2 anti-FLAG antibody affinity resin by gravity flow. The resin was

washed with 30 column volumes of wash buffer (10 mM HEPES, pH 7.5 containing 1 mM $MgCl_2$, 2

mM $CaCl_2$, 100 mM NaCl, and 0.1 % (w/v) LMNG-CHS), and the protein was eluted by the

application of wash buffer containing 100 µg/mL of 3XFLAG peptide. The eluted protein was

concentrated using a 100 kDa MWCO centrifugal concentrator (Amicon).

Nb35 and human $G\beta_1\gamma_2$ were purified according to previously published protocols[32].

ADGRF1-miniG$\alpha_{s/q}$, Nb35 and $G\beta_1\gamma_2$ were mixed in a 1:1.15:1.15 ratio and incubated overnight at

4 °C with rotation. The protein mixture was centrifuged at 21,000 g for 20 minutes at 4 °C, and

then spun through a Durapore® PVDF 0.1 µm column (Millipore). The protein mixture was injected

onto a Superdex S200 10/300 GL column equilibrated in SEC buffer (10 mM HEPES pH 7.5

containing 1 mM $MgCl_2$, 2 mM $CaCl_2$, 100 mM NaCl, and 0.005 % (w/v) LMNG-CHS). The peak

fraction containing the complex was collected and concentrated using a 100 kDa MWCO

centrifugal concentrator to approximately 8 mg/mL.

Cryo-EM sample preparation and image acquisition

The complex was vitrified on QUANTIFOIL® holey carbon grids (400-mesh, copper, R1.2/1.3,

Electron Microscopy Sciences) using a FEI Vitrobot Mark IV (FEI, Hillsboro). Grids were glow-

discharged, and 3 µL of sample loaded to the grid in a chamber at 22 °C and 100 % humidity.

Samples were applied at a force of 15 and blotted for 5-7 seconds before plunge-freezing in liquid

ethane.

Data were collected using a FEI Titan Krios at 300 kV with a Gatan Quantum Image Filter

with K3 Summit direct electron detection camera in counting mode with a total exposure dose of

approximately 54 e$^-$/Å^{-2}. 50 frames per movie were collected at magnification of 105,000x,

corresponding to 0.825 Å per pixel. Micrographs were collected at defocus values ranging from -

0.5 to -1.5 um. The processing scheme is summarised in Extended Data Figure 4, in brief, motion

corrected and dose-weighted using the RELION motion correction implementation[33] and contrast

transfer function parameters estimated by CTFFIND4[34]. Low quality micrographs were removed

using micassess[35] prior to particle picking in topaz [36]. ~4.9M particles were initially identified,

leading to ~2.9M particles after 2D classification in RELION [37]. Two independent 3D classifications

in RELION were then performed with different initial references, with the best classes from each

subsequently combined and duplicates removed to leave ~1.2M particles corresponding to

ADGRF1. Following further 3D classification in RELION ~660k particles underwent particle polishing

in RELION [33] and further 2D classification followed by masked 3D classification with no alignment

using a T value of 20 in RELION. 73,903 particles with strong transmembrane density were

identified which resulted in a 3.6Å reconstruction from CryoSPARC non-uniform refinement with

contrast transfer function refinement [38]. Subsequent local non-uniform refinement with a mask

excluding the micelle improved the reconstruction to 3.44Å.

Atomic modelling and model refinement

Model building was carried out in Coot using overlaid non-uniform refinement and local filtered

maps generated in CryoSPARC[39], as well as deepEMhancer[40] maps. The ADGRF1-miniGα_s

coordinate file (PDB: 7WU3) was used as an initial model. The miniGα_s sequence was changed in

Coot to generate miniG$\alpha_{s/q}$[24]. The coordinates were then manually rebuilt using Coot, and refined

using ISOLDE[41], Phenix Real-Space Refine[42] and Servalcat[43]. The final models were evaluated using

MolProbity. Statistics of the map reconstruction and model refinement are presented in

Supplementary Table 1. Structural biology applications used in this project (except CryoSPARC)

were compiled and configured by SBGrid[44].

Cell Culture

HEK293T and HEKΔ6 cells were cultured in DMEM supplemented with 10% FBS and 1% penicillin-

streptomycin at 37°C in a 5% CO_2 humidified incubator.

Dual luciferase reporter assays

HEK293T cells were seeded in DMEM in clear-bottom white 96-well plates pre-coated with poly-

D-lysine the day before transfection. At roughly 70 % cell confluency, the medium in each well was

replaced with fresh DMEM. Cells were transfected using Lipofectamine™ 2,000 (Invitrogen) at a

ratio of 3 μL per 1 μg DNA in Opti-MEM using the manufacturer's instructions. Each well was co-

transfected with 50 ng firefly reporter plasmid, 1 ng pRL-TK plasmid, and the indicated amounts

of receptor DNA. Cells were cultured in the transfection mixture for 6 hours before the media was

replaced with complete media. 24 hours after transfection, the media was replaced with fresh

DMEM, and the cells were incubated for another 8 hours prior to assay readout. For SRF-RE and

SRE read-outs the media was replaced with DMEM lacking FBS. Before measuring luminescence,

cells were washed once with PBS, 20 μL of 1X Passive Lysis Buffer was added to each well, and the

lysates were incubated with shaking for 15 minutes. Cell lysates were serially pipetted, and the

assay plate was centrifuged at 400 g for 1 minute. Dual-Glo luciferase measurements (Promega)

were read on a GloMax luminometer. The ratio of Firefly: Renilla signal was calculated for each

well and normalized to the mean of the EV-transfected controls to give Relative Luminescence

Units (RLU). Each assay was performed in technical triplicate wells and then repeated in biological

triplicate on different days.

For peptide experiments, transfection was performed as above. Peptides were dissolved

in DMSO to give 10 mM stocks and diluted to 10x concentration in Opti-MEM immediately before

use. The final DMSO concentration was 1 % (v/v) in all wells. Peptides were added *in trans* at the

initial media change 6 hours after transfection, and luminescence readings were acquired 24 hours

after transfection. Each assay was performed in technical triplicate wells and then repeated in

biological duplicate on different days.

For Gα rescue assays, HEKΔ6 cells were seeded as above the day before transfection.

Varying amounts of Gα subunit plasmid (0.3 ng of $G\alpha_{12}$ and $G\alpha_{13}$; 20 ng of $G\alpha_s$, $G\alpha_q$, $G\alpha_{11}$, and

$G\alpha_{16}$) were co-transfected with 30 ng firefly reporter plasmid and 0.6 ng pRL-TK plasmid (SRF-RE),

and total DNA load per well was balanced with pcDNA3.1. Media was changed 6 hours after

transfection mixes were added to the wells as in the standard assay format. At 24 hours after

transfection, the media was replaced with DMEM without FBS, and the cells were incubated for

another 8 hours prior to assay readout. Each assay was performed in technical triplicate wells and

then repeated in biological duplicate on different days.

G-protein NanoBiT recruitment assays

HEK293T cells were seeded in DMEM in clear-bottom white 96-well plates pre-coated with poly-D-lysine the day before transfection. Cells were transfected using GeneJuice™ (Sigma-Aldrich) at a ratio of 3 µL per 1 µg DNA in Opti-MEM using manufacturer's instructions. Equal amounts of receptor and miniG DNA (50 ng) were used per well. Cells were incubated for 48 hours and washed with PBS prior to assay readout. Cells were resuspended in opti-MEM and the Nano-Glo® Luciferase Assay was read out on a GloMax luminometer. Each assay was performed in biological triplicate.

Author Contributions:

D.T.D.J. and S.C.B. conceived of the research. D.T.D.J., M.M.B. and A.N.D. designed and cloned constructs. M.M.B. performed mutagenesis of TA and binding site residues. M.M.B., A.N.D. and D.T.D.J. performed signaling assays. D.T.D.J. performed NanoBiT recruitment assays. D.T.D.J. expressed, purified and formed the ADGRF1-MiniG$\alpha_{s/q}$-G$\beta_1\gamma_2$-Nb35 complex. D.T.D.J. and C.H.L. vitrified ADGRF1 complex on grids. S.D.R. and D.T.D.J. processed cryo-EM data, and S.D.R. produced the final map. D.T.D.J. built the ADGRF1 complex model. D.T.D.J. and S.C.B. wrote the manuscript.

Acknowledgements:

We thank members of the Blacklow and Kruse laboratories for helpful discussions. Cryo-EM data were collected at the Harvard Cryo-EM Centre for Structural Biology at Harvard Medical School. We thank Sarah Sterling and Megan Meyer for helpful comments and advice during grid screening, and Megan Meyer for data collection.

Funding:

This work was supported by R35 CA220340 (SCB), and a grant from the Warren Alpert Foundation (SCB). A.N.D. was also supported in part by a Fujifilm fellowship.

Competing Interests:

S.C.B. receives funding for an unrelated project from Novartis, is on the scientific advisory board

for Erasca, Inc., is an advisor to MPM Capital, and is a consultant for IFM, Scorpion Therapeutics

and Ayala Pharmaceuticals for unrelated projects.

References

1.   Bjarnadóttir, T. K. *et al.* The human and mouse repertoire of the adhesion family of G-
protein-coupled receptors. *Genomics* **84**, 23–33 (2004).

2.   Bondarev, A. D. *et al.* Opportunities and challenges for drug discovery in modulating
Adhesion G protein-coupled receptor (GPCR) functions.
https://doi.org/10.1080/17460441.2020.1791075 1291–1307 (2020).
doi:10.1080/17460441.2020.1791075

3.   Weston, M. D., Luijendijk, M. W. J., Humphrey, K. D., Möller, C. & Kimberling, W. J.
Mutations in the VLGR1 gene implicate G-protein signalling in the pathogenesis of Usher
syndrome type II. *Am. J. Hum. Genet.* **74**, 357–366 (2004).

4.   Boyden, S. E. *et al.* Vibratory Urticaria Associated with a Missense Variant in ADGRE2 . *N.*
Engl. J. Med. **374**, 656–663 (2016).

5.   Krasnoperov, V. G. *et al.* α-Latrotoxin Stimulates Exocytosis by the Interaction with a
Neuronal G-Protein-Coupled Receptor. *Neuron* **18**, 925–937 (1997).

6.   Araç, D. *et al.* A novel evolutionarily conserved domain of cell-adhesion GPCRs mediates
autoproteolysis. *EMBO J.* **31**, 1364–1378 (2012).

7.   Liebscher, I., Schö, J., Monk, K. R. & Schö Neberg Correspondence, T. A Tethered Agonist
within the Ectodomain Activates the Adhesion G Protein-Coupled Receptors GPR126 and
GPR133. (2014). doi:10.1016/j.celrep.2014.11.036

8.   Stoveken, H. M., Hajduczok, A. G., Xu, L. & Tall, G. G. Adhesion G protein-coupled
receptors are activated by exposure of a cryptic tethered agonist. *Proc. Natl. Acad. Sci. U.*
S. A. **112**, 6194–6199 (2015).

9.   Ping, Y. Q. *et al.* Structures of the glucocorticoid-bound adhesion receptor GPR97–Go
complex. *Nat. 2020 5897843* **589**, 620–626 (2021).

10.   Xiao, P. *et al.* Tethered peptide activation mechanism of the adhesion GPCRs ADGRG2 and
ADGRG4. *Nat. 2022 6047907* **604**, 771–778 (2022).

11.   Ping, Y.-Q. *et al.* Structural basis for the tethered peptide activation of adhesion GPCRs.
Nat. 2022 6047907 **604**, 763–770 (2022).

12.   Barros-Álvarez, X. *et al.* The tethered peptide activation mechanism of adhesion GPCRs.

Nat. 2022 6047907 **604**, 757–762 (2022).

13.   Qu, X. *et al.* Structural basis of tethered agonism of the adhesion GPCRs ADGRD1 and

ADGRF1. *Nat. 2022* **604**, 1–8 (2022).

14.   Fredriksson, R., Lagerström, M. C., Höglund, P. J. & Schiöth, H. B. Novel human G protein-

coupled receptors with long N-terminals containing GPS domains and Ser/Thr-rich regions.

FEBS Lett. **531**, 407–414 (2002).

15.   Lum, A. M. *et al.* Orphan receptor GPR110, an oncogene overexpressed in lung and

prostate cancer. *BMC Cancer* **10**, 40 (2010).

16.   Lee, J. W. *et al.* Orphan GPR110 (ADGRF1) targeted by N-docosahexaenoylethanolamine in

development of neurons and cognitive function. *Nat. Commun.* **7**, (2016).

17.   Kwon, H. *et al.* Ligand-Induced GPR110 Activation Facilitates Axon Growth after Injury. *Int.*

J. Mol. Sci. **22**, (2021).

18.   Stoveken, H. M., Larsen, S. D., Smrcka, A. V. & Tall, G. G. Gedunin- and khivorin-derivatives

are small-molecule partial agonists for adhesion G protein-coupled receptors

GPR56/ADGRG1 and GPR114/ADGRG5. *Mol. Pharmacol.* **93**, 477–488 (2018).

19.   Demberg, L. M. *et al.* Activation of Adhesion G Protein-coupled Receptors: AGONIST

SPECIFICITY OF STACHEL SEQUENCE-DERIVED PEPTIDES*. *J. Biol. Chem.* **292**, 4383 (2017).

20.   Dates, A. N. *et al.* A Fusion Protein Platform for Analyzing Tethered Agonism in the

Adhesion Family of G Protein-Coupled Receptors. *bioRxiv* 2022.07.14.500097 (2022).

doi:10.1101/2022.07.14.500097

21.   Demberg, L. M. *et al.* Activation of adhesion G protein-coupled receptors: Agonist

specificity of Stachel sequence-derived peptides. *J. Biol. Chem.* **292**, 4383–4394 (2017).

22.   Grundmann, M. *et al.* Lack of beta-arrestin signalling in the absence of active G proteins.

Nat. Commun. 2018 91 **9**, 1–16 (2018).

23.   Hauser, A. S. *et al.* Common coupling map advances GPCR-G protein selectivity. *Elife* **11**,

(2022).

24.   Nehmea, R. *et al.* Mini-G proteins: Novel tools for studying GPCRs in their active

conformation. *PLoS One* **12**, e0175642 (2017).

25.   Carpenter, B. & Tate, C. G. Engineering a minimal G protein to facilitate crystallisation of G

protein-coupled receptors in their active conformation. *Protein Eng. Des. Sel.* **29**, 583

(2016).

26. Hall, M. P. *et al.* Engineered luciferase reporter from a deep sea shrimp utilizing a novel

imidazopyrazinone substrate. *ACS Chem. Biol.* **7**, 1848–1857 (2012).

27. Wan, Q. *et al.* Mini G protein probes for active G protein–coupled receptors (GPCRs) in

live cells. *J. Biol. Chem.* **293**, 7466 (2018).

28. Hauser, A. S. *et al.* Common coupling map advances GPCR-G protein selectivity. *Elife* **11**,

(2022).

29. Dates, A. N. *et al.* A Fusion Protein Platform for Analyzing Tethered Agonism in the

Adhesion Family of G 5 Protein-Coupled Receptors 6 7 8 9.

doi:10.1101/2022.07.14.500097

30. Dixon, A. S. *et al.* NanoLuc Complementation Reporter Optimized for Accurate

Measurement of Protein Interactions in Cells. *ACS Chem. Biol.* **11**, 400–408 (2016).

31. Liu, H. & Naismith, J. H. An efficient one-step site-directed deletion, insertion, single and

multiple-site plasmid mutagenesis protocol. *BMC Biotechnol.* **8**, 91 (2008).

32. Rasmussen, S. G. F. *et al.* Crystal structure of the β2 adrenergic receptor–Gs protein

complex. *Nat. 2011 4777366* **477**, 549–555 (2011).

33. Zivanov, J., Nakane, T. & Scheres, S. H. W. A Bayesian approach to beam-induced motion

correction in cryo-EM single-particle analysis. *urn:issn:2052-2525* **6**, 5–17 (2019).

34. Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from electron

micrographs. *J. Struct. Biol.* **192**, 216 (2015).

35. Li, Y., Cash, J. N., Tesmer, J. J. G. & Cianfrocco, M. A. High-Throughput Cryo-EM Enabled by

User-Free Preprocessing Routines. *Structure* **28**, 858-869.e3 (2020).

36. Bepler, T. *et al.* Positive-unlabeled convolutional neural networks for particle picking in

cryo-electron micrographs. *Nat. Methods 2019 1611* **16**, 1153–1160 (2019).

37. Scheres, S. H. W. RELION: Implementation of a Bayesian approach to cryo-EM structure

determination. *J. Struct. Biol.* **180**, 519 (2012).

38. Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization

improves single-particle cryo-EM reconstruction. *Nat. Methods 2020 1712* **17**, 1214–1221

(2020).

39.   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid

507   unsupervised cryo-EM structure determination. *Nat. Methods 2017 143* **14**, 290–296

508   (2017).

509   40.   Sanchez-Garcia, R. *et al.* DeepEMhancer: a deep learning solution for cryo-EM volume

510   post-processing. *Commun. Biol. 2021 41* **4**, 1–8 (2021).

511   41.   Croll, T. I. ISOLDE: A physically realistic environment for model building into low-resolution

512   electron-density maps. *Acta Crystallogr. Sect. D Struct. Biol.* **74**, 519–530 (2018).

513   42.   Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography.

514   *urn:issn:2059-7983* **74**, 531–544 (2018).

515   43.   Yamashita, K., Palmer, C. M., Burnley, T. & Murshudov, G. N. Cryo-EM single-particle

516   structure refinement and map calculation using Servalcat. *Acta Crystallogr. Sect. D, Struct.*

517   *Biol.* **77**, 1282–1291 (2021).

44.   Morin, A. *et al.* Collaboration gets the most out of software. *Elife* **2**, (2013).

Figure 1 – ADGRF1 signals through all G-protein classes. (a) Transcription factor reporter assay
responses for ADGRF1 FL, M-CTF or MBP-CTF relative to empty vector for CRE (top left), NFAT
(top right), SRF-RE (bottom left) and SRE (bottom right). Cells were transfected with 30 ng
DNA/well for SRE and SRF-RE, and 10 ng DNA/well for CRE and NFAT. Data are normalized to
empty vector as relative luminescence units (RLU). Data are presented as mean ± s.d. of three
biological replicates. One-way analysis of variance (ANOVA) was used with Dunnett's multiple-
comparison post-hoc test to compare the difference between samples to each reporter's
empty vector control (*P≤0.05, **P≤0.01, and ****P≤0.0001). (b) Gα transient transfection
complementation assay. Activity of the SRF-RE reporter was measured in HEKΔ6 cells
transfected with a combination of ADGRF1 and different Gα subunits. The Gα subunit or Gα
empty vector control was co-transfected with receptor empty vector, ADGRF1 FL or MBP-CTF.
Values are reported as the ratio of the raw luminescence values of firefly and renilla luciferases.
Data are presented as mean ± s.d. of two biological replicates. Each point represents the mean
value of three technical replicates. In all panels, One-way analysis of variance (ANOVA) was
used with Dunnett's multiple-comparison post-hoc test to compare the difference between
samples to each Gα empty vector control (*P≤0.05, **P≤0.01, ***P≤0.001 and ****P≤0.0001).

Figure 2 – ADGRF1-miniGα$_{s/q}$-Nb35 complex purification and cryo-EM structure. (a) ADGRF1 miniGα$_{s/q}$-Gβ$_1$γ$_2$-Nb35 purification. Top: elution profile
of the complex after size-exclusion chromatography as a final purification step. Bottom: peak fractions (bar above chromatogram) analyzed by
SDS-PAGE. (b) Structure of the ADGRF1-Gα$_{s/q}$-β$_1$γ$_2$-Nb35 complex. Left: cryo-EM map of the ADGRF1 complex. ADGRF1 (light blue), the TA (green),
Gα$_{s/q}$ (orange), Gβ$_1$ (teal), Gγ$_2$ (grey) and Nb35 (pink) are coloured as indicated. The density threshold was set at five standard deviations from the
mean. Right: cartoon rendering of the structure, with residue side chains of the tethered agonist rendered as sticks. The density threshold was set
at three standard deviations from the mean for the TA. The LPC lipid (yellow) molecule is shown with a ball-and-stick model.

Figure 3 – The ADGRF1-miniGα$_{s/q}$ interface and comparison to ADGRF1-miniGα$_s$. (a) Close-up of the ADGRF1 (blue) interface with Gα$_{s/q}$ (orange)
along the Cα5 helix of Gα$_{s/q}$. Hydrogen bonding interactions between the receptor and Gα are indicated with yellow dashed lines. (b) Comparison
of the ADGRF1 (blue) - Gα$_{s/q}$ (orange) complex with the ADGRF1 (pink) - miniGα$_s$ (cyan) complex, aligned by least-squares superposition of the
ADGRF1 subunits. (c) Zoomed in view focusing on D842$^{8.47}$ in the turn connecting helices VII and VIII. The hydrogen bond between N392 and the
carboxylate of D842$^{8.47}$ is shown with a yellow dotted line. (d) Zoomed in view focusing on residues L593$^{1.47}$ and S596$^{1.50}$. Arrows indicate positional
shifts of these residues in the miniGα$_{s/q}$ structure relative to their positions in miniGα$_s$. The hydrogen bond between S596$^{1.50}$ and the backbone
carbonyl of A828$^{7.46}$ is shown with a yellow dotted line.

Figure 4 – Tethered agonism signalling determinants through NFAT and CRE. (a) Pose of the TA in the orthosteric site. TA sidechain atoms and
main-chain atoms making polar contacts are shown as sticks. Side-chains of 7TM residues making polar contacts (yellow dashes) to the TA are also
shown. TA superscript labels refer to the TA position relative to the GPS autoproteolysis site. Receptor residue superscript labels refer to Wootten
numbering. (b) Scanning alanine mutagenesis of the TA. NFAT and CRE reporter data were acquired using ADGRF1 CTF as a reference sequence.
Cells were transfected with 20 ng of receptor DNA/well. Data are normalized to empty vector and reported as relative luminescence units (RLU).
The mean for the CTF is indicated as a horizontal dotted line. *Figure legend continued overleaf.*

(c) α-FLAG Western blot comparing steady-state amounts of wild-type and mutated ADGRF1 CTF proteins tested in panel b. An α-GAPDH blot was
used as a loading control. (d) NFAT signalling data and (e) α-FLAG Western blot of wild-type and P3' mutated forms (L, M, R, K, D and E) of the
ADGRF1 CTF. (f) NFAT reporter gene activity of the ADGRF1 CTFΔstalk protein stimulated by *trans* addition of a 12-residue TA peptide (P12). The
response to the wild-type peptide or to peptides with alanine mutations at the P3', P6' or P7' position are shown at doses of 10 or 100 μM. (g) α-
FLAG Western blot comparing steady-state amounts of wild-type and Δstalk ADGRF1 CTF proteins tested in panel f. An α-GAPDH blot was used as
a loading control. For panel (b), data are presented as mean ± s.d. of three biological replicates. Each point represents the mean value of three
technical replicates. For panels (d and f) data are presented as mean ± s.d. of two biological replicates. Each point represents the mean value of
three technical replicates. One-way analysis of variance (ANOVA) was used with Dunnett's multiple-comparison post-hoc test to compare the
difference between CTF to all other samples individually in b and d, and to DMSO in f (**$P<0.01$;***$P<0.001$;****$P<0.0001$).

Figure 5 – Binding site signalling determinants through NFAT and CRE. (a) NFAT and CRE signalling data for TA binding site mutants in the context of the ADGRF1 CTF protein. Cells were transfected with 20 ng of receptor DNA/well. Data are normalized to empty vector and reported as relative luminescence units (RLU). Data are grouped according to their effects on NFAT and CRE signals relative to the wild-type ADGRF1 CTF protein. Mutations without significant effects on either reporter are grey, mutations that significantly reduce both NFAT and CRE signals are cyan, and mutations that significantly reduce only the CRE signal are magenta. Data are presented as mean ± s.d. of three biological replicates. The mean for the CTF is indicated as a horizontal dotted line. Each point represents the mean value of three technical replicates. One-way analysis of variance (ANOVA) was used with Dunnett's multiple-comparison post-hoc test to compare the difference between CTF to all other samples individually (*P≤0.05, **P≤0.01, ***P≤0.001 and ****P≤0.0001). *Figure legend continued overleaf.*

(b) Top view of the TA in the orthosteric site. TA sidechain atoms and main-chain atoms making
polar contacts (yellow dashes) are shown as sticks. Side-chains of 7TM residues that affect
signalling when mutated to alanine are shown as sticks and are coloured according to the
grouping in (a). (c) View from the P7' end and (d) the P3' end of the TA. Side-chains of 7TM
residues that affect signalling when mutated to alanine are shown as sticks, coloured according
to the grouping in (a), and annotated by residue with Wooten superscripts.

Figure 6 – Mutations of the TA and associated binding site cluster by G-protein class
preference. ADGRF1 coupled to miniG$\alpha_{s/q}$ represented as tubed helices, colored light grey and
orange respectively. Residues that reduced signaling activity are shown, with TA residues
rendered as sticks and with Cα atoms of residues in the TA binding site rendered as spheres.
Residues that reduce reduce both Gα_s and Ga$_q$ signalling are colored cyan, whereas residues
that only reduce Gα_s signaling are magenta.
591

Extended Data Figure 2 – NanoBiT complementation of GPR110-CTF with miniGα proteins. (a) NanoBiT recruitment assays for assessment of
receptor-miniG coupling efficiency. Adherent HEK293T cells were transfected with equal amounts of DNA for receptor and miniG proteins and
luminescence recorded after the addition of substrate. Full-length (FL) and C-terminal fragment (CTF) variants of the receptor were combined with
either functional miniG or non-functional miniG CΔ5 controls. Luminescence resulting from luciferase reconstitution is plotted, with bars grouped
and coloured by miniG class. One-way analysis of variance (ANOVA) was used with Dunnett's multiple-comparison post-hoc test to compare the
difference between FL and CΔ5 to all other samples within each miniG group individually (*P≤0.05, **P≤0.01, ***P≤0.001 and ****P≤0.0001). (b)
Suspension Expi293T cells were transfected with equal amounts of DNA for receptor and miniG proteins in whole cell (left) and solubilised formats
(right). Luminescence resulting from luciferase reconstitution is plotted, with bars coloured by miniG class as in (a). Unpaired two-tailed T-test was
used to compare the difference between FL and CTF for whole cell luminescence (***P≤0.001 and ****P≤0.0001). (c) Coomassie blue staining
after SDS-PAGE of ADGRF1-CTF complexes recovered on magnetic beads, assessing coupling efficiency to miniG proteins.

Extended Data Figure 3 – Cryo-EM data quality. (Top, left) A representative image of ADGRF1 complexes in vitreous ice visualized by cryo-EM on
a Titan Krios microscope equipped with Gatan K3 detector. (Top, right) 2D-class averages showing a range of particle orientations. (Bottom, left)
Local resolution map generated in Cryosparc (GS-FSC = 0.143 cut-off). (Bottom, centre) GS-FSC curves with default Cryosparc masks. (Bottom,
right) Particle orientation distribution map of ADGRF1-miniG$\alpha_{s/q}$ complexes used for structure determination.

7310 micrographs

Micrograph curation

6598 micrographs

Particle picking
4,960,830 particles

2D classification

2,844,289 particles

Independent 3D classification

Recombine + deduplication
1,166,910 particles

659,412 particles

Iterative Particle polishing + 2D classification

376,206 particles

73,903 particles
CTF Refinement
Non-Uniform Refinement (3.6Å)
Local Non-Uniform Refinement (3.44Å)

611   **Extended Data Figure 4 – Cryo-EM processing scheme for ADGRF1 reconstruction.** Selected
612   classes are blue and discarded classes are grey.

Extended data Figure 6 – Comparison of ADGRF1 bound to miniGα_{s/q} and miniGα_s. a) Close-up of the ADGRF1 (pink) interface with Gα_s (cyan)

along the Cα5 helix of Gα_s. Hydrogen bonds between the receptor and Gα are indicated. (b) Left: side-view, and Right: top-slice view of ADGRF1

bound to miniGα_s or miniGα_{s/q}. The receptors are aligned by least-squares superposition of TM helices I and VII.

Extended Data Figure 7 – Amount of ADGRF1 CTF protein expressed by wild-type and alanine scanning mutants. Protein was detected by Western
blotting with an α-ADGRF1 C-terminal antibody.

Extended Data Figure 8 – Amount of ADGRF1 CTF protein expressed by wild-type and alanine scanning mutants of the binding site. Protein was
detected by Western blotting with an α-FLAG antibody, which recognizes an N-terminal FLAG tag. A Western blot with an anti-GAPDH antibody
was used as a loading control. A Western blot with an anti-GAPDH antibody was used as a loading control.

 Table 1 | Cryo-EM data collection, refinement and validation statistics.

ADGRF1	
(EMD-XXX, PDB XXX)	
Data collection and processing	
Voltage (kV)	300
Electron exposure (e⁻/Å⁻²)	~53.5
Defocus range (μm)	0.5 – 1.5
Pixel size (Å)	0.825
Symmetry imposed	C1
Initial particle images (no.)	4,960,830
Final particle images (no.)	73,903
Map Resolution (Å)	3.44
FSC threshold	0.143
Map resolution range (Å)	3.1 - 6.0
Refinement	
Initial model used	7wu3
Model resolution (Å)	3.10
FSC threshold	0.143
Map-sharpening B-factor (Å²)	-111.2
Model composition	
Non-hydrogen atoms	6,671
Protein residues	891
B factors (Å²)	
Protein	74.17
Ligand	97.56
R.M.S.D. deviations	
Bond lengths (Å)	0.003
Bond angles (°)	0.594
Validation	
MolProbity score	1.63
Clashscore	5.21
Poor rotamers (%)	0.00
Ramachandran plot	
Favoured (%)	94.88
Allowed (%)	5.12
Disallowed (%)	0.00