# Regulatory and evolutionary annotation identifies KAN1, MYB44 and PIF4 as regulators of Arabidopsis lincRNAs expressed in root

Li Liu[1,2], Michel Heidecker[3,4], Thomas Depuydt[1,2], Nicolas Manosalva Perez[1,2], Martin Crespi[3,4], Thomas Blein[3,4*], Klaas Vandepoele[1,2,4*]

(1) Ghent University, Department of Plant Biotechnology and Bioinformatics, Technologiepark 71, 9052 Ghent, Belgium

(2) VIB Center for Plant Systems Biology, Technologiepark 71, 9052 Ghent, Belgium

(3) Université Paris-Saclay, CNRS, INRAE, Université Evry, Institute of Plant Sciences Paris-Saclay (IPS2), 91190 Gif-sur-Yvette, France

(4) Université Paris Cité, CNRS, INRAE, Institute of Plant Sciences Paris-Saclay (IPS2), 91190 Gif-sur-Yvette, France

(5) Bioinformatics Institute Ghent, Ghent University, Technologiepark 71, 9052 Ghent, Belgium

* Shared last author

Corresponding author: klaas.vandepoele@psb.vib-ugent.be

Running title: TF regulatory annotation of Arabidopsis root lincRNAs

**ABSTRACT**

Thousands of long intergenic noncoding RNAs (lincRNAs) have been identified in plant genomes. While some lincRNAs have been characterized as important regulators in different biological processes, little is known about the transcriptional regulation for most plant lincRNAs. Through the integration of eight annotation resources, we defined 6,599 high-confidence lincRNA loci in *Arabidopsis thaliana*. For lincRNAs belonging to different evolutionary age categories, we identified major differences in sequence and chromatin features, as well as in the level of conservation and selection. Spatiotemporal gene expression profiles combined with transcription factor (TF) chromatin immunoprecipitation data was used to construct a TF-lincRNA regulatory network containing 2,659 lincRNAs and 43,233 interactions. We experimentally confirmed the role of three TFs, KAN1, MYB44, and PIF4, as key regulators controlling root-specific lincRNA expression, demonstrating the predictive power of our network. Furthermore, we identified 58 lincRNAs regulated by these TFs showing strong root cell-type specific expression and that are linked with GWAS genetic associations related to root system development and growth. The multi-omics approach applied in this study sheds light on the global regulatory complexity of plant lincRNA networks and pinpoints a role of specific TFs in lincRNA regulation in roots.

## Introduction

Genomes are widely transcribed and produce thousands of long non-coding RNAs (lncRNAs), which are an abundant class of transcripts longer than 200 nucleotides with low protein coding capacity (Wu et al., 2017). LncRNAs are generally transcribed by RNA polymerase II (Pol II), are poly-adenylated and may contain introns as coding RNA. LncRNAs modulate gene expression through a wide range of mechanisms, including chromatin structure remodeling, transcription regulation in cis/trans, fine-tuning of miRNA activity, alternative splicing (AS) regulation, and the control of mRNA stability and translation (Sanchita et al., 2020; Bhogireddy et al., 2021; Lucero et al., 2021). One class of lncRNAs are the primary mRNAs containing miRNA precursors or pri-miRNAs. These are rapidly and generally processed into miRNAs (Jones-Rhoades et al., 2006; Shafiq et al., 2016). However, the large majority of lncRNAs are able to act without being further processed such as the lncRNAs controlling the epigenetic regulation the *Flowering Locus C* (*FLC*) gene expression and mediating plant vernalization, i.e., *COOLAIR*, *COLDAIR* and *COLDWRAP* (Liu et al., 2010; Heo and Sung, 2011; Kim and Sung, 2017). A subgroup of lncRNAs derived from intergenic regions is defined as long intergenic non-coding RNAs (lincRNAs) and have been identified in a wide range of eukaryotes including model and non-model plant species (Wu et al., 2017; Chen et al., 2021). In contrast to antisense lncRNAs, whose sequence evolution is constrained by the overlapping coding genes, the transcription and evolution of lincRNAs are independent of the surrounding genes.

While the low expression levels and tissue-specific expression patterns of lincRNAs in plants initially raised concerns (Liu et al., 2012; Bu et al., 2015), increasing experimental evidence supports the functional activity of lincRNAs. In plants, few lincRNAs have been experimentally validated (Chen et al., 2021), showing their involvement in various biological contexts such as in regulating flowering time (Chen et al., 2020) and root growth and development (Roule et al., 2022a), or influencing germination (Kramer et al., 2022). For example, the Arabidopsis *FLINC* lincRNA has been reported to regulate ambient temperature-

mediated flowering time (Severing et al., 2018). Arabidopsis lateral root
development is regulated by the *Alternative Splicing COmpetitor* (*ASCO*) lincRNA,
which modulates AS by interacting with the multiple splicing factors (Bardou et al.,
2014; Rigo et al., 2020), and the *AUXIN-REGULATED PROMOTER LOOP*
(*APOLO*) lincRNA, which influences the local chromatin conformation and the
activity of several Auxin-Responsive genes (Ariel et al., 2020). Plants lacking
*CONSERVED INBRASSICA RAPA1* (*lncCOBRA1*) were found to show a
delayed germination and were overall smaller than wild-type plants (Kramer et al.,
2022). Many lincRNAs are differentially expressed in various stress responses,
including drought (Qi et al., 2013; Shuai et al., 2014; Li et al., 2017; Qin et al.,
2017), cold (Li et al., 2017; Zhao et al., 2018; Shea et al., 2019), salinity (Deng et
al., 2018; Fukuda et al., 2019), and nutrient deficiency (Fukuda et al., 2019),
implying that lincRNAs may be involved in plant stress resilience (Jha et al.,
2020). Interestingly, some of the confirmed functional lincRNAs interact with
transcription factors (TFs) to activate or repress the expression of associated
genes, such as *APOLO* that interacts with WRKY42 to form a regulatory hub that
controls the activity of *RHD6* and promotes the expansion of root hair cell at low
temperatures (Moison et al., 2021; Pacheco et al., 2021).

Recently, a lot of attention has been placed on the evolutionary conservation of
lincRNAs, which is generally associated with functionality (Ulitsky, 2016;
Szczesniak et al., 2021). The conservation of noncoding transcripts can be
considered at the level of the primary sequence, position, splice sites, RNA
structure, and transcriptional level (Ulitsky, 2016; Szczesniak et al., 2021).
However, most lincRNA sequences undergo rapid evolution and are poorly
conserved (Ransohoff et al., 2018). In a study by Wang et al., only 5% of 2,281
rice lincRNAs had sequence similarity to maize lincRNAs. It was also found that
the positional conservation of lincRNAs was much higher than their sequence
conservation (Wang et al., 2015b). Nelson and co-workers reported that 22% of
1180 Arabidopsis lincRNA loci were conserved in 10 Brassicaceae genomes
(Nelson et al., 2016). These conserved lincRNAs exhibited higher expression
levels, stress-responsiveness and their gene body overlapped with conserved

noncoding sequences (CNSs), suggesting a role of their conserved sequence in

a genomic context (Nelson et al., 2016).

While different studies have reported on the identification and expression of

lincRNAs in plants (Wang et al., 2015a; Nelson et al., 2016; Ke et al., 2019; He et

al., 2021), a comprehensive overview of the different genomic features

contributing to the expression, regulation and evolutionary conservation of plant

lincRNAs is missing. How lincRNAs are embedded in transcriptional networks

controlling different biological processes remains largely unknown. Furthermore,

prioritizing lincRNAs for downstream functional analysis is not straightforward

without knowing the regulatory network where they are involved in. Here, we

integrated different Arabidopsis lincRNAs gene annotations and explored various

functional genomics datasets to characterize lincRNA expression in a biologically

relevant context. We leveraged large-scale expression datasets and protein-DNA

interaction data to study the molecular networks controlling lincRNA gene activity.

Combined with evolutionary conservation analysis, we explored the global

transcriptional regulatory properties of different evolutionary age categories and,

through regulatory network analysis, identified specific TFs controlling lincRNA

regulation in roots.

**RESULTS**

**Integration of lincRNA annotations in *Arabidopsis***

A substantial number of lncRNA transcripts in *Arabidopsis* have been identified

and several publicly available resources for the annotation of lncRNAs have been

developed (Jha et al., 2020). In contrast to antisense lncRNAs which generally

regulate the overlapping coding gene, much less is known about the potential

targets of lincRNAs, so we focus our study on these transcripts. Indeed,

lincRNAs transcription and evolution are independent of the surrounding genes,

in contrast to antisense lncRNAs that are constrained by the coding genes they

overlap with. To integrate and unify previously identified lincRNAs, annotations

were collected from different resources, including Araport11 (Cheng et al., 2017),

CANTATAdb (Szczesniak et al., 2016), NONCODEv5 (Fang et al., 2018),
PLNlncRbase (Xuan et al., 2015), key lncRNA research articles (Liu et al., 2012;
Nelson et al., 2017; Zhao et al., 2018), and new predictions based on root related
stranded RNA-seq (Material and Methods). Next, a pipeline was designed to
define a unified set of high-confidence lincRNAs by discarding transcripts with
length below 200 bp, removing transcripts that overlapped with protein-coding
genes (antisense lncRNAs), re-evaluating the coding potential of the transcripts,
and merging the remaining transcripts from various resources (see Materials and
Methods, Supplemental Figure S1A-B). In total, we identified 7,706 lincRNA
transcripts covering 6,599 high-confidence lincRNA loci (see Supplemental Data
Set S1). To explore the overlap of this high-confidence lincRNA gene set with the
individual annotations, we assessed the overlap between the different resources
(Figure 1A). A total of 4,955 (75.1%) lincRNA loci were supported by only one
resource and the remaining 1,644 (24.9%) lincRNA loci were derived from two or
more resources (Figure 1B). Araport11 contained the highest number of shared
loci and Liu et al. (2012) contained the highest number of unique loci (Chen et al.,
2021). Next, the genomic features of the high-confidence lincRNA transcripts
were compared to those of protein-coding transcripts. 6,428 (83%) lincRNA
transcripts and 6,250 (13%) protein-coding transcripts contained single exons,
while 1,278 (17%) lincRNA transcripts and 42,109 (87%) protein-coding
transcripts contained multiple exons. Furthermore, a higher frequency of multi-
exon transcripts was found in lincRNAs supported by two or more resources (974,
36%) than in those supported by a unique resource (304, 6%) (Figure 1C). The
transcript length distribution for lincRNAs showed a U-shape curve with the
majority of transcripts being 200-300bp long (Figure 1D).

163
**Figure 1. Overlap and gene features of Arabidopsis lincRNA annotations. (A)**
Upset plot showing the intersection of lincRNA annotation in the eight resources.
Each row represents a resource, reporting in parenthesis its total number of
lincRNA transcripts before merging. LincRNA annotations unique to a single
resource are represented as a single circle while circles connected by lines
represent the intersection of lincRNA loci shared between various resources. The
bar chart indicates the number of unique lincRNA loci and intersectional lincRNA
loci, displaying only intersections that contain at least ten lincRNA loci. More
complex overlapping patterns are not shown. **(B)** The pie chart shows the

proportion of lincRNA loci supported by one or more resources. **(C)** The

distribution of exon number for all lincRNA transcripts (purple), protein-coding

transcripts (green), transcripts of lincRNAs supported by single resource (purple)

and multiple resources (purple). Single exon and multiple exons are shown in

dark and light colors, respectively. **(D)** The distribution of transcript length for

lincRNAs (purple) and protein-coding genes (green).

**Contrasting patterns of sequence conservation for lincRNAs belonging to**

**different evolutionary age categories**

To assess the evolutionary conservation of Arabidopsis lincRNAs within flowering

plants, DNA sequence similarity searches were performed by comparing our set

of lincRNAs with the genomes of 40 plant species (see Materials and Methods,

Supplemental Table S1). Among the 6,599 lincRNAs, 2,480 lincRNAs were

restricted to Arabidopsis and named Arabidopsis-specific lincRNAs. The other

lincRNAs were classified into four evolutionary age categories according to the

presence of homologs in closely and more distantly related species (Figure 2A).

We found 81 lincRNAs with at least one homolog in eudicots and in monocots

and therefore conserved during 180 million years (MY) of evolution (Beilstein et

al., 2010; Zhang et al., 2020), defined as angiosperm-conserved lincRNAs. Forty-

two lincRNAs were conserved in eudicots with at least one homolog in rosids and

asterids, but without homologs in monocots. Similarly, 44 lincRNAs were

identified only having homologs in rosids species, outside the Brassicaceae

family. As the lincRNAs conserved in eudicots and rosids showed highly similar

conservation patterns for gene body and promoter (P<0.376, Mann–Whitney *U*

test), we combined these genes in one category, called Eudicot/rosid-conserved

lincRNAs (86 genes). 1,671 Brassicaceae_I_II-conserved lincRNAs were present

in the common ancestor of Brassicaceae lineages I and II, without homologs

outside the Brassicaceae. Lastly, 2,281 lincRNAs were identified only having

homologs in the Brassicaceae I lineage (Brassicaceae_I-conserved lincRNAs).

The monotonous decrease in the number of lincRNAs for the older age

categories suggests there is a continuous birth of lincRNA loci at the species
level, with only a small fraction showing deep conservation in other plant families
or orders.
206

207
**Figure 2. Sequence conservation analysis for lincRNAs of different**
**evolutionary age categories. (A)** Simplified species tree reporting the number
of lincRNAs found for the different age categories, which are also indicated by
the grey circle sizes. Numbers in parenthesis report the number of genomes
included per clade to assess sequence similarity and define a lincRNA's age
category. **(B)** Sequence conservation (measured by the average phastCons
score over the gene body) for different lincRNA age categories and gene types
(lincRNAs, pri-miRNAs, and protein-coding genes). PhastCons score ranges
from 0 (not under selection) to 1 (strong negative selection). P-values for pairwise
Mann–Whitney *U* test are shown using the horizontal lines connecting the series.
**(C)** Boxplot showing the average phastCons score for promoter regions (2kb
upstream of transcription start site) of different lincRNA age categories and gene
types (lincRNAs, pri-miRNAs, protein-coding genes). P-values for pairwise
Mann–Whitney *U* test are shown using the horizontal lines connecting the series.
222

Apart from detecting lincRNA homologs, we also evaluated the evolutionary
conservation of lincRNA loci, pri-miRNAs, where globally the 21-24 miRNA
sequence is the only conserved region at the nucleotide level, and protein-coding
genes using phastCons scores (Siepel et al., 2005), which report the probability
for each nucleotide to evolve neutrally or under negative, or purifying, selection
(low or high phastCons score, respectively). In contrast to the age categories
which are based on finding similar homologous sequences in other plant
genomes, phastCons works by fitting a two-state hidden Markov model to a
genome-wide multiple sequence alignment and predicting, based on the pattern
of nucleotide substitutions, conserved elements representing sites under
purifying selection (Siepel et al., 2005). For lincRNA, pri-miRNA, and protein-
coding gene loci, we compared phastCons scores for the gene body and the
promoter region (2kb upstream of transcription start site) (Supplemental Data Set
S2). In general, for gene bodies, lincRNAs are significantly less conserved than
pri-miRNAs and protein-coding genes (Figure 2B, purple, yellow, and green
series). However, we found that Angiosperm- and Eudicot/rosid-conserved
lincRNAs were as conserved as protein-coding genes. Brassicaceae_I_II-
conserved lincRNAs showed even significantly higher conservation levels than
protein-coding genes, indicating high levels of purifying selection. As expected,
Arabidopsis-specific lincRNAs show the lowest phastCons scores.
In contrast to the differences in gene body, the promoter scores are comparable
for lincRNA, pri-miRNA, and protein-coding genes (Figure 2C, purple, yellow and
green series). The level of purifying selection on promoter regions is similar in
Angiosperm- and Eudicot/rosid-conserved lincRNAs as well as in protein-coding
genes. We observed that conservation levels were again significantly higher in
the Brassicaceae_I_II-conserved lincRNA promoters than protein-coding gene
promoters. Taken together, these results indicate that Brassicaceae_I_II-
conserved lincRNAs stand out in the level of purifying selection acting on these
loci, both in the gene body and in their promoter, suggesting that the primary
sequence of the RNA and its promoter regulation is a critical element for lincRNA
function.

**Large-scale transcriptome analysis reveals highly-specific lincRNA expression in roots**

Increasing evidence supports the tissue or cell-type specific role of lincRNA gene functions (Liu et al., 2012; Li et al., 2016; Cheng et al., 2017; Rai et al., 2019). To characterize spatiotemporal Arabidopsis gene expression patterns also considering lincRNAs, we curated, processed, and integrated 791 RNA-seq samples to construct a genome-wide gene expression atlas covering a wide range of tissues, developmental stages, and stress conditions (Supplemental Table S2 and Supplemental Data Set S3). To find a threshold of detectable expression above background, we normalized all data to establish detectable expression levels for protein-coding genes and lincRNAs (Ramskold et al., 2009; Li et al., 2016) (see Materials and Methods, Supplemental Figure S2A). A normalized transcripts per million (TPM) value >= 0.2 was considered to define 5,586 (84.6%) expressed lincRNAs and 284 (87.4%) pri-miRNAs, whereas a TPM >= 2 was used to define 26,254 (94.9%) expressed protein-coding genes. Globally, the expression breadth, defined as the number of samples in which a gene is expressed, was lower for lincRNAs (median: 12/791 samples) compared to pri-miRNAs (median: 35.5/791 samples) and protein-coding genes (median: 648/791 samples) (Figure 3A). Although the identification of lincRNAs is known to be impacted by the sequencing depth (Cabili et al., 2011; Liu et al., 2012; Sun et al., 2017), we did not observe a clear correlation between the expression breadth and sequencing depth (Pearson correlation coefficient, r=0.04, p=0.86) (Supplemental Figure S2B). This result indicates the reported expression values are not strongly biased by study-specific technical parameters. As previously reported, the expression level of lincRNAs is generally lower than that of pri-miRNAs and protein-coding genes (Figure 3B). The lincRNA loci part of the two "older" categories (Angiosperm and Eudicot/rosid-conserved lincRNAs) showed significantly higher expression levels compared with the three "younger" categories of lincRNAs (Brassicaceae_I_II-conserved, Brassicaceae_I-conserved

and Arabidopsis-specific lincRNA). We also observed a six-fold difference in the fraction of expressed genes depending on their annotation source: lincRNAs annotated in only one resource where more frequently found as undetectable compared to lincRNAs annotated in two or more resources (20% versus 3%, respectively).

To identify groups of samples with similar expression patterns, 769 RNA-Seq samples were split into 22 expression clusters using meta-data curation (22 samples were discarded, see Materials and Methods). Globally, these clusters group Arabidopsis (Col-0) samples according to the organ or tissue considered and the stress, either biotic or abiotic, applied (Supplemental Data Set S3). As lincRNAs are proposed to exert their role in a tissue-specific manner (Liu et al., 2015), the tissue-specificity index (tau) (Yanai et al., 2005) for each lincRNA, pri-miRNA and protein-coding gene was calculated to estimate expression specificity between clusters (Figure 3C). LincRNAs were more specifically expressed than pri-miRNAs and protein-coding genes, with the median tau scores of 0.97, 0.95 and 0.74, respectively ($P<0.001$, Mann–Whitney U-test). Forty-six percent of all expressed lincRNAs (2,573/5,563) with a tau score ≥ 0.97 were defined as lincRNAs showing highly-specific expression (Supplemental Data Set S4). Even though we did not observe a uniform distribution of these highly-specific lincRNA over the 22 clusters (Figure 3D), especially cluster 19 (root), 18 (cell line light induction) and 3 (seedling heat) have the largest number of highly-specific lincRNAs. Between 28 and 42% of the highly-specific expressed lincRNAs were present in the different evolutionary categories, with the highest fraction in the Arabidopsis-specific lincRNAs (Figure 3E). Noteworthy, expression cluster 19, containing 66 samples covering both whole root and specific root cell types (Li et al., 2016), contains the highest fraction (14-17%) of highly-specific lincRNAs in the younger age categories (Figure 3E), hinting towards a role of these lincRNAs in root tissues.

**Figure 3. Expression analysis of lincRNAs. (A)** Line chart showing the distribution of expression breadth for lincRNAs, pri-miRNAs and protein-coding genes across all samples **(B)** Distribution of the maximum TPM expression levels for different lincRNA age categories and gene types (lincRNAs, pri-miRNAs and protein-coding genes). **(C)** Distribution of tissue specificity tau scores for different lincRNA age categories and gene types. Tau scores range from zero to one, where zero means widely expressed, and one means very specifically expressed

(detectable in only one cluster). The black dotted line represents a tau score of 0.97. **(D)** Bar chart reporting the number of expressed and highly-specific expressed lincRNAs for the different expression clusters. Cluster numbers and descriptions are shown below the chart, with numbers in parenthesis indicating the number of samples present per cluster. **(E)** Heatmap showing the proportion of highly-specific expressed lincRNAs in each cluster for each lincRNA age category. Cluster descriptions are the same as in panel D. Numbers in parenthesis report the number of genes per age category together with the fraction of lincRNAs showing highly-specific expression.

To validate these tissue-specific expression patterns, we verified the expression of previously characterized Arabidopsis lincRNAs (Supplemental Table S3). The lincRNA *SVALKA*, which was identified in a cold-sensitive region of the Arabidopsis genome, was maximally expressed in the cold stress-related cluster 5 in our study (Kindgren et al., 2018). In addition, the lincRNA *MARS*, which is involved in the response to abscisic acid, seed germination and root growth under osmotic stress (Roule et al., 2022b), was found to be expressed at high levels in root cluster 19 as well as cluster 11 (root bacteria) and cluster 17 (seed). The lincRNA *ASCO*, reported to be involved in lateral root formation and response to pathogens (Bardou et al., 2014; Rigo et al., 2020), was found to be widely expressed, including cluster 11, which contained samples reporting bacterial flagellin stress responses. The Arabidopsis lncCOBRA1, also conserved in *Brassica rapa* (Kramer et al., 2022) and playing a role in seed germination, was found in our set of Brassicaceae_I_II-conserved lincRNA and showed the highest expression in cluster 17 (seed) and cluster 3 (seedling heat).
Among the highly-specific lincRNAs, the vast majority (66%) were derived from one resource, with lincRNAs identified in Liu et al. 2012 being most abundant (30%, Supplemental Figure S3A-B). Taken together, these findings indicate that the expression clusters offer a good starting point for the context-specific characterization of known and novel lincRNAs. Furthermore, young age categories, including the Brassicaceae-conserved and Arabidopsis-specific

352  lincRNAs, show a bias for expression specificity in the root samples, with more
353  than half (1,448/2,573) of the highly-specific lincRNAs active in root, suggesting a
354  potentially diverse role of lincRNA networks in root growth and development.
355

**Experimental TF-lincRNA regulatory networks reveal active and complex**
356  
**gene regulation of Arabidopsis lincRNA genes**

To integrate lincRNAs into epigenetic and transcriptional networks, we compared
the chromatin states inferred by (Liu et al., 2018; Hazarika et al., 2022) for the
different lincRNA gene sets delineated in our study. These chromatin states
report a combination of multiple epigenetic marks along the genome and offer
detailed insights in the locations and functions of regulatory regions and genes.
Globally, lincRNAs show enrichment for vastly different chromatin states
compared to protein-coding genes (Figure 4A). Chromatin states 33-34, typically
associated with DNA methylation, repressive histone modifications and
transposable elements, were strongly overrepresented in angiosperm-conserved
lincRNAs, and to a lesser extent in lincRNAs showing highly-specific expression
patterns. Chromatin states 31-32, also associated with DNA methylation and
repressive histone modifications were strongly overrepresented in Arabidopsis-
specific lincRNAs. Interestingly, these repressive chromatin marks were less
enriched in the Brassicaceae-I_II-conserved lincRNAs, where chromatin states
13, 19 and 20 were most strongly overrepresented. State 13 refers to Polycomb
group mediated deposition of trimethylation of the lysine 27 of histone H3
(H3K27me3) while states 19-20 denote accessible chromatin and TF ChIP
binding. Chromatin states 19-20, but also states 31-32 and 34, were enriched in
highly-specific lincRNAs. Although we found no evidence of a correlation
between lincRNA age categories and chromatin states, our results revealed that
a significant number of Brassicaceae_I_II-conserved lincRNAs have epigenetic
signatures associated with Polycomb regulation and TF binding in accessible
chromatin.

**Figure 4. Chromatin state and TF ChIP-Seq peak annotation for different**

**gene types (A)** Dendrogram showing the enrichment for different lincRNA gene

sets (x-axis) towards different chromatin states (CS) (y-axis). The values report

the product of -log10(q-value) and the enrichment fold. Only significant

enrichment values are reported (q-value < 0.05). **(B)** The proportion of peak-gene

pairs present in single replicate (blue) and two or more ChIP-Seq replicates

(grey). **(C)** The percentage of three gene types assigned to peaks in two or more

ChIP-Seq replicates. **(D)** Distributions of the number of TF bindings for lincRNA

evolutionary age categories and gene types (lincRNAs, pri-miRNAs and protein-

coding genes).

Based on the specific expression profiles for different lincRNA genes, the
chromatin state information, as well as the high levels of promoter conservation,
we next decided to integrate TF chromatin immunoprecipitation (ChIP) data to
further identify the regulators controlling lincRNAs. Before we used TF ChIP data
to characterize the organization, complexity, and evolution of TF binding for
protein-coding genes (Heyndrickx et al., 2014), hence we here re-processed
publicly available ChIP-Seq to identify TF binding events potentially controlling
lincRNA gene expression. A total of 114 TF ChIP-Seq datasets, covering 45 TFs
with at least two biological replicates, were reprocessed (Supplemental Table S4).
Starting from our genome annotation containing protein-coding genes, pri-
miRNAs and the high-confidence lincRNAs, ChIP-Seq peaks were assigned to
the closest genes. A gene was defined as a potential target gene for a profiled TF
if it was the closest to at least one ChIP-Seq peak of that TF (see Material and
Methods). Based on all 471,078 TF-target gene interactions identified from the
114 ChIP-Seq datasets, 70% of the peak-gene pairs were confirmed by two or
more replicates and were kept to construct a robust TF- target gene regulatory
network (Figure 4B). While most peaks were associated with protein-coding
genes, we identified 48,279 (14.6%) TF peak – lincRNA regulatory interactions
(Figure 4C). Given the strong localization bias for TF binding in Arabidopsis
(Heyndrickx et al., 2014; Yu et al., 2016), we only considered target genes
localized relative to the peak midpoint within a 2kb window (Supplemental Figure
S4A), retaining 95.4% of the binding events (43,233 TF-lincRNA interactions, see
Supplemental Data Set S5). Globally, 2,659 lincRNAs had one or more TF peaks.
Twenty-two out of the 45 TFs, belonging to the bHLH, HD-Zip, bZIP, C2H2,
GRAS, MYB, NAC, NF-YB and NF-YC TF families, were each associated with at
least 200 lincRNAs (Supplemental Table S5).

(Haudry et al., 2013) identified over 90,000 conserved non-coding sequences
(CNS) in Arabidopsis that show evidence of transcriptional and post-
transcriptional regulation. Comparing these CNS with the ChIP-Seq peaks of
target genes revealed that most of the TF binding events close to lincRNAs
(78.1%) and protein-coding genes (73.7%) overlapped with a CNS

(Supplemental Figure S4B). This fraction was much lower for pri-miRNAs
(57.0%). The highest fraction of ChIP-Seq peaks containing a CNS was detected
for Brassicaceae_I_II-conserved lincRNAs loci (91.8%), indicating that these TF
binding events are evolutionary constrained and potentially functional.
Considering the different gene types, the median number of TF binding events
per locus was higher for protein-coding genes and lincRNAs (median of 4 TFs)
compared to pri-miRNAs, suggesting these genes are differently controlled
(Figure 4D). Brassicaceae-conserved lincRNAs have more TF binding events
than lincRNAs in any other age categories (Figure 4D). Furthermore, we
observed a positive correlation between TF binding frequency and the
conservation of lincRNA gene body (r=0.27, P=8.11e-92) (Supplemental Figure
S4C). More precisely, a large faction of lincRNAs without any TF binding event
also has very low phastCons scores close to zero, indicating that these genes
are not under purifying selection. We observed a neglectable correlation between
the number of TF binding events and tissue specificity (r=0.02, p=1.69e-01).
Altogether, lincRNAs experiencing stronger levels of purifying selection are
bound, and potentially regulated, by more TFs, independent of their tissue
specificity (Supplemental Figure S4D). Taken together, we generated a multi-
level genome-wide characterization covering chromatin state information,
promoter conservation, ChIP-based TF binding and CNSs, for all detectable
lincRNA across >700 expression samples, permitting to rapidly define the
biological context and relevance of lincRNAs in Arabidopsis regulatory networks.
447

**MYB44, PIF4 and KAN1 regulate Arabidopsis lincRNAs in different root cell**
**types**

To identify TF-lincRNA regulatory interactions active in specific cellular contexts,
we used the previously defined expression clusters to combine TFs regulation
and lincRNA expression in specific organs, tissues, or stress conditions.
Considering all expressed lincRNAs and all TF peak-based regulatory
interactions described above, we tested if specific expression clusters are

overrepresented for lincRNAs controlled by specific TFs (Y-axis and X-axis in

Figure 5A-B, respectively). We observed a significant overrepresentation for

KANADI1 (KAN1) binding to lincRNA loci in 11 clusters, of which clusters 19

(root), 18 (cell line light induction), 14 (leaf *P. syringae*) and 6 (leaf wounding)

showed the most significant overlaps (Figure 5A). Comparing the different

clusters revealed that root cluster 19 contained numerous enriched TFs,

including Arabidopsis Zinc-Finger protein 1 (AZF1), JAGGED (JAG),

Phytochrome-Interacting Factor 5 (PIF5), Repressor of GA (RGA), PIF1, MYB3,

KAN1 and HAT22. The observed patterns of TF-binding in different expression

clusters were unique for lincRNAs and highly dissimilar compared to TF binding

enrichment for protein-coding genes (Supplemental Figure S5A).

**Figure 5. Overview of TF-lincRNA regulatory interactions in different**

**expression clusters and age categories.** The dot sizes represent the number

of the lincRNAs while the color represents the statistical significance. Cluster

descriptions are the same as in Figure 3D. **(A)** Bubble chart showing the

enrichment of TF binding for expressed lincRNAs in different expression clusters.

**(B)** Bubble chart showing the enrichment of TF binding for highly-expressed

lincRNAs in different expression clusters. **(C)** Bubble chart showing the

enrichment of TF binding for expressed lincRNAs in different age categories.

Focusing on the set of 2,573 lincRNAs showing highly-specific expression, most

of the TF enrichments for root cluster 19 remained (Figure 5B). PIF1, PIF4, KAN1,

HAT22, MYB3, MYB44, ZAT6 and RGA showed the largest overlap and all these

TFs, apart from MYB3 and PIF1, contained >160 lincRNA target genes (Figure

5B and Supplemental Table S6). We confirmed that these eight TFs were all

expressed in one or more samples of the root expression cluster 19

(Supplemental Figure S5B) and earlier studies have reported that these TFs,

apart from HAT22, are involved in root development in Arabidopsis (Hawker and

Bowman, 2004; Devaiah et al., 2007; Moubayidin et al., 2016; Tominaga-Wada

and Wada, 2016; Zhao et al., 2016; Li et al., 2022). Comparing TF binding for the

different age categories revealed that the Brassicaceae_I_II-conserved lincRNAs

were most strongly enriched for these TFs (Figure 5C).

To validate the ChIP-based regulatory interactions for the identified TFs, we used

reverse transcription quantitative PCR (RT-qPCR) analysis in roots of lines

affected in TF expression such as overexpression or T-DNA inactivation (called

TF perturbation lines). We used an inducible line for KAN1 (*KAN1-GR*),

overexpression lines for MYB4 and PIF4 (*35S::MYB4*, 35S*::PIF4*) or quadruple

mutant of the *pif1*, *pif2*, *pif3*, and *pif4* (*pifq*), and a knockout mutant for RGA

(*rga28* T-DNA line). We then selected 27 potentially regulated lincRNAs showing

high expression levels in the root cluster 19 and that were targeted by several of

the selected TFs. For example, *LincRNA5331* and *LincRNA1119* were predicted

to be regulated by the four TFs. We could detect deregulation for *LincRNA5331*

and *LincRNA1119* in *35S::MYB44* and *pifq* roots, whereas the former was also

deregulated in *KAN1-GR* and the latter in *35S::PIF4* (Figure 6A, Supplemental

Figure S6). Overall, out of the 74 inferred regulatory interactions investigated, 36

were confirmed, meaning that for the tested TFs, a significant deregulation of the

lincRNA was found in comparison to the control ("Has a peak and is DE", in

Table 1, Supplemental Table S7). For 23 out of the 27 tested lincRNAs, we

confirmed one or more regulatory interactions (Figure 6A). The precision (i.e. the

positive predictive value), varied between 27-65% while the average accuracy

(i.e. the proportion of correct predictions among all genes examined) was 59%.

For eight interactions the TF peak annotation to the lincRNA was unclear ("Has a

putative peak and is DE" in Table 1), meaning these deregulated lincRNAs might

also resemble confirmed interactions. Lastly, while the 10 interactions where we

observed deregulation in the absence of a peak could indicate false predictions,

they might also represent cases of indirect regulation controlled by the perturbed

TF, influencing the expression of the profiled lincRNAs.

**Figure 6. Experimental validation and characterization of TF-lincRNA regulatory interactions. (A)** Heat map showing log2 fold change (FC) of lincRNA relative expression levels in transcription factor (TF) overexpressing

lines (35S::*MYB44*, 35S::*PIF4* and *KAN1-GR*) or TF knockout lines (*rga28* and

*pifq* (*pif1/pif2/pif3/pif4* quadruple mutants)) vs. control wild-type lines in 14-day

old roots. Expression values were determined by RT-qPCR. Asterisks indicate

statistically significant differences (*$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$) in an

unpaired two-tailed Student's t-Test (n =3). Solid boxes indicate TF ChIP peaks

<2kb from the lincRNA gene. Dashed boxes indicate TF ChIP peaks were

identified only in one ChIP-Seq replicate or the lincRNA is not the closest gene to

the TF peak. **(B)** Heatmap showing KAN1 ChIP-Seq peak-based target genes in

the cluster 19 root expression atlas. The first line shows the expression profile of

the TF under investigation while the other lines show groups of target genes with

similar expression. Next to the group number, the number of target genes is

reported and in parenthesis the fraction of lincRNAs per group is indicated. The

numbers in squares and circles report the number of GWAS root-related and RT-

qPCR experimentally validated lincRNA genes, respectively. **(C)** Heatmap

showing MYB44 ChIP-Seq peak-based target genes in the cluster 19 root

expression atlas. Group information is the same as in panel B. **(D)** Heatmap

showing PIF4 ChIP-Seq peak-based target genes in the cluster 19 root

expression atlas. Group information is the same as in panel B, but here triangles

and circles report the RT-qPCR experimentally validated lincRNA genes for the

*pifq* and *35S::PIF4* lines, respectively.

Based on the experimental validation results confirming that several of the root-

specific lincRNAs are controlled by the TFs inferred using the ChIP-based peaks,

we next integrated genotype-phenotype relationships from the AraGWAS catalog.

The aim of this analysis was to verify if root-related phenotypes have been

reported for regions containing lincRNAs in genome-wide association studies

(GWAS). After processing the significant associations from all GWAS studies

present in the catalog and only retaining associations overlapping with lincRNA

gene bodies or 2kb promoter regions (see Material and Methods), we identified

2,865 single nucleotide polymorphisms (SNPs) overlapping with 1,124 lincRNAs,

covering 147 different studies. While 61.6% of the lincRNAs had only significant

associations via SNPs in the promoter region, 38.4% had associations via SNPs

in the gene body (194 and 238 lincRNAs with significant associations only in the

gene body and in both gene body and promoter, respectively). After parsing and

summarizing phenotype information from the available trait ontology annotations,

we could link 23 lincRNAs to abiotic stress-related traits, 133 lincRNAs to flower-

related traits, 33 lincRNAs to leaf-related traits, 376 lincRNAs to root-related traits,

and 27 lincRNAs to seed-related traits (see Supplemental Data Set S6). The

reprocessed GWAS data indicate that hundreds of regions overlapping with

lincRNAs loci are significantly associated with different plant traits and can be

used to prioritize lincRNAs likely involved in specific biological processes.

To further support the relevance of the identified regulatory network, we focused

on KAN1, MYB44, and PIF4, as these regulators had several root-expressed

lincRNAs target genes that showed differential expression in the TF perturbation

lines (Table 1). Starting from the protein-coding genes and lincRNAs expressed

in root cluster 19 and annotated with a TF peak, we first grouped target genes

with similar expression in the root expression atlas. Secondly, for each group we

determined the fraction of lincRNAs and verified if any of these were overlapping

with our RT-qPCR experimentally validated lincRNAs or had root-related traits in

the GWAS catalog. Finally, we verified if the tissue- or cell-type specific

expression of the different groups was in agreement with the expression of the

regulatory TF. For KAN1, we identified 14 groups that fulfilled these criteria

(Figure 6B), of which eight contained two or more lincRNAs with root-related

traits and several showed strong cell-type specific expression. Group 14,

showing high expression in stele contained 27 lincRNAs (87%) of which five were

associated with GWAS root traits. The high expression of KAN1 in the

meristematic zone agreed with groups 1-2, 8-10, and 25, which together

contained 12 confirmed lincRNAs, either through root-related traits or our

experimental validations. For MYB44, we identified multiple groups containing

GWAS root-related lincRNAs showing specific expression patterns in different

cell types: group 4 contained multiple confirmed target genes expressed in

mature endodermis while group 7 contained 3 confirmed lincRNAs (one GWAS

root-related and two experimentally validated genes) showing high expression in
phloem and companion cells (Figure 6C). For PIF4, which showed highest
expression in mature xylem pole, groups 8 and 10 contained confirmed lincRNA
target genes showing similar cell-type specific expression (Figure 6D). While
other groups contained GWAS root-related lincRNAs showing strong expression
in stele (groups 11, 14 and 16), our expression data does not confirm that PIF4 is
actively expressed in the root stele. All GWAS root-related (58) or experimentally
validated (21) lincRNAs regulated by KAN1, MYB44 or PIF4 can be found in
Supplemental Table S8. This analysis further supports that KAN1, MYB44 and
PIF4 are controlling lincRNA genes showing highly specific expression in
different root tissues and cell types. Additionally, the overlap with genetic
associations hints to a role for many of these TF-controlled lincRNA loci in root
growth and development.
593
594
595
**DISCUSSION**

**Comprehensive annotation of Arabidopsis lincRNAs using transcriptomics**
**and evolutionary genomics**

In contrast to protein-coding genes, the characterization of lincRNAs is more
challenging as we lack highly curated annotations and extensive experimental
observations. Furthermore, the low levels of sequence conservation for the
majority of lincRNA loci makes it difficult to translate biological knowledge learned
in one species to another. Apart from co-expression network analysis, reporting
putative (in)direct associations between lincRNAs and other genes, information
about TF regulation of lincRNAs is scarce. Embedding lincRNAs in biological
networks has great potential to define lincRNAs linked to specific cellular or
morphological phenotypes.
Through the integration of eight lincRNA annotation resources, as well as
mapping various conservation, chromatin, and expression features, we presented

a global view on gene regulation for 5,586 expressed Arabidopsis lincRNAs. We

strongly focused on using replicated samples when processing high-throughput

datasets to obtain high-confidence gene information (Ponting and Haerty, 2022).

Combined with comparative genome analysis yielding information about age

categories and selection acting on gene bodies and promoter regions, we found

that different subsets of lincRNAs have distinct molecular properties. While the

high tissue-specificity and low levels of primary sequence conservation

corroborate previous findings about lincRNAs (Necsulea et al., 2014; Ma et al.,

2019; Palos et al., 2022), the analysis of lincRNA expression using a genome-

wide gene expression atlas covering nearly 800 samples revealed that lincRNA

expression is widespread in different organs and not restricted to stress

conditions (Jha et al., 2020). Focusing on lincRNAs exhibiting highly-specific

expression revealed that the root expression cluster contained a unique set of

1448 lincRNAs, which is the highest number among all expression clusters we

studied. When intersecting the different age categories with the expression

clusters, Brassicaceae_I_II-conserved lincRNAs covered a large fraction of these

root-expressed lincRNAs. While Palos and colleagues also identified a large

number of context-specific lincRNAs expressed in root tip or meristem (Palos et

al., 2022), our age category analysis revealed that many lincRNAs that originated

in the common ancestor of Brassicaceae lineages I and II, showed cell-type

specific expression in various root cell types.

We observed a clear trend of increasing levels of lincRNA gene expression when

going from young to old age categories. However, quantifying selection levels

acting on different loci revealed that the oldest age categories, showing the most

wide-spread and highest expression, are not undergoing the highest levels of

purifying selection. While older categories like angiosperm-conserved and

Eudicot/rosid-conserved lincRNAs had similar median phastCons scores as

protein-coding genes, Brassicaceae_I_II-conserved lincRNAs showed the

highest levels of purifying selection, both in their gene body and promoter. In

animals, a significant increase of dynamically expressed genes and higher levels

of purifying selection has been reported for older age categories (Necsulea et al.,

2014; Sarropoulos et al., 2019). Furthermore, massively parallel reporter assays

surveying thousands of human promoters revealed that tissue-specific lincRNAs

had fewer TF motifs compared to ubiquitously expressed genes (Mattioli et al.,

2019). In contrast to Arabidopsis protein-coding genes, this pattern was not found

for lincRNAs in our analysis (correlation between tau score and TF binding

frequency = 0.02). These results suggest, based on the genome-wide TF binding

data available in Arabidopsis, that the complexity in TF control of lincRNAs is

different between plants and animals and that the observed pattern of highly-

specific expression and purifying selection for Brassicaceae_I_II-conserved

lincRNAs deviates from the global trends observed in animals. Therefore, these

unique properties suggest that these lincRNAs, which only emerged 42 million

years ago, are integrated in a different manner in plant networks when compared

to young lincRNAs in animals.

**Multi-omics regulatory annotation of Arabidopsis lincRNAs**

The regulatory annotation using genome-wide chromatin state information

revealed that protein-coding genes and lincRNAs have distinct chromatin

signatures. For the latter, the enriched chromatin states were largely variable

between, and sometimes within, the different age categories. While states

associated with DNA methylation and repressive histone modifications were most

strongly overrepresented in the youngest (Arabidopsis-specific) and oldest

(angiosperm-conserved) lincRNA age categories, states denoting polycomb

group mediated deposition of H3K27me3 and accessible chromatin were most

enriched for Brassicaceae_I_II-conserved lincRNAs. H3K27me3 is a repressive

covalent histone modification resulting from the activity of Polycomb repressive

complexes. It was recently shown that a reduction in H3K27me3 levels leads to a

decrease in the interactions within Polycomb-associated repressive domains,

resulting in a global reconfiguration of chromatin architecture and transcriptional

reprogramming during plant development (Huang et al., 2021). Chromatin

accessibility is a hallmark of regulatory DNA as it allows sequence-specific

binding of TFs, key components of transcriptional regulatory networks (Schmitz

et al., 2022). The association of these chromatin states with specific sets of

lincRNAs strongly indicates active transcriptional regulation.

To identify context-specific TF regulation potentially driving the highly-specific

expression observed for many lincRNAs, we reprocessed, filtered and annotated

114 ChIP-Seq experiments covering 45 TFs, yielding a TF-lincRNA gene

regulatory network containing 2,659 lincRNAs and 43,233 interactions. To assess

the potential functionality of these inferred regulatory interactions, we overlapped

CNSs identified using nine Brassicaceae genomes, which confirmed that TF

peaks close to lincRNAs show similarly high levels of sequence constraint

compared to protein-coding genes (74-78%). Furthermore, TF binding events

close to Brassicaceae_I_II-conserved lincRNAs showed the highest levels of

CNS conservation (92%), which agrees with the very high phastCons promoter

scores we observed for this age category. While Palos and co-workers reported

that CNSs significantly correlated with gene bodies of Brassicaceae-conserved

lincRNAs (Palos et al., 2022), our results revealed that also lincRNA promoters

and TF binding sites are strongly conserved for Brassicaceae_I-II-conserved

lincRNAs. Ultraconserved CNSs, frequently associating with TF binding sites for

key plant regulators controlling essential biological processes, have been

identified for thousands of protein-coding genes (Van de Velde et al., 2016). Our

results suggest that such deep conservation of cis-regulatory elements is

extremely rare for lincRNAs, as only a small number of lincRNA genes show

deep evolutionary conservation. Such deeply conserved CNSs for protein-coding

genes frequently occur in divergent gene pairs, where they form mini-regulons

representing conserved transcriptional units of co-regulated and co-expressed

neighboring genes (Van de Velde et al., 2016). It is currently unclear if such

conserved transcriptional regulons also exist for lincRNA loci and could explain

the observed patterns of positionally-conserved but sequence-diverged lncRNAs

(Mohammadin et al., 2015).

The integrated TF binding information showed that promotors of lincRNA differ

strongly from those of protein-coding genes, but also revealed high levels of

heterogeneity among the different age categories. In animals, promoters of
protein-coding genes contain more TF binding sites than those of lincRNAs,
suggesting a stronger and more complex transcriptional regulation of the former
(Necsulea et al., 2014; Sarropoulos et al., 2019). When comparing the number of
TF binding events for the different gene types in Arabidopsis, we observed no
difference in TF binding frequency for protein-coding genes and lincRNAs,
indicating that lincRNAs are also regulated in a complex manner in plants. While
no correlation between lincRNA expression tissue-specificity and TF binding
frequency was found, a positive correlation between TF binding frequency and
the level of purifying selection was observed. Again, Brassicaceae_I_II-
conserved lincRNAs stood out having the highest number of binding TFs,
suggesting that neither the age nor the expression of a lincRNA, but its
importance for plant fitness is a major factor in determining its regulatory
complexity. While our results confirm that broadly expressed protein-coding
genes, showing high expression breadth, are positively correlated with the
number of regulating TFs (Heyndrickx et al., 2014), we did not observe this global
trend for lincRNAs, indicating that the regulatory properties of TF control for plant
protein-coding genes and lincRNAs are different. Although the number of
Arabidopsis TFs profiled using ChIP-Seq may be considered as limited, future
research will have to address whether the complexity of TF control varies for
lincRNAs, as well as protein-coding genes, active in different organs, tissues, or
stress conditions.
724

**A TF-lincRNA gene regulatory network identifies KAN1, MYB44 and PIF4 as**
**regulators controlling root lincRNAs**

Through integration of our spatiotemporal expression clusters and the TF-
lincRNA gene regulatory network, we identified eight TF regulators showing a
significant enrichment for TF binding close to lincRNAs specifically expressed in
roots. While the overlap between TF ChIP-Seq peaks and CNSs gave an indirect
indication of the potential functionality of TF binding sites close to hundreds of

lincRNA loci, we experimentally validated a set of inferred regulatory interactions,

focusing on 27 root-expressed lincRNAs and 5 TF perturbation lines. The number

of confirmed regulatory interactions as well as the positive prediction values

found for the tested TFs and lincRNAs here (27-65%) are 3 to 4 times higher

than the fraction of TF-bound protein-coding genes also showing deregulation

reported for a set of TF regulators involved in flowering (7-22%) (O'Maoileidigh et

al., 2014). Compared to the discovery rates obtained for large-scale phenotypic

screens of insertional lines (1.3%) (Ransbotyn et al., 2015), our discovery rates

for deregulation are 20-50 higher. Globally, for 85% of the profiled lincRNAs one

or more confirmed regulatory interactions were found. These findings indicate

that the ChIP peak-based inference of TF regulation is a promising approach to

characterize TF regulation of lincRNAs. Additional deregulated lincRNAs lacking

a TF peak were also identified, which might be due to an indirect effect caused

by crosstalk between different regulators in the Arabidopsis root as well as the

type of mutations chosen in the perturbed TF lines (e.g. partners lacking in

overexpressing lines, compensatory effects in gene families) (Heyndrickx et al.,

2014).

While detailed functional characterizations of lincRNA genes are scarce, the

integration of GWAS information allowed us to identify genetic associations for

1,124 lincRNAs covering various traits. While this number, corresponding to a

frequency of 9%, is slightly lower compared to that of protein-coding genes

associated with a specific trait in the AraGWAS catalog (3,030/27,655 = 11%), it

does confirm the great potential of this largely untapped resource to biologically

characterize lincRNAs potentially controlling different plant traits. As shown for

KAN1, MYB44 and PIF4, multiple groups of co-expressed lincRNAs were

identified bound by one of these TFs. Most of these groups showed strong cell-

type specific expression and contained lincRNAs that were annotated with root-

related traits.

Taken together, the integration of different gene annotations combined with

information about evolutionary conservation, selection, expression, TF regulation

and GWAS data yielded new insights on the biological relevance of hundreds of

Arabidopsis lincRNAs and offers a promising strategy to identify lincRNAs

involved in different aspects of plant biology.

**MATERIALS AND METHODS**

**Prediction of lincRNAs from in-house dataset**

Paired-end RNA-seq datasets with high sequencing depth conducted in previous

projects in the group at the Institute of Plant Sciences Paris-Saclay were used to

predicte additional lincRNAs. All data came from experiments carried out in the

Col0 ecotypes and involved nsra/b mutant in response to NPA/NAA treatment

GSE65717 and GSE116923 (Tran Vdu et al., 2016; Bazin et al., 2018)

modification of expression of the *ASCO* lincRNA GSE135376 (Rigo et al., 2020),

root tip submitted to a short phosphate starvation kinetic GSE128250 (Blein et al.,

2020) and a lateral root initiation kinetic from a binding essay of five time points

without replicates (6h, 12h, 24h, 36h and 48h after binding, unpublished). All

reads were quality trimmed using Trimmomatic. For each library independently,

cleaned reads were mapped on TAIR9 genome sequence with STAR (version

2.7.2a) using Araport11 as a guided annotation with the following additional

parameters: --alignIntronMin 20 --alignIntronMax 3000. For each alignment file,

StringTie (version 2.1.4) was used to predict transcripts using Araport11

annotations as a guide (additional parameters: -c 2.5 -j 10). GFFcompare

(v0.12.6) was then used to isolate the new transcripts in comparison to Araport11

gene annotation (retaining only class code k, m and j). The different transcripts

prediction were then combined using StringTie in merge (additional parameters: -

F 0 -T 0 -c 0 -f 0 -g 0 -i). The final set of transcripts were compared against

Araport11 annotation with GFFcompare removing all transcripts with a class code

of =, c, e, s or m. Coding potential was then assets using COME (Hu et al., 2017),

Coding Potential Calculator CPC (v0.9-r2), CPC2 (Kang et al., 2017) and infernal

(v1.1.2) (Nawrocki and Eddy, 2013) against Rfam v14.1 (Kalvari et al., 2021) with

default parameters. Non-coding transcripts were the ones predicted by CPC,

CPC2 and COME as non-coding and having no hits against tRNA, rRNA, snRNA

or snoRNA genes in Rfam.

**Integration of lincRNA gene annotations**

*Arabidopsis* lincRNAs were collected from public databases: Araport11 (Cheng et

al., 2017), CANTATAdb (Szczesniak et al., 2016), NONCODEv5 (Fang et al.,

2018), PLNlncRbase (Xuan et al., 2015), three publications(Liu et al., 2012;

Nelson et al., 2017; Zhao et al., 2018), and the in-house data.

The pipeline used for the identification of putative lincRNA is described in

Supplemental Figure S1A. (1) LincRNA transcripts with a length of at least 200

bp were retained. (2) Only transcripts that were at least 500 bp away from any

protein-coding gene were retained and considered as intergenic (Liu et al., 2012;

Yamada, 2017). (3) Transcripts lacking strand information were discarded. (4)

The coding potential of transcripts was assessed using the Coding-Non-Coding

Index (CNCI; Version 2) (Sun et al., 2013), CPC2, and Pfam-scan (PFAM) (Finn

et al., 2016), and only those transcripts fulfilling the CPC2 (cutoff < 0), CNCI

(cutoff < 0) and PFAM (E-value 1e-5) criteria were retained. (6) All candidate

intergenic transcripts were assigned to lincRNA loci using the GFFcompare

program (Pertea and Pertea, 2020). The number of transcripts retained after

each filtering step is reported in Supplemental Figure S1B.

**Evolutionary conservation analysis**

Our set of Arabidopsis lincRNAs was classified into distinct evolutionary age

categories based on sequence similarity. The sequences of 6599 lincRNA loci

were extracted using BEDTools getfasta v2.30.0 (Quinlan and Hall, 2010).

Genome sequence data for 40 plant species were obtained from PLAZA 5.0

Dicots and PLAZA 5.0 Monocot (Van Bel et al., 2021), representing 26 eudicots

(20 rosids and 6 asterids ) and 14 monocots. The 20 rosids contain 10

Brassicaceae species. LincRNA homologs were identified using sequence

similarity searches against these 40 genomes using BLASTn (Altschul et al.,

1990) and applying an E-value cutoff of 1e–10 (Nelson et al., 2017).

Classification rules were defined to construct five evolutionary age categories. A
lincRNA was deemed conserved in the Angiosperm evolutionary age category
when at least one homolog was found in eudicots and one homolog in monocots.
LincRNA was defined as a eudicot/rosid-conserved lincRNA with at least one
homolog in rosids, one homolog in asterids, and no homolog in monocots, in
addition with one homolog only in rosids. The lincRNA was assigned as a
Brassicaceae_I_II-conserved lincRNA with at least one homolog in Brassicaceae
lineage I, one homolog in Brassicaceae lineage II, and no homologs outside the
Brassicaceae species. A lincRNA that had at least one homolog in a
Brassicaceae lineage I species, apart from Arabidopsis, was defined as
Brassicaceae_I-conserved lincRNA. The last category, defined as Arabidopsis-
specific lincRNA, were restricted to Arabidopsis lincRNAs without homologs in
any of the other species.

We downloaded the GFF annotation file for 27,655 protein-coding genes and 325
pri-miRNAs from the Araport11 genome release (Cheng et al., 2017) and
extracted gene body sequences using BEDTools getfasta v2.30.0. We also
extracted the promoter region of 2kb upstream of the transcription start site for
lincRNAs, pri-miRNAs, and protein-coding genes. Noted that the promoter of a
gene was shortened when it overlapped with a nearby coding sequence. The
sequence conservation of gene body and promoter region per gene type was
evaluated by average phastCons scores, which were calculated using the
alignments of 20 angiosperm plant genomes (Hupalo and Kern, 2013). Genes
without phastCons score, for example due to missing genome alignment
information, were assigned a score of zero.

**Expression analysis of lincRNAs**

To generate an expression atlas for lincRNAs, pri-miRNAs and protein-coding
genes, we used Curse (Vaneechoutte and Vandepoele, 2019) to search and
curate relevant RNA-seq experiments. Details of the 18 RNA-seq experiments
across all 791 samples are shown in Supplemental Table S2 and Supplemental
Data Set S3. Next, we imported the expression metadata to the Prose tool

(Vaneechoutte and Vandepoele, 2019), which downloads the raw data from SRA
by using the SRA toolkit, performs quality control and adapter detection,
trimmomatic to perform adapter clipping and quality trimming and finally Kallisto
(Pertea et al., 2015) for quantifying transcript expression to normalized
transcripts per million (TPM) values. We downloaded the transcript FASTA file for
27,655 protein-coding genes and 325 pri-miRNAs from the Araport11 genome
release (Cheng et al., 2017) and retrieved transcript sequences for 6,599
lincRNAs using gffread (Pertea and Pertea, 2020). Gene-level atlases were
created by summing the TPM values of all transcripts. We retained TPM value
per gene across the biological replicates and took an average of TPM values per
gene from technical replicates, resulting in an expression atlas covering 791
samples.
We used a simulation experiment (Ramskold et al., 2009; Li et al., 2016) to
determine thresholds for detectable expression in TPM for protein-coding genes
and lincRNAs. The protein-coding genes were used as true positives and the
lincRNAs were used as true negatives. We calculated the false-positive rate and
false-negative rate at different TPM thresholds for each sample. The threshold
TPM of 0.5 balances the numbers of false positives and false negatives.
LincRNAs and pri-miRNAs with TPM >= 0.2 in at least one sample were
considered as expressed. A more stringent threshold TPM >= 2 was used to
define the expression of protein-coding genes.
To identify the clusters containing samples with related gene expression
information, a $\log_2$-transformed gene expression matrix (TPM+1) was used for t-
SNE clustering with perplexity 30 and n_iter = 1000 (Pedregosa et al., 2011). 22
clusters were retained, the largest of which contained 113 samples and the
smallest contained 5 samples. Twenty-two samples were not clustered and
removed (Supplemental Data Set S3). Expression specificity per gene type, for
the gene expression matrix described above, was measured using tissue-
specificity index (tau), defined by the following equation (Yanai et al., 2005):

$$\tau = \frac{\sum_{i=1}^{n}(1-\hat{x}_i)}{n-1}, \hat{X}_i = \frac{xi}{max_{1\leq i \leq n}(xi)}$$

Where $x_i$ was defined as the average TPM value of per cluster and $n$
corresponds to the number of clusters analyzed.

Groups of genes showing similar expression profiles reported in Figure 6 B-D
were identified using CAST clustering and only clusters with at least 10 genes
were retained (Vandepoele et al., 2009).

**Chromatin states analysis and TF peak annotation**

The enrichment of the lincRNA sets in different chromatin states (CSs) of
Arabidopsis from (Liu et al., 2018) was done by shuffling the lincRNA genome
coordinates 1000 times over the whole Arabidopsis genome. Then, we compared
the distribution of the number of lincRNAs expected to overlap by chance with
each CS with the real number of overlaps, and we used these values to calculate
enrichment statistics: p-value as the number of times the real overlap was higher
than the overlap with any of the 1000 shuffled lincRNA sets, and enrichment fold
as the real overlap divided by the median of overlap expected by chance (median
of the 1000 shuffling events). The p-value was adjusted for multiple testing using
Benjamini-Hochberg correction (significance level 0.05). For visualization
purposes, the two enrichment metrics were combined into the π-value, which is
the -log10(p-value)*enrichment fold.

TF ChIP-Seq peak coordinates were retrieved from the PlantPAN 3.0 database
(Chow et al., 2019) and (Song et al., 2016). The original ChIP-Seq of KAN1,
MYB44 and PIF4 were derived from these studies (Merelo et al., 2013; Pfeiffer et
al., 2014; Song et al., 2016). For each peak the closest gene was identified and
only peaks confirmed in two or more replicates and within a 2kb window of the
gene body were retained. Starting from all TF – target gene pairs (either a
protein-coding gene or a lincRNAs; Supplemental Data Set S5), enrichment
analysis was performed to identify enriched TFs in different expression clusters
or age categories. For all enrichment analyses the hypergeometric distribution
was applied and the q-value of enrichment was determined using the Benjamini–
Hochberg correction for multiple hypotheses testing.

**Identification of GWAS-associated genes**

GWAS data were collected from AraGWAS (Togninalli et al., 2020) and
overlapped with the gene body and promoter 2kb sequences of lincRNAs to
associate with the phenotype of interest. All significant associations (Permutation
threshold <0.05 and FDR <0.05) were retained for screening for minor allele
frequency>0.01, resulting in 1,124 lincRNAs, covering 147 different studies.
These significantly associated lincRNAs were classified into five main traits by
ontological annotation, including root, seed, flower, leaf, and abiotic-related traits
(see Supplemental Data Set S6).

**Validation of TF-lincRNA regulatory interactions using reverse**
919
**transcription-quantitative PCR**
920

Total RNA from five to eight 14-day roots grown vertically on solid MS ½ media
was extracted using TRI Reagent (Sigma-Aldrich) and digested with RNase-free
DNase (Fermentas) following the manufacturer's recommendations. cDNA was
synthetized using Maxima Reverse Transcriptase (Thermo Scientific). Expression
analysis by RT-qPCR was performed using SYBR Green master I (Roche) and
the LightCycler® 96 system following a standard protocol (40 cycles, 60°C
annealing). Data were analyzed using the ΔΔCt method with PP2A (PROTEIN
PHOSPHATASE 2A SUBUNIT A3 (AT1G13320)) as reference transcript for
normalization of RT-qPCR data. WT Col-0 plants grown at the same time were
used as sample reference. For the dexamethasone inducible KAN1-GR
expression lines analysis, after 14 days on MS media, the plants were transferred
for one day either on dexamethasone containing plates (10µM) or only DMSO
(dexamethasone solvent) as control. Primers used are listed in Supplemental
Table S9. Three biological replicates were performed per condition. Statistical
analyses were performed using the unpaired two-tailed Student's t-Test
(GraphPad prism).

**SUPPLEMENTAL DATA**

**Supplemental Figure S1.** Integration of lincRNA resources.

**Supplemental Figure S2.** Definition of expressed genes and influence of
sequencing depth.

**Supplemental Figure S3.** Resource annotation for highly-specific expressed
lincRNAs.

**Supplemental Figure S4.** Regulatory properties of lincRNAs.

**Supplemental Figure S5.** TF binding and expression.

**Supplemental Figure S6.** RT-qPCR validation results.

**Supplemental Table S1.** The number of homologous lincRNAs aligned to the
genomes of 40 species.

**Supplemental Table S2.** Summary of RNA sequencing data.

**Supplemental Table S3.** List of lincRNAs with known functions.

**Supplemental Table S4.** Overview of TFs ChIP-Seq data used in this study.

**Supplemental Table S5.** TF and the absolute number of target genes per gene
type.

**Supplemental Table S6.** Overview of TFs showing significant enrichment for
binding to highly-specific expressed lincRNAs for the different expression clusters
(corresponds to Figure 5B).

**Supplemental Table S7.** Overview of lincRNA peak annotation and RT-qPCR
validation.

**Supplemental Table S8.** RT-qPCR confirmed/associated with GWAS root traits
lincRNAs expression information for the different samples of expression cluster
19 (root) (corresponds to Figure 6B-D).

**Supplemental Table S9.** RT-qPCR primer sequences used in this study.

**Supplemental Data Set S1.** Full list of putative lincRNAs integrated from
different resources.

**Supplemental Data Set S2.** PhastCons Scores for lincRNA loci and evolutionary
age category.

**Supplemental Data Set S3.** Metadata for 791 samples of RNA-seq and cluster
information.

**Supplemental Data Set S4.** The list of the highly-specific lincRNAs.

**Supplemental Data Set S5.** The table of TF-lincRNA interactions.

**Supplemental Data Set S6.** The lincRNA loci containing significant GWAS hits.

L.L., M.C., T.B., and K.V., designed the research. L.L., T.D., and T.B., performed

data analysis. N.M.P. performed chromatin state analysis. M.H. performed

experimental work. L.L., T.B., and K.V., wrote the manuscript.

**ACKNOWLEDGEMENTS**

We thank Mariek Dubois, Andrés Ritter, Tereza Vavrdova, Freya De Winter,

Rebecca De Clercq for kindly providing the Arabidopsis lines. We thank Olivier

Martin and Federico Ariel for providing comments on the manuscript. The IPS2 is

benefited from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007).

L.L. is supported by the China Scholarship Council for a PhD fellowship

(201808530499). M.H. is supported by a PhD fellowship from the Fondation pour

la Recherche Médicale (ECO202106013730).

**FIGURE LEGENDS: see inline figure captions**

**TABLES**

**Table 1. Summary of qPCR validation for TF-lincRNA gene pairs.**

| Line | Total number of validated lincRNAs (1) | Has a peak and is DE (2) | No peak and not DE | Has a peak and is not DE | No peak and is DE | Has a putative peak and is DE (3) | Accuracy (4) | Precision | Recall |
|------|------|------|------|------|------|------|------|------|------|
| 35S:*MYB44* | 24 | 11 | 5 | 6 | 0 | 2 | 0.73 | 0.65 | 1.00 |
| 35S:*PIF4* | 26 | 8 | 4 | 9 | 4 | 1 | 0.48 | 0.47 | 0.67 |
| KAN1-GR | 24 | 6 | 6 | 7 | 3 | 2 | 0.55 | 0.46 | 0.67 |
| Pifq | 25 | 8 | 6 | 8 | 1 | 2 | 0.61 | 0.50 | 0.89 |
| rga-28 | 25 | 3 | 11 | 8 | 2 | 1 | 0.58 | 0.27 | 0.60 |

(1) In total 27 lincRNAs were tested in one or more lines.

(2) DE refers to differential expression, indicating deregulation in a TF

perturbation line.

(3) These peak-lincRNA pairs do not satisfy our stringent peak definition (present

in two or more replicates and present within 2kb of the gene body). Here a peak

is only identified in one ChIP-Seq replicate or the lincRNA is not the closest

target gene to this peak.

(4) The genes present in the category "Has a putative peak and is DE" were

excluded to compute Accuracy, Precision and Recall.

1000

# REFERENCES

**Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J.** (1990). Basic local alignment search tool. J Mol Biol **215,** 403-410.

**Ariel, F., Lucero, L., Christ, A., Mammarella, M.F., Jegu, T., Veluchamy, A., Mariappan, K., Latrasse, D., Blein, T., Liu, C., Benhamed, M., and Crespi, M.** (2020). R-Loop Mediated trans Action of the APOLO Long Noncoding RNA. Mol Cell **77,** 1055-1065 e1054.

**Bardou, F., Ariel, F., Simpson, C.G., Romero-Barrios, N., Laporte, P., Balzergue, S., Brown, J.W., and Crespi, M.** (2014). Long noncoding RNA modulates alternative splicing regulators in Arabidopsis. Dev Cell **30,** 166-176.

**Bazin, J., Romero, N., Rigo, R., Charon, C., Blein, T., Ariel, F., and Crespi, M.** (2018). Nuclear Speckle RNA Binding Proteins Remodel Alternative Splicing and the Non-coding Arabidopsis Transcriptome to Regulate a Cross-Talk Between Auxin and Immune Responses. Frontiers in Plant Science **9**.

**Beilstein, M.A., Nagalingum, N.S., Clements, M.D., Manchester, S.R., and Mathews, S.** (2010). Dated molecular phylogenies indicate a Miocene origin for Arabidopsis thaliana. Proc Natl Acad Sci U S A **107,** 18724-18728.

**Bhogireddy, S., Mangrauthia, S.K., Kumar, R., Pandey, A.K., Singh, S., Jain, A., Budak, H., Varshney, R.K., and Kudapa, H.** (2021). Regulatory non-coding RNAs: a new frontier in regulation of plant biology. Functional & Integrative Genomics **21,** 313-330.

**Blein, T., Balzergue, C., Roule, T., Gabriel, M., Scalisi, L., Francois, T., Sorin, C., Christ, A., Godon, C., Delannoy, E., Martin-Magniette, M.L., Nussaume, L., Hartmann, C., Gautheret, D., Desnos, T., and Crespi, M.** (2020). Landscape of the Noncoding Transcriptome Response of Two Arabidopsis Ecotypes to Phosphate Starvation. Plant Physiol **183,** 1058-1072.

**Bu, D.C., Luo, H.T., Jiao, F., Fang, S.S., Tan, C.F., Liu, Z.Y., and Zhao, Y.** (2015). Evolutionary annotation of conserved long non-coding RNAs in major mammalian species. Science China-Life Sciences **58,** 787-798.

**Cabili, M.N., Trapnell, C., Goff, L., Koziol, M., Tazon-Vega, B., Regev, A., and Rinn, J.L.** (2011). Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. Genes Dev **25,** 1915-1927.

**Chen, L., Zhu, Q.H., and Kaufmann, K.** (2020). Long non-coding RNAs in plants: emerging modulators of gene activity in development and stress responses. Planta **252,** 92.

**Chen, Q.S., Liu, K., Yu, R., Zhou, B.L., Huang, P.P., Cao, Z.X., Zhou, Y.Q., and Wang, J.H.** (2021). From "Dark Matter" to "Star": Insight Into the Regulation Mechanisms of Plant Functional Long Non-Coding RNAs. Frontiers in Plant Science **12**.

**Cheng, C.Y., Krishnakumar, V., Chan, A.P., Thibaud-Nissen, F., Schobel, S., and Town, C.D.** (2017). Araport11: a complete reannotation of the Arabidopsis thaliana reference genome. Plant J **89,** 789-804.

**Chow, C.N., Lee, T.Y., Hung, Y.C., Li, G.Z., Tseng, K.C., Liu, Y.H., Kuo, P.L., Zheng, H.Q., and Chang, W.C.** (2019). PlantPAN3.0: a new and updated resource for reconstructing transcriptional regulatory networks from ChIP-seq experiments in plants. Nucleic Acids Research **47,** D1155-D1163.

**Deng, F., Zhang, X., Wang, W., Yuan, R., and Shen, F.** (2018). Identification of Gossypium hirsutum long non-coding RNAs (lncRNAs) under salt stress. BMC Plant Biol **18,** 23.

**Devaiah, B.N., Nagarajan, V.K., and Raghothama, K.G.** (2007). Phosphate homeostasis and root development in Arabidopsis are synchronized by the zinc finger transcription factor ZAT6. Plant Physiol **145,** 147-159.

**Fang, S.S., Zhang, L.L., Guo, J.C., Niu, Y.W., Wu, Y., Li, H., Zhao, L.H., Li, X.Y., Teng, X.Y., Sun, X.H., Sun, L., Zhang, M.Q., Chen, R.S., and Zhao, Y.** (2018). NONCODEV5: a comprehensive annotation database for long non-coding RNAs. Nucleic Acids Research **46,** D308-D314.

**Finn, R.D., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Mistry, J., Mitchell, A.L., Potter, S.C., Punta, M., Qureshi, M., Sangrador-Vegas, A., Salazar, G.A., Tate, J., and Bateman, A.** (2016). The Pfam protein families database: towards a more sustainable future. Nucleic Acids Res **44,** D279-285.

**Fukuda, M., Nishida, S., Kakei, Y., Shimada, Y., and Fujiwara, T.** (2019). Genome-Wide Analysis of Long Intergenic Noncoding RNAs Responding to Low-Nutrient Conditions in Arabidopsis thaliana: Possible Involvement of Trans-Acting siRNA3 in Response to Low Nitrogen. Plant Cell Physiol **60,** 1961-1973.

**Haudry, A., Platts, A.E., Vello, E., Hoen, D.R., Leclercq, M., Williamson, R.J., Forczek, E., Joly-Lopez, Z., Steffen, J.G., Hazzouri, K.M., Dewar, K., Stinchcombe, J.R., Schoen, D.J., Wang, X., Schmutz, J., Town, C.D., Edger, P.P., Pires, J.C., Schumaker, K.S., Jarvis, D.E., Mandakova, T., Lysak, M.A., van den Bergh, E., Schranz, M.E., Harrison, P.M., Moses, A.M., Bureau, T.E., Wright, S.I., and Blanchette, M.** (2013). An atlas of over 90,000 conserved noncoding sequences provides insight into crucifer regulatory regions. Nat Genet **45,** 891-898.

**Hawker, N.P., and Bowman, J.L.** (2004). Roles for Class III HD-Zip and KANADI genes in Arabidopsis root development. Plant Physiol **135,** 2261-2270.

**Hazarika, R.R., Serra, M., Zhang, Z., Zhang, Y., Schmitz, R.J., and Johannes, F.** (2022). Molecular properties of epimutation hotspots. Nat Plants **8,** 146-156.

**He, H., Zhou, Y.F., Yang, Y.W., Zhang, Z., Lei, M.Q., Feng, Y.Z., Zhang, Y.C., Chen, Y.Q., Lian, J.P., and Yu, Y.** (2021). Genome-Wide Analysis Identified a Set of Conserved lncRNAs Associated with Domestication-Related Traits in Rice. International Journal of Molecular Sciences **22**.

**Heo, J.B., and Sung, S.** (2011). Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. Science **331,** 76-79.

**Heyndrickx, K.S., Van de Velde, J., Wang, C., Weigel, D., and Vandepoele, K.** (2014). A functional and evolutionary perspective on transcription factor binding in Arabidopsis thaliana. Plant Cell **26,** 3894-3910.

**Hu, L., Xu, Z., Hu, B., and Lu, Z.J.** (2017). COME: a robust coding potential calculation tool for lncRNA identification and characterization based on multiple features. Nucleic Acids Res **45,** e2.

**Huang, Y., Sicar, S., Ramirez-Prado, J.S., Manza-Mianza, D., Antunez-Sanchez, J., Brik-Chaouche, R., Rodriguez-Granados, N.Y., An, J., Bergounioux, C., Mahfouz, M.M., Hirt, H., Crespi, M., Concia, L., Barneche, F., Amiard, S., Probst, A.V., Gutierrez-Marcos, J., Ariel, F., Raynaud, C., Latrasse, D., and Benhamed, M.** (2021). Polycomb-dependent differential chromatin compartmentalization determines gene coregulation in Arabidopsis. Genome Research **31,** 1230-+.

**Hupalo, D., and Kern, A.D.** (2013). Conservation and functional element discovery in 20 angiosperm plant genomes. Mol Biol Evol **30,** 1729-1744.

Jha, U.C., Nayyar, H., Jha, R., Khurshid, M., Zhou, M., Mantri, N., and Siddique, K.H.M. (2020). Long non-coding RNAs: emerging players regulating plant abiotic stress response and adaptation. BMC Plant Biol **20,** 466.

Jones-Rhoades, M.W., Bartel, D.P., and Bartel, B. (2006). MicroRNAs and their regulatory roles in plants. Annual Review of Plant Biology **57,** 19-53.

Kalvari, I., Nawrocki, E.P., Ontiveros-Palacios, N., Argasinska, J., Lamkiewicz, K., Marz, M., Griffiths-Jones, S., Toffano-Nioche, C., Gautheret, D., Weinberg, Z., Rivas, E., Eddy, S.R., Finn, R.D., Bateman, A., and Petrov, A.I. (2021). Rfam 14: expanded coverage of metagenomic, viral and microRNA families. Nucleic Acids Res **49,** D192-D200.

Kang, Y.J., Yang, D.C., Kong, L., Hou, M., Meng, Y.Q., Wei, L., and Gao, G. (2017). CPC2: a fast and accurate coding potential calculator based on sequence intrinsic features. Nucleic Acids Res **45,** W12-W16.

Ke, L.L., Zhou, Z.W., Xu, X.W., Wang, X., Liu, Y.L., Xu, Y.T., Huang, Y., Wang, S.T., Deng, X.X., Chen, L.L., and Xu, Q. (2019). Evolutionary dynamics of lincRNA transcription in nine citrus species. Plant Journal **98,** 912-927.

Kim, D.H., and Sung, S. (2017). Vernalization-Triggered Intragenic Chromatin Loop Formation by Long Noncoding RNAs. Dev Cell **40,** 302-312 e304.

Kindgren, P., Ard, R., Ivanov, M., and Marquardt, S. (2018). Transcriptional read-through of the long non-coding RNA SVALKA governs plant cold acclimation. Nat Commun **9,** 4561.

Kramer, M.C., Kim, H.J., Palos, K.R., Garcia, B.A., Lyons, E., Beilstein, M.A., Nelson, A.D.L., and Gregory, B.D. (2022). A Conserved Long Intergenic Non-coding RNA Containing snoRNA Sequences, lncCOBRA1, Affects Arabidopsis Germination and Development. Front Plant Sci **13,** 906603.

Li, Q.Q., Zhang, Z., Zhang, C.X., Wang, Y.L., Liu, C.B., Wu, J.C., Han, M.L., Wang, Q.X., and Chao, D.Y. (2022). Phytochrome-interacting factors orchestrate hypocotyl adventitious root initiation in Arabidopsis. Development **149**.

Li, S., Yamada, M., Han, X., Ohler, U., and Benfey, P.N. (2016). High-Resolution Expression Map of the Arabidopsis Root Reveals Alternative Splicing and lincRNA Regulation. Dev Cell **39,** 508-522.

Li, S.X., Yu, X., Lei, N., Cheng, Z.H., Zhao, P.J., He, Y.K., Wang, W.Q., and Peng, M. (2017). Genome-wide identification and functional prediction of cold and/or drought-responsive lncRNAs in cassava. Scientific Reports **7**.

Liu, F., Marquardt, S., Lister, C., Swiezewski, S., and Dean, C. (2010). Targeted 3' processing of antisense transcripts triggers Arabidopsis FLC chromatin silencing. Science **327,** 94-97.

Liu, J., Wang, H., and Chua, N.H. (2015). Long noncoding RNA transcriptome of plants. Plant Biotechnol J **13,** 319-328.

Liu, J., Jung, C., Xu, J., Wang, H., Deng, S.L., Bernad, L., Arenas-Huertero, C., and Chua, N.H. (2012). Genome-Wide Analysis Uncovers Regulation of Long Intergenic Noncoding RNAs in Arabidopsis. Plant Cell **24,** 4333-4345.

Liu, Y., Tian, T., Zhang, K., You, Q., Yan, H., Zhao, N., Yi, X., Xu, W., and Su, Z. (2018). PCSD: a plant chromatin state database. Nucleic Acids Res **46,** D1157-D1167.

Lucero, L., Ferrero, L., Fonouni-Farde, C., and Ariel, F. (2021). Functional classification of plant long noncoding RNAs: a transcript is known by the company it keeps. New Phytologist **229,** 1251-1260.

Ma, J.C., Bai, X.T., Luo, W.C., Feng, Y.N., Shao, X.M., Bai, Q.X., Sun, S.J., Long, Q.M., and Wan, D.S. (2019). Genome-Wide Identification of Long Noncoding

RNAs and Their Responses to Salt Stress in Two Closely Related Poplars.
Frontiers in Genetics **10**.

**Mattioli, K., Volders, P.J., Gerhardinger, C., Lee, J.C., Maass, P.G., Mele, M., and**
**Rinn, J.L.** (2019). High-throughput functional analysis of lncRNA core promoters
elucidates rules governing tissue specificity. Genome Research **29**, 344-355.

**Merelo, P., Xie, Y.K., Brand, L., Ott, F., Weigel, D., Bowman, J.L., Heisler, M.G., and**
**Wenkel, S.** (2013). Genome-Wide Identification of KANADI1 Target Genes. Plos
One **8**.

**Mohammadin, S., Edger, P.P., Pires, J.C., and Schranz, M.E.** (2015). Positionally-
conserved but sequence-diverged: identification of long non-coding RNAs in the
Brassicaceae and Cleomaceae. BMC Plant Biol **15**, 217.

**Moison, M., Pacheco, J.M., Lucero, L., Fonouni-Farde, C., Rodriguez-Melo, J.,**
**Mansilla, N., Christ, A., Bazin, J., Benhamed, M., Ibanez, F., Crespi, M.,**
**Estevez, J.M., and Ariel, F.** (2021). The lncRNA APOLO interacts with the
transcription factor WRKY42 to trigger root hair cell expansion in response to
cold. Mol Plant **14**, 937-948.

**Moubayidin, L., Salvi, E., Giustini, L., Terpstra, I., Heidstra, R., Costantino, P., and**
**Sabatini, S.** (2016). A SCARECROW-based regulatory circuit controls
Arabidopsis thaliana meristem size from the root endodermis. Planta **243**, 1159-
1168.

**Nawrocki, E.P., and Eddy, S.R.** (2013). Infernal 1.1: 100-fold faster RNA homology
searches. Bioinformatics **29**, 2933-2935.

**Necsulea, A., Soumillon, M., Warnefors, M., Liechti, A., Daish, T., Zeller, U., Baker,**
**J.C., Grutzner, F., and Kaessmann, H.** (2014). The evolution of lncRNA
repertoires and expression patterns in tetrapods. Nature **505**, 635-+.

**Nelson, A.D.L., Devisetty, U.K., Palos, K., Haug-Baltzell, A.K., Lyons, E., and**
**Beilstein, M.A.** (2017). Evolinc:A Tool for the Identification and Evolutionary
Comparison of Long Intergenic Non-coding RNAs. Frontiers in Genetics **8**.

**Nelson, A.D.L., Forsythe, E.S., Devisetty, U.K., Clausen, D.S., Haug-Batzell, A.K.,**
**Meldrum, A.M.R., Frank, M.R., Lyons, E., and Beilstein, M.A.** (2016). A
Genomic Analysis of Factors Driving lincRNA Diversification: Lessons from
Plants. G3-Genes Genomes Genetics **6**, 2881-2891.

**O'Maoileidigh, D.S., Graciet, E., and Wellmer, F.** (2014). Gene networks controlling
Arabidopsis thaliana flower development. New Phytol **201**, 16-30.

**Pacheco, J.M., Mansilla, N., Moison, M., Lucero, L., Gabarain, V.B., Ariel, F., and**
**Estevez, J.M.** (2021). The lncRNA APOLO and the transcription factor WRKY42
target common cell wall EXTENSIN encoding genes to trigger root hair cell
elongation. Plant Signal Behav **16**, 1920191.

**Palos, K., Dittrich, A.C.N., Yu, L.A., Brock, J.R., Railey, C.E., Wu, H.Y.L.,**
**Sokolowska, E., Skirycz, A., Hsu, P.Y., Gregory, B.D., Lyons, E., Beilstein,**
**M.A., and Nelson, A.D.L.** (2022). Identification and functional annotation of long
intergenic non-coding RNAs in Brassicaceae. Plant Cell.

**Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O.,**
**Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos,**
**A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E.** (2011).
Scikit-learn: Machine Learning in Python. Journal of Machine Learning Research
**12**, 2825-2830.

**Pertea, G., and Pertea, M.** (2020). GFF Utilities: GffRead and GffCompare. F1000Res **9**.

**Pfeiffer, A., Shi, H., Tepperman, J.M., Zhang, Y., and Quail, P.H.** (2014). Combinatorial Complexity in a Transcriptionally Centered Signaling Hub in Arabidopsis. Molecular Plant **7,** 1598-1618.

**Ponting, C.P., and Haerty, W.** (2022). Genome-Wide Analysis of Human Long Noncoding RNAs: A Provocative Review. Annu Rev Genomics Hum Genet.

**Qi, X., Xie, S., Liu, Y., Yi, F., and Yu, J.** (2013). Genome-wide annotation of genes and noncoding RNAs of foxtail millet in response to simulated drought stress by deep sequencing. Plant Mol Biol **83,** 459-473.

**Qin, T., Zhao, H., Cui, P., Albesher, N., and Xiong, L.** (2017). A Nucleus-Localized Long Non-Coding RNA Enhances Drought and Salt Stress Tolerance. Plant Physiol **175,** 1321-1336.

**Quinlan, A.R., and Hall, I.M.** (2010). BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics **26,** 841-842.

**Rai, M.I., Alam, M., Lightfoot, D.A., Gurha, P., and Afzal, A.J.** (2019). Classification and experimental identification of plant long non-coding RNAs. Genomics **111,** 997-1005.

**Ramskold, D., Wang, E.T., Burge, C.B., and Sandberg, R.** (2009). An abundance of ubiquitously expressed genes revealed by tissue transcriptome sequence data. PLoS Comput Biol **5,** e1000598.

**Ransbotyn, V., Yeger-Lotem, E., Basha, O., Acuna, T., Verduyn, C., Gordon, M., Chalifa-Caspi, V., Hannah, M.A., and Barak, S.** (2015). A combination of gene expression ranking and co-expression network analysis increases discovery rate in large-scale mutant screens for novel Arabidopsis thaliana abiotic stress genes. Plant Biotechnol J **13,** 501-513.

**Ransohoff, J.D., Wei, Y., and Khavari, P.A.** (2018). The functions and unique features of long intergenic non-coding RNA. Nat Rev Mol Cell Biol **19,** 143-157.

**Rigo, R., Bazin, J., Romero-Barrios, N., Moison, M., Lucero, L., Christ, A., Benhamed, M., Blein, T., Huguet, S., Charon, C., Crespi, M., and Ariel, F.** (2020). The Arabidopsis lncRNA ASCO modulates the transcriptome through interaction with splicing factors. EMBO Rep **21,** e48977.

**Roule, T., Crespi, M., and Blein, T.** (2022a). Regulatory long non-coding RNAs in root growth and development. Biochem Soc Trans **50,** 403-412.

**Roule, T., Christ, A., Hussain, N., Huang, Y., Hartmann, C., Benhamed, M., Gutierrez-Marcos, J., Ariel, F., Crespi, M., and Blein, T.** (2022b). The lncRNA MARS modulates the epigenetic reprogramming of the marneral cluster in response to ABA. Mol Plant **15,** 840-856.

**Sanchita, Trivedi, P.K., and Asif, M.H.** (2020). Updates on plant long non-coding RNAs (lncRNAs): the regulatory components. Plant Cell Tissue and Organ Culture **140,** 259-269.

**Sarropoulos, I., Marin, R., Cardoso-Moreira, M., and Kaessmann, H.** (2019). Developmental dynamics of lncRNAs across mammalian organs and species. Nature **571,** 510-+.

**Schmitz, R.J., Grotewold, E., and Stam, M.** (2022). Cis-regulatory sequences in plants: Their importance, discovery, and future challenges. Plant Cell **34,** 718-741.

**Severing, E., Faino, L., Jamge, S., Busscher, M., Kuijer-Zhang, Y., Bellinazzo, F., Busscher-Lange, J., Fernandez, V., Angenent, G.C., Immink, R.G.H., and Pajoro, A.** (2018). Arabidopsis thaliana ambient temperature responsive lncRNAs. BMC Plant Biol **18,** 145.

**Shafiq, S., Li, J.R., and Sun, Q.W.** (2016). Functions of plants long non-coding RNAs. Biochimica Et Biophysica Acta-Gene Regulatory Mechanisms **1859,** 155-162.

**Shea, D.J., Nishida, N., Takada, S., Itabashi, E., Takahashi, S., Akter, A., Miyaji, N.,**
**Osabe, K., Mehraj, H., Shimizu, M., Seki, M., Kakizaki, T., Okazaki, K.,**
**Dennis, E.S., and Fujimoto, R.** (2019). Long noncoding RNAs in Brassica rapa
L. following vernalization. Sci Rep **9,** 9302.
**Shuai, P., Liang, D., Tang, S., Zhang, Z.J., Ye, C.Y., Su, Y.Y., Xia, X.L., and Yin, W.L.**
(2014). Genome-wide identification and functional prediction of novel and
drought-responsive lincRNAs in Populus trichocarpa. J Exp Bot **65,** 4975-4983.
**Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K.,**
**Clawson, H., Spieth, J., Hillier, L.W., Richards, S., Weinstock, G.M., Wilson,**
**R.K., Gibbs, R.A., Kent, W.J., Miller, W., and Haussler, D.** (2005).
Evolutionarily conserved elements in vertebrate, insect, worm, and yeast
genomes. Genome Res **15,** 1034-1050.
**Song, L., Huang, S.S.C., Wise, A., Castanon, R., Nery, J.R., Chen, H.M., Watanabe,**
**M., Thomas, J., Bar-Joseph, Z., and Ecker, J.R.** (2016). A transcription factor
hierarchy defines an environmental stress response network. Science **354**.
**Sun, L., Luo, H., Bu, D., Zhao, G., Yu, K., Zhang, C., Liu, Y., Chen, R., and Zhao, Y.**
(2013). Utilizing sequence intrinsic composition to classify protein-coding and
long non-coding transcripts. Nucleic Acids Research **41**.
**Sun, Z., Nair, A., Chen, X., Prodduturi, N., Wang, J., and Kocher, J.P.** (2017).
UClncR: Ultrafast and comprehensive long non-coding RNA detection from RNA-
seq. Sci Rep **7,** 14196.
**Szczesniak, M.W., Rosikiewicz, W., and Makalowska, I.** (2016). CANTATAdb: A
Collection of Plant Long Non-Coding RNAs. Plant Cell Physiol **57,** e8.
**Szczesniak, M.W., Kubiak, M.R., Wanowska, E., and Makalowska, I.** (2021).
Comparative genomics in the search for conserved long noncoding RNAs.
Essays Biochem **65,** 741-749.
**Togninalli, M., Seren, U., Freudenthal, J.A., Monroe, J.G., Meng, D., Nordborg, M.,**
**Weigel, D., Borgwardt, K., Korte, A., and Grimm, D.G.** (2020). AraPheno and
the AraGWAS Catalog 2020: a major database update including RNA-Seq and
knockout mutation data for Arabidopsis thaliana. Nucleic Acids Res **48,** D1063-
D1068.
**Tominaga-Wada, R., and Wada, T.** (2016). The ectopic localization of CAPRICE LIKE
MYB3 protein in Arabidopsis root epidermis. J Plant Physiol **199,** 111-115.
**Tran Vdu, T., Souiai, O., Romero-Barrios, N., Crespi, M., and Gautheret, D.** (2016).
Detection of generic differential RNA processing events from RNA-seq data. RNA
Biol **13,** 59-67.
**Ulitsky, I.** (2016). Evolution to the rescue: using comparative genomics to understand
long non-coding RNAs. Nature Reviews Genetics **17,** 601-614.
**Van Bel, M., Silvestri, F., Weitz, E.M., Kreft, L., Botzki, A., Coppens, F., and**
**Vandepoele, K.** (2021). PLAZA 5.0: extending the scope and power of
comparative and functional genomics in plants. Nucleic Acids Res.
**Van de Velde, J., Van Bel, M., Vaneechoutte, D., and Vandepoele, K.** (2016). A
Collection of Conserved Noncoding Sequences to Study Gene Regulation in
Flowering Plants. Plant Physiol **171,** 2586-2598.
**Vandepoele, K., Quimbaya, M., Casneuf, T., De Veylder, L., and Van de Peer, Y.**
(2009). Unraveling Transcriptional Control in Arabidopsis Using cis-Regulatory
Elements and Coexpression Networks. Plant Physiology **150,** 535-546.
**Vaneechoutte, D., and Vandepoele, K.** (2019). Curse: building expression atlases and
co-expression networks from public RNA-Seq data. Bioinformatics **35,** 2880-2881.

**Wang, H., Niu, Q.W., Wu, H.W., Liu, J., Ye, J., Yu, N., and Chua, N.H.** (2015a).
Analysis of non-coding transcriptome in rice and maize uncovers roles of
conserved lncRNAs associated with agriculture traits. Plant Journal **84,** 404-416.
**Wang, H., Niu, Q.W., Wu, H.W., Liu, J., Ye, J., Yu, N., and Chua, N.H.** (2015b).
Analysis of non-coding transcriptome in rice and maize uncovers roles of
conserved lncRNAs associated with agriculture traits. Plant J **84,** 404-416.
**Wu, H., Yang, L., and Chen, L.L.** (2017). The Diversity of Long Noncoding RNAs and
Their Generation. Trends in Genetics **33,** 540-552.
**Xuan, H.D., Zhang, L.Z., Liu, X.S., Han, G.M., Li, J., Li, X., Liu, A.G., Liao, M.Z., and**
**Zhang, S.H.** (2015). PLNlncRbase: A resource for experimentally identified
lncRNAs in plants. Gene **573,** 328-332.
**Yamada, M.** (2017). Functions of long intergenic non-coding (linc) RNAs in plants. J
Plant Res **130,** 67-73.
**Yanai, I., Benjamin, H., Shmoish, M., Chalifa-Caspi, V., Shklar, M., Ophir, R., Bar-**
**Even, A., Horn-Saban, S., Safran, M., Domany, E., Lancet, D., and Shmueli,**
**O.** (2005). Genome-wide midrange transcription profiles reveal expression level
relationships in human tissue specification. Bioinformatics **21,** 650-659.
**Yu, C.P., Lin, J.J., and Li, W.H.** (2016). Positional distribution of transcription factor
binding sites in Arabidopsis thaliana. Sci Rep **6,** 25164.
**Zhang, L., Chen, F., Zhang, X., Li, Z., Zhao, Y., Lohaus, R., Chang, X., Dong, W., Ho,**
**S.Y.W., Liu, X., Song, A., Chen, J., Guo, W., Wang, Z., Zhuang, Y., Wang, H.,**
**Chen, X., Hu, J., Liu, Y., Qin, Y., Wang, K., Dong, S., Liu, Y., Zhang, S., Yu, X.,**
**Wu, Q., Wang, L., Yan, X., Jiao, Y., Kong, H., Zhou, X., Yu, C., Chen, Y., Li, F.,**
**Wang, J., Chen, W., Chen, X., Jia, Q., Zhang, C., Jiang, Y., Zhang, W., Liu, G.,**
**Fu, J., Chen, F., Ma, H., Van de Peer, Y., and Tang, H.** (2020). The water lily
genome and the early evolution of flowering plants. Nature **577,** 79-84.
**Zhao, Q., Li, M., Jia, Z., Liu, F., Ma, H., Huang, Y., and Song, S.** (2016). AtMYB44
Positively Regulates the Enhanced Elongation of Primary Roots Induced by N-3-
Oxo-Hexanoyl-Homoserine Lactone in Arabidopsis thaliana. Mol Plant Microbe
Interact **29,** 774-785.
**Zhao, X.Y., Li, J.R., Lian, B., Gu, H.Q., Li, Y., and Qi, Y.J.** (2018). Global identification
of Arabidopsis lncRNAs reveals the regulation of MAF4 by a natural antisense
RNA. Nat Commun **9**.
1328

**Figure 1. Overlap and gene features of Arabidopsis lincRNA annotations. (A)** Upset plot showing the intersection of lincRNA annotation in the eight resources. Each row represents a resource, reporting in parenthesis the total number of lincRNA transcripts before merging. LincRNA annotations unique to a single resource are represented as a single circle while circles connected by lines represent the intersection of lincRNA loci shared between various resources. The bar chart indicates the number of unique lincRNA loci and intersectional lincRNA loci, displaying only intersections that contain at least ten lincRNA loci. More complex overlap patterns are not shown to maintain figure clarity. **(B)** The pie chart shows the proportion of lincRNA loci supported by one or more resources. **(C)** The distribution of exon number for all lincRNA transcripts (purple), protein-coding transcripts (green), transcripts of lincRNAs supported by single resource (purple) and multiple resources (purple). Single exon and multiple exons are shown in dark and light colors, respectively. **(D)** The distribution of transcript length for lincRNAs (purple) and protein-coding genes (green).

**Figure 2. Sequence conservation analysis for lincRNAs of different evolutionary age categories. (A)** Simplified species tree reporting the number of lincRNAs found for the different age categories, which are also indicated by the grey circle sizes. Numbers in parenthesis report the number of genomes included per clade to assess sequence similarity and define a lincRNA's age category. **(B)** Sequence conservation (measured by the average phastCons score over the gene body) for different lincRNA age categories and gene types (lincRNAs, pri-miRNAs, and protein-coding genes). PhastCons score ranges from 0 (not under selection) to 1 (strong negative selection). P-values for pairwise Mann–Whitney U test are shown using the horizontal lines connecting the series. **(C)** Boxplot showing the average phastCons score for promoter regions (2kb upstream of transcription start site) of different lincRNA age categories and gene types (lincRNAs, pri-miRNAs, protein-coding genes). P-values for pairwise Mann–Whitney U test are shown using the horizontal lines connecting the series.

**Figure 3. Expression analysis of lincRNAs. (A)** Line chart showing the distribution of expression breadth for lincRNAs, pri-miRNAs and protein-coding genes across all samples **(B)** Distribution of the maximum TPM expression levels for different lincRNA age categories and gene types (lincRNAs, pri-miRNAs and protein-coding genes). **(C)** Distribution of tissue specificity tau scores for different lincRNA age categories and gene types. Tau scores range from zero to one, where zero means widely expressed, and one means very specifically expressed (detectable in only one cluster). The black dotted line represents a tau score of 0.97. **(D)** Bar chart reporting the number of expressed and highly-specific expressed lincRNAs for the different expression clusters. Cluster numbers and descriptions are shown below the chart, with numbers in parenthesis indicating the number of samples present per cluster. **(E)** Heatmap showing the proportion of highly-specific expressed lincRNAs in each cluster for each lincRNA age category. Cluster descriptions are the same as in panel D. Numbers in parenthesis report the number of genes per age category together with the fraction of lincRNAs showing highly-specific expression.

**Figure 4. Chromatin state and TF ChIP-Seq peak annotation for different gene types. (A)** Dendrogram showing the enrichment for different lincRNA gene sets (x-axis) towards different chromatin states (CS) (y-axis). The values report the product of -log10(q-value) and the enrichment fold. Only significant enrichment values are reported (q-value < 0.05). **(B)** The proportion of peak-gene pairs present in single replicate (blue) and two or more ChIP-Seq replicates (grey). **(C)** The percentage of three gene types assigned to peaks in two or more ChIP-Seq replicates. **(D)** Distributions of the number of TF bindings for lincRNA evolutionary age categories and gene types (lincRNAs, pri-miRNAs and protein-coding genes).

**Figure 5. Overview of TF-lincRNA regulatory interactions in different expression clusters and age categories.** The dot sizes represent the number of the lincRNAs while the color represents the statistical significance. Cluster descriptions are the same as in Figure 3D. **(A)** Bubble chart showing the enrichment of TF binding for expressed lincRNAs in different expression clusters. **(B)** Bubble chart showing the enrichment of TF binding for highly-expressed lincRNAs in different expression clusters. **(C)** Bubble chart showing the enrichment of TF binding for expressed lincRNAs in different age categories.

**Figure 6. Experimental validation and characterization of TF-lincRNA regulatory interactions.** **(A)** Heat map showing log2 fold change (FC) of lincRNA relative expression levels in transcription factor (TF) overexpressing lines (35S::*MYB44*, 35S::*PIF4* and *KAN1-GR*) or TF knockout lines (*rga28* and *pifq* (*pif1/pif2/pif3/pif4* quadruple mutants)) vs. control wild-type lines in 14-day old roots. Expression values were determined by RT-qPCR. Asterisks indicate statistically significant differences (*p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001) in an unpaired two-tailed Student's t-Test (n =3). Solid boxes indicate TF ChIP peaks <2kb from the lincRNA gene. Dashed boxes indicate TF ChIP peaks were identified only in one ChIP-Seq replicate or the lincRNA is not the closest gene to the TF peak. **(B)** Heatmap showing KAN1 ChIP-Seq peak-based target genes in the cluster 19 root expression atlas. The first line shows the expression profile of the TF under investigation while the other lines show groups of target genes with similar expression. Next to the group number, the number of target genes is reported and in parenthesis the fraction of lincRNAs per group is indicated. The numbers in squares and circles report the number of GWAS root-related and RT-qPCR experimentally validated lincRNA genes, respectively. **(C)** Heatmap showing MYB44 ChIP-Seq peak-based target genes in the cluster 19 root expression atlas. Group information is the same as in panel B. **(D)** Heatmap showing PIF4 ChIP-Seq peak-based target genes in the cluster 19 root expression atlas. Group information is the same as in panel B, but here triangles and circles report the RT-qPCR experimentally validated lincRNA genes for the pifq and 35S::*PIF4* lines, respectively.

## Parsed Citations

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic local alignment search tool. J Mol Biol 215, 403-410.
Google Scholar: Author Only Title Only Author and Title

Ariel, F., Lucero, L., Christ, A, Mammarella, M.F., Jegu, T., Veluchamy, A, Mariappan, K., Latrasse, D., Blein, T., Liu, C., Benhamed, M., and Crespi, M. (2020). R-Loop Mediated trans Action of the APOLO Long Noncoding RNA. Mol Cell 77, 1055-1065 e1054.
Google Scholar: Author Only Title Only Author and Title

Bardou, F., Ariel, F., Simpson, C.G., Romero-Barrios, N., Laporte, P., Balzergue, S., Brown, J.W., and Crespi, M. (2014). Long noncoding RNA modulates alternative splicing regulators in Arabidopsis. Dev Cell 30, 166-176.
Google Scholar: Author Only Title Only Author and Title

Bazin, J., Romero, N., Rigo, R., Charon, C., Blein, T., Ariel, F., and Crespi, M. (2018). Nuclear Speckle RNA Binding Proteins Remodel Alternative Splicing and the Non-coding Arabidopsis Transcriptome to Regulate a Cross-Talk Between Auxin and Immune Responses. Frontiers in Plant Science 9.
Google Scholar: Author Only Title Only Author and Title

Beilstein, M.A, Nagalingum, N.S., Clements, M.D., Manchester, S.R., and Mathews, S. (2010). Dated molecular phylogenies indicate a Miocene origin for Arabidopsis thaliana. Proc Natl Acad Sci U S A 107, 18724-18728.
Google Scholar: Author Only Title Only Author and Title

Bhogireddy, S., Mangrauthia, S.K., Kumar, R., Pandey, A.K., Singh, S., Jain, A, Budak, H., Varshney, R.K., and Kudapa, H. (2021). Regulatory non-coding RNAs: a new frontier in regulation of plant biology. Functional & Integrative Genomics 21, 313-330.
Google Scholar: Author Only Title Only Author and Title

Blein, T., Balzergue, C., Roule, T., Gabriel, M., Scalisi, L., Francois, T., Sorin, C., Christ, A, Godon, C., Delannoy, E., Martin-Magniette, M.L., Nussaume, L., Hartmann, C., Gautheret, D., Desnos, T., and Crespi, M. (2020). Landscape of the Noncoding Transcriptome Response of Two Arabidopsis Ecotypes to Phosphate Starvation. Plant Physiol 183, 1058-1072.
Google Scholar: Author Only Title Only Author and Title

Bu, D.C., Luo, H.T., Jiao, F., Fang, S.S., Tan, C.F., Liu, Z.Y., and Zhao, Y. (2015). Evolutionary annotation of conserved long non-coding RNAs in major mammalian species. Science China-Life Sciences 58, 787-798.
Google Scholar: Author Only Title Only Author and Title

Cabili, M.N., Trapnell, C., Goff, L., Koziol, M., Tazon-Vega, B., Regev, A, and Rinn, J.L. (2011). Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. Genes Dev 25, 1915-1927.
Google Scholar: Author Only Title Only Author and Title

Chen, L., Zhu, Q.H., and Kaufmann, K. (2020). Long non-coding RNAs in plants: emerging modulators of gene activity in development and stress responses. Planta 252, 92.
Google Scholar: Author Only Title Only Author and Title

Chen, Q.S., Liu, K., Yu, R., Zhou, B.L., Huang, P.P., Cao, Z.X., Zhou, Y.Q., and Wang, J.H. (2021). From "Dark Matter" to "Star": Insight Into the Regulation Mechanisms of Plant Functional Long Non-Coding RNAs. Frontiers in Plant Science 12.
Google Scholar: Author Only Title Only Author and Title

Cheng, C.Y., Krishnakumar, V., Chan, A.P., Thibaud-Nissen, F., Schobel, S., and Town, C.D. (2017). Araport11: a complete reannotation of the Arabidopsis thaliana reference genome. Plant J 89, 789-804.
Google Scholar: Author Only Title Only Author and Title

Chow, C.N., Lee, T.Y., Hung, Y.C., Li, G.Z, Tseng, K.C., Liu, Y.H., Kuo, P.L., Zheng, H.Q., and Chang, W.C. (2019). PlantPAN3.0: a new and updated resource for reconstructing transcriptional regulatory networks from ChIP-seq experiments in plants. Nucleic Acids Research 47, D1155-D1163.
Google Scholar: Author Only Title Only Author and Title

Deng, F., Zhang, X., Wang, W., Yuan, R., and Shen, F. (2018). Identification of Gossypium hirsutum long non-coding RNAs (lncRNAs) under salt stress. BMC Plant Biol 18, 23.
Google Scholar: Author Only Title Only Author and Title

Devaiah, B.N., Nagarajan, V.K., and Raghothama, K.G. (2007). Phosphate homeostasis and root development in Arabidopsis are synchronized by the zinc finger transcription factor ZAT6. Plant Physiol 145, 147-159.
Google Scholar: Author Only Title Only Author and Title

Fang, S.S., Zhang, L.L., Guo, J.C., Niu, Y.W., Wu, Y., Li, H., Zhao, L.H., Li, X.Y., Teng, X.Y., Sun, X.H., Sun, L., Zhang, M.Q., Chen, R.S., and Zhao, Y. (2018). NONCODEV5: a comprehensive annotation database for long non-coding RNAs. Nucleic Acids Research 46, D308-D314.
Google Scholar: Author Only Title Only Author and Title

Finn, R.D., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Mistry, J., Mitchell, A.L., Potter, S.C., Punta, M., Qureshi, M., Sangrador-Vegas,

A, Salazar, G.A, Tate, J., and Bateman, A. (2016). The Pfam protein families database: towards a more sustainable future. Nucleic Acids Res 44, D279-285.
Google Scholar: Author Only Title Only Author and Title

Fukuda, M., Nishida, S., Kakei, Y., Shimada, Y., and Fujiwara, T. (2019). Genome-Wide Analysis of Long Intergenic Noncoding RNAs Responding to Low-Nutrient Conditions in Arabidopsis thaliana: Possible Involvement of Trans-Acting siRNA3 in Response to Low Nitrogen. Plant Cell Physiol 60, 1961-1973.
Google Scholar: Author Only Title Only Author and Title

Haudry, A, Platts, A.E., Vello, E., Hoen, D.R., Leclercq, M., Williamson, R.J., Forczek, E., Joly-Lopez, Z, Steffen, J.G., Hazzouri, K.M., Dewar, K., Stinchcombe, J.R., Schoen, D.J., Wang, X., Schmutz, J., Town, C.D., Edger, P.P., Pires, J.C., Schumaker, K.S., Jarvis, D.E., Mandakova, T., Lysak, M.A, van den Bergh, E., Schranz, M.E., Harrison, P.M., Moses, A.M., Bureau, T.E., Wright, S.I., and Blanchette, M. (2013). An atlas of over 90,000 conserved noncoding sequences provides insight into crucifer regulatory regions. Nat Genet 45, 891-898.
Google Scholar: Author Only Title Only Author and Title

Hawker, N.P., and Bowman, J.L. (2004). Roles for Class III HD-Zip and KANADI genes in Arabidopsis root development. Plant Physiol 135, 2261-2270.
Google Scholar: Author Only Title Only Author and Title

Hazarika, R.R., Serra, M., Zhang, Z, Zhang, Y., Schmitz, R.J., and Johannes, F. (2022). Molecular properties of epimutation hotspots. Nat Plants 8, 146-156.
Google Scholar: Author Only Title Only Author and Title

He, H., Zhou, Y.F., Yang, Y.W., Zhang, Z, Lei, M.Q., Feng, Y.Z., Zhang, Y.C., Chen, Y.Q., Lian, J.P., and Yu, Y. (2021). Genome-Wide Analysis Identified a Set of Conserved lncRNAs Associated with Domestication-Related Traits in Rice. International Journal of Molecular Sciences 22.
Google Scholar: Author Only Title Only Author and Title

Heo, J.B., and Sung, S. (2011). Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. Science 331, 76-79.
Google Scholar: Author Only Title Only Author and Title

Heyndrickx, K.S., Van de Velde, J., Wang, C., Weigel, D., and Vandepoele, K. (2014). A functional and evolutionary perspective on transcription factor binding in Arabidopsis thaliana. Plant Cell 26, 3894-3910.
Google Scholar: Author Only Title Only Author and Title

Hu, L., Xu, Z, Hu, B., and Lu, Z.J. (2017). COME: a robust coding potential calculation tool for lncRNA identification and characterization based on multiple features. Nucleic Acids Res 45, e2.
Google Scholar: Author Only Title Only Author and Title

Huang, Y., Sicar, S., Ramirez-Prado, J.S., Manza-Mianza, D., Antunez-Sanchez, J., Brik-Chaouche, R., Rodriguez-Granados, N.Y., An, J., Bergounioux, C., Mahfouz, M.M., Hirt, H., Crespi, M., Concia, L., Barneche, F., Amiard, S., Probst, A.V., Gutierrez-Marcos, J., Ariel, F., Raynaud, C., Latrasse, D., and Benhamed, M. (2021). Polycomb-dependent differential chromatin compartmentalization determines gene coregulation in Arabidopsis. Genome Research 31, 1230-+.
Google Scholar: Author Only Title Only Author and Title

Hupalo, D., and Kern, A.D. (2013). Conservation and functional element discovery in 20 angiosperm plant genomes. Mol Biol Evol 30, 1729-1744.
Google Scholar: Author Only Title Only Author and Title

Jha, U.C., Nayyar, H., Jha, R., Khurshid, M., Zhou, M., Mantri, N., and Siddique, K.H.M. (2020). Long non-coding RNAs: emerging players regulating plant abiotic stress response and adaptation. BMC Plant Biol 20, 466.
Google Scholar: Author Only Title Only Author and Title

Jones-Rhoades, M.W., Bartel, D.P., and Bartel, B. (2006). MicroRNAs and their regulatory roles in plants. Annual Review of Plant Biology 57, 19-53.
Google Scholar: Author Only Title Only Author and Title

Kalvari, I., Nawrocki, E.P., Ontiveros-Palacios, N., Argasinska, J., Lamkiewicz, K., Marz, M., Griffiths-Jones, S., Toffano-Nioche, C., Gautheret, D., Weinberg, Z, Rivas, E., Eddy, S.R., Finn, R.D., Bateman, A, and Petrov, A.I. (2021). Rfam 14: expanded coverage of metagenomic, viral and microRNA families. Nucleic Acids Res 49, D192-D200.
Google Scholar: Author Only Title Only Author and Title

Kang, Y.J., Yang, D.C., Kong, L., Hou, M., Meng, Y.Q., Wei, L., and Gao, G. (2017). CPC2: a fast and accurate coding potential calculator based on sequence intrinsic features. Nucleic Acids Res 45, W12-W16.
Google Scholar: Author Only Title Only Author and Title

Ke, L.L., Zhou, Z.W., Xu, X.W., Wang, X., Liu, Y.L., Xu, Y.T., Huang, Y., Wang, S.T., Deng, X.X., Chen, L.L., and Xu, Q. (2019). Evolutionary dynamics of lincRNA transcription in nine citrus species. Plant Journal 98, 912-927.
Google Scholar: Author Only Title Only Author and Title

Kim, D.H., and Sung, S. (2017). Vernalization-Triggered Intragenic Chromatin Loop Formation by Long Noncoding RNAs. Dev Cell 40, 302-312 e304.
  Google Scholar: Author Only Title Only Author and Title

Kindgren, P., Ard, R., Ivanov, M., and Marquardt, S. (2018). Transcriptional read-through of the long non-coding RNA SVALKA governs plant cold acclimation. Nat Commun 9, 4561.
  Google Scholar: Author Only Title Only Author and Title

Kramer, M.C., Kim, H.J., Palos, K.R., Garcia, B.A, Lyons, E., Beilstein, M.A, Nelson, A.D.L., and Gregory, B.D. (2022). A Conserved Long Intergenic Non-coding RNA Containing snoRNA Sequences, lncCOBRA1, Affects Arabidopsis Germination and Development. Front Plant Sci 13, 906603.
  Google Scholar: Author Only Title Only Author and Title

Li, Q.Q., Zhang, Z., Zhang, C.X., Wang, Y.L., Liu, C.B., Wu, J.C., Han, M.L., Wang, Q.X., and Chao, D.Y. (2022). Phytochrome-interacting factors orchestrate hypocotyl adventitious root initiation in Arabidopsis. Development 149.
  Google Scholar: Author Only Title Only Author and Title

Li, S., Yamada, M., Han, X., Ohler, U., and Benfey, P.N. (2016). High-Resolution Expression Map of the Arabidopsis Root Reveals Alternative Splicing and lincRNA Regulation. Dev Cell 39, 508-522.
  Google Scholar: Author Only Title Only Author and Title

Li, S.X., Yu, X., Lei, N., Cheng, Z.H., Zhao, P.J., He, Y.K., Wang, W.Q., and Peng, M. (2017). Genome-wide identification and functional prediction of cold and/or drought-responsive lncRNAs in cassava. Scientific Reports 7.
  Google Scholar: Author Only Title Only Author and Title

Liu, F., Marquardt, S., Lister, C., Swiezewski, S., and Dean, C. (2010). Targeted 3' processing of antisense transcripts triggers Arabidopsis FLC chromatin silencing. Science 327, 94-97.
  Google Scholar: Author Only Title Only Author and Title

Liu, J., Wang, H., and Chua, N.H. (2015). Long noncoding RNA transcriptome of plants. Plant Biotechnol J 13, 319-328.
  Google Scholar: Author Only Title Only Author and Title

Liu, J., Jung, C., Xu, J., Wang, H., Deng, S.L., Bernad, L., Arenas-Huertero, C., and Chua, N.H. (2012). Genome-Wide Analysis Uncovers Regulation of Long Intergenic Noncoding RNAs in Arabidopsis. Plant Cell 24, 4333-4345.
  Google Scholar: Author Only Title Only Author and Title

Liu, Y., Tian, T., Zhang, K., You, Q., Yan, H., Zhao, N., Yi, X., Xu, W., and Su, Z. (2018). PCSD: a plant chromatin state database. Nucleic Acids Res 46, D1157-D1167.
  Google Scholar: Author Only Title Only Author and Title

Lucero, L., Ferrero, L., Fonouni-Farde, C., and Ariel, F. (2021). Functional classification of plant long noncoding RNAs: a transcript is known by the company it keeps. New Phytologist 229, 1251-1260.
  Google Scholar: Author Only Title Only Author and Title

Ma, J.C., Bai, X.T., Luo, W.C., Feng, Y.N., Shao, X.M., Bai, Q.X., Sun, S.J., Long, Q.M., and Wan, D.S. (2019). Genome-Wide Identification of Long Noncoding RNAs and Their Responses to Salt Stress in Two Closely Related Poplars. Frontiers in Genetics 10.
  Google Scholar: Author Only Title Only Author and Title

Mattioli, K., Volders, P.J., Gerhardinger, C., Lee, J.C., Maass, P.G., Mele, M., and Rinn, J.L. (2019). High-throughput functional analysis of lncRNA core promoters elucidates rules governing tissue specificity. Genome Research 29, 344-355.
  Google Scholar: Author Only Title Only Author and Title

Merelo, P., Xie, Y.K., Brand, L., Ott, F., Weigel, D., Bowman, J.L., Heisler, M.G., and Wenkel, S. (2013). Genome-Wide Identification of KANADI1 Target Genes. Plos One 8.
  Google Scholar: Author Only Title Only Author and Title

Mohammadin, S., Edger, P.P., Pires, J.C., and Schranz, M.E. (2015). Positionally-conserved but sequence-diverged: identification of long non-coding RNAs in the Brassicaceae and Cleomaceae. BMC Plant Biol 15, 217.
  Google Scholar: Author Only Title Only Author and Title

Moison, M., Pacheco, J.M., Lucero, L., Fonouni-Farde, C., Rodriguez-Melo, J., Mansilla, N., Christ, A, Bazin, J., Benhamed, M., Ibanez, F., Crespi, M., Estevez, J.M., and Ariel, F. (2021). The lncRNA APOLO interacts with the transcription factor WRKY42 to trigger root hair cell expansion in response to cold. Mol Plant 14, 937-948.
  Google Scholar: Author Only Title Only Author and Title

Moubayidin, L., Salvi, E., Giustini, L., Terpstra, I., Heidstra, R., Costantino, P., and Sabatini, S. (2016). A SCARECROW-based regulatory circuit controls Arabidopsis thaliana meristem size from the root endodermis. Planta 243, 1159-1168.
  Google Scholar: Author Only Title Only Author and Title

Nawrocki, E.P., and Eddy, S.R. (2013). Infernal 1.1: 100-fold faster RNA homology searches. Bioinformatics 29, 2933-2935.
Google Scholar: Author Only Title Only Author and Title

Necsulea, A, Soumillon, M., Warnefors, M., Liechti, A, Daish, T., Zeller, U., Baker, J.C., Grutzner, F., and Kaessmann, H. (2014). The evolution of lncRNA repertoires and expression patterns in tetrapods. Nature 505, 635-+.
Google Scholar: Author Only Title Only Author and Title

Nelson, A.D.L., Devisetty, U.K., Palos, K., Haug-Baltzell, A.K., Lyons, E., and Beilstein, M.A (2017). Evolinc:A Tool for the Identification and Evolutionary Comparison of Long Intergenic Non-coding RNAs. Frontiers in Genetics 8.
Google Scholar: Author Only Title Only Author and Title

Nelson, A.D.L., Forsythe, E.S., Devisetty, U.K., Clausen, D.S., Haug-Batzell, A.K., Meldrum, A.M.R., Frank, M.R., Lyons, E., and Beilstein, M.A (2016). A Genomic Analysis of Factors Driving lincRNA Diversification: Lessons from Plants. G3-Genes Genomes Genetics 6, 2881-2891.
Google Scholar: Author Only Title Only Author and Title

O'Maoileidigh, D.S., Graciet, E., and Wellmer, F. (2014). Gene networks controlling Arabidopsis thaliana flower development. New Phytol 201, 16-30.
Google Scholar: Author Only Title Only Author and Title

Pacheco, J.M., Mansilla, N., Moison, M., Lucero, L., Gabarain, V.B., Ariel, F., and Estevez, J.M. (2021). The lncRNA APOLO and the transcription factor WRKY42 target common cell wall EXTENSIN encoding genes to trigger root hair cell elongation. Plant Signal Behav 16, 1920191.
Google Scholar: Author Only Title Only Author and Title

Palos, K., Dittrich, A.C.N., Yu, L.A, Brock, J.R., Railey, C.E., Wu, H.Y.L., Sokolowska, E., Skirycz, A, Hsu, P.Y., Gregory, B.D., Lyons, E., Beilstein, M.A., and Nelson, A.D.L. (2022). Identification and functional annotation of long intergenic non-coding RNAs in Brassicaceae. Plant Cell.
Google Scholar: Author Only Title Only Author and Title

Pedregosa, F., Varoquaux, G., Gramfort, A, Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A, Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E. (2011). Scikit-learn: Machine Learning in Python. Journal of Machine Learning Research 12, 2825-2830.
Google Scholar: Author Only Title Only Author and Title

Pertea, G., and Pertea, M. (2020). GFF Utilities: GffRead and GffCompare. F1000Res 9.
Google Scholar: Author Only Title Only Author and Title

Pfeiffer, A, Shi, H., Tepperman, J.M., Zhang, Y., and Quail, P.H. (2014). Combinatorial Complexity in a Transcriptionally Centered Signaling Hub in Arabidopsis. Molecular Plant 7, 1598-1618.
Google Scholar: Author Only Title Only Author and Title

Ponting, C.P., and Haerty, W. (2022). Genome-Wide Analysis of Human Long Noncoding RNAs: A Provocative Review. Annu Rev Genomics Hum Genet.
Google Scholar: Author Only Title Only Author and Title

Qi, X., Xie, S., Liu, Y., Yi, F., and Yu, J. (2013). Genome-wide annotation of genes and noncoding RNAs of foxtail millet in response to simulated drought stress by deep sequencing. Plant Mol Biol 83, 459-473.
Google Scholar: Author Only Title Only Author and Title

Qin, T., Zhao, H., Cui, P., Albesher, N., and Xiong, L. (2017). A Nucleus-Localized Long Non-Coding RNA Enhances Drought and Salt Stress Tolerance. Plant Physiol 175, 1321-1336.
Google Scholar: Author Only Title Only Author and Title

Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 26, 841-842.
Google Scholar: Author Only Title Only Author and Title

Rai, M.I., Alam, M., Lightfoot, D.A, Gurha, P., and Afzal, A.J. (2019). Classification and experimental identification of plant long non-coding RNAs. Genomics 111, 997-1005.
Google Scholar: Author Only Title Only Author and Title

Ramskold, D., Wang, E.T., Burge, C.B., and Sandberg, R. (2009). An abundance of ubiquitously expressed genes revealed by tissue transcriptome sequence data. PLoS Comput Biol 5, e1000598.
Google Scholar: Author Only Title Only Author and Title

Ransbotyn, V., Yeger-Lotem, E., Basha, O., Acuna, T., Verduyn, C., Gordon, M., Chalifa-Caspi, V., Hannah, M.A., and Barak, S. (2015). A combination of gene expression ranking and co-expression network analysis increases discovery rate in large-scale mutant screens for novel Arabidopsis thaliana abiotic stress genes. Plant Biotechnol J 13, 501-513.
Google Scholar: Author Only Title Only Author and Title

Ransohoff, J.D., Wei, Y., and Khavari, P.A. (2018). The functions and unique features of long intergenic non-coding RNA. Nat Rev Mol Cell Biol 19, 143-157.
  Google Scholar: Author Only Title Only Author and Title

Rigo, R., Bazin, J., Romero-Barrios, N., Moison, M., Lucero, L., Christ, A., Benhamed, M., Blein, T., Huguet, S., Charon, C., Crespi, M., and Ariel, F. (2020). The Arabidopsis lncRNA ASCO modulates the transcriptome through interaction with splicing factors. EMBO Rep 21, e48977.
  Google Scholar: Author Only Title Only Author and Title

Roule, T., Crespi, M., and Blein, T. (2022a). Regulatory long non-coding RNAs in root growth and development. Biochem Soc Trans 50, 403-412.
  Google Scholar: Author Only Title Only Author and Title

Roule, T., Christ, A., Hussain, N., Huang, Y., Hartmann, C., Benhamed, M., Gutierrez-Marcos, J., Ariel, F., Crespi, M., and Blein, T. (2022b). The lncRNA MARS modulates the epigenetic reprogramming of the marneral cluster in response to ABA. Mol Plant 15, 840-856.
  Google Scholar: Author Only Title Only Author and Title

Sanchita, Trivedi, P.K., and Asif, M.H. (2020). Updates on plant long non-coding RNAs (lncRNAs): the regulatory components. Plant Cell Tissue and Organ Culture 140, 259-269.
  Google Scholar: Author Only Title Only Author and Title

Sarropoulos, I., Marin, R., Cardoso-Moreira, M., and Kaessmann, H. (2019). Developmental dynamics of lncRNAs across mammalian organs and species. Nature 571, 510-+.
  Google Scholar: Author Only Title Only Author and Title

Schmitz, R.J., Grotewold, E., and Stam, M. (2022). Cis-regulatory sequences in plants: Their importance, discovery, and future challenges. Plant Cell 34, 718-741.
  Google Scholar: Author Only Title Only Author and Title

Severing, E., Faino, L., Jamge, S., Busscher, M., Kuijer-Zhang, Y., Bellinazzo, F., Busscher-Lange, J., Fernandez, V., Angenent, G.C., Immink, R.G.H., and Pajoro, A. (2018). Arabidopsis thaliana ambient temperature responsive lncRNAs. BMC Plant Biol 18, 145.
  Google Scholar: Author Only Title Only Author and Title

Shafiq, S., Li, J.R., and Sun, Q.W. (2016). Functions of plants long non-coding RNAs. Biochimica Et Biophysica Acta-Gene Regulatory Mechanisms 1859, 155-162.
  Google Scholar: Author Only Title Only Author and Title

Shea, D.J., Nishida, N., Takada, S., Itabashi, E., Takahashi, S., Akter, A., Miyaji, N., Osabe, K., Mehraj, H., Shimizu, M., Seki, M., Kakizaki, T., Okazaki, K., Dennis, E.S., and Fujimoto, R. (2019). Long noncoding RNAs in Brassica rapa L. following vernalization. Sci Rep 9, 9302.
  Google Scholar: Author Only Title Only Author and Title

Shuai, P., Liang, D., Tang, S., Zhang, Z.J., Ye, C.Y., Su, Y.Y., Xia, X.L., and Yin, W.L. (2014). Genome-wide identification and functional prediction of novel and drought-responsive lincRNAs in Populus trichocarpa. J Exp Bot 65, 4975-4983.
  Google Scholar: Author Only Title Only Author and Title

Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K., Clawson, H., Spieth, J., Hillier, L.W., Richards, S., Weinstock, G.M., Wilson, R.K., Gibbs, R.A., Kent, W.J., Miller, W., and Haussler, D. (2005). Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Res 15, 1034-1050.
  Google Scholar: Author Only Title Only Author and Title

Song, L., Huang, S.S.C., Wise, A., Castanon, R., Nery, J.R., Chen, H.M., Watanabe, M., Thomas, J., Bar-Joseph, Z., and Ecker, J.R. (2016). A transcription factor hierarchy defines an environmental stress response network. Science 354.
  Google Scholar: Author Only Title Only Author and Title

Sun, L., Luo, H., Bu, D., Zhao, G., Yu, K., Zhang, C., Liu, Y., Chen, R., and Zhao, Y. (2013). Utilizing sequence intrinsic composition to classify protein-coding and long non-coding transcripts. Nucleic Acids Research 41.
  Google Scholar: Author Only Title Only Author and Title

Sun, Z., Nair, A., Chen, X., Prodduturi, N., Wang, J., and Kocher, J.P. (2017). UClncR: Ultrafast and comprehensive long non-coding RNA detection from RNA-seq. Sci Rep 7, 14196.
  Google Scholar: Author Only Title Only Author and Title

Szczesniak, M.W., Rosikiewicz, W., and Makalowska, I. (2016). CANTATAdb: A Collection of Plant Long Non-Coding RNAs. Plant Cell Physiol 57, e8.
  Google Scholar: Author Only Title Only Author and Title

Szczesniak, M.W., Kubiak, M.R., Wanowska, E., and Makalowska, I. (2021). Comparative genomics in the search for conserved

long noncoding RNAs. Essays Biochem 65, 741-749.
Google Scholar: Author Only Title Only Author and Title

Togninalli, M., Seren, U., Freudenthal, J.A, Monroe, J.G., Meng, D., Nordborg, M., Weigel, D., Borgwardt, K., Korte, A, and Grimm, D.G. (2020). AraPheno and the AraGWAS Catalog 2020: a major database update including RNA-Seq and knockout mutation data for Arabidopsis thaliana. Nucleic Acids Res 48, D1063-D1068.
Google Scholar: Author Only Title Only Author and Title

Tominaga-Wada, R., and Wada, T. (2016). The ectopic localization of CAPRICE LIKE MYB3 protein in Arabidopsis root epidermis. J Plant Physiol 199, 111-115.
Google Scholar: Author Only Title Only Author and Title

Tran Vdu, T., Souiai, O., Romero-Barrios, N., Crespi, M., and Gautheret, D. (2016). Detection of generic differential RNA processing events from RNA-seq data. RNA Biol 13, 59-67.
Google Scholar: Author Only Title Only Author and Title

Ulitsky, I. (2016). Evolution to the rescue: using comparative genomics to understand long non-coding RNAs. Nature Reviews Genetics 17, 601-614.
Google Scholar: Author Only Title Only Author and Title

Van Bel, M., Silvestri, F., Weitz, E.M., Kreft, L., Botzki, A, Coppens, F., and Vandepoele, K. (2021). PLAZA 5.0: extending the scope and power of comparative and functional genomics in plants. Nucleic Acids Res.
Google Scholar: Author Only Title Only Author and Title

Van de Velde, J., Van Bel, M., Vaneechoutte, D., and Vandepoele, K. (2016). A Collection of Conserved Noncoding Sequences to Study Gene Regulation in Flowering Plants. Plant Physiol 171, 2586-2598.
Google Scholar: Author Only Title Only Author and Title

Vandepoele, K., Quimbaya, M., Casneuf, T., De Veylder, L., and Van de Peer, Y. (2009). Unraveling Transcriptional Control in Arabidopsis Using cis-Regulatory Elements and Coexpression Networks. Plant Physiology 150, 535-546.
Google Scholar: Author Only Title Only Author and Title

Vaneechoutte, D., and Vandepoele, K. (2019). Curse: building expression atlases and co-expression networks from public RNA-Seq data. Bioinformatics 35, 2880-2881.
Google Scholar: Author Only Title Only Author and Title

Wang, H., Niu, Q.W., Wu, H.W., Liu, J., Ye, J., Yu, N., and Chua, N.H. (2015a). Analysis of non-coding transcriptome in rice and maize uncovers roles of conserved lncRNAs associated with agriculture traits. Plant Journal 84, 404-416.
Google Scholar: Author Only Title Only Author and Title

Wang, H., Niu, Q.W., Wu, H.W., Liu, J., Ye, J., Yu, N., and Chua, N.H. (2015b). Analysis of non-coding transcriptome in rice and maize uncovers roles of conserved lncRNAs associated with agriculture traits. Plant J 84, 404-416.
Google Scholar: Author Only Title Only Author and Title

Wu, H., Yang, L., and Chen, L.L. (2017). The Diversity of Long Noncoding RNAs and Their Generation. Trends in Genetics 33, 540-552.
Google Scholar: Author Only Title Only Author and Title

Xuan, H.D., Zhang, L.Z, Liu, X.S., Han, G.M., Li, J., Li, X., Liu, A.G., Liao, M.Z., and Zhang, S.H. (2015). PLNlncRbase: A resource for experimentally identified lncRNAs in plants. Gene 573, 328-332.
Google Scholar: Author Only Title Only Author and Title

Yamada, M. (2017). Functions of long intergenic non-coding (linc) RNAs in plants. J Plant Res 130, 67-73.
Google Scholar: Author Only Title Only Author and Title

Yanai, I., Benjamin, H., Shmoish, M., Chalifa-Caspi, V., Shklar, M., Ophir, R., Bar-Even, A, Horn-Saban, S., Safran, M., Domany, E., Lancet, D., and Shmueli, O. (2005). Genome-wide midrange transcription profiles reveal expression level relationships in human tissue specification. Bioinformatics 21, 650-659.
Google Scholar: Author Only Title Only Author and Title

Yu, C.P., Lin, J.J., and Li, W.H. (2016). Positional distribution of transcription factor binding sites in Arabidopsis thaliana. Sci Rep 6, 25164.
Google Scholar: Author Only Title Only Author and Title

Zhang, L., Chen, F., Zhang, X., Li, Z, Zhao, Y., Lohaus, R., Chang, X., Dong, W., Ho, S.Y.W., Liu, X., Song, A, Chen, J., Guo, W., Wang, Z, Zhuang, Y., Wang, H., Chen, X., Hu, J., Liu, Y., Qin, Y., Wang, K., Dong, S., Liu, Y., Zhang, S., Yu, X., Wu, Q., Wang, L., Yan, X., Jiao, Y., Kong, H., Zhou, X., Yu, C., Chen, Y., Li, F., Wang, J., Chen, W., Chen, X., Jia, Q., Zhang, C., Jiang, Y., Zhang, W., Liu, G., Fu, J., Chen, F., Ma, H., Van de Peer, Y., and Tang, H. (2020). The water lily genome and the early evolution of flowering plants. Nature 577, 79-84.
Google Scholar: Author Only Title Only Author and Title

**Zhao, Q., Li, M., Jia, Z, Liu, F., Ma, H., Huang, Y., and Song, S. (2016). AtMYB44 Positively Regulates the Enhanced Elongation of Primary Roots Induced by N-3-Oxo-Hexanoyl-Homoserine Lactone in Arabidopsis thaliana. Mol Plant Microbe Interact 29, 774-785.**
  Google Scholar: Author Only Title Only Author and Title

**Zhao, X.Y., Li, J.R., Lian, B., Gu, H.Q., Li, Y., and Qi, Y.J. (2018). Global identification of Arabidopsis lncRNAs reveals the regulation of MAF4 by a natural antisense RNA. Nat Commun 9.**
  Google Scholar: Author Only Title Only Author and Title