Using machine learning to predict protein-protein interactions between a zombie ant

fungus and its carpenter ant host

Ian Will [1]*, William C. Beckerson [1], Charissa de Bekker [1,2]*

[1] University of Central Florida, Department of Biology, 4110 Libra Drive, 32816, Orlando, FL,

USA

[2] Utrecht University, Department of Biology, Microbiology, Padualaan 8, 3584 CH, Utrecht, NL

* Corresponding authors: Ian Will (ian.will@knights.ucf.edu), Charissa de Bekker

(a.m.debekker@uu.nl)

ORCID: I.W. - 0000-0002-2396-6778, W.C.B. - 0000-0001-7132-7578, C.dB - 0000-0002-

7325-172X

Abstract

Parasitic fungi produce proteins that modulate virulence, alter host physiology, and

trigger host responses. These proteins, classified as a type of "effector," often act via protein-

protein interactions (PPIs). The fungal parasite *Ophiocordyceps camponoti-floridani* (zombie ant

fungus) manipulates *Camponotus floridanus* (carpenter ant) behavior to promote transmission.

The most striking aspect of this behavioral change is a summit disease phenotype where infected

hosts ascend and attach to an elevated position. Plausibly, interspecific PPIs drive aspects of

Ophiocordyceps infection and host manipulation. Machine learning PPI predictions offer high-

throughput methods to produce mechanistic hypotheses on how this behavioral manipulation

occurs. Using D-SCRIPT to predict host-parasite PPIs, we found ca. 6,000 interactions involving

2,083 host proteins and 129 parasite proteins, which are encoded by genes upregulated during

manipulated behavior. We identified multiple overrepresentations of functional annotations

among these proteins. The strongest signals in the host highlighted neuromodulatory G-protein

coupled receptors and oxidation-reduction processes. We also detected *Camponotus* structural

and gene-regulatory proteins. In the parasite, we found enrichment of *Ophiocordyceps* proteases

and frequent involvement of novel small secreted proteins with unknown functions. From these

results, we provide new hypotheses on potential parasite effectors and host targets underlying

zombie ant behavioral manipulation.

Introduction

Fungal parasites employ a diverse array of secreted molecules to defend themselves,

promote infection, and modify their hosts. In certain cases, infection can even lead to parasitic

manipulation of host behavior. Often termed "effectors," secreted fungal molecules play a

critical role in host-parasite dynamics that include both widely shared and highly specific

mechanisms [1–4]. Parasite effectors have been suggested to play key roles during the infection of

insects by entomopathogenic fungi through their interactions with host nucleic acids,

carbohydrates, lipids, small metabolites, and proteins [3,5–7]. Bioinformatic exploration of effector

protein biology offers a high-throughput method to develop mechanistic hypotheses of how some

parasites can modify the behavior of their hosts. Here, we use such approaches to investigate the

behavior-manipulating fungus *Ophiocordyceps camponoti-floridani* (Florida zombie ant fungus)

and its insect host *Camponotus floridanus* (Florida carpenter ant).

Myrmecophilous *Ophiocordyceps* fungi are typically species-specific parasites that have

co-evolved with their ant hosts over millions of years [8]. This close relationship has resulted in

fungus-ant interactions that alter host behaviors in ways that are adaptive for the parasite.

Manipulated host ants succumb to a summit disease, affixing themselves to elevated positions

and dying at locations that promote fungal growth and transmission [9–13]. Behavioral changes

preceding this final summit could include host-adaptive responses, parasite-adaptive

manipulations, or general symptoms of disease that span hyperactivity, uncoordinated foraging,

decreased nestmate communication, and convulsions [9,14–17]. While these modified ant behaviors

have been observed in nature for some time [18], explorations of the fungal effectors involved are

relatively more recent endeavors [14,19–25]. These multiple "-omics" studies have provided

hypotheses about the parasite and host molecules that play a role in establishing the behavioral

phenotypes observed, but leave their potential interactions open to interpretation.

A large number of fungal molecules and ant pathways may be involved in

Ophiocordyceps behavioral manipulation, including neuro-modulators and -protectants, insect

hormones, feeding, locomotion, circadian rhythms and light-sensing, and muscular hyperactivity

[14,17,19,21,22,25–29]. Fungal proteins such as bacterial-like enterotoxins, protein tyrosine

phosphatases, peptidases (e.g., S8 subtilisin-like serine proteases), and undescribed small

secreted proteins (uSSPs) have been proposed as mechanisms of manipulation that presumably

function by PPIs. These hypotheses are supported by strong gene upregulation of these

candidates during active manipulation of the ant and genomic comparisons among the

Ophiocordyceps [14,19,24]. Numerous other such proteins are supported as plausible effectors as

well. Testing them *in vivo* will be a costly and laborious process in an emerging, non-traditional

model organism such as *O. camponoti-floridani*. New evidence linking previously hypothesized

candidate proteins and pathways to host-parasite protein-protein interactions (PPIs) would give

strong support to select top predictions for functional validation.

Leveraging big-data and computational methods to explore host-parasite biology is a

continuing and multifaceted effort. Yet, examples of using these assets in PPI prediction remain

limited [30,31]. Proteome-scale bioinformatic prediction of PPIs has largely been employed to

describe protein interaction networks within a single organism, as was done for the host species,

C. floridanus [32]. However, PPI predictions in some interspecific host-parasite relationships have

been made, including animal or plant hosts and bacterial, viral, or fungal parasites [33–36].

Predicting PPIs is typically done using either an interolog or machine learning method.

Interologs are orthologous interactions conserved across species; each protein in one PPI has

orthologs in the second PPI [37]. While a useful prediction tool, detecting an interolog in a new

system requires established reference PPIs from other systems. This constraint by available data

would result in a very limited number of predictions when analyzing interactions between non-

model organisms. Machine-learning methods offer a more flexible alternative. In particular,

neural network deep learning tools for PPI prediction have become increasingly available in

recent years and continue to be an active field of development [38–41]. Here, predicted PPIs are still

informed by a known set of PPIs used for training data. However, once trained, the program is

able to predict novel interactions outside the original dataset [35,42]. The deep learning tool D-

SCRIPT has shown an uncommon flexibility in cross-species PPI predictions when trained on

one organism but tested on a different species [35]. Protein embeddings that encode putative

structural information are used to train D-SCRIPT models, which have produced predicted

protein-protein contact maps that largely agree with experimentally measured docked proteins

[35,36,43]. Although other machine learning methods can outperform D-SCRIPT in same-species

contexts, the cross-species generalizability of D-SCRIPT's predictive power is an important step

for research on non-model or multi-species systems [35,38]. Notably, *Ophiocordyceps* and other

fungi secrete a range of taxonomically distinct uSSPs that are often thought to be effectors in

host-parasite interactions [14,19,24,44–46]. Therefore, bioinformatic techniques that hinge upon well

described protein annotations or are subject to strong species-specific biases can only have

limited success in assigning uSSPs to PPIs.

For our analyses between *O. camponoti-floridani* and *C. floridanus* proteins, we anchor

our reporting on group-level functional enrichments of predicted positive PPIs. We chose this

approach in consideration of the reported D-SCRIPT default model error rates on insect

(*Drosophila*) and fungal (*Saccharomyces*) species. In these tests, predicted positive PPIs

contained 71-79% true-positives (precision rate) and captured 22-36% of all true-positives (recall

rate) [35], which indicate this tool is strongest in broad characterizations rather than validating

specific, previously hypothesized PPIs. Similarly, we benchmarked D-SCRIPT on

experimentally supported fungus-animal interspecific PPIs and found comparable low true-

positive recall coupled with higher precision. The authors of D-SCRIPT have also suggested that

this tool can outperform an earlier deep learning method in reproducing gene ontology (GO)

term enrichments of experimentally identified virus-human PPIs [35].

We used D-SCRIPT to predict possible *O. camponoti-floridani* and *C. floridanus* PPIs

involved in behavioral manipulation. We filtered these predictions into subsets we considered

most likely to include relevant interactions. As such, our study focuses on PPIs involving

putatively secreted *O. camponoti-floridani* proteins encoded by genes that were previously found

to be upregulated during manipulated summiting [19]. These fungal proteins are plausibly exported

to the host environment during manipulation, providing support for hypotheses implicating such

proteins in manipulation of host behavior.

We highlight aspects of functional enrichments after filtering out *Camponotus* protein

interactions with fungi that do not naturally infect or manipulate this host ("aspecific" fungi).

These aspecific fungi cover different lifestyles and phylogenetic relationships to

Ophiocordyceps: (*i*) *Cordyceps bassiana* (i.e., *Beauveria bassiana*), a generalist entomopathogen

in the order Hypocreales (which includes *Ophiocordyceps*), (*ii*) *Trichoderma reesei*, a plant-

degrading saprophyte in the order Hypocreales (although the genus also includes

entomopathogens), and (*iii*) *Saccharomyces cerevisiae*, a phylogenetically distant saprophytic

yeast in the order Saccharomycetales with a smaller secretome. This approach brought the

unique *Ophiocordyceps* PPIs to the forefront but does not discount the aspecific predictions.

Rather, it brings additional attention to a narrower set of specific PPIs that could include co-

evolved adaptations underlying *Ophiocordyceps*-manipulated *Camponotus* behavior.

Materials and Methods

Fungus-animal PPI benchmarking: The Host-Pathogen Interaction Database (v 3.0)

provided experimentally supported PPIs between 5 fungal and 12 animal species. We used this

data to benchmark D-SCRIPT in predicting interspecific PPIs. The types of experimental support

we selected included direct interactions and reactions (e.g., "phosphorylation reaction") but

excluded associations, colocalizations, and genetic effects (e.g., "additive genetic interaction").

We only included PPIs with proteins $\leq$ 2,000 amino acids, for computational reasons. We further

limited similar PPIs to include only one representative interaction by clustering all proteins in the

dataset by 40% similarity using MMseqs2 (v. 9-d36de) (single step easy-clustering with

minimum sequence identity 0.4) [47]. For any PPIs that connected the same two homologous

protein clusters, we selected one of those PPIs at random and removed the others. In total, this

produced 567 experimentally verified fungus-animal PPIs.

In addition to these verified, positive fungus-animal PPIs, we constructed a set of

negative interactions to use in evaluating D-SCRIPT. Previously, D-SCRIPT evaluation datasets

have used randomly paired proteins from the tested organism as negative PPI examples [35]. In this

vein, we created artificial interspecific fungus-animal PPIs using random yeast (*S. cerevisiae*)

and fly (*Drosophila melanogaster*) proteins collected from the STRING database (v. 11.0) [48].

Rather than randomly pairing proteins from all organisms in the fungus-animal dataset, we chose

these two organisms because they both: (*i*) were represented in the positive PPIs, (*ii*) had robust

PPI data available for removing known positive interactions from the random pairings, (*iii*) were

used in the original D-SCRIPT evaluations, and (*iv*) match *Ophiocordyceps* and *Camponotus* as

a fungus-insect pair [35]. Using the aforementioned 40% clustering method, we filtered out any

random yeast-fly PPIs that were similar to positive fungus-animal interactions. We also removed

any that were similar to known, high-confidence, experimentally-supported fly or yeast PPIs

listed in the STRING database ("experiments score" > 0 and "combined score" ≥ 700) [48]. Then,

we kept only one representative of PPIs that were similar to each other. After filtering, we

selected 5,670 random PPIs (10-fold the number of positive cases) to use as negative examples in

the evaluation dataset.

We used D-SCRIPT (v 0.1.5) with the default human-protein pretrained model and

settings in evaluation mode with this fungus-animal dataset (567 positive and 5,670 negative

PPIs) [35]. D-SCRIPT assigns each tested protein interaction an edge value (i.e., a confidence score

to rank possible PPIs) with a default threshold for positive prediction of ≥ 0.5, on a scale of zero

to one. These parameters have been previously used to predict PPIs within fly and yeast [35].

Protein sequences and initial selection: To predict PPIs between *O. camponoti-*

floridani secreted proteins and *C. floridanus* proteins, we retrieved sequence information for both

organisms from high-quality genome assemblies integrating long-read technology (GenBank

accessions GCA_012980515.1 and GCA_003227725.1, respectively) [19,49]. For *Camponotus*

proteins, we used the sequence represented by the single longest isoform (n = 12,512). We

predicted *Ophiocordyceps* proteins to have a localization outside of the cell by combining eight

annotation tools (SignalP v6.0, TMHMM, Phobius, Prosite, PredGPI, NucPred, and TargetP

v2.0) [50–56] following the relatively conservative approaches outlined in Beckerson et al., 2019 [1].

We focused on secreted proteins (n = 586) and excluded transmembrane proteins (n = 1,301) in

our downstream analyses, as we considered these to most plausibly constitute effectors.

Moreover, our preliminary findings showed that transmembrane proteins were generally less

specific in their binding (Supplementary Discussion S1). All remaining non-secreted, non-

transmembrane proteins were considered intracellular (n = 5,568). We annotated secreted

proteins as uSSPs if they were under 300 amino acids in length and lacked any BLAST

description, GO term, or PFAM domain (n = 154) [19]. We used proteins ≤ 2,000 amino acids due

to computational constraints. Limiting the length of input proteins removed 63 (1.1%)

intracellular *Ophiocordyceps* proteins and 299 *Camponotus* proteins (2.4%).

PPI predictions: We used D-SCRIPT (v 0.1.5) in prediction mode with the default

human-protein pretrained model and settings to predict PPIs (as above for fungus-animal

benchmarking) [35]. We tested every PPI combination between secreted *O. camponoti-floridani*

proteins and the entire *C. floridanus* proteome (Fig. 1 step 1).

To identify self-interacting *Ophiocordyceps* PPIs, we tested secreted *Ophiocordyceps*

proteins against intracellular *Ophiocordyceps* proteins (Fig. 1 step 2). We then identified

reciprocal homologs (putative orthologs) between *Ophiocordyceps* and *Camponotus* proteins

using Proteinortho (v 5.0) with default settings [57]. We filtered out PPIs that contained a fungal

secreted protein predicted to bind both one of *Ophiocordyceps*' own intracellular proteins and a

Camponotus protein that was orthologous to that fungal intracellular protein (Fig. 1 step 3). We

reasoned that an interspecific PPI would more likely indicate an interaction related to infection or

manipulation when the *Ophiocordyceps* protein was predicted to only bind a *Camponotus*

protein, without also binding a fungal ortholog to that ant protein. We considered this to be a

favorable tradeoff with possibly overlooking parasite effectors that may target well-conserved

biological processes.

We narrowed the remaining PPIs to those involving secreted fungal proteins encoded by

differentially expressed genes (DEGs) (Fig. 1 step 4). These genes were upregulated during

manipulated summiting behavior in *C. floridanus* ants experimentally infected with *O.*

camponoti-floridani, as compared to their expression in culture [19]. By incorporating empirical

gene expression data, we anchored our computational analyses in predictions that are most

biologically relevant during manipulation. Although the chance exists that some parasite

effectors act during manipulation without increased gene transcription, we presumed that in most

cases, gene expression would be informative.

Hypergeometric enrichment analyses: We performed hypergeometric enrichment

analyses on proteins from PPIs with secreted, upregulated *Ophiocordyceps* proteins that only had

predicted interactions with the host. For enrichment analyses of the fungal proteins in these PPIs,

we used the *O. camponoti-floridani* secretome as the background. For enrichment analyses of the

ant proteins in these PPIs, we used the *C. floridanus* proteome as the background (Fig. 1, step 5).

We examined annotated GO terms (biological processes and molecular functions

combined) [58], PFAM domains [59], and weighted gene co-expression network analysis (WGCNA)

[60] module memberships reported in the transcriptomics study from which we also obtained the

above-mentioned DEGs [19]. These WGCNA modules correlated expression of gene-networks to

sample types in that study, i.e., (*i*) healthy ant controls, (*ii*) *in vitro* fungus controls, (*iii*) active

fungus-manipulated ants, and (*iv*) dying fungus-manipulated ants. These modules were

additionally correlated directly between *Ophiocordyceps* gene modules and *Camponotus* gene

modules [19]. We referenced their previously assigned names following the convention of ant-

module-1 (A1) and fungus-module-1 (F1) as used in that study [19]. For *Camponotus* proteins, we

also tested for enrichments of DEG class (i.e., upregulated or downregulated gene expression

from healthy controls to active manipulated ants) [19]. For *Ophiocordyceps* proteins, we

additionally tested for enrichment of uSSP annotations.

We performed the hypergeometric enrichment analyses with the R package

timecourseRNAseq (v 0.0.9000) using default settings (significant enrichment at FDR $\leq$ 0.05) in

R studio (v 2021.09.2) with R (v 4.1) [61–63]. We plotted GO terms in semantic space to cluster

related terms based on R code generated from REVIGO but did not remove any GO terms (R

package ggplot2, v 3.3.5) [64,65].

Aspecific fungus-*Camponotus* PPI combinations: For more selective analyses of

species-specific *Ophiocordyceps-Camponotus* PPIs, we produced aspecific PPIs from the

secretomes of other fungi tested against the *C. floridanus* proteome. Removal of more common

PPIs, leaving only those potentially unique to *Ophiocordyceps*, provided a view that was easier

to interpret in the context of specialized parasite-host interactions, such as behavioral

manipulation. However, a PPI need not be unique to a certain interspecific interaction to be

relevant [1]. Additionally, computationally predicted protein binding does not indicate that all such

interactions occur in dynamic physiological environments. In and around the cell, protein

localization, transcriptional control, post-translational modification, competing PPIs, and

biochemical conditions (e.g., pH) could affect the likelihood of a predicted PPI to occur. The

cellular environments within fungal cells and insect cells could plausibly differ in many ways,

shaping protein activity. Therefore, we used the removal of aspecific interactions to help us focus

on some of the, likely, more important PPIs involved in establishing *Ophiocordyceps*

manipulation of ant behavior, but still anchored our analyses and discussion on the larger dataset.

We chose three fungi based on lifestyle, phylogeny, and availability of well-annotated

genomes to generate aspecific PPIs: *C. bassiana* (GenBank accession GCA_000280675.1), *T.

reesei* (GenBank accession GCA_000167675.2), and *S. cerevisiae* (GenBank accession

GCF_000146045.2) [7,66,67]. As for *Ophiocordyceps* (see above), we predicted each fungal

secretome and removed secreted proteins over 2,000 amino acids long, which only removed one

T. reesei protein (0.002%). The secretomes of *C. bassiana*, *T. reesei*, and *S. cerevisiae* fungi

included 833, 505, and 153 proteins, respectively. These proteins were then paired with the

Camponotus proteome for D-SCRIPT predictions (see above).

To find orthologous proteins between *Ophiocordyceps* and each of the aspecific fungi we

used Proteinortho (v 6.0) with default settings while enforcing blastp+ and single-best reciprocal

hits (i.e., -sim=1) [57]. Using this information, we identified shared PPIs between *Ophiocordyceps*

and *Camponotus*, and aspecific fungi and *Camponotus*; i.e., interactions between orthologous

fungal proteins and the same ant protein (Supplementary Fig. S1). More stringently, we also

applied a filter that eliminated shared host proteins with aspecific fungal interactions, regardless

of orthology. This resulted in a set of *Ophiocordyceps* PPIs where the *Camponotus* protein was

never predicted in any aspecific PPI (Supplementary Fig. S1). After removing either shared PPIs

or shared host proteins, we again performed enrichment analyses (see above) to investigate

overrepresented annotations among the *Ophiocordyceps*-specific PPIs (Fig. 1 step 6). To

visualize changes in PPI connectivity and abundance that resulted from these aspecific filters, we

used Cytoscape (v 3.8.2) to produce networks of host proteins and their respective

Ophiocordyceps binding partners [68].

Results

Fungus-animal benchmarking PPIs: To evaluate D-SCRIPT's performance on

predicting interspecific PPIs between fungi and animals, we used experimentally supported

interactions from the Host-Pathogen Interaction Database [69]. Our evaluation dataset included 567

of these verified, positive interactions from a diversity of fungal and animal species. We

combined these with a 10-fold negative background (5,670 PPIs) constructed by randomly

pairing fungal (*S. cerevisiae*) and insect (*D. melanogaster*) proteins. For this evaluation with a 1

positive:10 negative ratio test data, the model performance metric, area under precision-recall

curve (AUPR), has a baseline of 0.091. D-SCRIPT produces results well above this level for fly-

only [35], yeast-only [35], and these fungus-animal data (AUPR 0.552, 0.405, and 0.373,

respectively) (Table 1). Although interspecific interactions indeed appear to be challenging to

predict, our benchmarking data (Table 1) suggest D-SCRIPT still provides meaningful insights

in interspecific contexts.

Ophiocordyceps-Camponotus **PPIs:** We tested a total of 7,156,818 potential interspecific

PPIs between the parasite secretome and host proteome (Fig. 1 step 1). Of these possible protein-

protein combinations, 0.33% were positively predicted as PPIs (n = 23,629 PPIs) (Table 2,

Supplementary Fig. S2). This result is in line with a previous 0.95% positive D-SCRIPT

prediction rate from a single-species dataset testing 50 million potential PPIs [35].

From these positively predicted interspecific interactions, we removed 2,850 PPIs that

were homologous to within-*Ophiocordyceps* interactions (Fig. 1 step 2 and 3, Table 2). We then

selected those involving *Ophiocordyceps* proteins encoded by upregulated DEGs during

manipulation. This resulted in 6,011 predicted PPIs. We used this set of putative effector-host

PPIs, containing 129 unique *Ophiocordyceps* proteins and 2,083 unique *Camponotus* proteins,

for further analyses (Fig. 1 step 4, Table 2, Supplementary File S1).

PPIs between aspecific fungi and *Camponotus* in comparison to *Ophiocordyceps*: The

proportions of positive PPI predictions between aspecific fungi and *C. floridanus* (range 0.26-

0.28%) and secreted proteins predicted to be in at least one PPI (range 68-76%) were similar to

the values observed for *Ophiocordyceps* (0.33% and 69%, respectively) (Table 2, Supplementary

Table S1). Eliminating orthologous PPIs between *Ophiocordyceps* and any of the aspecific fungi

(i.e., shared PPIs) removed 1,776 of the 6,011 (30%) *Ophiocordyceps-Camponotus* PPIs (Table

3, Supplementary Fig. S1). Eliminating PPIs involving host proteins that were also predicted to

be bound by aspecific fungi, regardless of orthology of the fungal binding partner (i.e., shared

host proteins), removed 1,936 of 2,083 (93%) host proteins, leaving only 157 *Ophiocordyceps*-

Camponotus PPIs (3%) (Table 3, Supplementary Fig. S1). In line with secretome size,

phylogeny, and entomopathogenic lifestyle, *C. bassiana* shared the most overlap with

Ophiocordyceps predictions (Supplementary Fig. S1). As expected, removing PPIs involving

shared host proteins led to major changes in enrichment results while removing shared PPIs led

to more modest changes (Table 3, Supplementary File S2). We primarily report the results based

on all three fungi that we included in this study. However, those based on *C. bassiana* alone were

largely similar (Supplementary Fig. S1, Supplementary File S2). While these filters assisted us in

ordering and emphasizing results (i.e., signals maintained through the strictest filters are being

prioritized), we did not use this approach to wholly select PPIs of interest. Results presented

below only use aspecific filters where stated.

Ophiocordyceps enrichment signals: We found that *Ophiocordyceps* proteins in PPIs

with *Camponotus* proteins were enriched for one PFAM domain and three WGCNA modules

(Fig. 2, Table 3).

S8 serine peptidases – The enriched PFAM domain was a "peptidase S8 domain,"

indicating an overrepresentation of fungal serine proteases that could cleave host proteins. The

five fungal peptidase S8 proteins in this enrichment were predicted to interact with 34 host

proteins across 39 PPIs (Supplementary File S1). Among these host proteins, there were seven

kinesin-like proteins (motor proteins), five nuclear pore proteins, and a (pro-)resilin that is an

insect cuticle and connective tissue protein [70]. Even after removing PPIs that were shared with

aspecific fungi, the enrichment signal for S8 peptidases was retained. However, the much more

stringent removal of shared host targets eliminated the S8 peptidase enrichment result (Fig. 1

step 6, Table 3).

Fungal manipulation WGCNA modules – We found fungal WGCNA modules F1, F2,

and F3 to be enriched (Fig. 2). We previously reported modules F1 and F2 to have significant

positive correlations to manipulation. This means that most of the fungal genes in those modules

were upregulated, in a similar co-expressed manner, during manipulated summiting behavior. As

such, these modules contained multiple genes hypothesized to mediate infection or manipulation

processes ("uSSPs and hypothesized effectors" modules) (Fig. 2). With fungal gene expression

clearly negatively correlated with control cultures, module F3 was modestly positively correlated

to fungi in both manipulated hosts and dying hosts. This module largely contained genes related

to "core cell processes" in the fungus (e.g., DNA packaging) (Fig. 2). All three fungal modules

also had significant direct correlations to ant WGCNA modules, with F1 being negatively

correlated to ant modules A14 and A15 – both associated with host neuronal processes (Fig. 2).

Core cell processes module F3 was positively correlated with neuronal process module A15 [19].

The overrepresentations of all three fungal WGCNA modules was retained after eliminating

shared PPIs, but uSSP and effectors module F1 was lost after removing shared host proteins

(Table 3).

Camponotus **enrichment signals:** We found that *Camponotus* PPI proteins were

enriched for 50 GO terms, 74 PFAM domains, and nine WGCNA modules (Table 3). As GO

term and PFAM domain enrichments often indicated similar functions, we present results largely

from the GO term perspective. We highlight PFAM domains for emphasis or to discuss

biologically interesting results not well captured by GO terms alone.

G protein-coupled receptor signaling – We detected GO term enrichments for "G

protein-coupled receptor (GPCR) activity" and "GPCR signaling pathway" (Fig. 3). These

enrichments resulted from 71 GPCR-related PPIs with 16 unique *Ophiocordyceps* proteins

(Supplementary File S1). All but two receptor proteins in these PPIs carried the PFAM domain

"7 transmembrane receptor (rhodopsin family)." We found 33 unique GPCR-related protein

genes (of 128 possible with these GO terms), with a range of putative ligands: 11 neuropeptides

(n = 12 receptors), four biogenic monoamines (dopamine, serotonin, octopamine, tyramine) (n =

11 receptors), acetylcholine (n = 2 receptors), a spider venom toxin protein (alpha-latrotoxin) (n

= 1 receptor), and opsin blue- and ultraviolet-sensitive receptors (n = 2 receptors). Fourteen of

these 33 *Camponotus* receptors (or receptor subunits) belonged in ant-neuronal processes

WGCNA modules A14 and A15, which showed overall reduced expression during manipulation

compared to healthy ants (i.e., the modules were negatively correlated to manipulation) [19]. Host

GPCR-related PPI proteins were predicted to bind 16 fungal proteins that included five uSSPs, a

carboxylesterase, a glycosyl hydrolase, a CAP cysteine-rich secretory protein, and a protein

tyrosine phosphatase. Most of these fungal proteins were previously assigned to fungal uSSPs

and effectors module F2 (n = 9) but included some in modules F1-4 (Supplementary File S1),

which all were also correlated to infection and manipulation of hosts.

The enrichment signals for GPCRs were retained after we removed both PPIs and host

proteins shared with aspecific fungi (Fig. 4). Twenty-eight of the 33 GPCR host proteins

remained after removing shared PPIs. However, only seven remained after filtering out shared

host proteins. These host proteins, predicted to be uniquely bound by *Ophiocordyceps* during

manipulation, were a putative dopamine receptor 2, Oamb octopamine receptor, methuselah-like

10 receptor, orexin receptor, trissin receptor, cAMP-dependent protein kinase subunit, and

receptor without an informative BLAST description ("uncharacterized protein"). The four

Ophiocordyceps proteins binding these host receptors were predicted in many PPIs (range = 101

to 458). The dopamine receptor 2, Oamb octopamine receptor, and uncharacterized receptor

shared the same fungal uSSP partner. The orexin and trissin receptors also shared an unannotated

secreted fungal protein (larger than an uSSP).

Oxidation-reduction – We also found the enrichment of oxidation-reduction functions

among *Camponotus* proteins in PPIs (six supporting GO terms) (Fig. 3, Supplementary File S1).

Altogether, 159 host proteins had oxidation-reduction annotations, forming 321 PPIs with 48

parasite proteins. A plurality of the host proteins were encoded by cytochrome P450 genes (70

unique genes, 15 putative subtypes). This abundance of cytochrome P450, a hemoprotein, also

contributed to the enrichment of "heme-binding" and "iron ion binding" GO terms (Fig. 3) and

the PFAM domain "cytochrome P450" (Supplementary File S1). These *Camponotus* cytochrome

P450 proteins paired with 10 unique *Ophiocordyceps* genes, consisting of uSSPs and fungal

cytochrome P450s. Other notable contributors to the host enrichment of oxidation reduction

related functions were peroxiredoxins, laccases, dehydrogenases, and a vat-1 synaptic vesicle

membrane-like protein (Supplementary File S1). The 48 fungal proteins included cytochrome

P450s, oxidases and oxidoreductases, amidases, peptidases, and an enterotoxin.

Enrichment signals for four of the six oxidation-reduction GO terms were retained after

we removed PPIs that were shared with the aspecific fungi. Only two remained after we applied

the more stringent shared host protein filter (Supplementary File S2). Additionally, indicators of

cytochromes, heme and iron binding GO terms and the "cytochrome P450" PFAM domain,

continued to be enriched after either filter (Supplementary File S2). Most host proteins passed

the shared PPI filter (140 of 159), but only 19 remained after removing shared host proteins (Fig.

4). Cytochrome P450 proteins shifted from a plurality to a dominant majority after shared host

protein filtering (17 of 19 proteins), with eight cytochrome P450 subtypes: 3A19, 315a1, 4C1,

49a1, 6a13, 6a14, 6k1, and 9e2.

Transcription and DNA-binding – Multiple enrichments that can be associated to gene

transcription and regulation were enriched. We found 204 host proteins annotated for any of

"DNA binding," "DNA-binding transcription factor activity," "sequence-specific DNA binding,"

"DNA helicase activity," "regulation of transcription, DNA-templated," and "mRNA binding"

GO terms (Fig. 3). These proteins similarly contributed to the high enrichment of PFAM

domains such as "helix-loop-helix DNA-binding domain," "zinc Finger C4 type," "'paired box'

domain," and "bZIP transcription factor" (Supplementary File S1). These 204 host proteins

interacted with 43 fungal proteins to generate 922 PPIs. Among the 204 host proteins, we found

transcription factors and regulators with many functions, which included examples associated

with insect behavior and neuronal function such as Mushroom body large type Kenyon specific

protein 1 (e.g., elevated in worker bees, foraging behavior), Gooseberry (e.g., neuromuscular

junction function), Hairy (e.g., juvenile hormone mediated gene repression), photoreceptor

specific nuclear receptor (light sensing), CLOCK (circadian clocks), and steroid and ecdysone

nuclear receptors and induced proteins [71–75]. Other proteins, most clearly understood from the

neuronal development perspective thus far, produce atypical behavioral or locomotor phenotypes

when dysregulated and include Jim Lovell, Dead ringer, Hairy/enhancer-of-split related,

Achaete-scute complex proteins, and Forkhead [76–81]. These 34 proteins taken as biologically

relevant examples account for 186 of the 922 gene regulation PPIs, interacting with 20

Ophiocordyceps proteins covering a similar breadth of protein types as the 43 parasite proteins

overall (Supplementary File S1). The 43 fungal proteins included five uSSPs that were also

found in the PPIs reported above (Fig. 4A), three that were not, and various hydrolases,

oxidoreductases, peptidases, and carboxylesterases.

All of the aforementioned GO terms and PFAM domains continued to be enriched after

removing shared PPIs (191 of 204 host proteins retained), but none of these annotations

continued to be enriched after removing shared aspecific host proteins (five host proteins

retained) (Fig. 4, Supplementary File S2).

Cuticular proteins – We found 28 ant proteins that shared the enriched GO term

"structural constituent of cuticle" (Fig. 3) and PFAM domain "insect cuticle protein"

(Supplementary File S1). We predicted these ant proteins interacted with 22 fungal proteins to

form 138 PPIs (Fig. 4A). These host proteins included (pro-)resilin, endocuticle structural

proteins, and a range of "cuticle proteins" (e.g., "cuticle protein 7"). The fungal proteins included

a CAP protein and a carboxylesterase also predicted to bind GPCRs (see above). Four fungal

uSSPs were shared with the transcription and DNA-binding host proteins. We additionally

identified a glycosyl hydrolase, a disintegrin/reprolysin-like peptidase, a S8 subtilisin-like serine

peptidase, M43 peptidases, an uSSP, and an unannotated secreted protein (larger than an uSSP).

These enrichments were only kept under the aspecific shared PPI filter (with all 28 original

cuticular proteins) and lost with the shared host protein filter (none of the 28 proteins were

uniquely bound by *Ophiocordyceps*) (Fig. 4, Supplementary File S2).

Ant DEGs and WGCNA modules – *Camponotus* genes that were previously

determined to be differentially expressed during manipulation, as well as host gene modules

correlated to manipulation, were enriched among PPI proteins (Supplementary File S1). Both up-

and downregulated host DEGs were enriched. The upregulated host DEGs were enriched for

gene/DNA regulatory functions and the downregulated DEGs for oxidation-reduction

(Supplementary File S1). Three ant WGCNA modules were enriched that had significant

correlations to manipulation, with an additional six that did not (of 22 possible modules total).

"Neuronal processes" module A14 was negatively correlated to manipulation and fungal uSSP

and effectors module F1 (Fig. 2). This host module included many genes putatively associated

with neuronal function, including GPCRs reported above [19]. The other host neuronal processes

module, A15, was not enriched for PPIs. Module A10 was positively correlated to manipulation

and contained enrichment signals for GO terms indicating "signal transduction and transcription"

[19] (Fig. 2). Module A4 had a positive correlation to manipulation and contained enrichment

signals for "proteasomes and odor detection" [19] (Fig. 2). All three of these host modules passed

the shared PPI filter but did not persist after removing shared host targets with aspecific fungi.

Discussion

The interspecific molecular interactions during manipulation of *C. floridanus* by *O.*

camponoti-floridani are likely mediated by parasite effector proteins that target host proteins. To

predict such PPIs, we used the machine learning tool D-SCRIPT, which infers structural

relationships between proteins without relying on protein-wide sequence homology to known

interactions. This flexibility allows D-SCRIPT models to extend predictions across organisms

and beyond previously documented PPIs. We benchmarked D-SCRIPT against a dataset

including experimentally supported fungus-animal PPIs. This evaluation indicated D-SCRIPT

can produce meaningful predictions on interspecific PPIs, although it performs better on single-

species data. D-SCRIPT predicted several thousand PPIs between *Camponotus* host proteins and

upregulated, secreted *Ophiocordyceps* parasite proteins. Even with low recall (identification of

true PPIs) functional enrichments and trends at the group level can plausibly be detected. Correct

PPIs predictions reflect a true biological signal while, most likely, incorrect-prediction noise

would not frequently produce coherent enrichments signals.

Due to the unfamiliar terrain of leveraging this approach between non-model species, we

sought to propose hypotheses balancing biological plausibility and the possibility of novel

interactions. On the one hand, we considered PPI predictions in the light of previous knowledge

of protein and cellular function. However, since the molecular aspects of parasitic manipulation

are still largely to be uncovered, we also considered that undescribed or unusual mechanisms

may underlie these interactions. Often, we found that single proteins, especially from

Ophiocordyceps, were in multiple host-parasite PPIs. Although in some cases these could be

errant predictions, *Ophiocordyceps* effectors may indeed bind many host proteins across

different pathways. This is a commonly observed phenomenon in microbe-plant interactions, and

is possibly even the norm [2,4]. Moreover, we found that only the S8 peptidase annotation term was

enriched among *Ophiocordyceps* secreted, upregulated proteins that were in PPIs with

Camponotus proteins. This finding indicates that the PPI proteins are not a sharply distinct subset

from the fungal secretome (i.e., background proteins of the enrichment analysis), which is

functionally enriched for pathogenesis and proteolytic activity overall [19]. In contrast, we detected

many enriched annotations for the host side of the interactions.

***Ophiocordyceps* proteolysis of core intracellular host proteins:** Fungal PPI proteins

were enriched for S8 peptidase domains, indicating that a significant number of these proteins

had proteolytic functions with possibly diverse targets [82]. Cuticular host proteins are a canonical

target for such effectors as we discuss below. Additionally, multiple host kinesin motor proteins

were in PPIs with these *Ophiocordyceps* proteases. Kinesins play a role in mitotic and

intracellular transport processes – with key functions in vesicle transport towards the periphery

of neurons [83]. Defects in kinesin activity can lead to dysregulated neurotransmitter release at

synapses, thereby contributing to impaired neurological functions and development [83,84]. From

these predicted S8 peptidase-kinesin interactions, we infer that the fungus may be dysregulating

core host cell processes. The effects of which possibly impair neuronal signaling and

neurotransmitter release to contribute to modified behavioral phenotypes. We also predicted

these peptidases to interact with multiple *Camponotus* nuclear pore proteins. Although some

components of the nuclear pore undergo proteolytic post-translational modification to remove

SUMOylation [85,86], those peptidases are in a different family than S8 peptidases. If the predicted

S8 peptidase-nucleopore proteins interactions are genuine, this could mean the parasite alters

transport of molecules in and out of the host nucleus. Speculatively, this could facilitate entry of

the fungal proteins that bind host transcription factors.

***Camponotus* G protein-coupled receptors as targets to modify behavior:** We detected

over 30 ant GPCRs or GPCR subunits in PPIs with fungal proteins. Most GPCRs were from the

rhodopsin/A family and included receptors related to biogenic monoamines, acetylcholine, and

neuropeptides. These receptors offer a wide range of hypothetical connections to manipulated ant

behavior. Commonly, these receptors and their ligands have been linked to locomotion, feeding,

and circadian rhythms – processes that have been previously highlighted in hypotheses of

Ophiocordyceps and other fungus-insect summit-disease behavioral manipulations [10,14,19,87,88].

The monoamine neurotransmitters dopamine, serotonin, octopamine (analogous to

vertebrate norepinephrine), and tyramine have been implicated in modulating locomotor,

foraging, learning, social, reproductive, and aggressive behaviors in many insects, with

experimental evidence in social insects, including ants [89–94]. Compared to the aspecific fungi

tested, both a *C. floridanus* dopamine and octopamine receptor were uniquely bound by

Ophiocordyceps upregulated secreted proteins. This interaction hypothetically implicates these host receptors in specific *Ophiocordyceps* interactions such as behavioral manipulation. We additionally detected two muscarinic acetylcholine receptors associated with sensory and motor neurons in non-insect animals [95–97]. In bees, excessive activation of acetylcholine receptors can cause changes in locomotor, navigational, foraging, and social behaviors [98]. Changes in such behaviors have also been observed in *C. floridanus* ants infected with *O. camponoti-floridani*. Locomotor behavior is enhanced in these ants, while social behaviors are diminished, daily timed foraging activities become arrhythmic and navigation capabilities appear less effective [17].

The neuropeptide GPCRs in PPIs had a range of putative ligands linking parasite interference with behavior-regulating processes such as circadian rhythm, nutritional signaling, insect hormones, and odor reception. These GPCRs included a host trissin receptor and an allatotropin receptor (annotated as an "orexin" receptor, but likely an allatotropin receptor in insects), which, compared to the aspecific fungi, only *Ophiocordyceps* was predicted to bind. Between these two receptors, dysregulation of feeding and circadian control of locomotor activity, juvenile hormone production, immune activation, and muscle stimulation are possible effects [99–105]. Circadian rhythm is related to ant foraging behavior (and, perhaps indirectly, feeding) and is affected by *Ophiocordyceps* infection [17]. The synchronized time-of-day summiting and biting behaviors further suggests a role for circadian control of activity [15,19,23]. Interlaced with feeding and foraging processes in ants, juvenile hormone mediates insulin/feeding related signaling and social caste identity (e.g., foraging workers) [28,106–108]. Among the host neuropeptide GPCR targets shared with the aspecific fungi, we predicted binding of receptors also associated with locomotion and activity levels, feeding, aggression, muscle control, juvenile hormone, reproductive behavior, or pheromones that included those

activated by: allatostatin-A [109–112], adipokinetic hormone or corazonin (i.e., a "gondaotropin-

releasing hormone" receptor) [113], cholecystokinin-like peptides [114], CCHamide-2 [115], and

pyrokinin, pheromone biosynthesis activating neuropeptide (PBAN), and/or capa type

neuropeptides [116–118]. Odor and pheromone detection are key elements of ant communication,

behavioral regulation, and social interactions [119–121]. GPCRs typically involved in odorant

response may become dysregulated by fungal effectors to dampen nestmate interaction and/or

induce aberrant behavior without normal cues. Taken together, we found multiple PPIs involving

ant GPCRs putatively involved in signaling systems that could control which and when certain

behaviors are performed by manipulated ants.

We also predicted PPIs involving a putative host cAMP-dependent protein kinase

catalytic subunit, found only between *Ophiocordyceps* and *Camponotus*, and a host G-protein

subunit alpha, both of which serve important roles as components of GPCRs. This alpha subunit

was also found previously to be downregulated in manipulated *Camponotus* [19]. *Ophiocordyceps*

interference with host GPCR signaling has been previously hypothesized to be mediated, at least

in part, by enterotoxins and other fungal protein toxins that interfere with cAMP levels and

signal transduction [14,19,24,25].

Given the signal for GPCR interference, with both our broadest and narrowest analyses,

and that many of these receptors/ligands overlap in their behavioral associations, we suggest that

this provides evidence for host-parasite GPCR PPIs as one mechanism of behavioral

manipulation used by *Ophiocordyceps*. The fungal counterparts in these host GPCR-parasite

effector PPIs were often predicted in many interactions. These fungal proteins included five

uSSPs upregulated during manipulation and correlated to manipulation in WGCNA modules [19].

Given that these fungal proteins lack annotations, we cannot yet determine if they might be

agonists or antagonists, nor whether they bind the ligand-site or elsewhere on the receptor. As

these uSSPs were not predicted to bind only these host proteins, their interaction with the

receptors may be less likely mediated by specific receptor-ligand binding sites than more generic

contact sites elsewhere on the host protein.

Host-pathogen interactions involving oxidation-reduction: We predicted over 300

PPIs that contributed to an enrichment of oxidation-reduction processes among the host proteins.

Oxidation-reduction processes can be involved in a range of metabolic, developmental, and host-

parasite interaction pathways that have been previously hypothesized to play a role in

Ophiocordyceps infections [14,19,122,123]. Approximately half of the host proteins were putative

cytochrome P450s, with enrichment signals for cytochrome P450s maintained after aspecific

fungi filtering. In insects, cytochrome P450 proteins are often involved in hormone metabolism

(e.g. ecdysone and juvenile hormone), stress response, xenobiotic detoxification processes, and

cuticular development [124–130]. For example, we found multiple cytochrome p450 proteins bound

uniquely by *Ophiocordyceps* of the 4C1 subtype that has been implicated in stress response (e.g.,

social isolation, temperature, UV, or starvation) and insecticide/xenobiotic detoxification, and is

regulated by hormones including juvenile hormone [126,131–134]. In turn, D-SCRIPT predicted that

many of those host cytochrome P450s interacted with fungal cytochrome P450s. Although

cytochrome P450 proteins often act independently, homo- and hetero-dimer formation of

cytochrome P450 proteins can modulate protein function in positive, negative, and substrate-

specific manners [135]. Oxidation-reduction related PPIs might most strongly represent the

physiological stress and antagonistic cellular processes between host and parasite. However,

fungal interference with these pathways may also be critical elements of creating a host

susceptible to manipulation.

Alteration of *Camponotus* gene regulation: We found signs that suggest

Ophiocordyceps could interfere with *Camponotus* gene regulation via proteins that bind host

transcription factors and DNA- or mRNA-binding proteins. Using the host's own cells to

dysregulate behavioral pathways could be metabolically favorable for the parasite. Some of the

fungal proteins in these PPIs were uSSPs and we, therefore, cannot predict the functional result

of those PPIs. However, several parasite proteins were hydrolases. We suggest that these

hydrolases most plausibly cleave host proteins or their post-translational modifications.

Depending on the role of the protein and specific bond broken, parasite hydrolase activity could

increase or decrease transcription of host genes. Among the ca. 200 host proteins in the

nucleotide-binding category, we found many tied to behavioral activity via changes in

locomotion, feeding/foraging, gravitaxis, light perception, circadian clocks, development and

social caste related hormones, and neuronal maintenance. Covering a similar breadth of

processes as the host GPCRs in PPIs, transcriptional interference may be a complementary

parasite strategy to modify their hosts. However, very few of these putative targets passed the

strictest shared aspecific host protein filter. As such, specific hypotheses about individual

transcription factors are difficult to parse. However, as a broader functional category, many of

these host proteins intersect with processes that appear relevant to manipulated ant behavior.

Destruction of *Camponotus* structural proteins: During manipulation, the fungus also

secretes proteins predicted to bind host cuticle and connective tissue proteins. Being an

endoparasite, *Ophiocordyceps* has direct access to internal host connective tissues, the

endocuticle layer (once the epidermis is breached), and the cuticular lining of the gut and

trachea, which include tissues that contain chitin [70,136]. However, the fungal cell wall also

contains chitin. As such, we cannot fully discount that some of these secreted fungal proteins

may simply be cell wall modifying proteins that have falsely been predicted to bind chitin-

associated ant proteins. Nevertheless, the fungus could be degrading cuticular proteins for

nutrition, detaching muscles, or in preparation for emerging hyphae to later grow out of the host

cadaver. Corroborating evidence for the disruption of the musculature stems from detailed

microscopy work on *Ophiocordyceps*-infected ants at the time of manipulated summiting and

biting behavior. These reports describe visual signs of fungal invasion, tissue degradation, and

atrophy of manipulated ant mandible muscles [15,22,29,137].

The predicted fungal interactors with host structural proteins included multiple proteases

such as a M43-like metalloprotease and S8 subtilisin-like serine peptidase. S8 peptidases can

degrade insect cuticle and are found in many fungi [138,139]. We also detected unannotated secreted

proteins in cuticular protein PPIs that could have similar or different functions. The aspecific

fungi included a generalist entomopathogen capable of degrading a wide variety of insect tissue

(*C. bassiana*); consistent with this, we found no *Ophiocordyceps*-specific cuticular-protein host

targets.

Gene networks and expression broadly implicate PPIs in manipulation:

Ophiocordyceps PPI proteins were enriched for WGCNA modules transcriptionally associated

with manipulation and containing uSSPs and other previously hypothesized effectors (F1 and F2)

[19]. In *Camponotus*, gene modules associated with neuronal processes and GPCRs (A14) and

signaling and transcription (A10) further support the involvement of these processes with

manipulation [19]. Assuming "guilt-by-association," in some cases this could indicate these PPIs

are related to infection and manipulation processes even if we yet lack clear mechanistic

hypotheses of how (e.g., PPIs with uSSPs) [140–142]. Additionally, both upregulated and

downregulated host DEGs [19] were enriched in predicted PPIs, possibly indicating homeostatic

responses to fungal modulation of functional protein levels. As such, the proteome-level

predictions that we made in this study are well in line with previous transcriptomics work.

Therefore, this work serves as an additional line of evidence for certain host processes and

narrows the pool of well-supported effector candidates involved with behavioral manipulation.

Conclusion: Faced with the challenge of uncovering how a parasitism with two non-

model species may operate at the molecular level, we combined multiple datatypes and tools to

predict host-parasite PPIs. These approaches are feasible in many study systems, even for those

that may lack the tools for sophisticated functional testing. We employed a structurally aware

machine learning program, D-SCRIPT, to predict PPIs beyond well-documented interactions. By

combining genomic and transcriptomic data we selected PPIs of interest with multiple

bioinformatic tools to: (*i*) annotate secreted fungal proteins, (*ii*) filter interactions predicted

across different species, and (*iii*) select proteins linked to increased gene expression during

infection. Using the resulting PPIs, we performed enrichment analyses to identify functional

categories that should be robust to computational error rates at the individual PPI level. Taken

together, we produced multiple hypotheses connecting host-parasite interspecific PPIs to

processes related to infection and manipulation of *Camponotus* by *Ophiocordyceps*. We

highlighted evidence for PPIs involving fungal S8 proteases and PPIs involving host oxidation-

reduction processes, cuticular proteins, gene regulation, and GPCRs. Especially, with

analytically robust results and direct biological links to behavior, we emphasize parasitic

dysregulation of GPCR signaling as a mechanism of parasitic behavioral manipulation.

Acknowledgements

We would like to thank Biplabendu Das, who was involved in discussions of project

goals and approaches, and Anna Savage, who provided constructive comments on a draft of this

manuscript.

Competing Interests

638   The authors declare no competing interests.

Data Availability

The genome assemblies used for this study are available on NCBI: *O. camponoti-*

floridani (GCA_012980515.1), *C. floridanus* (GCA_003227725.1), *C. bassiana*

(GCA_000280675.1), *T. reesei* (GCA_000167675.2), and *S. cerevisiae* (GCF_000146045.2)

[7,19,49,66,67]. Transcriptomic sequence data for *O. camponoti-floridani* and *C. floridanus* are also

hosted on NCBI (BioProject PRJNA600972) [19]. Analyzed transcriptomic data used here are

available on figshare, as supplemental information associated with their original publication

(https://doi.org/10.25387/g3.12121659) [19]. Prediction and enrichment results are available in

Supplementary Files S1 and S2 published with this article.

References

1.   Beckerson, W. C. *et al.* Cause and effectors: Whole-genome comparisons reveal shared

but rapidly evolving effector sets among host-specific plant-castrating fungi. *MBio* **10**,

(2019).

2.   Hogenhout, S. A., Van Der Hoorn, R. A. L., Terauchi, R. & Kamoun, S. Emerging

concepts in effector biology of plant-associated organisms. *Mol. Plant-Microbe Interact.*

22, 115–122 (2009).

3.   Wang, H., Peng, H., Li, W., Cheng, P. & Gong, M. The Toxins of Beauveria bassiana and
        the Strategies to Improve Their Virulence to Insects. *Front. Microbiol.* **12**, 2375 (2021).

4.   Win, J. *et al.* Effector Biology of Plant-Associated Organisms: Concepts and Perspectives.
        Cold Spring Harb. Symp. Quant. Biol. **77**, 235–247 (2012).

5.   Cen, K., Li, B., Lu, Y., Zhang, S. & Wang, C. Divergent LysM effectors contribute to the
        virulence of Beauveria bassiana by evasion of insect immune defenses. *PLOS Pathog.* **13**,
        e1006604 (2017).

6.   Wang, C. & Wang, S. Insect Pathogenic Fungi: Genomics, Molecular Interactions, and
        Genetic Improvements. *Annu. Rev. Entomol.* **62**, 73–90 (2017).

7.   Xiao, G. *et al.* Genomic perspectives on the evolution of fungal entomopathogenicity in
        Beauveria bassiana. *Sci. Rep.* **2**, 483 (2012).

8.   Araújo, J. P. M. & Hughes, D. P. Zombie-Ant Fungi Emerged from Non-manipulating,
        Beetle-Infecting Ancestors. *Curr. Biol.* **29**, 1–4 (2019).

9.   Andersen, S. B. *et al.* The Life of a Dead Ant: The Expression of an Adaptive Extended
        Phenotype. *Am. Nat.* **174**, 424–433 (2009).

10.   de Bekker, C., Beckerson, W. C. & Elya, C. Mechanisms behind the Madness: How Do
        Zombie-Making Fungal Entomopathogens Affect Host Behavior To Increase
        Transmission? *MBio* **12**, (2021).

11.   Lovett, B., St. Leger, R. J. & de Fine Licht, H. H. Going gentle into that pathogen-induced
        goodnight. *J. Invertebr. Pathol.* **174**, 107398 (2020).

12.   Andriolli, F. S. *et al.* Do zombie ant fungi turn their hosts into light seekers? *Behav. Ecol.*
        30, 609–616 (2019).

13.   Will, I., Linehan, S., Jenkins, D. G. & de Bekker, C. Natural history and ecological effects
on the establishment and fate of Florida carpenter ant cadavers infected by the parasitic
manipulator Ophiocordyceps camponoti-floridani. *Funct. Ecol.* **00**, 1–14 (2022).

14.   de Bekker, C. *et al.* Gene expression during zombie ant biting behavior reflects the
complexity underlying fungal parasitic behavioral manipulation. *BMC Genomics* **16**, 620
(2015).

15.   Hughes, D. P. *et al.* Behavioral mechanisms and morphological symptoms of zombie ants
dying from fungal infection. *BMC Ecol.* **11**, 13 (2011).

16.   Pontoppidan, M. B., Himaman, W., Hywel-Jones, N. L., Boomsma, J. J. & Hughes, D. P.
Graveyards on the move: The spatio-temporal distribution of dead Ophiocordyceps-
infected ants. *PLoS One* **4**, e4835 (2009).

17.   Trinh, T., Ouellette, R. & de Bekker, C. Getting lost: the fungal hijacking of ant foraging
behaviour in space and time. *Anim. Behav.* **181**, 165–184 (2021).

18.   Evans, H. C. Entomogenous fungi in tropical forest ecosystems: an appraisal. *Ecol.*
Entomol. 7–60 (1982).

19.   Will, I. *et al.* Genetic Underpinnings of Host Manipulation by Ophiocordyceps as
Revealed by Comparative Transcriptomics. *G3 (Bethesda).* **10**, 2275–2296 (2020).

20.   Beckerson, W. C., Krider, C., Mohammad, U. A. & Bekker, C. de. 28 Minutes Later:
Investigating the role of aflatrem-like compounds in Ophiocordyceps parasite
manipulation of zombie ants. *bioRxiv* (2023) doi:10.1101/2022.09.08.507134.

21.   Loreto, R. G. & Hughes, D. P. The metabolic alteration and apparent preservation of the
zombie ant brain. *J. Insect Physiol.* **118**, 103918 (2019).

22.   Zheng, S. *et al.* Specialist and Generalist Fungal Parasites Induce Distinct Biochemical

Changes in the Mandible Muscles of Their Host. *Int. J. Mol. Sci.* **20**, (2019).

23.   de Bekker, C. *et al.* Species-specific Ant Brain Manipulation by a Specialized Fungal

Parasite. *BMC Evol. Biol.* **14**, 166 (2014).

24.   de Bekker, C., Ohm, R. A., Evans, H. C., Brachmann, A. & Hughes, D. P. Ant-infecting

Ophiocordyceps genomes reveal a high diversity of potential behavioral manipulation

genes and a possible major role for enterotoxins. *Sci. Rep.* **7**, 12508 (2017).

25.   Kobmoo, N. *et al.* A genome scan of diversifying selection in Ophiocordyceps zombie-ant

fungi suggests a role for enterotoxins in co-evolution and host specificity. *Mol. Ecol.* **27**,

3582–3598 (2018).

26.   de Bekker, C. Ophiocordyceps–ant interactions as an integrative model to understand the

molecular basis of parasitic behavioral manipulation. *Curr. Opin. Insect Sci.* **33**, 19–24

(2019).

27.   de Bekker, C., Merrow, M. & Hughes, D. P. From behavior to mechanisms: An

integrative approach to the manipulation by a parasitic fungus (Ophiocordyceps

unilateralis s.l.) of its host ants (Camponotus spp.). *Integr. Comp. Biol.* **52**, 166–176

(2014).

28.   de Bekker, C. & Das, B. Hijacking time: How Ophiocordyceps fungi could be using ant

host clocks to manipulate behavior. *Parasite Immunol.* **44**, e12909 (2022).

29.   Mangold, C. A., Ishler, M. J., Loreto, R. G., Hazen, M. L. & Hughes, D. P. Zombie ant

death grip due to hypercontracted mandibular muscles. *J. Exp. Biol.* **222**, jeb200683

(2019).

30.   Agany, D. D. M. M., Pietri, J. E. & Gnimpieba, E. Z. Assessment of vector-host-pathogen

relationships using data mining and machine learning. *Comput. Struct. Biotechnol. J.* **18**,

1704 (2020).

31.   Soyemi, J., Isewon, I., Oyelade, J. & Adebiyi, E. Inter-Species/Host-Parasite Protein

727   Interaction Predictions Reviewed. *Curr. Bioinform.* **13**, 396 (2018).

728   32.   Gupta, S. K., Srivastava, M., Osmanoglu, Ö. & Dandekar, T. Genome-wide inference of

729   the Camponotus floridanus protein-protein interaction network using homologous

730   mapping and interacting domain profile pairs. *Sci. Rep.* **10**, 2334 (2020).

731   33.   Loaiza, C. D., Duhan, N., Lister, M. & Kaundal, R. In silico prediction of host–pathogen

732   protein interactions in melioidosis pathogen Burkholderia pseudomallei and human

733   reveals novel virulence factors and their targets. *Brief. Bioinform.* **22**, 1–18 (2021).

734   34.   Ma, S. *et al.* Prediction of protein–protein interactions between fungus (Magnaporthe

735   grisea) and rice (Oryza sativa L.). *Brief. Bioinform.* **20**, 448–456 (2019).

736   35.   Sledzieski, S., Singh, R., Cowen, L. & Berger, B. D-SCRIPT translates genome to

737   phenome with sequence-based, structure-aware, genome-scale predictions of protein-

738   protein interactions. *Cell Syst.* **12**, 969-982.e6 (2021).

739   36.   Chen, H. *et al.* Systematic evaluation of machine learning methods for identifying human–

740   pathogen protein–protein interactions. *Brief. Bioinform.* **22**, 1–21 (2021).

741   37.   Walhout, A. J. M. *et al.* Protein interaction mapping in C. elegans Using proteins involved

742   in vulval development. *Science (80-.).* **287**, 116–122 (2000).

743   38.   Hu, X., Feng, C., Ling, T. & Chen, M. Deep learning frameworks for protein–protein

744   interaction prediction. *Comput. Struct. Biotechnol. J.* **20**, (2022).

745   39.   Song, B. *et al.* Learning spatial structures of proteins improves protein–protein interaction

746   prediction. *Brief. Bioinform.* **23**, 1–11 (2022).

40.   Singh, R., Devkota, K., Sledzieski, S., Berger, B. & Cowen, L. Topsy-Turvy: integrating a

global view into sequence-based PPI prediction. *Bioinformatics* **38**, i264–i272 (2022).

41.   Yang, S., Feng, D., Cheng, P., Liu, Y. & Wang, S. Exploring the Knowledge of An

Outstanding Protein to Protein Interaction Transformer. *bioRxiv* (2023)

doi:10.1101/2023.02.09.527848.

42.   Chen, M. *et al.* Multifaceted protein–protein interaction prediction based on Siamese

residual RCNN. *Bioinformatics* **35**, i305–i314 (2019).

43.   Hwang, H., Vreven, T., Janin, J. & Weng, Z. Protein–protein docking benchmark version

4.0. *Proteins Struct. Funct. Bioinforma.* **78**, 3111–3114 (2010).

44.   Cheng, Q. *et al.* Discovery of a novel small secreted protein family with conserved N-

terminal IGY motif in Dikarya fungi. *BMC Genomics* **15**, 1–12 (2014).

45.   Feldman, D., Yarden, O. & Hadar, Y. Seeking the Roles for Fungal Small-Secreted

Proteins in Affecting Saprophytic Lifestyles. *Front. Microbiol.* **11**, 455 (2020).

46.   Fischer, R. & Requena, N. Small-secreted proteins as virulence factors in nematode-

trapping fungi. *Trends Microbiol.* (2022) doi:10.1016/J.TIM.2022.03.005.

47.   Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for

the analysis of massive data sets. *Nat. Biotechnol. 2017 3511* **35**, 1026–1028 (2017).

48.   Szklarczyk, D. *et al.* The STRING database in 2023: protein–protein association networks

and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids*

Res. **51**, D638 (2023).

49.   Shields, E. J., Sheng, L., Weiner, A. K., Garcia, B. A. & Bonasio, R. High-Quality

Genome Assemblies Reveal Long Non-coding RNAs Expressed in Ant Brains. *Cell Rep.*

23, 3078–3090 (2018).

50.   Teufel, F. *et al.* SignalP 6.0 predicts all five types of signal peptides using protein

language models. *Nat. Biotechnol. 2022 407* **40**, 1023–1025 (2022).

51. Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E. L. . Predicting transmembrane protein topology with a hidden markov model: application to complete genomes. *J. Mol. Biol.* **305**, 567–580 (2001).

52. Käll, L., Krogh, A. & Sonnhammer, E. L. L. A Combined Transmembrane Topology and Signal Peptide Prediction Method. *J. Mol. Biol.* **338**, 1027–1036 (2004).

53. Sigrist, C. J. A. *et al.* PROSITE: A documented database using patterns and profiles as motif descriptors. *Brief. Bioinform.* **3**, 265–274 (2002).

54. Pierleoni, A., Martelli, P. & Casadio, R. PredGPI: A GPI-anchor predictor. *BMC Bioinformatics* **9**, 1–11 (2008).

55. Brameier, M., Krings, A. & MacCallum, R. M. NucPred—Predicting nuclear localization of proteins. *Bioinformatics* **23**, 1159–1160 (2007).

56. Emanuelsson, O., Nielsen, H., Brunak, S. & Von Heijne, G. Predicting Subcellular Localization of Proteins Based on their N-terminal Amino Acid Sequence. *J. Mol. Biol.* **300**, 1005–1016 (2000).

57. Lechner, M. *et al.* Proteinortho: Detection of (Co-)orthologs in large-scale analysis. *BMC Bioinformatics* **12**, 1–9 (2011).

58. Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25–29 (2000).

59. Finn, R. D. *et al.* Pfam: the protein families database. *Nucleic Acids Res.* **42**, D222–D230 (2014).

60. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559 (2008).

794   61.   Das, B. timecourseRnaseq: Analyses And Visualisation Of Timecourse RNASeq Data. at

795   https://github.com/biplabendu/timecourseRnaseq (2022).

796   62.   R Core Team. R: A language and environment for statistical computing. at https://www.r-

797   project.org/ (2021).

798   63.   RStudio Team. RStudio: Integrated Development for R. at http://www.rstudio.com/

799   (2015).

800   64.   Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO Summarizes and Visualizes

801   Long Lists of Gene Ontology Terms. *PLoS One* **6**, e21800 (2011).

802   65.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, 2016).

803   66.   Engel, S. R. *et al.* New data and collaborations at the Saccharomyces Genome Database:

804   updated reference genome, alleles, and the Alliance of Genome Resources. *Genetics* **220**,

805   (2022).

806   67.   Martinez, D. *et al.* Genome sequencing and analysis of the biomass-degrading fungus

807   Trichoderma reesei (syn. Hypocrea jecorina). *Nat. Biotechnol. 2008 265* **26**, 553–560

808   (2008).

809   68.   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of

810   biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

811   69.   Ammari, M. G., Gresham, C. R., McCarthy, F. M. & Nanduri, B. HPIDB 2.0: a curated

812   database for host–pathogen interactions. *Database* **2016**, 103 (2016).

813   70.   Burrows, M. & Sutton, G. P. Locusts use a composite of resilin and hard cuticle as an

814   energy store for jumping and kicking. *J. Exp. Biol.* **215**, 3501–3512 (2012).

815   71.   Guo, Z., Qin, J., Zhou, X. & Zhang, Y. Insect Transcription Factors: A Landscape of

816   Their Structures and Biological Functions in Drosophila and beyond. *Int. J. Mol. Sci.* **19**,

(2018).

72.   Liu, F. *et al.* Mblk-1 regulates sugar responsiveness in honey bee (Apis mellifera)

foragers. *Insect Sci.* **29**, 683–690 (2022).

73.   Marie, B., Pym, E., Bergquist, S. & Davis, G. W. Synaptic Homeostasis Is Consolidated

by the Cell Fate Gene gooseberry, a Drosophila pax3/7 Homolog. *J. Neurosci.* **30**, 8071–

8082 (2010).

74.   Saha, T. T. *et al.* Hairy and Groucho mediate the action of juvenile hormone receptor

Methoprene-tolerant in gene repression. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E735–E743

(2016).

75.   Velarde, R. A., Robinson, G. E. & Fahrbach, S. E. Nuclear receptors of the honey bee:

annotation and expression in the adult brain. *Insect Mol. Biol.* **15**, 583–595 (2006).

76.   Bjorum, S. M. *et al.* The Drosophila BTB Domain Protein Jim Lovell Has Roles in

Multiple Larval and Adult Behaviors. *PLoS One* **8**, 61270 (2013).

77.   Cheng, Y. C. *et al.* The transcription factor hairy/E(spl)-related 2 induces proliferation of

neural progenitors and regulates neurogenesis and gliogenesis. *Dev. Biol.* **397**, 116–128

(2015).

78.   García-Bellido, A. & De Celis, J. F. The Complex Tale of the achaete–scute Complex: A

Paradigmatic Case in the Analysis of Gene Organization and Function During

Development. *Genetics* **182**, 631 (2009).

79.   Häcker, U. *et al.* The Drosophila fork head domain protein crocodile is required for the

establishment of head structures. *EMBO J.* **14**, 5306 (1995).

80.   Shandala, T., Takizawa, K. & Saint, R. The dead ringer/retained transcriptional regulatory

gene is required for positioning of the longitudinal glia in the Drosophila embryonic CNS.

Development **130**, 1505–1513 (2003).

81.   Tang, B. *et al.* Forkhead box protein p1 is a transcriptional repressor of immune signaling

in the CNS: implications for transcriptional dysregulation in Huntington disease. *Hum.*

Mol. Genet. **21**, 3097 (2012).

82.   Muszewska, A. *et al.* Fungal lifestyle reflected in serine protease repertoire. *Sci. Reports*

2017 71 **7**, 1–12 (2017).

83.   Klinman, E. & Holzbaur, E. L. F. Walking Forward with Kinesin. *Trends Neurosci.* **41**,

555–556 (2018).

84.   Liu, L., Downs, M., Guidry, J. & Wojcik, E. J. Inter-organelle interactions between the

ER and mitotic spindle facilitates Zika protease cleavage of human Kinesin-5 and results

in mitotic defects. *iScience* **24**, 102385 (2021).

85.   Hang, J. & Dasso, M. Association of the Human SUMO-1 Protease SENP2 with the

Nuclear Pore. *J. Biol. Chem.* **277**, 19961–19966 (2002).

86.   Zhang, H., Saitoh, H. & Matunis, M. J. Enzymes of the SUMO modification pathway

localize to filaments of the nuclear pore complex. *Mol. Cell. Biol.* **22**, 6498–6508 (2002).

87.   de Bekker, C., Will, I., Hughes, D. P., Brachmann, A. & Merrow, M. Daily rhythms and

enrichment patterns in the transcriptome of the behavior-manipulating parasite

Ophiocordyceps kimflemingiae. *PLoS One* **12**, (2017).

88.   Elya, C. *et al.* Robust manipulation of the behavior of Drosophila melanogaster by a

fungal pathogen in the laboratory. *Elife* **7**, 1–34 (2018).

89.   Adamo, S. A. Norepinephrine and octopamine: linking stress and immune function across

phyla. *ISJ* **5**, 12–19 (2008).

90.   Adamo, S. A., Linn, C. E. & Beckage, N. E. Parasites: evolution's neurobiologists. *J. Exp.*

Biol. **216**, 3–10 (2013).

91.   Aonuma, H. & Watanabe, T. Changes in the content of brain biogenic amine associated

with early colony establishment in the Queen of the ant, Formica japonica. *PLoS One* **7**,

e43377–e43377 (2012).

92.   Finetti, L., Roeder, T., Calò, G. & Bernacchia, G. The Insect Type 1 Tyramine Receptors:

From Structure to Behavior. *Insects* **12**, 315 (2021).

93.   Kamhi, J. F., Arganda, S., Moreau, C. S. & Traniello, J. F. A. Origins of aminergic

regulation of behavior in complex insect social systems. *Front. Syst. Neurosci.* **11**, 74

(2017).

94.   Verlinden, H. Dopamine signalling in locusts and other insects. *Insect Biochem. Mol. Biol.*

97, 40–52 (2018).

95.   Lee, Y. S. *et al.* Characterization of GAR-2, a Novel G Protein-Linked Acetylcholine

Receptor from Caenorhabditis elegans. *J. Neurochem.* **75**, 1800–1809 (2000).

96.   Stankiewicz, M. *et al.* Effects of a centipede venom fraction on insect nervous system, a

native Xenopus oocyte receptor and on an expressed Drosophila muscarinic receptor.

Toxicon **37**, 1431–1445 (1999).

97.   Tsentsevitsky, A. N., Kovyazina, I. V., Nurullin, L. F. & Nikolsky, E. E. Muscarinic

cholinoreceptors (M1-, M2-, M3- and M4-type) modulate the acetylcholine secretion in

the frog neuromuscular junction. *Neurosci. Lett.* **649**, 62–69 (2017).

98.   Grünewald, B. & Siefert, P. Acetylcholine and Its Receptors in Honeybees: Involvement

in Development and Impairments by Neonicotinoids. *Insects* **10**, (2019).

99.   Hernaândez-Martõânez, S. *et al.* Allatotropin: A pleiotropic neuropeptide that elicits

mosquito immune responses. *PLoS One* **12**, e0175759 (2017).

100.   Lismont, E. *et al.* Molecular cloning and characterization of the allatotropin precursor and

receptor in the desert locust, Schistocerca gregaria. *Front. Neurosci.* **9**, 84 (2015).

101.   Nagata, S., Matsumoto, S., Mizoguchi, A. & Nagasawa, H. Identification of cDNAs

encoding allatotropin and allatotropin-like peptides from the silkworm, Bombyx mori.

Peptides **34**, 98–105 (2012).

102.   Nässel, D. R. & Zandawala, M. Recent advances in neuropeptide signaling in Drosophila,

from genes to physiology and behavior. *Prog. Neurobiol.* **179**, 101607 (2019).

103.   Petri, B., Homberg, U., Loesel, R. & Stengl, M. Evidence for a role of GABA and Mas-

allatotropin in photic entrainment of the circadian clock of the cockroach Leucophaea

maderae. *J. Exp. Biol.* **205**, 1459–1469 (2002).

104.   Abruzzi, K. C. *et al.* RNA-seq analysis of Drosophila clock and non-clock neurons reveals

neuron-specific cycling and novel candidate neuropeptides. *PLOS Genet.* **13**, e1006613

(2017).

105.   Roller, L. *et al.* Molecular cloning, expression and identification of the promoter

regulatory region for the neuropeptide trissin in the nervous system of the silkmoth

Bombyx mori. *Cell Tissue Res.* **364**, 499–512 (2016).

106.   Schwinghammer, M. A., Zhou, X., Kambhampati, S., Bennett, G. W. & Scharf, M. E. A

novel gene from the takeout family involved in termite trail-following behavior. *Gene*

474, 12–21 (2011).

107.   Ament, S. A., Corona, M., Pollock, H. S. & Robinson, G. E. Insulin signaling is involved

in the regulation of worker division of labor in honey bee colonies. *Proc. Natl. Acad. Sci.*

U. S. A. **105**, 4226–31 (2008).

108.   Chandra, V. *et al.* Social regulation of insulin signaling and the evolution of eusociality in

ants. *Science (80-.).* **361**, 398–402 (2018).

109.   Aguilar, R. *et al.* Allatostatin gene expression in brain and midgut, and activity of

synthetic allatostatins on feeding-related processes in the cockroach Blattella germanica.

Regul. Pept. **115**, 171–177 (2003).

110.   Chen, J. *et al.* Allatostatin A Signalling in Drosophila Regulates Feeding and Sleep and Is

Modulated by PDF. *PLOS Genet.* **12**, e1006346 (2016).

111.   Hergarden, A. C., Tayler, T. D. & Anderson, D. J. Allatostatin-A neurons inhibit feeding

behavior in adult Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 3967–3972 (2012).

112.   Tsao, C. H., Chen, C. C., Lin, C. H., Yang, H. Y. & Lin, S. Drosophila mushroom bodies

integrate hunger and satiety signals to control innate food-seeking behavior. *Elife* **7**,

(2018).

113.   Sakai, T. *et al.* Invertebrate gonadotropin-releasing hormone-related peptides and their

receptors: An update. *Front. Endocrinol. (Lausanne).* **8**, (2017).

114.   Nässel, D. R. & Williams, M. J. Cholecystokinin-Like Peptide (DSK) in Drosophila, Not

Only for Satiety Signaling. *Front. Endocrinol. (Lausanne).* **5**, (2014).

115.   Ren, G. R. *et al.* CCHamide-2 Is an Orexigenic Brain-Gut Peptide in Drosophila. *PLoS*

One **10**, e0133017 (2015).

116.   Choi, M.-Y., Vander Meer, R. K., Coy, M. & Scharf, M. E. Phenotypic impacts of PBAN

RNA interference in an ant, Solenopsis invicta, and a moth, Helicoverpa zea. *J. Insect*

Physiol. **58**, 1159–1165 (2012).

117.   Jurenka, R. & Nusawardani, T. The pyrokinin/ pheromone biosynthesis-activating

neuropeptide (PBAN) family of peptides and their receptors in Insecta: evolutionary trace

indicates potential receptor ligand-binding domains. *Insect Mol. Biol.* **20**, 323–334 (2011).

118.   Ragionieri, L. *et al.* Identification of mature peptides from pban and capa genes of the

moths Heliothis peltigera and Spodoptera littoralis. *Peptides* **94**, 1–9 (2017).

119.   Chang, H. *et al.* Pheromone binding proteins enhance the sensitivity of olfactory receptors

to sex pheromones in Chilo suppressalis. *Sci. Rep.* **5**, 13093 (2015).

120.   Ferguson, S. T., Park, K. Y., Ruff, A. A., Bakis, I. & Zwiebel, L. J. Odor coding of

nestmate recognition in the eusocial ant Camponotus floridanus. *J. Exp. Biol.* **223**, (2020).

121.   Gotzek, D., Shoemaker, D. D. & Ross, K. G. Molecular variation at a candidate gene

implicated in the regulation of fire ant social behavior. *PLoS One* **2**, e1088 (2007).

122.   Dubovskii, I. M. *et al.* Generation of reactive oxygen species and activity of antioxidants

in hemolymph of the moth larvae Galleria mellonella (L.) (Lepidoptera: Piralidae) at

development of the process of encapsulation. *J. Evol. Biochem. Physiol. 2010 461* **46**, 35–

43 (2010).

123.   Iwanicki, N. S. A., Delalibera, I., Eilenberg, J. & de Fine Licht, H. H. Comparative

RNAseq Analysis of the Insect-Pathogenic Fungus Metarhizium anisopliae Reveals

Specific Transcriptome Signatures of Filamentous and Yeast-Like Development. *G3*

Genes|Genomes|Genetics **10**, 2141–2157 (2020).

124.   Iga, M. & Kataoka, H. Recent Studies on Insect Hormone Metabolic Pathways Mediated

by Cytochrome P450 Enzymes. *Biol. Pharm. Bull.* **35**, 838–843 (2012).

125.   Scott, J. G. & Wen, Z. Cytochromes P450 of insects: the tip of the iceberg. *Pest Manag.*

Sci. **57**, 958–967 (2001).

126.   Su, L., Yang, C., Meng, J., Zhou, L. & Zhang, C. Comparative transcriptome and

metabolome analysis of Ostrinia furnacalis female adults under UV-A exposure. *Sci.*

Reports | **11**, 6797 (2021).

127.   Sztal, T. *et al.* A Cytochrome P450 Conserved in Insects Is Involved in Cuticle Formation.

956   *PLoS One* **7**, e36544 (2012).

957   128.   Wang, H. *et al.* CYP6AE gene cluster knockout in Helicoverpa armigera reveals role in

958   detoxification of phytochemicals and insecticides. *Nat. Commun. 2018 91* **9**, 1–8 (2018).

959   129.   Xing, X. *et al.* Cytochrome P450s Are Essential for Insecticide Tolerance in the

960   Endoparasitoid Wasp Meteorus pulchricornis (Hymenoptera: Braconidae). (2021)

961   doi:10.3390/insects.

962   130.   Zhang, B. zhong *et al.* Insecticide induction of O-demethylase activity and expression of

963   cytochrome P450 genes in the red imported fire ant (Solenopsis invicta Buren). *J. Integr.*

964   *Agric.* **15**, 135–144 (2016).

965   131.   Kim Lien, N. T. *et al.* Transcriptome Sequencing and Analysis of Changes Associated

966   with Insecticide Resistance in the Dengue Mosquito (Aedes aegypti) in Vietnam. *Am. J.*

967   *Trop. Med. Hyg.* **100**, 1240 (2019).

968   132.   Lu, K.-H. H., Bradfield, J. Y. & Keeley, L. L. Juvenile hormone inhibition of gene

969   expression for cytochrome P4504C1 in adult females of the cockroach, Blaberus

970   discoidalis. *Insect Biochem. Mol. Biol.* **29**, 667–673 (1999).

971   133.   Scharf, I. *et al.* Social isolation causes downregulation of immune and stress response

972   genes and behavioural changes in a social insect. *Mol. Ecol.* **30**, 2378–2389 (2021).

973   134.   Shen, X., Liu, W., Wan, F., Lv, Z. & Guo, J. The Role of Cytochrome P450 4C1 and

974   Carbonic Anhydrase 3 in Response to Temperature Stress in Bemisia tabaci. *Insects* **12**,

975   (2021).

976   135.   Kandel, S. E. & Lampe, J. N. Role of Protein–Protein Interactions in Cytochrome P450-

977   Mediated Drug Metabolism and Toxicity. *Chem. Res. Toxicol.* **27**, 1474 (2014).

136. Davies, R. G. Insect structure and function. in *Outlines of Entomology* 7–96 (Springer,

Dordrecht, 1988). doi:10.1007/978-94-009-1189-5_2.

137. Fredericksen, M. A. *et al.* Three-dimensional visualization and a deep-learning model

reveal complex fungal parasite networks in behaviorally manipulated ants. *Proc. Natl.*

Acad. Sci. **114**, 201711673 (2017).

138. Arnesen, J. A. *et al.* Early Diverging Insect-Pathogenic Fungi of the Order

Entomophthorales Possess Diverse and Unique Subtilisin-Like Serine Proteases. *G3* **8**,

3311–3319 (2018).

139. St Leger, R., Joshi, L., Bidochka, M. J. & Roberts, D. W. Construction of an improved

mycoinsecticide overexpressing a toxic protease. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 6349–

54 (1996).

140. Carlson, M. R. J. *et al.* Gene connectivity, function, and sequence conservation:

Predictions from modular yeast co-expression networks. *BMC Genomics* **7**, 1–15 (2006).

141. Gillis, J. & Pavlidis, P. "Guilt by Association" Is the Exception Rather Than the Rule in

Gene Networks. *PLOS Comput. Biol.* **8**, e1002444 (2012).

142. Stuart, J. M., Segal, E., Koller, D. & Kim, S. K. A gene-coexpression network for global

discovery of conserved genetic modules. *Science (80-.).* **302**, 249–255 (2003).

Author Contributions

997   I.W., W.C.B., and C.dB. conceived the project and wrote the manuscript. W.C.B.

998 performed the secretome annotation. I.W. performed PPI predictions, enrichment analyses, and

999 visualization. This work was supported by the National Science Foundation (CAREER IOS-

1000 1941546 to C.dB).

1001

Tables

Table 1. Fungus-animal PPI D-SCRIPT benchmark.

We evaluated D-SCRIPT on interspecific fungus-animal PPIs using positive interactions from the Host-Pathogen

Interaction Database (n = 567) and negative interactions from randomly paired yeast and fly proteins (n = 5,670).

For this dataset and those published with D-SCRIPT [35] with 1 postive:10 negatives, the baseline AUPR is 0.091.

Recall is the proportion of true positive examples in the dataset that the model correctly predicted to interact.

Precision is the proportion of predicted positives that were indeed correct. Based on these benchmarking data,

interspecific PPIs are more difficult for D-SCRIPT to predict, but, we suggest, they still can be detected at rates

useful for initial biological investigation and characterization of group-level trends.

Evaluation dataset	AUPR	Precision	Recall
Fly, as published [35]	0.552	0.798	0.359
Yeast, as published [35]	0.405	0.706	0.223
Fungus-animal	0.373	0.720	0.169

1011

1012 **Table 2. PPI sets overview.**

1013 We tested 7,156,818 combinations of secreted *O. camponoti-floridani* (*O.*) proteins (n = 586) with every *C.*

1014 *floridanus* (*C.*) protein (n = 12,213). Predicted PPIs were combinations returned with an edge value of ≥ 0.5. We

1015 then created PPI subsets based on homologous interaction between secreted and intracellular fungal proteins and

1016 fungal gene upregulation during manipulation.

PPI set	Predicted PPIs (% of protein combinations tested)	*Ophiocordyceps* proteins in PPIs	*Camponotus* proteins in PPIs
All secreted *O. & C.*	23,629 (0.33%)	402	3289
no homologous fungal-self PPIs	20,779 (0.29%)	399	3143

| no homologous fungal-self PPIs | 6,011 (0.08%) | 129 | 2083 |
| & *O.* upregulated | | | |

1017

Table 3. Annotation enrichments of PPI proteins.

We performed enrichment analyses on interspecific PPIs without homology to *Ophiocordyceps* self-interactions and

involving upregulated, secreted *Ophiocordyceps* proteins and any *Camponotus* proteins (Fig. 1 step 5) with or

without additional filtering based on shared aspecific fungal interactions (Fig. 1 step 6). We found no enrichment

signal for uSSPs in the fungus. Only in the unfiltered analysis were *Camponotus* DEGs overrepresented – for both

upregulated and downregulated genes. The number of gene co-expression modules enriched are given under column

WGCNA. While typically more stringent filtering reduced the number of enrichments detected, it increased the

number of enriched GO terms for *Ophiocordyceps*. These enrichments included oxidation-reduction terms and

various general binding terms (e.g., "binding," "catalytic activity," "cofactor binding," or "ion binding").

Test proteins (n)	GO	PFAM	WGCNA
Ophiocordyceps (129)	0	1	3
after shared PPIs removed (125)	1	1	3
after shared host proteins removed (26)	11	0	2
Camponotus (2,083)	50	74	9
after shared PPIs removed (1,901)	44	64	9
after shared host proteins removed (147)	12	3	2

1027

 Figure legends

 Figure 1. Conceptual framework for PPI testing, selection, and analysis.

 Step 1) We tested every secreted *O. camponoti-floridani* protein (secretome) with every *C. floridanus* host protein

 (proteome). Steps 2, 3) If an interspecific PPI and an *Ophiocordyceps* self-interaction PPI shared a common fungal

 protein paired with a protein homologous between both species, we removed that PPI from our predictions. Step 4)

 Subsequently, we filtered predicted PPIs to focus on those that involved fungal proteins encoded by genes that were

 upregulated during manipulation. Step 5) We analyzed host and parasite proteins from these PPIs with

 hypergeometric enrichment analyses. For enrichment analyses, the background protein set was either the parasite

 secretome (*Ophiocordyceps*) or host proteome (*Camponotus*). Step 6) We also performed enrichment analyses after

 removing PPIs or host proteins that were predicted in interactions between aspecific fungi and *C. floridanus*. Fungal

 proteins are shown as circles and ant proteins as squares. Proteins are color-coded, with *Ophiocordyceps*

 upregulated, secreted proteins in interspecific PPIs indicated in red and their interacting *Camponotus* proteins in

 bright blue. Shades of orange represent other secreted fungal protein categories while blue-gray indicates other host

 proteins in PPIs. All other proteins are shown in gray. Proteins "A" are *Ophiocordyceps* and *Camponotus* homologs,

 and protein "B" is a single fungal protein.

 Figure 2. Enriched WGCNA modules and their correlations to manipulation and each other.

 Module membership, correlations, and functional summary of modules were based on previous work [19]. Each

 module is a mutually exclusive network of genes that are coexpressed across control, manipulated ant, and dying ant

 conditions. Here, we only depict modules enriched among PPIs and only their correlations to the manipulated state

 or between host-parasite modules directly. Core cell processes module F3 was clearly negatively correlated to

 control conditions, with modest positive correlations divided between both live manipulation and dying ants [19], as

 indicated by "+" here. While module F1 and F3 were also notably correlated to neuronal process module A15, this

 ant module is not depicted here as it was not enriched among the PPIs detected in this study.

 Figure 3. Enrichments for *C. floridanus* proteins in PPIs with upregulated secreted fungal proteins. GO terms

 are plotted in semantic space, which lacks any inherent unit or value other than to cluster terms by functional

 similarity. Some labels were omitted for readability and when not relevant to our discussion of the results.

1056

Figure 4. PPI network with host proteins contributing to selected GO term enrichments.

Host proteins (blue square nodes) are clustered and labeled by their general GO term functional category and

connected (gray line edges) to their fungal PPI partners that are either secreted proteins (small pink circle nodes) or

uSSPs (large red triangle nodes) that were found to be upregulated during *Ophiocordyceps* manipulation of

Camponotus behavior. Proteins may have PPI connections outside of those depicted here. a) PPI network without

aspecific PPI filtering. b) Network of remaining PPIs after removing shared PPIs with aspecific fungi. Although

nearly 30% of PPIs (edges) were removed, most proteins (nodes) and enrichment results were kept. c) Network of

remaining PPIs after removing PPIs with all host proteins that also interacted with aspecific fungi, regardless of

orthology of the fungal binding partner. GO terms related to G-protein coupled receptor signaling and some

oxidation-reduction enrichments persisted after this filter. Enrichment signals for transcription and DNA-binding

were lost, but some individual proteins remained.

1068

Supplementary Information

Discussion S1: A Word document briefly describing preliminary analysis of transmembrane

fungal proteins.

Figure S1: Filtering aspecific PPIs and host proteins shared between *Ophiocordyceps* and

alternative fungi.

Figure S2: D-SCRIPT edge values for tested protein pairings between extracellular fungal

proteins and the ant proteome.

Table S1: Aspecific fungi and *Ophiocordyceps* PPI overview.

File S1: *Ophiocordyceps-Camponotus* PPI protein and enrichment data in a single Excel file

with multiple tabs.

File S2: Aspecific fungi PPI data in a single Excel file with multiple tabs.

Ophiocordyceps

Camponotus

Proteome

Secretome

Secreted in PPIs

Upregulated, secreted in PPIs

Proteome

In PPIs with secreted fungal proteins

In PPIs with upregulated secreted fungal proteins

1. *Ophiocordyceps-Camponotus* **initial interspecific PPI predictions:** fungal secretome against host proteome

2. *Ophiocordyceps* **self-PPI predictions:** secreted against intracellular proteins

3. Remove: homologous PPIs

4. Select: upregulated fungal protein PPIs

5. Enrichment analyses

6. "Aspecific" fungi filters: Remove shared PPIs or host proteins

6. "Aspecific" fungi filters: Remove shared PPIs or host proteins

a

Oxidation-
reduction

GPCR
signaling

Cuticular
proteins

Transcription & DNA-binding

b

Oxidation-
reduction

GPCR
signaling

Transcription & DNA-binding

c

Oxidation-
reduction

GPCR
signaling

Transcription &
DNA-binding